Some identities for the Narayana polynomial and its derivatives

Nguyen Tien Dung^{*†}

June 18, 2022

Abstract

In this note, by employing a nice property of semicircular distributions, we derive some identities for the Narayana polynomial and its derivatives.

Keywords: Narayana polynomials, Semicircular distribution. 2020 Mathematics Subject Classification: 11B83, 60C05.

1 Introduction

The Narayana numbers N(r, k) and the Narayana polynomials $\mathcal{N}_r(z)$ are defined by

$$N(r,k) = \frac{1}{r} \binom{r}{k-1} \binom{r}{k}, \qquad (1.1)$$

$$\mathcal{N}_{r}(z) = \sum_{k=1}^{r} N(r,k) z^{k-1}.$$
(1.2)

Because of their applications in Combinatorics theory, the Narayana numbers and the Narayana polynomials have been investigated by several authors. There is a large number of combinatorial properties can be found in the literature, see [1, 2, 3, 4] and references therein.

The main result of this paper can be formulated as follows.

Theorem 1.1. For every $r \geq 3$, we have

$$\mathcal{N}_{r}(z) = \frac{2r-1}{r+1}(1+z)\mathcal{N}_{r-1}(z) - \frac{r-2}{r+1}(1-z)^{2}\mathcal{N}_{r-2}(z), \qquad (1.3)$$

$$\mathcal{N}_{r}'(z) = \frac{r-1}{2z} \mathcal{N}_{r}(z) + \frac{(r-1)(z-1)}{2z} \mathcal{N}_{r-1}(z), \qquad (1.4)$$

$$\mathcal{N}_{r}''(z) = -\frac{r-1}{z^{2}}\mathcal{N}_{r}(z) + \left(\frac{(r-1)^{2}}{z} + \frac{r-1}{z^{2}}\right)\mathcal{N}_{r-1}(z).$$
(1.5)

*Department of Mathematics, VNU University of Science, Vietnam National University, Hanoi, 334 Nguyen Trai, Thanh Xuan, Hanoi, 084 Vietnam. Email: dung@hus.edu.vn

[†]Department of Mathematics, FPT University, Hoa Lac High Tech Park, Hanoi, Vietnam.

The proof of the above theorem is deferred to Section 2. Let us end up this Section with some remarks.

Remark 1.1. In his paper [3], Lasalle established the following recurrence

$$(z+1)\mathcal{N}_r(z) - \mathcal{N}_{r+1}(z) = \sum_{n\geq 1} (-z)^n \binom{r-1}{2n-1} A_n \mathcal{N}_{r-2n+1}(z), \tag{1.6}$$

where the numbers A_n satisfies the recurrence

$$(-1)^{n-1}A_n = C_n + \sum_{j=1}^{n-1} (-1)^j \binom{2n-1}{2j-1} A_j C_{n-j},$$

with $A_1 = 1$ and $C_n = \frac{1}{n+1} {\binom{2n}{n}}$ the Catalan number. Interestingly, our relation (1.3) is simpler than (1.6).

Remark 1.2. From (1.3) and (1.4) we obtain the following recurrence for the first order derivative

$$\mathcal{N}_{r}'(z) = \frac{r-1}{2r+2} \frac{r-2+3rz}{z} \mathcal{N}_{r-1}(z) - \frac{(r-1)(r-2)}{2r+2} \frac{(1-z)^{2}}{z} \mathcal{N}_{r-2}(z).$$

Similarly, we can express $\mathcal{N}_{r}''(z)$ in term of $\mathcal{N}_{r-1}(z)$ and $\mathcal{N}_{r-2}(z)$.

Remark 1.3. It is known that $C_r := \mathcal{N}_r(1), r \ge 1$ is the sequence of the Catalan numbers. Hence, by choosing z = 1, we obtain from (1.3), (1.4) and (1.5) the following identities

$$C_r = \frac{4r - 2}{r + 1}C_{r-1},$$
$$\sum_{k=1}^r (k - 1)N(r, k) = \frac{r - 1}{2}C_r,$$
$$\sum_{k=1}^r (k - 1)(k - 2)N(r, k) = -(r - 1)C_r + r(r - 1)C_{r-1}.$$

Similarly, by choosing z = 2, we also obtain the identities for the sequence of the large Schröder numbers $S_r := \frac{1}{2} \mathcal{N}_r(2), r \ge 1$.

2 Proofs

Let X be a random variable with the semicircular distribution, i.e. X has a density function f(x) given by

$$f(x) = \begin{cases} \frac{2}{\pi}\sqrt{1-x^2} & \text{if } -1 \le x \le 1\\ 0 & \text{otherwise.} \end{cases}$$
(2.1)

It is known from [1] that the Narayana polynomial can be expressed as follows

$$\mathcal{N}_r(z) = \mathbb{E}[(1+z+2\sqrt{z}X)^{r-1}],$$
 (2.2)

where \mathbb{E} denotes the expectation operator.

In order to prove Theorem 1.1, we need to use the following property of semicircular distributions.

Lemma 2.1. For any continuously differentiable function h, we have

$$\mathbb{E}[h(X)X] = \frac{1}{3}\mathbb{E}[h'(X)(1-X^2)].$$

Proof. We observe that

$$F(x) := \int_{x}^{1} yf(y)dy = \frac{2}{\pi} \int_{x}^{1} y\sqrt{1-y^{2}}dy = \frac{2}{3\pi}(1-x^{2})\sqrt{1-x^{2}}, \ -1 \le x \le 1.$$

Hence, by integrating by parts, we obtain

$$\mathbb{E}[h(X)X] = \int_{-1}^{1} h(x)xf(x)dx = -\int_{-1}^{1} h(x)dF(x)$$
$$= \int_{-1}^{1} F(x)h'(x)dx = \int_{-1}^{1} \frac{1}{3}(1-x^2)h'(x)f(x)dx$$
$$= \frac{1}{3}\mathbb{E}[h'(X)(1-X^2)].$$

The proof of the lemma is complete.

We now are ready to present the proof of Theorem 1.1. Proof of (1.3). Using the representation (2.2), we have

$$\mathcal{N}_{r}(z) = \mathbb{E}[(1+z+2\sqrt{z}X)^{r-2}(1+z+2\sqrt{z}X)]$$

= $(1+z)\mathcal{N}_{r-1}(z) + 2\sqrt{z}\mathbb{E}[(1+z+2\sqrt{z}X)^{r-2}X]$

and hence,

$$\mathcal{N}_{r}(z) - (1+z)\mathcal{N}_{r-1}(z) = 2\sqrt{z}\mathbb{E}[(1+z+2\sqrt{z}X)^{r-2}X].$$
(2.3)

Furthermore, by using Lemma 2.1 and the relation (2.3), we deduce

$$\mathcal{N}_{r}(z) - (1+z)\mathcal{N}_{r-1}(z) = \frac{4(r-2)}{3}z\mathbb{E}[(1+z+2\sqrt{z}X)^{r-3}(1-X^{2})].$$

This gives us the following

$$z\mathbb{E}[(1+z+2\sqrt{z}X)^{r-3}X^2] = -\frac{3}{4(r-2)}\left(\mathcal{N}_r(z) - (1+z)\mathcal{N}_{r-1}(z)\right) + z\mathcal{N}_{r-2}(z).$$
(2.4)

Once again we use the representation (2.2) to get

$$\mathcal{N}_{r}(z) = \mathbb{E}[(1+z+2\sqrt{z}X)^{r-3}(1+z+2\sqrt{z}X)^{2}]$$

= $(1+z)^{2}\mathcal{N}_{r-2}(z) + 4(1+z)\sqrt{z}\mathbb{E}[(1+z+2\sqrt{z}X)^{r-3}X] + 4z\mathbb{E}[(1+z+2\sqrt{z}X)^{r-3}X^{2}]$

which, together with (2.3) and (2.4), yields

$$\mathcal{N}_{r}(z) = (1+z)^{2} \mathcal{N}_{r-2}(z) + 2(1+z) \left(\mathcal{N}_{r-1}(z) - (1+z) \mathcal{N}_{r-2}(z) \right) - \frac{3}{r-2} \left(\mathcal{N}_{r}(z) - (1+z) \mathcal{N}_{r-1}(z) \right) + 4z \mathcal{N}_{r-2}(z).$$

So it holds that

$$\frac{r+1}{r-2}\mathcal{N}_r(z) = \frac{2r-1}{r-2}(1+z)\mathcal{N}_{r-1}(z) - (1-z)^2\mathcal{N}_{r-2}(z)$$

and (1.3) follows.

Proof of (1.4). It follows from the representation (2.2) that

$$\mathcal{N}'_r(z) = (r-1)\mathbb{E}\left[(1+z+2\sqrt{z}X)^{r-2} \left(1+\frac{X}{\sqrt{z}}\right) \right],$$

which implies that

$$\mathcal{N}'_{r}(z) - (r-1)\mathcal{N}_{r-1}(z) = \frac{r-1}{\sqrt{z}} \mathbb{E}\left[(1+z+2\sqrt{z}X)^{r-2}X \right].$$
(2.5)

Combining (2.3) and (2.5) yields

$$\mathcal{N}'_{r}(z) - (r-1)\mathcal{N}_{r-1}(z) = \frac{r-1}{2z} \left(\mathcal{N}_{r}(z) - (1+z)\mathcal{N}_{r-1}(z) \right).$$

So (1.4) follows.

Proof of (1.5). We have

$$\begin{split} N_r''(z) &= (r-1)(r-2)\mathbb{E}\left[(1+z+2\sqrt{z}X)^{r-3} \left(1+\frac{X}{\sqrt{z}}\right)^2 \right] \\ &- \frac{r-1}{2}\mathbb{E}\left[(1+z+2\sqrt{z}X)^{r-2}\frac{X}{z\sqrt{z}} \right] \\ &= (r-1)(r-2)\mathbb{E}\left[(1+z+2\sqrt{z}X)^{r-3} \left(1+\frac{2X}{\sqrt{z}}+\frac{X^2}{z}\right) \right] \\ &- \frac{r-1}{2}\mathbb{E}\left[(1+z+2\sqrt{z}X)^{r-2}\frac{X}{z\sqrt{z}} \right]. \end{split}$$

Hence, recalling the relations (2.3) and (2.4), we obtain

$$\mathcal{N}_{r}''(z) = (r-1)(r-2) \left(\mathcal{N}_{r-2}(z) + \frac{1}{z} \left(\mathcal{N}_{r-1}(z) - (1+z)\mathcal{N}_{r-2}(z) \right) - \frac{3}{4(r-2)z^{2}} \left(\mathcal{N}_{r}(z) - (1+z)\mathcal{N}_{r-1}(z) \right) + \frac{1}{z}\mathcal{N}_{r-2}(z) \right) - \frac{r-1}{4z^{2}} \left(\mathcal{N}_{r}(z) - (1+z)\mathcal{N}_{r-1}(z) \right).$$

Rearranging the terms we get

$$\mathcal{N}_{r}''(z) = -\frac{r-1}{z^{2}}\mathcal{N}_{r}(z) + \left(\frac{(r-1)^{2}}{z} + \frac{r-1}{z^{2}}\right)\mathcal{N}_{r-1}(z).$$

Thus the proof of (1.5) is complete.

References

- [1] T. Amdeberhan, V. H. Moll, C. Vignat, A probabilistic interpretation of a sequence related to Narayana polynomials. *Online J. Anal. Comb.* No. 8 (2013), 25 pp.
- [2] W. Y. C. Chen, L. X. W. Wang, A. L. B. Yang, Schur positivity and the q-log-convexity of the Narayana polynomials. J. Algebraic Combin. 32 (2010), no. 3, 303–338.
- [3] M. Lassalle, Two integer sequences related to Catalan numbers. J. Combin. Theory Ser. A 119 (2012), no. 4, 923–935.
- [4] T. Mansour, Y. Sun, Identities involving Narayana polynomials and Catalan numbers. Discrete Math. 309 (2009), no. 12, 4079–4088.