ON DEFORMATIONS OF ISOLATED SINGULARITY FUNCTIONS

AURELIO MENEGON AND MIRIAM DA SILVA PEREIRA ´

Abstract. We study multi-parameters deformations of isolated singularity functiongerms on either a subanalytic set or a complex analytic spaces. We prove that if such a deformation has no coalescing of singular points, then it has constant topological type. This extends some classical results due to Lê $\&$ Ramanujam (1976) and Parusiński (1999), as well as a recent result due to Jesus-Almeida and the first author. It also provides a sufficient condition for a one-parameter family of complex isolated singularity surfaces in \mathbb{C}^3 to have constant topological type. On the other hand, for complex isolated singularity families defined on an isolated determinantal singularity, we prove that μ -constancy implies constant topological type.

INTRODUCTION

The study of deformations of maps and spaces plays a central role in Singularity Theory and in Complex Geometry. For holomorphic isolated singularity function-germs on a complex affine space \mathbb{C}^n , the topological behavior of small perturbations is related to the Milnor number, which is an analytic invariant with a very clear topological and algebraic meaning. In fact, when $n \neq 3$ there is a classical result due to Lê & Ramanujam [\[4\]](#page-7-0) in 1976, which states that a family $f_s: (\mathbb{C}^n,0) \to (\mathbb{C},0)$ is topologically trivial if and only if the Milnor number $\mu(f_s)$ is constant, for $s \in \mathbb{C}$ sufficiently small.

The case $n = 3$ is one of the main open problems in Singularity Theory, known as the μ -constant conjecture. Here, topological triviality means that for each s small enough there exists a homeomorphism of germs $h_s : (\mathbb{C}^n,0) \to (\mathbb{C}^n,0)$ that takes the germ of $V(f_s) := f_s^{-1}(0)$ at the origin onto the germ of $V(f_0) := f_0^{-1}(0)$ at the origin.

In 1999, Parusiński $[6]$ extended Lê & Ramanujam's result for K-analytic isolated singularity families $f_s: (\mathbb{K}^n,0) \to (\mathbb{K},0)$, with $\mathbb{K} = \mathbb{C}$ or \mathbb{R} , even when $n = 3$, provided that the family depends linearly on the parameter $s \in \mathbb{K}$. This means that it has the form $f_s = f_0 + s\varphi$, for some K-analytic isolated singularity function-germ $\varphi: (\mathbb{K}^n,0) \to (\mathbb{K},0).$

Recently, Jesus-Almeida and the first author [\[5\]](#page-7-1) studied K-analytic families f_s : $(X,0) \to (\mathbb{K},0)$ depending linearly on one parameter $s \in \mathbb{K}$, where $(X,0) \subset (\mathbb{K}^N,0)$ is the germ of a subanalytic set, when $\mathbb{K} = \mathbb{R}$, or a complex analytic space, when $\mathbb{K} = \mathbb{C}$. In particular, it was proved that if such a family has no coalescing of singular points, then it has constant topological type. Here, having no coalescing of singular points means that there exist real numbers $\delta > 0$ and $\epsilon > 0$ such that $\Sigma(f_s) \cap \mathbb{B}_{\epsilon} = \{0\}$ for every $s \in \mathbb{D}_{\delta}$, where $\Sigma(f_s)$ denotes the singular locus of f_s (in the stratified sense, with respect to a given (w)-regular stratification) and \mathbb{B}_{ϵ} denotes the closed ball around 0 in K^N. Also, having constant topological type means that for each $s \in (\mathbb{K}, 0)$ there exists a homeomorphism of germs $h_s: (V(f_s), 0) \to (V(f_0), 0)$. Clearly, this is weaker than having topological triviality in the sense of $\text{Le } \& \text{Ramanujam}.$

The first author was supported by the *Conselho Nacional de Desenvolvimento Científico e Tec* $nológico$ (CNPq), Brazil.

The first goal of this paper is to study K-analytic isolated singularity families $f_{s_1,...,s_k}$: $(X, 0) \to (\mathbb{K}, 0)$ depending linearly on k-many parameters $s_1, \ldots, s_k \in \mathbb{K}$, with $k > 1$. Then this will allow us to study one-parameter K-analytic isolated singularity families $f_s: (X,0) \to (\mathbb{K},0)$ that depend polynomially on $s \in \mathbb{K}$, which means it has the form

$$
\mathbf{f}_s = f_0 + s^{a_1}\varphi_1 + s^{a_2}\varphi_2 + \cdots + s^{a_k}\varphi_k,
$$

with $0 < a_1 < \cdots < a_k$. This will be done by setting

$$
f_{s_1,...,s_k} := f_0 + s_1\varphi_1 + s_2\varphi_2 + \cdots + s_k\varphi_k,
$$

so that $\mathbf{f}_s = f_{s^{a_1},...,s^{a_k}}$.

Our main results are the following:

In Section 1, we give an easy criterium to assure that a K-analytic family $f_{s_1,...,s_k}$: $(X,0) \to (\mathbb{K},0)$ has isolated singularity or even no coalescing of critical points. This is Proposition [1.2.](#page-3-0) In Section 2, we prove:

Theorem 0.1. Let $(X,0) \subset (\mathbb{K}^N,0)$ be the germ of either a subanalytic set, when $\mathbb{K} = \mathbb{R}$, or a complex analytic space, when $\mathbb{K} = \mathbb{R}$. Let S be a (w)-regular stratification of a representative X of $(X,0)$. Let (f_{s_1,\dots,s_k}) be a family of K-analytic isolated singularity function-germs on $(X, 0)$ depending linearly on the parameters $s_1, \ldots, s_k \in \mathbb{K}$. If (f_{s_1,\ldots,s_k}) has no coalescing of singular points (with respect to S) then it has constant topological type.

As an important consequence, we have:

Theorem 0.2. Let $f_s : (X, 0) \to (\mathbb{K}, 0)$ be a one-parameter \mathbb{K} -analytic isolated singularity family depending polynomially on $s \in \mathbb{K}$. Let $f_{s_1,...,s_k} : (X,0) \to (\mathbb{K},0)$ be the multi-parameter family associated to (f_s) , defined as above. If (f_{s_1,\dots,s_k}) has no coalescing of singular points, then (f_s) has constant topological type.

Corollary 0.2 relates to the μ -constant conjecture, as it provides a sufficient condition for a one-parameter family of complex isolated singularity surfaces in \mathbb{C}^3 to have constant topological type. Precisely, we have:

Corollary 0.3. Let $f_s : (\mathbb{C}^3, 0) \to (\mathbb{C}, 0)$ be an analytic isolated singularity family depending polynomially on $s \in \mathbb{C}$. Let $f_{s_1,...,s_k} : (\mathbb{C}^3,0) \to (\mathbb{C},0)$ be the multi-parameter family associated to (\mathbf{f}_s) , as above. If $(f_{s_1,...,s_k})$ is μ -constant, then (\mathbf{f}_s) has constant topological type.

The case when the family $f_s : (\mathbb{C}^3, 0) \to (\mathbb{C}, 0)$ depends analytically on $s \in \mathbb{C}$ is analogous, by a well-known result of Samuel $[7]$ (see also $[4]$, Theorem 1.7).

In Section 3, we restrict our attention to the particular case of complex analytic isolated singularity families $f_{s_1,\dots,s_k} : (X,0) \to (\mathbb{C},0)$ depending linearly on the parameter $s_1, \ldots, s_k \in \mathbb{C}$, with $k \geq 1$, where $(X,0) \subset (\mathbb{C}^N,0)$ is the germ of a complex isolated determinantal singularity (IDS). In this case, there is a well-defined Milnor number $\mu(f_s)$ in the sense of [\[1\]](#page-7-2).

We prove:

Theorem 0.4. Let $(X,0) \subset (\mathbb{C}^N,0)$ be the germ of a complex isolated determinantal singularity. Let $f_{s_1,...,s_k} : (X,0) \to (\mathbb{C},0)$ be a family of isolated singularity functiongerms depending linearly on $s_1, \ldots, s_k \in \mathbb{C}$, with $k \geq 1$. If (f_{s_1,\ldots,s_k}) is μ -constant, then it has constant topological type.

As a consequence, like before, we have:

Corollary 0.5. Let $(X,0) \subset (\mathbb{C}^N,0)$ be the germ of a complex isolated determinantal singularity. Let $f_s : (X, 0) \to (\mathbb{C}, 0)$ be an analytic isolated singularity family depending polynomially on $s \in \mathbb{C}$, and let $f_{s_1,\dots,s_k} : (X,0) \to (\mathbb{C},0)$ be the multi-parameter family associated to (\mathbf{f}_s) , as above. If $(f_{s_1,...,s_k})$ is μ -constant, then (\mathbf{f}_s) has constant topological type.

1. Isolated singularity families and coalescing of singular points

Let $(X, 0) \subset (\mathbb{K}^N, 0)$ be as above, and let $\mathcal{S} = (\mathcal{S}_{\alpha})_{\alpha \in \Lambda}$ be a (w) -regular stratification of a representative X of $(X, 0)$, in the sense of $[8]$. Without losing generality, we can suppose that $\{0\}$ is a stratum of S.

Let $\tilde{f}_0 : (\mathbb{K}^N, 0) \to (\mathbb{K}, 0)$ be an analytic function-germ and let $f_0 : (X, 0) \to (\mathbb{K}, 0)$ be the restriction of \tilde{f}_0 to $(X, 0)$. Clearly, the singular locus $\Sigma(f_0)$ of f_0 is formed by $X \cap \Sigma(\tilde{f}_0)$, the intersection of X with the critical set of \tilde{f}_0 , together with the points $x \in X \setminus \Sigma(\tilde{f}_0)$ such that the smooth manifold $(\tilde{f}_0)^{-1}(\tilde{f}_0(x))$ intersects $\mathcal{S}_{\alpha(x)}$ not transversally at x in \mathbb{K}^N , where $\mathcal{S}_{\alpha(x)}$ denotes the stratum of the stratification S that contains the point x.

Notice that since we chose a stratification S such that $\{0\}$ is a stratum, the origin $0 \in X \subset \mathbb{K}^N$ is a singular point of f_0 , even if \tilde{f}_0 is a submersion at $0 \in \mathbb{K}^N$.

We say that $f_0 : (X, 0) \to (\mathbb{K}, 0)$ has an *isolated singularity* if there exists a positive real number $\epsilon > 0$ such that $\Sigma(f_0) \cap \mathbb{B}_{\epsilon} = \{0\}$. This means that the restriction of f_0 to each stratum $S_{\alpha} \neq \{0\}$ is a submersion.

Now, if $\tilde{f}_s : (\mathbb{K}^N, 0) \to (\mathbb{K}, 0)$ is a deformation of \tilde{f}_0 depending continuously on the parameter $s \in (\mathbb{K}^k, 0)$, it induces a deformation $f_s : (X, 0) \to (\mathbb{K}, 0)$ of f_0 that obviously depends continuously on s . In the case when f_0 has an isolated singularity, the next lemma relates the critical set of f_s with the critical set of \tilde{f}_s inside some small neighborhood of $0 \in \mathbb{K}^N$, whenever $||s||$ is small enough.

We have:

Lemma 1.1. Suppose that $f_0 : (X, 0) \to (\mathbb{K}, 0)$ has an isolated singularity, and let $\epsilon > 0$ be such that $\Sigma(f_0) \cap \overline{\mathbb{B}}_{\epsilon} = \{0\}$. Then there exists $\delta > 0$ sufficiently small such that for every $s \in \mathbb{D}_{\delta}$ one has that

$$
\Sigma(f_s) \cap \overline{\mathbb{B}}_{\epsilon} = \left(\Sigma(\tilde{f}_s) \cap X \cap \overline{\mathbb{B}}_{\epsilon}\right) \cup \{0\}.
$$

Proof. Clearly, one has that $\Sigma(\tilde{f}_s) \cap X \subset \Sigma(f_s)$, so one inclusion is obvious. On the other hand, for every $x \in X \cap \overline{\mathbb{B}}_{\epsilon}$ fixed, one has that either x is the origin or $(\tilde{f}_0)^{-1}(\tilde{f}_0(x))$ is smooth at x and it intersects $\mathcal{S}_{\alpha(x)}$ transversally at x in \mathbb{K}^N . Since transversality is a stable property, there exists $\delta(x) > 0$ sufficiently small such that for every $s \in \mathbb{D}_{\delta(x)}$ one has that either $x \in \Sigma(\tilde{f}_s) \cup \{0\}$ or $(\tilde{f}_s)^{-1}(\tilde{f}_s(x))$ is smooth at x and it intersects $\mathcal{S}_{\alpha(x)}$ transversally at x in \mathbb{K}^N . Moreover, there exists a small neighborhood V_x of x in $X \cap \overline{\mathbb{B}}_{\epsilon}$ such that for every $x' \in V_x$ and for every $s \in \mathbb{D}_{\delta(x)}$ one has that either $x' \in \Sigma(\tilde{f}_s) \cup \{0\}$ or $(\tilde{f}_s)^{-1}(\tilde{f}_s(x'))$ is smooth at x' and it intersects $\mathcal{S}_{\alpha(x)}$ transversally at x' in \mathbb{K}^N . Therefore, since $X \cap \overline{\mathbb{B}}_{\epsilon}$ is compact, there exists $\delta > 0$ such that for every $x \in X \cap \overline{\mathbb{B}}_{\epsilon}$ and for every $s \in \mathbb{D}_{\delta}$ one has that either $x \in \Sigma(\tilde{f}_s) \cup \{0\}$ or $x \notin \Sigma(f_s)$. Hence $\Sigma(f_s) \cap \overline{\mathbb{B}}_{\epsilon} \subset \left(\Sigma(\tilde{f}_s) \cap X \cap \overline{\mathbb{B}}_{\epsilon}\right) \cup \{0\}.$

We say that a family (f_s) as above is an *isolated singularity family* if for each $s \in$ K with $||s||$ sufficiently small, the function-germ $f_s : (X,0) \to (\mathbb{K},0)$ has isolated singularity. That is, if there exists $\delta > 0$ such that for every $s \in \mathbb{D}_{\delta}$ there is $\epsilon(s) > 0$ such that $\Sigma(f_s) \cap \mathbb{B}_{\epsilon(s)} = \{0\}.$

An isolated singularity family (f_s) as above is said to have no coalescing of singular points if there exist real numbers $\delta > 0$ and $\epsilon > 0$ such that $\Sigma(f_s) \cap \mathbb{B}_{\epsilon} = \{0\}$ for every $s \in \mathbb{D}_{\delta}$.

As an immediate consequence of Lemma [1.1,](#page-2-0) we have:

Proposition 1.2. Suppose that $f_0 : (X, 0) \to (\mathbb{K}, 0)$ has isolated singularity. Then:

- (i) If (\tilde{f}_s) is an isolated singularity family, then (f_s) is also an isolated singularity family.
- (ii) If (\tilde{f}_s) has no coalescing of critical points, then (f_s) has no coalescing of singular points.

Example 1.3. Consider $X = \{x^2 - y^2 = 0\} \subset \mathbb{K}^3$ with the stratification given by $\mathcal{S}_0 = X \setminus \{x = y = 0\}$ and $\mathcal{S}_1 = \{x = y = 0\}$. Consider the family of function-germs $f_s: (X,0) \to (\mathbb{K},0)$ given by

$$
f_s(x, y, z) = x^3 + y^4 + z^2 + sx^4 + s^2y^5.
$$

Since f_0 has an isolated singularity and since the family (\tilde{f}_s) on \mathbb{K}^3 is an isolated singularity family with no coalescing of critical points, it follows from Theorem [1.2](#page-3-0) that (f_s) is an isolated singularity family with no coalescing of singular points.

2. One-parameter families of isolated singularities

Let $(X, 0) \subset (\mathbb{K}^N, 0)$ and $\mathcal{S} = (\mathcal{S}_{\alpha})_{\alpha \in \Lambda}$ be as above. Let $g_s : (X, 0) \to (\mathbb{K}, 0)$ be an isolated singularity family of K-analytic function-germs depending linearly on the parameter $s \in \mathbb{K}$. This means that each g_s has the form

$$
g_s(z) = g_0(z) + s\varphi(z) ,
$$

where $g_0: (X,0) \to (\mathbb{K},0)$ and $\varphi: (X,0) \to (\mathbb{K},0)$ are K-analytic isolated singularity function-germs.

Define the map-germ

$$
\mathcal{F}: (X,0) \rightarrow (\mathbb{K}^2,0)
$$

$$
z \mapsto (g_0(z),\varphi(z)) ,
$$

and for each $s \in \mathbb{K}$ define the line \mathcal{H}_s in \mathbb{K}^2 given by

$$
\mathcal{H}_s := \{ (y_0, y_1) \in \mathbb{K}^2; \ y_0 + sy_1 = 0 \}.
$$

Notice that

$$
(\mathcal{F})^{-1}(\mathcal{H}_s)=V(g_s).
$$

Following [\[5\]](#page-7-1) (Definition 2.1), we say that the family (g_s) is Δ -regular (with respect to the stratification S) if the line \mathcal{H}_0 is not a limit of secant lines of the discriminant set $\Delta(\mathcal{F})$ of $\mathcal F$ at $0 \in \mathbb{K}^2$. This means that there exist a neighborhood U of 0 in \mathbb{K}^2 and a real number $\delta > 0$ such that

$$
\mathcal{H}_s \cap \Delta(\mathcal{F}) \cap U \subset \{0\}
$$

whenever $||s|| < \delta$.

Now we give:

Definition 2.1. Let (g_s) be an isolated singularity family as above, and let $\epsilon > 0$ be a small real number. We say that a real number $\delta > 0$ is a good parameter-radius for (g_s) in \mathbb{B}_{ϵ} if for every $s \in \mathbb{D}_{\delta}$ one has that:

- (i) $\Sigma(g_s) \cap \mathbb{B}_{\epsilon} = \{0\};$
- (ii) $V(g_s) \setminus V(\varphi)$ intersects the sphere \mathbb{S}_{ϵ} transversally in \mathbb{K}^n , in the stratified sense; (iii) $\mathcal{H}_s \cap \Delta(\mathcal{F}) \cap U \subset \{0\}.$

It follows from [\[5\]](#page-7-1) (Lemma 3.2 together with Proposition 4.4) that if (g_s) has no coalescing of singular points, then for every $\epsilon > 0$ small enough there exists a good parameter-radius $\delta > 0$ for (g_s) in \mathbb{B}_{ϵ} .

Then the next proposition easily follows from the proof of $([5]$ $([5]$, Theorem 1.1).

Proposition 2.2. Let (g_s) be an isolated singularity family as above. If $\epsilon > 0$ is small enough and if $\delta > 0$ is a good parameter-radius for (g_s) in \mathbb{B}_{ϵ} , then for each $s \in \mathbb{D}_{\delta}$ there exists a homeomorphism

$$
h_s: V(g_s) \cap \mathbb{B}_{\epsilon} \to V(g_0) \cap \mathbb{B}_{\epsilon}
$$

that fixes the origin.

3. Multi-parameters families of isolated singularities

Let $(X,0) \subset (\mathbb{K}^N,0)$ and S be as above. Let $f_{s_1,\dots,s_k}:(X,0) \to (\mathbb{K},0)$ be an isolated singularity family of K-analytic function-germs depending linearly on the parameters s_1, \ldots, s_k , with $k > 1$. This means that each f_{s_1,\ldots,s_k} has the form

$$
f_{s_1,...,s_k}(z) = f_0(z) + s_1 \varphi_1(z) + \cdots + s_k \varphi_k(z),
$$

where $f_0: (X,0) \to (\mathbb{K},0)$ and $\varphi_i: (X,0) \to (\mathbb{K},0)$ for $i=1,\ldots,k$ are K-analytic isolated singularity function-germs.

If $\mathbb{K} = \mathbb{R}$ set $\xi := 1$, and if $\mathbb{K} = \mathbb{C}$ set $\xi := 2$. Then let $\mathbb{S}^{\xi k-1}$ denote the unitary sphere around 0 in \mathbb{K}^k .

Now, for each point $r = (r_1, \ldots, r_k) \in \mathbb{S}^{\xi k-1}$ fixed, consider the one-parameter isolated singularity family (g_s^r) given by

$$
g_s^r(z) := f_{sr_1,\ldots, sr_k}(z) .
$$

As before, we define the map-germ

$$
\mathcal{F}^r: (X,0) \rightarrow (K^2,0)
$$

$$
z \mapsto (f_0(z), r_1\varphi_1(z) + \cdots + r_k\varphi_k(z)) ,
$$

so that

$$
(\mathcal{F}^r)^{-1}(\mathcal{H}_s) = V(g_s^r).
$$

We say that the family $(f_{s_1,...,s_k})$ is Δ -regular if for each $r \in \mathbb{S}^{\xi k-1}$ the corresponding one-parameter family (g_s^r) is Δ -regular. This means that for each $r \in \mathbb{S}^{\xi k-1}$ there exist a neighborhood U_r of 0 in \mathbb{K}^2 and a real number $\delta_r > 0$ such that

$$
\mathcal{H}_s \cap \Delta(\mathcal{F}^r) \cap U_r \subset \{0\}
$$

whenever $||s|| < \delta_r$. We have:

Lemma 3.1. If the family (f_{s_1,\dots,s_k}) is Δ -regular, then there exist a neighborhood U of 0 in \mathbb{K}^2 and a real number $\delta > 0$ such that for every $r \in \mathbb{S}^{\xi k-1}$ and for every $s \in \mathbb{K}$ with $||s|| < \delta$ one has

$$
\mathcal{H}_s \cap \Delta(\mathcal{F}^r) \cap U \subset \{0\}.
$$

Proof. By continuity, for each $r \in \mathbb{S}^{\xi k-1}$ fixed, there exists some small open neighborhood N_r of r in $\mathbb{S}^{\xi k-1}$ such that, for every $w \in N_r$, one has that $\mathcal{H}_s \cap \Delta(\mathcal{F}^w) \cap U_r \subset \{0\}$ whenever $||s|| < \delta_r$. Then the result follows from the fact that $\mathbb{S}^{\xi k-1}$ is compact.

Now suppose that the family (f_{s_1,\ldots,s_k}) has no coalescing of singular points. Then it is clear that there exists $\lambda > 0$ sufficiently small such that for every $r \in \mathbb{S}^{\xi k-1}$ and for every $||s|| \leq \lambda$ one has that $\Sigma(g_s^r) \cap \mathbb{B}_{\epsilon} = \{0\}.$

Moreover, it is not difficult to see that $\lambda > 0$ can be chosen small enough so that it is also true that $V(g_s^r) \setminus V(r_1\varphi_1 + \cdots + r_k\varphi_k)$ intersects the sphere \mathbb{S}_{ϵ} transversally in \mathbb{K}^n , in the stratified sense, for every $s \in \mathbb{D}_{\lambda}$ and for every $r = (r_1, \ldots, r_k) \in \mathbb{S}^{\xi k-1}$.

So if we set $\delta' := \min\{\delta, \lambda\},\$ it follows that for every $\epsilon > 0$ small enough, δ' is a good parameter-radius for (g_s^r) in \mathbb{B}_{ϵ} , for every $r \in \mathbb{S}^{\xi k-1}$. Then it follows from Proposition [2.2](#page-4-0) that for each $s = (s_1, \ldots, s_k) \in \mathbb{K}^k$ with $||s|| < \delta'$ there exists a homeomorphism

$$
h_s: V(f_{s_1,\ldots,s_k}) \cap \mathbb{B}_{\epsilon} \to V(f_0) \cap \mathbb{B}_{\epsilon}
$$

that fixes the origin.

This proves Theorem [0.1,](#page-1-1) and Theorem [0.2](#page-1-0) follows easily.

Example 3.2. Recall the one-parameter isolated singularity family

$$
\mathbf{f}_s(x, y, z) = x^3 + y^4 + z^2 + sx^4 + s^2y^5
$$

on $X = \{x^2 - y^2 = 0\} \subset \mathbb{R}^3$, as in Example [1.3.](#page-3-1) Consider the two-parameters family $f_{s_1,s_2}(x,y,z) = x^3 + y^4 + z^2 + s_1x^4 + s_2y^5$

on X. Using Theorem [1.2,](#page-3-0) one can easily see that the family (f_{s_1,s_2}) has no coalescing of singular points. So it follows from Theorem [0.2](#page-1-0) that the family (f_s) has constant topological type.

Finally, Corollary [0.3](#page-1-2) follows from Theorem [0.2](#page-1-0) together with the following:

Lemma 3.3. In the case when $(X, 0) = (\mathbb{C}^n, 0)$, an isolated singularity family $(f_{s_1,...,s_k})$ has no coalescing of singular (critical) points if and only if it is μ -constant.

Proof. First notice that in the case when $k = 1$ the result is well-known (see [\[3\]](#page-7-3) for instance). So, in our case, where $k > 1$, it follows that (f_{s_1,\ldots,s_k}) has no coalescing of critical points if and only for each $r \in \mathbb{S}^{2k-1}$ there exists a real number $\lambda_r > 0$ such that $\mu(g_s^r)$ is constant for every $s \in \mathbb{C}$ with $||s|| < \lambda$. But since \mathbb{S}^{2k-1} is compact, that happens if and only if there is a real number $\lambda > 0$ such that $\mu(f_{s_1,\dots,s_k})$ is constant for every $s \in \mathbb{C}^k$ with $||s|| < \lambda$.

Example 3.4. Consider the two-parameters isolated singularity family (f_{s_1,s_2}) on \mathbb{C}^2 given by

$$
f_{s_1,s_2}(x,y) := x^3 + y^2 + x^2y + s_1(x^2 - x^2y) + s_2x^2.
$$

Note that it is not μ -constant. In fact, $\mu(f_0) = 2$, while $\mu(f_{0,s_2}) = 1$ for $s_2 \neq 0$. However, the one-parameter family (g_s) given by

$$
g_s(x, y) := f_{s_1, -s_1}(x, y) = x^3 + y^2 + x^2y + s_1(-x^2y)
$$

is clearly µ-constant.

Clearly, if the multi-parameters family (f_{s_1,\dots,s_k}) associated to a one-parameter family (f_s) on \mathbb{C}^n is μ -constant, then (f_s) is also μ -constant. Thus our criterium for constancy of the topological type of (f_s) is weaker then the criterium suggested by Le-Ramanujam's Conjecture when $n = 3$. So the natural question is:

Question 3.5. Is there an example of a μ -constant isolated singularity family (f_s) on \mathbb{C}^3 such that the corresponding multi-parameters family $(f_{s_1,...,s_k})$ is not μ -constant?

A negative answer to that question would imply that the μ -constant conjecture is true.

Now let $(X,0) \subset (\mathbb{C}^N,0)$ be the germ of a complex analytic isolated determinantal singularity (IDS), and let $\tilde{f}_s : (\mathbb{C}^N, 0) \to (\mathbb{C}, 0)$ be a family of complex function-germs depending holomorphically on the parameter $s = (s_1, \ldots, s_k) \in \mathbb{C}^k$. This means that the map $\tilde{F}: (\mathbb{C}^N \times \mathbb{C}^k, (0,0)) \to (\mathbb{C}, 0)$ defined by $F(x, s) := \tilde{f}_s(x)$ is holomorphic. The restriction of \tilde{F} to $(X, 0)$ defines a family of complex function-germs

$$
f_s:(X,0)\to(\mathbb{C},0).
$$

If (f_s) is an isolated singularity family, then for each $s \in (\mathbb{C}^k, 0)$ there is a well-defined Milnor number $\mu(f_s)$ in the sense of [\[1\]](#page-7-2). It is the number of points in the critical locus of a morsification f'_{s} $s'_{s}: X' \to \mathbb{C}$ of f_{s} , where X' is a smoothing of X near the origin.

In the particular case when X is an ICIS, it is well-know that $\mu(f_s) \leq \mu(f_0)$ whenever $\|s\|$ is small enough. Moreover, the family (f_s) has constant Milnor number if and only if it has no coalescing of singular points $([2]$ $([2]$ and $[3]$). This means that there exist positive real numbers $\epsilon > 0$ and $\delta > 0$ such that 0 is the only critical point of f_s in \mathbb{B}_{ϵ} , for every $s \in \mathbb{D}_{\delta}$. These two facts come from the fact that when X is an ICIS, the Milnor number $\mu(f_0, 0)$ of f_0 at 0 has an algebraic expression as the dimension of a quotient ring that is Cohen-Macaulay. Then the principle of conservation of number gives that

(1)
$$
\mu(f_0, 0) = \sum_{x \in \Sigma(f_s)} \mu(f_s, x),
$$

where $\mu(f_s, x)$ denotes the Milnor number of f_s at the singular point $x \in \Sigma(f_s)$.

When X is an arbitrary IDS there is no such algebraic expression for the Milnor number of $f_s: (X,0) \to (\mathbb{C},0)$, so one cannot apply the same argument. However, a simple topological argument gives the same equality. Precisely, we have:

Lemma 4.1. Let $f_s : (X,0) \to (\mathbb{C},0)$ be an isolated singularity family defined on an IDS. Then there exists $\delta > 0$ such that for every $s \in \mathbb{D}_{\delta}$ equality (1) above holds.

Proof. Let f'_s $S'_{s}: X' \to \mathbb{C}$ be a morsification (in family) of f_{s} . Since X is an IDS and since f_0 has an isolated singularity, we can choose $\epsilon > 0$ be small enough such that $X \cap \mathbb{B}_{\epsilon} \setminus \{0\}$ is smooth and the restriction of f_0 to $X \cap \mathbb{B}_{\epsilon} \setminus \{0\}$ is a submersion. We can also assume that f'_0 has exactly $\mu(f_0)$ -many (Morse) critical points in $X' \cap \mathbb{B}_{\epsilon}$. Then there exists $\delta > 0$ sufficiently small such that for every $s \in \mathbb{D}_{\delta}$ one has that f'_{s} has exactly $\mu(f_0)$ -many critical points in $X' \cap \mathbb{B}_{\epsilon}$. This follows from the fact that Morse points are stable under small perturbations. Thus equality (1) above follows. \Box

As an immediate consequence, we have:

Proposition 4.2. Let $f_s : (X,0) \to (\mathbb{C},0)$ be an isolated singularity family defined on an IDS. Then there exists $\delta > 0$ such that:

- (i) for every $s \in \mathbb{D}_{\delta}$ one has $\mu(f_s) \leq \mu(f_0)$.
- (ii) $\mu(f_s) = \mu(f_0)$ for every $s \in \mathbb{D}_{\delta}$ if and only if (f_s) has no coalescing of singular points.

On the one hand, Proposition [4.2](#page-6-0) gives a characterization for the μ -constancy of deformations of functions on an IDS, in the spirit of [\[3\]](#page-7-3) and [\[2\]](#page-7-4).

On the other hand, if $(X,0) \subset (\mathbb{C}^N,0)$ is an IDS and if $f_s : (X,0) \to (\mathbb{C},0)$ is an isolated singularity family depending linearly on the parameters $s_1, \ldots, s_k \in \mathbb{C}$, Proposition [4.2](#page-6-0) together with Theorem [0.2](#page-1-0) give Theorem [0.4.](#page-1-3)

Notice that Proposition [4.2](#page-6-0) and Proposition [1.1](#page-2-0) combined give the following Corollary, which is a good criterium to decide if the family (f_s) is μ -constant:

Corollary 4.3. Let $\tilde{f}_s : (\mathbb{C}^N, 0) \to (\mathbb{C}, 0)$ be an isolated singularity family depending holomorphically on $s \in \mathbb{C}^k$. Let $(X,0) \subset (\mathbb{C}^N,0)$ be an IDS and suppose that the family $f_s: (X,0) \to (\mathbb{C},0)$ given by restriction to $(X,0)$ is also an isolated singularity family. If (\tilde{f}_s) is μ -constant then (f_s) is μ -constant.

Example 4.4. Let $X \subset \mathbb{C}^4$ be the 2-dimensional IDS defined by the matrix

$$
\begin{pmatrix} w & y & x \ z & w & y \end{pmatrix}.
$$

Consider the μ -constant family $\tilde{f}_s : (\mathbb{C}^4,0) \to (\mathbb{C},0)$ given by

$$
\tilde{f}_s(x, y, z, w) := x^2 + y^2 + z^2 + w^2 + sx^3,
$$

and let $f_s: (X,0) \to (\mathbb{C},0)$ be its restriction to X. For each $s \in \mathbb{C}$ fixed, the singular points of f_s are those points in X at which all the 3×3 -minors of the following matrix vanish:

$$
\begin{pmatrix} \alpha(x) & 2y & 2z & 2w \\ 0 & z & y & -2w \\ z & -w & x & -y \\ w & -2y & 0 & x \end{pmatrix},
$$

where $\alpha(x) := 2x + 3sx^2$. Then some calculation shows that the singular locus of f_s is the finite set

$$
\Sigma(f_s) = \{ \alpha(x) = y = z = w = 0 \} = \left\{ (0, 0, 0, 0), \left(-\frac{2}{3s}, 0, 0, 0 \right) \right\}.
$$

Hence the family (f_s) has no coalescing of singular points. Thus if follows from Propo-sition [4.2](#page-6-0) that f_s is a μ -constant deformation of the function-germ $f_0 : (X,0) \to (\mathbb{C},0)$.

On the other hand, consider the family $g_s: (X,0) \to (\mathbb{C},0)$ given by restriction to X of the family $\tilde{g}_s : (\mathbb{C}^4,0) \to (\mathbb{C},0)$ defined by

$$
\tilde{g}_s(x, y, z, w) := x^3 + y^2 + z^2 + w^2 + sx^2,
$$

which has non-constant Milnor number. Setting $\beta(x) := 3x^2 + 2sx$ one has that

$$
\Sigma(g_s) = \{ \beta(x) = y = z = w = 0 \} = \left\{ (0, 0, 0, 0), \left(-\frac{2s}{3}, 0, 0, 0 \right) \right\}.
$$

Then the family (g_s) has coalescing of singular points and thus it is not a μ -constant deformation.

REFERENCES

- [1] J. J. Nu˜no-Ballesteros, B. Orefice-Okamoto and J. N. Tomazella, The vanishing Euler characteristic of an isolated determinantal singularity. Israel J. of Math. 197 (2013), 475-495. [2,](#page-1-4) [7](#page-6-1)
- [2] S. Carvalho, B. Orefice-Okamoto and J. N. Tomazella, µ-constant deformations of functions on an ICIS. J. of Singularities, Vol. 19, 163-176 (2019). [7](#page-6-1)
- [3] G. M. Greuel, Constant Milnor number implies constant multiplicity for quasihomogeneous singularities. Manuscripta Math. 56 (1986), 159-166. [6,](#page-5-0) [7](#page-6-1)
- [4] D. T. Lê and C. P. Ramanujam, The invariance of Milnor's number implies the invariance of the topological type. Am. J. Math. 98, 67-78 (1976). [1,](#page-0-0) [2](#page-1-4)
- [5] A. Menegon and J. P. de Jesus-Almeida, Topological triviality in families of functions on subanalytic sets and complex analytic spaces. To appear in Int. Math. Research Notices (https://doi.org/10.1093/imrn/rnac008). [1,](#page-0-0) [4,](#page-3-2) [5](#page-4-1)
- [6] A. Parusiński, *Topological Triviality of* μ *-constant deformations of type* $f(x) + tg(x)$ *.* Bull. London Math. Soc. 31 (1999), no. 6, 686-692. [1](#page-0-0)
- [7] P. Samuel, Algébricité de certains points singuliers algebroides. J. Math. Pures Appl., 35 (1956), 1-6. [2](#page-1-4)
- [8] J. L. Verdier, Stratifications de Whitney et théorème de Bertini-Sard. Inv. Math. 36, 295-312 (1976). [3](#page-2-1)

AURÉLIO MENEGON: DEPARTAMENTO DE MATEMÁTICA - UNIVERSIDADE FEDERAL DA PARAÍBA - Brazil.

Email address: aurelio.menegon@academico.ufpb.br

Miriam da Silva Pereira: Departamento de Matematica - Universidade Federal da ´ PARAÍBA - BRAZIL.

Email address: msp@academico.ufpb.br