

BEHAVIOR OF KREISS BOUNDED C_0 -SEMIGROUPS ON A HILBERT SPACE

L. ARNOLD

ABSTRACT. Let $0 < \alpha \leq 1$. We prove that a α -Kreiss bounded C_0 semigroup $(T_t)_{t \geq 0}$ on a Hilbert space has asymptotics $\|T_t\| = \mathcal{O}(t^\alpha / \sqrt{\log(t)})$. Then, we give an application to perturbed wave equation.

Let $-A$ be the generator of a C_0 -semigroup $(T_t)_{t \geq 0}$ on a complex Banach space X . For $\alpha > 0$, we say that $(T_t)_{t \geq 0}$ is α -Kreiss bounded (or simply Kreiss bounded when $\alpha = 1$) if $\sigma(A) \subset \overline{\mathbb{C}_+} = \{\lambda \in \mathbb{C} : \operatorname{Re}(\lambda) \geq 0\}$ and

$$(1) \quad \|R(\lambda, A)\| \leq \frac{C}{(-\operatorname{Re}(\lambda))^\alpha}, \quad \forall \operatorname{Re}(\lambda) < 0.$$

According to [5, (2.6)] (see also [8, Theorem 1.1.(1)]), if $(T_t)_{t \geq 0}$ is an α -Kreiss bounded C_0 -semigroup on a Hilbert space for $\alpha > 0$, then $\|T_t\| = O(t^\alpha)$. Let us fix $\gamma \in (0, 1)$. In [3], the authors give an example (see [3, Example 4.4.]) of Kreiss bounded C_0 -semigroup for which there exists a constant $C > 0$ such that $\|T_t\| \geq Ct^\gamma$ for $t \geq 1$. A natural question (listed as an open question in [7, Appendix B]) is whether the estimate $\|T_t\| = O(t^\alpha)$, for α -Kreiss bounded C_0 -semigroup $(T_t)_{t \geq 0}$ on a Hilbert space, is sharp. In the following we answer this question for $\alpha \in (0, 1]$.

Theorem 1. *Let $0 < \alpha \leq 1$ and let $-A$ be the generator of α -Kreiss bounded C_0 -semigroup $(T_t)_{t \geq 0}$ on a Hilbert space. Then*

$$(2) \quad \|T_t\| = \mathcal{O}_{t \rightarrow \infty} \left(t^\alpha / \sqrt{\log(t)} \right).$$

The proof of the Theorem 1 is based on techniques given in [2]. We first state and prove two lemmas. It is known (see [4] or [9]) that a C_0 -semigroup $(T_t)_{t \geq 0}$ on a Hilbert space, with negative generator A , is bounded if and only if

$$\|R(-r + i \cdot, A)x\|_{L^2(\mathbb{R}, H)}^2 \leq \frac{C}{r} \|x\|^2, \quad r > 0$$

and

$$\|R(-r + i \cdot, A^*)x^*\|_{L^2(\mathbb{R}, H)}^2 \leq \frac{C}{r} \|x^*\|^2, \quad r > 0.$$

The following lemma gives an analogous necessary condition for a C_0 -semigroup to be α -Kreiss bounded, for $\alpha > 0$.

Lemma 2. *Let $\alpha > 0$ and $(T_t)_{t \geq 0}$ be a C_0 -semigroup on a Hilbert space whose negative generator A . Assume $(T_t)_{t \geq 0}$ is α -Kreiss bounded then*

$$(3) \quad \|R(-r + i \cdot, A)x\|_{L^2(\mathbb{R}, H)}^2 \leq \frac{C(1+r^\alpha)^2}{r^{2\alpha}} \|x\|^2, \quad r > 0$$

2020 *Mathematics Subject Classification.* Primary 47D06 ; Secondary 34D05, 35B45.
Key words and phrases. C_0 -semigroup, Kreiss condition, perturbed wave equation.

and

$$(4) \quad \|R(-r + i\cdot, A^*)x^*\|_{L^2(\mathbb{R}, H)}^2 \leq \frac{C(1 + r^\alpha)^2}{r^{2\alpha}} \|x^*\|^2, \quad r > 0.$$

Proof. Let $x \in H$. First, we by [6, formula (7.1) section 1.7], we have

$$R(-r + i\beta, A)x = - \int_0^\infty e^{i\beta s} e^{-rs} T_s x ds, \quad r > 0, \beta \in \mathbb{R}.$$

This means that for $r > 0$,

$$(5) \quad R(-r + i\beta, A)x = -\mathcal{F}^{-1}(s \mapsto e^{-rs} T_s x)(\beta),$$

where $\mathcal{F} : L^2(\mathbb{R}, H) \mapsto L^2(\mathbb{R}, H)$ is the Fourier-Plancherel operator. By Fourier-Plancherel Theorem

$$(6) \quad \|\mathcal{F}^{-1}(s \mapsto e^{-rs} T_s x)(\cdot)\|_{L^2(\mathbb{R}, H)} = \frac{1}{\sqrt{2\pi}} \|e^{-r\cdot} T_{(\cdot)} x\|_{L^2(\mathbb{R}_+, H)}.$$

The Gearhart-Prüss Theorem (see [1, Theorem 5.2.1.]) says that for each $r > 0$, $\sup_{t>0} \|e^{-rt} T_t\| < \infty$. This implies that there exists $K > 0$ such that

$$\sup_{a>1} \|e^{-a\cdot} T_{(\cdot)} x\|_{L^2(\mathbb{R}_+, H)} \leq K \|x\|$$

and then

$$(7) \quad \sup_{a>1} \|R(-a + i\beta, A)x\|_{L^2(\mathbb{R}_+, H)} \leq \frac{K}{\sqrt{2\pi}} \|x\|.$$

Now, let $r > 0$ and $a = r + 1 > 1$. By the resolvent identity, we have, for $\beta \in \mathbb{R}$,

$$\begin{aligned} R(-r + i\beta, A)x &= (I + (r - a)R(-r + i\beta, A))R(-a + i\beta, A)x \\ &= (I - R(-r + i\beta, A))R(-a + i\beta, A)x. \end{aligned}$$

Using (1) and (7):

$$\begin{aligned} \|R(-r + i\cdot, A)x\|_{L^2(\mathbb{R}, H)}^2 &\leq (1 + \sup_{\beta \in \mathbb{R}} \|R(-r + i\beta, A)\|)^2 \sup_{a' < -1} \|R(-a' + i\cdot, A)x\|_{L^2(\mathbb{R}_+, H)}^2 \\ &\leq 2\pi K^2 \left(1 + \frac{C}{r^\alpha}\right)^2 \|x\|^2 \\ &\leq C' \frac{(1 + r^\alpha)^2}{r^{2\alpha}} \|x\|^2. \end{aligned}$$

In the same way we obtain (4). □

The next lemma says that if $(T_t)_{t \geq 0}$ and $(T_t^*)_{t \geq 0}$ satisfy a kind of Cesàro condition then $\|T_t\|$ satisfy (2).

Lemma 3. *Let $\alpha \in (0, 1)$ and $(T_t)_{t \geq 0}$ be a C_0 -semigroup on a Banach space X . Assume that there exists $C > 0$ such that for each $t > 1$,*

$$\int_0^t \|T_s x\|^2 ds \leq C t^{2\alpha} \|x\|^2, \quad x \in X,$$

and

$$\int_0^t \|T_s^* x^*\|^2 ds \leq C t^{2\alpha} \|x^*\|^2, \quad x^* \in X^*.$$

Then, $\|T_t\| = \mathcal{O}(t^\alpha / \sqrt{\log(t)})$.

Proof. Let $t > 2$ and $1 < P < Q < t$. We have

$$\begin{aligned} (Q - P)^2 |\langle T_t x, x^* \rangle|^2 &= \left| \int_P^Q \langle T_{t-s} x, T_s^* x^* \rangle ds \right|^2 \\ &\leq \left(\int_P^Q \|T_{t-s} x\|^2 ds \right) \left(\int_P^Q \|T_s^* x^*\|^2 ds \right) \\ &\leq \left(\int_P^Q \|T_{t-s} x\|^2 ds \right) \left(\int_0^Q \|T_s^* x^*\|^2 ds \right) \\ &\leq CQ^{2\alpha} \|x^*\|^2 \left(\int_{t-Q}^{t-P} \|T_s x\|^2 ds \right) \end{aligned}$$

Finally taking the supremum over $\{\|x^*\| = 1\}$ one obtains, since $\alpha \in (0, 1]$,

$$(8) \quad \begin{aligned} \frac{(Q - P)^2}{Q^2} \|T_t x\|^2 &\leq CQ^{2(\alpha-1)} \int_{t-Q}^{t-P} \|T_s x\|^2 ds \\ &\leq C \int_{t-Q}^{t-P} \|T_s x\|^2 ds \end{aligned}$$

Now we set $L := \lfloor \log(t)/\log(2) \rfloor$. For $0 \leq l \leq L - 1$, $Q = 2^{l+1}$ and $P = 2^l$ we have

$$\|T_t x\|^2 \leq 4C \int_{t-2^{l+1}}^{t-2^l} \|T_s x\|^2 ds$$

Therefore

$$L \|T_t x\|^2 = \sum_{l=0}^{L-1} \|T_t x\|^2 \leq \sum_{l=0}^{L-1} 4C \int_{t-2^{l+1}}^{t-2^l} \|T_s x\|^2 ds \leq 4C \int_0^t \|T_s x\|^2 ds \leq 4C^2 t^{2\alpha}$$

Hence for $t > 2$,

$$\|T_t x\| \leq \frac{2Ct^\alpha}{\sqrt{L}} \|x\| \leq \frac{C't^\alpha}{\sqrt{\log(t)}} \|x\|.$$

□

We are now ready to prove Theorem 1.

Proof of Theorem 1. Let $x \in H$ and $r > 0$. By (5), (6) and Lemma 2,

$$\|e^{-r \cdot} T_{(\cdot)} x\|_{L^2(\mathbb{R}_+, H)}^2 = \|R(-r + i \cdot, A)x\|_{L^2(\mathbb{R}, H)}^2 \leq \frac{C(1 + r^\alpha)^2}{r^{2\alpha}} \|x\|^2.$$

Furthermore

$$\|e^{-r \cdot} T_{(\cdot)} x\|_{L^2(\mathbb{R}_+, H)}^2 = \int_{\mathbb{R}} e^{-2rs} \|T_s x\|^2 ds \geq e^{-2rt} \int_0^t \|T_s x\|^2 ds.$$

Taking $t = \frac{1}{r}$ we obtain

$$\int_0^t \|T_s x\|^2 ds \leq C e^{2t^{2\alpha}} \left(1 + \frac{1}{t^\alpha}\right)^2 \|x\|^2.$$

Hence, for $t > 1$,

$$\int_0^t \|T_s x\|^2 ds \leq C' t^{2\alpha} \|x\|^2,$$

where $C' = 4e^2C$. Similarly, since $(T_t^*)_{t \geq 0}$ is Kreiss bounded C_0 -semigroup on a Hilbert space, for $t > 1$ and $x^* \in H$, we have

$$\int_0^t \|T_s^* x^*\|^2 ds \leq C' t^{2\alpha} \|x^*\|^2.$$

Finally, Lemma 3 allows us to conclude that $\|T_t\| = \mathcal{O}(t^\alpha/\sqrt{\log(t)})$. \square

Remark 4. (1) When $\alpha > 1$, (8) with $Q = 2$ and $P = 1$ gives

$$\|T_t x\|^2 \leq C 2^\alpha \int_{t-2}^{t-1} \|T_s x\|^2 ds \leq C' \int_0^t \|T_s x\|^2 ds \leq C'' t^{2\alpha} \|x\|^2.$$

Then $\|T_t\| = \mathcal{O}(t^\alpha)$, which gives another proof of [8, Theorem 1.1.(1) with $g(s) = s^\alpha$ for $\alpha > 1$].

(2) Let us notice that if $(T_t)_{t \geq 0}$ satisfies the α -Kreiss bounded condition only for $\operatorname{Re}(\lambda) \in (0, 1)$ that is

$$\|R(\lambda, A)\| \leq \frac{C}{-\operatorname{Re}(\lambda)^\alpha}, \quad \forall \operatorname{Re}(\lambda) \in (-1, 0),$$

then the conclusion $\|T_t\| = \mathcal{O}(t^\alpha/\sqrt{\log(t)})$ remains true.

Let give an application of the Theorem 1. Let $W^{2,s}(\mathbb{T}^2)$ be the second order Sobolev space equipped with the standard norm. Let A be the operator with domain $D(A) = W^{2,2}(\mathbb{T}^2) \times W^{2,1}(\mathbb{T}^2)$ defined by

$$A = \begin{pmatrix} 0 & -1 \\ -\Delta - M \frac{\partial}{\partial x} & 0 \end{pmatrix},$$

where Δ is the Laplacian with $D(\Delta) = W^{2,2}(\mathbb{T}^2)$ and $M : L^2(\mathbb{T}^2) \rightarrow L^2(\mathbb{T}^2)$ is the multiplication operator given by $M(h)(x, y) = e^{iy}h(x, y)$. Then, it is known (see [8, section 4] for details) that $-A$ generates a C_0 -group $(T_t)_{t \in \mathbb{R}}$ on the Hilbert space $H = W^{1,2}(\mathbb{T}^2) \times L^2(\mathbb{T}^2)$, satisfying $\|T_t\| = \mathcal{O}_{t \rightarrow \pm\infty}(|t|e^{|t|/2})$. With Theorem 1, we are able to improve this estimate:

Proposition 5. Let $(T_t)_{t \in \mathbb{R}}$ be the C_0 -group on H generated by $-A$. Then

$$(9) \quad \|T_t\| = \mathcal{O}_{t \rightarrow \pm\infty} \left(\frac{|t|}{\sqrt{\log(|t|)}} e^{|t|/2} \right).$$

Proof. According to [8, Lemma 4.4.], for $\lambda \in \mathbb{C}_+$, with $\operatorname{Re}(\lambda) \in (0, 1)$,

$$\left\| R\left(\lambda, A + \frac{1}{2}\right) \right\| \leq \frac{C}{-\operatorname{Re}(\lambda)} \quad \text{and} \quad \left\| R\left(\lambda, -A + \frac{1}{2}\right) \right\| \leq \frac{C}{-\operatorname{Re}(\lambda)}.$$

Then, according to Remark 4 (2), the semigroup $(e^{-1/2t}T_t)_{t \geq 0}$ (resp. $(e^{-1/2t}T_{-t})_{t \geq 0}$) which is generated by $-(A + \frac{1}{2})$ (resp. $A + \frac{1}{2}$), satisfies $\|e^{-1/2t}T_t\| = \mathcal{O}(t/\sqrt{\log(t)})$ (resp. $\|e^{-1/2t}T_{-t}\| = \mathcal{O}(t/\sqrt{\log(t)})$). This yields the estimate (9). \square

Acknowledgements. Author was supported by the ERC grant *Rigidity of groups and higher index theory* under the European Union's Horizon 2020 research and innovation program (grant agreement no. 677120-INDEX).

REFERENCES

- [1] A. Arendt, C. Batty, M. Hierber, and F. Neubrander. *Vector-valued Laplace Transforms and Cauchy Problems : Second Edition*. Monograph in Mathematics 96, Springer Basel AG, 2011.
- [2] G Cohen, C Cuny, T Eisner, and M Lin. Resolvent conditions and growth of powers of operators. *J. Math. Anal. Appl.*, 487, 2020.
- [3] T Eisner and H Zwart. Continuous-time kreiss resolvent condition on infinite-dimensional spaces. *Math. Comput.*, 75:1971–1985, 2006.
- [4] A.M. Gomilko. Conditions on the generator of a uniformly bounded C_0 -semigroup. *Funct. anal. Appl.* 33, pages 294–296, 1999.
- [5] B Helffer and J Sjostrand. From resolvent bounds to semigroup bounds. <https://arxiv.org/pdf/1001.4171v1.pdf>, 2010.
- [6] A. Pazy. *Semigroups of Linear Operators and Applications to PDEs*. Springer, 1992.
- [7] J Rozendaal. Operator-valued (l^p, l^q) fourier multipliers and stability theory for evolution equations. <https://arxiv.org/pdf/2204.02223.pdf>, 2022.
- [8] J Rozendaal and M Veraar. Sharp growth rates for semigroups using resolvent bounds. *Journal of Evolution Equations*, 18:1721–1744, 2018.
- [9] D.-H. Shi and Feng D.X. Characteristic conditions of the generation of C_0 -semigroups in a hilbert space. *J. Math. Anal. Appl.* 247 : 356-376, 2000.

(L. Arnold) JANA I JEDRZEJA ŚNIADECKICH 8, 00-656 WARSZAWA, POLAND
Email address: larnold@impan.pl