

Prescribed graphon symmetries and flavors of rigidity

Alexandru Chirvasitu

Abstract

We prove that an arbitrary compact metrizable group can be realized as the automorphism group of a graphing; this is a continuous analogue to Frucht’s theorem recovering arbitrary finite groups as automorphism groups of finite graphs.

The paper also contains a number of results the persistence of transitivity of a compact-group action upon passing to a limit of graphons. Call a compact group \mathbb{G} graphon-rigid if, whenever it acts transitively on each member Γ_n of a convergent sequence of graphons, it also acts transitively on the limit $\lim_n \Gamma$. We show that for a compact Lie group \mathbb{G} graphon rigidity is equivalent to the identity component \mathbb{G}_0 being semisimple; as a partial converse to a result of Lovász and Szegedy, this is also equivalent to weak randomness: the property that the group have only finitely many irreducible representations in each dimension. Similarly, call a compact group \mathbb{G} image-rigid if for every compact Lie group \mathbb{H} the images of morphisms $\mathbb{G} \rightarrow \mathbb{H}$ form a closed set (of closed subgroups, in the natural topology). We prove that graphon rigidity implies image rigidity for compact groups that are either connected or profinite, and the two conditions are equivalent (and also equivalent to being torsion) for profinite abelian groups.

Key words: graphon; graphing; compact group; metrizable; Borel space; standard; kernel; Lie group

MSC 2020: 22C05; 28A33; 60B05; 22F50; 20B27; 54E35; 54E45; 54E50; 54E70

Contents

1 Preliminaries	3
2 Prescribed graphon symmetries	5
2.1 Revisiting a definition	6
3 Rigidity	8
3.1 Graphon-rigid Lie groups	8
3.2 Stable compact-group images	18
4 Graphing automorphisms	23
References	27

Introduction

Graphons are continuous and/or probabilistic analogues of graphs, intended to capture the latter’s long-term, limiting behavior. One of several slight variations (e.g. [26, §2.1]; we review standard spaces briefly in Section 1):

Definition 0.1 A *graphon* $\Gamma = (X, W, \mu)$ consists of a standard probability space (X, μ) and a measurable symmetric function $W : X \times X \rightarrow [0, 1]$. \blacklozenge

Aspects of the rich, multi-faceted ensuing theory are discussed in [24, 25, 39, 26] (and their references) as well as the excellent [23]. The topic of interest here is that of symmetries (or automorphisms) of a graphon. The automorphism group $\text{Aut}(\Gamma)$ is introduced in passing in [23, §13.5], and discussed more extensively in [26]. Most importantly for the discussion below, it is always compact [26, Theorem 10].

One direction the discussion below takes in the general spirit of an “inverse” symmetry problem: given a group (enjoying various properties, as appropriate), realize it as the automorphism group of a structure (here, graphons). The pattern recurs frequently:

- Every finite group is the automorphism group of some finite graph by a classical result of Frucht’s [10].
- That result can be improved by imposing various constraints on the graphs realizing the given group (connectivity, chromatic number, etc.) [33, Theorem 1.2].
- And in fact *arbitrary* (possibly infinite) groups can be realized as graph automorphism groups [34, Introduction, Theorem].
- Adjacent to graphs, (convex) *polytopes* also realize arbitrary finite groups: see [7, Theorem 1.1], [36, Theorems 1 and 2], [35, Theorem 3.1].
- There are variations on this theme: the group might have a distinguished involution which is required to act as the central symmetry in a *symmetric* convex polytope; again, arbitrary data of this nature can be realized [4, Theorem 2.1].
- And then there is of course the celebrated *inverse Galois problem*, of realizing finite groups as Galois groups of finite field extensions of \mathbb{Q} (still unsolved in full generality): [37].

The graphon analogue of all of this is Theorem 2.5 below:

Theorem *Every compact metrizable group is the automorphism group of some graphon.* \blacksquare

Or in other words, since [26, Theorem 10] says that the automorphism group of a graphon is compact (and it is obviously metrizable), that result has as strong a converse as one could possibly hope for.

Another point of interest is the limiting behavior of graphon automorphism groups. The issue comes up in [26], where [26, Examples 34 and 35] show that if

$$\Gamma_n \xrightarrow[n]{} \Gamma$$

is a convergent sequence of graphons with Γ_n all admitting transitive actions by a group \mathbb{G} , the limit Γ , though again transitive under $\text{Aut}(\Gamma)$, need not admit a transitive action by the *same* group \mathbb{G} .

By contrast, [26, Theorem 37] identifies (following [39, §1.8]) a class of groups for which transitivity *does* transport over from the individual Γ_n to the limit $\Gamma = \lim_n \Gamma_n$: the *weakly random* compact groups, i.e. those having only finitely many irreducible representations in every dimension.

To simplify the awkward language, we refer to compact groups \mathbb{G} for which transitive actions transfer over from the members Γ_n of a convergent sequence to the limit $\Gamma = \lim_n \Gamma_n$ as *graphon-rigid* (see Definition 3.1). Weak randomness and graphon rigidity are both (as the latter’s name

suggests) rigidity properties of sorts. In light of the above the question naturally arises of whether (and to what extent) weak randomness is *necessary* for graphon rigidity. The two turn out to coincide for Lie groups, and to in fact be equivalent to yet another (much more?) familiar property in that case. To paraphrase the more verbose [Theorem 3.12](#):

Theorem *A compact Lie group is graphon-rigid precisely when it is weakly random or, equivalently, when its identity connected component is semisimple. ■*

Being Lie is essential: [Example 3.14](#) shows that it is perfectly possible for infinite *abelian* compact groups to be graphon-rigid. Being abelian, these are maximally far from weak randomness: *all* of their irreducible representations are 1-dimensional.

The various (counter)examples suggest a related notion of rigidity ([Definition 3.21](#)): call a compact group \mathbb{G} *image-rigid* if for every compact Lie group \mathbb{H} , the images of the morphisms $\mathbb{G} \rightarrow \mathbb{H}$ form a closed set in the natural compact Hausdorff topology ([Remark 3.7](#)) on the space $\mathcal{CLG}(\mathbb{H})$ of closed subgroups of \mathbb{H} .

This property, in some cases, appears to be more pertinent to graphon rigidity than weak randomness. Aggregating [Theorem 3.28](#) and [Proposition 3.29](#):

Theorem *Let \mathbb{G} be a compact group.*

- *If \mathbb{G} is graphon-rigid then it is image-rigid provided it is either connected or profinite.*
- *If furthermore \mathbb{G} is abelian and profinite, then graphon and image rigidity are equivalent. ■*

In the short [Section 4](#) we give a possible definition of the automorphism group of a *graphing*, in partial answer to a question asked in [[22](#), §5]. That group will not, in general, be compact, but the definition does make it invariant under taking *full subgraphings* in the sense of [[22](#), §2].

Acknowledgements

This work is partially supported by NSF grant DMS-2001128.

I am grateful for helpful comments and pointers to the literature from L. Lovasz.

1 Preliminaries

We assume some basic background on measure theory, as covered in, say, [[32](#), [2](#)]; the references will be more precise when needed. As is customary in the literature (e.g. [[2](#), §2.2]), for a measure μ we abbreviate *almost everywhere* or μ -almost everywhere to *a.e.* or μ -*a.e.*. Recall also ([[2](#), Definition 2.1.3]) that a function

$$\varphi : (X, \mathcal{B}_X) \rightarrow (Y, \mathcal{B}_Y)$$

between spaces equipped with σ -algebras [[2](#), Definition 1.2.2] is *measurable* when

$$\varphi^{-1}(S) \in \mathcal{B}_X, \quad \forall S \in \mathcal{B}_Y.$$

We work virtually exclusively with *probability* measures, and all probability spaces (X, \mathcal{B}, μ) are *complete* ([[32](#), §1.1] or [[2](#), Definition 1.5.10]): subsets of measurable sets $N \subset X$ with $\mu(N) = 0$ are again measurable. The following objects feature prominently in the sequel (see e.g. [[2](#), Definitions 9.2.1 and 9.4.6]).

Definition 1.1 (a) Two probability spaces $(X, \mathcal{B}_X, \mu_X)$ and $(Y, \mathcal{B}_Y, \mu_Y)$ are *isomorphic mod 0* if

- there are sets of $M \subset X$ and $N \subset Y$ of measure zero;
- and a measurable, measure-preserving bijection

$$X \setminus M \rightarrow Y \setminus N$$

- whose inverse is also measurable.

(b) A *standard* probability space is a (complete) probability space that is isomorphic mod 0 to the unit interval with its usual Lebesgue measure appropriately scaled, together, perhaps, with a sequence of point masses. \blacklozenge

Remark 1.2 Standard probability spaces are also (essentially)

- the *Lebesgue spaces* of [32, §1.4, Definition 4.5];
- the *Lebesgue-Rohlin spaces* of [2, Definition 9.4.6 and Theorem 9.4.7];
- the *L.R. spaces* of [11, Definition 7] (see [11, Proposition 6 and Remarks following it]).

The terminology also matches [23, §A.3.1]. \blacklozenge

Remark 1.3 The concept of mod-0 morphism

$$(X, \mathcal{B}_X, \mu_X) \rightarrow (Y, \mathcal{B}_Y, \mu_Y)$$

is potentially ambiguous, with at least three possible interpretations. It might mean

- a measurable, measure-preserving map defined on a full-measure set $X \setminus M$ as in Definition 1.1, with two such maps identified if they agree a.e.;
- a measurable, measure-preserving map defined *everywhere* on X , again identifying two such maps if they agree everywhere;
- or, finally, a measure-preserving morphism

$$\mathcal{B}_Y \rightarrow \mathcal{B}_X$$

of σ -algebras [32, §1.4.C] (note the direction reversal: such a map is to be thought of as the preimage function φ^{-1} induced by φ as in (a) or (b)). Or, in the language of [32, §1.4.C], a morphism of *measure algebras*.

For standard spaces the ambiguity is only apparent:

- On the one hand the notions listed above are clearly progressively “weaker” as listed, in the sense that an object as in (a) produces one as in (b), etc. Being standard plays no role here.
- On the other hand, for standard spaces a σ -algebra morphism as in (c) arises from (b) by [32, §1.4, Theorem 4.7].
- And the passage from (b) back to (a) is illustrated in the proof of [11, Proposition 6].

As all of our spaces are standard unless noted otherwise, we can make free use of these mod-0 concepts in whatever form is most convenient in any given situation. The discussion also applies to *isomorphisms*, *automorphisms*, etc. For instance, for a standard probability space (X, \mathcal{B}, μ) we can speak of its *automorphism group* $\text{Aut}(X, \mathcal{B}, \mu)$: it consists of mod-0 classes of automorphisms in the sense of the preceding discussion. \blacklozenge

2 Prescribed graphon symmetries

First, recall from [26, §2.3] (also [23, §13.3]):

Definition 2.1 The *neighborhood distance* r_W attached to a graphon (X, W, μ) is

$$r_W(x, y) := \|W(x, -) - W(y, -)\|_1 = \int_X |W(x, z) - W(y, z)| \, d\mu(z).$$

A graphon (J, W, μ) is *pure* if it is complete under the neighborhood distance and its probability measure μ has full support. \blacklozenge

We work with metric probability spaces (X, d, μ) as graphons, as explained in [23, Example 13.17]. These turn out to automatically be pure, as follows from the following simple remark.

Proposition 2.2 A graphon $\Gamma = (X, W, \mu)$ with

- X compact Hausdorff second-countable;
- W continuous and

$$X \ni x \mapsto W(x, -) \in C(X) := \text{continuous functions on } X \quad (2-1)$$

one-to-one.

- and μ fully supported.

is pure.

Proof For compact Hausdorff spaces being second-countable is equivalent to metrizable [30, §34, Exercise 3], and the continuity of W then entails its *uniform* continuity [30, Theorem 27.6]. It follows that (2-1) is continuous when equipping the right-hand side with the uniform topology, so the identity map is continuous from the original (compact Hausdorff) topology on X and the r_W -metric topology.

Note also that r_W is indeed a metric (rather than a pseudometric) because we are assuming that (2-1) is injective. It follows that

$$\text{id} : X \rightarrow (X, r_W)$$

is a bijective and continuous map from a compact space to a Hausdorff space, and must thus be a homeomorphism. This then implies that the metric space (X, r_W) is complete (being compact [30, Theorem 45.1]), and purity follows from the additional assumption that μ is fully supported. \blacksquare

In particular, taking for the kernel W the distance function d of a compact metric space (X, d) as in [23, Example 13.17], we obtain

Corollary 2.3 For a compact metric space (X, d) with fully-supported probability measure μ the resulting graphon (X, d, μ) is pure. \blacksquare

The following observation will be implicit in some of the discussion below.

Lemma 2.4 Under the hypotheses of Proposition 2.2 the automorphism group of Γ acts continuously on X in the latter's original compact Hausdorff topology.

Proof Indeed, it acts continuously on the metric space (X, r_W) , and we saw in the course of the proof that that metric space is homeomorphic to the original X . ■

The following result is a graphon analogue of fact that arbitrary groups can be realized as graph automorphism groups [34, p.64, Theorem].

Theorem 2.5 *A compact group is the automorphism group of a graphon if and only if it is metrizable.*

Proof On the one hand, [26, Theorem 10] shows that the automorphism group of a graphon is compact, while [26, discussion following Definition 7] explains how to metrize it.

Conversely, according to [27, Theorem 1.2] shows that an arbitrary compact metrizable group \mathbb{G} is of the form $\mathbb{G} \cong \text{Aut}(X, d)$ for a compact metric space (X, d) . Now recast (X, d) as a pure graphon, as follows:

- rescale d so that it takes values in $[0, 1]$, making it into a *kernel* [23, §13.1];
- and consider a fully-supported probability measure μ on X , invariant under \mathbb{G} : one can always be produced by averaging *any* fully-supported measure (extant, as X is second-countable compact Hausdorff) against the Haar measure of \mathbb{G} .

Our graphon is now (X, μ, d) ; it is pure by Corollary 2.3 and its automorphism group is precisely the original \mathbb{G} . ■

2.1 Revisiting a definition

The approach taken in [26, Definition 7] to defining the automorphism group of a graphon is different from that suggested by Remark 1.3: rather than define the automorphisms as classes modulo zero (and perhaps modulo automorphisms that are in other ways “trivial”), loc.cit. first normalizes the graphon to a pure one and then isolates concrete automorphisms thereof:

Definition 2.6 Let (X, W, μ) be a pure graphon. Its *automorphism group* $\text{Aut}_{LS}(X, W, \mu)$ consists of those measure-preserving bijections $\varphi : X \rightarrow X$ such that

$$\forall x \in X, W(\varphi x, \varphi \cdot) = \varphi(x, \cdot) \text{ almost everywhere.} \quad \blacklozenge$$

The ‘LS’ subscript (for Lovász-Szegedy) is temporary; we will see that the definition specializes to something more akin to [42, §1].

Definition 2.7 Let $\Gamma := (X, W, \mu)$ be a graphon (pure or not). Its *automorphism group*

$$\text{Aut}(\Gamma) = \text{Aut}(V, \mathcal{B}, E, \mu)$$

is defined as follows.

- Consider the group $\text{Aut}(X, \mu)$ of mod-0 equivalence classes of measure-preserving automorphisms of the standard probability space (X, μ) .
- Set

$$\widetilde{\text{Aut}}(\Gamma) := \{g \in \text{Aut}(X, \mu) \mid W(g \cdot, g \cdot) = W \text{ almost everywhere}\}.$$

We equip $\widetilde{\text{Aut}}(\Gamma)$ with the *weak topology* (e.g. [12, Weak Topology]).

- Consider, next, the closed normal subgroup

$$\widetilde{\text{Aut}}(\Gamma) \supseteq \widetilde{\text{Aut}}_{triv}(\Gamma) := \left\{ g \in \widetilde{\text{Aut}}(\Gamma) \mid (g \cdot, \cdot) = \text{almost everywhere} \right\}.$$

Finally, set

$$\text{Aut}(\Gamma) := \widetilde{\text{Aut}}(\Gamma) / \widetilde{\text{Aut}}_{triv}(\Gamma)$$

with the quotient topology. ◆

For pure graphons much of the quotienting (by trivial automorphisms, etc.) is not necessary:

Lemma 2.8 *If $\Gamma = (X, W, \mu)$ is a pure graphon then $\widetilde{\text{Aut}}_{triv}(\Gamma)$ is trivial.*

Proof Let $\varphi : X \rightarrow X$ be a measure-preserving and a.e. W -preserving bijection with

$$W(-, -) = W(\varphi-, -) \text{ almost everywhere.} \quad (2-2)$$

We then have

$$\begin{aligned} \int_X r_W(x, \varphi x) \, d\mu(x) &= \int_X \int_X |W(x, y) - W(\varphi x, y)| \, d\mu(y) \, d\mu(x) \quad \text{by Definition 2.1} \\ &= 0 \quad \text{by (2-2).} \end{aligned}$$

Since r_W is assumed to be a metric and μ is fully supported, we have $x = \varphi x$ almost everywhere. In $\widetilde{\text{Aut}}_{triv}(\Gamma)$ we identify (by definition) automorphisms modulo measure zero, so we are done: $\varphi = \text{id}$. ■

Furthermore, the reason why $\text{Aut}_{LS}(\Gamma)$ functions so well is that it provides a section for the quotient by bijections that are trivial modulo measure zero.

Proposition 2.9 *Let $\Gamma = (X, W, \mu)$ be a pure graphon and $\varphi : X \rightarrow X$ a measure-preserving bijection with*

$$W(\varphi-, \varphi-) = W \text{ almost everywhere.}$$

φ is then congruent modulo zero to a unique element of $\text{Aut}_{LS}(\Gamma)$.

Proof We are making two claims: existence and uniqueness. We will drop Γ in the notation for automorphism groups: $\text{Aut}_{LS} = \text{Aut}_{LS}(\Gamma)$.

(1) **Uniqueness.** The assertion to prove is that if two elements of Aut_{LS} coincide modulo 0 then they coincide period. Or equivalently: if $\psi \in \text{Aut}_{LS}$ is trivial mod 0 then it is the identity.

For $x \in X$ we have

$$W(x, -) = W(\varphi x, \varphi-) \text{ a.e.} \quad (2-3)$$

because $\varphi \in \text{Aut}_{LS}$, and the assumption is that

$$\varphi = \text{id}_X \text{ a.e.} \quad (2-4)$$

We then have

$$\begin{aligned} r_W(x, \varphi x) &= \int_X |W(x, y) - W(\varphi x, y)| \, d\mu(y) \quad \text{by Definition 2.1} \\ &= \int_X |W(x, y) - W(\varphi x, \varphi y)| \, d\mu(y) \quad \text{by (2-4)} \\ &= \int_X |W(x, y) - W(x, y)| \, d\mu(y) \quad \text{by (2-3)} \\ &= 0. \end{aligned}$$

Because r_W is a metric, we must have $x = \varphi x$ and we are done: $x \in X$ was arbitrary.

(2) **Existence.** The hypothesis implies that there is a full-measure (hence also dense) $Y \subseteq X$ such that φ preserves r_W -distances between points in Y . But then, because (X, r_W) is by assumption the completion of (Y, r_W) , the restriction $\varphi|_Y$ extends uniquely to an r_W isometry φ' on X .

We have now

- modified φ mod zero to φ'
- so that the latter still preserves W almost everywhere and is an r_W -isometry.

The conclusion that $\varphi' \in \text{Aut}_L S$ follows from [26, Lemma 9] (or rather from its simpler analogue for r_W rather than \bar{r}_W).

This finishes the proof. ■

In summary:

Corollary 2.10 *For a pure graphon $\Gamma = (X, W, \mu)$ the automorphism groups $\text{Aut}_{LS}(\Gamma)$ and $\text{Aut}(\Gamma)$ of Definitions 2.6 and 2.7 coincide.* ■

3 Rigidity

3.1 Graphon-rigid Lie groups

[26, Theorem 37] notes that *weakly random* compact groups (i.e. those that have finitely many isomorphism classes of irreducible representations of dimension $\leq d$ for every d) have the following “permanence” property.

Definition 3.1 A compact metrizable group \mathbb{G} is *graphon-rigid* if the limit of every convergent sequence of Cayley graphons on \mathbb{G} is again a Cayley graphon on \mathbb{G} . ◆

By contrast, [26, Example 35] gives an example of Cayley graphons on the circle converging to a Cayley graphon on a 2-torus; essentially the same example is also discussed in [40, text preceding Theorem 5], and slightly adapted it reads as follows.

Example 3.2 Let $\chi : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ be a generating character for the Pontryagin dual $\mathbb{Z} \cong \widehat{\mathbb{S}^1}$, and consider the Cayley graphons Γ_n with

- underlying Haar probability space (\mathbb{S}^1, μ) ;
- and respective kernels

$$W_n(x, y) = f_n(xy^{-1}), \quad f_n := \Re \frac{2 + \chi + \chi^n}{4}$$

(‘ \Re ’ denoting the real part).

[26, Example 35] observes that this sequence converges to the Cayley graphon on the torus

$$\mathbb{T}^2 = \mathbb{S}^1 \times \mathbb{S}^1$$

with kernel

$$W(x, y) = f(xy^{-1}), \quad f := \Re \frac{2 + \chi_1 + \chi_2}{4},$$

where χ_i , $i = 1, 2$ are the characters generating the Pontryagin duals of the two \mathbb{S}^1 Cartesian factors. That agrees with [40, discussion preceding Theorem 5]; we give one possible proof below, that will also help fit the example into a broader family. ◆

Remark 3.3 [26, Example 35] actually uses the functions

$$\Im \frac{1 + \chi + \chi^n}{2} \quad \text{in place of} \quad \Re \frac{2 + \chi + \chi^n}{4}$$

(i.e. *imaginary* rather than real parts), but this seems to not quite meet the requirements:

- Per [26, Definition 30] one needs “symmetric” functions f_n in the sense that $f_n(x^{-1}) = f_n(x)$. The real parts of the characters have this property, but their imaginary parts do not.
- And additionally, the kernels are meant to take values in $[0, 1]$, whereas both the imaginary and the real part of $1 + \chi + \chi^n$ will, in general, take some negative values. \blacklozenge

To unpack [Example 3.2](#), note first the following simple procedure for producing convergent graphon sequences; recall [2, §8.1] that nets $(\mu_\alpha)_\alpha$ of measures on topological spaces *converge weakly* to measures μ if

$$\int_X f \, d\mu_\alpha \xrightarrow{\alpha} \int_X f \, d\mu$$

for every bounded continuous function f on X .

Lemma 3.4 *Let X be a compact metrizable space and $W : X^2 \rightarrow [0, 1]$ a symmetric continuous function. If a sequence μ_n of probability measures converges weakly to μ , then we correspondingly have graphon convergence*

$$\Gamma_n := (X, W, \mu_n) \xrightarrow{n} (X, W, \mu) =: \Gamma.$$

Proof This is obvious from the characterization of convergence via morphism densities [26, §2.1]:

$$t(F, W_n) \xrightarrow{n} f(F, W)$$

for every finite simple graph $F = (V, E)$. Plainly, the hypothesis ensures that

$$t(F, W_n) := \int_{X^V} \prod_{ij \in E} W(x_i, x_j) \prod_{i \in V} d\mu_n(x_i)$$

converges to the analogous quantity with μ in place of μ_n . \blacksquare

A slightly different take on [Lemma 3.4](#) would be

Lemma 3.5 *Let*

$$\varphi_n : (X_n, \mu_n) \rightarrow (X, \mu)$$

be continuous measure-preserving functions between compact metrizable probability spaces and (X, W, μ) a graphon structure.

Weak measure convergence

$$\varphi_{n*} \mu_n \xrightarrow{n} \mu$$

then implies graphon convergence

$$(X_n, W \circ (\varphi_n^{\times 2}), \mu_n) \xrightarrow{n} (X, W, \mu).$$

Proof Upon recalling ([26, §2.1]) the weak isomorphism

$$(X_n, W \circ (\varphi_n^{\times 2}), \mu_n) \cong (X, W, \varphi_n * \mu_n),$$

the conclusion follows from Lemma 3.4. ■

We can now place Example 3.2 in broader context: it is an instance of Example 3.6, with $k = 1$.

Example 3.6 Let χ be a non-trivial character on \mathbb{S}^1 , as in Example 3.2 (it need not be generating; any non-trivial character will do) and consider, for any

$$n_i \in \mathbb{Z}_{>0}, 1 \leq i \leq k,$$

the function

$$f_{(n_i)} := \Re \frac{1}{2(k+1)} (k+1 + \chi + \chi^{n_1} + \chi^{n_1 n_2} + \cdots + \chi^{n_1 \cdots n_k}). \quad (3-1)$$

Clearly, the $f_{(n_i)}$ are $[0, 1]$ -valued and symmetric, in the sense that they take equal values on $x \in \mathbb{S}^1$ and x^{-1} . They thus induce kernels

$$W_{(n_i)}(x, y) := f_{(n_i)}(xy^{-1})$$

on \mathbb{S}^1 . I claim that as the n_i all increase to infinity, the Haar-measure graphons

$$\Gamma_{(n_i)} := (\mathbb{S}^1, W_{(n_i)}, \mu)$$

converge to the $(k+1)$ -torus-based graphon $\Gamma = (\mathbb{T}^{k+1}, W, \mu_{k+1})$, where μ_k is the Haar measure again and

$$W(x, y) = f(xy^{-1})$$

with f being the symmetric function on \mathbb{T}^{k+1} constructed as in (3-1):

$$f := \Re \frac{1}{2(k+1)} (k+1 + \chi_1 + \cdots + \chi_{k+1}),$$

the summands being $k+1$ non-trivial characters on the $k+1$ factors of the torus.

To verify the claim, note first that it does not matter *which* $\chi_i \in \widehat{\mathbb{T}^{k+1}}$ we select: all choices transform back to the one where all χ_i are generating via a component-wise measure preserving transformation of the torus, which will not alter the weak isomorphism class of the graphon. For that reason, we might as well take φ_i to be the i^{th} -component copy of χ .

Next, observe that the functions $W_{(n_i)}$ are pullbacks of W through the maps

$$\mathbb{S}^1 \ni z \xrightarrow{\varphi_{(n_i)}} (z, z^{n_1}, z^{n_1 n_2}, \dots, z^{n_1 \cdots n_k}) \in \mathbb{T}^{k+1}.$$

It is not difficult to check now that the conditions of Lemma 3.5 are met (except that we are working here with multi-sequences indexed by (n_i) in place of a singly-indexed sequence): the images

$$\varphi_{(n_i)}(\mathbb{S}^1) \subset \mathbb{T}^{k+1}$$

converge to all of \mathbb{T}^{k+1} in the *Hausdorff metric* [3, Definition 7.3.1] on closed subsets of the latter, so any limit point of

$$\varphi_{(n_i)} * \mu \in \text{Prob}(\mathbb{T}^{k+1})$$

will be invariant under translations by the torus. The only such creature is the Haar measure μ_{k+1} , hence the conclusion (that the hypotheses of Lemma 3.5 hold). ◆

Remark 3.7 Defining the Hausdorff metric on the space $\mathcal{CL}(X)$ of closed subsets of a compact space X technically requires the ambient space to be metrized, but the metric will make no difference to the induced (compact [3, Theorem 7.3.8]) topology on $\mathcal{CL}(X)$.

One way to see this is to observe that having fixed a metric (X, d) inducing the underlying topology, convergence in $\mathcal{CL}(X)$ with respect to the induced Hausdorff metric can be defined in terms of just the *uniformity* induced by d ([17, Definition 7.1 and pp.88-89]). Because X is compact Hausdorff that uniformity is in turn uniquely determined by the topology of X [17, Proposition 8.16].

For an alternative take on the matter, note that the Hausdorff-metric topology on $\mathcal{CL}(X)$ (associated to any metric topologizing X) can also be recovered intrinsically, with no reference to a metric, as the topology introduced by Fell on [9, p.472] (see also e.g. [1, Chapter II, Section 2] for a recollection): a subbasis of open sets consists of the sets

$$\{Y \subseteq X \text{ closed} \mid Y \cap K = \emptyset, Y \cap U \neq \emptyset, \forall U \in \mathcal{F}\}.$$

as $K \subseteq X$ ranges over compact subsets and \mathcal{F} over finite collections of open subsets of X ,

The definition makes sense for arbitrary topological spaces X , and makes $\mathcal{CL}(X)$ compact Hausdorff whenever X is *locally* compact Hausdorff [9, Theorem 1]. This is not the case, in general, for Hausdorff-metric topologies, but the various topologies *do* agree when X is compact (rather than just locally so); this being a simple enough exercise, we omit the proof.

In any event, when X is furthermore a compact *group*, the collection $\mathcal{CLG}(X)$ of closed subgroups is closed and hence also compact [9, p.474 (IV)]. In fact, [9, Theorem 1] makes it reasonable to work with $\mathcal{CL}(X)$ (or \mathcal{CLG} , when the ambient space is a group) even when X is compact but *not* metrizable. \blacklozenge

We need a few auxiliary observations for later.

Lemma 3.8 *Let \mathbb{G} be a compact second-countable group. There is a continuous symmetric function $f : \mathbb{G} \rightarrow [0, 1]$ such that*

$$\mathbb{G} \ni x \mapsto f(x \cdot)$$

is injective.

Proof The topology of \mathbb{G} is induced by a metric d , bi-invariant in the sense that

$$d(gx, gy) = d(x, y) = d(xg, yg), \quad \forall g, x, y \in \mathbb{G}$$

([16, Corollary A4.19]). To conclude, set

$$f(x) := d(x, 1).$$

For $x \neq y \in \mathbb{G}$ the translates $f(x \cdot)$ and $f(y \cdot)$ take the value 0 at x^{-1} and y^{-1} respectively, and nowhere else. \blacksquare

And a consequence thereof:

Lemma 3.9 *Every second-countable compact group has a Cayley-graphon structure meeting the requirements of Proposition 2.2.*

Proof Indeed: construct one using a function f as in Lemma 3.8. \blacksquare

It will be useful to distill a general principle operative in Example 3.6.

Lemma 3.10 Let $\Gamma := (\mathbb{G}, W, \mu_{\mathbb{G}})$ be a Cayley graphon on a compact metrizable group and

$$\varphi_n : \mathbb{G}_n \rightarrow \mathbb{G}$$

morphisms of compact metrizable groups.

If $\varphi_n(\mathbb{G}_n)$ converge to \mathbb{G} in the Hausdorff distance induced by any metric topologizing \mathbb{G} then we have convergence

$$\Gamma_n := (\mathbb{G}_n, W \circ \varphi_n^{\times 2}, \mu_{\mathbb{G}_n}) \xrightarrow[n]{} \Gamma$$

of Cayley graphons.

Proof As in Example 3.6: any limit-point of the sequence of pushforward probabilities $\varphi_{*n}\mu_{\mathbb{G}_n}$ will be translation-invariant under the Hausdorff limit

$$\mathbb{G} = \lim_n \varphi_n(\mathbb{G}_n),$$

so must be the Haar measure $\mu_{\mathbb{G}}$. We can then apply Lemma 3.5 to conclude. ■

Incidentally, we can supplement Example 3.6 with non-Lie analogues.

Example 3.11 Let p be a prime number, and denote by

$$\mathbb{Z}_p := \varprojlim_n \mathbb{Z}/p^n \tag{3-2}$$

the profinite group of p -adic integers [16, Example 1.28].

Equip the circle \mathbb{S}^1 with a Cayley-graphon structure (\mathbb{S}^1, W, μ) as in Lemma 3.9, and take for the maps φ_n of Lemma 3.10 the morphisms

$$\mathbb{Z}_p \rightarrow \mathbb{Z}/p^n \hookrightarrow \mathbb{S}^1$$

obtained by first surjecting along the canonical map provided by (3-2) and then embedding the finite cyclic group as the $(p^n)^{th}$ roots of unity.

The hypothesis of Lemma 3.10 clearly holds, and hence so does its conclusion. We thus have \mathbb{Z}_p -Cayley graphons converging to a circle-based Cayley graphon. The latter is not \mathbb{Z}_p -Cayley though: the p -adics would have to act transitively (and continuously: Lemma 2.4) on the manifold \mathbb{S}^1 , which would entail that the action factor through a Lie ([41, Proposition 1.6.5]) and hence finite quotient. Naturally, finite groups do not act transitively on the circle. ◆

To make sense of Theorem 3.12, recall [16, Definition 9.5] that a compact connected group \mathbb{G} is *semisimple* if it equals its *commutator subgroup* \mathbb{G}' ; this is the subgroup generated by commutators

$$[x, y] := xyx^{-1}y^{-1}, \quad x, y \in \mathbb{G}.$$

According to [16, Theorem 9.2] it is also just the *set* of such commutators, and is always connected and closed in \mathbb{G} .

We also remind the reader that *simple* compact groups [16, Definition 9.88] are those having no proper, non-trivial normal subgroups (or equivalently [16, Theorem 9.90], no such *closed* subgroups).

For compact *Lie* groups, it turns out that [26, Theorem 37] in fact provides a characterization of graphon-rigidity. The following result collects that along with a number of other such characterizations.

Theorem 3.12 *Let \mathbb{G} be a compact Lie group and \mathbb{G}_0 its connected identity component. The following conditions are equivalent.*

- (a) \mathbb{G} is graphon-rigid.
- (b) \mathbb{G}_0 is graphon-rigid.
- (c) \mathbb{G}_0 is semisimple.
- (d) \mathbb{G}_0 has finite center.
- (e) \mathbb{G}_0 does not admit a surjection $\mathbb{G} \rightarrow \mathbb{S}^1$ onto the circle group.
- (f) Modulo a finite central subgroup, \mathbb{G}_0 is a product of finitely many compact connected simple Lie groups.
- (g) More precisely, there are compact connected simple Lie groups \mathbb{S}_i , $1 \leq i \leq n$, finite central subgroups $\mathbb{N}_i \leq \mathbb{S}_i$, and surjections

$$\prod_{i=1}^n \mathbb{S}_i \longrightarrow \mathbb{G}_0 \longrightarrow \prod_{i=1}^n \mathbb{S}_i/\mathbb{N}_i. \quad (3-3)$$

- (h) \mathbb{G} is weakly random.
- (i) \mathbb{G}_0 is weakly random.

We first dispose of the relationship between conditions (a) and (b), in slightly more general form than needed. Wherever compact groups act transitively on graphons (X, W, μ) , we will assume the latter pure; it follows from [26, Corollary 16] (and its proof) that they are also compact, with topology induced by either of the metrics r_W or \bar{r}_W .

Proposition 3.13 *Let \mathbb{G} be a compact metrizable group whose connected component $\mathbb{G}_0 \leq \mathbb{G}$ has finite index. If \mathbb{G} is graphon-rigid so is \mathbb{G}_0 .*

Proof Let

$$\Gamma_n \xrightarrow[n]{} \Gamma, \quad \Gamma_n \text{ } \mathbb{G}_0\text{-Cayley}$$

be a convergent sequence of graphons.

As all $\Gamma_n := (X_n, W_n, \mu_n)$ come equipped with transitive (measure-preserving) \mathbb{G}_0 -actions, we can form the *induced* probability \mathbb{G} -spaces in the sense of [20, Appendix G]:

- Set

$$\overline{X}_n := \text{Ind}_{\mathbb{G}_0}^{\mathbb{G}} X_n := X_n \times \mathbb{G}/\mathbb{G}_0$$

as a space.

- The probability measure is

$$\overline{\mu}_n := \mu_n \times (\text{normalized counting measure on } \mathbb{G}/\mathbb{G}_0).$$

- The kernel \overline{W}_n is a copy of the original W_n on each individual copy of X_n . For points

$$(x, s) \in X_n \times \{s\}, (x', s') \in X_n \times \{s'\}, s \neq s' \in \mathbb{G}/\mathbb{G}_0$$

we can set, for instance,

$$\overline{W}_n((x, s), (x', s')) = 0;$$

this will do (making the resulting induced graphon pure) provided X_n was not a singleton with $W \equiv 0$. We assume X_n and X are at any rate not singletons, since otherwise there would be (almost) nothing to prove.

- The (also transitive) \mathbb{G} -action is as explained in [20, Appendix G]: having chosen a finite set $S \subset \mathbb{G}$ of representatives for the cosets \mathbb{G}/\mathbb{G}_0 , the action of $g \in \mathbb{G}$ is

$$g(x, s) = (hx, s') \text{ for the unique } gs = s'h, h \in \mathbb{G}_0, s, s' \in S.$$

The analogous construction applies to $\Gamma := (X, W, \mu)$ to produce

$$\overline{\Gamma} := \text{Ind}_{\mathbb{G}_0}^{\mathbb{G}} \Gamma = (\overline{X}, \overline{W}, \overline{\mu}),$$

and it is easy to see that this induction procedure is continuous:

$$\overline{\Gamma}_n \xrightarrow[n]{} \overline{\Gamma}.$$

The hypothesis says that the limit $\overline{\Gamma}$ is \mathbb{G} -Cayley. The orbits of the *connected* compact group \mathbb{G}_0 are each contained within single “slices”

$$X \cong X \times \{s\}, s \in S \cong \mathbb{G}/\mathbb{G}_0, \tag{3-4}$$

so each orbit has measure $\leq \frac{1}{[\mathbb{G}:\mathbb{G}_0]}$. But the orbits also cover X (by \mathbb{G} -transitivity), and there must be at least $[\mathbb{G}:\mathbb{G}_0]$ of them. It follows that each \mathbb{G}_0 -orbit constitutes a single leaf (3-4); as those are isomorphic to the original Γ , we are done. \blacksquare

Proof of Theorem 3.12 We do this in stages.

(Part 1): (c) through (g) are mutually equivalent. This is standard compact-group structure theory: according to [16, Theorem 9.24], say, an arbitrary compact connected \mathbb{G}_0 can be sandwiched as in (3-3), with finite central kernels, except that perhaps there are additional torus factors on the two sides (apart from the semisimple factors). Those torus factors are absent precisely when \mathbb{G}_0 is semisimple [16, Theorem 9.19, Corollary 9.20].

(Part 2): (h) \Leftrightarrow (i). The identity component $\mathbb{G}_0 \leq \mathbb{G}$ has finite index because \mathbb{G} is compact and Lie. The usual “Mackey machine” applies ([18, Theorems 4.64 and 4.65]) to conclude that

- every irreducible \mathbb{G} -representation is a sum of at most $[\mathbb{G}:\mathbb{G}_0]$ conjugates under \mathbb{G} of an irreducible \mathbb{G}_0 -representation;
- and every irreducible \mathbb{G}_0 -representation arises in this fashion, as a summand of an irreducible \mathbb{G} -representation.

This gives a relation between irreducible \mathbb{G} - and \mathbb{G}_0 -representations: two such are related precisely when one contains the other. Related representations have dimensions differing by a factor of at most $[\mathbb{G}:\mathbb{G}_0]$, and the relation is at most $[\mathbb{G}:\mathbb{G}_0]$ -to-one.

It follows, then, that a cap of $N \in \mathbb{Z}_{>0}$ on the number of irreducible \mathbb{G} -representations of dimension $\leq d$ will give a cap of $[\mathbb{G} : \mathbb{G}_0]N$ on the number of irreducible \mathbb{G}_0 -representations of dimension $\leq \frac{d}{[\mathbb{G} : \mathbb{G}_0]}$ and vice versa.

(Part 3): (a) \Rightarrow (b). This follows from Proposition 3.13, since for compact *Lie* groups the identity component is also open and hence has finite index.

This, so far, breaks up the various conditions into three aggregates:

- (1) (a) \Rightarrow (b).
- (2) (c) up to (g), all mutually equivalent.
- (3) (h) and (i), all mutually equivalent.

To establish *inter*-group equivalence, note first that (3) implies (1) by [26, Theorem 37], so only two other implications are needed.

(Part 4): (2) \Rightarrow (3). This says that semisimple compact connected Lie groups are weakly random; the claim follows easily, say, from familiar Lie-group/algebra representation theory via, say, the *Weyl dimension formula* for representations: [38, Theorem IX.6.1].

(Part 5): (1) \Rightarrow (2). Or the contrapositive: if \mathbb{G}_0 *does* admit a surjection onto \mathbb{S}^1 , then it cannot be graphon-rigid.

By [16, Theorem 9.24], \mathbb{G}_0 surjects with finite central kernel onto a Lie group of the form

$$\mathbb{T}^\ell \times \prod_{i=1}^n \mathbb{S}_i, \quad \ell \geq 1, \mathbb{S}_i \text{ simple.} \quad (3-5)$$

That finite kernel will make no material difference to the argument, so we will simplify matters and assume \mathbb{G}_0 is (3-5) to begin with.

Now consider a somewhat larger group

$$\mathbb{H} := \mathbb{T}^{\ell+1} \times \prod_{i=1}^n \mathbb{S}_i$$

(note the higher torus dimension), and equip it with a Cayley graphon structure $\Gamma := (\mathbb{H}, W, \mu)$ as in Lemma 3.9, so that the conditions of Proposition 2.2 hold.

Next, consider maps

$$\varphi_n : \mathbb{G}_0 \rightarrow \mathbb{H}$$

that

- operate as the identity on the \mathbb{S}_i and all but one of the ℓ circle factors in \mathbb{T}^ℓ ;
- and wind one of the circle factors ever more tightly around \mathbb{T}^2 via, say,

$$\mathbb{S}^1 \ni z \xrightarrow{\varphi_n} (z, z^n) \in \mathbb{T}^2$$

(as in Example 3.6). It follows from Lemma 3.10 that the Cayley graphons

$$\Gamma_n := (\mathbb{G}_0, W \circ \varphi_n^{\times 2}, \mu_{\mathbb{G}_0}), \quad n \in \mathbb{Z}_{>0}$$

converge to Γ . I claim, however, that \mathbb{G}_0 cannot act transitively on their limit Γ .

Indeed, such an action would be continuous on the compact metric space (\mathbb{H}, r_W) (Lemma 2.4), so would identify \mathbb{H} homeomorphically with the homogeneous space

$$\mathbb{G}_0/\mathbb{G}_{0,p} \text{ (for some } p \in \mathbb{H}\text{)}$$

by the (closed, hence Lie [21, Theorem 20.12]) stabilizer group $\mathbb{G}_{0,p} \leq \mathbb{G}_0$ of a point $p \in \mathbb{H}$. But that homogeneous space is a topological manifold of dimension

$$\dim \mathbb{G}_0 - \dim \mathbb{G}_{0,p} \leq \dim \mathbb{G}_0$$

[21, Theorem 21.17], while \mathbb{H} is a manifold of strictly larger dimension $\dim \mathbb{G}_0 + 1$. That the two cannot be homeomorphic then follows from dimension invariance [21, Theorem 17.26].

This concludes the proof. ■

It is perhaps worth noting at this stage that for general (non-Lie) compact metrizable groups graphon-rigidity does *not* imply weak randomness: the latter is thus the strictly stronger condition.

Example 3.14 Let $\mathbb{G} = (\mathbb{Z}/2)^{\aleph_0}$, the product of countably infinitely many copies of $\mathbb{Z}/2$. Naturally, \mathbb{G} is not weakly random: being abelian *all* of its (infinitely many) irreducible representations are 1-dimensional. I claim that it is nevertheless graphon-rigid.

To see this, note that by the argument employed in the proof of [26, Theorem 39], any limit graphon of a sequence of \mathbb{G} -Cayley graphons will be acted upon transitively by an abelian group \mathbb{A} of exponent ≤ 2 (i.e. such that $x^2 = 1$ for all elements x).

Any compact group of exponent ≤ 2 is a product of copies of $\mathbb{Z}/2$, and since all groups in sight are metrizable, at most countably many copies at that. \mathbb{G} will then surject onto \mathbb{A} , hence the conclusion. ◆

Recall the following useful device from [26, §4.2].

Definition 3.15 Let $\Gamma := (X, W, \mu)$ be a graphon and $r > 0$. The *truncation* $[\Gamma]_r$ is the graphon obtained from Γ as follows:

- Regard W as a *Hilbert-Schmidt operator* [13, §4] on $L^2 := L^2(X, \mu)$ by

$$(Wf)(x) := \int_X W(x, y)f(y) \, d\mu(y), \quad f \in L^2.$$

- Let λ . Denoting by P_λ the orthogonal projection onto the (finite-dimensional) λ -eigenspace of W , the *truncation* $[W]_r$ is

$$\sum_{|\lambda| \geq r} \lambda P_\lambda.$$

It is again a kernel, as explained in [26, §4.2].

- Then set

$$[\Gamma]_r := (X, [W]_r, \mu). \quad \text{◆}$$

Remark 3.16 The truncation procedure of Definition 3.15 gives a kind of canonical “finite-dimensional approximation” of the graphon Γ . For $r > 0$ the purification of the truncated $[\Gamma]_r$ can be realized as a subspace of

$$V_r := \bigoplus_{|\lambda| \geq r} (\lambda\text{-eigenspace of } W),$$

equipped with the kernel

$$V_r^2 \ni (x, y) \mapsto \sum_i \lambda_i x_i y_i$$

for an enumeration (λ_i) of the relevant eigenvalues $|\lambda| \geq r$ and choice of coordinates

$$V_r \ni x = (x_i)_i \in \mathbb{R}^{\dim V_r}.$$

Furthermore, it follows from [26, Lemma 17] that if $r > 0$ is “generic enough” (specifically, not an eigenvalue of Γ), then for any convergent sequence $\Gamma_n \rightarrow \Gamma$ the truncations $[\Gamma_n]_r$ also converge to $[\Gamma]_r$, the measures associated with a subsequence thereof converge weakly to that attached to $[\Gamma]_r$, etc. \blacklozenge

Example 3.14 replicates for *connected* compact groups as well.

Example 3.17 Consider the Pontryagin dual $\widehat{\mathbb{Q}}$ of the additive group of rationals. Since \mathbb{Q} is a filtered colimit (in the category of discrete abelian groups) of copies of \mathbb{Z} , we have a description

$$\widehat{\mathbb{Q}} \cong \varprojlim \mathbb{S}^1 \tag{3-6}$$

as a limit (in the category of compact abelian groups), where the connecting maps are $z \mapsto z^n$ for various n .

This time the group will be an infinite product $\mathbb{G} = \widehat{\mathbb{Q}}^{\mathbb{N}_0}$. Once more, being abelian, it has only 1-dimensional irreducible representations. We argue that it too is graphon-rigid however, much along the same lines as in Example 3.14:

Let $\Gamma := (X, W, \mu)$ be the limit of a sequence of \mathbb{G} -Cayley graphons. Γ is the limit of its truncations $[\Gamma]_\alpha$ (for $\alpha > 0$), as in Definition 3.15. These are in turn limits of the corresponding truncations $[\Gamma_n]_\alpha$ of the Γ_n .

The $[\Gamma_n]_\alpha$ are acted upon transitively by Lie-group quotients of \mathbb{G} , i.e. tori \mathbb{T}_n . Assuming, as in the proof of [26, Theorem 39], that $[\Gamma_n]_\alpha$ as well as $[\Gamma]_\alpha$ are all supported in the same Euclidean space \mathbb{R}^d , a limit (perhaps of some subsequence)

$$\lim_n \mathbb{T}_n =: \mathbb{T} \leq \text{orthogonal group } \mathbb{O}(d)$$

in the Hausdorff topology of Remark 3.7 on (the compact space $\mathcal{CLG}(\mathbb{O}(d))$) of closed subgroups of $\mathbb{O}(d)$ will act transitively on the underlying space of $[\Gamma]_\alpha$. But such a limit \mathbb{T} is again a torus (being Lie connected and abelian [6, Theorem 4.2.4], properties easily seen to be preserved upon passage to limits in \mathcal{CLG}).

Letting $\alpha \rightarrow 0$, the inverse limit \mathbb{H} of these groups \mathbb{T} will act transitively on Γ . That limit is Pontryagin-dual to an infinite torsion-free discrete abelian group, which embeds into

$$\mathbb{Q}^{\oplus \mathbb{N}_0} \cong \widehat{\mathbb{G}};$$

for that reason, \mathbb{G} surjects onto \mathbb{H} .

Conclusion: a limit of \mathbb{G} -Cayley graphons admits a transitive \mathbb{G} -action. \blacklozenge

A familiar phenomenon, when studying symmetries of objects parametrized by a space, is that of the *upper semicontinuity* of their groups of automorphisms. A case in point is that of Riemannian structures: consider a compact smooth manifold M and the space $\text{RIEM}(M)$ of Riemannian structures, appropriately topologized [8, §1]. The diffeomorphism group \mathbb{D} of M acts on $\text{RIEM}(M)$, and the isotropy group

$$\mathbb{D}_{(M,g)} := \{\gamma \in \mathbb{D} \mid \gamma \text{ fixes } (M, g) \in \text{RIEM}(M)\}$$

of a point (i.e. Riemannian structure) is nothing but the isometry group $\text{Aut}(M, g)$: always compact and Lie. It turns out that for points (M, g') close to (M, g) the automorphism groups $\text{Aut}(M, g')$ are “almost contained” in $\text{Aut}(M, g)$ [8, Theorem 8.1]: they are conjugate in \mathbb{D} to subgroups of $\text{Aut}(M, g)$ by elements close to the identity. This upper semicontinuity phenomenon does *not* obtain for graphons (at least not in anything like the same fashion):

Example 3.18 Consider *any* compact probability metric space (X, d, μ) as a graphon, and scale the distance towards 0. This produces a net converging to a singleton equipped with the zero kernel; all members (X, cd, μ) of the net (for scalars $c > 0$) have the same automorphism group, which may well be non-trivial, while naturally, the limit has trivial automorphism group. \blacklozenge

Or more interestingly:

Example 3.19 According to Theorem 2.5 there is a graphon $\Gamma := (X, W, \mu)$ having $\widehat{\mathbb{Q}}$ as an automorphism group:

$$\text{Aut}(\Gamma) \cong \widehat{\mathbb{Q}}.$$

The eigenvalue-truncations of Γ (as in [26, §§4.2 and 4.4] and Example 3.17) all have automorphism groups that are

- compact Lie;
- and quotients of $\text{Aut}(\Gamma)$.

The only compact Lie quotients of $\widehat{\mathbb{Q}}$ are also connected and abelian, so they must be tori [15, Chapter II, Exercise C.2]. But they cannot be *higher-dimensional* tori \mathbb{T}^d , $d \geq 2$ because

$$\widehat{\mathbb{T}}^d \cong \mathbb{Z}^d, \quad d \geq 2$$

does not embed into $\mathbb{Q} \cong \widehat{\widehat{\mathbb{Q}}}$ [16, Theorem 7.63].

In short: Γ , with automorphism group $\widehat{\mathbb{Q}}$, is a limit of graphons with automorphism groups isomorphic to \mathbb{S}^1 . It is enough to note now that the latter cannot embed into $\widehat{\mathbb{Q}}$, because its dual \mathbb{Z} is not a quotient of \mathbb{Q} . \blacklozenge

3.2 Stable compact-group images

By way of motivation for the subsequent material:

Remark 3.20 Starting with a $\widehat{\mathbb{Q}}$ -Cayley graphon, its truncations are \mathbb{S}^1 -Cayley (arguing as in Example 3.19: they admit transitive actions by Lie quotients of $\widehat{\mathbb{Q}}$, etc.). It follows, then, that the circle \mathbb{S}^1 fails to be graphon-rigid through two qualitatively distinct mechanisms:

- (a) it can “wrap around” higher-dimensional tori increasingly densely, as in Example 3.6 (and [26, Example 35]);
- (b) and it self-surjects

$$\mathbb{S}^1 \ni z \mapsto z^n \in \mathbb{S}^1$$

with increasingly large kernel, giving rise to the inverse limit (3-6).

Both of these phenomena provide examples of \mathbb{S}^1 -Cayley graphons converging to a limit not acted upon transitively by \mathbb{S}^1 . \blacklozenge

The following notion is intended to isolate and address the type of behavior noted in item (a) of Remark 3.20.

Definition 3.21 A compact group \mathbb{G} is *image-rigid* if for every compact Lie group \mathbb{H} the images of morphisms $\mathbb{G} \rightarrow \mathbb{H}$ constitute a closed set of groups in the space $\mathcal{CLG}(\mathbb{H})$ of Remark 3.7. \blacklozenge

An equivalent formulation for image-rigidity, occasionally useful:

Lemma 3.22 A compact group \mathbb{G} is image-rigid in the sense of Definition 3.21 if and only if it satisfies the following property:

\mathbb{G} surjects onto every compact Lie group \mathbb{H} belonging to the closure in $\mathcal{CLG}(\mathbb{H})$ of images of morphisms $\mathbb{G} \rightarrow \mathbb{H}$.

Proof That Definition 3.21 is formally stronger than the present requirement is clear, so only the other implication is interesting.

Assume, then, that \mathbb{G} has the property in the statement, and consider morphisms

$$\varphi_n : \mathbb{G} \rightarrow \mathbb{K}, \quad \mathbb{K} \text{ compact Lie}$$

with

$$\varphi_n(\mathbb{G}) \xrightarrow[n]{} \mathbb{H} \leq \mathbb{K} \quad \text{in } \mathcal{CLG}(\mathbb{K}).$$

By [28, Theorem 1 and its Corollary], for sufficiently large n we can find

$$\mathbb{K} \ni x_n \xrightarrow[n]{} 1, \quad x_n \varphi_n(\mathbb{G}) x_n^{-1} \leq \mathbb{H}.$$

Replacing φ_n with their respective conjugates $x_n \cdot \varphi(-) \cdot x_n^{-1}$, we can place ourselves entirely within \mathbb{H} ; the property in the statement then applies to deliver the conclusion (that \mathbb{G} surjects onto \mathbb{H}). \blacksquare

Examples will be familiar by now.

Example 3.23 As noted repeatedly, a circle \mathbb{S}^1 is not image-rigid: the limit in $\mathcal{CLG}(\mathbb{T}^2)$ of the images of

$$\mathbb{S}^1 \ni z \mapsto (z, z^n) \in \mathbb{T}^2 \text{ as } n \rightarrow \infty$$

is the full 2-dimensional torus, but of course *it* is not an image of the circle.

By the same token, none of the tori \mathbb{T}^n are image-rigid: they all wrap around larger-dimensional tori in the same fashion. \blacklozenge

In fact, for Lie groups we have just recovered another facet of Theorem 3.12.

Proposition 3.24 For a compact Lie group \mathbb{G} the following conditions are equivalent (and hence also equivalent to those of Theorem 3.12):

- (a) \mathbb{G} has semisimple connected component \mathbb{G}_0 .
- (b) For every compact Lie group \mathbb{H} the space of morphisms $\mathbb{G} \rightarrow \mathbb{H}$ is compact in the compact-open topology.
- (c) \mathbb{G} is image-rigid.

Before moving on to the proof, we pause to note that things are again somewhat more complicated in the non-Lie case.

Example 3.25 An *infinite*-dimensional torus \mathbb{T}^{\aleph_0} is image-rigid: its images $\mathbb{T}^{\aleph_0} \rightarrow \mathbb{H}$ in a compact Lie group \mathbb{H} are all compact and connected and hence (finite-dimensional) tori [6, Theorem 4.2.4], as are limits thereof in $\mathcal{CLG}(\mathbb{H})$; \mathbb{T}^{\aleph_0} surjects onto every finite-dimensional torus, hence the claim.

On the other hand, \mathbb{T}^{\aleph_0} is not graphon-rigid. We noted in Example 3.19 and Remark 3.20 that $\widehat{\mathbb{Q}}$ -Cayley graphons will be limits of \mathbb{S}^1 -Cayley ones, which in turn are also \mathbb{T}^{\aleph_0} -Cayley. On the other hand, the infinite torus cannot act continuously and transitively on $\widehat{\mathbb{Q}}$.

One way to verify this last claim is to observe that \mathbb{T}^{\aleph_0} is path-connected, while $\widehat{\mathbb{Q}}$. These claims, in turn, follow from [16, Theorem 8.62]: according to that result the quotient of a compact abelian group \mathbb{G} by the (possibly non-closed) path component of the identity is the *ext group* [16, Definition A1.51] $\text{Ext}(\widehat{\mathbb{G}}, \mathbb{Z})$. Now,

- For $\mathbb{G} = \mathbb{T}^{\aleph_0}$ we have

$$\text{Ext}(\widehat{\mathbb{G}}, \mathbb{Z}) \cong \text{Ext}(\mathbb{Z}^{\oplus \aleph_0}, \mathbb{Z}) = \{0\}$$

(e.g. by [16, Proposition A1.14]), hence path connectedness.

- On the other hand, for $\mathbb{G} = \widehat{\mathbb{Q}}$ we have

$$\text{Ext}(\widehat{\mathbb{G}}, \mathbb{Z}) \cong \text{Ext}(\mathbb{Q}, \mathbb{Z}) \cong \mathbb{R} \neq \{0\}$$

by [43]. ◆

Some preparation will help simplify the argument.

Lemma 3.26 *For any compact group \mathbb{G} the set of closed subgroups with abelian connected component is closed in $\mathcal{CLG}(\mathbb{G})$.*

Proof Expressing an arbitrary compact group as a limit of compact *Lie* groups [16, Corollary 2.43], it is enough to focus on these: throughout the proof, then, \mathbb{G} will be Lie.

Every closed subgroup $\mathbb{H} \leq \mathbb{G}$ with abelian identity component is a product

$$\mathbb{H} = \mathbb{H}_0 \cdot \mathbb{F}, \quad \mathbb{F} \leq \mathbb{H} \text{ finite}$$

by [16, Theorem 6.10], so a limit of a net \mathbb{H}_n can be recovered (perhaps after passing to a subnet) as the product between the (connected, abelian) limit of $\mathbb{H}_{n,0}$ and that of finite subgroups $\mathbb{F}_n \leq \mathbb{H}_n$. It thus suffices to prove the main claim for *finite* groups.

That, in turn, follows from the fact that \mathbb{G} being Lie and hence a matrix group [16, Corollary 9.58], there is a positive integer N such that every finite subgroup thereof has an abelian subgroup of index $\leq N$ [5, Theorem 36.13]. The limit of a convergent sequence of finite groups \mathbb{F}_n will again have an abelian subgroup of index $\leq N$, and we are done. ■

The lemma in turn helps with the following result, intended to reduce (part of) the problem to groups with abelian identity component. As in Theorem 3.12 (and the discussion preceding it), denote by \mathbb{G}_0 the identity component of a compact group \mathbb{G} and by $\mathbb{G}' \leq \mathbb{G}$ the (algebraic) commutator subgroup; it will often be closed automatically, e.g. when \mathbb{G} is connected [16, Theorem 9.2] or Lie [16, Theorem 6.11].

Proposition 3.27 *If a compact group \mathbb{G} is image-rigid in the sense of Definition 3.21, so are \mathbb{G}/\mathbb{G}'_0 and $\mathbb{G}_{ab} := \mathbb{G}/\mathbb{G}'$.*

Compact Lie groups are very close to connected: their connected components are open and hence cofinite. At the other extreme stand *profinite* groups: inverse limits of finite groups or, equivalently, totally disconnected compact groups [16, Theorem 1.34]. For abelian ones, at least, the two notions of rigidity converge (together with a complete characterization of those that enjoy that property):

Theorem 3.28 *For a profinite abelian group \mathbb{G} the following conditions are equivalent.*

(a) \mathbb{G} is image-rigid.

(b) We have

$$\mathbb{G} \cong \prod_{i \in I} \mathbb{Z}/n_i, \quad (n_i)_i \text{ bounded.} \quad (3-7)$$

(c) \mathbb{G} is graphon-rigid.

Proof We complete an implication cycle.

(a) \Rightarrow (b). Having an isomorphism (3-7) is equivalent to the discrete Pontryagin dual $\widehat{\mathbb{G}}$ being of the form

$$\widehat{\mathbb{G}} \cong \bigoplus_i \mathbb{Z}/n_i, \quad (n_i)_i \text{ bounded.} \quad (3-8)$$

The dual $\widehat{\mathbb{G}}$ is in any event torsion (being dual to a profinite group [16, Corollary 8.5]), and (3-8) is in turn equivalent (by [19, Theorem 6], say) to $\widehat{\mathbb{G}}$ being of *bounded order*: there is an upper bound on the order of an element of $\widehat{\mathbb{G}}$.

Suppose, now, that $\widehat{\mathbb{G}}$ is *not* of bounded order. This means that we can find characters

$$\chi : \mathbb{G} \rightarrow \mathbb{S}^1$$

of arbitrarily large order, and hence with arbitrarily large image. But then images of such characters will converge in $\mathcal{CLG}(\mathbb{S}^1)$ to the whole circle, which of course is not an image of the profinite group \mathbb{G} (images of compact profinite groups are profinite [16, Exercise E1.13]). Image rigidity is thus violated.

(b) \Rightarrow (c). This is a slight elaboration on (and adaptation of) Example 3.14: a limit of Cayley graphons over (3-7) will be acted upon transitively by an inverse limit of quotients of that group, but it is easy to see that the product in question surjects onto any such inverse limit.

(c) \Rightarrow (a). The claim goes through without the abelianness assumption; we relegate it to Proposition 3.29. ■

Proposition 3.29 *A compact graphon-rigid group \mathbb{G} is image-rigid if it is either*

(a) *profinite;*

(b) *or connected.*

Proof In evaluating image rigidity, Lemma 3.22 provides a compact Lie group \mathbb{H} and morphisms

$$\varphi_n : \mathbb{G} \rightarrow \mathbb{H}$$

whose images converge to all of \mathbb{H} in $\mathcal{CLG}(\mathbb{H})$. The argument now bifurcates.

(a): **profinite groups.** Failing image rigidity, Lemma 3.26 allows us to assume that \mathbb{H} has abelian non-trivial identity component \mathbb{H}_0 . But then the argument of Example 3.11 applies to produce \mathbb{G} -Cayley graphons converging to a \mathbb{H} -Cayley graphon, while at the same time showing that the latter does not admit a transitive \mathbb{G} -action.

(b): **connected groups.** This time \mathbb{H} must be connected. We consider two cases.

(I) \mathbb{H} is **non-abelian**. It is then the isometry group of a left-invariant metric (\mathbb{H}, d) by [31, Theorem 1.3]. The Haar measure μ will then complete a graphon structure

$$\Gamma := (\mathbb{H}, d, \mu)$$

(perhaps after rescaling d so that it takes values in the unit interval), and the morphisms φ_n give us a sequence of \mathbb{G} -Cayley graphons converging to Γ as in Lemma 3.10. Graphon rigidity means that we have a morphism

$$\mathbb{G} \rightarrow \text{Aut}(\Gamma) \cong \mathbb{H}.$$

Its image must act transitively on \mathbb{H} by translation, so the morphism in question is onto; hence the conclusion.

(II) \mathbb{H} is **abelian**. So that it must be a torus

$$\mathbb{T}^\ell \cong \mathbb{R}^\ell / \mathbb{Z}^\ell, \ell \in \mathbb{Z}_{>0}$$

(being compact, connected, abelian and Lie [6, Theorem 4.2.4]). Now equip \mathbb{T}^ℓ with its metric d induced by the standard Euclidean norm on \mathbb{R}^ℓ and its Haar measure μ . It is easy to see that the \mathbb{T}^ℓ -translations constitute precisely the identity component $\text{Aut}(\Gamma)_0$ of

$$\text{Aut}(\Gamma), \quad \Gamma := (\mathbb{T}^\ell, d, \mu).$$

Now conclude as previously: the morphisms φ_n generate \mathbb{G} -Cayley graphon structures converging to Γ , so that graphon rigidity will give a morphism

$$\varphi : \mathbb{G} \rightarrow \text{Aut}(\Gamma)$$

whose image

- on the one hand is contained in $\text{Aut}(\Gamma)_0 \cong \mathbb{T}^\ell$ because \mathbb{G} is connected;
- and on the other hand, operates transitively on the torus by translation.

In short, φ is onto. ■

4 Graphing automorphisms

Recall the *graphings* of [22, §2]:

Definition 4.1 A *graphing* $\Gamma = (V, \mathcal{B}, E, \mu)$ consists of a standard probability space (V, \mathcal{B}, μ) and a graph (V, E) with no loops such that

- $E \subset V \times V$ is a Borel set;
- the degree of (E, V) is uniformly bounded by some positive integer D ;
- and the degree-symmetry condition

$$\int_A \text{deg}_B(x) \, d\mu(x) = \int_B \text{deg}_A(x) \, d\mu(x) \tag{4-1}$$

holds for all Borel sets $A, B \in \mathcal{A}$, where $\text{deg}_B(x)$ denotes the number of elements in B connected to x . ◆

Following [22, §2], the *edge measure* of a graphing $\Gamma = (V, \mathcal{B}, E, \mu)$ is defined by

$$\eta(A \times B) = \eta_\Gamma(A \times B) := \int_A \deg_B(x) \, d\mu(x)$$

for $A, B \in \mathcal{B}$.

In light of Remark 1.3 (specifically, part (c)), the most straightforward route to defining the automorphism group of a graphing is via the Borel measure algebra:

Definition 4.2 Let $\Gamma = (V, \mathcal{B}, E, \mu)$ be a graphing. Its *automorphism group* $\text{Aut}(\Gamma)$ consists of those automorphisms of the measure algebra (\mathcal{B}, μ) that also preserve the edge measure η_Γ on $\mathcal{B} \times \mathcal{B}$. \blacklozenge

By contrast to the automorphism group of a graphon, which is compact by [26, Theorem 10], that of a graphing need not be:

Example 4.3 If the edge-set E is empty, the automorphism group $\text{Aut}(V, \mathcal{B}, E, \mu)$ is nothing but the full automorphism group of the standard probability space (V, \mathcal{B}, μ) . If that space is non-atomic (and hence can be identified with the unit interval with its Lebesgue measure), or even as soon as its non-atomic component is non-empty, that group is certainly non-compact. \blacklozenge

Recall [22, §2]:

Definition 4.4 A *full subgraphing* of a graphing (V, \mathcal{B}, E, μ) is a full subgraph supported by a full-measure Borel subset of V . \blacklozenge

It is immediate that full subgraphings retain the automorphism groups of their larger ambient graphings:

Proposition 4.5 For a full subgraphing Γ' of a graphing $\Gamma = (V, \mathcal{B}, E, \mu)$ we have an isomorphism $\text{Aut}(\Gamma') \cong \text{Aut}(\Gamma)$. \blacksquare

As explained in [22, §2], full subgraphings are both *locally-globally equivalent* and plain *locally equivalent* to the original graphings in the sense of [14, Definition 3.3]. Local equivalence alone does not suffice to preserve the automorphism group:

Example 4.6 The graph(ing) consisting of a single vertex and no edges is locally equivalent to the two-vertex discrete graph assigning probability $\frac{1}{2}$ to each of the vertices (by [14, Example 3.5], for instance). Clearly though, the first graph has trivial automorphism group whereas in the second we can interchange the vertices (measure-preservingly). \blacklozenge

References

- [1] Louis Auslander and Calvin C. Moore. Unitary representations of solvable Lie groups. *Mem. Amer. Math. Soc.*, 62:199, 1966.
- [2] V. I. Bogachev. *Measure theory. Vol. I, II*. Springer-Verlag, Berlin, 2007.
- [3] Dmitri Burago, Yuri Burago, and Sergei Ivanov. *A course in metric geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.

- [4] Alexandru Chirvasitu, Frieder Ladisch, and Pablo Soberón. Finite groups as prescribed polytopal symmetries. *Israel J. Math.*, 245(1):75–91, 2021.
- [5] Charles W. Curtis and Irving Reiner. *Representation theory of finite groups and associative algebras*. AMS Chelsea Publishing, Providence, RI, 2006. Reprint of the 1962 original.
- [6] Anton Deitmar and Siegfried Echterhoff. *Principles of harmonic analysis*. Universitext. Springer, Cham, second edition, 2014.
- [7] Jean-Paul Doignon. Any finite group is the group of some binary, convex polytope. *Discrete Comput. Geom.*, 59(2):451–460, 2018.
- [8] David G. Ebin. The manifold of Riemannian metrics. In *Global Analysis (Proc. Sympos. Pure Math., Vol. XV, Berkeley, Calif., 1968)*, pages 11–40. Amer. Math. Soc., Providence, R.I., 1970.
- [9] J. M. G. Fell. A Hausdorff topology for the closed subsets of a locally compact non-Hausdorff space. *Proc. Amer. Math. Soc.*, 13:472–476, 1962.
- [10] R. Frucht. Herstellung von Graphen mit vorgegebener abstrakter Gruppe. *Compositio Math.*, 6:239–250, 1939.
- [11] Jean Haezendonck. Abstract Lebesgue-Rohlin spaces. *Bull. Soc. Math. Belg.*, 25:243–258, 1973.
- [12] Paul R. Halmos. *Lectures on ergodic theory*. Chelsea Publishing Co., New York, 1960.
- [13] Paul Richard Halmos and Viakalathur Shankar Sunder. *Bounded integral operators on L^2 spaces*, volume 96 of *Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]*. Springer-Verlag, Berlin-New York, 1978.
- [14] Hamed Hatami, László Lovász, and Balázs Szegedy. Limits of locally-globally convergent graph sequences. *Geom. Funct. Anal.*, 24(1):269–296, 2014.
- [15] Sigurdur Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 34 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001. Corrected reprint of the 1978 original.
- [16] Karl H. Hofmann and Sidney A. Morris. *The structure of compact groups—a primer for the student—a handbook for the expert*, volume 25 of *De Gruyter Studies in Mathematics*. De Gruyter, Berlin, [2020] ©2020. Fourth edition [of 1646190].
- [17] I. M. James. *Topological and uniform spaces*. Undergraduate Texts in Mathematics. Springer-Verlag, New York, 1987.
- [18] Eberhard Kaniuth and Keith F. Taylor. *Induced representations of locally compact groups*, volume 197 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2013.
- [19] Irving Kaplansky. *Infinite abelian groups*. University of Michigan Press, Ann Arbor, 1954.
- [20] Alexander S. Kechris. *Global aspects of ergodic group actions*, volume 160 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.

- [21] John M. Lee. *Introduction to smooth manifolds*, volume 218 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2013.
- [22] L. Lovász. Compact graphings. *Acta Math. Hungar.*, 161(1):185–196, 2020.
- [23] László Lovász. *Large networks and graph limits*, volume 60 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2012.
- [24] László Lovász and Balázs Szegedy. Limits of dense graph sequences. *J. Combin. Theory Ser. B*, 96(6):933–957, 2006.
- [25] László Lovász and Balázs Szegedy. Regularity partitions and the topology of graphons. In *An irregular mind*, volume 21 of *Bolyai Soc. Math. Stud.*, pages 415–446. János Bolyai Math. Soc., Budapest, 2010.
- [26] László Lovász and Balázs Szegedy. The automorphism group of a graphon. *J. Algebra*, 421:136–166, 2015.
- [27] Julien Melleray. Compact metrizable groups are isometry groups of compact metric spaces. *Proc. Amer. Math. Soc.*, 136(4):1451–1455, 2008.
- [28] Deane Montgomery and Leo Zippin. A theorem on Lie groups. *Bull. Amer. Math. Soc.*, 48:448–452, 1942.
- [29] Calvin C. Moore. Extensions and low dimensional cohomology theory of locally compact groups. I, II. *Trans. Amer. Math. Soc.*, 113:40–63; *ibid.* 113 (1964), 64–86, 1964.
- [30] James R. Munkres. *Topology*. Prentice Hall, Inc., Upper Saddle River, NJ, 2000. Second edition of [MR0464128].
- [31] Piotr Niemiec. Isometry groups of proper metric spaces. *Trans. Amer. Math. Soc.*, 366(5):2597–2623, 2014.
- [32] Karl Petersen. *Ergodic theory*, volume 2 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1989. Corrected reprint of the 1983 original.
- [33] Gert Sabidussi. Graphs with given group and given graph-theoretical properties. *Canadian J. Math.*, 9:515–525, 1957.
- [34] Gert Sabidussi. Graphs with given infinite group. *Monatsh. Math.*, 64:64–67, 1960.
- [35] Egon Schulte, Pablo Soberón, and Gordon Ian Williams. Prescribing symmetries and automorphisms for polytopes. In *Polytopes and discrete geometry*, volume 764 of *Contemp. Math.*, pages 221–233. Amer. Math. Soc., [Providence], RI, [2021] ©2021.
- [36] Egon Schulte and Gordon Ian Williams. Polytopes with preassigned automorphism groups. *Discrete Comput. Geom.*, 54(2):444–458, 2015.
- [37] Jean-Pierre Serre. *Topics in Galois theory*, volume 1 of *Research Notes in Mathematics*. A K Peters, Ltd., Wellesley, MA, second edition, 2008. With notes by Henri Darmon.
- [38] Barry Simon. *Representations of finite and compact groups*, volume 10 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1996.

- [39] Balázs Szegedy. Limits of kernel operators and the spectral regularity lemma. *European J. Combin.*, 32(7):1156–1167, 2011.
- [40] Balázs Szegedy. On higher order fourier analysis, 2012. <http://arxiv.org/abs/1203.2260v1>.
- [41] Terence Tao. *Hilbert’s fifth problem and related topics*, volume 153 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2014.
- [42] A. M. Vershik and U. Haböck. Compactness of the congruence group of measurable functions in several variables. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 334(Chisl. Metody i Vopr. Organ. Vychisl. 19):57–67, 267–268, 2006.
- [43] James Wiegold. $\text{Ext}(Q, Z)$ is the additive group of real numbers. *Bull. Austral. Math. Soc.*, 1:341–343, 1969.

DEPARTMENT OF MATHEMATICS, UNIVERSITY AT BUFFALO, BUFFALO, NY 14260-2900, USA
E-mail address: achirvas@buffalo.edu