

Graph Similarity Based on Matrix Norms

Timo Gervens  

RWTH Aachen University, Germany

Martin Grohe  

RWTH Aachen University, Germany

Abstract

Quantifying the similarity between two graphs is a fundamental algorithmic problem at the heart of many data analysis tasks for graph-based data. In this paper, we study the computational complexity of a family of similarity measures based on quantifying the mismatch between the two graphs, that is, the “symmetric difference” of the graphs under an optimal alignment of the vertices. An important example is similarity based on graph edit distance. While edit distance calculates the “global” mismatch, that is, the number of edges in the symmetric difference, our main focus is on “local” measures calculating the maximum mismatch per vertex.

Mathematically, our similarity measures are best expressed in terms of the adjacency matrices: the mismatch between graphs is expressed as the difference of their adjacency matrices (under an optimal alignment), and we measure it by applying some matrix norm. Roughly speaking, global measures like graph edit distance correspond to entrywise matrix norms like the Frobenius norm and local measures correspond to operator norms like the spectral norm.

We prove a number of strong NP-hardness and inapproximability results even for very restricted graph classes such as bounded-degree trees.

2012 ACM Subject Classification Mathematics of computing → Graph theory; Theory of computation → Complexity theory and logic

Keywords and phrases graph similarity, approximate graph isomorphism, graph matching

Funding *Timo Gervens*: This work was supported by the DFG grant GR 1492/20-1.

1 Introduction

Graphs are basic models ubiquitous in science and engineering. They are used to describe a diverse range of objects and processes from chemical compounds to social interactions. To understand and classify graph models, we need to compare graphs. Since data and models are not always guaranteed to be exact, it is essential to understand what makes two graphs similar or dissimilar, and to be able to compute similarity efficiently. There are many different approaches to similarity, for example, based on edit-distance (e.g. [3, 6, 28]), spectral similarity (e.g. [17, 30, 31]), optimal transport (e.g. [5, 24, 26]), or behavioral equivalence (e.g. [29, 32]). This is only natural, because the choice of a “good” similarity measure will usually depend on the application. While graph similarity has received considerable attention in application areas such as computer vision (see, for example, [7, 9]) and network science (see, for example, [8]), theoretical computer scientists have not explored similarity systematically; only specific “special cases” such as isomorphism [13] and bisimilarity [29] have been studied to great depth. Yet it seems worthwhile to develop a *theory of graph similarity* that compares different similarity measures, determines their algorithmic and semantic properties, and thus gives us a better understanding of their suitability for various kinds of applications. We see our paper as one contribution to such a theory.

Maybe the simplest graph similarity measure is based on graph edit distance: the *edit distance* between graphs G, H (for simplicity of the same order) is the minimum number of edges that need to be added and deleted from G to obtain a graph isomorphic to H .

In this paper, we study the computational complexity of a natural class of similarity

measures that generalize the similarity derived from edit distance of graphs. In general, we view similarity as proximity with respect to some metric. A common way of converting a graph metric d into a similarity measure s is to let $s(G, H) := \exp(-\beta \cdot d(G, H))$ for some constant $\beta > 0$. For our considerations the transformation between distance and similarity is irrelevant, so we focus directly on the metrics. In terms of computational complexity, computing similarity tends to be a hard algorithmic problem. It is known that computing edit distance, exactly or approximately, is NP-hard [3, 12, 19, 28] even on very restricted graph classes. In fact, the problem is closely related to the quadratic assignment problem [23, 27], which is notorious for being a very hard combinatorial optimization problem also for practical, heuristic approaches. Within the spectrum of similarities, the “limit” case of graph isomorphism shows that overall the complexity of graph similarity is far from trivial.

The metrics we study in this paper are based on minimizing the mismatch between two graphs. For graphs G, H with the same vertex set V , we define their *mismatch graph* $G - H$ to be the graph with vertex set V and edge set $E(G) \Delta E(H)$, the symmetric difference between the edge sets of the graphs. We assign *signs* to edges of the mismatch graph to indicate which graph they come from, say, a positive sign to the edges in $E(G) \setminus E(H)$ and a negative sign to the edges in $E(H) \setminus E(G)$. To quantify the mismatch between our graphs we introduce a *mismatch norm* μ on signed graphs that satisfies a few basic axioms such as subadditivity as well as invariance under permutations and under flipping the signs of all edges. Now for graphs G, H , not necessarily with the same vertex set, but for simplicity of the same order,¹ we define the distance $\text{dist}_\mu(G, H)$ to be the minimum of $\mu(G^\pi - H)$, where π ranges over all bijective mappings from $V(G)$ to $V(H)$ and G^π is the image of G under π .

The simplest mismatch norm μ_{ed} just counts the number of edges of the mismatch graph $G - H$, ignoring their signs. Then the associated distance $\text{dist}_{\text{ed}}(G, H)$ is the edit distance between G and H . (Note that we write dist_{ed} instead of the clunky $\text{dist}_{\mu_{\text{ed}}}$; we will do the same for other mismatch norms discussed here.) Another simple yet interesting mismatch norm is μ_{deg} measuring the maximum degree of the mismatch graph, again ignoring the signs of the edges. Then $\text{dist}_{\text{deg}}(G, H)$ measures how well we can align the two graphs in order to minimize the “local mismatch” at every node. Hence an alignment where at every vertex there is a mismatch of one edge yields a smaller dist_{deg} than an alignment that is perfect at all nodes except one, where it has a mismatch of, say, $n/2$, where n is the number of vertices. For edit distance it is the other way round. Depending on the application, one or the other may be preferable. Another well-known graph metric that can be described via the mismatch graph is Lovász’s *cut distance* (see [20, Chapter 8] and Section 3 of this paper). And, last but not least, for the mismatch norm μ_{iso} defined to be 0 if the mismatch graph has no edges and 1 otherwise, $\text{dist}_{\text{iso}}(G, H)$ is 0 if G and H are isomorphic, and 1 otherwise, so computing the distance between two graphs amounts to deciding if they are isomorphic.

Mathematically, the framework of mismatch norms and the associated distances is best described in terms of the adjacency matrices of the graphs; the adjacency matrix A_{G-H} of the mismatch graph (viewed as a matrix with entries 0, +1, -1 displaying the signs of the edges) is just the difference $A_G - A_H$. Then mismatch norms essentially are just matrix norms applied to A_{G-H} . It turns out that “global norms” such as edit distance and direct generalizations correspond to entrywise matrix norms (obtained by applying a vector norm to the flattened matrix), and “local norms” such as μ_{deg} correspond to operator matrix norms

¹ A general definition that also applies to graphs of distinct order can be found in Section 3, but for the hardness results we prove in this paper we can safely restrict our attention to graphs of the same order; this only makes the results stronger.

(see Section 3). Cut distance corresponds to the cut norm of matrices. Instead of adjacency matrices, we can also consider the Laplacian matrices, exploiting that $L_{G-H} = L_G - L_H$, and obtain another interesting family of graph distances.

For every mismatch norm μ , we are interested in the problem DIST_μ of computing $\text{dist}_\mu(G, H)$ for two given graphs G, H . Note that this is a minimization problem where the feasible solutions are the bijective mappings between the vertex sets of the two graphs. It turns out that the problem is hard for most mismatch norms μ , in particular almost all the natural choices discussed above. The single exception is μ_{iso} related to the graph isomorphism problem. Furthermore, the hardness results usually hold even if the input graphs are restricted to be very simple, for example trees or bounded degree graphs. For edit distance and the related distance based on entrywise matrix norms this was already known [3, 12]. Our focus in this paper is on operator norms. We prove a number of hardness results for different graph classes. One of the strongest ones is the following (see Theorem 19). Here DIST_p denotes the distance measure derived from the ℓ_p -operator norm.

► **Theorem 1.** *For $1 \leq p \leq \infty$ there is a constant $c > 1$ such that, unless $\text{P} = \text{NP}$, DIST_p has no factor- c approximation algorithm even if the input graphs are restricted to be trees of bounded degree.*

For details and additional results, we refer to Section 4 and 5.

Initially we aimed for a general hardness result that applies to all mismatch norms satisfying some additional natural conditions. However, we found that the hardness proofs, while following the same general strategies, usually have some intricacies exploiting special properties of the specific norms. Furthermore, for cut distance, none of these strategies seemed to work. Nevertheless, we were able to give a hardness proof for cut distance (Theorem 23) that is simply based on the hardness of computing the cut norm of the matrix [1]. This is remarkable in so far as usually the hard part of computing the distance is to find an optimal alignment π , whereas computing $\mu(G^\pi - H)$ is usually easy. For cut norm, it is even hard to compute $\mu(G^\pi - H)$ for a fixed alignment π .

1.1 Related Work

Graph similarity has mostly been studied in specific application areas, most importantly computer vision, data mining, and machine learning (see the references above). Of course not all similarity measures are based on mismatch. For example, metrics derived from vector embedding or graph kernels in machine learning (see [18]) provide a completely different approach (see [11] for a broader discussion). Of interest compared to our work (specifically for the ℓ_2 -operator norm a.k.a. spectral norm) is the spectral approach proposed by Kolla, Koutis, Madan, and Sinop [17]. Intuitively, instead of the “difference” of two graphs that is described by our mismatch graphs, their approach is based on taking a “quotient”.

The complexity of similarity, or “approximate graph isomorphism”, or “robust graph isomorphism” has been studied in [2, 3, 12, 16, 17, 19, 28], mostly based on graph edit distance and small variations. Operator norms have not been considered in this context.

2 Preliminaries

We denote the class of real numbers by \mathbb{R} and the nonnegative and positive reals by $\mathbb{R}_{\geq 0}, \mathbb{R}_{> 0}$, respectively. By $\mathbb{N}, \mathbb{N}_{> 0}$ we denote the sets of nonnegative resp. positive integers. For every $n \in \mathbb{N}_{> 0}$ we let $[n] := \{1, \dots, n\}$.

We will consider matrices with real entries and with rows and columns indexed by arbitrary finite sets. Formally, for finite sets V, W , a $V \times W$ matrix is a function $A: V \times W \rightarrow \mathbb{R}$. A standard $m \times n$ -matrix is just an $[m] \times [n]$ -matrix. We denote the set of all $V \times W$ matrices by $\mathbb{R}^{V \times W}$, and we denote the entries of a matrix A by A_{vw} .

For a square matrix $A \in \mathbb{R}^{V \times V}$ and injective mapping $\pi: V \rightarrow W$, we let A^π be the $V^\pi \times V^\pi$ -matrix with entries $A_{v^\pi w^\pi} := A_{vw}$. Note that we apply the mapping π from the right and denote the image of v under π by v^π . If $\rho: W \rightarrow X$ is another mapping, we denote the composition of π and ρ by $\pi\rho$. We typically use this notation for mappings between matrices and between graphs.

We use a standard graph theoretic notation. We denote the vertex set of a graph G by $V(G)$ and the edge set by $E(G)$. Graphs are finite, simple, and undirected, that is, $E(G) \subseteq \binom{V(G)}{2}$. We denote edges by vw instead of $\{v, w\}$. The *order* of a graph is $|G| := |V(G)|$. The *adjacency matrix* $A_G \in \{0, 1\}^{V(G) \times V(G)}$ is defined in the usual way. We denote the class of all graphs by \mathcal{G} .

Let G be a graph with vertex set $V := V(G)$. For a mapping $\pi: V \rightarrow W$ we let G^π be the graph with vertex set $\{v^\pi \mid v \in V\}$ and edge set $\{v^\pi w^\pi \mid vw \in E(G) \text{ with } v^\pi \neq w^\pi\}$. Then $A_{G^\pi} = A_G^\pi$.

For graphs G, H , we denote the set of all injective mappings $\pi: V(G) \rightarrow V(H)$ by $\text{Inj}(G, H)$. Graphs G and H are *isomorphic* (we write $G \cong H$) if there is an $\pi \in \text{Inj}(G, H)$ such that $G^\pi = H$. We think of the mappings in $\text{Inj}(G, H)$, in particular if $|G| = |H|$ and they are bijective, as *alignments* between the graphs. Intuitively, to measure the distance between two graphs, we will align them in an optimal way to minimize the mismatch.

3 Graph Metrics Based on Mismatches

A *graph metric* is a function $\delta: \mathcal{G} \times \mathcal{G} \rightarrow \mathbb{R}_{\geq 0}$ such that

(GM0) $\delta(G, H) = \delta(G', H')$ for all G, G', H, H' such that $G \cong G'$ and $H \cong H'$;

(GM1) $\delta(G, G) = 0$ for all G ;

(GM2) $\delta(G, H) = \delta(H, G)$ for all G, H ;

(GM3) $\delta(F, H) \leq \delta(F, G) + \delta(G, H)$ for all F, G, H .

Note that we do not require $\delta(G, H) > 0$ for all $G \neq H$, not even for $G \not\cong H$, so strictly speaking this is just a *pseudometric*. We are interested in the complexity of the following problem:

DIST $_\delta$

Instance: Graphs G, H , $p, q \in \mathbb{N}_{>0}$

Problem: Decide if $\delta(G, H) \geq \frac{p}{q}$

A *signed graph* is a weighted graph with edge weights $-1, +1$, and for every edge e of a signed graph we denote its *sign* by $\text{sg}(e)$. For a signed graph Δ , we let $E_+(\Delta) := \{e \in E(\Delta) \mid \text{sg}(e) = +1\}$ and $E_-(\Delta) := \{e \in E(\Delta) \mid \text{sg}(e) = -1\}$. *Isomorphisms* of signed graphs must preserve signs. A signed graph Δ is a subgraph of a signed graph Γ (we write $\Delta \subseteq \Gamma$) if $V(\Delta) \subseteq V(\Gamma)$, $E_+(\Delta) \subseteq E_+(\Gamma)$, and $E_-(\Delta) \subseteq E_-(\Gamma)$. We denote the class of all signed graphs by \mathcal{S} .

For every $\Delta \in \mathcal{S}$, we let $-\Delta$ be the signed graph obtained from Δ by flipping the signs of all edges. We define the *sum* of $\Delta, \Gamma \in \mathcal{S}$ to be the signed graph $\Delta + \Gamma$ with vertex set $V(\Delta + \Gamma) = V(\Delta) \cup V(\Gamma)$ and signed edge sets $E_+(\Delta + \Gamma) = (E_+(\Delta) \cup E_+(\Gamma)) \setminus (E_-(\Delta) \cup E_-(\Gamma))$ and

$E_-(\Delta + \Gamma) = (E_-(\Delta) \cup E_-(\Gamma)) \setminus (E_+(\Delta) \cup E_+(\Gamma))$. The adjacency matrix A_Δ of a signed graph Δ displays the signs of the edges, so $A_\Delta \in \{0, 1, -1\}^{V(\Delta) \times V(\Delta)}$ with $(A_\Delta)_{vw} = \text{sg}(vw)$ if $vw \in E(\Delta)$ and $(A_\Delta)_{vw} = 0$ otherwise. Note that $A_{-\Delta} = -A_\Delta$ for all $\Delta \in \mathcal{S}$ and $A_{\Delta+\Gamma} = A_\Delta + A_\Gamma$ for all $\Delta, \Gamma \in \mathcal{S}$ with $V(\Delta) = V(\Gamma)$ and $E_+(\Delta) \cap E_+(\Gamma) = \emptyset$ and $E_-(\Delta) \cap E_-(\Gamma) = \emptyset$.

The *mismatch graph* of two graphs G, H is the signed graph $G - H$ with vertex set $V(G - H) := V(G) \cup V(H)$ and signed edge set $E_+(G - H) := E(G) \setminus E(H)$, $E_-(G - H) := E(H) \setminus E(G)$. Note that if $V(G) = V(H)$ then for the adjacency matrices we have $A_{G-H} = A_G - A_H$. Observe that every signed graph Δ is the mismatch graph of the graphs $\Delta_+ := (V(\Delta), E_+(\Delta))$ and $\Delta_- := (V(\Delta), E_-(\Delta))$.

A *mismatch norm* is a function $\mu : \mathcal{S} \rightarrow \mathbb{R}_{\geq 0}$ satisfying the following conditions:

(MN0) $\mu(\Delta) = \mu(\Gamma)$ for all $\Delta, \Gamma \in \mathcal{S}$ such that $\Delta \cong \Gamma$;

(MN1) $\mu(\Delta) = 0$ for all $\Delta \in \mathcal{S}$ with $E(\Delta) = \emptyset$;

(MN2) $\mu(\Delta) = \mu(-\Delta)$ for all Δ ;

(MN3) $\mu(\Delta + \Gamma) \leq \mu(\Delta) + \mu(\Gamma)$ for all $\Delta, \Gamma \in \mathcal{S}$ with $V(\Delta) = V(\Gamma)$ and $E_+(\Delta) \cap E_+(\Gamma) = \emptyset$ and $E_-(\Delta) \cap E_-(\Gamma) = \emptyset$.

(MN4) $\mu(\Delta) = \mu(\Gamma)$ for all $\Delta, \Gamma \in \mathcal{S}$ with $E_+(\Delta) = E_+(\Gamma)$ and $E_-(\Delta) = E_-(\Gamma)$;

For every mismatch norm μ , we define $\text{dist}_\mu : \mathcal{G} \times \mathcal{G} \rightarrow \mathbb{R}_{\geq 0}$ by

$$\text{dist}_\mu(G, H) := \begin{cases} \min_{\pi \in \text{Inj}(G, H)} \mu(G^\pi - H) & \text{if } |G| \leq |H|, \\ \min_{\pi \in \text{Inj}(H, G)} \mu(G - H^\pi) & \text{if } |H| < |G|. \end{cases}$$

We write DIST_μ instead of $\text{DIST}_{\text{dist}_\mu}$ to denote the algorithmic problem of computing dist_μ for two graphs G, H .

► **Lemma 2.** *For every mismatch norm μ the function dist_μ is a graph metric.*

Proof. Conditions (GM0), (GM1), (GM2) follow from (MN0), (MN1), (MN2), respectively. To prove (GM3), let F, G, H be graphs. Padding the graphs with isolated vertices, by (MN4) we may assume that $|F| = |G| = |H|$. By (MN0) we may further assume that $V(F) = V(G) = V(H) := V$. Choose $\pi, \rho \in \text{Inj}(V, V)$ such that $\text{dist}_\mu(F, G) = \mu(F^\pi - G)$ and $\text{dist}_\mu(G, H) = \mu(G^\rho - H)$.

Then by (MN0) we have

$$\mu(F^{\pi\rho} - G^\rho) = \mu((F^\pi - G)^\rho) = \mu(F^\pi - G) = \text{dist}_\mu(F, G).$$

Now consider the two mismatch graphs $\Delta := F^{\pi\rho} - G^\rho$ and $\Gamma := G^\rho - H$. We have $E_+(\Delta) = E(F^{\pi\rho}) \setminus E(G^\rho)$ and $E_+(\Gamma) = E(G^\rho) \setminus E(H)$. Thus $E_+(\Delta) \cap E_+(\Gamma) = \emptyset$ and $E_-(\Delta) \cap E_-(\Gamma) = \emptyset$. Moreover, $\Delta + \Gamma = F^{\pi\rho} - H$, because

$$A_{\Delta+\Gamma} = A_\Delta + A_\Gamma = (A_F^{\pi\rho} - A_G^\rho) + (A_G^\rho - A_H) = A_F^{\pi\rho} - A_H = A_{F^{\pi\rho} - H}.$$

Thus by (MN3),

$$\text{dist}_\mu(F, H) \leq \mu(F^{\pi\rho} - H) = \mu(\Delta + \Gamma) \leq \mu(\Delta) + \mu(\Gamma) = \text{dist}_\mu(F, G) + \text{dist}_\mu(G, H).$$

This proves (GM3). ◀

► **Remark 3.** None of the five conditions (MN0)–(MN4) on a mismatch norm μ can be dropped if we want to guarantee that dist_μ is a graph metric, but of course we could replace them by other conditions. While (MN0)–(MN3) are very natural and directly correspond to

conditions (GM0)–(GM3) for graph metrics, condition (MN4) is may be less so. We chose (MN4) as the simplest condition that allows us to compare graphs of different sizes.

Having said this, it is worth noting that (MN0), (MN1) and (MN3) imply (MN4) for graphs Δ, Γ with $|\Delta| = |\Gamma|$. For graphs Δ, Γ with $|\Delta| < |\Gamma|$ they only imply $\mu(\Delta) \geq \mu(\Gamma)$. Thus as long as we only compare graphs of the same order, (MN4) is not needed. In particular, since our hardness results always apply to graphs of the same order, (MN4) is inessential for the rest of the paper.

However, it is possible to replace (MN4) by other natural conditions. For example, Lovász’s metric based on a scaled cut norm [21] does not satisfy (MN4) and instead uses a blowup of graphs to a common size to compare graphs of different sizes.

Let us now consider a few examples of mismatch norms.

► **Example 4 (Isomorphism).** The mapping $\iota := \mathcal{S} \rightarrow \mathbb{R}_{\geq 0}$ defined by $\iota(\Delta) := 0$ if $E(\Delta) = \emptyset$ and $\iota(\Delta) := 1$ otherwise is a mismatch norm. Under the metric dist_ι , all nonisomorphic graphs have distance 1 (and isomorphic graphs have distance 0, as they have under all graph metrics).

► **Example 5 (Matrix Norms).** Recall that a matrix (pseudo) norm $\|\cdot\|$ associates with every matrix A (say, with real entries) an nonnegative real $\|A\|$ in such a way that $\|N\| = 0$ for matrices N with only 0-entries, $\|aA\| = |a| \cdot \|A\|$ for all matrices A and reals $a \in \mathbb{R}$, and $\|A + B\| \leq \|A\| + \|B\|$ for all matrices A, B of the same dimensions.

Actually, we are only interested in square matrices here. We call a matrix norm $\|\cdot\|$ *permutation invariant* if for all $A \in \mathbb{R}^{V \times V}$ and all bijective $\pi : V \rightarrow V$ we have $\|A\| = \|A^\pi\|$. It is easy to see that for every permutation invariant matrix norm $\|\cdot\|$, the mapping $\mu_{\|\cdot\|} : \mathcal{S} \rightarrow \mathbb{R}_{\geq 0}$ defined by $\mu_{\|\cdot\|}(\Delta) := \|A_\Delta\|$ satisfies (MN0)–(MN3).

We call a permutation invariant matrix norm $\|\cdot\|$ *paddable* if it is invariant under extending matrices by zero entries, that is, $\|A\| = \|A'\|$ for all $A \in \mathbb{R}^{V \times V}$, $A' \in \mathbb{R}^{V' \times V'}$ such that $V' \supseteq V$, $A_{vw} = A'_{vw}$ for all $v, w \in V$, and $A'_{v'w'} = 0$ if $v \in V' \setminus V$ or $w \in V' \setminus V$. A paddable matrix norm also satisfies (MN4).

The following common matrix norms are paddable (and thus by definition also invariant). Let $1 \leq p \leq \infty$ and $A \in \mathbb{R}^{V \times V}$.

1. *Entrywise p -norm:* $\|A\|_{(p)} := \left(\sum_{v,w \in V} |A_{vw}|^p \right)^{\frac{1}{p}}$. The best-known special case is the *Frobenius norm* $\|\cdot\|_F := \|\cdot\|_{(2)}$.
2. *ℓ_p -operator norm:* $\|A\|_p := \sup_{\mathbf{x} \in \mathbb{R}^V \setminus \{\mathbf{0}\}} \frac{\|A\mathbf{x}\|_p}{\|\mathbf{x}\|_p}$, where the vector p -norm is defined by $\|\mathbf{a}\|_p := \left(\sum_{v \in V} a_v^p \right)^{\frac{1}{p}}$. In particular, $\|A\|_2$ is known as the *spectral norm*.
3. *Absolute ℓ_p -operator norm:* $\|A\|_{|p|} := \|\text{abs}(A)\|_p$, where abs takes entrywise absolute values. For the mismatch norm, taking entrywise absolute values means that we ignore the signs of the edges.
4. *Cut norm:* $\|A\|_{\square} := \max_{S, T \subseteq V} \left| \sum_{v \in S, w \in T} A_{vw} \right|$.

► **Example 6 (Laplacian Matrices).** Recall that the *Laplacian matrix* of a weighted graph G with vertex set $V := V(G)$ is the $V \times V$ matrix L_G with off-diagonal entries $(L_G)_{vw}$ being the negative weight of the edge $vw \in E(G)$ if there is such an edge and 0 otherwise and diagonal entries $(L_G)_{vv}$ being the sum of the weights of all edges incident with v . For an unweighted graph we have $L_G = D_G - A_G$, where D_G is the diagonal matrix with the vertex degrees as diagonal entries.

Observe that for a signed graph $\Delta = G - H$ we have $L_\Delta = L_G - L_H$.

It is easy to see that for every paddable matrix norm $\|\cdot\|$, the function $\mu_{\|\cdot\|}^L : \mathcal{S} \rightarrow \mathbb{R}_{\geq 0}$ defined by $\mu_{\|\cdot\|}^L(\Delta) := \|L_\Delta\|$ is a mismatch norm.

Clearly, DIST_μ is not a hard problem for every mismatch norm. For example, DIST_ν is trivial for the trivial mismatch norm ν defined by $\nu(\Delta) := 0$ for all Δ , and DIST_ι for ι from Example 4 is equivalent to the graph isomorphism problem and hence in quasipolynomial time [4].

However, for most natural matrix norms the associated graph distance problem is NP-hard. In particular, for every $p \in \mathbb{R}_{>0}$, this holds for the metric $\text{dist}_{(p)}$ based on the entrywise p -norm $\|\cdot\|_{(p)}$.

► **Theorem 7** ([12]). *For $p \in \mathbb{R}_{>0}$, $\text{DIST}_{(p)}$ is NP-hard even if restricted to trees or bounded-degree graphs.*

The proof in [12] is only given for the Frobenius norm, that is, $\text{DIST}_{(2)}$, but it actually applies to all p . In the rest of the paper, we study the complexity of DIST_p , $\text{DIST}_{|p|}$ and DIST_\square .

4 Complexity for Operator Norms

In this section we investigate the complexity of DIST_p and $\text{DIST}_{|p|}$ for $1 \leq p \leq \infty$. However, as $\mu_{|p|}$ would only be a special case within the upcoming proofs, we omit to mention it explicitly. We also omit to specify the possible values for p .

Given graphs G and H , an alignment from G to H , and a node $v \in V(G)$ we refer to the nodes whose adjacency is not preserved by π as the π -mismatches of v . If π is clear from the context we might omit it. We call $G^\pi - H$ the *mismatch graph* of π .

For all ℓ_p -operator norms the value of $\mu_p(G^\pi - H)$ strongly depends on the maximum degree in the mismatch graph of π . We capture this property with the following definition.

► **Definition 8.** *Let G, H be graphs of the same order and π an alignment from G to H . The π -mismatch count (π -MC) of a node $v \in V(G)$ is defined as:*

$$\text{MC}(v, \pi) := |\{w \in V(G) \mid w \text{ is a } \pi\text{-mismatch of } v\}|.$$

We use MC for nodes in H analogously. The *maximum mismatch count* (π -MMC) of π is defined as:

$$\text{MMC}(\pi) := \max_{v \in V(G)} \text{MC}(v, \pi).$$

Again we might drop the π if it is clear from the context. Note that the MMC corresponds to the maximum degree in the mismatch graph and that we use a slightly abbreviated notation in which we assume the graphs are given by the alignment.

The ℓ_1 -operator norm and the ℓ_∞ -operator norm measure exactly the maximum mismatch count. Due to the relation between the ℓ_p -operator norms we can derive an upper bound for μ_p . The proof of Lemma 9 can be found in Appendix A.1.

► **Lemma 9.** *Let G, H be graphs of the same order and π an alignment from G to H . Then*

$$\begin{aligned} \mu_1(G^\pi - H) &= \mu_\infty(G^\pi - H) = \text{MMC}(\pi), \\ \mu_p(G^\pi - H) &\leq \text{MMC}(\pi). \end{aligned}$$

Next, we observe that μ_p is fully determined by the connected component of the mismatch graph with the highest mismatch norm. The proof of Lemma 10 can be found in Appendix A.1.

► **Lemma 10.** *Let G, H be graphs of the same order, π an alignment from G to H , and \mathcal{C} the set of all connected components in $G^\pi - H$. Then*

$$\mu_p(G^\pi - H) = \max_{C \in \mathcal{C}} \mu_p(C).$$

For the sake of readability, we introduce a function as abbreviation for our upcoming bounds.

► **Definition 11.** *For all $1 \leq p \leq \infty$ we define the function bound_p as follows:*

$$\text{bound}_p(c) := \max\left(c^{1/p}, c^{1-1/p}\right).$$

In particular, $\text{bound}_2(c) = \sqrt{c}$. Now we derive our lower bound. The proof of Lemma 12 can be found in Appendix A.1.

► **Lemma 12.** *Let G, H be graphs of the same order, π an alignment from G to H , and $v \in V(G)$.*

If v has at least c mismatches, then

$$\mu_p(G^\pi - H) \geq \text{bound}_p(c).$$

If $G^\pi - H$ is a star, then

$$\mu_p(G^\pi - H) = \text{bound}_p(\text{MMC}(\pi)).$$

While μ_1 and μ_∞ simply measure the MMC, μ_2 also considers the connectedness of the mismatches around the node with the highest MC. As Lemma 9 and Lemma 12 tell us, μ_2 ranges between the $\sqrt{\text{MMC}}$ and the MMC. In fact, the lower bound is tight for stars and the upper bound is tight for regular graphs. This is intuitive as these are the extreme cases in which no mismatch can be removed/added without decreasing/increasing the MMC, respectively. Other ℓ_p -operator norms interpolate between μ_2 and μ_1 / μ_∞ in terms of how much they value the connectedness within the mismatch component of the node with the highest MC.

The lower bound for $\mu_p(G^\pi - H)$ gives us lower bound for $\text{dist}_p(G, H)$.

► **Lemma 13.** *Let G, H be graphs of the same order and π an alignment from G to H . If all alignments from G to H have a node with at least c mismatches, then*

$$\text{dist}_p(G, H) \geq \text{bound}_p(c).$$

Proof. This follows directly from the first claim of Lemma 12. ◀

The following upper bound might seem to have very restrictive conditions but is actually used in several hardness proofs.

► **Lemma 14.** *Let G, H be graphs of the same order and π an alignment from G to H . If the mismatch graph of π consists only of stars, then*

$$\text{dist}_p(G, H) \leq \mu_p(G^\pi - H) = \text{bound}_p(\text{MMC}(\pi)).$$

Proof. This follows directly from Lemma 10 and the second claim of Lemma 12. ◀

The last tool we need to prove the hardness is that we can distinguish two alignments by their MMC as long as the mismatch graph of the alignment with the lower MMC consists only of stars. The proof of Lemma 15 can be found in Appendix A.1.

► **Lemma 15.** *For all $c, d \in \mathbb{N}$ with $c < d$ it holds that $\text{bound}_p(c) < \text{bound}_p(d)$.*

The graph isomorphism problem becomes solvable in polynomial time if restricted to graphs of bounded degree [22]. In contrast to this, DIST_F is NP-hard even under this restriction [12]. We show that the same applies to DIST_p .

The reduction in the hardness proof works for any mismatch norm which can, said intuitively, distinguish the mismatch norm of the 1-regular graph of order n from any other n -nodes mismatch graph in which every node has at least one -1 edge but at least one node has an additional -1 edge and $+1$ edge. In particular, the construction also works for $\text{DIST}_{(p)}$. However, it does not work for the cut-distance, for which we independently prove the hardness in Section 6.

► **Theorem 16.** *DIST_p and $\text{DIST}_{|p|}$ are NP-hard for $1 \leq p \leq \infty$ even if both graphs have bounded degree.*

Proof. The proof is done by reduction from the NP-hard Hamiltonian Cycle problem in 3-regular graphs (HAM-CYCLE) [10]. Given a 3-regular graph G of order n as an instance of HAM-CYCLE, the reduction uses the n -nodes cycle C_n and G as inputs for DIST_p . We claim G has a Hamiltonian cycle if and only if $\text{dist}_p(C_n, G) \leq \text{bound}_p(1)$.

Assume that G has a Hamiltonian cycle. Then there exists a bijection $\pi : V(C_n) \rightarrow V(G)$ that aligns the cycle C_n perfectly with the Hamiltonian cycle in G . Each node in G has three neighbors, two of which are matched correctly by π as they are part of the Hamiltonian cycle. Hence, each node has a π -mismatch count of 1 and obviously the MMC of π is 1 as well. According to Lemma 14, we get $\text{dist}_p(C_n, G) \leq \text{bound}_p(1)$.

Conversely, assume that G has no Hamiltonian cycle. Then, for any alignment π' from C_n to G , there exists at least one edge vw in C_n that is not mapped to an edge in G . Hence, only one of the three edges incident to $\pi'(v)$ in G can be matched correctly. In total, v has at least one mismatch from C_n to G and two mismatches from G to C_n , which implies $\text{MC}(v, \pi') \geq 3$. Using Lemma 13, we get $\text{dist}_p(C_n, G) \geq \text{bound}_p(3)$. And then $\text{dist}_p(C_n, G) > \text{bound}_p(1)$ according to Lemma 15. ◀


Next, we modify the construction to get an even stronger NP-hardness result. The proof can be found in Appendix A.1.

► **Theorem 17.** *DIST_p and $\text{DIST}_{|p|}$ are NP-hard for $1 \leq p \leq \infty$ even if restricted to a path and a graph of maximum degree 3.*

Similar to bounded degree input graphs, restricting graph isomorphism to trees allows it to be solved in polynomial time [25] but DIST_F is NP-hard for trees [12]. We show that DIST_p remains NP-hard for trees even when applying the bounded degree restriction simultaneously.

► **Theorem 18.** *DIST_p and $\text{DIST}_{|p|}$ are NP-hard for $1 \leq p \leq \infty$ even if restricted to bounded-degree trees.*

Proof. The proof is done by reduction from the NP-hard THREE-PARTITION problem [10] that is defined as follows. Given the integers A and a_1, \dots, a_{3m} in unary representation, such that $\sum_{i=1}^{3m} a_i = mA$ and $A/4 < a_i < A/2$ for $1 \leq i \leq 3m$, decide whether there exists a partition of a_1, \dots, a_{3m} into m groups of size 3 such that the elements in each group sum up to exactly A . For technical reasons, we restrict the reduction to $A \geq 8$. However, the ignored cases are trivial. Precisely, for $A \in \{6, 7\}$ the answer is always YES and for $A \in [5]$ there exist no valid instances.



■ **Figure 1** Example of the construction in the proof of Theorem 18 where $m=2$, $A=10$, $a_1=a_2=4$ and $a_3=a_4=a_5=a_6=3$. Best viewed in color.

Given an instance of THREE-PARTITION with $A \geq 8$, we construct two trees T_1 and T_2 such that the given instance has answer YES, if and only if $\text{dist}_p(T_1, T_2) \leq \text{bound}_p(2)$. Figure 1 shows an example of T_1 and T_2 for $m = 2$ and $A = 10$. For illustrative reasons we assign each node a color during the construction. The colors are used in the example and we will refer to certain nodes by their color later in the proof. However, they do not restrict the possible alignments in any way.

Initialize T_1 as the disjoint union of paths p_1^1, \dots, p_{3m}^1 such that p_i^1 has a_i black nodes; initialize T_2 as the disjoint union of paths p_1^2, \dots, p_m^2 consisting of A black nodes each. In the following we refer to one endpoint of p_i^k as $e_1(p_i^k)$ and the other endpoint as $e_2(p_i^k)$. We attach three new red leaves and one new orange leaf to each black node in both T_1 and T_2 . Next we modify the graphs into trees by connecting the paths. For $1 \leq i \leq 3m - 1$ we add an edge between the orange leaf adjacent to $e_1(p_i^1)$ and the orange leaf adjacent to $e_2(p_{i+1}^1)$. For $1 \leq i \leq m - 1$ we add an edge between one of the red leaves adjacent to $e_1(p_i^2)$ and one of the red leaves adjacent to $e_2(p_{i+1}^2)$.

Next we attach two new pink leaves to each inner (non-endpoint) path node in T_1 and attach a new blue leaf to each pink node. Then we add the same number of blue nodes to T_2 and connect each blue node to one of the red nodes with degree 1. Finally, we attach a new pink leaf to each blue node. Note that both T_1 and T_2 are trees with bounded degree.

Precisely, the highest degree in T_1 is 8 and 6 in T_2 independent of the problem instance.

Intuitively, the construction ensures that every inner path node has already 2 mismatches just because of the degree difference. If there is no partition, at least one path in T_1 cannot be mapped contiguously into a path in T_2 which raises the MC of some inner path node to at least 3. Simultaneously, the construction ensures that there is an alignment for which the mismatch graph consists only of stars with maximum degree 2 if there is an alignment.

The formal continuation of this proof can be found in Appendix A.1. ◀

5 Approximability for Operator Norms

In this section we investigate the approximability of DIST_p and $\text{DIST}_{|p|}$ for $1 \leq p \leq \infty$. Again, we omit specifying the possible values for p and mentioning $\text{DIST}_{|p|}$ explicitly as the proofs work the same for it. We consider the following two possibilities to measure the error of an approximation algorithm for a minimization problem. An algorithm \mathcal{A} has *multiplicative error* $\alpha > 1$, if for any instance \mathcal{I} of the problem with an optimum $\text{OPT}(\mathcal{I})$, \mathcal{A} outputs a solution with value $\mathcal{A}(\mathcal{I})$ such that $\text{OPT}(\mathcal{I}) \leq \mathcal{A}(\mathcal{I}) \leq \alpha \text{OPT}(\mathcal{I})$. In this case we call \mathcal{A} an α -*approximation algorithm*. An algorithm \mathcal{B} has *additive error* $\varepsilon > 0$, if $\text{OPT}(\mathcal{I}) \leq \mathcal{B}(\mathcal{I}) \leq \text{OPT}(\mathcal{I}) + \varepsilon$ for any instance \mathcal{I} .

Approximating DIST_p with multiplicative error is at least as hard as the graph isomorphism problem (GI) since such an approximation algorithm \mathcal{A} could be used to decide GI considering that $\mathcal{A}(G, H) = 0$ if and only if G is isomorphic to H .

Furthermore, we can deduce thresholds under which the α -approximation is NP-hard using the gap between the decision values of the reduction in each hardness proof from Section 4.

► **Theorem 19.** *For $1 \leq p \leq \infty$ and any $\varepsilon > 0$, unless $\text{P} = \text{NP}$, there is no polynomial time approximation algorithm for DIST_p or $\text{DIST}_{|p|}$ with a multiplicative error guarantee of*

1. $\text{bound}_p(3) - \varepsilon$, even if both input graphs have bounded degree,
2. $\text{bound}_p(2) - \varepsilon$, even if one input graph is a path and the other one has bounded degree,
3. $\frac{\text{bound}_p(3)}{\text{bound}_p(2)} - \varepsilon$, even if both input graphs are trees with bounded degree.

Proof. We recall the proof of Theorem 16. If G has a Hamiltonian cycle, then $\text{dist}_p(C_n, G) \leq \text{bound}_p(1) = 1$. Otherwise $\text{dist}_p(C_n, G) \geq \text{bound}_p(3)$. Assume there is a polynomial time approximation algorithm \mathcal{A} with a multiplicative error guarantee of $\text{bound}_p(3) - \varepsilon$ for $\varepsilon > 0$. Then we can distinguish the two cases by checking whether $\mathcal{A}(C_n, G) < \text{bound}_p(3)$ and therefore decide HAM-CYCLE on 3-regular graphs in polynomial time. The same argument can be used for the other bounds using the proofs of Theorem 17 and Theorem 18, respectively. ◀

In particular, this implies that there is no polynomial time approximation scheme (PTAS) for DIST_p or $\text{DIST}_{|p|}$ under the respective restrictions.

Next we show the additive approximation hardness by scaling up the gap between the two decision values of the reduction in the proof of Theorem 16. For this we replace each edge with a colored gadget and then modify the graph so that an optimal alignment has to be color-preserving. The proof of Theorem 20 can be found in Appendix A.2.

► **Theorem 20.** *For $1 \leq p \leq \infty$ there is no polynomial time approximation algorithm for DIST_p with any constant additive error guarantee unless $\text{P} = \text{NP}$.*

However, the approximation of DIST_p becomes trivial once we restrict the input to graphs of bounded degree, although DIST_p stays NP-hard under this restriction. The proof of Theorem 21 can be found in Appendix A.2.

► **Theorem 21.** For $1 \leq p \leq \infty$, if one graph has maximum degree d , then there is a polynomial time approximation algorithm for DIST_p and $\text{DIST}_{|p|}$ with

1. a constant additive error guarantee $2d$,
2. a constant multiplicative error guarantee $1 + 2d$.

6 Complexity for Cut Norm

Finally, we show the hardness for DIST_\square which corresponds to the cut distance $\hat{\delta}_\square$ (see [20, Chapter 8]). For any signed graph G and $V \subseteq V(G)$ the induced subgraph $G[V]$ is the signed graph with vertex set V , $E_+(G[V]) = \{vw \in E_+(G) \mid v, w \in V\}$, and $E_-(G[V]) = \{vw \in E_-(G) \mid v, w \in V\}$.

► **Lemma 22.** Let Δ be a signed graph and $W \subseteq V(\Delta)$. Then $\mu_\square(\Delta) \geq \mu_\square(\Delta[W])$.

Proof. Let $V := V(\Delta)$, $A := A_\Delta$, $B := A_{\Delta[W]}$ and $S', T' := \text{argmax}_{S, T \subseteq W} \left| \sum_{v \in S, w \in T} B_{vw} \right|$. Then

$$\|\Delta[W]\|_\square = \left| \sum_{v \in S', w \in T'} B_{vw} \right| = \left| \sum_{v \in S', w \in T'} A_{vw} \right| \leq \max_{S, T \subseteq V} \left| \sum_{v \in S, w \in T} A_{vw} \right| = \|\Delta\|_\square. \quad \blacktriangleleft$$

Our hardness proof for DIST_\square is based on the hardness of computing the cut norm. With a construction similar to the one in Lemma 25 we enforce a specific alignment. The proof can be found in Appendix A.3.

► **Theorem 23.** The problem DIST_\square is NP-hard.

7 Concluding Remarks

We study the computational complexity of a class of graph metrics based on mismatch norms, or equivalently, matrix norms applied to the difference of the adjacency matrices of the input graphs under an optimal alignment between the vertex sets. We find that computing the distance between graphs under these metrics (at least for the standard, natural matrix norms) is NP-hard, often already on simple input graphs such as trees. This was essentially known for entrywise matrix norms. We prove it for operator norms and also for the cut norm.

We leave it open to find (natural) general conditions on a mismatch norm such that the corresponding distance problem becomes hard. Maybe more importantly, we leave it open to find meaningful tractable relaxations of the distance measures.

Measuring similarity via mismatch norms is only one approach. There are several other, fundamentally different ways to measure similarity. We are convinced that graph similarity deserves a systematic and general theory that compares the different approaches and studies their semantic as well as algorithmic properties. Our paper is one contribution to such a theory.

References

- 1 Noga Alon and Assaf Naor. Approximating the cut-norm via grothendieck's inequality. *SIAM J. Comput.*, 35(4):787–803, apr 2006. doi:10.1137/S0097539704441629.
- 2 S. Arora, A. Frieze, and H. Kaplan. A new rounding procedure for the assignment problem with applications to dense graph arrangement problems. *Mathematical Programming*, 92(1):1–36, 2002.

- 3 V. Arvind, J. Köbler, S. Kuhnert, and Y. Vasudev. Approximate graph isomorphism. In *Proceedings of the 37th International Symposium on Mathematical Foundations of Computer Science (MFCS 2012)*, volume 7464, pages 100–111, 2012.
- 4 László Babai. Graph isomorphism in quasipolynomial time [extended abstract]. In Daniel Wichs and Yishay Mansour, editors, *Proceedings of the 48th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2016, Cambridge, MA, USA, June 18-21, 2016*, pages 684–697. ACM, 2016. doi:10.1145/2897518.2897542.
- 5 Amélie Barbe, Marc Sebban, Paulo Gonçalves, Pierre Borgnat, and Rémi Gribonval. Graph diffusion wasserstein distances. In Frank Hutter, Kristian Kersting, Jeffrey Lijffijt, and Isabel Valera, editors, *Machine Learning and Knowledge Discovery in Databases - European Conference, ECML PKDD 2020, Ghent, Belgium, September 14-18, 2020, Proceedings, Part II*, volume 12458 of *Lecture Notes in Computer Science*, pages 577–592. Springer, 2020. URL: https://doi.org/10.1007/978-3-030-67661-2_34, doi:10.1007/978-3-030-67661-2_34.
- 6 H. Bunke. On a relation between graph edit distance and maximum common subgraph. *Pattern Recognition Letters*, 18:689–694, 1997.
- 7 D. Conte, P. Foggia, C. Sansone, and M. Vento. Thirty years of graph matching in pattern recognition. *International journal of pattern recognition and artificial intelligence*, 18(03):265–298, 05 2004.
- 8 F. Emmert-Streib, M. Dehmer, and Y. Shi. Fifty years of graph matching, network alignment and network comparison. *Information Sciences*, 346:180–197, 2016.
- 9 P. Foggia, G. Percannella, and M. Vento. Graph matching and learning in pattern recognition in the last 10 years. *International Journal of Pattern Recognition and Artificial Intelligence*, 28(1), 2014.
- 10 M. R. Garey and D. S. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. Freeman, San Francisco, 1979.
- 11 M. Grohe. word2vec, node2vec, graph2vec, x2vec: Towards a theory of vector embeddings of structured data. In D. Suciú, Y. Tao, and Z. Wei, editors, *Proceedings of the 39th ACM SIGMOD-SIGACT-SIGART Symposium on Principles of Database Systems*, pages 1–16, 2020. doi:10.1145/3375395.3387641.
- 12 M. Grohe, G. Rattan, and G. J. Woeginger. Graph similarity and approximate isomorphism. In *43rd International Symposium on Mathematical Foundations of Computer Science (MFCS 2018)*, volume 117, pages 20:1–20:16, 2018.
- 13 Martin Grohe and Pascal Schweitzer. The graph isomorphism problem. *Communications of the ACM*, 63(11):128–134, 2020. doi:10.1145/3372123.
- 14 N. J. Higham. Estimating the matrix p-norm. *Numer. Math.*, 62:511–538, 1992.
- 15 T. Kato. *Perturbation Theory for Linear Operators; 2nd ed.* Grundlehren Math. Wiss. Springer, Berlin, 1976.
- 16 P. Keldenich. Random robust graph isomorphism. Master’s thesis, Department of Computer Science, RWTH Aachen University, 2015.
- 17 A. Kolla, I. Koutis, V. Madan, and A. K. Sinop. Spectrally Robust Graph Isomorphism. In *45th International Colloquium on Automata, Languages, and Programming (ICALP 2018)*, volume 107, pages 84:1–84:13, 2018.
- 18 N.M. Kriege, F.D Johansson, and C. Morris. A survey on graph kernels. *ArXiv*, arXiv:1903.11835 [cs.LG], 2019.
- 19 C.-L. Lin. Hardness of approximating graph transformation problem. In D.-T. Du and X.-S. Zhang, editors, *Proceedings of the 5th International Symposium on Algorithms and Computation*, volume 834 of *Lecture Notes in Computer Science*, pages 74–82. Springer, 1994.
- 20 L. Lovász. *Large Networks and Graph Limits*. American Mathematical Society, 2012.
- 21 László Lovász. *Large Networks and Graph Limits*, volume 60 of *Colloquium Publications*. American Mathematical Society, 2012. URL: <http://www.ams.org/bookstore-getitem/item=COLL-60>.

- 22 E. M. Luks. Isomorphism of graphs of bounded valence can be tested in polynomial time. *Journal of Computer and System Sciences*, 25(1):42 – 65, 1982.
- 23 K. Makarychev, R. Manokaran, and M. Sviridenko. Maximum quadratic assignment problem: Reduction from maximum label cover and lp-based approximation algorithm. *ACM Transactions on Algorithms*, 10(4):18, 2014.
- 24 Hermina Petric Maretic, Mireille El Gheche, Giovanni Chierchia, and Pascal Frossard. GOT: an optimal transport framework for graph comparison. In Hanna M. Wallach, Hugo Larochelle, Alina Beygelzimer, Florence d’Alché-Buc, Emily B. Fox, and Roman Garnett, editors, *Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019, NeurIPS 2019, December 8-14, 2019, Vancouver, BC, Canada*, pages 13876–13887, 2019. URL: <https://proceedings.neurips.cc/paper/2019/hash/fdd5b16fc8134339089ef25b3cf0e588-Abstract.html>.
- 25 D. W. Matula. Subtree isomorphism in $o(n^{5/2})$. In *Algorithmic Aspects of Combinatorics*, volume 2, pages 91 – 106. 1978.
- 26 Facundo Mémoli. Gromov-wasserstein distances and the metric approach to object matching. *Found. Comput. Math.*, 11(4):417–487, 2011. doi:10.1007/s10208-011-9093-5.
- 27 V. Nagarajan and M. Sviridenko. On the maximum quadratic assignment problem. In *Proceedings of the twentieth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 516–524, 2009.
- 28 R. O’Donnell, J. Wright, C. Wu, and Y. Zhou. Hardness of robust graph isomorphism, lasserre gaps, and asymmetry of random graphs. In *Proceedings of the 25th Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1659–1677, 2014.
- 29 D. Sangiorgi. *Introduction to Bisimulation and Coinduction*. Cambridge University Press, 2012.
- 30 A. Tsitsulin, D. Mottin, P. Karras, A. Bronstein, and E. Müller. Netlsd: Hearing the shape of a graph. In Y. Guo and F. Farooq, editors, *Proceedings of the 24th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, pages 2347–2356, 2018.
- 31 S. Umeyama. An eigendecomposition approach to weighted graph matching problems. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 10(5):695–703, 1988.
- 32 R.J. van Glabbeek. The linear time - branching time spectrum I. The semantics of concrete, sequential processes. In J.A. Bergstra, A. Ponse, and S.A. Smolka, editors, *Handbook of Process Algebra*, chapter 1, pages 3–99. Elsevier, 2001.

A Appendix

A.1 Missing Proofs from Section 4

Proof of Lemma 9. By definition we have $\mu_1(G^\pi - H) = \|A_G^\pi - A_H\|_1$. It is well known that the ℓ_1 -operator norm boils down to the maximum absolute column sum:

$$\|A\|_1 = \max_{\|x\|_1=1} \|Ax\|_1 = \max_{j \in [n]} \sum_{i=1}^m |A_{ij}|.$$

The v -th column of the matrix $A_G^\pi - A_H$ has an entry with value $+1$ or -1 for each mismatch of the node v and all its other entries are 0. Therefore, the MC of v is equal to the absolute sum of all entries in the v -th column. It follows that the MMC is equal to the maximum absolute column sum. Similarly, the ℓ_∞ -operator norm boils down to the maximum absolute row sum:

$$\|A\|_\infty = \max_{\|x\|_\infty=1} \|Ax\|_\infty = \max_{i \in [m]} \sum_{j=1}^n |A_{ij}|.$$

Due to the symmetry of the matrix $A_G^\pi - A_H$ we can apply the same argument for μ_∞ . Since all ℓ_p -operator norms can be estimated in terms of the ℓ_1 -operator norm and ℓ_∞ -operator norm [14; 15, page 29] by

$$\|A\|_p \leq \|A\|_1^{1/p} \cdot \|A\|_\infty^{1-1/p},$$

we get the upper bound

$$\begin{aligned} \mu_p(G^\pi - H) &= \|A_G^\pi - A_H\|_p \\ &\leq \|A_G^\pi - A_H\|_1^{1/p} \cdot \|A_G^\pi - A_H\|_\infty^{1-1/p} \\ &= \text{MMC}(\pi)^{1/p} \cdot \text{MMC}(\pi)^{1-1/p} \\ &= \text{MMC}(\pi). \end{aligned} \quad \blacktriangleleft$$

Proof of Lemma 10. Let M be the adjacency matrix of $G^\pi - H$ and \mathcal{S} be the set of submatrices of M corresponding to the connected components in \mathcal{C} . We show $\|M\|_p = \max_{S \in \mathcal{S}} \|S\|_p$ by induction over $n := |\mathcal{S}|$. For $n = 1$ we have $\mathcal{S} = \{S\}$ with $S = M$. For the induction step ($n \rightarrow n + 1$) let S_1 be the $(n + 1)$ -th submatrix of M and S_2 the smallest submatrix of M that contains the first n submatrices of M . We can permute M to the form

$$M^\tau = \begin{pmatrix} S_1 & 0 \\ 0 & S_2 \end{pmatrix}.$$

Due to the permutation invariance of ℓ_p -operator norms it holds that $\|M\|_p = \|M^\tau\|_p$. Now

16 Graph Similarity Based on Matrix Norms

let $x_1 := \operatorname{argmax}_{\|x\|_p=1} \|S_1 x\|_p$ and $x_2 := \operatorname{argmax}_{\|x\|_p=1} \|S_2 x\|_p$. Then

$$\begin{aligned}
\|M\|_p &= \max_{\|x\|_p=1} \|Mx\|_p \\
&= \max_{|\alpha_1|^p + |\alpha_2|^p = 1} \left\| \begin{pmatrix} S_1 \cdot \alpha_1 x_1 \\ S_2 \cdot \alpha_2 x_2 \end{pmatrix} \right\|_p \\
&= \max_{|\alpha_1|^p + |\alpha_2|^p = 1} \left(\|\alpha_1 S_1\|_p^p + \|\alpha_2 S_2\|_p^p \right)^{1/p} \\
&= \max_{|\alpha_1|^p + |\alpha_2|^p = 1} \left(|\alpha_1|^p \|S_1\|_p^p + |\alpha_2|^p \|S_2\|_p^p \right)^{1/p} \\
&= \max_{\alpha \in [0,1]} \left(\alpha^p \|S_1\|_p^p + (1 - \alpha^p) \|S_2\|_p^p \right)^{1/p} \\
&= \max_{\alpha \in [0,1]} \left(\alpha^p \left(\|S_1\|_p^p - \|S_2\|_p^p \right) \right)^{1/p}.
\end{aligned}$$

Now we compute α as

$$\begin{aligned}
\operatorname{argmax}_{\alpha \in [0,1]} \left(\alpha^p \left(\|S_1\|_p^p - \|S_2\|_p^p \right) \right)^{1/p} &= \operatorname{argmax}_{\alpha \in [0,1]} \left(\alpha^p \left(\|S_1\|_p^p - \|S_2\|_p^p \right) \right) \\
&= \operatorname{argmax}_{\alpha \in [0,1]} \left(\alpha \left(\|S_1\|_p^p - \|S_2\|_p^p \right) \right) \\
&= \begin{cases} 1, & \text{if } \|S_1\|_p^p \geq \|S_2\|_p^p, \\ 0, & \text{else,} \end{cases} \\
&= \begin{cases} 1, & \text{if } \|S_1\|_p \geq \|S_2\|_p, \\ 0, & \text{else.} \end{cases}
\end{aligned}$$

Hence, $\|M\|_p = \max \left(\|S_1\|_p, \|S_2\|_p \right)$, which concludes the induction step. \blacktriangleleft

Proof of Lemma 12. Let M be the adjacency matrix of $G^\pi - H$. First we show $\|M\|_p \geq c^{1/p}$. Let $x \in \mathbb{R}^n$ with $x_v = 1$ and $x_i = 0$ for $i \neq v$, then:

$$\|M\|_p \geq \|Mx\|_p = \|M_{*v}\|_p = \left(|M_{1v}|^p + \dots + |M_{nv}|^p \right)^{1/p} = \left(\underbrace{1^p + \dots + 1^p}_c \right)^{1/p} = c^{1/p}.$$

In the second to last step we used the fact that there are exactly c entries equal to 1 or -1 in the v -th column of M while all other entries in that column are 0. Next, we define $x' \in \mathbb{R}^n$ as

$$x'_i = \begin{cases} c^{-1/p}, & \text{if } M_{vi} = 1 \text{ (} i \text{ is a mismatch of } v\text{),} \\ 0, & \text{else.} \end{cases}$$

Note that $\|x'\|_p = 1$. We use x' to show the other lower bound:

$$\begin{aligned} \|M\|_p &\geq \|Mx'\|_p = \left\| \begin{pmatrix} \sum_i M_{1i}x'_i \\ \vdots \\ \sum_i M_{ni}x'_i \end{pmatrix} \right\|_p = \left(\left| \sum_i M_{1i}x'_i \right|^p + \cdots + \left| \sum_i M_{ni}x'_i \right|^p \right)^{1/p} \\ &\geq \left(\left| \sum_i M_{vi}x'_i \right|^p \right)^{1/p} = \left| \sum_i M_{vi}x'_i \right| = \left| \sum_{\substack{i \text{ with} \\ |M_{vi}|=1}} M_{vi}c^{-1/p} \right| \\ &= \left| c^{-1/p} \sum_{\substack{i \text{ with} \\ |M_{vi}|=1}} M_{vi} \right| = \left| c^{-1/p} \sum_{\substack{i \text{ with} \\ |M_{vi}|=1}} 1 \right| = |c^{-1/p}c| = c^{1-1/p}. \end{aligned}$$

Now, we assume $G^\pi - H$ is a star and show $\mu_p(G^\pi - H) = \text{bound}_p(\text{MMC}(\pi))$. M has exactly c entries with value 1 or -1 in the column and the row corresponding to v , which represent the c mismatches; all other entries are 0. Hence, we can permute M to the form

$$M^\tau = \begin{pmatrix} 0 & 1 & \dots & 1 & -1 & \dots & -1 \\ 1 & & & & & & \\ \vdots & & & & & & \\ 1 & & & & & & \\ -1 & & & 0 & & & \\ \vdots & & & & & & \\ -1 & & & & & & \end{pmatrix} \in \mathbb{R}^{(c+1) \times (c+1)}.$$

Due to the permutation invariance of ℓ_p -operator norms it holds that $\|M\|_p = \|M^\tau\|_p$. Using the same argument as in the proof of Lemma 10 we get

$$\|M\|_p = \max \left(\|V^T\|_p, \|V\|_p \right), \quad V := (M_{1,2}, \dots, M_{1,c+1}) = (1, \dots, 1, -1, \dots, -1) \in \mathbb{R}^{1 \times c}.$$

The computation of the first maximization term is straightforward because x is one-dimensional:

$$\underbrace{\|V^T\|_p}_{\text{matrix norm}} = \max_{\|x\|_p=1} \underbrace{\|V^T x\|_p}_{\text{vector norm}} = \underbrace{\|V^T(1)\|_p}_{\text{vector norm}} = \underbrace{\|V^T\|_p}_{\text{vector norm}} = \underbrace{(1^p + \dots + 1^p)}_c^{1/p} = c^{1/p}.$$

The optimal solution for the other maximization term is exactly x' :

$$\begin{aligned} \operatorname{argmax}_{\|x\|_p=1} \|Vx\|_p &= \operatorname{argmax}_{\|x\|_p=1} \sqrt[p]{(x_1 + \dots + x_c)^p} \\ &= \operatorname{argmax}_{\|x\|_p=1} x_1 + \dots + x_c \\ &= \operatorname{argmax}_{|x_1|^p + \dots + |x_c|^p = 1} x_1 + \dots + x_c \\ &= x'. \end{aligned}$$

Therefore, we know that $\|V\|_p = \|Vx'\|_p = c^{1-1/p}$. ◀

Proof of Lemma 15. If $p \leq 2$ then

$$\text{bound}_p(c) = \max \left(c^{1-1/p}, c^{1/p} \right) = c^{1/p} < d^{1/p} = \max \left(d^{1-1/p}, d^{1/p} \right) = \text{bound}_p(d).$$

Otherwise

$$\text{bound}_p(c) = \max\left(c^{1-1/p}, c^{1/p}\right) = c^{1-1/p} < d^{1-1/p} = \max\left(d^{1-1/p}, d^{1/p}\right) = \text{bound}_p(d). \blacktriangleleft$$

Proof of Theorem 17. We modify the proof of Theorem 16 so that the comparison graph is a path instead of the cycle C_n . Given a 3-regular graph G of order n as an instance of HAM-CYCLE, let P_{n+2} be the path of length $n+2$ and v be a node randomly chosen from G . We remove one of the edges incident to v and attach a new leaf to each of the two separated nodes. Now we create another copy of G with the same modification applied to a different edge incident to v and call the new graphs G_1 and G_2 . One of the two has a Hamiltonian path if and only if G has a Hamiltonian cycle.

The proof follows from the following claim: G_i has a Hamiltonian path if and only if $\text{dist}_p(P_{n+2}, G_i) \leq \text{bound}_p(1)$. Indeed, if G_i has a Hamiltonian path, the natural bijection $\pi : V(P_{n+2}) \rightarrow V(G_i)$, which aligns the path P_{n+2} with the Hamiltonian path in G_i , will cause exactly an MC of 1 on each original node (nodes from G) and an MC of 0 on the two newly inserted leaves.

Conversely, assume $\text{dist}_p(P_{n+2}, G_i) \leq \text{bound}_p(1)$. Again, this implies every node has at most one mismatch. Then, the path end nodes in P_{n+2} can only be mapped to the new nodes in G_i because the degree of the original nodes in G_i differs by 2. Therefore, inner path nodes are mapped to nodes from G . Since one of their incident edges in G is always mismatched due to the degree difference, the other two incident edges have to be matched correctly. Altogether, every edge in P_{n+2} is matched correctly, which is a contradiction to the assumption. \blacktriangleleft

Continuation of the proof of Theorem 18. We have to show that the THREE-PARTITION instance has answer YES if and only if $\text{dist}_p(T_1, T_2) \leq \text{bound}_p(2)$. First we show via contradiction that $\text{dist}_p(T_1, T_2) \geq \text{bound}_p(3)$ if the THREE-PARTITION instance has answer NO. According to Lemma 15 this implies $\text{dist}_p(T_1, T_2) > \text{bound}_p(2)$. So, we assume that the THREE-PARTITION instance has answer NO and there exists an alignment π from T_1 to T_2 such that $\mu_p(T_1^\pi - T_2) < \text{bound}_p(3)$. According to Lemma 13 the MMC of π has to be smaller than 3. In particular, nodes with a degree d can only be mapped to nodes with a degree between $d-2$ and $d+2$.

Inner path nodes in T_1 have degree 8 and can only be mapped to inner path nodes in T_2 , since there are no other nodes in T_2 that have a degree of at least $8-2=6$. Using the analog degree argument, path endpoints in T_1 can only be mapped to path nodes in T_2 . However, π needs to be a bijection and the inner path nodes are already matched, so path endpoints can only be mapped to path endpoints. All in all, we have shown that under our assumption that the MMC of π is smaller than 3, black nodes can only be mapped to black nodes.

The degree of the inner path nodes in T_1 is at least by 2 higher than the degree of any path node in T_2 . Hence, each inner path node has already two mismatches from T_1 to T_2 . If two adjacent path nodes in T_1 would be mapped to two non-adjacent path nodes in T_2 , at least one of them is an inner path node and would then have a third mismatch, so the MMC would be at least 3. Intuitively, this means the internal structure of the paths cannot be destroyed by the alignment. Formally, for each path p_i^2 in T_2 there have to be three paths $p_{i_1}^1, p_{i_2}^1, p_{i_3}^1$ in T_1 such that the subgraph of T_2 induced by $\pi(p_{i_1}^1 \cup p_{i_2}^1 \cup p_{i_3}^1)$ is equal to p_i^2 . It follows that $|p_i^2| = |p_{i_1}^1| + |p_{i_2}^1| + |p_{i_3}^1|$, so $A = a_{i_1} + a_{i_2} + a_{i_3}$ for $i \in [n]$. This corresponds exactly to the existence of a partition satisfying the requirements of THREE-PARTITION, which is a contradiction to our assumption.

Note that our argument relies on the property $A/4 < a_i < A/2$ for $1 \leq i \leq 3m$ of the THREE-PARTITION instance because otherwise more or less than three paths in T_1 might be

packed into one of the paths in T_2 . Also, instances with $a_i = 2$ would be allowed without our restriction to $A \geq 8$. In such a case the corresponding path i would not have an inner path node.

Now we assume the desired partition exists and show that we can find an alignment π with $\mu_p(T_1^\pi - T_2) \leq \text{bound}_p(2)$. Note that our choice of π will map nodes in T_1 only to nodes that have the same color in T_2 . For each group in the partition we pack the three corresponding paths in T_1 into one of the paths in T_2 . Formally, let $\{j, k, l\}$ be the i -th group and p_j^1, p_k^1, p_l^1 the corresponding paths in T_1 . Then the subgraphs of T_2 induced by $\pi(p_j^1)$, $\pi(p_k^1)$, $\pi(p_l^1)$, and $\pi(p_j^1 \cup p_k^1 \cup p_l^1)$ are paths consisting only of black nodes.

Each blue node in T_1 is mapped to any of the blue nodes in T_2 . By definition π is required to be a bijection. It is possible to find such a π because the number of blue nodes in T_1 is the same as in T_2 . Each pink node in T_1 is mapped to the pink node in T_2 that has the “same” blue neighbor. Formally, let p^1 be a pink node in T_1 and b^1 its blue neighbor. Let p^2 be the pink node adjacent to $\pi(b^1)$ in T_2 . Then $\pi(p^1) = p^2$ holds.

Each red node in T_1 is mapped to a red node in T_2 that has the “same” black neighbor. Formally, let r_1^1, r_2^1, r_3^1 be the red nodes adjacent to a path node b^1 in T_1 . Let r_1^2, r_2^2, r_3^2 be the red nodes adjacent to $\pi(b^1)$ in T_2 . Then $\pi(r_1^1) = r_1^2$, $\pi(r_2^1) = r_2^2$, and $\pi(r_3^1) = r_3^2$ holds. Each orange node in T_1 is mapped to the orange node in T_2 that has the “same” black neighbor.

According to Lemma 14, it suffices to show that the MMC of π is 2 and that the mismatched neighbors of a node with an MC of 2 have an MC of 1. The second condition is called *star condition* in the following.

Blue nodes have degree 1 in T_1 and degree 2 in T_2 . Since their pink neighbor is matched correctly by definition of π , we know that their MC is exactly 1. Pink nodes have degree 2 in T_1 and degree 1 in T_2 . Since their blue neighbor is matched correctly, their MC is exactly 1. Red nodes are mapped next to the same black neighbor. Some red nodes have an additional edge to a blue node or to another red node in T_2 . Hence, their MC can be 0 or 1. Orange nodes are mapped to orange nodes with the same black neighbor. However, some orange nodes have an additional edge to another orange node in T_1 . So their MC can be 0 or 1.

Path endpoints have no mismatches. Inner path nodes have an MC of 2. Since their mismatches, which are pink nodes, have an MC of at most 1, the star condition is fulfilled.

Using Lemma 14 we finally conclude that $\text{dist}_p(T_1, T_2) \leq \text{bound}_p(2)$ if the THREE-PARTITION instance has answer YES and therefore prove the theorem. \blacktriangleleft

A.2 Missing Proofs from Section 5

For the proof of Theorem 20 we use node coloring which we introduce in the following definition.

► Definition 24. *Let G and H be two colored graphs of order n with the respective colorings $c_G : V(G) \rightarrow \mathbb{N}$ and $c_H : V(H) \rightarrow \mathbb{N}$ such that G and H have the same number of nodes of each color. The problem COLOR-DIST_p is the task to compute the following distance function for a given ℓ_p -operator norm:*

$$\begin{aligned} \text{C-dist}_p(G, H) &:= \min_{\pi \in \Pi(G, H)} \|A_G^\pi - A_H\|_p, \\ \Pi(G, H) &:= \{\pi \in S_n \mid c_G(v) = c_H(v^\pi) \text{ for all } v \in V(G)\}. \end{aligned}$$

The following lemma allows us to color nodes if a polynomial increase of the input graph size is acceptable.

► **Lemma 25.** *Every instance (G, H) of COLOR-DIST_p can be converted to an instance (G', H') of DIST_p such that $\text{C-dist}_p(G, H) = \text{dist}_p(G', H')$ and the order of G' is polynomial in the order of G .*


Proof. First, we compute an upper bound for $\text{C-dist}_p(G, H)$. Let n denote the order of the input graphs. Obviously, the MMC of any alignment π from G to H cannot be greater than $n - 1$. According to Lemma 9 it follows that $\text{C-dist}_p(G, H) \leq \mu_p(G^\pi - H) \leq n - 1$.

Now we want to modify G and H such that any alignment, that does not preserve the coloring, has a mismatch norm of at least $n - 1$, which is at least as high as the mismatch norm of any color-preserving alignment. Then we can drop the coloring without affecting the graph distance. Let c be the number of colors that are used for G and H and let the colors be represented by the integers $1, \dots, c$. For $1 \leq i \leq c$ we attach $i \cdot n^2$ new leaves to each node in G and H that has the color i . Then we drop the coloring and denote the resulting graphs as G_2 and H_2 .

Any alignment π from G to H can be converted to an alignment τ from G_2 to H_2 with $\mu_p(G^\pi - H) = \mu_p(G_2^\tau - H_2)$ as follows. For each non-leaf node v in G_2 we set $v^\tau = v^\pi$ assuming that the original nodes of G and H are not renamed within the modification to G_2 and H_2 . Each leaf in G_2 is mapped adjacency-preserving. Precisely, let w be the neighbor of a leaf v in G_2 , then v is mapped to a leaf in H_2 that is adjacent to w^τ . Hence, there are no additional mismatches due to the leaves in G_2 and so $\mu_p(G^\pi - H) = \mu_p(G_2^\tau - H_2)$.

Now assume there exists an alignment τ from G_2 to H_2 that does not preserve the original coloring and has $\mu_p(G_2^\tau - H_2) < n - 1$. Each non-leaf node in G_2 has degree n^2 or more, while each leaf in H_2 has degree 1. Hence, if τ maps a non-leaf node to a leaf, then $\text{MMC}(\tau) \geq n^2 - 1$ and according to Lemma 13 it follows that $\mu_p(G_2^\tau - H_2) \geq \text{bound}_p(n^2 - 1)$. However, $\text{bound}_p(n^2 - 1) = \max((n^2 - 1)^{1/p}, (n^2 - 1)^{1-1/p}) \geq \sqrt{n^2 - 1} \geq n - 1$ for every ℓ_p -operator norm, which is a contradiction to our assumption. So, τ can only map leaves to leaves and non-leaves to non-leaves.

If τ does not preserve the original coloring, then it also maps some node v in G_2 to a node w in H_2 such that the color of v in G is greater than the color of w in H . This means that v has at least n^2 more leaves in G_2 than w in H_2 . We get $\text{MMC}(\tau) \geq n^2$ and following the same argument as before this is a contradiction to our assumption. All in all, we conclude that the value of $\text{dist}_p(G_2, H_2)$ is determined by an alignment that does preserve the original coloring, so $\text{dist}_p(G_2, H_2) = \text{C-dist}_p(G, H)$. Note that the order of G_2 and H_2 is polynomial in the order of G because we add at most $c \cdot n^2 \leq n^3$ leaves to each node in G . ◀



■ **Figure 2** Two black nodes connected by the gadget E from Theorem 20 with $m = 3$ and $o = 5$.

Proof of Theorem 20. We recall the proof of Theorem 16. If the given 3-regular graph G of order n has a Hamiltonian cycle, then there is an alignment with an MMC of 1 from G to the

n -nodes cycle C_n , so $\text{dist}_p(G, C_n) \leq 1$. Otherwise, any alignment has an MMC of at least 3, so $\text{dist}_p(G, C_n) \geq \text{bound}_p(3)$. We show that we can distinguish the two cases even if we can only compute dist_p up to a constant additive error ε . For this we modify G and C_n into colored graphs G' and C'_n (see Definition 24 in Appendix A.2) so that $\text{C-dist}_p(G', C'_n) + \varepsilon < \text{bound}_p(2m)$ for some m if G has a Hamiltonian cycle, and $\text{C-dist}_p(G', C'_n) \geq \text{bound}_p(2m)$ otherwise. According to Lemma 25 (see Appendix A.2), we can transform G' and C'_n into the non-colored graphs G'' and C''_n such that $\text{C-dist}_p(G', C'_n) = \text{dist}_p(G'', C''_n)$. Hence, we can decide HAM-CYCLE using an approximation algorithm for DIST_p with an additive error guarantee of ε .

First, we define the colored gadget E that is visualized in Figure 2 and used to replace edges in G and C_n as part of their modification. To construct E we create two paths p^1 and p^2 with m blue nodes each. Then, for $1 \leq i < m$ we remove the path edges $p_i^1 p_{i+1}^1$ and $p_i^2 p_{i+1}^2$, add o new red nodes to the graph, and connect each of them to p_i^1, p_{i+1}^1, p_i^2 , and p_{i+1}^2 . In total E consists of $2m$ blue nodes and $(m-1) \cdot o$ red nodes.

Now we modify the graphs G and C_n . We assign the black color to all existing nodes and replace each edge vw in G and C_n by a gadget E connecting v and w as visualized in Figure 2. Precisely, we remove the edge vw and connect each blue node in E to both v and w . In addition, we add n isolated gadgets to the graph C_n . The resulting graphs are denoted as G' and C'_n in the following. Note that both G' and C'_n contain exactly $3n$ gadgets.

We choose m so that $\text{bound}_p(2m) > \text{bound}_p(m) + \varepsilon$ and o so that $\text{bound}_p(\lceil o/2 \rceil) \geq \text{bound}_p(2m)$. Both G' and C'_n are well-defined because the function bound_p is strictly increasing according to Lemma 15. The order of G' and C'_n is polynomial in n and the following claim holds for our construction.

▷ **Claim 26.** If G has a Hamiltonian cycle then $\text{C-dist}_p(G', C'_n) \leq \text{bound}_p(m)$ and otherwise $\text{C-dist}_p(G', C'_n) \geq \text{bound}_p(2m)$.

Proof. First we assume that G has a Hamiltonian cycle. Then there is an alignment π with an MMC of 1 from G to the n -nodes cycle C_n as we showed in the proof of Theorem 16. We convert π into an alignment π' from G' to C'_n as follows. For each black node v in G' we set $v^{\pi'} := v^\pi$ assuming that the original nodes of G are not renamed within the modification to G' . Each gadget in C'_n is mapped to a gadget in G' as follows. If π maps an edge vw in G to an edge $v^\pi w^\pi$ in C_n , then π' maps the gadget that connects the nodes v and w in G' to the gadget that connects the nodes v^π and w^π in C'_n in the natural way. If π maps an edge vw in G to a non-edge $v^\pi w^\pi$ in C_n , then π' maps the gadget that connects the nodes v and w in G' to an isolated gadget in C'_n in the natural way. This is possible because the 3-regular graph G has n edges more than C_n and there are n isolated gadgets in C'_n .

Each node in G is adjacent to exactly one edge that is mapped to a non-edge. Hence, each black node in G' has exactly m blue mismatched neighbors. These blue nodes do not have any further mismatches themselves, so their MC is 1. Using Lemma 14 we conclude $\text{C-dist}_p(G', C'_n) \leq \text{bound}_p(m)$.

Now assume that G has no Hamiltonian cycle and $\text{C-dist}_p(G', C'_n) < \text{bound}_p(2m)$. Then there exists an alignment τ from G' to C'_n such that $\mu_p(G'^\tau - H') < \text{bound}_p(2m)$. First we show that τ has to preserve the structure of the gadgets. If τ maps two blue nodes v and w in G that are connected to the same o red nodes to two blue nodes v^τ and w^τ in C'_n that are not connected to the same o red nodes, then either v or w has a MC of at least $\lceil o/2 \rceil$, so $\mu_p(G'^\tau - H') \geq \text{bound}_p(\lceil o/2 \rceil) \geq \text{bound}_p(2m)$ according to Lemma 13, which is a contradiction to our assumption.

Let π be the alignment from G to C_n such that $v^\pi = v^\tau$ for all nodes v in G . Since every node in G has degree 3 and every node in C_n has degree 2, we know that one edge of each

node in G is mapped to a non-edge in C_n . This means for each node v in G' that the blue nodes of one gadget attached to v are mapped to the blue nodes of a gadget in C'_n that is not attached to v^π . Hence, every black node in G' has at already m τ -mismatches. Since G does not have a Hamiltonian cycle, there has to be one additional edge vw in G that is not mapped to an edge in C_n . This means that the gadget connecting v and w in G' cannot be connected to both v^τ and w^τ in C'_n . Hence, either v or w has a τ -MC of at least $2m$, so $\mu_p(G'^\tau - H') \geq \text{bound}_p(2m)$ according to Lemma 13, which is a contradiction to our assumption. \triangleleft

Transforming G' and C'_n into the non-colored graphs G'' and C''_n as described in Lemma 25 leads to the following results. The order of G'' and C''_n is polynomial in $n = |V(G)|$. If G does not have a Hamiltonian cycle, then $\text{dist}_p(G'', C'') = \text{C-dist}_p(G', C') \geq \text{bound}_p(2m)$. If G has a Hamiltonian cycle then $\text{dist}_p(G'', C'') = \text{C-dist}_p(G', C') \leq \text{bound}_p(m)$, so $\text{dist}_p(G'', C'') + \varepsilon \leq \text{bound}_p(m) + \varepsilon < \text{bound}_p(2m)$ using the condition for the choice of m . All in all we have shown that we can decide the NP-hard problem HAM-CYCLE on 3-regular graphs in polynomial time if we have a polynomial time approximation algorithm for DIST_p with constant additive error. \blacktriangleleft

Proof of Theorem 21. Let G and H be graphs such that H has maximum degree d . We consider the algorithm \mathcal{A} that simply returns $\|A_G\|_p + \|A_H\|_p$. First, we show that \mathcal{A} is correct using the triangle inequality:

$$\begin{aligned} \text{dist}_p(G, H) &= \min_{\pi} \|A_G^\pi - A_H\|_p \\ &\leq \min_{\pi} \|A_G^\pi\|_p + \|-A_H\|_p \\ &= \|A_G\|_p + \|A_H\|_p \\ &= \mathcal{A}(G, H). \end{aligned}$$

Now we show that \mathcal{A} has constant additive error. Choosing $\tau = \text{argmin}_{\pi} \|A_G^\pi - A_H\|$ we get

$$\begin{aligned} \mathcal{A}(G, H) &= \|A_G\|_p + \|A_H\|_p \\ &= \|A_G^\tau\|_p + \|A_H\|_p \\ &= \|A_G^\tau - A_H + A_H\|_p + \|A_H\|_p \\ &\leq \|A_G^\tau - A_H\|_p + 2\|A_H\|_p \\ &\stackrel{(*)}{\leq} \|A_G^\tau - A_H\|_p + 2d \\ &= \min_{\pi} \|A_G^\pi - A_H\|_p + 2d \\ &= \text{dist}_p(G, H) + 2d. \end{aligned}$$

Equation (*) follows from the argument in the proof of Lemma 12 applied to A_H instead of $A_G^\pi - A_H$.

The approximation algorithm \mathcal{A}' compares the maximum degree of G to d . If they are equal, \mathcal{A}' checks whether G and H are isomorphic. This can be done in polynomial time since both graphs have bounded degree [22]. In the case of isomorphy we know that $\text{dist}_p(G, H) = 0$, so \mathcal{A}' returns 0. Otherwise, \mathcal{A}' returns $\mathcal{A}(G, H)$. We compute the following

upper bound of the multiplicative error for the latter case:

$$\frac{\mathcal{A}(G, H)}{\text{dist}_p(G, H)} \leq \frac{\text{dist}_p(G, H) + 2d}{\text{dist}_p(G, H)} \quad (1)$$

$$= 1 + \frac{2d}{\text{dist}_p(G, H)} \quad (2)$$

$$\leq 1 + d. \quad (3)$$

Equation 3 follows from Lemma 13 since $\text{bound}_p(c) \geq 1$ for $c \geq 1$ and any alignment from G to H has at least one mismatch if the graphs have a different maximum degree or the isomorphism test failed. \blacktriangleleft

A.3 Missing Proofs from Section 6

Proof of Theorem 23. The proof is done by reduction from the NP-hard MAX-CUT problem on unweighted graphs [10]. First, we recall how Alan and Naor [1] construct a matrix A for any graph G so that $\|A\|_{\square} = \text{MAXCUT}(G)$. Orient G in an arbitrary manner, let $V(G) = \{v_1, v_2, \dots, v_n\}$ and $E(G) = \{e_1, e_2, \dots, e_m\}$. Then A is the $2m \times n$ matrix defined as follows. For each $1 \leq i \leq m$, if e_i is oriented from v_j to v_k , then $A_{2i-1, j} = A_{2i, k} = 1$ and $A_{2i-1, k} = A_{2i, j} = -1$. The rest of the entries in A are all 0.

Next, we observe that the matrix

$$B = \begin{pmatrix} 0 & A \\ A^T & 0 \end{pmatrix}$$

has the property $\|B\|_{\square} = 2\|A\|_{\square} = 2 \cdot \text{MAXCUT}(G)$ since the cut norm is invariant under transposition and the two submatrices A, A^T have no common rows or columns in B .

We interpret B as the adjacency matrix of a signed graph Δ' and construct the two unsigned graphs $F' := (V(\Delta), E_+(\Delta))$, $H' := (V(\Delta), E_-(\Delta))$. Then $\mu_{\square}(F' - H') = \mu_{\square}(\Delta') = \|B\|_{\square} = 2 \cdot \text{MAXCUT}(G)$. Next, we modify F', H' into the graphs F, H by adding $i \cdot \left(\left\lceil \frac{n^2}{4} \right\rceil + n\right)$ leaves to node v_i for $1 \leq i \leq n$. The reduction follows from the claim that $\text{dist}_{\square}(F, H) = 2 \cdot \text{MAXCUT}(G)$, which we prove in the following.

Let π be an alignment that maps v_i to v_i for $1 \leq i \leq n$ and each leaf in F to a leaf in H so that its adjacency is preserved; let Δ be the mismatch graph of π . Then $E(\Delta) = E(\Delta')$ and therefore $\mu_{\square}(F^{\pi} - H) = 2 \cdot \text{MAXCUT}(G)$. We conclude $\text{dist}_{\square}(F, H) \leq 2 \cdot \text{MAXCUT}(G)$.

It remains to show that no alignment can lead to a lower mismatch norm. First, let σ be any alignment that maps v_i to v_i for $1 \leq i \leq n$; let Λ be the mismatch graph of σ . Then $\Lambda[\{v_1, \dots, v_n\}] = \Delta'$ and we get $\mu_{\square}(\Lambda) \geq 2 \cdot \text{MAXCUT}(G)$ from Lemma 22.

Conversely, let ρ be any alignment that maps v_i to v_j for $i \neq j$ and $i, j \leq n$; let Γ be the mismatch graph of ρ . The number of leaves adjacent to v_i in F and to v_j in H differs at least by $\left\lceil \frac{n^2}{4} \right\rceil + n$. Without restriction we can assume there are at least $l := \left\lceil \frac{n^2}{4} \right\rceil$ leaves w_1, \dots, w_l adjacent to v_i in F that are mismatched by ρ . Let $S := \{v_i^{\rho}, w_1^{\rho}, \dots, w_l^{\rho}\}$. Then $\Gamma[S]$ has exactly l edges all of which have the same sign. It is easy to see that $\mu_{\square}(\Gamma[S]) = 2l$ which implies $\mu_{\square}(\Gamma) \geq 2l$ according to Lemma 22. We chose l so that $2l \geq \frac{n^2}{2} \geq 2 \cdot \text{MAXCUT}(G)$. After considering all alignments, we get $\text{dist}_{\square}(F, G) \geq 2 \cdot \text{MAXCUT}(G)$. This proves our claim and therefore concludes the reduction. \blacktriangleleft