

CLASSES OF F -HYPERIDEALS IN A KRASNER $F^{(m,n)}$ -HYPERRING

M. ANBARLOEI

ABSTRACT. Krasner $F^{(m,n)}$ -hyperrings were introduced and investigated by Farshi and Davvaz. In this paper, our purpose is to define and characterize three particular classes of F -hyperideals in a Krasner $F^{(m,n)}$ -hyperring, namely prime F -hyperideals, maximal F -hyperideals and primary F -hyperideals, which extend similar concepts of ring context. Furthermore, we examine the relations between these structures. Then a number of major conclusions are given to explain the general framework of these structures.

1. INTRODUCTION

As it is well known, the notion of a fuzzy set was introduced by Zadeh in 1965 [32]. After the pioneering work of Zadeh, the fuzzy sets have been used in the reconsideration of classical mathematics. A number of articles have applied fuzzy concepts to algebraic structures. Rosenfeld introduced and studied fuzzy sets in the context of group theory and formulated the notion of a fuzzy subgroup of a group [29]. The notions of fuzzy subrings and ideals were defined by Liu [21]. A considerable amount of work has been done on fuzzy ideals. In particular, many papers were written on prime fuzzy ideals.

Hyperstructure theory was born in 1934 when Marty [23], a French mathematician, defined the concept of a hypergroup as a generalization of groups. Many papers and books concerning hyperstructure theory have appeared in literature. For instance, you can see the papers [7, 8, 9, 27, 30]. The simplest algebraic hyperstructures which possess the properties of closure and associativity are called semihypergroups. n -ary semigroups and n -ary groups are algebras with one n -ary operation which is associative and invertible in a generalized sense. The idea of investigations of n -ary algebras goes back to Kasner's lecture [16] at the 53rd annual meeting of the American Association of the Advancement of Science in 1904. In 1928, Dorente wrote the first paper concerning the theory of n -ary groups [12]. Later on, Crombez and Timm [5, 6] defined the notion of the (m, n) -rings and their quotient structures. The n -ary hyperstructures have been studied in [17, 18, 19, 22, 28]. In [10], Davvaz and Vougiouklis introduced a generalization of the notion of a hypergroup in the sense of Marty and a generalization of an n -ary group, which is called n -ary hypergroup. Mirvakili and Davvaz [25] defined (m, n) -hyperrings and obtained several results in this respect. One important class of hyperrings was introduced by Krasner, where the addition is a hyperoperation, while the multiplication is an ordinary binary operation, which is called Krasner hyperring. In [24], a generalization of the Krasner hyperrings, which is a subclass of

Key words and phrases. prime F -hyperideal, maximal F -hyperideal, primary F -hyperideal, Krasner $F^{(m,n)}$ -hyperring.

(m, n) -hyperrings, was defined by Mirvakili and Davvaz. It is called Krasner (m, n) -hyperring. A Krasner (m, n) -hyperring is an algebraic hyperstructure (R, f, g) , or simply R , which satisfies the following axioms: (1) (R, f) is a canonical m -ary hypergroup; (2) (R, g) is a n -ary semigroup; (3) the n -ary operation g is distributive with respect to the m -ary hyperoperation f , i.e., for every $a_1^{i-1}, a_{i+1}^n, x_1^m \in R$, and $1 \leq i \leq n$, $g(a_1^{i-1}, f(x_1^m), a_{i+1}^n) = f(g(a_1^{i-1}, x_1, a_{i+1}^n), \dots, g(a_1^{i-1}, x_m, a_{i+1}^n))$; (4) 0 is a zero element (absorbing element) of the n -ary operation g , i.e., for every $x_2^n \in R$ we have $g(0, x_2^n) = g(x_2, 0, x_3^n) = \dots = g(x_2^n, 0) = 0$. Ameri and Norouzi in [1] introduced some important hyperideals such as Jacobson radical, n -ary prime and primary hyperideals, nilradical, and n -ary multiplicative subsets of Krasner (m, n) -hyperrings. For more study on Krasner (m, n) -hyperring refer to [2, 3, 4, 15, 24, 26, 28, 31].

The connections between hyperstructures and fuzzy sets is considered as many noticeable researches. The concept of F -polygroups was introduced and analyzed by Zahedi and Hasankhani in [34, 35]. The fuzzy hyperring notion was defined and studied in [20]. In this regards Motameni and et. al. continued the study of the notion of fuzzy hyperideals of a fuzzy hyperring. They defined and characterized prime fuzzy hyperideals and maximal fuzzy hyperideals and studied the hyperideal transfer through a fuzzy hyperring homomorphism. Zhan and et. al. in [36] concentrated on the quasi-coincidence of a fuzzy interval value with an interval valued fuzzy set. Davvaz in [27] introduced the notion of a fuzzy hyperideal of a Krasner (m, n) -hyperring and to extended the fuzzy results to Krasner (m, n) -hyperring. Let G be an arbitrary set and $L = [0, 1]$ be the unit interval. Let L^G (resp. L_*^G) be the set of all fuzzy subsets of G . An F -hyperoperation on G is a function \circ from $G \times G$ into L_*^G . If $\mu, \gamma \in L_*^G$ and $x \in G$, then $x \circ \mu = \bigcup_{a \in \text{supp}(\mu)} x \circ a$ and $\mu \circ \gamma = \bigcup_{a \in \text{supp}(\mu), b \in \text{supp}(\gamma)} a \circ b$ such that $\text{supp}(\mu) = \{a \in G \mid \mu(a) \neq 0\}$. The couple (G, \circ) is called an F -polygroup if the following conditions are satisfied: (1) $(a \circ b) \circ c = a \circ (b \circ c)$ for all $a, b, c \in G$, (2) there exists $e \in G$ with $a \in \text{supp}(a \circ e \cap e \circ a)$, for all $a \in G$, (3) for each $a \in G$, there exists a unique element $a^{-1} \in G$ with $e \in \text{supp}(a \circ a^{-1} \cap a^{-1} \circ a)$, (4) $c \in \text{supp}(a \circ b)$ implies that $a \in \text{supp}(c \circ b^{-1})$ implies that $b \in \text{supp}(a^{-1} \circ c)$, for all $a, b, c \in G$. Indeed, a fuzzy hyperoperation assigns to each pair of elements of G a non-zero fuzzy subset of G , while a hyperoperation assigns to each pair of elements of G a non-empty subset of G .

The concepts of Krasner $F^{(m, n)}$ -hyperrings and F -hyperideals were defined in [14] by Farshi and Davvaz. In this paper, we continue the study of F -hyperideals of a Krasner $F^{(m, n)}$ -hyperring, initiated in [14]. We define and analyze there particular types of F -hyperideals in a Krasner $F^{(m, n)}$ -hyperring, maximal F -hyperideals and primary F -hyperideals. We investigate the connections between them. Moreover, we introduce the concepts of F -radical, quotient Krasner $F^{(m, n)}$ -hyperring and Jacobson radical. The overall framework of these structures is then explained. It is shown (Theorem 4.6) that if \mathfrak{Q} is an primary F -hyperideal of a Krasner $F^{(m, n)}$ -hyperring (\mathfrak{A}, f, g) , then $\sqrt{\mathfrak{Q}}^F$ is a prime F -hyperideal of \mathfrak{A} .

2. PRELIMINARIES

In this section we recall some basic terms and definitions from [14] which we need to develop our paper.

A fuzzy subset of G is a function $\mu : G \rightarrow L$ such that L is the unit interval $[0, 1] \subseteq \mathbb{R}$. The set of all fuzzy subsets of G is denoted by L^G . Let $\mu, \gamma \in L^G$ and $\{\mu_\alpha \mid \alpha \in \Lambda\} \subseteq L^G$. We define the fuzzy subsets $\mu \cup \gamma$ and $\bigcup_{\alpha \in \Lambda} \mu_\alpha$ as follows:

$$(\mu \cup \gamma)(a) = \max\{\mu(a), \gamma(a)\}$$

and

$$\left(\bigcup_{\alpha \in \Lambda} \mu_\alpha\right)(a) = \bigvee_{\alpha \in \Lambda} \{\mu_\alpha(a)\}$$

For all $a \in G$. The set, $\{a \in G \mid \mu(a) \neq 0\}$ is called the support of μ and is denoted by $\text{supp}(\mu)$. When $H \subseteq G$ and $t \in L$, we define $H_t \in L^G$ as follows:

$$H_t(a) = \begin{cases} t & \text{if } a \in H, \\ 0 & \text{if } a \notin H. \end{cases}$$

In particular, if H is a singleton, say $\{x\}$, then $\{x\}_t$ is referred to as fuzzy point and, sometimes, denoted by x_t . The characteristic function of set H is denoted by χ_H . Let $L_*^G = L^G - \{0\}$. For a positive integer n , an F^n -hyperoperation on G is a mapping f from G^n to L_*^G . This means that for any $a_1, \dots, a_n \in G$, $f(a_1, \dots, a_n)$ is a non-zero fuzzy subset of G . If for all $a_1, \dots, a_n \in G$, $\text{supp}(f(a_1, \dots, a_n))$ is singleton, then f is called an F^n -operation.

Notice that the sequence a_i, a_{i+1}, \dots, a_j will be denoted by a_i^j . For $j < i$, a_i^j is the empty symbol. Using this notation,

$$f(a_1, \dots, a_i, b_{i+1}, \dots, b_j, c_{j+1}, \dots, c_n)$$

will be written as $f(a_1^i, b_{i+1}^j, c_{j+1}^n)$. The expression will be written in the form $f(a_1^i, b^{(j-i)}, c_{j+1}^n)$, when $b_{i+1} = \dots = b_j = b$.

For $\mu_1^n \in L_*^G$, we define $f(\mu_1^n)$ as follows:

$$f(\mu_1^n) = \bigcup_{a_i \in \text{supp}(\mu_i)} f(a_1^n)$$

. Let $a_1^n, a \in G$, $H \in P^*(G)$ and $\mu_1^n, \mu \in L_*^G$. Then, for $1 \leq i \leq n$

- (1) $f(a_1^{i-1}, \mu, a_{i+1}^n)$ denotes $f(\chi_{\{a_1\}}, \dots, \chi_{\{a_{i-1}\}}, \mu, \chi_{\{a_{i+1}\}}, \dots, \chi_{\{a_n\}})$,
- (2) $f(a_1^{i-1}, H, a_{i+1}^n)$ denotes $f(\chi_{\{a_1\}}, \dots, \chi_{\{a_{i-1}\}}, \chi_H, \chi_{\{a_{i+1}\}}, \dots, \chi_{\{a_n\}})$,
- (3) $f(\mu_1^{i-1}, a, \mu_{i+1}^n)$ denotes $f(\mu_1^{i-1}, \chi_{\{a\}}, \mu_{i+1}^n)$,
- (4) $f(\mu_1^{i-1}, H, \mu_{i+1}^n)$ denotes $f(\mu_1^{i-1}, \chi_H, \mu_{i+1}^n)$.

If for every $1 \leq i < j \leq n$ and all $a_1^{2n-1} \in G$,

$$f(a_1^{i-1}, f(a_i^{n+i-1}), a_{n+i}^{2n-1}) = f(a_1^{j-1}, f(a_j^{n+j-1}), a_{n+j}^{2n-1}),$$

then the F^n -hyperoperation (F^n -operation) f is called associative. G with the associative F^n -hyperoperation (F^n -operation) is called F^n -semihypergroup (F^n -semigroup).

Definition 2.1. Let (G, f) be a F^m -semihypergroup. Suppose that G is equipped with a unitary operation $^{-1} : G \rightarrow G$. The couple (G, f) is called a canonical F^m -hypergroup, if

- (1) G has an F -identity element, i.e., there exists an element $e \in G$ such that for every $a \in G$, $\text{supp}(f(a, e^{(n-1)})) = \{a\}$,
- (2) $a \in \text{supp}(f(a_1^m))$ implies $a_i \in \text{supp}(f(a_1^{-1}, \dots, a_{i-1}^{-1}, a, a_{i+1}^{-1}, \dots, a_n^{-1}))$, for all $a_1^m, a \in G$ and $1 \leq i \leq n$,
- (3) for all a_1^m and for all $\sigma \in \mathbb{S}_m$, $f(a_1^m) = f(a_{\sigma(1)}^{\sigma(m)})$.

Definition 2.2. A Krasner $F^{(m,n)}$ -hyperring is an algebraic hyperstructure (\mathfrak{A}, f, g) , or simply \mathfrak{A} , which satisfies the following axioms:

- (i) (\mathfrak{A}, f) is a canonical F^m -hypergroup,
- (ii) (\mathfrak{A}, g) is an F^n -semigroup,
- (iii) for every $x_1^{i-1}, x_{i+1}^n, a_1^m \in \mathfrak{A}$ and $1 \leq i \leq n$,

$$g(x_1^{i-1}, f(a_1^m), x_{i+1}^n) = f(g(x_1^{i-1}, a_1, x_{i+1}^n), \dots, g(x_1^{i-1}, a_m, x_{i+1}^n)),$$

- (iv) for every $a_2^n \in \mathfrak{A}$, $\text{supp}(g(e, a_2^n)) = \{e\}$ where e is the F -identity element of (\mathfrak{A}, f) .

\mathfrak{A} is called commutative Krasner $F^{(m,n)}$ -hyperring if $g(a_1^n) = g(x_{\sigma(1)}^{\sigma(n)})$, for all $a_1^n \in \mathfrak{A}$ and for every $\sigma \in \mathbb{S}_n$. In the sequel, we assume that all Krasner $F^{(m,n)}$ -hyperrings are commutative. We say that \mathfrak{A} is with scalar F -identity if there exists an element e' such that $\text{supp}(g(a, e'^{(n-1)})) = \{a\}$ for all $a \in \mathfrak{A}$.

Definition 2.3. Let (\mathfrak{A}, f, g) be a Krasner $F^{(m,n)}$ -hyperring. A non-empty subset S of \mathfrak{A} is support closed under f and g whenever for all $a_1^m, b_1^n \in S$, $\text{supp}(f(a_1^m)) \subseteq S$ and $\text{supp}(g(b_1^n)) \subseteq S$. S is called a Krasner F -subhyperring of (\mathfrak{A}, f, g) if (S, f, g) is itself a Krasner $F^{(m,n)}$ -hyperring. A Krasner F -subhyperring \mathfrak{J} of (\mathfrak{A}, f, g) is said to be an F -hyperideal if $\text{supp}(g(a_1^{i-1}, \mathfrak{J}, a_{i+1}^n)) \subseteq \mathfrak{J}$ for all $a_1^n \in \mathfrak{A}$ and $1 \leq i \leq n$.

Definition 2.4. Let $(\mathfrak{A}_1, f_1, g_1)$ and $(\mathfrak{A}_2, f_2, g_2)$ be two Krasner (m, n) -hyperrings. A mapping $h : \mathfrak{A}_1 \rightarrow \mathfrak{A}_2$ is called a homomorphism if for all $x_1^m \in \mathfrak{A}_1$ and $y_1^n \in \mathfrak{A}_1$ we have

- (1) $h(e_{\mathfrak{A}_1}) = e_{\mathfrak{A}_2}$ such that $e_{\mathfrak{A}_1}$ and $e_{\mathfrak{A}_2}$ are F -identity elements of \mathfrak{A}_1 and \mathfrak{A}_2 , respectively,
- (2) $h(\text{supp}(f_1(x_1, \dots, x_m))) = \text{supp}(f_2(h(x_1), \dots, h(x_m)))$
- (3) $h(\text{supp}(g_1(y_1, \dots, y_n))) = \text{supp}(g_2(h(y_1), \dots, h(y_n)))$.

3. PRIME F -HYPERIDEALS AND MAXIMAL F -HYPERIDEALS

We start this section by introducing the concept of prime F -hyperideals of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) .

Definition 3.1. An F -hyperideal \mathfrak{P} of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) is called a prime F -hyperideal if for all $\mu_1^n \in L_*^{\mathfrak{A}}$, $\text{supp}(g(\mu_1^n)) \subseteq \mathfrak{P}$ implies that $\text{supp}(\mu_i) \subseteq \mathfrak{P}$ for some $1 \leq i \leq n$.

Our first theorem characterizes the prime F -hyperideals of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) .

Theorem 3.2. *Let \mathfrak{P} is an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) . Then \mathfrak{P} is a prime F -hyperideal if and only if for all $a_1^n \in \mathfrak{A}$, $\text{supp}(g(a_1^n)) \subseteq \mathfrak{P}$ implies that $a_i \in \mathfrak{P}$ for some $1 \leq i \leq n$.*

Proof. \implies Let \mathfrak{P} is a prime F -hyperideal of \mathfrak{A} . Let for $a_1^n \in \mathfrak{A}$, $\text{supp}(g(a_1^n)) \subseteq \mathfrak{P}$. Then $\text{supp}(g(\chi_{\{a_i\}}, \dots, \chi_{\{a_n\}})) \subseteq \mathfrak{P}$. Since \mathfrak{P} is a prime F -hyperideal of \mathfrak{A} , then we have $\text{supp}(\chi_{\{a_i\}}) \subseteq \mathfrak{P}$ for some $1 \leq i \leq n$. This implies that $\{a_i\} \subseteq \mathfrak{P}$ and so $a_i \in \mathfrak{P}$, as needed.

\impliedby Let $\text{supp}(g(\mu_1^n)) \subseteq \mathfrak{P}$ for some $\mu_1^n \in L_*^{\mathfrak{A}}$. Let for all $1 \leq i \leq n$, $\text{supp}(\mu_i) \not\subseteq \mathfrak{P}$. Then for each $1 \leq i \leq n$, there exist an element $a_i \in \text{supp}(\mu_i)$ such that $a_i \notin \mathfrak{P}$. By Proposition 3.2 in [14], $\text{supp}(g(a_1^n)) \subseteq \text{supp}(g(\mu_1^n))$. Therefore we have

$\text{supp}(g(a_1^n)) \subseteq \mathfrak{P}$. By the hypothesis, there exists $1 \leq i \leq n$ such that $a_i \in \mathfrak{P}$ and this is a contradiction. Thus \mathfrak{P} is a prime F -hyperideal of \mathfrak{A} . \square

Example 3.3. Let $(\mathbb{Z}, +, \cdot)$ be the ring of integers and $t_1, t_2 \in (0, 1]$. We define an F^m -operation f and an F^n -operation g on \mathbb{Z} as follows:

$$\begin{aligned} f(a_1, \dots, a_m) &= (a_1 + \dots + a_m)_{t_1} \quad \text{for all } a_1, \dots, a_m \in \mathbb{Z} \\ g(a_1, \dots, a_n) &= (a_1 \cdot \dots \cdot a_n)_{t_2} \quad \text{for all } a_1, \dots, a_n \in \mathbb{Z}. \end{aligned}$$

It is easy to verify that (\mathbb{Z}, f, g) is a Krasner $F^{(m,n)}$ -hyperring. All F -hyperideals $p\mathbb{Z}$, where p is a prime natural number are prime F -hyperideals of Krasner $F^{(m,n)}$ -hyperring (\mathbb{Z}, f, g) .

Example 3.4. Consider the Krasner $F^{(m,n)}$ -hyperring (G, f, g) , given in Example 3.3 in [14]. Let $a \in G$. Then $\{e, a\}$ is a hyperideal of G but it is not a prime F -hyperideal of G .

Let \mathfrak{J} be a normal F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) . The set $[\mathfrak{A} : \mathfrak{J}^*] = \{\mathfrak{J}^*[x] \mid x \in \mathfrak{A}\}$ is a Krasner (m, n) -hyperring with m -hyperoperation $f|_{I^*}$ and n -operation $g|_{I^*}$ as follows:

$$\begin{aligned} f|_{I^*}(I^* \begin{bmatrix} x_n \\ \vdots \\ x_1 \end{bmatrix}) &= \{I^*[x]\}, \quad \forall x \in \text{supp}(f(x_1^m)) \\ g|_{I^*}(I^* \begin{bmatrix} x_n \\ \vdots \\ x_1 \end{bmatrix}) &= \{I^*[\text{supp}(g(x_1^n))]\}. \end{aligned}$$

Thus $[\mathfrak{A} : \mathfrak{J}^*]_i$ is a Krasner $F^{(m,n)}$ -hyperring (for more details refer to [14]). Now, we determine when the F -hyperideal $[\mathfrak{J} : \mathfrak{J}^*]_i$ of $[\mathfrak{A} : \mathfrak{J}^*]_i$ is prime.

Theorem 3.5. *Let \mathfrak{J} be a normal F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) and let \mathfrak{J} be an F -hyperideal of (\mathfrak{A}, f, g) . If \mathfrak{J} is a prime F -hyperideal of \mathfrak{A} , then $[\mathfrak{J} : \mathfrak{J}^*]_i$ is a prime F -hyperideal of $[\mathfrak{A} : \mathfrak{J}^*]_i$.*

Proof. By Theorem 5.10 in [14], $[\mathfrak{J} : \mathfrak{J}^*]_i$ is an F -hyperideal of $[\mathfrak{A} : \mathfrak{J}^*]_i$. Let for $I^* \begin{bmatrix} x_n \\ \vdots \\ x_1 \end{bmatrix} \in [\mathfrak{A} : \mathfrak{J}^*]_i$, $g|_{I^*}(I^* \begin{bmatrix} x_n \\ \vdots \\ x_1 \end{bmatrix}) \subseteq [\mathfrak{J} : \mathfrak{J}^*]_i$. Then $I^*[\text{supp}(g(x_1^n))] \subseteq [\mathfrak{J} : \mathfrak{J}^*]_i$. This implies that $\text{supp}(g(x_1^n)) \subseteq \mathfrak{J}$. Since \mathfrak{J} is a prime F -hyperideal of \mathfrak{A} , then there exist $1 \leq i \leq n$ such that $x_i \in \mathfrak{J}$. This means that $I^*[x_i] \in [\mathfrak{J} : \mathfrak{J}^*]_i$. Thus $[\mathfrak{J} : \mathfrak{J}^*]_i$ is a prime F -hyperideal of $[\mathfrak{A} : \mathfrak{J}^*]_i$. \square

The following result investigates the stability of prime F -hyperideals property under a transfer.

Theorem 3.6. *Let $(\mathfrak{A}_1, f_1, g_1)$ and $(\mathfrak{A}_2, f_2, g_2)$ be two Krasner $F^{(m,n)}$ -hyperrings and $h : \mathfrak{A}_1 \rightarrow \mathfrak{A}_2$ be a homomorphism. If \mathfrak{P} is a prime F -hyperideal of \mathfrak{A}_2 , then $h^{-1}(\mathfrak{P})$ is a prime F -hyperideal of \mathfrak{A}_1 .*

Proof. Suppose that $(\mathfrak{A}_1, f_1, g_1)$ and $(\mathfrak{A}_2, f_2, g_2)$ are two Krasner $F^{(m,n)}$ -hyperrings and $h : \mathfrak{A}_1 \rightarrow \mathfrak{A}_2$ is a homomorphism. Let $\text{supp}(g_2(a_1^n)) \subseteq h^{-1}(\mathfrak{P})$ for some $a_1^n \in \mathfrak{A}_1$. Then we obtain $h(\text{supp}(g_1(a_1^n))) \subseteq \mathfrak{P}$ which implies $\text{supp}(g_1(a_1^n)) \subseteq h^{-1}(\mathfrak{P})$. Since \mathfrak{P} is a prime F -hyperideal of \mathfrak{A}_2 , there exists some $1 \leq i \leq n$ such that $h(a_i) \in \mathfrak{P}$. Therefore $a_i \in h^{-1}(\mathfrak{P})$, as needed. \square

Let $(\mathfrak{A}_1, f_1, g_1)$ and $(\mathfrak{A}_2, f_2, g_2)$ be two Krasner $F^{(m,n)}$ -hyperrings and $\mathfrak{A}_1 \times \mathfrak{A}_2 = \{(a, b) \mid a \in \mathfrak{A}_1, b \in \mathfrak{A}_2\}$. Then, by Proposition 5.6 in [14], $(\mathfrak{A}_1 \times \mathfrak{A}_2, f_{\otimes}, g_{\otimes})$ is a Krasner $F^{(m,n)}$ -hyperring, where

$$f_{\otimes}((a_1, b_1), \dots, (a_m, b_m))(a, b) = \min\{f_1(a_1^m)(a), f_2(b_1^m)(b)\}$$

$$g_{\otimes}((a_1, b_1), \dots, (a_n, b_n))(a, b) = \min\{g_1(a_1^n)(a), g_2(b_1^n)(b)\}$$

Now, we establish the following result.

Theorem 3.7. *Let $(\mathfrak{R}_1, f_1, g_1)$ and $(\mathfrak{R}_2, f_2, g_2)$ be two Krasner $F^{(m,n)}$ -hyperrings. If \mathfrak{P}_1 is a prime F -hyperideal of \mathfrak{R}_1 , then $\mathfrak{P}_1 \times \mathfrak{R}_2$ is a prime F -hyperideal of $\mathfrak{R}_1 \times \mathfrak{R}_2$.*

Proof. We presume $(a_1, b_1), \dots, (a_n, b_n) \in \mathfrak{R}_1 \times \mathfrak{R}_2$ with

$$\text{supp}(g_{\otimes}((a_1, b_1), \dots, (a_n, b_n))) \subseteq \mathfrak{P}_1 \times \mathfrak{R}_2.$$

This means that

$$\{(a, b) \mid g_{\otimes}((a_1, b_1), \dots, (a_n, b_n))(a, b) \neq 0\} \subseteq \mathfrak{P}_1 \times \mathfrak{R}_2$$

and so

$$\{(a, b) \mid \min\{g_1(a_1^n)(a), g_2(b_1^n)(b)\} \neq 0\} \subseteq \mathfrak{P}_1 \times \mathfrak{R}_2.$$

This implies that $\text{supp}(g_1(a_1^n)) \subseteq \mathfrak{P}_1$. Since \mathfrak{P}_1 is a prime F -hyperideal of \mathfrak{R}_1 , we get $a_i \in \mathfrak{P}_1$ for some $1 \leq i \leq n$. Thus we have $(a_i, b_i) \in \mathfrak{P}_1 \times \mathfrak{R}_2$. Consequently, $\mathfrak{P}_1 \times \mathfrak{R}_2$ is a prime F -hyperideal of $\mathfrak{R}_1 \times \mathfrak{R}_2$. \square

Let \mathfrak{J} be a F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{R}, f, g) . Then the set

$$\mathfrak{R}/\mathfrak{J} = \{\text{supp}(f(a_1^{i-1}, \mathfrak{J}, a_{i+1}^m)) \mid a_1^{i-1}, a_{i+1}^m \in \mathfrak{R}\}$$
endowed with F^m -hyperoperation f which for all $a_{11}^{1m}, \dots, a_{m1}^{mm} \in \mathfrak{R}$

$$f(\text{supp}(f(a_{11}^{1(i-1)}, \mathfrak{J}, a_{1(i+1)}^{1m})), \dots, \text{supp}(f(a_{m1}^{m(i-1)}, \mathfrak{J}, a_{m(i+1)}^{mm}))))$$

$$= \text{supp}(f(\text{supp}(f(a_{11}^{m1}), \dots, \text{supp}(f(a_{1(i-1)}^{m(i-1)}, \mathfrak{J}, \text{supp}(f(a_{1(i+1)}^{m(i+1)})), \dots, \text{supp}(f(a_{1m}^{mm}))))$$
and with F^n -operation g which for all $a_{11}^{1m}, \dots, a_{n1}^{nm} \in \mathfrak{R}$

$$g(\text{supp}(f(a_{11}^{1(i-1)}, \mathfrak{J}, a_{1(i+1)}^{1m})), \dots, \text{supp}(f(a_{n1}^{n(i-1)}, \mathfrak{J}, a_{n(i+1)}^{nm}))))$$

$$= \text{supp}(f(\text{supp}(g(a_{11}^{n1}), \dots, \text{supp}(g(a_{1(i-1)}^{n(i-1)}, \mathfrak{J}, \text{supp}(g(a_{1(i+1)}^{n(i+1)})), \dots, \text{supp}(g(a_{1m}^{nm}))))$$
construct a Krasner $F^{(m,n)}$ -hyperring, and $(\mathfrak{R}/\mathfrak{J}, f, g)$ is called the quotient Krasner $F^{(m,n)}$ -hyperring of R by \mathfrak{J} .

Theorem 3.8. *Let \mathfrak{J} be an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{R}, f, g) . Then the natural map $\pi : \mathfrak{R} \rightarrow \mathfrak{R}/\mathfrak{J}$, by $\pi(a) = \text{supp}(f(a, \mathfrak{J}, e^{(m-2)}))$ is an epimorphism.*

Proof. Let \mathfrak{J} be an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{R}, f, g) . It is clear that π is a projection map. We have to show that π is a homomorphism.

$\pi(e_R) = \text{supp}(f(e_R, \mathfrak{J}, e^{(m-2)})) = \mathfrak{J} = e_{R/\mathfrak{J}}$, by Lemma 3.14 in [14]. For all $a_1^m \in \mathfrak{R}$,

$$\begin{aligned} \pi(\text{supp}(f(a_1^m))) &= \text{supp}(f(\text{supp}(f(a_1^m)), \mathfrak{J}, \{e\}^{(m-2)})) \\ &= \text{supp}(f(\text{supp}(f(a_1^m)), \mathfrak{J}, (\text{supp}(f(e^m))^{(m-2)}))) \\ &= \text{supp}(f(\text{supp}(f(a_1, \mathfrak{J}, e^{(m-2)})), \dots, \text{supp}(f(a_m, \mathfrak{J}, e^{(m-2)}))) \\ &= \text{supp}(f(\pi(a_1), \dots, \pi(a_m))). \end{aligned}$$

Furthermore, for all $b_1^n \in \mathfrak{R}$ we have

$$\begin{aligned} \pi(\text{supp}(g(b_1^n))) &= \text{supp}(f(\text{supp}(g(b_1^n)), \mathfrak{J}, \{e\}^{(m-2)})) \\ &= \text{supp}(f(\text{supp}(g(b_1^n)), \mathfrak{J}, (\text{supp}(g(e^n))^{(m-2)}))) \\ &= \text{supp}(g(\text{supp}(f(b_1, \mathfrak{J}, e^{(m-2)})), \dots, \text{supp}(f(b_n, \mathfrak{J}, e^{(m-2)})))) \\ &= \text{supp}(g(\pi(b_1), \dots, \pi(b_n))). \end{aligned}$$

Hence, π is a homomorphism. \square

Definition 3.9. A Krasner $F^{(m,n)}$ -hyperring (\mathfrak{R}, f, g) is called a hyperintegral F -domain, if for all $a_1^n \in \mathfrak{R}$, $\text{supp}(g(a_1^n)) = \{e\}$ implies that $a_i = e$ for some $1 \leq i \leq n$.

The next theorem characterizes prime F -hyperideals in the sense of quotient Krasner $F^{(m,n)}$ -hyperrings .

Theorem 3.10. *Let \mathfrak{P} be an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{R}, f, g) . Then \mathfrak{P} is prime if and only if $\mathfrak{R}/\mathfrak{P}$ is a hyperintegral F -domain.*

Proof. \implies Let \mathfrak{P} be a F -hyperideal of \mathfrak{R} and for all $a_{11}^{1m}, \dots, a_{n1}^{nm}$,

$$\text{supp}(f(a_{11}^{1(i-1)}, \mathfrak{P}, a_{1(i+1)}^{1m}), \dots, \text{supp}(f(a_{n1}^{n(i-1)}, \mathfrak{P}, a_{n(i+1)}^{nm})) \in \mathfrak{R}/\mathfrak{P}$$

such that

$$g(\text{supp}(f(a_{11}^{1(i-1)}, \mathfrak{J}, a_{1(i+1)}^{1m})), \dots, \text{supp}(f(a_{n1}^{n(i-1)}, \mathfrak{J}, a_{n(i+1)}^{nm}))) = \mathfrak{P} = \{e_{\mathfrak{R}/\mathfrak{J}}\}.$$

Then we get

$$\text{supp}(f(\text{supp}(g(a_{11}^{n1})), \dots, \text{supp}(g(a_{1(i-1)}^{n(i-1)})), \mathfrak{J}, \text{supp}(g(a_{1(i+1)}^{n(i+1)})), \dots, \text{supp}(g(a_{1m}^{nm})))) = \mathfrak{P}$$

and so

$$\text{supp}(f(\text{supp}(g(a_{11}^{n1})), \dots, \text{supp}(g(a_{1(i-1)}^{n(i-1)})), e, \text{supp}(g(a_{1(i+1)}^{n(i+1)})), \dots, \text{supp}(g(a_{1m}^{nm})))) \subseteq \mathfrak{P}.$$

This means that

$$g(\text{supp}(f(a_{11}^{1(i-1)}, e, a_{1(i+1)}^{1m})), \dots, \text{supp}(f(a_{n1}^{n(i-1)}, e, a_{n(i+1)}^{nm}))) \subseteq \mathfrak{P}$$

which implies

$$g(\chi_{\text{supp}(f(a_{11}^{1(i-1)}, e, a_{1(i+1)}^{1m}))}, \dots, \chi_{\text{supp}(f(a_{n1}^{n(i-1)}, e, a_{n(i+1)}^{nm}))}) \subseteq \mathfrak{P}.$$

Since \mathfrak{P} is a prime F -hyperideal of \mathfrak{R} , we obtain $\text{supp}(\chi_{\text{supp}(f(a_{j1}^{j(i-1)}, e, a_{j(i+1)}^{jm}))}) \subseteq \mathfrak{P}$

for some $1 \leq j \leq n$. Therefore $\text{supp}(f(a_{j1}^{j(i-1)}, e, a_{j(i+1)}^{jm})) \subseteq \mathfrak{P}$. Then we get $\text{supp}(f(a_{j1}^{j(i-1)}, \mathfrak{P}, a_{j(i+1)}^{jm})) = \mathfrak{P}$, by Lemma 3.14 (1) in [14]. Consequently, $\mathfrak{R}/\mathfrak{P}$ is a hyperintegral F -domain.

\Leftarrow Let $\mathfrak{R}/\mathfrak{P}$ be a hyperintegral F -domain. Suppose that $\text{supp}(g(a_1^n)) \subseteq \mathfrak{P}$ for all $a_1^n \in \mathfrak{R}$. Then we have $\text{supp}(f(\text{supp}(g(a_1^n)), \mathfrak{P}, e^{(m-2)})) = \mathfrak{P}$ by Lemma 3.14 (1) in [14]. Therefore

$$\text{supp}(f(\text{supp}(g(a_1^n)), \mathfrak{P}, (\text{supp}(g(e^n))^{(m-2)}))) = \mathfrak{P}.$$

Then by the definition of the quotient Krasner $F^{(m,n)}$ -hyperring we have

$$g(\text{supp}(f(a_1, \mathfrak{P}, e^{(m-2)})), \dots, \text{supp}(f(a_n, \mathfrak{P}, e^{(m-2)}))) = \mathfrak{P} = \{e_{\mathfrak{R}/\mathfrak{J}}\}.$$

Since $\mathfrak{R}/\mathfrak{P}$ is a hyperintegral F -domain, then we get $\text{supp}(f(a_i, \mathfrak{P}, e^{(m-2)})) = \mathfrak{P}$ for some $1 \leq i \leq n$ which implies $a_i \in \mathfrak{P}$. consequently, \mathfrak{P} is a prime F -hyperideal of \mathfrak{R} . \square

Theorem 3.11. *Let (\mathfrak{R}, f, g) be a Krasner $F^{(m,n)}$ -hyperring with the scalar identity e' . Then for all $x \in \mathfrak{R}$, the set $\{a \in \text{supp}(g(r, x, e'^{(n-2)})) \mid r \in \mathfrak{R}\}$ is an F -hyperideal of \mathfrak{R} . We say that the F -hyperideal is the hyperideal generated by x and it is denoted by $\langle x \rangle_F$.*

Proof. We first show $\langle x \rangle_F$ is support closed under f and g . Let $a_1^m \in \langle x \rangle_F$. Then for each $1 \leq i \leq m$ there exist $r_i \in \mathfrak{R}$ such that $a_i \in \text{supp}(g(r_i, x, e'^{(n-2)}))$. By Proposition 3.2 in [14], we get

$$\begin{aligned} \text{supp}(f(a_1^m)) &= \text{supp}(f(g(r_1, x, e'^{(n-2)}), \dots, g(r_m, x, e'^{(n-2)}))) \\ &= \text{supp}(g(f(r_1, \dots, r_m), x, e'^{(n-2)})) \\ &= \bigcup_{r \in \text{supp}(f(r_1, \dots, r_m))} \text{supp}(g(r, x, e'^{(n-2)})). \end{aligned}$$

Since $r \in \text{supp}(f(r_1, \dots, r_m)) \subseteq \mathfrak{R}$, then $\text{supp}(f(a_1^m)) \subseteq \langle x \rangle_F$. Let $b_1^n \in \langle x \rangle_F$. Then for each $1 \leq i \leq n$ there exist $r_i \in \mathfrak{R}$ such that $b_i \in \text{supp}(g(r_i, x, e'^{(n-2)}))$. Therefore we have

$$\begin{aligned}
\text{supp}(g(b_1^n)) &= \text{supp}(g(g(r_1, x, e'^{(n-2)}), \dots, g(r_n, x, e'^{(n-2)}))) \\
&= \text{supp}(g(g(r_1, \dots, r_n), x, e'^{(n-2)})) \\
&= \text{supp}(g(r, x, e'^{(n-2)}))
\end{aligned}$$

such that $r \in \text{supp}(g(r_1, \dots, r_n)) \subseteq \mathfrak{A}$. Hence $\text{supp}(g(b_1^n)) \subseteq \langle x \rangle_F$. It is easy to see that $(\langle x \rangle_F, f, g)$ is a F -subhyperring of \mathfrak{A} . Now we show that for all $r_1^n \in \mathfrak{A}$ and $1 \leq i \leq n$, $\text{supp}(g(r_1^{i-1}, \langle x \rangle, r_{i+1}^n)) \subseteq \langle x \rangle_F$. Let $r_1^n \in \mathfrak{A}$. Then

$$\begin{aligned}
\text{supp}(g(r_1^{i-1}, \langle x \rangle_F, r_{i+1}^n)) &= \text{supp}(g(r_1^{i-1}, \chi_{\langle x \rangle_F}, r_{i+1}^n)) \\
&= \bigcup_{a \in \text{supp}(\chi_{\langle x \rangle_F})} \text{supp}(g(r_1^{i-1}, a, r_{i+1}^n)) \\
&= \bigcup_{a \in \langle x \rangle_F} \text{supp}(g(r_1^{i-1}, a, r_{i+1}^n)) \\
&\subseteq \bigcup_{r \in \mathfrak{A}} \text{supp}(g(r_1^{i-1}, g(r, x, e'^{(n-2)}), r_{i+1}^n)) \\
&= \bigcup_{r \in \mathfrak{A}} \text{supp}(g(g(r_1^{i-1}, r, r_{i+1}^n), x, e'^{(n-2)})) \\
&= \bigcup_{r \in \mathfrak{A}} \bigcup_{r' \in \text{supp}(g(r_1^{i-1}, r, r_{i+1}^n))} \text{supp}(g(r', x, e'^{(n-2)})) \\
&\subseteq \langle x \rangle_F.
\end{aligned}$$

Thus F -subhyperring $\langle x \rangle_F = \{a \in \text{supp}(g(r, x, e'^{(n-2)})) \mid r \in \mathfrak{A}\}$ of \mathfrak{A} is an F -hyperideal of \mathfrak{A} . \square

Definition 3.12. An F -hyperideal \mathfrak{M} of a Krasner $F^{(m,n)}$ -hyperring is called maximal if for every F -hyperideal \mathfrak{N} of \mathfrak{A} , $\mathfrak{M} \subseteq \mathfrak{N} \subseteq \mathfrak{A}$ implies that $\mathfrak{N} = \mathfrak{M}$ or $\mathfrak{N} = \mathfrak{A}$.

The intersection of all maximal F -hyperideals of \mathfrak{A} is called Jacobson radical of \mathfrak{A} and it is denoted by $J(\mathfrak{A})$. If \mathfrak{A} does not have any maximal F -hyperideal, we let $J(\mathfrak{A}) = \mathfrak{A}$.

Example 3.13. Let us consider the Krasner $F^{(m,2)}$ -hyperring (G, f, g) , given in Example 3.4 in [14]. Then $\{e\}$ is a maximal F -hyperideal of G .

The following theorem ensure that there is always a sufficient supply of the maximal F -hyperideals.

Theorem 3.14. Every Krasner $F^{(m,2)}$ -hyperring \mathfrak{A} with scalar F -identity e' , has at least one maximal F -hyperideal.

Proof. The proof is the same as the proof in the classical context of maximal ideals of rings. \square

We say that an element $x \in \mathfrak{A}$ is F -invertible if there exists $y \in \mathfrak{A}$ such that $\text{supp}(g(x, y, e'^{(n-2)})) = \{e'\}$. Also, the subset U of \mathfrak{A} is F -invertible if and only if every element of U is F -invertible.

Theorem 3.15. Let \mathfrak{J} be an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring with the scalar F -identity e' . If every $a \in \mathfrak{A} - \mathfrak{M}$ is F -invertible in \mathfrak{A} , then \mathfrak{M} is the only maximal hyperideal of \mathfrak{A} .

Proof. Straight forward \square

The Jacobson radical of a Krasner $F^{(m,n)}$ -hyperring \mathfrak{A} can be characterized as follows:

Theorem 3.16. Let \mathfrak{J} be an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring with the scalar F -identity e' . Then every element of $\text{supp}(f(e', \mathfrak{J}, e'^{(m-2)}))$ is F -invertible if and only if $\mathfrak{J} \subseteq J(\mathfrak{A})$.

Proof. \implies Suppose that every element of $\text{supp}(f(e', \mathfrak{J}, e^{(n-2)}))$ is F -invertible. Let $\mathfrak{J} \not\subseteq J(\mathfrak{R})$. Then there exists a maximal F -hyperideal \mathfrak{M} of \mathfrak{R} such that $\mathfrak{J} \not\subseteq \mathfrak{M}$. Let $x \in \mathfrak{J}$ but $x \notin \mathfrak{M}$. By Lemma 3.12 in [14], $\text{supp}(f(\mathfrak{M}, \langle x \rangle_F, e^{(m-2)}))$ is an F -hyperideal of \mathfrak{R} . Since $\mathfrak{M} \subseteq \text{supp}(f(\mathfrak{M}, \langle x \rangle_F, e^{(m-2)}))$ and \mathfrak{M} is a maximal F -hyperideal of \mathfrak{R} , we have $\text{supp}(f(\mathfrak{M}, \langle x \rangle_F, e^{(m-2)})) = \mathfrak{R}$ and so $e' \in \text{supp}(f(\mathfrak{M}, \langle x \rangle_F, e^{(m-2)}))$. This means that there exist $m \in \text{supp}(\chi_{\mathfrak{M}}) = \mathfrak{M}$ and $a \in \text{supp}(\chi_{\langle x \rangle_F}) = \langle x \rangle_F$ such that $e' \in \text{supp}(f(m, a, e^{(m-2)}))$. Since (\mathfrak{R}, f) is a canonical F^m -hypergroup, then $m \in \text{supp}(f(e', -a, e^{(m-2)}))$. Since $\text{supp}(f(e', -a, e^{(m-2)})) \subseteq \text{supp}(f(e', -g(r, x, e'), e^{(m-2)}))$ for some $r \in \mathfrak{R}$, we have $m \in \text{supp}(f(e', g(r, x, e'), e^{(m-2)}))$ which implies $m \in \text{supp}(f(e', \mathfrak{J}, e^{(m-2)}))$. This m is F -invertible, a contradiction.

\impliedby Suppose that $\mathfrak{J} \subseteq J(\mathfrak{R})$. Let $a \in \text{supp}(f(e', \mathfrak{J}, e^{(m-2)}))$ is not F -invertible. Then there exists $x \in \mathfrak{J}$ such that $a \in \text{supp}(f(e', x, e^{(m-2)}))$. We have $a \in \mathfrak{M}$ for some maximal F -hyperideal \mathfrak{M} , because a is not F -invertible. From $a \in \text{supp}(f(e', x, e^{(m-2)}))$, it follows that $e' \in \text{supp}(f(a, -x, e^{(m-2)})) \subseteq \mathfrak{M}$, a contradiction. Thus every element of $\text{supp}(f(e', \mathfrak{J}, e^{(m-2)}))$ is F -invertible. \square

In view of Theorem 3.16, we have the following result.

Corollary 3.17. Let \mathfrak{M} be a maximal F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring with the scalar F -identity e' . If every element of $\text{supp}(f(e', \mathfrak{M}, e^{(m-2)}))$ is F -invertible, then \mathfrak{M} is the only maximal hyperideal of \mathfrak{R} .

Theorem 3.18. Suppose that T is a non-empty subset of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{R}, f, g) that is support closed under g and \mathfrak{J} is an F -hyperideal of \mathfrak{R} such that $\mathfrak{J} \cap T = \emptyset$. Then there exists an F -hyperideal \mathfrak{P} which is maximal in the set of all hyperideals of \mathfrak{R} disjoint from T containing \mathfrak{J} . Furthermore any such F -hyperideal is prime.

Proof. Let Σ be the set of all hyperideals of \mathfrak{R} disjoint from T containing \mathfrak{J} . Since $\mathfrak{J} \in \Sigma$, then $\Sigma \neq \emptyset$. Thus Σ is a partially ordered set with respect to set inclusion relation. Then there is an F -hyperideal \mathfrak{P} which is maximal in Σ , by Zorn's lemma. Our task now is to show that \mathfrak{P} is a prime F -hyperideal of \mathfrak{R} . Let $\text{supp}(g(a_1^n)) \subseteq \mathfrak{P}$ for some $a_1^n \in \mathfrak{R}$ such that for all $1 \leq i \leq n$, $a_i \notin \mathfrak{P}$. Then for each $1 \leq i \leq n$, $\mathfrak{P} \subseteq \text{supp}(f(\mathfrak{P}, \langle a_i \rangle, e^{(m-2)}))$. By maximality of \mathfrak{P} , we conclude that $\text{supp}(f(\mathfrak{P}, \langle a_i \rangle, e^{(m-2)})) \cap T \neq \emptyset$. Hence there exist $p_1^n \in \mathfrak{P}$ and $x_i \in \langle a_i \rangle$ such that $\text{supp}(f(p_1, x_i, e^{(m-2)})) \cap T \neq \emptyset$ for each $1 \leq i \leq n$. Since T is support closed under g , then there exist $r_i \in \text{supp}(f(p_1, x_i, e^{(m-2)}))$ for each $1 \leq i \leq n$ such that $\text{supp}(g(r_1^n)) \cap T \neq \emptyset$. Therefore

$$\begin{aligned} \text{supp}(g(r_1^n)) &\subseteq \text{supp}(g(f(p_1, x_1, e^{(m-2)}), \dots, f(p_n, x_n, e^{(m-2)}))) \\ &= \text{supp}(f(g(p_1^n), g(p_1^{n-1}, x_n), \dots, g(p_1, x_2^n), \dots, g(x_1^n), e^{(m-2^n)})) \\ &\subseteq \mathfrak{P}. \end{aligned}$$

This means that $\mathfrak{P} \cap T \neq \emptyset$ which is contradiction with $\mathfrak{P} \in \Sigma$. Thus, by Theorem 3.2, \mathfrak{P} is a prime F -hyperideal of \mathfrak{R} . \square

Definition 3.19. Let \mathfrak{J} be an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{R}, f, g) with scalar F -identity e' . The intersection of all prime F -hyperideals of \mathfrak{R} containing \mathfrak{J} is called F -radical of \mathfrak{J} , being denoted by $\sqrt{\mathfrak{J}}^F$. If \mathfrak{R} does not have any prime F -hyperideal containing \mathfrak{J} , we define $\sqrt{\mathfrak{J}}^F = \mathfrak{R}$.

The following theorem gives an alternative definition of $\sqrt{\mathfrak{J}}^F$.

Theorem 3.20. \mathfrak{J} be an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) with scalar F -identity e' . Then

$$\sqrt{\mathfrak{J}}^F = \left\{ a \in \mathfrak{A} \mid \left\{ \begin{array}{l} \text{supp}(g(a^{(s)}, e'^{(n-s)})) \subseteq \mathfrak{J}, \quad s \leq n \\ \text{supp}(g_{(l)}(a^{(s)})) \subseteq \mathfrak{J} \quad s > n, s = l(n-1) + 1 \end{array} \right\} \right\}.$$

Proof. Let $a \in \sqrt{\mathfrak{J}}^F$ and let \mathfrak{P} be a prime F -hyperideal of \mathfrak{A} with $\mathfrak{J} \subseteq \mathfrak{P}$. Thus there exists $s \in \mathbb{N}$ with $\text{supp}(g(a^{(s)}, e'^{(n-s)})) \subseteq \mathfrak{J}$ for $s \leq n$, or $\text{supp}(g_{(l)}(a^{(s)})) \subseteq \mathfrak{J}$ for $s = l(n-1) + 1$. In the first case, we have $\text{supp}(g(a^{(s)}, e'^{(n-s)})) \subseteq \mathfrak{P}$ and so $\text{supp}(g(a, g(a^{(s-1)}, e'^{(n-s+1)}), e'^{(n-2)})) \subseteq \mathfrak{P}$. Since \mathfrak{P} is a prime F -hyperideal of \mathfrak{A} , we get $a \in \mathfrak{P}$ or $\text{supp}(g(a^{(s-1)}, e'^{(n-s+1)})) \subseteq \mathfrak{P}$. From $\text{supp}(g(a^{(s-1)}, e'^{(n-s+1)})) \subseteq \mathfrak{P}$, it follows that $\text{supp}(g(a, g(a^{(s-2)}, e'^{(n-s+2)}), e'^{(n-2)})) \subseteq \mathfrak{P}$ which implies $a \in \mathfrak{P}$ or $\text{supp}(g(a^{(s-2)}, e'^{(n-s+2)})) \subseteq \mathfrak{P}$. By continuing this process, we obtain $a \in \mathfrak{P}$. Hence we have $a \in \mathfrak{P}$ for all $\mathfrak{J} \subseteq \mathfrak{P}$ and so $a \in \bigcap_{\mathfrak{J} \subseteq \mathfrak{P}} \mathfrak{P}$. This means $\sqrt{\mathfrak{J}}^F \subseteq \bigcap_{\mathfrak{J} \subseteq \mathfrak{P}} \mathfrak{P}$. In the second case, we get $\text{supp}(g(g(\dots g(a^{(n)}), a^{(n-1)}), \dots), a^{(n-1)}) \subseteq \mathfrak{P}$. By using a similar argument, we get $\sqrt{\mathfrak{J}}^F \subseteq \bigcap_{\mathfrak{J} \subseteq \mathfrak{P}} \mathfrak{P}$. Now, suppose that $a \in \bigcap_{\mathfrak{J} \subseteq \mathfrak{P}} \mathfrak{P}$ but $a \notin \sqrt{\mathfrak{J}}^F$. Hence we conclude that for every $s \in \mathbb{N}$, $\text{supp}(g(a^{(s)}, e'^{(n-s)})) \not\subseteq \mathfrak{J}$. Let $T = \{e', a\} \cup \{r \in \text{supp}(g(a^{(t)}, e'^{(n-t)})) \mid 2 \leq t\}$. Clearly, T is a subset of \mathfrak{A} that is support closed under g and $T \cap \mathfrak{J} = \emptyset$. By Theorem 3.18, there exists a prime F -hyperideal \mathfrak{P} with $\mathfrak{J} \subseteq \mathfrak{P}$ and $T \cap \mathfrak{P} = \emptyset$. This means $a \notin \mathfrak{P}$. This is contradiction as $a \in \bigcap_{\mathfrak{J} \subseteq \mathfrak{P}} \mathfrak{P}$. Thus $a \in \sqrt{\mathfrak{J}}^F$. Consequently, $\sqrt{\mathfrak{J}}^F = \bigcap_{\mathfrak{J} \subseteq \mathfrak{P}} \mathfrak{P}$. \square

4. PRIMARY F -HYPERIDEALS

In this section, we aim to present the definition of primary F -hyperideals in a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) and give some basic properties of them.

Definition 4.1. An F -hyperideal \mathfrak{Q} of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) (with scalar F -identity e') is called a primary F -hyperideal if for all $\mu_1^n \in L_*^{\mathfrak{A}}$, $\text{supp}(g(\mu_1^n)) \subseteq \mathfrak{Q}$ implies that $\text{supp}(\mu_i) \subseteq \mathfrak{Q}$ or $\text{supp}(g(\mu_1^{i-1}, \chi_{\{e'\}}, \mu_{i+1}^n)) \subseteq \sqrt{\mathfrak{Q}}^F$ for some $1 \leq i \leq n$.

Theorem 4.2. Let \mathfrak{Q} be an F -hyperideal of a Krasner $F^{(m,2)}$ -hyperring (\mathfrak{A}, f, g) (with scalar F -identity e'). Then \mathfrak{Q} is primary if and only if for all $a_1^2 \in \mathfrak{A}$, $\text{supp}(g(a_1^2)) \subseteq \mathfrak{Q}$ implies that $a_1 \in \mathfrak{Q}$ or $a_2 \in \sqrt{\mathfrak{Q}}^F$.

Proof. \implies Let \mathfrak{Q} is a primary F -hyperideal of \mathfrak{A} . Suppose that $\text{supp}(g(a_1^2)) \subseteq \mathfrak{Q}$ for some $a_1^2 \in \mathfrak{A}$. Since $\text{supp}(g(a_1^2)) = \text{supp}(g(\chi_{\{a_1\}}, \chi_{\{a_2\}}))$, then we have $\text{supp}(g(\chi_{\{a_1\}}, \chi_{\{a_2\}})) \subseteq \mathfrak{Q}$. Since \mathfrak{Q} is a primary F -hyperideal of \mathfrak{A} , we get $\text{supp}(\chi_{\{a_1\}}) \subseteq \mathfrak{Q}$ or $\text{supp}(\chi_{\{a_2\}}) \subseteq \sqrt{\mathfrak{Q}}^F$. This means $a_1 \in \mathfrak{Q}$ or $a_2 \in \sqrt{\mathfrak{Q}}^F$. \implies Let $\text{supp}(g(\mu_1^2)) \subseteq \mathfrak{Q}$ for some $\mu_1^2 \in L_*^{\mathfrak{A}}$ such that neither $\text{supp}(\mu_1) \subseteq \mathfrak{Q}$ nor $\text{supp}(\mu_2) \subseteq \sqrt{\mathfrak{Q}}^F$. Suppose that $x_1 \in \text{supp}(\mu_1) - \mathfrak{Q}$ and $x_2 \in \text{supp}(\mu_2) - \sqrt{\mathfrak{Q}}^F$. Clearly, $\text{supp}(g(x_1^2)) \subseteq \text{supp}(g(\mu_1^2)) \subseteq \mathfrak{Q}$. By the hypothesis, we get $x_1 \in \mathfrak{Q}$ or $x_2 \in \sqrt{\mathfrak{Q}}^F$, a contradiction. \square

Corollary 4.3. Let \mathfrak{Q} be an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{A}, f, g) (with scalar F -identity e'). Then \mathfrak{Q} is primary if and only if for all $a_1^n \in \mathfrak{A}$,

$\text{supp}(g(a_1^n)) \subseteq \mathfrak{Q}$ implies that $a_i \in \mathfrak{Q}$ or $\text{supp}(g(a_1^{i-1}, e', a_{i+1}^n)) \subseteq \sqrt{\mathfrak{Q}}^F$ for some $1 \leq i \leq n$.

The following is a direct consequence and can be proved easily and so the proof is omitted.

Theorem 4.4. *If \mathfrak{P} is a prime F -hyperideal of \mathfrak{R} , then \mathfrak{P} is a primary F -hyperideal of \mathfrak{R} .*

The next example shows that the inverse of Theorem 4.4 is not true, in general.

Example 4.5. Suppose that $R = [0, 1]$ and $t \in (0, 1]$. Then (R, f, g) is a Krasner $F^{2,3}$ -hyperring, where f and g defined by

$$f(a, b) = \begin{cases} \chi_{\max\{a,b\}} & \text{if } a \neq b \\ \chi_{[0,a]} & \text{if } a = b \end{cases}$$

and

$$g(a, b, c) = (a.b.c)_t$$

for all $a, b, c \in R$. The F -hyperideal $I = [0, 0.5]$ is a primary F -hyperideal of R . But it is not a prime F -hyperideal of R as the fact that $\text{supp}(g(0.8, 0.7, 0.6)) \subseteq I$ but $0.8, 0.7, 0.6 \notin I$.

In next theorem, we establish a relationship between prime F -hyperideals and primary F -hyperideals of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{R}, f, g) .

Theorem 4.6. *Let \mathfrak{Q} be an F -hyperideal of a Krasner $F^{(m,n)}$ -hyperring (\mathfrak{R}, f, g) (with scalar F -identity e'). If \mathfrak{Q} is primary, then $\sqrt{\mathfrak{Q}}^F$ is a prime F -hyperideal of \mathfrak{R} .*

Proof. Let $\text{supp}(g(x_1^n)) \subseteq \sqrt{\mathfrak{Q}}^F$ for some $x_1^n \in \mathfrak{R}$ such that $x_1^{i-1}, x_{i+1}^n \notin \sqrt{\mathfrak{Q}}^F$. Our task now is to show that $a_i \in \sqrt{\mathfrak{Q}}^F$. Let $x \in \text{supp}(g(x_1^n))$. From $\text{supp}(g(x_1^n)) \subseteq \sqrt{\mathfrak{Q}}^F$, it follows that there exists $s \in \mathbb{N}$ such that if $s \leq n$, then $\text{supp}(g(x^{(s)}, e'^{(n-s)})) \subseteq \mathfrak{Q}$. Therefore we have $\text{supp}(g(g(x_1^n)^{(s)}, e'^{(n-s)})) \subseteq \mathfrak{Q}$, because $\text{supp}(g(x^{(s)}, e'^{(n-s)})) = \text{supp}(g(g(x_1^n)^{(s)}, e'^{(n-s)}))$. Thus $\text{supp}(g(x_i^{(s)}, g(x_1^{i-1}, e', x_{i+1}^n)^{(s)}, e'^{(n-2s)})) \subseteq \mathfrak{Q}$ and so $\text{supp}(g(g(x_i^{(s)}, e'^{(n-s)}), g(g(x_1^{i-1}, e', x_{i+1}^n)^{(s)}, e'^{(n-s)}), e'^{(n-2)})) \subseteq \mathfrak{Q}$. Since \mathfrak{Q} is a primary F -hyperideal of \mathfrak{R} , we get $\text{supp}(g(g(x_1^{i-1}, e', x_{i+1}^n)^{(s)}, e'^{(n-s)})) \subseteq \mathfrak{Q}$ or $\text{supp}(g(x_i^{(s)}, e'^{(n-s)})) \subseteq \sqrt{\mathfrak{Q}}^F$. Suppose that $\text{supp}(g(g(x_1^{i-1}, e', x_{i+1}^n)^{(s)}, e'^{(n-s)})) \subseteq \mathfrak{Q}$. Then we get $\text{supp}(g(g(x_1^{(s)}, e'^{(n-s)}), g(x_2^{i-1}, e'^2, x_{i+1}^n)^{(s)}, e'^{(n-s-1)})) \subseteq \mathfrak{Q}$ which means $\text{supp}(g(x_1^{(s)}, e'^{(n-s)})) \subseteq \mathfrak{Q}$ or $\text{supp}(g(g(x_2^{i-1}, e'^2, x_{i+1}^n)^{(s)}, e'^{(n-s)})) \subseteq \sqrt{\mathfrak{Q}}^F$. Since $x_1 \notin \sqrt{\mathfrak{Q}}^F$, then $\text{supp}(g(g(x_2^{i-1}, e'^2, x_{i+1}^n)^{(s)}, e'^{(n-s)})) \subseteq \sqrt{\mathfrak{Q}}^F$. By continuing this process, since $x_1^{i-1}, x_{i+1}^n \notin \sqrt{\mathfrak{Q}}^F$, we have $\text{supp}(g(x_i^{(s)}, e'^{(n-s)})) \subseteq \sqrt{\mathfrak{Q}}^F$. By using the definition of F -radical of \mathfrak{Q} , we conclude that $a_i \in \sqrt{\mathfrak{Q}}^F$. If $s = l(n-1) + 1$, then by using a similar argument, one can easily complete the proof. \square

5. CONCLUSION

In this paper, our purpose is to extend the study initiated in [14] about Krasner $F^{(m,n)}$ -hyperrings by Farshi and Davvaz. We defined prime F -hyperideals, maximal F -hyperideals and primary F -hyperideals of a Krasner $F^{(m,n)}$ -hyperring \mathfrak{R} . We obtain many specific results explaining the structures. The future work can be on

defining the concept of δ -primary F -hyperideals unifying the notions of the prime and primary F -hyperideal in a frame.

REFERENCES

- [1] R. Ameri, M. Norouzi, Prime and primary hyperideals in Krasner (m, n) -hyperrings, *European Journal Of Combinatorics*, (2013) 379-390.
- [2] M. Anbarloei, n -ary 2-absorbing and 2-absorbing primary hyperideals in Krasner (m, n) -hyperrings, *Matematicki Vesnik*, **71** (3) (2019) 250-262.
- [3] M. Anbarloei, Unifying the prime and primary hyperideals under one frame in a Krasner (m, n) -hyperring, *Comm. Algebra*, **49** (2021) 3432-3446.
- [4] A. Asadi, R. Ameri, Direct limit of Krasner (m, n) -hyperrings, *Journal of Sciences*, **31** (1) (2020) 75-83.
- [5] G. Crombez, On (m, n) -rings, *Abh. Math. Semin. Univ.*, Hamburg, **37** (1972) 180-199.
- [6] G. Crombez, J. Timm, On (m, n) -quotient rings, *Abh. Math. Semin. Univ.*, Hamburg, **37** (1972) 200-203.
- [7] S. Corsini, Prolegomena of hypergroup theory, *Second edition*, Aviani editor, Italy, (1993).
- [8] S. Corsini, V. Leoreanu, Applications of hyperstructure theory, *Advances in Mathematics*, vol. 5, Kluwer Academic Publishers, (2003).
- [9] B. Davvaz, V. Leoreanu-Fotea, Hyperring Theory and Applications, *International Academic Press, Palm Harbor, USA*, (2007).
- [10] B. Davvaz, T. Vougiouklis, n -ary hypergroups, *Iran. J. Sci. Technol.*, **30** (A2) (2006) 165-174.
- [11] B. Davvaz, Fuzzy Krasner (m, n) -hyperrings, *Computers and Mathematics with Applications*, **59** (2010) 3879-3891.
- [12] W. Dorente, Untersuchungen über einen verallgemeinerten Gruppenbegriff, *Math. Z.*, **29** (1928) 1-19.
- [13] B. Fahid, Z. Dongsheng, 2-Absorbing δ -primary ideals of commutative rings. *Kyungpook Mathematical Journal*, **57** (2017) 193-198.
- [14] M. Farshi, B. Davvaz, Krasner $F^{(m, n)}$ -hyperrings, *Iranian Journal of Fuzzy Systems*, **11** (6) (2014) 67-88.
- [15] K. Hila, K. Naka, B. Davvaz, On (k, n) -absorbing hyperideals in Krasner (m, n) -hyperrings, *Quarterly Journal of Mathematics*, **69** (2018) 1035-1046.
- [16] E. Kasner, An extension of the group concept (reported by L.G. Weld), *Bull. Amer. Math. Soc.*, **10** (1904) 290-291.
- [17] V. Leoreanu, Canonical n -ary hypergroups, *Ital. J. Pure Appl. Math.*, **24**(2008).
- [18] V. Leoreanu-Fotea, B. Davvaz, n -hypergroups and binary relations, *European J. Combin.*, **29** (2008) 1027-1218.
- [19] V. Leoreanu-Fotea, B. Davvaz, Roughness in n -ary hypergroups, *Inform. Sci.*, **178** (2008) 4114-4124.
- [20] V. Leoreanu-Fotea, B. Davvaz, Fuzzy hyperrings, *Fuzzy Sets Syst.*, **160** (2009) 2366-2378.
- [21] W. J. Liu, Fuzzy invariant subgroups and fuzzy ideals, *Fuzzy Sets Syst.* **8** (1982) 133-139.
- [22] X. Ma, J. Zhan, B. Davvaz, Applications of rough soft sets to Krasner (m, n) -hyperrings and corresponding decision making methods, *Filomat*, **32** (2018) 6599-6614.
- [23] F. Marty, Sur une generalization de la notion de groupe, *8th Congress Math. Scandnaves, Stockholm*, (1934) 45-49.
- [24] S. Mirvakili, B. Davvaz, Relations on Krasner (m, n) -hyperrings, *European J. Combin.*, **31**(2010) 790-802.
- [25] S. Mirvakili, B. Davvaz, Constructions of (m, n) -hyperrings, *Matematicki Vesnik*, **67** (1) (2015) 1-16.
- [26] M. Norouzi, R. Ameri, V. Leoreanu-Fotea, Normal hyperideals in Krasner (m, n) -hyperrings, *An. St. Univ. Ovidius Constanta* **26** (3) (2018) 197-211.
- [27] S. Omid, B. Davvaz, Contribution to study special kinds of hyperideals in ordered semihyperrings, *J. Taibah Univ. Sci.*, **11** (2017) 1083-1094.
- [28] S. Ostadhadi-Dehkordi, B. Davvaz, A Note on Isomorphism Theorems of Krasner (m, n) -hyperrings, *Arabian Journal of Mathematics*, **5** (2016) 103-115.
- [29] A. Rosenfeld, Fuzzy groups. *J. Math. Anal. Appl.*, **35** (1971) 512-517.
- [30] T. Vougiouklis, Hyperstructures and their representations, *Hadronic Press Inc., Florida*, (1994).

- [31] A. Yassine, M.J. Nikmehr, R. Nikandish, n -Ary k -absorbing hyperideals in krasner (m, n) -hyperrings, *Afrika Matematika* (2022) 19-33. doi.org/10.1007/s13370-022-00961-6.
- [32] L. A. Zadeh, Fuzzy sets, *Inform Control* **8**(1965) 338-353.
- [33] M. M. Zahedi, R. Ameri, On the prime, primary and maximal subhypermodules, *Ital. J. Pure Appl. Math.*, **5** (1999) 61-80.
- [34] M. M. Zahedi, A. Hasankhani, F -polygroups I, *J. Fuzzy Math.*, **4**(3) (1996) 533-548.
- [35] M. M. Zahedi, A. Hasankhani, F -polygroups II, *Information Sciences*, **89**(3-4) (1996) 225-243.
- [36] J. Zhan, B. Davvaz, K.P. Shum, Generalized fuzzy hyperideals of hyperrings, *Computers and Mathematics with Applications*, **56** (2008) 1732-1740.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, IMAM KHOMEINI INTERNATIONAL UNIVERSITY, QAZVIN, IRAN.

Email address: m.anbarloei@sci.ikiu.ac.ir