

Derivatives of Complete Weight Enumerators and New Balance Principle of Binary Self-Dual Codes

Vassil Yorgov

Department of Mathematics and Computer Science
Fayetteville State University
1200 Murchison Rd
Fayetteville, NC 28311

July 5, 2022

Abstract

Let $K = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$. It is known that the complete weight enumerator W of a binary self-dual code of length n is an eigenvector corresponding to an eigenvalue 1 of the Kronecker power $K^{[n]}$. For every integer t , $0 \leq t \leq n$, we define the t -th derivative $W_{\langle t \rangle}$ of W in such a way that $W_{\langle t \rangle}$ is in the eigenspace of 1 of the matrix $K^{[n-t]}$. For large values of t , $W_{\langle t \rangle}$ contains less information about the code but has smaller length while $W_{\langle 0 \rangle} = W$ completely determines the code. We compute the derivative of order $n-5$ for the extended Golay code of length 24, the extended quadratic residue code of length 48, and the putative $[72, 24, 12]$ code and show that they are in the eigenspace of 1 of the matrix $K^{[5]}$. We use the derivatives to prove a new balance equation which involves the number of code vectors of given weight having 1 in a selected coordinate position. As an example, we use the balance equation to eliminate some candidates for weight enumerators of binary self-dual codes of length eight.

1 Introduction

In this work, we use the standard notation of error correcting codes. An appropriate reference is the book [4]. Let C be a binary linear self-dual code of length n . Thus C is a linear subspace of dimension $n/2$ of the vector space F^n , where F is the binary field, and C is equal to its orthogonal code with respect to the usual dot product. The exact weight enumerator of C is a row vector W of length 2^n with integer entries labeled with the vectors from F^n listed in lexicographic order. The entry $W[v]$ labeled with a vector v is defined with

$$W[v] = \begin{cases} 1 & \text{if } v \in C \\ 0 & \text{if } v \notin C. \end{cases}$$

Thus W belongs to the vector space Q^{2^n} where Q is the rational field. Clearly the exact weight enumerator W determines completely the code C .

Let $K = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$, $K^{[2]} = K \otimes K$, $K^{[3]} = K^{[2]} \otimes K$, and let $K^{[n]} = K^{[n-1]} \otimes K$, be the n -th Kronecker power of K . A summary of the properties of Kronecker products of matrices is given in [10]. Let R be the real number field. The matrix K has eigenvalues 1 and -1 . The vectors $(1, \sqrt{2}-1)$ and $(1, -\sqrt{2}-1)$ are eigenvectors of K corresponding to 1 and -1 . Therefore, K is simple, meaning that there exist a basis of R^2 consisting of eigenvectors of K . The rows of the matrix

$$B = \begin{bmatrix} 1 & \sqrt{2}-1 \\ 1 & -\sqrt{2}-1 \end{bmatrix}$$

form such basis of R^2 . It follows that the matrix $K^{[n]}$ is also simple and has eigenvalues 1 and -1 , [5] page 413. The rows of the Kronecker power $B^{[n]}$ form a basis of R^{2^n} . We label the rows of $B^{[n]}$ with the vectors from F^n listed in lexicographic order.

Lemma 1 *The rows of $B^{[n]}$ with even weight labels form a basis of the eigenspace of 1 of $K^{[n]}$.*

Proof. We have $BK = DB$ where

$$D = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Using the properties of Kronecker power, we obtain $(BK)^{[n]} = (DB)^{[n]}$ and $B^{[n]}K^{[n]} = D^{[n]}B^{[n]}$. Since the diagonal matrix $D^{[n]}$ has 1 on the rows labeled with even weight vectors and -1 on the rows labeled with odd weight vectors, the lemma follows from the last equality. ■

The next lemma is a particular case of a result proved in [8].

Lemma 2 [8] *The complete weight enumerator W of the binary self-dual code C belongs to the eigenspace of 1 of the matrix $K^{[n]}$. Moreover, any nontrivial $(0,1)$ vector from the eigenspace of 1 of the matrix $K^{[n]}$ determines a self-dual binary code.*

Since 2^n increases very rapidly with n , this lemma is not practical when the code length n increases. In the next section we define derivatives of W which contain less information about the code C but have smaller size.

2 Derivatives of the Complete Weight Enumerator

Let $\rho = \sqrt{2}-1$ and $\mu = -\sqrt{2}-1$. The quadratic extension $Q(\rho)$ has automorphism group of order two generated by the automorphism $\Phi : Q(\rho) \rightarrow Q(\rho)$ defined with $\Phi(\rho) = \mu$.

Definition 3 For every integer t , $0 \leq t \leq n$, the t -th derivative of W is the row vector $W_{\langle t \rangle}$ of length 2^{n-t} having an entry with label $v \in F^{n-t}$ determined by

$$W_{\langle t \rangle}[v] = \sum_{u \in F^t} \rho^{wt(u)} W[uv]$$

where $wt(u)$ is the Hamming weight of u and uv is the concatenation of u and v .

For example, if $u = (1, 1, 0, 1)$ and $v = (1, 0)$, then $uv = (1, 1, 0, 1, 1, 0)$. It follows that every derivative $W_{\langle t \rangle}$ has nonnegative entries from the quadratic extension $Q(\rho)$. Particularly, $W_{\langle n \rangle}$ is determined by the weight distribution of C .

Lemma 4

$$W_{\langle n \rangle} = \sum_{u \in F^n} \rho^{wt(u)} W[u] = \sum_{k=0}^n A_k \rho^k$$

where A_k is the number of weight k vectors in C .

Thus $W_{\langle n \rangle} \in Q(\rho)$ and contains less information about C than the weight distribution of C . On the other end, $W_{\langle 0 \rangle} = W$ determines the code C completely.

Lemma 5 For every integer t , $0 \leq t \leq n-1$ and for every $v \in F^{n-t-1}$, we have $W_{\langle t+1 \rangle}[v] = W_{\langle t \rangle}[(0)v] + \rho W_{\langle t \rangle}[(1)v]$.

Proof. The proof is straightforward:

$$\begin{aligned} W_{\langle t+1 \rangle}[v] &= \sum_{u' \in F^{t+1}} \rho^{wt(u')} W[u'v] \\ &= \sum_{u \in F^t} \rho^{wt(u(0))} W[u(0)v] + \sum_{u \in F^t} \rho^{wt(u(1))} W[u(1)v] \\ &= \sum_{u \in F^t} \rho^{wt(u)} W[u(0)v] + \rho \sum_{u \in F^t} \rho^{wt(u)} W[u(1)v] \\ &= W_{\langle t \rangle}[(0)v] + \rho W_{\langle t \rangle}[(1)v]. \end{aligned}$$

■

Less formally, $W_{\langle t+1 \rangle} = W_{\langle t \rangle}[0] + \rho W_{\langle t \rangle}[1]$ where $W_{\langle t \rangle}[0]$ is the first half and $W_{\langle t \rangle}[1]$ is the second half of $W_{\langle t \rangle}$.

The next lemma shows that every entry of the second half of $W_{\langle t \rangle}$ depends only on a corresponding entry from the first half of $W_{\langle t \rangle}$.

Lemma 6 For every integer t , $0 \leq t \leq n-1$ and for every $v \in F^{n-t-1}$, we have $W_{\langle t \rangle}[(1)\bar{v}] = (-1)^{wt(v)} \rho^t \Phi(W_{\langle t \rangle}[(0)v])$ where \bar{v} is the complementary vector of v and Φ is the automorphism of $Q(\rho)$ defined above.

Proof. As the all one vector is in the code C , for any $u \in F^t$ we have $W[\bar{u}(1)\bar{v}] = W[u(0)v]$ where \bar{u} and \bar{v} are the complementary vectors of u and v . By Definition 3,

$$\begin{aligned}
W_{\langle t \rangle}[(1)\bar{v}] &= \sum_{\bar{u} \in F^t} \rho^{wt(\bar{u})} W[\bar{u}(1)\bar{v}] \\
&= \sum_{\bar{u} \in F^t} \rho^{wt(\bar{u})} W[u(0)v] \\
&= \sum_{u \in F^t} \rho^{t-wt(u)} W[u(0)v] \\
&= \rho^t \sum_{u \in F^t} \rho^{-wt(u)} W[u(0)v]
\end{aligned}$$

We check that $\rho^{-1} = -\mu$. Because C contains only even weight vectors, $W[u(0)v] = 0$ when $wt(u)$ and $wt(v)$ have different parities. Therefore

$$\begin{aligned}
W_{\langle t \rangle}[(1)\bar{v}] &= \rho^t \sum_{u \in F^t} (-\mu)^{wt(u)} W[u(0)v] \\
&= \rho^t (-1)^{wt(v)} \sum_{u \in F^t} \mu^{wt(u)} W[u(0)v] \\
&= \rho^t (-1)^{wt(v)} \Phi \left(\sum_{u \in F^t} \rho^{wt(u)} W[u(0)v] \right) \\
&= (-1)^{wt(v)} \rho^t \Phi(W_{\langle t \rangle}[(0)v])
\end{aligned}$$

■

Theorem 7 For every integer t , $0 \leq t \leq n$, the t -th derivative $W_{\langle t \rangle}$ belongs to the eigenspace of 1 of $K^{[n-t]}$.

Proof. The statement of the theorem is true for $W_{\langle 0 \rangle} = W$ from Lemma 2. Thus $WK^{[n]} = W$. Since $W = (W[0]W[1])$ (recall that this is the concatenation of $W[0]$ and $W[1]$) and

$$K^{[n]} = K \otimes K^{[n-1]} = \frac{1}{\sqrt{2}} \begin{bmatrix} K^{[n-1]} & K^{[n-1]} \\ K^{[n-1]} & -K^{[n-1]} \end{bmatrix}$$

we have

$$\begin{aligned}
\frac{1}{\sqrt{2}}(W[0] + W[1])K^{[n-1]} &= W[0] \\
\frac{1}{\sqrt{2}}(W[0] - W[1])K^{[n-1]} &= W[1].
\end{aligned}$$

By adding and then subtracting these equations, we obtain

$$\begin{aligned}
W[0]K^{[n-1]} &= \frac{1}{\sqrt{2}}(W[0] + W[1])K^{[n-1]} \\
W[1]K^{[n-1]} &= \frac{1}{\sqrt{2}}(W[0] - W[1])K^{[n-1]}.
\end{aligned}$$

Now it is easy to check that

$$\begin{aligned}
(W[0] + (\sqrt{2} - 1)W[1])K^{[n-1]} &= \frac{1}{\sqrt{2}}(W[0] + W[1]) + \frac{\sqrt{2} - 1}{\sqrt{2}}(W[0] - W[1]) \\
&= W[0] + \frac{2 - \sqrt{2}}{\sqrt{2}}W[1] \\
&= W[0] + (\sqrt{2} - 1)W[1].
\end{aligned}$$

Hence, $W_{\langle 1 \rangle} = W_{\langle 0 \rangle}[0] + \rho W_{\langle 0 \rangle}[1] = W[0] + (\sqrt{2} - 1)W[1]$ is in the eigenspace of 1 of $K^{[n-1]}$.

From this, we proof similarly that $W_{\langle 2 \rangle} = W_{\langle 1 \rangle}[0] + \rho W_{\langle 1 \rangle}[1]$ is in the eigenspace of 1 of $K^{[n-2]}$.

Using Lemma 5, we complete the proof by induction on t . ■

3 A Balance Principle for Self-Dual Codes

Theorem 8 *Let t be a selected coordinate position of C , let $\delta \in F = \{0, 1\}$, and let $A_{k,\delta}$ be the cardinality of the set*

$$\{v \in C \mid wt(v) = k \text{ and } v[t] = \delta\}.$$

Then $\sum_{k=0}^n A_{k,1}\rho^{k-1} = \sum_{k=0}^n A_{k,0}\rho^{k+1}$ and $\sum_{k=0}^n A_{k,1}\rho^{k-1} = \frac{1+\rho}{4}W_{\langle n \rangle}$. Particularly, $\sum_{k=0}^n A_{k,1}\rho^{k-1}$ does not depend on the selection of t .

Proof. Let $t = n$. From Theorem 6, $W_{\langle n-1 \rangle} = (W_{\langle n-1 \rangle}(0)W_{\langle n-1 \rangle}(1))$ is in the eigenspace of 1 of K , which is generated by vector $(1, \rho)$. Therefore, $W_{\langle n-1 \rangle}(1) = \rho W_{\langle n-1 \rangle}(0)$. Since

$$\begin{aligned}
W_{\langle n-1 \rangle}(1) &= \sum_{u \in F^{n-1}} \rho^{wt(u)}W[u(1)] = \sum_{k=0}^n A_{k,1}\rho^{k-1} \\
W_{\langle n-1 \rangle}(0) &= \sum_{u \in F^{n-1}} \rho^{wt(u)}W[u(0)] = \sum_{k=0}^n A_{k,0}\rho^k
\end{aligned}$$

we obtain the first equality of the Theorem. From Lemma 5, $W_{\langle n \rangle} = W_{\langle n-1 \rangle}(0) + \rho W_{\langle n-1 \rangle}(1)$. Thus $W_{\langle n \rangle} = W_{\langle n-1 \rangle}(0) + \rho^2 W_{\langle n-1 \rangle}(0)$. Using Lemma 4, we obtain $\sum_{k=0}^n A_k \rho^k = (1 + \rho^2)W_{\langle n-1 \rangle}(0)$ and $W_{\langle n-1 \rangle}(0) = \frac{1}{1+\rho^2} \sum_{k=0}^n A_k \rho^k$. Then $W_{\langle n-1 \rangle}(1) = \frac{\rho}{1+\rho^2} \sum_{k=0}^n A_k \rho^k$. Since $\frac{\rho}{1+\rho^2} = \frac{1+\rho}{4}$, we obtain the second equality of the Theorem.

If $t \neq n$, we apply a permutation to the coordinates of C which sends t to n . Since equivalent codes have the same weight distribution, the statement of the Theorem holds for $t \neq n$. ■

4 Examples

4.1 Binary Self-Dual Codes of Length 8

There are two binary self-dual codes of length 8 [7]. They have weight enumerators

$W_a = (1, 0, 0, 0, 14, 0, 0, 0, 1)$ and $W_b = (1, 0, 4, 0, 6, 0, 4, 0, 1)$. Here the entry A_k in position $1+k$ is equal to the number of code vectors of weight k , $k = 0, 1, \dots, 8$.

Each of these weight enumerators satisfies the equation $XM = 2^4X$ where M is the transpose of the Krawtchouk matrix K_8 [8]. The entry of K_n in row i and column j is

$$\sum_{m=0}^i (-1)^m \binom{j}{m} \binom{n-j}{i-m}.$$

Using computer algebra system Magma [2], we determine all solutions of this equation with nonnegative integer entries. They are:

$(1, 0, 0, 0, 14, 0, 0, 0, 1)$, $(1, 0, 1, 0, 12, 0, 1, 0, 1)$,
 $(1, 0, 20, 10, 0, 20, 1)$, $(1, 0, 30, 8, 0, 30, 1)$,
 $(1, 0, 4, 0, 6, 0, 40, 1)$, $(1, 0, 5, 0, 4, 0, 5, 0, 1)$,
 $(1, 0, 6, 0, 20, 6, 0, 1)$, $(1, 0, 7, 0, 0, 0, 7, 0, 1)$.

Let $A_{2,0} = y$ for a self-dual binary code realizing some of these solutions. Then we can express the possible nonzero values of $A_{k,0}$ and $A_{k,1}$ with y . We have

k	0	2	4	6	8
$A_{k,0}$	1	y	$\frac{1}{2}A_4$	$A_2 - y$	0
$A_{k,1}$	0	$A_2 - y$	$\frac{1}{2}A_4$	y	1

For each of the eight possible weight enumerators we solve for y the balance equation from Theorem 7. The found values of y are $0, \frac{3}{4}, \frac{3}{2}, \frac{9}{4}, 3, \frac{15}{4}, \frac{9}{2}, \frac{21}{4}$. Since y must be a nonnegative integer, this result eliminates all possible weight enumerators except for W_a and W_b .

4.2 The Golay Code of Length 24

This well known code is the only binary self-dual code of length 24 with minimum weight 8 [6]. The nonzero entries of the weight distribution of this code are $A_0 = 1$, $A_8 = 759$, $A_{12} = 2576$, $A_{16} = 759$, $A_{24} = 1$. The supports of all code vectors of weight 8, 12, and 16 form 5-designs [1]. Hence we can compute all values $A_{k,0}$ and $A_{k,1}$. The result is given in the table

k	0	8	12	16	24
$A_{k,0}$	1	506	$\frac{2576}{2}$	$759 - 506$	0
$A_{k,1}$	0	$759 - 506$	$\frac{2576}{2}$	506	1

These values satisfy the balance equation of Theorem 7.

Using the three 5-design supported by the code words of weight 8, 12, and 16, we compute the derivative $W_{\langle 19 \rangle}$ and check that it satisfies the equation $W_{\langle 19 \rangle} K^{[5]} = W_{\langle 19 \rangle}$. Its entries are given below:

$$-1167936*p + 483776, 1202240*p - 497984,$$

1202240*p - 497984, -1180608*p + 489024,
 1202240*p - 497984, -1180608*p + 489024,
 -1180608*p + 489024, 1023936*p - 424128,
 1202240*p - 497984, -1180608*p + 489024,
 -1180608*p + 489024, 1023936*p - 424128,
 -1180608*p + 489024, 1023936*p - 424128,
 1023936*p - 424128, -750144*p + 310720,
 1202240*p - 497984, -1180608*p + 489024,
 -1180608*p + 489024, 1023936*p - 424128,
 -1180608*p + 489024, 1023936*p - 424128,
 1023936*p - 424128, -750144*p + 310720,
 -1180608*p + 489024, 1023936*p - 424128,
 1023936*p - 424128, -750144*p + 310720,
 1023936*p - 424128, -750144*p + 310720,
 -750144*p + 310720, 7081024*p - 2933056

4.3 The Extended Quadratic Residue Code of Length 48

This is the only $[48, 24, 12]$ binary self-dual code [3]. The supports of all code vectors of weight 12, 16, 20, 24, 28, 32, and 36 form 5-designs [1]. This allows us to compute the derivative $W_{\langle 43 \rangle}$ and to check that $W_{\langle 43 \rangle} K^{[5]} = W_{\langle 43 \rangle}$. The entries of $W_{\langle 43 \rangle}$ are

-1801295637184512*p + 746121082765312,
 1799773991403520*p - 745490796445696,
 1799773991403520*p - 745490796445696,
 -1787591301070848*p + 740444560883712,
 1799773991403520*p - 745490796445696,
 -1787591301070848*p + 740444560883712,
 -1787591301070848*p + 740444560883712,
 1749670572392448*p - 724737280770048,
 1799773991403520*p - 745490796445696,
 -1787591301070848*p + 740444560883712,
 -1787591301070848*p + 740444560883712,
 1749670572392448*p - 724737280770048,
 -1787591301070848*p + 740444560883712,
 1749670572392448*p - 724737280770048,
 1749670572392448*p - 724737280770048,
 -1666630947373056*p + 690341141872640,
 1799773991403520*p - 745490796445696,
 -1787591301070848*p + 740444560883712,
 -1787591301070848*p + 740444560883712,
 1749670572392448*p - 724737280770048,
 -1787591301070848*p + 740444560883712,
 1749670572392448*p - 724737280770048,
 1749670572392448*p - 724737280770048,
 -1666630947373056*p + 690341141872640,

$-1787591301070848^*p + 740444560883712,$
 $1749670572392448^*p - 724737280770048,$
 $1749670572392448^*p - 724737280770048,$
 $-1666630947373056^*p + 690341141872640,$
 $1749670572392448^*p - 724737280770048,$
 $-1666630947373056^*p + 690341141872640,$
 $-1666630947373056^*p + 690341141872640,$
 $11704454583615488^*p - 4848143828713472$

4.4 The Putative [72,36,16] Binary Self-Dual Code

The existence of this code is a long standing open question [9]. Its weight enumerator is known:

$[<0,1>, <16,249849>, <20,18106704>, <24,462962955>, <28,4397342400>, <32,16602715899>, <36,25756721120>, <40,16602715899>, <44,4397342400>, <48,462962955>, <52,18106704>, <56,249849>, <72,1>].$

The supports all code vectors of weight 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, and 56 form 5-designs [1]. Using this designs, we compute the derivative $W_{<67>}$ and check that $W_{<67>}K^{[5]} = W_{<67>}$. The entries of $W_{<67>}$ are

$-2766052144192871265730560^*p + 1145736312355867181711360,$
 $2764814255733869297795072^*p - 1145223562167443435552768,$
 $2764814255733869297795072^*p - 1145223562167443435552768,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $2764814255733869297795072^*p - 1145223562167443435552768,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$
 $2764814255733869297795072^*p - 1145223562167443435552768,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$
 $-2713300183161139309314048^*p + 1123885734654746797539328,$
 $2764814255733869297795072^*p - 1145223562167443435552768,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$
 $-2713300183161139309314048^*p + 1123885734654746797539328,$
 $-2759776102730699957600256^*p + 1143136690864219955920896,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$
 $2745563299524396139413504^*p - 1137249555016829104029696,$

-2713300183161139309314048*p + 1123885734654746797539328,
 2745563299524396139413504*p - 1137249555016829104029696,
 -2713300183161139309314048*p + 1123885734654746797539328,
 -2713300183161139309314048*p + 1123885734654746797539328,
 18297763639587213294960640*p - 7579181860614308846632960

The existence of this code remains an open question. If a code exists, there must be a chain of derivatives $W_{\langle 67 \rangle}, \dots, W_{\langle 0 \rangle}$ with $W_{\langle 0 \rangle}$ having only 0 and 1 entries. Lemma 5, Lemma 6, and Theorem 7 may be useful tools in a search for such chain.

Acknowledgement 9 *This copy of computer algebra system Magma [2] used in this work has been made available through a generous initiative of the Simons Foundation covering U.S. colleges, universities, nonprofit research entities, and their students, faculty, and staff.*

5 References

References

- [1] E. F. Assmus, Jr. and H. F. Mattson, Jr., New 5-designs, J. Combin. Theory, vol. 6, 122–151, 1969.
- [2] W. Bosma, J. Cannon, and C. Playoust, The Magma algebra system. I. The user language, J. Symbolic Comput., 24, 235–265, 1997.
- [3] S. K. Houghten, C. W. H. Lam, L. H. Thiel, J. A. Parker, The extended quadratic residue code is the only (48, 24, 12) self-dual doubly-even code, IEEE Trans. Inform. Theory, IT-49, 53–59 2003.
- [4] W. C. Huffman and V. Pless, Fundamentals of Error-Correcting Codes, Cambridge University Press, Cambridge, 2003.
- [5] P. Lancaster and M. Tismenetsky, The Theory of Matrices with Applications, Academic Press, New York, NY, USA, 1985.
- [6] V. Pless, On the uniqueness of the Golay codes, J. Combin. Theory 5, 215–228 1968.
- [7] V. Pless, A classification of self-orthogonal codes over GF(2), Discrete Math., vol. 3, 209-246, 1972.
- [8] J. Simonis, MacWilliams identities and coordinate partitions, Linear Algebra Appl., 216: 81-91, 1995.
- [9] N. J. A. Sloane, Is there a (72, 36) d=16 self-dual code, IEEE Trans. Inform. Theory, IT-19, 251, 1973.
- [10] H. Zhang and F. Ding, On the Kronecker products and their applications, J. of Applied Math., vol. 2013, 1-8.