

RADII CONSTANTS FOR FUNCTIONS WITH FIXED SECOND COEFFICIENT

SHALINI RANA, OM AHUJA, NAVEEN KUMAR JAIN

ABSTRACT. We introduce three classes of analytic functions with fixed second coefficient which are defined using the class \mathcal{P} of analytic functions with positive real part. The objective is to determine radii such that the three classes are contained in various subclasses of starlike functions. The radii estimated in the present investigation are better than the radii obtained earlier. Furthermore, connections with previous known results are shown.

1. INTRODUCTION

Let Δ denote the open unit disc in \mathbb{C} and the class \mathcal{A} be defined as collection of all analytic functions in Δ satisfying $f(0) = 1$ and $f'(0) = 1$. The class \mathcal{S} is defined to be collection of univalent functions in class \mathcal{A} . The well known Bieberbach theorem states that for a univalent function $f(z) = z + a_2z^2 + \dots$ the bound on second coefficient, that is $|a_2| \leq 2$, plays an important role in the study of univalent function theory. This bound attracted the interest of many mathematicians which led to the investigation of the class A_b consisting of the functions of the form $f(z) = z + a_2z^2 + \dots$, $|a_2| = b$ for a fixed b with $0 \leq b \leq 1$. For $n \in \mathbb{N}$ and $0 \leq b \leq 1$, let \mathcal{A}_{nb} be the class of analytic functions of the form $f(z) = z + nbz^2 + \dots$ for $z \in \Delta$ such that $\mathcal{A}_b := \mathcal{A}_{1b}$. The study of class A_b was initiated as early as 1920 by Gronwall [13]. He determined the growth and distortion estimates for the class of univalent functions with fixed second coefficients. In 2011, Ali *et al.* [5] obtained various results for the class of functions with fixed second coefficients by applying the theory of second order differential subordination. Later, Lee *et al.* [19] investigated certain applications of differential subordination for such functions. Kumar *et al.* [17] determined the best possible estimates on initial coefficients of Ma-Minda type univalent functions. Ali *et al.* [4] obtained sharp radii of starlikeness for certain classes of functions with fixed second coefficients. A survey on functions with fixed initial coefficient can be found in [1].

Let f and h be analytic functions in Δ . Then f is subordinate to h , that is $f \prec h$ if $f(z) = h(w(z))$ for some analytic function w , with $w(0) = 0$ and $|w(z)| < 1$. In particular, if $h \in \mathcal{S}$, then $f \prec h$ provided $f(0) = h(0)$ and $f(\Delta) \subset h(\Delta)$. Several well

2020 *Mathematics Subject Classification.* 30C45, 30C80.

Key words and phrases. Univalent functions . Subordination . Starlike functions . Radius of starlikeness . Radius constants.

known subclasses of starlike functions were characterized by subordination of zf'/f to some function in \mathcal{P} . Using a function ψ , univalent in unit disc Δ with $\operatorname{Re}(\psi(z)) > 0$, starlike with respect to $\psi(0) = 1$, symmetric about the real axis and $\psi'(0) > 0$, Ma and Minda [20] gave a unified treatment for several subclasses of starlike functions by studying the class $\mathcal{S}^*(\psi) = \{f \in \mathcal{A} : zf'(z)/f(z) \prec \psi(z)\}$. Recently, Anand *et al.* [7] also obtained some results for a class of analytic functions defined using the function ψ . Several authors considered the class $\mathcal{S}^*(\psi)$ for various choices of the function ψ . For $-1 < B < A \leq 1$, let $\psi(z) = (1 + Az)/(1 + Bz)$ be a convex function whose image is symmetric with respect to the real axis. For this function ψ , the class $\mathcal{S}^*(\psi)$ reduces to the class $\mathcal{S}^*[A, B]$ [15] (see also [38]) of Janowski starlike functions. Other well-known choices for $\psi(z)$ include $\sqrt{1+z}$, $\exp z$, $1 + (4/3)z + (2/3)z^2$, $1 + \sin(z)$, $z + \sqrt{1+z^2}$, $1 + z - z^3/3$, $2/(1 + e^{-z})$. For a brief survey of these classes, reader may refer to [12, 39].

Consider the class $\mathcal{P}(\alpha)$ of analytic functions $p(z) = 1 + b_1z + b_2z^2 + \dots$ satisfying $\operatorname{Re}\{p(z)\} > \alpha$ where $0 \leq \alpha < 1$. It can be noted that the class $\mathcal{P} = \mathcal{P}(0)$. For $p \in \mathcal{P}(\alpha)$ [29], we have $|b_1| \leq 2(1-\alpha)$. Denote by $\mathcal{P}_b(\alpha)$ the subclass of $\mathcal{P}(\alpha)$ consisting of functions of the form $p(z) = 1 + 2b(1-\alpha)z + \dots$ for $b \in [0, 1]$ such that $\mathcal{P}_b = \mathcal{P}_b(0)$. For any two subclasses \mathcal{M} and \mathcal{N} of \mathcal{A} , the \mathcal{N} radius for the class \mathcal{M} is the largest number $\rho \in (0, 1)$ such that $r^{-1}f(rz) \in \mathcal{N}$, for all $f \in \mathcal{M}$ and for $0 < r < \rho$. Among several studies available on radius problems, many results involving ratio between two classes of functions, where one of them belong to some particular subclasses of \mathcal{A} has been a center of major focus in geometric function theory and can be seen in [31, 32, 22, 23, 24]. Recently, Lecko and Ravichandran[18] estimated certain best possible radii for the classes of function $g \in \mathcal{A}$ satisfying the conditions (i) $g/h \in \mathcal{P}$ where $h/(zp) \in \mathcal{P}$ or $h/(zp) \in \mathcal{P}(1/2)$ (ii) $g/(zp) \in \mathcal{P}$. Motivated by above work, we define three classes of functions by making use of the classes \mathcal{A}_{6b} , \mathcal{A}_{4c} and \mathcal{P} as follows:

$$H_{b,c}^1 = \{f \in \mathcal{A}_{6b} : \frac{f}{g} \in \mathcal{P} \text{ and } \frac{g}{zp} \in \mathcal{P} \text{ where } g \in \mathcal{A}_{4c}, p \in \mathcal{P}\}$$

$$H_{b,c}^2 = \{f \in \mathcal{A}_{5b} : \frac{f}{g} \in \mathcal{P} \text{ and } \frac{g}{zp} \in \mathcal{P}(1/2) \text{ where } g \in \mathcal{A}_{3c}, p \in \mathcal{P}\}$$

and

$$H_b^3 = \{f \in \mathcal{A}_{4b} : \frac{f}{zp} \in \mathcal{P} \text{ where } p \in \mathcal{P}\}$$

where $b \in [0, 1]$ and $c \in [0, 1]$.

The results presented in this paper are nice extensions of radii estimates results in [18] along with some improved radii. It includes radii estimates for functions in the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 to belong to several subclasses of normalized analytic functions \mathcal{A} like starlike functions of order α , starlike functions associated with lemniscate of Bernoulli, parabolic starlike functions, exponential function, cardioid, sine function, lune, a particular rational function, nephroid and modified sigmoid function.

2. ANALYSIS AND MAPPING OF $zf'(z)/f(z)$ FOR $H_{b,c}^1$, $H_{b,c}^2$ AND H_b^3

In this section, we investigate extremal functions for all three classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 which demonstrates the fact that the classes are non empty. Furthermore, we obtain the disc centered at 1, containing the image of the disc Δ under the mapping zf'/f where f belongs to each of these classes. We begin by stating the following lemmas by McCarty:

Lemma 2.1. [27] *Let $b \in [0, 1]$ and $0 \leq \alpha < 1$. If $p \in \mathcal{P}_b(\alpha)$, then for $|z| = r < 1$,*

$$\left| \frac{zp'(z)}{p(z)} \right| \leq \frac{2(1-\alpha)r}{1-r^2} \frac{br^2 + 2r + b}{(1-2\alpha)r^2 + 2b(1-\alpha)r + 1}.$$

Lemma 2.2. [26] *Let $b \in [0, 1]$ and $0 \leq \alpha < 1$. If $p \in \mathcal{P}_b(\alpha)$, then for $|z| = r < 1$,*

$$\operatorname{Re} \left(\frac{zp'(z)}{p(z)} \right) \geq \begin{cases} \frac{-2(1-\alpha)r}{1+2\alpha br+(2\alpha-1)r^2} \frac{br^2+2r+b}{r^2+2br+1} & \text{if } R_\alpha \leq R_b \\ \frac{2\sqrt{\alpha C_1 - C_1 - \alpha}}{1-\alpha} & \text{if } R_\alpha \geq R_b \end{cases},$$

where $R_b = C_b - D_b$, $R_\alpha = \sqrt{\alpha C_1}$, and $r = |z| < 1$;

$$(1) \quad C_b = \frac{(1+br)^2 - (2\alpha-1)(b+r)^2 r^2}{(1+2br+r^2)(1-r^2)} \quad \text{and} \quad D_b = \frac{2(1-\alpha)(b+r)(1+br)r}{(1+2br+r^2)(1-r^2)}$$

2.1. The class $H_{b,c}^1$. Let the functions f and g whose Taylor series expansions are given by $f(z) = z + f_1 z^2 + \dots$ and $g(z) = z + g_1 z^2 + \dots$ be such that $\operatorname{Re}\{f/g\} > 0$ and $\operatorname{Re}\{g/(zp)\} > 0$ where $p \in \mathcal{P}$ is represented by $p(z) = 1 + qz + \dots$. Now consider, $g/(zp) = 1 + (g_1 - q)z + \dots$ where $|g_1 - q| \leq 2$ and $|q| \leq 2$ which gives $|g_1| \leq 4$. Also, $f/g = 1 + (f_1 - g_1)z + \dots$ where $|f_1 - g_1| \leq 2$ and hence $|f_1| \leq 6$. Hence, for $b, c \in [0, 1]$, we consider the class involving the function f and g with fixed second coefficient whose Taylor series expansion is given by $f(z) = z + 6bz^2 + \dots$ and $g(z) = z + 4cz^2 + \dots$ such that $f \in \mathcal{A}_{6b}$ and $g \in \mathcal{A}_{4c}$. If the function $f \in H_{b,c}^1$, then there exists an element $g \in \mathcal{A}_{4c}$ and $p \in \mathcal{P}$ such that $f/g \in \mathcal{P}$ and $g/(zp) \in \mathcal{P}$. Define

$$h(z) = \frac{f(z)}{g(z)} = 1 + (6b - 4c)z + \dots \quad \text{and} \quad k(z) = \frac{g(z)}{zp(z)} = 1 + (4c - q)z + \dots$$

where $|6b - 4c| \leq 2$ and $|4c - q| \leq 2$. Therefore, we observe that $h \in \mathcal{P}_{(6b-4c)/2}$, $k \in \mathcal{P}_{(4c-q)/2}$, $p \in \mathcal{P}_{q/2}$ and f can be expressed as $f(z) = zp(z)h(z)k(z)$. Then a calculation shows that

$$(2) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \left| \frac{zh'(z)}{h(z)} \right| + \left| \frac{zk'(z)}{k(z)} \right| + \left| \frac{zp'(z)}{p(z)} \right|.$$

For $d = |6b - 4c| \leq 2$, $s = |4c - q| \leq 2$ and $\alpha = 0$, using lemma 2.1, we get

$$(3) \quad \left| \frac{zh'(z)}{h(z)} \right| \leq \frac{r}{(1-r^2)} \frac{(dr^2 + 4r + d)}{(r^2 + dr + 1)}, \quad \left| \frac{zk'(z)}{k(z)} \right| \leq \frac{r}{(1-r^2)} \frac{(sr^2 + 4r + s)}{(r^2 + sr + 1)},$$

and

$$(4) \quad \left| \frac{zp'(z)}{p(z)} \right| \leq \frac{r}{(1-r^2)} \frac{(qr^2 + 4r + q)}{(r^2 + qr + 1)}.$$

By (2), (3) and (4), it follows that

$$(5) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \frac{r[(dr^2 + 4r + d)(r^2 + sr + 1)(r^2 + qr + 1) + (sr^2 + 4r + s)(r^2 + dr + 1)(r^2 + qr + 1) + (qr^2 + 4r + q)(r^2 + dr + 1)(r^2 + sr + 1)]}{(1-r^2)(r^2 + dr + 1)(r^2 + sr + 1)(r^2 + qr + 1)}.$$

Define the functions f_1 , g_1 and $p_1 : \Delta \rightarrow \mathbb{C}$ by

$$(6) \quad f_1(z) = \frac{z(1 - qz + z^2)(1 - (4c - q)z + z^2)(1 - (6b - 4c)z + z^2)}{(1 - z^2)^3},$$

$$(7) \quad g_1(z) = \frac{z(1 - qz + z^2)(1 - (4c - q)z + z^2)}{(1 - z^2)^2}, \quad \text{and} \quad p_1(z) = \frac{(1 - qz + z^2)}{(1 - z^2)},$$

$$\text{where } |6b - 4c| \leq 2, \quad |4c - q| \leq 2 \quad \text{and} \quad |q| \leq 2.$$

By (6) and (7), we have

$$\frac{f_1(z)}{g_1(z)} = \frac{(1 - (6b - 4c)z + z^2)}{(1 - z^2)} = \frac{1 + w_1(z)}{1 - w_1(z)}$$

$$\frac{g_1(z)}{zp_1(z)} = \frac{(1 - (4c - q)z + z^2)}{(1 - z^2)} = \frac{1 + w_2(z)}{1 - w_2(z)} \quad \text{and} \quad p_1(z) = \frac{(1 - qz + z^2)}{(1 - z^2)} = \frac{1 + w_3(z)}{1 - w_3(z)}$$

where

$$w_1(z) = \frac{z(z - (6b - 4c)/2)}{(1 - z(6b - 4c)/2)}, \quad w_2(z) = \frac{z(z - (4c - q)/2)}{(1 - z(4c - q)/2)} \quad \text{and} \quad w_3(z) = \frac{z(z - q/2)}{(1 - zq/2)}$$

which are analytic functions satisfying the conditions of Schwarz lemma in the unit disc Δ and hence $\text{Re}(f_1/g_1) > 0$, $\text{Re}(g_1/(zp_1)) > 0$ and $\text{Re}(p_1) > 0$. Thus, $f_1/g_1 \in \mathcal{P}_{(6b-4c)/2}$, $g_1/(zp_1) \in \mathcal{P}_{(4c-q)/2}$ and $p_1 \in \mathcal{P}_{(q/2)}$. Thus the function $f_1 \in H_{b,c}^1$ and the class $H_{b,c}^1$ is non-empty.

$$(8) \quad F_1(z) = \frac{z(1 - z^2)^3}{(1 - qz + z^2)(1 - (4c - q)z + z^2)(1 - (6b - 4c)z + z^2)}$$

and $f_1(z)$ are extreme functions for the class $H_{b,c}^1$ provided $q \leq 2$, $c \geq q/4$ and $b \geq 2c/3$.

2.2. The class $H_{b,c}^2$. Let f and g be functions with Taylor series expansions $f(z) = z + f_1z^2 + \dots$ and $g(z) = z + g_1z^2 + \dots$ such that $\text{Re}\{f/g\} > 0$ and $\text{Re}\{g/(zp)\} > 1/2$ where $p \in \mathcal{P}$ is represented by $p(z) = 1 + qz + \dots$. Now consider, $g/zp = 1 + (g_1 - q)z + \dots$ where $|g_1 - q| \leq 1$ (lemma 2, pg 33 et al [14]) and $|q| \leq 2$ which gives $|g_1| \leq 3$. Also, $f/g = 1 + (f_1 - g_1)z + \dots$ where $|f_1 - g_1| \leq 2$ and hence $|f_1| \leq 5$. Therefore, we consider the class involving the function f and g with fixed second coefficient whose Taylor series expansion is given by $f(z) = z + 5bz^2 + \dots$ and $g(z) = z + 3cz^2 + \dots$ where $b \in [0, 1]$ and

$c \in [0, 1]$ such that $f \in \mathcal{A}_{5b}$ and $g \in \mathcal{A}_{3c}$. If the function $f \in H_{b,c}^2$ then there exists an element $g \in \mathcal{A}_{3c}$ and $p \in \mathcal{P}$ such that $f/g \in \mathcal{P}$ and $g/(zp) \in \mathcal{P}(1/2)$. Define

$$h(z) = \frac{f(z)}{g(z)} = 1 + (5b - 3c)z + \dots$$

and

$$k(z) = \frac{g(z)}{zp(z)} = 1 + (3c - q)z + \dots$$

It is easy to see that $h \in \mathcal{P}_{(5b-3c)/2}$, $k \in \mathcal{P}_{(3c-q)}$, $p \in P_{q/2}$ and $f(z) = zp(z)h(z)k(z)$. Then a calculation shows that

$$(9) \quad \frac{zf'(z)}{f(z)} = 1 + \frac{zh'(z)}{h(z)} + \frac{zk'(z)}{k(z)} + \frac{zp'(z)}{p(z)}$$

so that

$$(10) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \left| \frac{zh'(z)}{h(z)} \right| + \left| \frac{zk'(z)}{k(z)} \right| + \left| \frac{zp'(z)}{p(z)} \right|.$$

Let $m = |5b - 3c| \leq 2$, $n = |3c - q| \leq 1$ and $\alpha = 0$. Using Lemma 2.1 for the functions h, p and k , we have

$$(11) \quad \left| \frac{zh'(z)}{h(z)} \right| \leq \frac{r}{(1-r^2)} \frac{(mr^2 + 4r + m)}{(r^2 + mr + 1)}, \quad \left| \frac{zk'(z)}{k(z)} \right| \leq \frac{r}{(1-r^2)} \frac{(nr^2 + 2r + n)}{(nr + 1)},$$

and

$$(12) \quad \left| \frac{zp'(z)}{p(z)} \right| \leq \frac{r}{(1-r^2)} \frac{(qr^2 + 4r + q)}{(r^2 + qr + 1)}.$$

Inequality (10) together with (11) and (12), gives

$$(13) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \frac{r}{(1-r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right).$$

Define the functions $f_2, g_2 : \Delta \rightarrow \mathbb{C}$ by

$$(14) \quad f_2(z) = \frac{z(1 - qz + z^2)(1 - (3c - q)z)(1 - (5b - 3c)z + z^2)}{(1 - z^2)^3}$$

and

$$(15) \quad g_2(z) = \frac{z(1 - qz + z^2)(1 - (3c - q)z)}{(1 - z^2)^2}, \text{ where } |5b - 3c| \leq 2 \text{ and } |3c - q| \leq 1.$$

It follows from (7), (14) and (15) that

$$\frac{f_2(z)}{g_2(z)} = \frac{(1 - (5b - 3c)z + z^2)}{(1 - z^2)} = \frac{1 + w_4(z)}{1 - w_4(z)}$$

and

$$\frac{g_2(z)}{zp_1(z)} = \frac{(1 - (3c - q)z)}{(1 - z^2)} = \frac{1 + w_5(z)}{1 - w_5(z)}$$

where

$$w_4(z) = \frac{z(z - (5b - 3c)/2)}{(1 - z(5b - 3c)/2)} \text{ and } w_5(z) = \frac{z(z - (3c - q))}{(2 - (3c - q)z - z^2)}$$

which are Schwarz functions in the unit disc Δ and hence $\operatorname{Re}(f_2/g_2) > 0$, $\operatorname{Re}(g_2/(zp_1)) > 1/2$ and $\operatorname{Re}(p_1) > 0$ (as shown for class $H_{b,c}^1$). Thus, $f_2/g_2 \in \mathcal{P}_{(5b-3c)/2}$, $g_2/(zp_1) \in \mathcal{P}_{(3c-q)}$ and $p_1 \in \mathcal{P}_{(q/2)}$. Thus the function $f_2 \in H_{b,c}^2$ and the class $H_{b,c}^2$ is non-empty. $f_2(z)$ is an extreme functions for the class $H_{b,c}^2$ provided $q \leq 2$, $c \geq q/3$ and $b \geq 3c/5$.

2.3. The class H_b^3 . Let the functions f and g whose Taylor series expansions are given by $f(z) = z + f_1z^2 + \dots$ be such that $\operatorname{Re}\{f/(zp)\} > 0$ where $p \in \mathcal{P}$ is represented by $p(z) = 1 + qz + \dots$. Now consider, $f/(zp) = 1 + (f_1 - q)z + \dots$ where $|f_1 - q| \leq 2$ and $|q| \leq 2$ which gives $|f_1| \leq 4$. Therefore, we consider the function f with fixed second coefficient whose Taylor series expansion is given by $f(z) = z + 4bz^2 + \dots$ where $b \in [0, 1]$ such that $f \in \mathcal{A}_{4b}$. If the function $f \in H_b^3$ then there exists $p \in \mathcal{P}$ such that $f/(zp) \in \mathcal{P}$. Define

$$h(z) = \frac{f}{zp}(z) = 1 + (4b - q)z + \dots$$

so that $h \in \mathcal{P}_{(4b-q)/2}$, $p \in \mathcal{P}_{q/2}$ and f can be expressed as $f(z) = zp(z)h(z)$. Then a calculation shows that

$$(16) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \left| \frac{zh'(z)}{h(z)} \right| + \left| \frac{zp'(z)}{p(z)} \right|.$$

From lemma 2.1, for $l = |4b - q| \leq 2$, $|q| \leq 2$ and $\alpha = 0$, we get

$$(17) \quad \left| \frac{zh'(z)}{h(z)} \right| \leq \frac{r}{(1-r^2)} \frac{(lr^2 + 4r + l)}{(r^2 + lr + 1)} \quad \text{and} \quad \left| \frac{zp'(z)}{p(z)} \right| \leq \frac{r}{(1-r^2)} \frac{(qr^2 + 4r + q)}{(r^2 + qr + 1)}.$$

Using (16) and (17), it is easy to show that

$$(18) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \frac{r[(lr^2 + 4r + l)(qr^2 + 4r + q) + (qr^2 + 4r + q)(r^2 + lr + 1)]}{(1-r^2)(r^2 + lr + 1)(r^2 + qr + 1)}.$$

Define the functions $f_3 : \Delta \rightarrow \mathbb{C}$ by

$$(19) \quad f_3(z) = \frac{z(1 - qz + z^2)(1 - (4b - q)z + z^2)}{(1 - z^2)^2}, \quad |4b - q| \leq 2.$$

Then, (17) together with (7) gives

$$\frac{f_3(z)}{zp_1(z)} = \frac{(1 - (4b - q)z + z^2)}{(1 - z^2)} = \frac{1 + w_6(z)}{1 - w_6(z)}, \quad \text{where } w_6(z) = \frac{z(z - (4b - q)/2)}{(1 - z(4b - q)/2)}$$

which is analytic function satisfying the conditions of Schwarz lemma in the unit disc Δ and hence $\operatorname{Re}(f_3/(zp_1)) > 0$ and $\operatorname{Re}(p_1) > 0$ (shown above in class $H_{b,c}^1$). Thus, $f_3/(zp_1) \in \mathcal{P}_{(6b-q)/2}$ and $p_1 \in \mathcal{P}_{(q/2)}$. Thus, the function $f_3 \in H_b^3$ and the class H_b^3 is non-empty.

$$(20) \quad F_3(z) = \frac{z(1 - z^2)^2}{(1 - qz + z^2)(1 - (4b - q)z + z^2)}, \quad |4b - q| \leq 2$$

and $f_3(z)$ are extreme functions for the class H_b^3 provided $q \leq 2$ and $b \geq q/4$. Using the information in the above subsections, we investigate several radius problems associated with functions in the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 in the next section.

3. RADIUS OF STARLIKENESS

In the present section, we determine the radii of the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 to belong to various Ma-Minda starlike classes of functions. Throughout this section we assume that $d = |6b - 4c| \leq 2$, $s = |4c - q| \leq 2$, $m = |5b - 3c| \leq 2$, $n = |3c - q| \leq 1$, $l = |4b - q| \leq 2$. For $0 \leq \alpha \leq 1$, the class $\mathcal{S}^*(\alpha) = \mathcal{S}^*[1 - 2\alpha, -1] = \{f \in \mathcal{A} : \operatorname{Re}(zf'(z)/f(z)) > \alpha\}$ is the class of starlike functions of order α .

Theorem 3.1. *The sharp $\mathcal{S}^*(\alpha)$ radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:*

(i) *For the class $H_{b,c}^1$, the sharp $\mathcal{S}^*(\alpha)$ radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where*

$$x_1(r) = \alpha - 1 + \alpha(d + s + q)r + (10 + 2\alpha + ds + \alpha ds + (1 + \alpha)(d + s)q)r^2 + ((10 + \alpha)(s + q) + d(10 + \alpha + (2 + \alpha)sq))r^3 + 8(3 + sq + d(s + q))r^4 + (-(-12 + \alpha)(s + q) - d(-12 + \alpha + (-4 + \alpha)sq))r^5 + (14 + 3sq + 3d(s + q) - \alpha(2 + sq + d(s + q)))r^6 - (-2 + \alpha)(d + s + q)r^7 + (1 - \alpha)r^8.$$

(ii) *For the class $H_{b,c}^2$, the sharp $\mathcal{S}^*(\alpha)$ radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where*

$$x_2(r) = -\frac{4}{1 - r^2} - \frac{1}{1 + nr} + \frac{2 + 2nr}{1 + 2nr + r^2} + \frac{2 + mr}{1 + r(m + r)} + \frac{2 + qr}{1 + r(q + r)} - \alpha.$$

(iii) *For the class H_b^3 , the sharp $\mathcal{S}^*(\alpha)$ radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where*

$$x_3(r) = \alpha - 1 + \alpha(l + q)r + (7 + \alpha + (1 + \alpha)lq)r^2 + 6(l + q)r^3 + (9 + 3lq - \alpha(1 + lq))r^4 - (-2 + \alpha)(l + q)r^5 + (1 - \alpha)r^6.$$

Proof. Radius estimate for respective classes is as follows:

(i) Note that $x_1(0) = (\alpha - 1) < 0$ and $x_1(1) = 6(2 + d)(2 + e)(2 + q) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. For $f \in H_{b,c}^1$, using (5), we have

$$(21) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \frac{r}{(1 - r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right)$$

which yields

$$(22) \quad \operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) \geq 1 - \frac{r}{(1 - r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \geq \alpha$$

whenever $x_1(r) \leq 0$. This shows that $\operatorname{Re}(zf'(z)/f(z)) \geq \alpha$ for $|z| = r \leq \rho_1$.

For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (22), the function $F_1(z)$ defined for the class $H_{b,c}^1$ in (8) at $z = -\rho_1$ satisfies the equality

$$\operatorname{Re} \left(\frac{zF_1'(z)}{F_1(z)} \right) = 1 - \frac{\rho_1}{(1 - \rho_1^2)} \left(\frac{u\rho_1^2 + 4\rho_1 + u}{\rho_1^2 + u\rho_1 + 1} + \frac{v\rho_1^2 + 4\rho_1 + v}{\rho_1^2 + v\rho_1 + 1} + \frac{q\rho_1^2 + 4\rho_1 + q}{\rho_1^2 + q\rho_1 + 1} \right) = \alpha.$$

Thus, the radius is sharp.

- (ii) A calculation shows that $x_2(0) = 1 - \alpha > 0$ and $x_2(1/3) = -(-60(-5 + (-12 + n)n) + 6(135 + n(148 + 21n))q + 3m(9(30 + 17q) + n(296 + 42n + 156q + 27nq)) + 2(10 + 3m)(3 + n)(5 + 3n)(10 + 3q)\alpha)/(2(10 + 3m)(3 + n)(5 + 3n)(10 + 3q)) < 0$. By intermediate value theorem, there exists a root $r \in (0, 1/3)$ of the equation $x_2(r) = 0$. Let $\rho_2 \in (0, 1/3)$ be the smallest root of the equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. An easy calculation shows that for $0 < r < 1/3$,

$$(23) \quad C_b - D_b - \sqrt{C_1/2} = \frac{-1 + 4r^2 + 2b^2r^2 + 8br^3 + r^4 + 2b^2r^4}{2(-1 + r)(1 + r)(1 + 2br + r^2)^2} > 0,$$

where C_b and D_b are given by (1). From (9) and (11), and using Lemma 2.2 together with (23), we have

(24)

$$\operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) \geq 1 - \left(\frac{r}{1 - r^2} \right) \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) - \frac{r(n + 2r + nr^2)}{(1 + nr)(1 + 2nr + r^2)} \geq \alpha$$

whenever $x_2(r) \leq 0$. Thus, $\operatorname{Re}(zf'(z)/f(z)) \geq \alpha$ for $|z| = r \leq \rho_2$. To prove the sharpness, consider the function $F_2, G_2 : \Delta \rightarrow \mathbb{C}$ defined by

$$F_2(z) = \frac{z(1 + 3cz - qz)(1 - z^2)^2}{(1 + 5bz - 3cz + z^2)(1 + 6cz - 2qz + z^2)(1 + qz + z^2)}$$

and

$$G_2(z) = z \frac{1 - z^2}{(1 + qz + z^2)(1 + 3cz - qz)},$$

where $|5b - 3c| \leq 2$ and $|3c - q| \leq 1$. Note that for $c = (1 + q)/3$,

$$\frac{F_2(z)}{G_2(z)} = \frac{(1 - z^2)}{1 - (1 - 5b + q)z + z^2} = \frac{1 + w_1(z)}{1 - w_1(z)}$$

and

$$\frac{G_2(z)}{z p_1(z)} = \frac{1}{1 + 3cz - qz} = \frac{1 + w_2(z)}{1 - w_2(z)},$$

where

$$w_1(z) = \frac{(1 - 5b + q - 2z)z}{2 + (-1 + 5b - q)z} \quad \text{and} \quad w_2(z) = \frac{(q - 3c)z}{2 + (3c - q)z}$$

which are Schwarz functions and hence $\operatorname{Re}(F_2/G_2) > 0$, $\operatorname{Re}(G_2/(zp_1)) > 0$ and $\operatorname{Re}(p_1) > 0$. For $5b - 3c \geq 0$, $3c - q = 1$ and $z = \rho_2$, it follows from (24) that

$$\begin{aligned} & \operatorname{Re} \left(\frac{zF_2'(z)}{F_2(z)} \right) \\ &= 1 - \left(\frac{\rho_2}{1 - \rho_2^2} \right) \left(\frac{m\rho_2^2 + 4\rho_2 + m}{\rho_2^2 + m\rho_2 + 1} + \frac{q\rho_2^2 + 4\rho_2 + q}{\rho_2^2 + q\rho_2 + 1} \right) - \frac{\rho_2(n + 2\rho_2 + n\rho_2^2)}{(1 + n\rho_2)(1 + 2n\rho_2 + \rho_2^2)} \\ &= \alpha. \end{aligned}$$

- (iii) It is easy to see that $x_3(0) = \alpha - 1 < 0$ and $x_3(1) = 4(2 + l)(2 + q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$. Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. From (18) it follows that for any $f \in H_b^3$,

$$(25) \quad \operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) \geq 1 - \frac{r}{1 - r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \geq \alpha$$

whenever $x_3(r) \leq 0$. This proves that $\operatorname{Re}(zf'(z)/f(z)) \geq \alpha$ for $|z| = r \leq \rho_3$. The result is sharp for the function F_3 defined for the class H_b^3 in (20). At $z = -\rho_3$, and for $u = 4b - q \geq 0$, it follows from (25) that

$$\operatorname{Re} \left(\frac{zF_3'(z)}{F_3(z)} \right) = 1 - \frac{\rho_3}{1 - \rho_3^2} \left(\frac{u\rho_3^2 + 4\rho_3 + u}{\rho_3^2 + u\rho_3 + 1} + \frac{q\rho_3^2 + 4\rho_3 + q}{\rho_3^2 + q\rho_3 + 1} \right) = \alpha.$$

□

Remark 3.1. For $b = 1$ and $c = 1$ and $q = 2$, Theorem 3.1 yields the result [18, Theorem 1, p. 6].

The class $\mathcal{S}_L^* = \mathcal{S}^*(\sqrt{1+z})$ and it represents the collection of functions in the class \mathcal{A} whose $zf'(z)/f(z)$ lies in the region bounded by the lemniscate of Bernoulli $|w^2 - 1| = 1$. Various studies on \mathcal{S}_L^* can be seen in [37, 25]. In the following result we obtain the sharp radii results for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 .

Theorem 3.2. The sharp \mathcal{S}_L^* radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:

- (i) For the class $H_{b,c}^1$, the sharp \mathcal{S}_L^* radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where
- $$x_1(r) = (1 - \sqrt{2}) - (-2 + \sqrt{2})(d + s + q)r + (-2(-7 + \sqrt{2}) - (-3 + \sqrt{2})(sq + d(s + q)))r^2 + (-(-12 + \sqrt{2})(s + q) - d(-12 + \sqrt{2} + (-4 + \sqrt{2})sq))r^3 + 8(3 + sq + d(s + q))r^4 + ((10 + \sqrt{2})(s + q) + d(10 + \sqrt{2} + (2 + \sqrt{2})sq))r^5 + (2(5 + \sqrt{2}) + (1 + \sqrt{2})(sq + d(s + q)))r^6 + \sqrt{2}(d + s + q)r^7 + (-1 + \sqrt{2})r^8.$$
- (ii) For the class $H_{b,c}^2$, the sharp \mathcal{S}_L^* radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where
- $$x_2(r) = (1 - \sqrt{2}) - (-2 + \sqrt{2})(m + n + q)r + (11 - \sqrt{2} - (-3 + \sqrt{2})(nq + m(n + q)))r^2 +$$

(iii) For the class H_b^3 , the sharp \mathcal{S}_L^* radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where

$$x_3(r) = (1 - \sqrt{2}) - (-2 + \sqrt{2})(l + q)r + (9 - \sqrt{2} - (-3 + \sqrt{2})lq)r^2 + 6(l + q)r^3 + (7 + \sqrt{2} + (1 + \sqrt{2})lq)r^4 + \sqrt{2}(l + q)r^5 + (-1 + \sqrt{2})r^6.$$

Proof. (i) Note that $x_1(0) = (1 - \sqrt{2}) < 0$ and $x_1(1) = 6(2 + d)(2 + q)(2 + s) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. Ali *et al.* [2, Lemma 2.2] had proved, for $2\sqrt{2}/3 < C < \sqrt{2}$, that

$$(26) \quad \left\{ w \in \mathbb{C} : |w - C| < \sqrt{2} - C \right\} \subset \left\{ w \in \mathbb{C} : |w^2 - 1| < 1 \right\}.$$

In view of (26) and the fact that the centre of the disc in (21) is 1, $f \in \mathcal{S}_L^*$ if

$$(27) \quad \frac{r}{(1 - r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \sqrt{2} - 1$$

which is equivalent to $f \in \mathcal{S}_L^*$ if $x_1(r) \leq 0$. Since $x_1(0) < 0$ and ρ_1 is the smallest root of the equation $x_1(r) = 0$, $x_1(r)$ is an increasing function on $(0, \rho_1)$. In view of this $f \in \mathcal{S}_L^*$ for $|z| = r \leq \rho_1$. For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (27), the function $f_1(z)$ defined for the class $H_{b,c}^1$ in (6) at $z = -\rho_1$, satisfies the following equality

$$\begin{aligned} & \left| \left(\frac{zf_1'(z)}{f_1(z)} \right)^2 - 1 \right| \\ &= \left| \left(1 - \frac{\rho_1}{(1 - \rho_1^2)} \left(\frac{u\rho_1^2 + 4\rho_1 + u}{\rho_1^2 + u\rho_1 + 1} + \frac{v\rho_1^2 + 4\rho_1 + v}{\rho_1^2 + v\rho_1 + 1} + \frac{q\rho_1^2 + 4\rho_1 + q}{\rho_1^2 + q\rho_1 + 1} \right) \right)^2 - 1 \right| = 1. \end{aligned}$$

Thus, the radius is sharp.

(ii) A calculation shows that $x_2(0) = 1 - \sqrt{2} < 0$ and $x_2(1) = 6(2 + m)(1 + n)(2 + q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_2(r) = 0$. Let $\rho_2 \in (0, 1)$ be the smallest root of the equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. As the centre of the disc in (13) is 1, by (26), $f \in \mathcal{S}_L^*$ if

$$(28) \quad \frac{r}{(1 - r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \sqrt{2} - 1$$

which is equivalent to $f \in \mathcal{S}_L^*$ if $x_2(r) \leq 0$. Since $x_2(0) < 0$ and ρ_2 is the smallest root of the equation $x_2(r) = 0$, $x_2(r)$ is an increasing function on $(0, \rho_2)$. Thus, $f \in \mathcal{S}_L^*$ for $|z| = r \leq \rho_2$. To prove the sharpness, consider the function f_2 defined

in (14). For $u = 5b - 3c \geq 0$, $v = 3c - q \geq 0$ and $z = -\rho_2$, it follows from (28) that

$$\begin{aligned} & \left| \left(\frac{zf'_2(z)}{f_2(z)} \right)^2 - 1 \right| \\ &= \left| \left(1 - \frac{\rho_2}{(1-\rho_2^2)} \left(\frac{u\rho_2^2 + 4\rho_2 + u}{\rho_2^2 + u\rho_2 + 1} + \frac{v\rho_2^2 + 2\rho_2 + v}{v\rho_2 + 1} + \frac{q\rho_2^2 + 4\rho_2 + q}{\rho_2^2 + q\rho_2 + 1} \right) \right)^2 - 1 \right| = 1 \end{aligned}$$

- (iii) It is easy to see that $x_3(0) = 1 - \sqrt{2} < 0$ and $x_3(1) = 4(2+l)(2+q) > 0$, by intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$. Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. From (16) and (17) it follows that for any $f \in H_b^3$

$$(29) \quad \left| \frac{zf'(z)}{f(z)} - 1 \right| \leq \frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right)$$

As the centre of the disc in (29) is 1, by (26), $f \in \mathcal{S}_L^*$ if

$$(30) \quad \frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \sqrt{2} - 1$$

which is equivalent to $f \in \mathcal{S}_L^*$ if $x_3(r) \leq 0$. Since $x_3(0) < 0$ and ρ_3 is the smallest root of the equation $x_3(r) = 0$, $x_3(r)$ is an increasing function on $(0, \rho_3)$. This proves that $f \in \mathcal{S}_L^*$ for $|z| = r \leq \rho_3$.

The result is sharp for the function f_3 defined for the class H_b^3 in (19). At $z = -\rho_3$, and for $u = 4b - q \geq 0$, it follows from (30) that

$$\left| \left(\frac{zf'_3(z)}{f_3(z)} \right)^2 - 1 \right| = \left| \left(\frac{\rho_3}{1-\rho_3^2} \left(\frac{u\rho_3^2 + 4\rho_3 + u}{\rho_3^2 + u\rho_3 + 1} + \frac{q\rho_3^2 + 4\rho_3 + q}{\rho_3^2 + q\rho_3 + 1} \right) \right)^2 - 1 \right| = 1.$$

□

Remark 3.2. For $b = 1$ and $c = 1$ and $q = 2$, Theorem 3.2 yields the result [18, Theorem 2, p. 8].

For $\phi_{PAR} = 1 + \frac{2}{\pi^2} \left(\log \left(\frac{1 + \sqrt{z}}{1 - \sqrt{z}} \right) \right)^2$ the class $\mathcal{S}_p^* := \mathcal{S}^*(\phi_{PAR})$ is the class of parabolic starlike functions. A function $f \in \mathcal{S}_p^*$ provided $zf'(z)/f(z)$ lies in the parabolic region given by $\{w \in \mathbb{C} : \operatorname{Re}(w) > |w - 1|\}$. For further reading refer to [6, 10, 21, 33]. The following theorem gives the radius of parabolic starlikeness of the three classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 .

Theorem 3.3. The \mathcal{S}_p^* radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:

- (i) For the class $H_{b,c}^1$, the sharp \mathcal{S}_p^* radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where

- $x_1(r) = -1 + (d+s+q)r + (22+3sq+3d(s+q))r^2 + (21(s+q)+d(21+5sq))r^3 + 16(3+sq+d(s+q))r^4 + (23(s+q)+d(23+7sq))r^5 + (26+5sq+5d(s+q))r^6 + 3(d+s+q)r^7 + r^8$.
- (ii) For the class $H_{b,c}^2$, \mathcal{S}_p^* radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where
- $x_2(r) = -1 + (m+q)r + (18+4mn+3mq+4nq)r^2 + (17m+36n+17q+8mnq)r^3 + (40+28mn+12mq+28nq)r^4 + (19m+40n+19q+12mnq)r^5 + (22+8mn+5mq+8nq)r^6 + (3m+4n+3q)r^7 + r^8$.
- (iii) For the class H_b^3 , the sharp \mathcal{S}_p^* radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where
- $x_3(r) = -1 + (l+q)r + 3(5+lq)r^2 + 12(l+q)r^3 + (17+5lq)r^4 + 3(l+q)r^5 + r^6$.

Proof. (i) Note that $x_1(0) = -1 < 0$ and $x_1(1) = 12(2+d)(2+q)(2+s) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. Shanmughan and Ravichandran [34, p. 321] had proved, for $1/2 < C < 3/2$, that

$$(31) \quad \{w \in \mathbb{C} : |w - C| < C - 1/2\} \subset \{w \in \mathbb{C} : \operatorname{Re}(w) > |w - 1|\}.$$

As the centre of the disc in (21) is 1, by (31), $f \in \mathcal{S}_p^*$ if

$$(32) \quad \frac{r}{(1-r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{1}{2}$$

which is equivalent to $f \in \mathcal{S}_p^*$ if $x_1(r) \leq 0$. Since $x_1(0) < 0$ and ρ_1 is the smallest root of the equation $x_1(r) = 0$, $x_1(r)$ is an increasing function on $(0, \rho_1)$. In view of this $f \in \mathcal{S}_p^*$ for $|z| = r \leq \rho_1$. For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (32), the function $F_1(z)$ defined for the class $H_{b,c}^1$ in (8) at $z = -\rho_1$, satisfies the following equality

$$\begin{aligned} \operatorname{Re} \left(\frac{zF_1'(z)}{F_1(z)} \right) &= \left| 1 - \frac{\rho_1}{(1-\rho_1^2)} \left(\frac{u\rho_1^2 + 4\rho_1 + u}{\rho_1^2 + u\rho_1 + 1} + \frac{v\rho_1^2 + 4\rho_1 + v}{\rho_1^2 + v\rho_1 + 1} + \frac{q\rho_1^2 + 4\rho_1 + q}{\rho_1^2 + q\rho_1 + 1} \right) \right| \\ &= \left| \frac{zF_1'(z)}{F_1(z)} - 1 \right|. \end{aligned}$$

Thus, the radius is sharp.

- (ii) A calculation shows that $x_2(0) = -1 < 0$ and

$$x_2 \left(\frac{1}{3} \right) = \frac{4(9m(190 + 89q + n(146 + 63q)) + 2(1250 + 855q + 3n(410 + 219q)))}{6561}$$

which is greater than 0. By intermediate value theorem, there exists a root $r \in (0, 1/3)$ of the equation $x_2(r) = 0$. Let $\rho_2 \in (0, 1/3)$ be the smallest root of the equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. From (9) and (11), and using Lemma 2.2 together with

(23), we have

$$\begin{aligned} & \operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) \\ & \geq 1 - \left(\frac{r}{1-r^2} \right) \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) - \frac{r(n + 2r + nr^2)}{(1 + nr)(1 + 2nr + r^2)} \\ & \geq \frac{r}{(1-r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \geq \left| \frac{zf'_2(z)}{f_2(z)} - 1 \right| \end{aligned}$$

whenever $x_2(r) \leq 0$. Since $x_2(0) < 0$ and ρ_2 is the smallest root of the equation $x_2(r) = 0$, $x_2(r)$ is an increasing function on $(0, \rho_2)$. Thus, $f \in \mathcal{S}_p^*$ for $|z| = r \leq \rho_2$.

- (iii) It is easy to see that $x_3(0) = 1 < 0$ and $x_3(1) = 8(2+l)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$. Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. In view of (31) and the fact that the centre of the disc in (29) is 1, $f \in \mathcal{S}_p^*$ if

$$(33) \quad \frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{1}{2}$$

which is equivalent to $f \in \mathcal{S}_p^*$ if $x_3(r) \leq 0$. Since $x_3(0) < 0$ and ρ_3 is the smallest root of the equation $x_3(r) = 0$, $x_3(r)$ is an increasing function on $(0, \rho_3)$. This proves that $f \in \mathcal{S}_p^*$ for $|z| = r \leq \rho_3$.

The result is sharp for the function F_3 defined for the class H_b^3 in (20). At $z = -\rho_3$, and for $u = 4b - q \geq 0$, it follows from (33) that

$$\operatorname{Re} \left(\frac{zF'_3(z)}{F_3(z)} \right) = \left| 1 - \frac{\rho_3}{1-\rho_3^2} \left(\frac{u\rho_3^2 + 4\rho_3 + u}{\rho_3^2 + u\rho_3 + 1} + \frac{q\rho_3^2 + 4\rho_3 + q}{\rho_3^2 + q\rho_3 + 1} \right) \right| = \left| \frac{zF'_3(z)}{F_3(z)} - 1 \right|. \quad \square$$

Remark 3.3. Putting $b = 1$ and $c = 1$ and $q = 2$ in Theorem 3.3, we get the result [18, Theorem 3, p. 9] with the part(ii) having improved radius ($= 0.0990195 > 0.0972$).

In 2015, the class of starlike functions associated with the exponential function as $\mathcal{S}_e^* = \mathcal{S}^*(e^z)$ was introduced by Mendiratta *et al.* [28]. It satisfies the condition $|\log(zf'(z)/f(z))| < 1$.

Theorem 3.4. The \mathcal{S}_e^* radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:

- (i) For the class $H_{b,c}^1$, the sharp \mathcal{S}_e^* radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where
- $$x_1(r) = (1-e) + (d+q+s)r + (2+10e+dq+deq+(1+e)(d+q)s)r^2 + (q+s+10e(q+s)+d(1+qs+2e(5+qs)))r^3 + 8e(3+qs+d(q+s))r^4 + ((-1+12e)(q+s)+d(-1-qs+4e(3+qs)))r^5 + (-2+14e-dq+3deq+(-1+3e)(d+q)s)r^6 + (-1+2e)(d+q+s)r^7 + (-1+e)r^8.$$
- (ii) For the class $H_{b,c}^2$, \mathcal{S}_e^* radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where
- $$x_2(r) = (1-e) + (m+n+q)r + (1+9e+(1+e)(nq+m(n+q)))r^2 + (8e(m+q) +$$

- $n(1 + mq + 2e(5 + mq))r^3 + (-1 - mq + e(13 + 8mn + 5mq + 8nq))r^4 + (-n - q + 4e(3n + q) + m(-1 + 4e)(1 + nq))r^5 + (-1 + 3e)(1 + n(m + q))r^6 + n(-1 + 2e)r^7$.
- (iii) For the class H_b^3 , the sharp \mathcal{S}_e^* radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where
- $$x_3(r) = (1 - e) + (l + q)r + (1 + 7e + (1 + e)lq)r^2 + 6e(l + q)r^3 + (-1 + 9e + (-1 + 3e)lq)r^4 + (-1 + 2e)(l + q)r^5 + (-1 + e)r^6.$$

Proof. (i) Note that $x_1(0) = 1 - e < 0$ and $x_1(1) = 6e(2 + d)(2 + q)(2 + s) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. Mendiratta *et al.*[28] proved, for $e^{-1} \leq C \leq (e + e^{-1})/2$, that

$$(34) \quad \{w \in \mathbb{C} : |w - C| < C - e^{-1}\} \subset \{w \in \mathbb{C} : |\log(w)| < 1\}.$$

As the centre of the disc in (21) is 1, by (34), $f \in \mathcal{S}_e^*$ if

$$(35) \quad \frac{r}{(1 - r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq 1 - \frac{1}{e}$$

which is equivalent to $f \in \mathcal{S}_e^*$ if $x_1(r) \leq 0$. Since $x_1(0) < 0$ and ρ_1 is the smallest root of the equation $x_1(r) = 0$, $x_1(r)$ is an increasing function on $(0, \rho_1)$. In view of this $f \in \mathcal{S}_e^*$ for $|z| = r \leq \rho_1$. For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (35), the function $F_1(z)$ defined for the class $H_{b,c}^1$ in (8) at $z = -\rho_1$, satisfies the following equality

$$\begin{aligned} & \left| \log \left(\frac{zF_1'(z)}{F_1(z)} \right) \right| \\ &= \left| \log \left(1 - \frac{\rho_1}{(1 - \rho_1^2)} \left(\frac{u\rho_1^2 + 4\rho_1 + u}{\rho_1^2 + u\rho_1 + 1} + \frac{v\rho_1^2 + 4\rho_1 + v}{\rho_1^2 + v\rho_1 + 1} + \frac{q\rho_1^2 + 4\rho_1 + q}{\rho_1^2 + q\rho_1 + 1} \right) \right) \right| \\ &= \left| \log \left(\frac{1}{e} \right) \right| = 1. \end{aligned}$$

Thus, the radius is sharp.

- (ii) A calculation shows that $x_2(0) = 1 - e < 0$ and $x_2(1) = 6e(2 + m)(1 + n)(2 + q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_2(r) = 0$. Let $\rho_2 \in (0, 1)$ be the smallest root of the equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. In view of (34) and the fact that the centre of the disc in (13) is 1, $f \in \mathcal{S}_e^*$ if

$$\frac{r}{(1 - r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq 1 - \frac{1}{e}$$

which is equivalent to $f \in \mathcal{S}_e^*$ if $x_2(r) \leq 0$. Since $x_2(0) < 0$ and ρ_2 is the smallest root of the equation $x_2(r) = 0$, $x_2(r)$ is an increasing function on $(0, \rho_2)$. Thus, $f \in \mathcal{S}_e^*$ for $|z| = r \leq \rho_2$.

- (iii) It is easy to see that $x_3(0) = 1 - e < 0$ and $x_3(1) = 4e(2 + l)(2 + q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$.

Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. Since the centre of the disc in (29) is 1, by (34), $f \in \mathcal{S}_e^*$ if

$$(36) \quad \frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq 1 - \frac{1}{e}$$

which is equivalent to $f \in \mathcal{S}_e^*$ if $x_3(r) \leq 0$. Since $x_3(0) < 0$ and ρ_3 is the smallest root of the equation $x_3(r) = 0$, $x_3(r)$ is an increasing function on $(0, \rho_3)$. This proves that $f \in \mathcal{S}_e^*$ for $|z| = r \leq \rho_3$.

The result is sharp for the function F_3 defined for the class H_b^3 in (20). At $z = -\rho_3$, and for $u = 4b - q \geq 0$, it follows from (36) that

$$\begin{aligned} \left| \log \left(\frac{zF_3'(z)}{F_3(z)} \right) \right| &= \left| \log \left(1 - \frac{\rho_3}{1-\rho_3^2} \left(\frac{u\rho_3^2 + 4\rho_3 + u}{\rho_3^2 + u\rho_3 + 1} + \frac{q\rho_3^2 + 4\rho_3 + q}{\rho_3^2 + q\rho_3 + 1} \right) \right) \right| \\ &= \left| \log \left(\frac{1}{e} \right) \right| = 1. \quad \square \end{aligned}$$

Remark 3.4. For $b = 1$ and $c = 1$ and $q = 2$, Theorem 3.4 reduces to the result [18, Theorem 4, p. 10].

The class $\mathcal{S}_c^* = \mathcal{S}^*(1 + (4/3)z + (2/3)z^2)$ is the class of starlike functions $f \in \mathcal{A}$ such that $zf'(z)/f(z)$ lies in the region bounded by the cardioid $\Omega_c = \{u + iv : (9u^2 + 9v^2 - 18u + 5)^2 - 16(9u^2 + 9v^2 - 6u + 1) = 0\}$. Sharma *et al.* [35] studied various properties of the class \mathcal{S}_c^* . The following theorem determines radii for starlike functions associated with the cardioid

Theorem 3.5. The \mathcal{S}_c^* radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:

- (i) For the class $H_{b,c}^1$, the sharp \mathcal{S}_c^* radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where

$$x_1(r) = -2 + (d+q+s)r + 4(8+qs+d(q+s))r^2 + (31(q+s)+d(31+7qs))r^3 + 24(3+qs+d(q+s))r^4 + (35(q+s)+d(35+11qs))r^5 + 8(5+qs+d(q+s))r^6 + 5(d+q+s)r^7 + 2r^8.$$
- (ii) For the class $H_{b,c}^2$, the \mathcal{S}_c^* radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where

$$x_2(r) = -2 + (m+n+q)r + 4(7+nq+m(n+q))r^2 + (31n+24q+m(24+7nq))r^3 + (38+24mn+24nq+14mq)r^4 + (35n+11q+11m(1+nq))r^5 + 8(1+n(m+q))r^6 + 5nr^7.$$
- (iii) For the class H_b^3 , the sharp \mathcal{S}_c^* radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where

$$x_3(r) = -2 + (l+q)r + (22+4lq)r^2 + 18(l+q)r^3 + (26+8lq)r^4 + 5(l+q)r^5 + 2r^6.$$

Proof. (i) Note that $x_1(0) = -2 < 0$ and $x_1(1) = 18(2+d)(2+q)(2+s) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. Sharma *et al.* [35] proved that for $1/3 < C \leq 5/3$,

$$(37) \quad \{w \in \mathbb{C} : |w - C| < (3C - 1)/3\} \subseteq \Omega_c.$$

As the centre of the disc in (21) is 1, by (37), $f \in \mathcal{S}_c^*$ if

$$(38) \quad \frac{r}{(1-r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{2}{3}$$

which is equivalent to $f \in \mathcal{S}_c^*$ if $x_1(r) \leq 0$. Since $x_1(0) < 0$ and ρ_1 is the smallest root of the equation $x_1(r) = 0$, $x_1(r)$ is an increasing function on $(0, \rho_1)$. In view of this $f \in \mathcal{S}_c^*$ for $|z| = r \leq \rho_1$. For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (38), the function $F_1(z)$ defined for the class $H_{b,c}^1$ in (8) at $z = -\rho_1$, satisfies the following equality

$$\left| \frac{zF_1'(z)}{F_1(z)} \right| = \left| 1 - \frac{\rho_1}{(1-\rho_1^2)} \left(\frac{u\rho_1^2 + 4\rho_1 + u}{\rho_1^2 + u\rho_1 + 1} + \frac{v\rho_1^2 + 4\rho_1 + v}{\rho_1^2 + v\rho_1 + 1} + \frac{q\rho_1^2 + 4\rho_1 + q}{\rho_1^2 + q\rho_1 + 1} \right) \right| = \frac{1}{3},$$

which belongs to boundary of the region Ω_c . Thus, the radius is sharp.

- (ii) A calculation shows that $x_2(0) = -2 < 0$ and $x_2(1) = 18(2+m)(1+n)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_2(r) = 0$. Let $\rho_2 \in (0, 1)$ be the smallest root of the equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. In view of (37) and the fact that the centre of the disc in (13) is 1, $f \in \mathcal{S}_c^*$ if

$$\frac{r}{(1-r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{2}{3}$$

which is equivalent to $f \in \mathcal{S}_c^*$ if $x_2(r) \leq 0$. Since $x_2(0) < 0$ and ρ_2 is the smallest root of the equation $x_2(r) = 0$, $x_2(r)$ is an increasing function on $(0, \rho_2)$. Thus, $f \in \mathcal{S}_c^*$ for $|z| = r \leq \rho_2$.

- (iii) It is easy to see that $x_3(0) = -2 < 0$ and $x_3(1) = 12(2+l)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$. Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. In view of the fact that the centre of the disc in (29) is 1, by (37), $f \in \mathcal{S}_c^*$ if

$$(39) \quad \frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{2}{3}$$

which is equivalent to $f \in \mathcal{S}_c^*$ if $x_3(r) \leq 0$. Since $x_3(0) < 0$ and ρ_3 is the smallest root of the equation $x_3(r) = 0$, $x_3(r)$ is an increasing function on $(0, \rho_3)$. This proves that $f \in \mathcal{S}_c^*$ for $|z| = r \leq \rho_3$.

The result is sharp for the function F_3 defined for the class H_b^3 in (20). At $z = -\rho_3$, and for $u = 4b - q \geq 0$, it follows from (39) that

$$\left| \frac{zF_3'(z)}{F_3(z)} \right| = \left| 1 - \frac{\rho_3}{1-\rho_3^2} \left(\frac{u\rho_3^2 + 4\rho_3 + u}{\rho_3^2 + u\rho_3 + 1} + \frac{q\rho_3^2 + 4\rho_3 + q}{\rho_3^2 + q\rho_3 + 1} \right) \right| = \frac{1}{3}. \quad \square$$

Remark 3.5. Putting $b = 1$ and $c = 1$ and $q = 2$ in Theorem 3.5, we get the result [18, Theorem 5, p. 11].

In 2019, Cho *et al.* [8] considered the class of starlike functions $\mathcal{S}_{\sin}^* = \{f \in \mathcal{A} : zf'(z)/f(z) \prec 1 + \sin(z) := q_0(z)\}$ associated with the sine function.

Theorem 3.6. *The sharp \mathcal{S}_{\sin}^* radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:*

- (i) *For the class $H_{b,c}^1$, the sharp \mathcal{S}_{\sin}^* radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where*

$$x_1(r) = -\sin(1) - (d+q+s)(-1+\sin(1))r - 2(-6+\sin(1) - (qs+d(q+s))(\sin(1)-2))r^2 + (11(q+s) + d(11+3qs) - (d+q+s+dqs)\sin(1))r^3 + 8(3+qs+d(q+s))r^4 + ((q+s)(11+\sin(1)) + d(11+\sin(1)+qs(3+\sin(1))))r^5 + ((qs+d(q+s))(2+\sin(1)) + 2(6+\sin(1)))r^6 + (d+q+s)(1+\sin(1))r^7 + \sin(1)r^8.$$
- (ii) *For the class $H_{b,c}^2$, the sharp \mathcal{S}_{\sin}^* radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where*

$$x_2(r) = -\sin(1) - (m+n+q)(\sin(1)-1)r + (10 - (nq+m(n+q))(-2+\sin(1)) - \sin(1))r^2 + (8m+11n+8q+3mnq - n(1+mq)\sin(1))r^3 + (12+8nq+\sin(1)+m(8n+q(4+\sin(1))))r^4 + (q(3+\sin(1))+m(1+nq)(3+\sin(1))+n(11+\sin(1)))r^5 + (1+n(m+q))(2+\sin(1))r^6 + n(1+\sin(1))r^7.$$
- (iii) *For the class H_b^3 , the sharp \mathcal{S}_{\sin}^* radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where*

$$x_3(r) = -\sin(1) - (l+q)(\sin(1)-1)r + (8-lq(-2+\sin(1)) - \sin(1))r^2 + 6(l+q)r^3 + (8+\sin(1)+lq(2+\sin(1)))r^4 + (l+q)(1+\sin(1))r^5 + \sin(1)r^6.$$

Proof. (i) Note that $x_1(0) = -\sin(1) < 0$ and $x_1(1) = 6(2+d)(2+q)(2+s) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. For $|C-1| \leq \sin(1)$, Cho *et al.* [8] established the following inclusion:

$$(40) \quad \{w \in \mathbb{C} : |w-C| < \sin(1) - |C-1|\} \subseteq \Omega_s,$$

where $\Omega_s := q_0(\Delta)$ is the image of the unit disc Δ under the mappings $q_0(z) = 1 + \sin(z)$. As the centre of the disc in (21) is 1, by (40), $f \in \mathcal{S}_{\sin}^*$ if

$$(41) \quad \frac{r}{(1-r^2)} \left(\frac{dr^2+4r+d}{r^2+dr+1} + \frac{sr^2+4r+s}{r^2+sr+1} + \frac{qr^2+4r+q}{r^2+qr+1} \right) \leq \sin(1)$$

which is equivalent to $f \in \mathcal{S}_{\sin}^*$ if $x_1(r) \leq 0$. Since $x_1(0) < 0$ and ρ_1 is the smallest root of the equation $x_1(r) = 0$, $x_1(r)$ is an increasing function on $(0, \rho_1)$. In view of this $f \in \mathcal{S}_{\sin}^*$ for $|z| = r \leq \rho_1$. For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (41), the function $f_1(z)$ defined for class $H_{b,c}^1$ in (6) at $z = -\rho_1$, satisfies the following equality

$$\begin{aligned} \left| \frac{zf_1'(z)}{f_1(z)} \right| &= \left| 1 + \frac{\rho_1}{(1-\rho_1^2)} \left(\frac{u\rho_1^2+4\rho_1+u}{\rho_1^2+u\rho_1+1} + \frac{v\rho_1^2+4\rho_1+v}{\rho_1^2+v\rho_1+1} + \frac{q\rho_1^2+4\rho_1+q}{\rho_1^2+q\rho_1+1} \right) \right| \\ &= 1 + \sin 1 = q_0(1), \end{aligned}$$

which belongs to the boundary of region Ω_s . Thus, the radius is sharp.

- (ii) A calculation shows that $x_2(0) = -\sin(1) < 0$ and $x_2(1) = 6(2+m)(1+n)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation

$x_2(r) = 0$. Let $\rho_2 \in (0, 1)$ be the smallest root of equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. In view of (40) and the fact that centre of the disc in (13) is 1, $f \in \mathcal{S}_{\sin}^*$ if

$$(42) \quad \frac{r}{(1-r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \sin(1)$$

which is equivalent to $f \in \mathcal{S}_{\sin}^*$ if $x_2(r) \leq 0$. Since $x_2(0) < 0$ and ρ_2 is the smallest root of the equation $x_2(r) = 0$, $x_2(r)$ is an increasing function on $(0, \rho_2)$. Thus, $f \in \mathcal{S}_{\sin}^*$ for $|z| = r \leq \rho_2$. To prove sharpness, consider the function f_2 defined in (14). For $u = 5b - 3c \geq 0$, $v = 3c - q \geq 0$ and $z := -\rho_2$, it follows from (42) that

$$\begin{aligned} \left| \frac{zf_2'(z)}{f_2(z)} \right| &= \left| 1 + \frac{\rho_2}{(1-\rho_2^2)} \left(\frac{u\rho_2^2 + 4\rho_2 + u}{\rho_2^2 + u\rho_2 + 1} + \frac{v\rho_2^2 + 2\rho_2 + v}{v\rho_2 + 1} + \frac{q\rho_2^2 + 4\rho_2 + q}{\rho_2^2 + q\rho_2 + 1} \right) \right| \\ &= 1 + \sin(1) \end{aligned}$$

which illustrates the sharpness.

- (iii) It is easy to see that $x_3(0) = -\sin(1) < 0$ and $x_3(1) = 4(2+l)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$. Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. Since the centre of the disc in (29) is 1, by (40), $f \in \mathcal{S}_{\sin}^*$ if

$$(43) \quad \frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \sin(1)$$

which is equivalent to $f \in \mathcal{S}_{\sin}^*$ if $x_3(r) \leq 0$. Since $x_3(0) < 0$ and ρ_3 is the smallest root of the equation $x_3(r) = 0$, $x_3(r)$ is an increasing function on $(0, \rho_3)$. This proves that $f \in \mathcal{S}_{\sin}^*$ for $|z| = r \leq \rho_3$.

The result is sharp for function f_3 defined for the class H_b^3 in (19). At $z = -\rho_3$ and for $u = 4b - q \geq 0$, it follows from (43) that

$$\left| \frac{zf_3'(z)}{f_3(z)} \right| = \left| 1 + \frac{\rho_3}{1-\rho_3^2} \left(\frac{u\rho_3^2 + 4\rho_3 + u}{\rho_3^2 + u\rho_3 + 1} + \frac{q\rho_3^2 + 4\rho_3 + q}{\rho_3^2 + q\rho_3 + 1} \right) \right| = 1 + \sin(1).$$

□

Remark 3.6. Substituting $b = 1$ and $c = 1$ and $q = 2$ in Theorem 3.6, we obtain the result [18, Theorem 6, p. 13].

In 2015, Raina and Sokól [30] introduced the class $\mathcal{S}_{\zeta}^* = \mathcal{S}^*(z + \sqrt{1+z^2})$. Geometrically, a function $f \in \mathcal{S}_{\zeta}^*$ if and only if $zf'(z)/f(z)$ lies in the region bounded by the lune shaped region $\{w \in \mathbb{C} : |w^2 - 1| < 2|w|\}$.

Theorem 3.7. The \mathcal{S}_{ζ}^* radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:

- (i) For the class $H_{b,c}^1$, the sharp \mathcal{S}_{ζ}^* radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where
- $$x_1(r) = \sqrt{2} - 2 + (\sqrt{2} - 1)(d + q + s)r + (2(4 + \sqrt{2}) + \sqrt{2}(qs + d(q + s)))r^2 + ((9 +$$

$\sqrt{2})(q+s) + d(9 + \sqrt{2} + (1 + \sqrt{2})qs)r^3 + 8(3 + qs + d(q+s))r^4 + (-(-13 + \sqrt{2})(q+s) - d(-13 + \sqrt{2} + (-5 + \sqrt{2})qs))r^5 + (-2(-8 + \sqrt{2}) - (-4 + \sqrt{2})(qs + d(q+s)))r^6 - (-3 + \sqrt{2})(d + q + s)r^7 + (2 - \sqrt{2})r^8$.

(ii) For the class $H_{b,c}^2$, the $\mathcal{S}_{\mathcal{C}}^*$ radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where

$x_2(r) = \sqrt{2} - 2 + (\sqrt{2} - 1)(m + n + q)r + (8 + \sqrt{2} + \sqrt{2}(nq + m(n+q)))r^2 + (8(m+q) + n(9 + \sqrt{2} + (1 + \sqrt{2})mq))r^3 + (14 - \sqrt{2} + 6mq - \sqrt{2}mq + 8n(m+q))r^4 + (-(-5 + \sqrt{2})(m+q) - n(-13 + \sqrt{2} + (-5 + \sqrt{2})mq))r^5 - (-4 + \sqrt{2})(1 + n(m+q))r^6 - (\sqrt{2} - 3)nr^7$.

(iii) For the class H_b^3 , the sharp $\mathcal{S}_{\mathcal{C}}^*$ radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where

$x_3(r) = \sqrt{2} - 2 + (\sqrt{2} - 1)(l + q)r + (6 + \sqrt{2} + \sqrt{2}lq)r^2 + 6(l + q)r^3 + (10 - \sqrt{2} - (-4 + \sqrt{2})lq)r^4 - (-3 + \sqrt{2})(l + q)r^5 + (2 - \sqrt{2})r^6$.

Proof. (i) Note that $x_1(0) = \sqrt{2} - 2 < 0$ and $x_1(1) = 6(2 + d)(2 + q)(2 + s) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. Gandhi and Ravichandran [9, Lemma 2.1] proved that for $\sqrt{2} - 1 < C \leq \sqrt{2} + 1$,

$$(44) \quad \{w \in \mathbb{C} : |w - C| < 1 - |\sqrt{2} - C|\} \subseteq \{w \in \mathbb{C} : |w^2 - 1| < 2|w|\}.$$

As the centre of the disc in (21) is 1, by (44), $f \in \mathcal{S}_{\mathcal{C}}^*$ if

$$(45) \quad \frac{r}{(1 - r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq 2 - \sqrt{2}$$

which is equivalent to $f \in \mathcal{S}_{\mathcal{C}}^*$ if $x_1(r) \leq 0$. Since $x_1(0) < 0$ and ρ_1 is the smallest root of the equation $x_1(r) = 0$, $x_1(r)$ is an increasing function on $(0, \rho_1)$. In view of this $f \in \mathcal{S}_{\mathcal{C}}^*$ for $|z| = r \leq \rho_1$. For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (45), the function $F_1(z)$ defined for the class $H_{b,c}^1$ in (8) at $z = -\rho_1$, satisfies the following equality

$$\begin{aligned} & \left| \left(\frac{zf_1'(z)}{f_1(z)} \right)^2 - 1 \right| \\ &= \left| \left(1 - \frac{\rho_1}{(1 - \rho_1^2)} \left(\frac{u\rho_1^2 + 4\rho_1 + u}{\rho_1^2 + u\rho_1 + 1} + \frac{v\rho_1^2 + 4\rho_1 + v}{\rho_1^2 + v\rho_1 + 1} + \frac{q\rho_1^2 + 4\rho_1 + q}{\rho_1^2 + q\rho_1 + 1} \right) \right)^2 - 1 \right| \\ &= |1 - (2 - \sqrt{2})^2 - 1| = 2(1 - \sqrt{2}) = 2 \left| \frac{zf_1'(z)}{f_1(z)} \right|. \end{aligned}$$

Thus, the radius is sharp.

(ii) A calculation shows that $x_2(0) = \sqrt{2} - 2 < 0$ and $x_2(1) = 6(2 + m)(1 + n)(2 + q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation

$x_2(r) = 0$. Let $\rho_2 \in (0, 1)$ be the smallest root of the equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. In view of (44) and the fact centre of the disc in (13) is 1, $f \in \mathcal{S}_{\mathbb{C}}^*$ if

$$\frac{r}{(1-r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq 2 - \sqrt{2}$$

which is equivalent to $f \in \mathcal{S}_{\mathbb{C}}^*$ if $x_2(r) \leq 0$. Since $x_2(0) < 0$ and ρ_2 is the smallest root of the equation $x_2(r) = 0$, $x_2(r)$ is an increasing function on $(0, \rho_2)$. Thus, $f \in \mathcal{S}_{\mathbb{C}}^*$ for $|z| = r \leq \rho_2$.

- (iii) It is easy to see that $x_3(0) = \sqrt{2} - 2 < 0$ and $x_3(1) = 4(2+l)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$. Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. Since the centre of the disc in (29) is 1, by (44), $f \in \mathcal{S}_{\mathbb{C}}^*$ if

$$(46) \quad \frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq 2 - \sqrt{2}$$

which is equivalent to $f \in \mathcal{S}_{\mathbb{C}}^*$ if $x_3(r) \leq 0$. Since $x_3(0) < 0$ and ρ_3 is the smallest root of the equation $x_3(r) = 0$, $x_3(r)$ is an increasing function on $(0, \rho_3)$. This proves that $f \in \mathcal{S}_{\mathbb{C}}^*$ for $|z| = r \leq \rho_3$.

The result is sharp for the function F_3 defined for the class H_b^3 in (20). At $z = -\rho_3$, and for $u = 4b - q \geq 0$, it follows from (46) that

$$\begin{aligned} \left| \left(\frac{zF_3'(z)}{F_3(z)} \right)^2 - 1 \right| &= \left| \left(1 - \frac{\rho_3}{1-\rho_3^2} \left(\frac{u\rho_3^2 + 4\rho_3 + u}{\rho_3^2 + u\rho_3 + 1} + \frac{q\rho_3^2 + 4\rho_3 + q}{\rho_3^2 + q\rho_3 + 1} \right) \right)^2 - 1 \right| \\ &= 2 \left| \frac{zF_3'(z)}{F_3(z)} \right|. \quad \square \end{aligned}$$

Remark 3.7. For $b = 1$ and $c = 1$ and $q = 2$, Theorem 3.7 yields the result [18, Theorem 7, p. 14].

Kumar *et al.*[16] introduced the class of starlike functions, defined by $\mathcal{S}_R^* = \mathcal{S}^*(\psi(z))$, consisting of functions associated with a rational function $\psi(z) = 1 + z(k+z)/(k(k-z))$, where $k = \sqrt{2} + 1$. The following theorem yields \mathcal{S}_R^* radii.

Theorem 3.8. The \mathcal{S}_R^* radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:

- (i) For the class $H_{b,c}^1$, the sharp \mathcal{S}_R^* radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where
- $$\begin{aligned} x_1(r) &= (2\sqrt{2} - 3) + 2(\sqrt{2} - 1)(d + q + s)r + (6 + 4\sqrt{2} + (2\sqrt{2} - 1)(qs + d(q + s)))r^2 + \\ &+ 2((4 + \sqrt{2})(q + s) + d(4 + \sqrt{2} + \sqrt{2}qs))r^3 + 8(3 + qs + d(q + s))r^4 - 2((-7 + \sqrt{2})(q + \\ &+ s) + d(-7 + \sqrt{2} + (-3 + \sqrt{2})qs))r^5 + (18 - 4\sqrt{2} + (5 - 2\sqrt{2})(qs + d(q + s)))r^6 - \\ &- 2((-2 + \sqrt{2})(d + q + s))r^7 + (3 - 2\sqrt{2})r^8. \end{aligned}$$

(ii) For the class $H_{b,c}^2$, the \mathcal{S}_R^* radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where

$$x_2(r) = (2\sqrt{2} - 3) + 2(\sqrt{2} - 1)(m + n + q)r + (7 + 2\sqrt{2} + (2\sqrt{2} - 1)(nq + m(n + q)))r^2 + (8(m + q) + 2n(4 + \sqrt{2} + \sqrt{2}mq))r^3 + (15 - 2\sqrt{2} + 7mq - 2\sqrt{2}mq + 8n(m + q))r^4 - 2((-7 + \sqrt{2})n + (-3 + \sqrt{2})q + (-3 + \sqrt{2})m(1 + nq))r^5 + (5 - 2\sqrt{2})(1 + n(m + q))r^6 - 2((-2 + \sqrt{2})n)r^7.$$

(iii) For the class H_b^3 , the sharp \mathcal{S}_R^* radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where

$$x_3(r) = (2\sqrt{2} - 3) + 2(\sqrt{2} - 1)(l + q)r + (5 + 2\sqrt{2} + (-1 + 2\sqrt{2})lq)r^2 + 6(l + q)r^3 + (11 - 2\sqrt{2} + (5 - 2\sqrt{2})lq)r^4 - 2((-2 + \sqrt{2})(l + q))r^5 + (3 - 2\sqrt{2})r^6.$$

Proof. (i) Note that $x_1(0) = 2\sqrt{2} - 3 < 0$ and $x_1(1) = 6(2 + d)(2 + q)(2 + s) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. For $2(\sqrt{2} - 1) < C \leq \sqrt{2}$, Kumar *et al.*[16] proved that

$$(47) \quad \{w \in \mathbb{C} : |w - C| < C - 2(\sqrt{2} - 1)\} \subseteq \psi(\Delta).$$

As the centre of the disc in (21) is 1, by (47) $f \in \mathcal{S}_{\zeta}^*$ if

$$(48) \quad \frac{r}{(1 - r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq 3 - 2\sqrt{2}$$

which is equivalent to $f \in \mathcal{S}_R^*$ if $x_1(r) \leq 0$. Since $x_1(0) < 0$ and ρ_1 is the smallest root of the equation $x_1(r) = 0$, $x_1(r)$ is an increasing function on $(0, \rho_1)$. In view of this $f \in \mathcal{S}_R^*$ for $|z| = r \leq \rho_1$. For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (48), the function $F_1(z)$ defined for the class $H_{b,c}^1$ in (8) at $z = -\rho_1$, satisfies the following equality

$$\begin{aligned} \left| \frac{zF_1'(z)}{F_1(z)} \right| &= \left| 1 - \frac{\rho_1}{(1 - \rho_1^2)} \left(\frac{u\rho_1^2 + 4\rho_1 + u}{\rho_1^2 + u\rho_1 + 1} + \frac{v\rho_1^2 + 4\rho_1 + v}{\rho_1^2 + v\rho_1 + 1} + \frac{q\rho_1^2 + 4\rho_1 + q}{\rho_1^2 + q\rho_1 + 1} \right) \right| \\ &= 2(\sqrt{2} - 1) = \psi(-1). \end{aligned}$$

Thus, the radius is sharp.

(ii) A calculation shows that $x_2(0) = 2\sqrt{2} - 3 < 0$ and $x_2(1) = 6(2 + m)(l + n)(2 + q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_2(r) = 0$. Let $\rho_2 \in (0, 1)$ be the smallest root of the equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. In view of (47) and the fact that the centre of the disc in (13) is 1, $f \in \mathcal{S}_R^*$ if

$$\frac{r}{(1 - r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq 3 - 2\sqrt{2}$$

which is equivalent to $f \in \mathcal{S}_R^*$ if $x_2(r) \leq 0$. Since $x_2(0) < 0$ and ρ_2 is the smallest root of the equation $x_2(r) = 0$, $x_2(r)$ is an increasing function on $(0, \rho_2)$. Thus, $f \in \mathcal{S}_R^*$ for $|z| = r \leq \rho_2$.

- (iii) It is easy to see that $x_3(0) = 2\sqrt{2} - 3 < 0$ and $x_3(1) = 4(2+l)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$. Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. Since the centre of the disc in (29) is 1, by (47), $f \in \mathcal{S}_R^*$ if

$$(49) \quad \frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq 3 - 2\sqrt{2}$$

which is equivalent to $f \in \mathcal{S}_R^*$ if $x_3(r) \leq 0$. Since $x_3(0) < 0$ and ρ_3 is the smallest root of the equation $x_3(r) = 0$, $x_3(r)$ is an increasing function on $(0, \rho_3)$. This proves that $f \in \mathcal{S}_R^*$ for $|z| = r \leq \rho_3$.

The result is sharp for the function F_3 defined for the class H_b^3 in (20). At $z = -\rho_3$, and for $u = 4b - q \geq 0$, it follows from (49) that

$$\left| \frac{zF_3'(z)}{F_3(z)} \right| = \left| 1 - \frac{\rho_3}{1-\rho_3^2} \left(\frac{u\rho_3^2 + 4\rho_3 + u}{\rho_3^2 + u\rho_3 + 1} + \frac{q\rho_3^2 + 4\rho_3 + q}{\rho_3^2 + q\rho_3 + 1} \right) \right| = \psi(-1). \quad \square$$

Remark 3.8. *Substituting $b = 1$ and $c = 1$ and $q = 2$ in Theorem 3.8, we get the result [18, Theorem 8, p. 15].*

In 2020, Wani and Swaminathan [40, Lemma 2.2] introduced the class $\mathcal{S}_{Ne}^* = \mathcal{S}^*(1 + z - (z^3/3))$ consisting of functions associated with a nephroid. So, a function $f \in \mathcal{S}_{Ne}^*$ if and only if zf'/f maps the open unit disc Δ onto the interior of a two cusped kidney shaped curve $\Omega_{Ne} := \{u + iv : ((u-1)^2 + v^2 - 4/9)^3 - 4v^2/3 < 0\}$.

Theorem 3.9. *The sharp \mathcal{S}_{Ne}^* radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:*

- (i) *For the class $H_{b,c}^1$, the sharp \mathcal{S}_{Ne}^* radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where*

$$x_1(r) = -2 + (d+q+s)r + 4(8+qs+d(q+s))r^2 + (31(q+s)+d(31+7qs))r^3 + 24(3+qs+d(q+s))r^4 + (35(q+s)+d(35+11qs))r^5 + 8(5+qs+d(q+s))r^6 + 5(d+q+s)r^7 + 2r^8.$$

(ii) *For the class $H_{b,c}^2$, the sharp \mathcal{S}_{Ne}^* radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where*

$$x_2(r) = -2 + (m+n+q)r + 4(7+nq+m(n+q))r^2 + (31n+24q+m(24+7nq))r^3 + (38+24mn+14mq+24nq)r^4 + (35n+11q+11m(1+nq))r^5 + 8(1+n(m+q))r^6 + 5nr^7.$$

(iii) *For the class H_b^3 , the sharp \mathcal{S}_{Ne}^* radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where*

$$x_3(r) = -2 + (2+l)r + (22+8l)r^2 + 18(2+l)r^3 + (26+16l)r^4 + 5(2+l)r^5 + 2r^6.$$

Proof. (i) Note that $x_1(0) = -2 < 0$ and $x_1(1) = 18(2+d)(2+q)(2+s) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. For $1 \leq C < 5/3$, Wani and Swaminathan [40, Lemma 2.2] had proved that

$$(50) \quad \{w \in \mathbb{C} : |w - C| < 5/3 - C\} \subseteq \Omega_{Ne}.$$

As the centre of the disc in (21) is 1, by (50), $f \in \mathcal{S}_{Ne}^*$ if

$$(51) \quad \frac{r}{(1-r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{2}{3}$$

which is equivalent to $f \in \mathcal{S}_{Ne}^*$ if $x_1(r) \leq 0$. Since $x_1(0) < 0$ and ρ_1 is the smallest root of the equation $x_1(r) = 0$, $x_1(r)$ is an increasing function on $(0, \rho_1)$. In view of this $f \in \mathcal{S}_{Ne}^*$ for $|z| = r \leq \rho_1$. For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (51), the function $f_1(z)$ defined for the class $H_{b,c}^1$ in (6) at $z = -\rho_1$, satisfies the following equality

$$\begin{aligned} \left| \frac{zf_1'(z)}{f_1(z)} \right| &= \left| 1 + \frac{\rho_1}{(1-\rho_1^2)} \left(\frac{u\rho_1^2 + 4\rho_1 + u}{\rho_1^2 + u\rho_1 + 1} + \frac{v\rho_1^2 + 4\rho_1 + v}{\rho_1^2 + v\rho_1 + 1} + \frac{q\rho_1^2 + 4\rho_1 + q}{\rho_1^2 + q\rho_1 + 1} \right) \right| \\ &= \frac{5}{3}. \end{aligned}$$

which belongs to the boundary of the region Ω_{Ne} . Thus, the radius is sharp.

- (ii) A calculation shows that $x_2(0) = -2 < 0$ and $x_2(1) = 18(2+m)(l+n)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_2(r) = 0$. Let $\rho_2 \in (0, 1)$ be the smallest root of the equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. In view of (50) and the fact that centre of the disc in (13) is 1, $f \in \mathcal{S}_{Ne}^*$ if

$$(52) \quad \frac{r}{(1-r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{2}{3}$$

which is equivalent to $f \in \mathcal{S}_{Ne}^*$ if $x_2(r) \leq 0$. Since $x_2(0) < 0$ and ρ_2 is the smallest root of the equation $x_2(r) = 0$, $x_2(r)$ is an increasing function on $(0, \rho_2)$. Thus, $f \in \mathcal{S}_{Ne}^*$ for $|z| = r \leq \rho_2$. To prove the sharpness, consider the function f_2 defined in (14). For $u = 5b - 3c \geq 0$, $v = 3c - q \geq 0$ and $z = -\rho_2$, it follows from (52) that

$$\begin{aligned} \left| \frac{zf_2'(z)}{f_2(z)} \right| &= \left| 1 + \frac{\rho_2}{(1-\rho_2^2)} \left(\frac{u\rho_2^2 + 4\rho_2 + u}{\rho_2^2 + u\rho_2 + 1} + \frac{v\rho_2^2 + 2\rho_2 + v}{v\rho_2 + 1} + \frac{q\rho_2^2 + 4\rho_2 + q}{\rho_2^2 + q\rho_2 + 1} \right) \right| \\ &= \frac{5}{3} \end{aligned}$$

which illustrates the sharpness.

- (iii) It is easy to see that $x_3(0) = -2 < 0$ and $x_3(1) = 12(2+l)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$. Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. Since the centre of the disc in (29) is 1, by (50), $f \in \mathcal{S}_{Ne}^*$ if

$$(53) \quad \frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{2}{3}$$

which is equivalent to $f \in \mathcal{S}_{Ne}^*$ if $x_3(r) \leq 0$. Since $x_3(0) < 0$ and ρ_3 is the smallest root of the equation $x_3(r) = 0$, $x_3(r)$ is an increasing function on $(0, \rho_3)$. This proves that $f \in \mathcal{S}_{Ne}^*$ for $|z| = r \leq \rho_3$.

The result is sharp for the function f_3 defined for the class H_b^3 in (19). At $z = -\rho_3$, and for $u = 4b - q \geq 0$, it follows from (53) that

$$\left| \frac{zf_3'(z)}{f_3(z)} \right| = \left| 1 - \frac{\rho_3}{1 - \rho_3^2} \left(\frac{u\rho_3^2 + 4\rho_3 + u}{\rho_3^2 + u\rho_3 + 1} + \frac{q\rho_3^2 + 4\rho_3 + q}{\rho_3^2 + q\rho_3 + 1} \right) \right| = \frac{5}{3}.$$

□

Remark 3.9. For $b = 1$ and $c = 1$ and $q = 2$, Theorem 3.9 reduces to the result [18, Theorem 10, p. 18].

In 2020, the class $\mathcal{S}_{SG}^* = \mathcal{S}^*(2/(1 + e^{-z}))$ that maps the open unit disc Δ onto a domain $\Delta_{SG} := \{w \in \mathbb{C} : |\log(w/(2 - w))| < 1\}$ was introduced by Goel and Kumar[11]. Some results for the class \mathcal{S}_{SG}^* can be seen in [36]. The following theorem gives the sharp \mathcal{S}_{SG}^* radii.

Theorem 3.10. The sharp \mathcal{S}_{SG}^* radii for the classes $H_{b,c}^1$, $H_{b,c}^2$ and H_b^3 are as follows:

- (i) For the class $H_{b,c}^1$, the sharp \mathcal{S}_{SG}^* radius $\rho_1 \in (0, 1)$ is the smallest root of the equation $x_1(r) = 0$, where

$$x_1(r) = (1 - e) + 2(d + q + s)r + (2(7 + 5e) + (3 + e)(qs + d(q + s)))r^2 + (2(6 + 5e)(q + s) + 2d(6 + 5e + (2 + e)qs))r^3 + 8(1 + e)(3 + qs + d(q + s))r^4 + 2((5 + 6e)(q + s) + d(5 + qs + 2e(3 + qs)))r^5 + (2(5 + 7e) + (1 + 3e)(qs + d(q + s)))r^6 + 2e(d + q + s)r^7 + (-1 + e)r^8.$$
- (ii) For the class $H_{b,c}^2$, the sharp \mathcal{S}_{SG}^* radius $\rho_2 \in (0, 1)$ is the smallest root of the equation $x_2(r) = 0$, where

$$x_2(r) = (1 - e) + 2(m + n + q)r + (11 + 9e + (3 + e)(nq + m(n + q)))r^2 + 2(4(1 + e)m + (6 + 5e)n + 4(1 + e)q + (2 + e)mnq)r^3 + (11 + 8mn + 3mq + 8nq + e(13 + 8mn + 5mq + 8nq))r^4 + 2((5 + 6e)n + q + 2eq + (1 + 2e)m(1 + nq))r^5 + (1 + 3e)(1 + n(m + q))r^6 + 2enr^7.$$
- (iii) For the class H_b^3 , the sharp \mathcal{S}_{SG}^* radius $\rho_3 \in (0, 1)$ is the smallest root of the equation $x_3(r) = 0$, where

$$x_3(r) = (1 - e) + 2(l + q)r + (9 + 7e + (3 + e)lq)r^2 + 6(1 + e)(l + q)r^3 + (7 + lq + 3e(3 + lq))r^4 + 2e(l + q)r^5 + (-1 + e)r^6.$$

Proof. (i) Note that $x_1(0) = 1 - e < 0$ and $x_1(1) = 6(1 + e)(2 + d)(2 + q)(2 + s) > 0$ and thus in view of the intermediate value theorem, there exists a root of the equation $x_1(r) = 0$ in the interval $(0, 1)$. Let $r = \rho_1 \in (0, 1)$ be the smallest root of the equation $x_1(r) = 0$. For $2/(1 + e) < C < 2e/(1 + e)$, Goel and Kumar[11] proved the following inclusion:

$$(54) \quad \{w \in \mathbb{C} : |w - C| < r_{SG}\} \subset \Delta_{SG}, \text{ where } r_{SG} = \left(\frac{e - 1}{e + 1} \right) - |C - 1|.$$

As the centre of the disc in (21) is 1, by (54), $f \in \mathcal{S}_{SG}^*$ if

$$(55) \quad \frac{r}{(1 - r^2)} \left(\frac{dr^2 + 4r + d}{r^2 + dr + 1} + \frac{sr^2 + 4r + s}{r^2 + sr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{e - 1}{e + 1}$$

which is equivalent to $f \in \mathcal{S}_{SG}^*$ if $x_1(r) \leq 0$. Since $x_1(0) < 0$ and ρ_1 is the smallest root of the equation $x_1(r) = 0$, $x_1(r)$ is an increasing function on $(0, \rho_1)$. In view of this $f \in \mathcal{S}_{SG}^*$ for $|z| = r \leq \rho_1$. For $u = 6b - 4c \geq 0$, $v = 4c - q \geq 0$, using (55), the function $f_1(z)$ defined for the class $H_{b,c}^1$ in (6) at $z = -\rho_1$, satisfies the following equality

$$(56) \quad \left| \log \left(\frac{\frac{zf_1'(z)}{f_1(z)}}{2 - \frac{zf_1'(z)}{f_1(z)}} \right) \right| = \left| \log \left(\frac{1 + \frac{e-1}{e+1}}{2 - \left(1 + \frac{e-1}{e+1}\right)} \right) \right| = |\log(e)| = 1.$$

Thus, the radius is sharp.

- (ii) A calculation shows that $x_2(0) = 1 - e < 0$ and $x_2(1) = 6(1+e)(2+m)(l+n)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_2(r) = 0$. Let $\rho_2 \in (0, 1)$ be the smallest root of the equation $x_2(r) = 0$ and $f \in H_{b,c}^2$. In view of (54) and the fact that the centre of the disc in (13) is 1, $f \in \mathcal{S}_{SG}^*$ if

$$(57) \quad \frac{r}{(1-r^2)} \left(\frac{mr^2 + 4r + m}{r^2 + mr + 1} + \frac{nr^2 + 2r + n}{nr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{e-1}{e+1}$$

which is equivalent to $f \in \mathcal{S}_{SG}^*$ if $x_2(r) \leq 0$. Since $x_2(0) < 0$ and ρ_2 is the smallest root of the equation $x_2(r) = 0$, $x_2(r)$ is an increasing function on $(0, \rho_2)$. Thus, $f \in \mathcal{S}_{SG}^*$ for $|z| = r \leq \rho_2$. To prove the sharpness, consider the function f_2 defined in (14). For $u = 5b - 3c \geq 0$, $v = 3c - q \geq 0$ and $z = -\rho_2$, the similar calculations as in (56) together with (57) yields the result is sharp.

- (iii) It is easy to see that $x_3(0) = 1 - e < 0$ and $x_3(1) = 4(1+e)(2+l)(2+q) > 0$. By intermediate value theorem, there exists a root $r \in (0, 1)$ of the equation $x_3(r) = 0$. Let $\rho_3 \in (0, 1)$ be the smallest root of the equation $x_3(r) = 0$. Since the centre of the disc in (29) is 1, by (54), $f \in \mathcal{S}_{SG}^*$ if

$$\frac{r}{1-r^2} \left(\frac{lr^2 + 4r + l}{r^2 + lr + 1} + \frac{qr^2 + 4r + q}{r^2 + qr + 1} \right) \leq \frac{e-1}{1+e}$$

which is equivalent to $f \in \mathcal{S}_{SG}^*$ if $x_3(r) \leq 0$. Since $x_3(0) < 0$ and ρ_3 is the smallest root of the equation $x_3(r) = 0$, $x_3(r)$ is an increasing function on $(0, \rho_3)$. This proves that $f \in \mathcal{S}_{SG}^*$ for $|z| = r \leq \rho_3$.

At $z = -\rho_3$, and for $u = 4b - q \geq 0$, a calculation as in part(i) shows that the result is sharp for the function F_3 defined for the class H_b^3 in (20)

□

Remark 3.10. Putting $b = 1$ and $c = 1$ and $q = 2$ in Theorem 3.10, we obtain the result [18, Theorem 11, p. 19].

DECLARATION

The authors declare that they have no conflict of interest.

REFERENCES

- [1] R. M. Ali, N. E. Cho, N. K. Jain and V. Ravichandran, Radii of starlikeness and convexity for functions with fixed second coefficient defined by subordination, *Filomat* **26** (2012), no. 3, 553–561.
- [2] R. M. Ali, N. K. Jain and V. Ravichandran, Radii of starlikeness associated with the lemniscate of Bernoulli and the left-half plane, *Appl. Math. Comput.* **218** (2012), no. 11, 6557–6565.
- [3] R. M. Ali, N. K. Jain and V. Ravichandran, On the radius constants for classes of analytic functions, *Bull. Malays. Math. Sci. Soc. (2)* **36** (2013), no. 1, 23–38.
- [4] R. M. Ali, V. Kumar, V. Ravichandran and S. S. Kumar, Radius of starlikeness for analytic functions with fixed second coefficient, *Kyungpook Math. J.* **57** (2017), no. 3, 473–492.
- [5] R. M. Ali, S. Nagpal and V. Ravichandran, Second-order differential subordination for analytic functions with fixed initial coefficient, *Bull. Malays. Math. Sci. Soc. (2)* **34** (2011), no. 3, 611–629.
- [6] R. M. Ali, and V. Ravichandran, Uniformly convex and uniformly starlike functions, *Ramanujan Mathematics Newsletter* **21** (2011), no. 1, 16–30.
- [7] S. Anand, N. K. Jain and S. Kumar, Certain estimates of normalized analytic functions, *Math. Slovaca* **72** (2022), no. 1, 85–102.
- [8] N. E. Cho, V. Kumar, S. S. Kumar and V. Ravichandran, Radius problems for starlike functions associated with the sine function, *Bull. Iranian Math. Soc.* **45** (2019), no. 1, 213–232.
- [9] S. Gandhi and V. Ravichandran, Starlike functions associated with a lune, *Asian-Eur. J. Math.* **10** (2017), no. 4, 1750064, 12 pp.
- [10] A. Gangadharan, V. Ravichandran and T. N. Shanmugam, Radii of convexity and strong starlikeness for some classes of analytic functions, *J. Math. Anal. Appl.* **211** (1997), no. 1, 301–313.
- [11] P. Goel and S. Sivaprasad Kumar, Certain class of starlike functions associated with modified sigmoid function, *Bull. Malays. Math. Sci. Soc.* **43** (2020), no. 1, 957–991.
- [12] A. W. Goodman, *Univalent functions. Vol. II*, Mariner Publishing Co., Inc., Tampa, FL, 1983.
- [13] T. H. Gronwall, On the distortion in conformal mapping when the second coefficient in the mapping function has an assigned value, *Nat. Acad. Proc.* **6** (1920), 300–302.
- [14] T. Hayami and S. Owa, Hankel determinant for p -valently starlike and convex functions of order α , *Gen. Math.* **17** (2009), no. 4, 29–44.
- [15] W. Janowski, Some extremal problems for certain families of analytic functions. I, *Ann. Polon. Math.* **28** (1973), 297–326.
- [16] S. Kumar and V. Ravichandran, A subclass of starlike functions associated with a rational function, *Southeast Asian Bull. Math.* **40** (2016), no. 2, 199–212.
- [17] S. Kumar, V. Ravichandran and S. Verma, Initial coefficients of starlike functions with real coefficients, *Bull. Iranian Math. Soc.* **43** (2017), no. 6, 1837–1854.
- [18] A. Lecko, V. Ravichandran and A. Sebastian, Starlikeness of certain non-univalent functions, *Anal. Math. Phys.* **11** (2021), no. 4, Paper No. 163, 23 pp.
- [19] S. K. Lee, V. Ravichandran and S. Supramaniam, Applications of differential subordination for functions with fixed second coefficient to geometric function theory, *Tamsui Oxf. J. Inf. Math. Sci.* **29** (2013), no. 2, 267–284.
- [20] W. C. Ma and D. Minda, A unified treatment of some special classes of univalent functions, in *Proceedings of the Conference on Complex Analysis (Tianjin, 1992)*, 157–169, Conf. Proc. Lecture Notes Anal., I, Int. Press, Cambridge, MA.
- [21] W. C. Ma and D. Minda, Uniformly convex functions, *Ann. Polon. Math.* **57** (1992), no. 2, 165–175.
- [22] T. H. MacGregor, The radius of convexity for starlike functions of order $\frac{1}{2}$, *Proc. Amer. Math. Soc.* **14** (1963), 71–76.
- [23] T. H. MacGregor, The radius of univalence of certain analytic functions, *Proc. Amer. Math. Soc.* **14** (1963), 514–520.
- [24] T. H. MacGregor, A class of univalent functions, *Proc. Amer. Math. Soc.* **15** (1964), 311–317.

- [25] V. Madaan, A. Kumar and V. Ravichandran, Starlikeness associated with lemniscate of Bernoulli, *Filomat* **33** (2019), no. 7, 1937–1955.
- [26] C. P. McCarty, Two radius of convexity problems, *Proc. Amer. Math. Soc.* **42** (1974), 153–160.
- [27] C. P. McCarty, Functions with real part greater than α , *Proc. Amer. Math. Soc.* **35** (1972), 211–216.
- [28] R. Mendiratta, S. Nagpal and V. Ravichandran, On a subclass of strongly starlike functions associated with exponential function, *Bull. Malays. Math. Sci. Soc.* **38** (2015), no. 1, 365–386.
- [29] Z. Nehari, *Conformal mapping*, McGraw-Hill Book Co., Inc., New York, 1952.
- [30] R. K. Raina and J. Sokół, Some properties related to a certain class of starlike functions, *C. R. Math. Acad. Sci. Paris* **353** (2015), no. 11, 973–978.
- [31] J. S. Ratti, The radius of convexity of certain analytic functions, *Indian J. Pure Appl. Math.* **1** (1970), no. 1, 30–36.
- [32] J. S. Ratti, The radius of univalence of certain analytic functions, *Math. Z.* **107** (1968), 241–248.
- [33] V. Ravichandran, F. Rønning and T. N. Shanmugam, Radius of convexity and radius of starlikeness for some classes of analytic functions, *Complex Variables Theory Appl.* **33** (1997), no. 1-4, 265–280.
- [34] T. N. Shanmugam and V. Ravichandran, Certain properties of uniformly convex functions, in *Computational methods and function theory 1994 (Penang)*, Ser. Approx. Decompos., 5, World Sci. Publ., River Edge, NJ, 1995, pp. 319–324.
- [35] K. Sharma, N. K. Jain and V. Ravichandran, Starlike functions associated with a cardioid, *Afr. Mat.* **27** (2016), no. 5-6, 923–939.
- [36] M. Sharma, S. Kumar and N. K. Jain, Differential subordinations for functions with positive real part using admissibility conditions, *Asian-Eur. J. Math.* **15** (2022), no. 4, Paper No. 2250066, pp 22.
- [37] J. Sokół and J. Stankiewicz, Radius of convexity of some subclasses of strongly starlike functions, *Zeszyty Nauk. Politech. Rzeszowskiej Mat. No. 19* (1996), 101–105.
- [38] H. M. Srivastava, M. Tahir, B. Khan, Q. Z. Ahmad and N. Khan, Some general families of q -starlike functions associated with the Janowski functions, *Filomat* **33** (2019), no. 9, 2613–2626.
- [39] H. M. Srivastava and S. Owa (eds.), *Current Topics in Analytic Function Theory*, World Scientific Publishing Co., Inc., River Edge, NJ, 1992.
- [40] L. A. Wani and A. Swaminathan, Starlike and convex functions associated with a nephroid domain, *Bull. Malays. Math. Sci. Soc.* **44** (2021), no. 1, 79–104.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI, DELHI-110 007, INDIA

Email address: shalinirana3010@gmail.com

DEPARTMENT OF MATHEMATICAL SCIENCES,, KENT STATE UNIVERSITY, OHIO, 44021, U.S.A

Email address: oahuja@kent.edu

DEPARTMENT OF MATHEMATICS, ARYABHATTA COLLEGE, DELHI, 110021, INDIA

Email address: naveenjain@aryabhatacollege.ac.in