

Water UV-shielding in the terrestrial planet-forming zone: Implications for Oxygen-18 Isotope Anomalies in H₂¹⁸O Infrared Emission and Meteorites

JENNY K. CALAHAN,¹ EDWIN A. BERGIN,¹ AND ARTHUR D. BOSMAN¹

¹University of Michigan, 323 West Hall, 1085 South University Avenue, Ann Arbor, MI 48109, USA

Submitted to AJL

ABSTRACT

An understanding of abundance and distribution of water vapor in the innermost region of protoplanetary disks is key to understanding the origin of habitable worlds and planetary systems. Past observations have shown H₂O to be abundant and a major carrier of elemental oxygen in disk surface layers that lie within the inner few au of the disk. The combination of high abundance and strong radiative transitions leads to emission lines that are optically thick across the infrared spectral range. Its rarer isotopologue H₂¹⁸O traces deeper into this layer and will trace the full content of the planet forming zone. In this work, we explore the relative distribution of H₂¹⁶O and H₂¹⁸O within a model that includes water self-shielding from the destructive effects of ultraviolet radiation. In this Letter we show that there is an enhancement in the relative H₂¹⁸O abundance high up in the warm molecular layer within 0.1-10 au due to self-shielding of CO, C¹⁸O, and H₂O. Most transitions of H₂¹⁸O that can be observed with *JWST* will partially emit from this layer, making it essential to take into account how H₂O self-shielding may effect the H₂O to H₂¹⁸O ratio. Additionally, this reservoir of H₂¹⁸O-enriched gas in combination with the vertical “cold finger” effect might provide a natural mechanism to account for oxygen isotopic anomalies found in meteoritic material in the solar system.

Keywords: protoplanetary disk, astrochemistry

1. INTRODUCTION

It is expected that the vast majority of solar-type stars host an exoplanet within 1 au (Johnson et al. 2010; Mulders et al. 2018). As a result, it can be assumed that the inner gaseous region of protoplanetary disks must be a hospitable environment for active planet formation. Typical temperatures place nearly all primary volatile carriers of carbon, nitrogen, and oxygen into the gas within surface layers providing a rich environment for astronomical observations and constraint (Pontoppidan et al. 2014). Volatiles (i.e. CO, H₂O, CO₂) are potential starting points for the creation of complex molecules and the ice-coatings of pre-planetary pebbles (Wang et al. 2005; Gundlach & Blum 2015). These simple pre-cursors have been widely observed towards gas-rich disks within the 0.1-10 au region around young stars using the *Spitzer* space telescope (i.e. Carr & Najita 2008; Pontoppidan et al. 2010; Salyk et al. 2011a). Mapping the spatial extent and abundance of volatile molecules within the inner disk is essential knowledge used to piece together

the connection between the chemical reservoir of a gas-rich disk and the resulting planets.

The *JWST* mission will provide improved spectral resolution access to the inner disk region (*JWST*-*MIRI*: $\lambda/\Delta\lambda = 2000 - 3000$; Rieke et al. 2015), as compared to *Spitzer*'s InfraRed Spectrograph ($\lambda/\Delta\lambda = 600$; Houck et al. 2004). Nearly fifty disks will be targeted in the first set of *JWST* observations and will be used to enhance the study of emissions from volatiles in the terrestrial planet-forming zone. One molecule of high scientific impact is H₂O. Based on previous observations and theoretical work, it is expected that H₂O will be in high enough abundance such that its emission is optically thick (Carr & Najita 2008). The lesser abundant isotopologue H₂¹⁸O is also observable with *JWST* instruments, and, with reduced optical depth for its emission lines, will be used to infer the abundance and distribution of H₂O using the ratio between ¹⁶O and ¹⁸O.

The most frequently used value for ¹⁶O/¹⁸O is 550. This value comes from Wilson (1999), and is the average

observed ratio for five local ISM sources. This ratio has been seen to vary across the Galaxy following

$$(^{16}\text{O}/^{18}\text{O}) = (58.8 \pm 11.8)D_{\text{GC}} + (37.1 \pm 82.6) \quad (1)$$

where D_{GC} is the distance of a source from the Galactic center (Wilson & Rood 1994). Within the environment of a protoplanetary disk, however, there are chemical processes that can alter this ratio.

One such process is molecular self-shielding. Self-shielding is a process via which molecules, such as CO, N_2 , and H_2 , can protect themselves further from the source of radiation from the destructive effects of photodissociation from ultraviolet photons (van Dishoeck & Black 1988). These molecules are dissociated via a line process, as opposed to molecules with photodissociation cross-sections that are continuous at UV wavelengths (Heays et al. 2017). Thus, molecules closer to the radiation source can absorb UV photons allowing for the lines to become optically thick and, hence, self-shield molecules downstream from the source of UV photons. CO self-shielding can occur in gas-rich disks leading to a relative overabundance of $^{12}\text{C}^{16}\text{O}$ compared to $^{12}\text{C}^{18}\text{O}$ and $^{12}\text{C}^{17}\text{O}$ in the warm disk region (Miotello et al. 2014) because the absorbing lines of the lesser abundant isotopologues have reduced opacity. This process can help explain relatively high $\text{C}^{16}\text{O}/\text{C}^{18}\text{O}$ ratios that have been observed in a handful of protoplanetary disks (i.e. Brittain et al. 2005; Smith et al. 2009) and potentially might contribute to isotopic anomalies detected in asteroids and comets as compared to our Sun (Lyons & Young 2005; Nittler & Gaidos 2012; Altwegg et al. 2020).

Bethell & Bergin (2009) demonstrated that, because of fast formation rates in hot (>400 K) gas, water can also self-shield. Further, this shielding is unique in that it not only shields itself, but shields over a wide range of wavelength space, similar to ozone in the Earth’s atmosphere. Thus, water might also operate as a shield for other molecules, a process we call water UV-shielding. The combination of CO self-shielding and H_2O UV-shielding could produce an environment where the H_2O , H_2^{18}O , and CO are protected while C^{18}O continues to photodissociate. In this ^{18}O -rich environment, H_2^{18}O will form and continue to be protected, raising the relative abundance of H_2^{18}O with respect to H_2O in specific surface layers. This could have a strong impact on both the $\text{CO}/\text{C}^{18}\text{O}$ and $\text{H}_2\text{O}/\text{H}_2^{18}\text{O}$ values, altering them drastically from what is expected within the average ISM. These ratios will be commonly used in future *JWST* observations due to the expected high optical depth of CO and H_2O within the inner disk. Addition-

ally, the spatial extent of any ^{18}O -rich environment relative to where observed H_2^{18}O and H_2O lines emit from will have a strong effect on the determination of H_2O abundance within protoplanetary disks. Having accurate isotopic conversation factors could be critical as it is possible that the inner disk could betray evidence of volatile enhancements due to pebble drift (Ciesla & Cuzzi 2006; Banzatti et al. 2020).

One final consideration to the chemical make-up of the inner disk is Lyman- α ($\text{Ly-}\alpha$) radiation. The $\text{Ly-}\alpha$ transition often contains the vast majority of the energy within the FUV range if present in a stellar spectrum of T Tauri stars (Herczeg et al. 2002; Schindhelm et al. 2012). It is readily observed towards young stars that are actively accreting and possess a gas-rich disk. The $\text{Ly-}\alpha$ transition occurs at 1216 Å and can photodissociate water (van Dishoeck et al. 2006). $\text{Ly-}\alpha$ is not often taken into account in radiation transfer codes, due to its added complexity of not only scattering off of dust but also atomic Hydrogen, causing the $\text{Ly-}\alpha$ photons to scatter isotropically below the hot atomic layer of the disk (Bethell & Bergin 2011). To explore the strength that both water and CO self-shielding will have on the oxygen isotopic ratio, $\text{Ly-}\alpha$ needs to be accounted for.

This paper is a companion to Bosman et al. (2022a), Bosman et al. (2022b), and Duval et al. (2022, in prep.) Bosman et al. (2022a) sets up protoplanetary disk models focused on the innermost region which take into account water UV-shielding and efficient chemical heating processes. These two processes are required in order for a thermo-chemical calculation to reproduce water line emission and observed excitation temperatures from *Spitzer*. Using the same modeling setup, Bosman et al. (2022b) shows that CO_2 -to- H_2O ratios can also be achieved, strengthened by additional depletion of excess oxygen beyond the water snowline. Duval et al. (in prep) will focus on the impact of water UV-shielding on the chemical composition of the terrestrial planet forming region.

In this paper we will explore the abundances of both H_2O and H_2^{18}O in the inner most few au of protoplanetary disks where the excitation conditions are favorable for detection of rotational and vibrational emissions at mid-infrared wavelengths. This exploration will use the *Spitzer* observational legacy as a constraint (Pontoppidan et al. 2010; Salyk et al. 2011a), but look forward to *JWST* observations which will be observing systems where both $\text{Ly-}\alpha$ radiation and water UV-shielding will be present.

2. METHODS

2.1. Thermo-chemical modeling: DALI

DALI (Dust And Lines) is a physical-chemical code that accounts for radiation transfer, thermal and chemical calculations throughout a disk (Bruderer et al. 2012; Bruderer 2013). It contains an isotope chemical network (Miotello et al. 2014) and accounts for molecular self-shielding. For this project, H₂O UV-shielding has been additionally accounted for, see Bosman et al. (2022a) for detail. We utilize a model derived from the AS 209 and corresponding stellar spectrum from Zhang et al. (2021), and explore the effect of H₂O UV-shielding and Ly- α radiation in flat vs. thick model and two levels of dust settling. We include the ¹⁸O isotopologue of every oxygen carrying species in our chemical network.

After an initial radiative transfer calculation, thermal balance, and chemical abundance distribution determination, we account for Ly- α radiation using another code, detailed in the following section. The results from the separate Ly- α radiative transfer calculation are combined with the initial disk radiation field as calculated by DALI. We then calculate a new thermal balance and chemical environment, containing the final distribution of H₂O and H₂¹⁸O.

Excitation data for both ortho- and para-H₂¹⁸O was compiled into a file identical to the format from the Leiden LAMDA database (Schöier et al. 2005).¹ Line excitation data was obtained via the HITRAN database (Gordon et al. 2022)² and collisional data from the H₂¹⁶O LAMDA file (Faure & Josselin 2008). This was used for ray-tracing calculations with DALI to determine which lines may be observable and from where in the disk they emit.

2.2. Inclusion of Lyman Alpha

The inclusion of the Ly- α is derived from a radiation transfer code described by Bethell & Bergin (2011). Briefly, this code calculates the transport of both FUV-continuum and Ly- α photons. FUV photons are solely affected by the dust distribution, while Ly- α is additionally affected by resonance line scattering. An initial distribution of H and H₂ is derived using the thermochemical calculation from DALI, and Ly- α propagates throughout the distribution of gas and dust, scattered by atomic H in the H-rich layer, and eventually scattered and absorbed by dust.

After this calculation, the Ly- α -affected radiation field is combined with the DALI radiation field. The effects of a separate stellar input spectrum are normalized out of the Ly- α radiative transfer results, and a depletion or enhancement factor across wavelengths covered by the

Ly- α line and its line wings is calculated. The original DALI radiation field contains Ly- α emission as it would exist if Ly- α -photons scattered normally off of small dust grains only, identically to continuum-UV photons.

3. RESULTS

3.1. H₂¹⁸O-Enhanced Region

We find an H₂¹⁸O-rich environment high up in the molecular region of the disk. Enhanced H₂¹⁸O exists in models that range in flaring angle and dust settling parameters. Four models were explored with two different values for scale height and dust settling, and these individual models are detailed in the Appendix of Bosman et al. (2022a). The following figures and discussion are based on the flattest and most highly settled model, which reproduce H₂O and CO₂ spectra simultaneously (see Bosman et al. 2022b). In this model, a region where H₂O/H₂¹⁸O dips to approximately 45% of the initial ISM ratio exists at a z/r of 0.16-0.2 at r=0.1-1 AU, as seen in Figure 1. The upper boundary of this layer corresponds to where CO becomes optically thick enough to self-shield, producing an environment where C¹⁸O continues to photodissociate while C¹⁶O does not, releasing free ¹⁸O. Much of the free ¹⁸O finds its way into water molecules, enhancing the H₂¹⁸O abundance, thus providing a lower ratio between H₂O and H₂¹⁸O (≈ 300). The most common ¹⁸O destruction mechanisms come in the form of interactions with H₂ to form ¹⁸OH and an extra H atom. ¹⁸OH can then interact with an H₂ atom to form H₂¹⁸O. The main cycle of ¹⁸O once the chemistry has reached equilibrium is shown in Figure 2. The ratio returns to an ISM level at z/r ≈ 0.16 when C¹⁸O starts to be shielded, cutting off the supply of free ¹⁸O. Near the same z/r, within 1 au, H₂O UV-shielding aids in shielding C¹⁸O to a point where its photodissociation rate is extremely low, many orders of magnitude lower than an environment where H₂O UV-shielding would be ignored.

Beyond ~ 1 AU, the water abundance drops, yet a non-ISM ratio still exists. CO self-shielding continues to dominate, continuing to produce a ¹⁸O enhanced environment, thus a continued low ratio of H₂O to H₂¹⁸O, however there exists very little water vapor in the gas phase beyond r=1 au. The H₂¹⁸O-enhanced region exists between a top-down vertical CO column of 10¹⁷ and 10¹⁹cm⁻², an H₂ column of 10¹⁹ and 10²²cm⁻², and below an H column of 10²¹cm⁻². This H₂¹⁸O-enhanced region co-exists with large temperature gradients. Within 1 au, the thermal range from the bottom of this layer to the top is 500-2,000 K. Beyond 1 au the gradient is larger, ranging from 100-4,000 K.

¹ <https://home.strw.leidenuniv.nl/~moldata/>

² <https://hitran.org/>




Figure 1. The H_2O , H_2^{18}O , CO , and C^{18}O abundance in units of $(\text{mol}/\text{cm}^{-3})/n_{\text{gas}}$ throughout the inner disk (left and center plots), and the ratio between H_2O , H_2^{18}O (top right) and CO and C^{18}O (bottom right). At $z/r \approx 0.2$ a unique region exists where H_2^{18}O is enhanced and C^{18}O is greatly depleted.




Figure 2. The evolution of ^{18}O in equilibrium chemistry in the H_2^{18}O -enhanced region. Most of the oxygen comes from the dark cloud material originating from molecules like H_2O , CO_2 , organics and silicates. In this region photodissociation of C^{18}O provides an elevated abundance of ^{18}O which reacts with H and H_2 to quickly form ^{18}OH and H_2^{18}O . Because C^{16}O is shielded, very little extra ^{16}O is included in this oxygen cycle, thus the relative increase in H_2^{18}O production.

We find that most $\text{Ly-}\alpha$ photons do not reach the H_2 -dominated zone of the disk, including where the H_2^{18}O -rich region exits. The $\text{Ly-}\alpha$ emission feature is 25 times less bright in the radiation field of the disk at the location of the H_2^{18}O -rich region than it would be if the transfer of $\text{Ly-}\alpha$ photons was disregarded. Despite the

decrease in overall UV flux, the H_2^{18}O -rich region exits over the same spatial extent of the disk compared to a model that does not account for $\text{Ly-}\alpha$ scattering off of hydrogen atoms. The limit at which scattered $\text{Ly-}\alpha$ photons have a decreasing contribution to the radiation field corresponds to a $z/r \approx 0.25$, or where the H-to-H_2 ratio is $\approx 10^{-5}$. While in this disk model $\text{Ly-}\alpha$ did not reach into the molecular disk, Bethell & Bergin (2011) found that the molecular disk can be enhanced with $\text{Ly-}\alpha$ relative to FUV photons using the detailed disk physical models by D'Alessio et al. (2006). In this study, we use the thermo-chemical calculation within DALI and a chemical network to determine the H-fraction throughout the disk, while Bethell & Bergin (2011) used an analytical expression taking into account photodissociation and self-shielding processes while leaving out chemical destruction or creation pathways for H_2 and H . This results in a H-to-H_2 transition within the work that exists higher in the disk as compared to Bethell & Bergin (2011). Additionally, in Bethell & Bergin (2011) the H-fraction drops to near zero while H_2 is dominant while in DALI there exists a population of free H atoms in the H_2 -dominated region of the disk due to destructive chemical reactions. Thus, in the DALI disk, there are scattering events deep in the disk where in Bethell & Bergin (2011) there were none. The additional scattering events greatly increase the chance for $\text{Ly-}\alpha$ photons to be absorbed by dust.

Table 1. Transitions of H₂¹⁸O Observable with JWST

Transition	Wavelength (μm)	E _{up} (K)	A (s ⁻¹)	Weighted ¹⁶ O/ ¹⁸ O Average
10 _(3,8) - 9 _(0,9)	19.08	2072.3	0.83	520
13 _(5,8) - 12 _(4,9)	19.78	3772.5	8.9	500
10 _(4,7) - 9 _(1,8)	20.01	2265.3	1.9	510
11 _(4,7) - 10 _(3,8)	22.03	2725.3	4.4	490
9 _(7,2) - 9 _(4,5)	22.77	2581.7	0.031	540
9 _(9,0) - 8 _(8,1)	23.17	3165.9	41.	420
9 _(6,3) - 8 _(5,4)	26.83	2329.5	15.	410
8 _(7,2) - 7 _(6,1)	26.99	2265.6	21.	400

NOTE—We quote each weighted average of the ¹⁶O-to-¹⁸O ratio with two significant figures following the level of certainty around the ISM value (assumed to be 550).

3.2. H₂¹⁸O Emission Spectrum

The bulk of the water content exists within a radius of 1 au. We use DALI to calculate the flux contribution per cell for each emitted line. In order to determine which water lines will be of high interest, we run a quick non-LTE ray tracing program (Bosman et al. 2017, Appendix B) that estimates the flux of water lines over the full extent of the MIRI wavelength range. We find that IR bright H₂O lines emit high up in the disk, proving to be optically thick across all IR wavelengths (see Bosman et al. 2022 in prep for detail). Isolated and bright H₂¹⁸O emission lines are rare, but a handful of lines will be accessible and distinguishable from H₂O with the MIRI instrument. We find eight distinct lines listed in Table 1 and highlighted in Figure 3. These lines exist between 19-27 μm ; notably a region in MIRI with a lower sensitivity compared to shorter wavelength ranges.

Each H₂¹⁸O line emits from distinct heights within the disk, and may partially emit from the H₂¹⁸O-enhanced region, thus when converting from H₂¹⁸O-emission to a H₂O abundance, a ratio below that of the ISM should be used. One of the brightest and most isolated lines is the H₂¹⁸O 8_(7,2) - 7_(6,1) (27 μm) transition. We find that this transition starts to become optically thick and emits primarily in the enriched ¹⁸O region. We find this to be true for seven out of the eight identified lines. The H₂¹⁸O 9_(7,2) - 9_(4,5) transition (22.77 μm) appears to primarily emit below the H₂¹⁸O-enriched region, thus if using this line to determine a H₂O abundance, the conversion factor will be closer to ISM value of 550.

4. DISCUSSION & ANALYSIS

4.1. Measuring H₂O abundance using H₂¹⁸O observations

A weighted ratio between H₂O and H₂¹⁸O must be used in order to extrapolate to an H₂O abundance and distribution using H₂¹⁸O observations. We have identified eight transitions of H₂¹⁸O that are observable with JWST, and calculate the weighted ratio of H₂O to H₂¹⁸O that should be assumed based on our AS 209 model, and the vertical layers where we predict the emission arises. We compute weighted averages of H₂O/H₂¹⁸O using the relative spatial contribution from where each line is emitting from, as calculated by DALI.

Our first example comes from the H₂¹⁸O 8_(7,2) - 7_(6,1) transition at 27 μm . As seen in Figure 4, much of the emission from this line originates from the H₂¹⁸O enhanced region. We use the following equation to calculate a weighted H₂O/H₂¹⁸O value:

$$W = \frac{\sum_{i=1}^n w_i \chi_i}{\sum_{i=1}^n w_i} \quad (2)$$

where n is the number of cells in our model, w_i is the relative emission contribution in a cell, and χ_i is the H₂O/H₂¹⁸O value in a cell. Using this formalism, we find the average H₂O/H₂¹⁸O coming from this transition is ~ 400 . The H₂¹⁸O at 22.77 μm appears to emit from a drastically different region in the disk, thus its average H₂O/H₂¹⁸O is 540, nearly identical within the margin of error to the typically used ISM estimate. We list the calculated weighted averaged determined for the eight identified bright and isolated H₂¹⁸O transitions in Table 1. There is a general trend that transitions from smaller wavelengths trend towards a weighted average of 550. This trend arises from the Einstein A -coefficient associated with each transition. Transitions with lower A values are more likely to emit from deeper within the disk at higher gas densities, thus farther away from the H₂¹⁸O-enhanced region. To reduce the uncertainty in the H₂O/H₂¹⁸O ratio, transitions associated with low Einstein A -coefficients could be preferentially used. These differences are small and up to a factor of two. However, chemical studies will be looking for enhanced abundances of water vapor in this region, perhaps supplied by pebble drift (Banzatti et al. 2020). To isolate these chemical changes we need to correct for effects that can be understood; water UV-shielding is one of these effects.

4.2. Corresponding CO Infrared Observations

In the H₂¹⁸O-enhanced region, C¹⁸O abundance drops and the ¹²CO-to-C¹⁸O ratio increases by upwards of four




Figure 3. The H_2O and H_2^{18}O spectra as predicted from an AS 209 model including water UV-shielding and $\text{Ly-}\alpha$ contributions with 0.1 km s^{-1} velocity bins. Blue lines correspond to H_2O and dark purple with H_2^{18}O . H_2^{18}O lines that are relatively bright and isolated from H_2O lines are highlighted.




Figure 4. Top: The overlapping spectra from H_2O and H_2^{18}O focusing on a narrow wavelength regions within the MIRI instrument on JWST, focused on the H_2^{18}O $27 \mu\text{m}$ line (left) and $22.77 \mu\text{m}$ (right) lines. The bottom plots show the 50% (yellow contour) and 90% (white contour) contribution regions for each line plotted over the H_2O -to- H_2^{18}O ratio.

orders of magnitude. Measuring the ^{12}CO -to- C^{18}O ratio using lines of CO that emit from this region could help confirm the presence of this ^{18}O -rich region and even calibrate uncertainties in the true ratio between H_2O -to- H_2^{18}O . The R and P branches for multiple vibrational transitions of CO and its isotopologues peak at $\sim 5 \mu\text{m}$ and are observable from the ground using high spectral resolution instruments such as *Keck* NIRSPEC and *VLT*'s CRIFRES. The C^{18}O column densities in the H_2^{18}O -rich region range from 10^{12} - 10^{15} cm^{-2} and

are predicted to produce optically thin C^{18}O emission in the infrared. As summarized and expanded upon in [Banzatti et al. \(2022\)](#), over 100 T Tauri and Herbig disks have been observed with high-spectral resolution in the wavelength range of CO's vibrational lines which emit from $\sim 1000 \text{ K}$, corresponding to the temperature of the H_2^{18}O -enhanced region. In these studies only CO and ^{13}CO observations were reported, making it currently impossible to use these observations to corroborate our predicted ^{18}O -rich region within 1 au. The only work

that set out to measure $^{12}\text{CO}/\text{C}^{18}\text{O}$ and $^{12}\text{CO}/\text{C}^{17}\text{O}$ is found in [Smith et al. \(2015\)](#). This study observed nine YSO systems, three of which were optically thick disks. These three disks show tentative increase in the $^{12}\text{CO}/\text{C}^{18}\text{O}$ well above the ISM-measured value, however it was noted at significant systematic uncertainties are associated with these measurements. A future study dedicated to deep observations of multiple disk systems could aid in upcoming MIRI observations of these regions, and put constraints on the ^{18}O -enhancement within the radial extent of planet-formation.

4.3. Impact of a different $^{16}\text{O}/^{18}\text{O}$ Ratio

Mass-independent fractionation of oxygen isotopes, with enrichment of the heavy isotopes, has been isolated within meteoritic material ([Clayton et al. 1973](#); [Thiemens & Heidenreich 1983](#)). The origin of these isotopic anomalies requires a specific mechanism that creates an ^{18}O (and ^{17}O)-rich environment as compared to the natal stellar nebula and envelope ([Clayton 1993](#)). This fractionation must originate in the solar nebular disk or within the natal molecular cloud ([Yurimoto & Kuramoto 2004](#); [Lyons & Young 2005](#); [Lee et al. 2008](#)). CO self-shielding presents a mechanism to enrich gas in heavy isotopes, particularly on surfaces exposed to ultraviolet, as it selectively photodissociates C^{18}O and C^{17}O relative to CO producing gas enriched in ^{18}O and ^{17}O ([Lyons & Young 2005](#); [Miotello et al. 2014](#)). In this work, we find that the addition of H₂O UV-shielding not only continues the ^{18}O enrichment within the inner disk, but enhances it compared to results solely based on CO self-shielding. We find a depletion of C^{18}O approaching a factor of 10^3 high up in the disk between 0.1-30 au at a $\sim z/r=0.2-0.3$. This is a reservoir rich with ^{18}O , enhancing the ^{18}O isotopologues for many different volatile species, including water vapor and ice.

Vertical mixing may act as a mechanism to deliver the ^{18}O -enhanced molecules to the planet-forming midplane. The ‘‘cold finger’’ effect ([Stevenson & Lunine 1988](#); [Meijerink et al. 2009](#)) has been proposed a mechanism that will transport water and other organics efficiently from the atmosphere of the disk to the midplane. Due to vertical mixing, water from the gaseous atmosphere can transport down to the midplane where it locked onto grains as ice. The cold finger effect has been used to reconcile models in which high water abundances are predicted, yet corresponding sub-mm observations exhibit non-detections ([Du et al. 2015](#); [Salyk et al. 2011b](#); [Carr et al. 2018](#); [Bosman & Bergin 2021](#); [Bosman et al. 2022b](#)). Thus, while the H₂¹⁸O-rich regions resides high up in the disk atmosphere, vertical

mixing can bring a portion of this reservoir down to the midplane and enrich meteoritic precursors.

[Lyons & Young \(2005\)](#) explored a solution in the outer tens of au that suggested that meteoritic ^{18}O enrichment could be the result of CO self-shielding. In this model, oxygen atoms (enriched with heavy isotopes) produced via isotopic-selective CO self-shielding, are mixed downwards forming water ice via grain surface chemistry. These ices will need to be transported to the inner few au, perhaps by drift ([Ciesla & Cuzzi 2006](#)). However, current meteoritic evidence suggests the inner solar system is chemically separated from the outer ($\gtrsim 5$ au) parts of the disk between 1 Myr and 3-4 Myr, potentially due to early formation of Jupiter’s core ([Kruijer et al. 2017](#)). Our model shows that this vertical layer of enhanced ^{18}O is placed directly into water in surface layers above the inner disk spatially closer to the region where meteoritic progenitors originate.

5. CONCLUSION

In this Letter, we implement H₂O UV-shielding and chemical heating in addition to CO self-shielding within a gas-rich disk environment. Focusing on the innermost region of the disk, a water vapor-rich reservoir exits within 1 au, and a H₂¹⁸O enhanced region exists where many water IR lines approach a $\tau = 1$. In this region, the H₂O-to-H₂¹⁸O ratio approaches 300, a significantly lower value than the assume ISM ratio of 550. We then seek use this context to provide insight into future JWST observations of H₂¹⁸O as it will act as a tracer for the abundance and distribution of gaseous H₂O in the main planet formation zone.

1. In our disk model, the H₂O-to-H₂¹⁸O ratio approaches 300 between a $z/r \approx 0.16-0.2$ and inside 1 au due to CO self-shielding.
2. We identify eight bright and isolated H₂¹⁸O lines that can be used to access the abundance and distribution of H₂O. Transitions associated with higher Einstein A-coefficients, such as H₂¹⁸O $8_{(7,2)} - 7_{(6,1)}$ at 27 μm emit, primarily in the H₂¹⁸O-rich region, thus the ratio between H₂O and H₂¹⁸O are less than the local ISM value of 550. Seven out of the eight identified transitions emit predominately from the H₂¹⁸O-enhanced region, thus most observable H₂¹⁸O transitions with JWST will have an additional factor of uncertainty when converting H₂¹⁸O abundance to H₂O.
3. In this model Ly- α radiation does not penetrate down to the molecular-rich region of the disk in the case of our relatively thin and 0.0045 M_⊙ disk.

4. Vertical diffusive mixing, or the “cold finger” effect, could transport a significant amount of ^{18}O -rich molecules to the planet-forming midplane to be incorporated into pre-planetary materials that reside in the terrestrial planet-forming zone.

6. ACKNOWLEDGMENTS

Software: DALI, (Bruderer et al. 2012; Bruderer 2013) scipy (Virtanen et al. 2020), Matplotlib (Hunter 2007), Astropy (Astropy Collaboration

et al. 2013, 2018), NumPy (Harris et al. 2020), RADMC-3D (Dullemond et al. 2012), radmc3dPy: [HTTPS://BITBUCKET.ORG/AT_JUHASZ/RADMC3DPY/](https://bitbucket.org/at_juhasz/radmc3dpy/), parallel (Tange 2018)

J.K.C. acknowledges support from the National Science Foundation Graduate Research Fellowship under Grant No. DGE 1256260 and the National Aeronautics and Space Administration FINESST grant, under Grant no. 80NSSC19K1534.

A.D.B. and E.A.B. acknowledge support from NSF Grant#1907653 and NASA grant XRP 80NSSC20K0259

REFERENCES

- Altwegg, K., Balsiger, H., Combi, M., et al. 2020, MNRAS, 498, 5855, doi: [10.1093/mnras/staa2701](https://doi.org/10.1093/mnras/staa2701)
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)
- Banzatti, A., Pascucci, I., Bosman, A. D., et al. 2020, ApJ, 903, 124, doi: [10.3847/1538-4357/abbc1a](https://doi.org/10.3847/1538-4357/abbc1a)
- Banzatti, A., Abernathy, K. M., Brittain, S., et al. 2022, AJ, 163, 174, doi: [10.3847/1538-3881/ac52f0](https://doi.org/10.3847/1538-3881/ac52f0)
- Bethell, T., & Bergin, E. 2009, Science, 326, 1675, doi: [10.1126/science.1176879](https://doi.org/10.1126/science.1176879)
- Bethell, T. J., & Bergin, E. A. 2011, ApJ, 739, 78, doi: [10.1088/0004-637X/739/2/78](https://doi.org/10.1088/0004-637X/739/2/78)
- Bosman, A. B., Bergin, E. A., Calahan, J. K., & Duval, S. 2022a, ApJL, submitted, xxx, xxx, doi: [xx.xxxxx](https://doi.org/xx.xxxxx)
- . 2022b, ApJL, submitted, xxx, xxx, doi: [xx.xxxxx](https://doi.org/xx.xxxxx)
- Bosman, A. D., & Bergin, E. A. 2021, ApJL, 918, L10, doi: [10.3847/2041-8213/ac1db1](https://doi.org/10.3847/2041-8213/ac1db1)
- Bosman, A. D., Bruderer, S., & van Dishoeck, E. F. 2017, A&A, 601, A36, doi: [10.1051/0004-6361/201629946](https://doi.org/10.1051/0004-6361/201629946)
- Brittain, S. D., Rettig, T. W., Simon, T., & Kulesa, C. 2005, ApJ, 626, 283, doi: [10.1086/429310](https://doi.org/10.1086/429310)
- Bruderer, S. 2013, A&A, 559, A46, doi: [10.1051/0004-6361/201321171](https://doi.org/10.1051/0004-6361/201321171)
- Bruderer, S., van Dishoeck, E. F., Doty, S. D., & Herczeg, G. J. 2012, A&A, 541, A91, doi: [10.1051/0004-6361/201118218](https://doi.org/10.1051/0004-6361/201118218)
- Carr, J. S., & Najita, J. R. 2008, Science, 319, 1504, doi: [10.1126/science.1153807](https://doi.org/10.1126/science.1153807)
- Carr, J. S., Najita, J. R., & Salyk, C. 2018, Research Notes of the American Astronomical Society, 2, 169, doi: [10.3847/2515-5172/aadfe7](https://doi.org/10.3847/2515-5172/aadfe7)
- Ciesla, F. J., & Cuzzi, J. N. 2006, Icarus, 181, 178, doi: [10.1016/j.icarus.2005.11.009](https://doi.org/10.1016/j.icarus.2005.11.009)
- Clayton, R. N. 1993, Annual Review of Earth and Planetary Sciences, 21, 115, doi: [10.1146/annurev.ea.21.050193.000555](https://doi.org/10.1146/annurev.ea.21.050193.000555)
- Clayton, R. N., Grossman, L., & Mayeda, T. K. 1973, Science, 182, 485, doi: [10.1126/science.182.4111.485](https://doi.org/10.1126/science.182.4111.485)
- D’Alessio, P., Calvet, N., Hartmann, L., Franco-Hernández, R., & Servín, H. 2006, ApJ, 638, 314, doi: [10.1086/498861](https://doi.org/10.1086/498861)
- Du, F., Bergin, E. A., & Hogerheijde, M. R. 2015, ApJL, 807, L32, doi: [10.1088/2041-8205/807/2/L32](https://doi.org/10.1088/2041-8205/807/2/L32)
- Dullemond, C. P., Juhasz, A., Pohl, A., et al. 2012, RADMC-3D: A multi-purpose radiative transfer tool. <http://ascl.net/1202.015>
- Faure, A., & Josselin, E. 2008, A&A, 492, 257, doi: [10.1051/0004-6361:200810717](https://doi.org/10.1051/0004-6361:200810717)
- Gordon, I. E., Rothman, L. S., Hargreaves, R. J., et al. 2022, JQSRT, 277, 107949, doi: [10.1016/j.jqsrt.2021.107949](https://doi.org/10.1016/j.jqsrt.2021.107949)
- Gundlach, B., & Blum, J. 2015, ApJ, 798, 34, doi: [10.1088/0004-637X/798/1/34](https://doi.org/10.1088/0004-637X/798/1/34)
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, Nature, 585, 357, doi: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2)
- Heays, A. N., Bosman, A. D., & van Dishoeck, E. F. 2017, A&A, 602, A105, doi: [10.1051/0004-6361/201628742](https://doi.org/10.1051/0004-6361/201628742)
- Herczeg, G. J., Linsky, J. L., Valenti, J. A., Johns-Krull, C. M., & Wood, B. E. 2002, ApJ, 572, 310, doi: [10.1086/339731](https://doi.org/10.1086/339731)
- Houck, J. R., Roellig, T. L., Van Cleve, J., et al. 2004, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 5487, Optical, Infrared, and Millimeter Space Telescopes, ed. J. C. Mather, 62–76, doi: [10.1117/12.550517](https://doi.org/10.1117/12.550517)
- Hunter, J. D. 2007, Computing in Science & Engineering, 9, 90, doi: [10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)

- Johnson, J. A., Aller, K. M., Howard, A. W., & Crepp, J. R. 2010, *PASP*, 122, 905, doi: [10.1086/655775](https://doi.org/10.1086/655775)
- Kruijer, T. S., Burkhardt, C., Budde, G., & Kleine, T. 2017, *Proceedings of the National Academy of Science*, 114, 6712, doi: [10.1073/pnas.1704461114](https://doi.org/10.1073/pnas.1704461114)
- Lee, J.-E., Bergin, E. A., & Lyons, J. R. 2008, *M&PS*, 43, 1351, doi: [10.1111/j.1945-5100.2008.tb00702.x](https://doi.org/10.1111/j.1945-5100.2008.tb00702.x)
- Lyons, J. R., & Young, E. D. 2005, *Nature*, 435, 317, doi: [10.1038/nature03557](https://doi.org/10.1038/nature03557)
- Meijerink, R., Pontoppidan, K. M., Blake, G. A., Poelman, D. R., & Dullemond, C. P. 2009, *ApJ*, 704, 1471, doi: [10.1088/0004-637X/704/2/1471](https://doi.org/10.1088/0004-637X/704/2/1471)
- Miotello, A., Bruderer, S., & van Dishoeck, E. F. 2014, *A&A*, 572, A96, doi: [10.1051/0004-6361/201424712](https://doi.org/10.1051/0004-6361/201424712)
- Mulders, G. D., Pascucci, I., Apai, D., & Ciesla, F. J. 2018, *AJ*, 156, 24, doi: [10.3847/1538-3881/aac5ea](https://doi.org/10.3847/1538-3881/aac5ea)
- Nittler, L. R., & Gaidos, E. 2012, *M&PS*, 47, 2031, doi: [10.1111/j.1945-5100.2012.01410.x](https://doi.org/10.1111/j.1945-5100.2012.01410.x)
- Pontoppidan, K. M., Salyk, C., Bergin, E. A., et al. 2014, in *Protostars and Planets VI*, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning, 363, doi: [10.2458/azu_uapress_9780816531240-ch016](https://doi.org/10.2458/azu_uapress_9780816531240-ch016)
- Pontoppidan, K. M., Salyk, C., Blake, G. A., et al. 2010, *ApJ*, 720, 887, doi: [10.1088/0004-637X/720/1/887](https://doi.org/10.1088/0004-637X/720/1/887)
- Rieke, G. H., Wright, G. S., Böker, T., et al. 2015, *PASP*, 127, 584, doi: [10.1086/682252](https://doi.org/10.1086/682252)
- Salyk, C., Pontoppidan, K. M., Blake, G. A., Najita, J. R., & Carr, J. S. 2011a, *ApJ*, 731, 130, doi: [10.1088/0004-637X/731/2/130](https://doi.org/10.1088/0004-637X/731/2/130)
- . 2011b, *ApJ*, 731, 130, doi: [10.1088/0004-637X/731/2/130](https://doi.org/10.1088/0004-637X/731/2/130)
- Schindhelm, R., France, K., Herczeg, G. J., et al. 2012, *ApJL*, 756, L23, doi: [10.1088/2041-8205/756/1/L23](https://doi.org/10.1088/2041-8205/756/1/L23)
- Schöier, F. L., van der Tak, F. F. S., van Dishoeck, E. F., & Black, J. H. 2005, *A&A*, 432, 369, doi: [10.1051/0004-6361:20041729](https://doi.org/10.1051/0004-6361:20041729)
- Smith, R. L., Pontoppidan, K. M., Young, E. D., & Morris, M. R. 2015, *ApJ*, 813, 120, doi: [10.1088/0004-637X/813/2/120](https://doi.org/10.1088/0004-637X/813/2/120)
- Smith, R. L., Pontoppidan, K. M., Young, E. D., Morris, M. R., & van Dishoeck, E. F. 2009, *ApJ*, 701, 163, doi: [10.1088/0004-637X/701/1/163](https://doi.org/10.1088/0004-637X/701/1/163)
- Stevenson, D. J., & Lunine, J. I. 1988, *Icarus*, 75, 146, doi: [10.1016/0019-1035\(88\)90133-9](https://doi.org/10.1016/0019-1035(88)90133-9)
- Tange, O. 2018, *GNU Parallel 2018 (Ole Tange)*, doi: [10.5281/zenodo.1146014](https://doi.org/10.5281/zenodo.1146014)
- Thiemens, M. H., & Heidenreich, J. E., I. 1983, *Science*, 219, 1073, doi: [10.1126/science.219.4588.1073](https://doi.org/10.1126/science.219.4588.1073)
- van Dishoeck, E. F., & Black, J. H. 1988, *ApJ*, 334, 771, doi: [10.1086/166877](https://doi.org/10.1086/166877)
- van Dishoeck, E. F., Jonkheid, B., & van Hemert, M. C. 2006, *Faraday Discussions*, 133, 231, doi: [10.1039/b517564j](https://doi.org/10.1039/b517564j)
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature Methods*, doi: <https://doi.org/10.1038/s41592-019-0686-2>
- Wang, H., Bell, R. C., Iedema, M. J., Tsekouras, A. A., & Cowin, J. P. 2005, *ApJ*, 620, 1027, doi: [10.1086/427072](https://doi.org/10.1086/427072)
- Wilson, T. L. 1999, *Reports on Progress in Physics*, 62, 143, doi: [10.1088/0034-4885/62/2/002](https://doi.org/10.1088/0034-4885/62/2/002)
- Wilson, T. L., & Rood, R. 1994, *ARA&A*, 32, 191, doi: [10.1146/annurev.aa.32.090194.001203](https://doi.org/10.1146/annurev.aa.32.090194.001203)
- Yurimoto, H., & Kuramoto, K. 2004, *Science*, 305, 1763, doi: [10.1126/science.1100989](https://doi.org/10.1126/science.1100989)
- Zhang, K., Booth, A. S., Law, C. J., et al. 2021, *arXiv e-prints*, arXiv:2109.06233, <https://arxiv.org/abs/2109.06233>