

Extending density functional theory with near chemical accuracy beyond pure water

SUHWAN SONG^{a,b}, STEFAN VUCKOVIC^{c,d}, YOUNGSAM KIM^a, HAYOUNG YU^a,
EUNJI SIM^{a,*} AND KIERON BURKE^{b,e}

^aDepartment of Chemistry, Yonsei University, 50 Yonsei-ro Seodaemun-gu, Seoul 03722, Korea

^bDepartment of Chemistry, University of California, Irvine, CA 92697, USA

^cInstitute for Microelectronics and Microsystems (CNR-IMM), Via Monteroni, Campus Unisalento, 73100 Lecce, Italy

^dDepartment of Chemistry & Pharmaceutical Sciences and Amsterdam Institute of Molecular and Life Sciences (AIMMS),
Faculty of Science, Vrije Universiteit, De Boelelaan 1083, 1081HV Amsterdam, The Netherlands

^eDepartments of Physics & Astronomy, University of California, Irvine, CA 92697, USA

February 10, 2023

Abstract

Density functional simulations of condensed phase water are typically inaccurate, due to the inaccuracies of approximate functionals. A recent breakthrough showed that the SCAN approximation can yield chemical accuracy for pure water in all its phases, but only when its density is corrected. This is a crucial step toward first-principles biosimulations. However, weak dispersion forces are ubiquitous and play a key role in noncovalent interactions among biomolecules, but are not included in the new approach. Moreover, naïve inclusion of dispersion in HF-SCAN ruins its high accuracy for pure water. Systematic application of the principles of density-corrected DFT yields a functional (HF- r^2 SCAN-DC₄) which recovers and not only improves over HF-SCAN for pure water, but also captures vital noncovalent interactions in biomolecules, making it suitable for simulations of solutions.

I. INTRODUCTION

The overwhelming importance of simulating water: The properties of water, such as the uniqueness of its phase diagram, never stop surprising scientific communities.[1] Given the vital importance of water in fields that vary from material science to biology, there has been a recent surge in the development and competition of different electronic structure methods for simulating water.[2, 3, 4, 5, 6, 7, 8, 9] As *ab initio* quantum-chemical methods are too expensive for large systems, Kohn-Sham density functional theory (KS-DFT) has become a workhorse of electronic structure methods for running water calculations.[10, 11, 12, 13, 14] But, despite an excellent accuracy to cost ratio, historically KS-DFT has been unable to deliver sufficiently high accuracy in water simulations to reproduce experimental data.[15, 16, 17, 18]

A recent breakthrough in this direction by Dasgupta *et al.* showed that the strongly constrained and appropriately normed (SCAN) functional, when used in tandem with 'density-corrected DFT' (DC-DFT), is a game changer for water simulation, because it brings KS-DFT close to chemical accuracy.[3, 4] The role of water in a chemical or biochemical reaction goes beyond providing an environment to help a reaction in an aqueous solvation and is often explicitly involved in the mechanism. For this reason, a complete understanding of the reaction is possible only when the interaction between water and other molecules is accurately described. Figure 1 shows how an integratively designed DC-DFT procedure, HF- r^2 SCAN-DC₄, describes not only the interactions between water-water, water-organic molecules, and water-biochemical molecules

in various situations, but also the interactions of noncovalent complexes at chemical accuracy or better.

The importance of the density: DC-DFT is a general framework that separates errors of any DFT calculations into a contribution coming from the approximate "D" (density) and the 'true' error coming from the approximate "F" (functional).[19, 20, 21, 22] In addition to being a rigorous exact theory, DC-DFT gives practical guidance on when and how it can be used to reduce errors in DFT simulation.[23, 24, 25, 26] Standard DFT calculations are performed self-consistently (SC). The simplest form of practical DC-DFT is HF-DFT, where density functionals are evaluated instead on Hartree-Fock (HF) densities and orbitals.[27, 28, 29, 30, 31] While in most cases, SC-DFT gives the best answer, in some errors in specific cases SC-DFT suffers from large energetic errors due to the approximate density (density-driven errors).[19, 22] In such cases, HF-DFT typically yields significant improvements over SC-DFT, and these include a number of chemical domains (barrier heights, some torsional barriers, halogen bonds, anions, etc.).[26]

The importance of the functional: SCAN is a non-empirical meta-GGA functional designed to satisfy 17 exact physical constraints, and to recover several nonbonded 'norms'. [32] Meta-GGA's use the KS kinetic energy density as an ingredient, but are *not* hybrid functionals like B3LYP[33, 34, 35, 36], which include some fraction of exact exchange from a HF calculation.[35] In terms of accuracy, SCAN is often on par with highly empirical more expensive density functionals designed for molecules. At the same time, it enjoys great successes for simulations of extended systems, making it one of the most-used general-purpose functionals developed over the last 10 years.[37, 38, 39, 40, 41] Earlier works have shown that

*esim@yonsei.ac.kr




Figure 1: Performance of HF- r^2 SCAN-DC₄ relative to HF-SCAN for various chemical reactions: (a) the interaction energy of various configurations of the stacked cytosine dimer, where HF-SCAN underbinds by 2-3 kcal/mol; (b) energies of water hexamer relative to the lowest-lying prism isomer, with HF-SCAN underestimating by almost 1 kcal/mol; (c) errors in binding energy of WATER27 complexes as a function of density sensitivity (how much a DFT energy changes when the density is changed), showing how large errors can be without using the HF density. One cluster, $H_3O^+(H_2O)_6OH^-$ (at x close to 4 kcal/mol) is an outlier argued to exhibit a significant multiconfigurational character[4]; (d) relative energies of water 20-mer isomers (not density sensitive) from WATER27, where self-consistent SC- r^2 SCAN-D₄ performs best, but using the HF density introduces little error; (e) errors in interaction energies in the water \cdots aspirin dimer structures from an MD simulation at $T=298.15$ K; (f) mean-absolute-errors (MAEs) for intra- and inter-molecular noncovalent interactions datasets from the GMTKN55 database. For more details, see the main text and supporting information.

standard (SC) DFT calculations of water clusters suffer badly from density-driven errors, which explains why HF-SCAN is much more accurate than its SC counterpart for simulations of water.[3, 25] In addition to water clusters, Dasgupta *et al.* used HF-SCAN in tandem with many-body potential energy function related to the highly popular MB-pol[10, 11, 12] to run molecular dynamics (MD) simulations of liquid water and obtained results in excellent agreement with the experimental data. These were the first successful DFT-based simulations able to correctly describe the condensation of water.

Nevertheless, the convergence of the SCAN functional can be painfully slow with respect to the size of molecular grids, due to either the size of a system or because it would require grids larger than those available in most of the standard-quantum chemical codes.[31] Larger grids also lead to longer computational times. To address these issues of SCAN, Perdew and co-workers developed the regularized-restored SCAN functional (r^2 SCAN), which regularizes SCAN but restores SCAN's adherence to exact constraints.[42] But, as we show below, a standalone version of HF- r^2 SCAN is much less accurate for water simulations than HF-SCAN.

The vital importance of dispersion: Despite enormous success in modelling water, HF-SCAN is not a panacea. In their water simulations, Dasgupta *et al.* used HF-SCAN without dispersion correction, as they found that the standard dispersion

corrections, such as those of Grimme[43], worsen the original results of HF-SCAN for water. But such dispersion corrections have long been known to be necessary for noncovalent interactions (NCIs).[44, 45, 46, 47, 48, 49, 50, 51, 52, 53] So, despite delivering a high accuracy for pure water simulations, HF-SCAN without a dispersion correction cannot describe accurately long-range dispersion interactions. For this reason, the errors of HF-SCAN are several times larger than those of DFT enhanced by a dispersion correction for the standard noncovalent datasets.[40] The challenge is then to construct an efficient density functional that correctly describes NCIs of different nature, while recovering or even improving the accuracy of HF-SCAN for water simulations.

HF- r^2 SCAN-DC₄, an integratively designed DC-DFT procedure: In the present paper, we resolve these issues by using the principles of DC-DFT to carefully parameterize a dispersion correction for HF- r^2 SCAN. This yields HF- r^2 SCAN-DC₄, which produces the following key results: (i) HF- r^2 SCAN-DC₄ improves upon HF-SCAN for pure water simulations, by up to 0.7 kcal/mol for relative energies of water hexamers, and up to 2.4 kcal/mol for those of water 20-mers; (ii) HF- r^2 SCAN-DC₄ is far more accurate than HF-SCAN for interactions of water with other molecules and for NCIs in general, because of the inclusion of explicit dispersion corrections; (iii) HF- r^2 SCAN-DC₄ can be routinely and efficiently used in calculations because, unlike HF-SCAN[31], HF- r^2 SCAN-

DC4 has no grid convergence issues. In our HF- r^2 SCAN-DC4, each of the three ingredients is vitally important: The "HF" part reduces density-driven errors, while r^2 SCAN fixes the grid issues of SCAN. But most importantly, the way in which we parametrize the D4 corrections by using the DC-DFT principles is vital, as an unwitting fitting of D4 ruins the accuracy for water simulations. If we drop any of those elements of HF- r^2 SCAN-DC4, at least one of its three appealing results will be lost.

To illustrate all these points, and how they work together, we created Figure 1. We show how HF- r^2 SCAN-DC4 is better than HF-SCAN for interactions of nucleobases [panel (a)], water molecules with one another [panels (b), (c), (d)], water with other molecules [panel (e)], and NCIs in general [panel (f)].

Stacking interactions in nucleobases are of vital importance in biology as their energetics is essential to describe the formation and stability of DNA and RNA.[55, 56] In Figure 1(a), we compare the accuracy of HF-SCAN and HF- r^2 SCAN-DC4 for interaction energies of stacked cytosine dimers at different configurations. As we can see from Figure 1(a), our HF- r^2 SCAN-DC4 essentially greatly reduces the errors of HF-SCAN that systematically underbinds these stacked complexes by about 2.5 kcal/mol. This demonstrates that despite its success for modeling water, HF-SCAN misses most of dispersion and thus cannot compete with our HF- r^2 SCAN-DC4 in modelling NCIs. This is especially the case for NCIs dominated by dispersion interactions as those present in stacked nucleobases. (See Figure S3 in the supporting information for the errors in interaction energies.) We note that the mean absolute error (MAE) of HF- r^2 SCAN-DC4 (0.4 kcal/mol) is very good relative to HF-SCAN, but not very impressive relative to B3LYP-D3(BJ) (less than 0.2 kcal/mol).[56] But such functionals include only a fraction of HF exchange, and so still suffer from large density-driven errors in water, and so have larger errors for pure water (as shown below).

Water hexamers, "the smallest drops of water"[57, 58], are important, as they represent the transition from two-dimensional to three-dimensional hydrogen-bonding networks.[59, 60, 61] The energy differences between two adjacent isomers of water hexamers are tiny, making even the ordering of isomers a very challenging test for quantum-chemical methods.[59, 62] In Figure 1(b), we compare the energies of water hexamer isomers relative to the energy of the prism, as the lowest-lying isomer.[59, 63, 64] Despite being more accurate for water hexamers than most DFT methods available on the market, HF-SCAN mistakes the ordering of the isomers, as it predicts too low energies of the chair isomer. Our HF- r^2 SCAN-DC4 is also here superior to HF-SCAN, as it not only gives the right ordering of isomers, but essentially reproduces the reference values for the relative energies of isomers. If D4 is fitted by not accounting for the DC-DFT principles (see below), the accuracy of HF- r^2 SCAN-DC4 for the water simulation is lost. This happened in Ref. [65] and will be discussed in the method section.

We use the WATER27 dataset to illustrate the importance (and subtlety) of DC-DFT for water simulations. WATER27 is a standard dataset for binding energies of water clusters. Density sensitivity, \tilde{S} , is a measure for how sensitive a given DFT simulation is to errors in densities (see Section S2 in the supporting information for further details and specific definitions).[23] Typically, the errors of SC-DFT calculations grow with \tilde{S} , indicating the presence of large density-driven errors.[26, 31, 66] DC-DFT reduces these large density-driven errors of SC-DFT and thus the errors of DC-DFT do not grow with \tilde{S} . In Figure 1(c) we plot WATER27 errors as a function of

density sensitivity. As the errors of SC- r^2 SCAN-D4 grow with \tilde{S} , so also does the energetic improvement of HF- r^2 SCAN-DC4 over SC- r^2 SCAN-D4. Furthermore, sometimes dispersion corrections worsen SC-DFT for cases with large density-driven errors.[25, 26] This is also the case here, as SC- r^2 SCAN-D4 significantly deteriorates the accuracy of SC- r^2 SCAN (see Figure S4). The errors of HF-SCAN are also substantially lower than those of SC- r^2 SCAN-D4, and for most of the binding energies of the WATER27 clusters, HF-SCAN is comparable to HF- r^2 SCAN-DC4. But, for the four clusters with the largest sensitivities, HF- r^2 SCAN-DC4 outperforms HF-SCAN by ~ 4 kcal/mol.

WATER27 is a part of the GMTKN55[40], a database that we use to train the D4 parameters in HF- r^2 SCAN-DC4 (see methods). But, according to the principles of DC-DFT, we exclude those WATER27 clusters that are density-sensitive, as their energetic errors are dominated by the errors in their densities.[26] Thus none of the clusters that are to the right of the vertical dashed line placed at $\tilde{S} = 2$ kcal/mol (see Method Section for the details on this reasoning) are used in the fitting, which means HF- r^2 SCAN-DC4 makes genuinely accurate predications for a vast majority of these water clusters. Not only does it recover HF-SCAN for binding energies of the water clusters, but also provides substantial improvements for the most challenging clusters.

An important question is whether or not one should *always* correct the density. The general principles of DC-DFT say that one should only correct the density in cases of substantial density-driven errors. In density insensitive cases, the effect of correcting the density should be small, and may actually worsen energetics. Figure 1(d) shows energies of water 20-mers relative to the energy of the lowest of the four 20-mers. Here SC- r^2 SCAN-D4 beats its DC counterpart every time. In contrast to large \tilde{S} for binding energies of the four 20-mers (the last four datapoints in Figure 1(c)), the sensitivities corresponding to their relative isomer energies are about twenty times smaller (see Figure S7). Thus the higher accuracy of SC- r^2 SCAN-D4 over HF- r^2 SCAN-DC4 does not come as a surprise. But the crucial point is that, even in this *low-sensitivity* scenario, the errors introduced by the HF density are far smaller than those of HF-SCAN, and remain tiny on a per molecule basis.

A crucial figure of merit is how accurate energetics are for water molecules in the vicinity of an organic molecule, especially if it is polar. In Figure 1(e), we show errors in the interaction energies between water and aspirin from structures that we extracted from an MD simulation at $T=298.15$ K (see Section S6 for further details on the MD simulation). The structures are sorted by the distance between the oxygen atom in water and the specified oxygen atom in the carboxyl group of aspirin. The errors of HF- r^2 SCAN-DC4 are much smaller than those of HF-SCAN. They are also substantially smaller than those of SC- r^2 SCAN-D4 (see Figure S8), demonstrating again the importance of both the D4 and DC components in our method.

Getting NCI right across a broad range of molecules is important, even in the absence of water. The GMTKN55 collection of 55 databases has become a standard benchmark[40] and includes many databases for NCIs. In Figure 1(f), we compare the MAEs of HF-SCAN and HF- r^2 SCAN-DC4 for the standard datasets with intra- and intermolecular NCIs[40]. Despite its high accuracy for water clusters, HF-SCAN does not capture long-ranged dispersion interactions. This is why it is far less accurate than HF- r^2 SCAN-DC4 for noncovalent datasets. We can see that HF- r^2 SCAN-DC4




Figure 2: Water dimer interaction energies for (a) Smith stationary points[54] and (b) MD simulated water dimers with the oxygen-oxygen distance. For (a), MAEs of each functional are (following the order in the legend) 0.25, 0.11, 0.09, 0.17, and 0.08 kcal/mol. DLPNO-CCSD(T)-F12 has been used as a reference. For (b), MAE of each functionals are 0.25, 0.08, 0.20, 0.30 and 0.08 kcal/mol. Figure S1 shows the corresponding density sensitivities and Figure S2 shows the errors of approximations for the Smith dimers and interaction energies for MD dimers.

is highly accurate here, and on average it beats SC-r²SCAN-D4 for both inter- and intramolecular NCI (see Table S1 in the supporting information comparing the metrics for overall performance).

II. RESULTS

i. Interaction energies for water dimers

As discussed already, HF-SCAN performs incredibly well for interactions in pure water. In this section, we look at select water dimers that are relevant to water simulations, and show how HF-r²SCAN-DC4 reproduces (or even exceeds) this accuracy. More importantly, we show how each aspect of its construction (density correction, regularization of SCAN, and dispersion correction) are vital to its accuracy for water. Later we will show that no other approximation at this level of cost comes close to this performance for water.

Figure 2 shows the interaction energies for many water dimers (the difference in the energies of a dimer and two monomers). (a) shows the interaction energies at Smith stationary points, some of which resemble geometries from dense ice structures.[18] (b) shows the errors of approximations in interaction energies for water dimers as a function of the distance between the two oxygen atoms. The underlying structures were extracted from an MD simulation at T=298.15K (see Section S6 for further details on the simulation). For the interaction energies of these water dimers, HF-SCAN without a dispersion correction already provides a very high accuracy (with MAEs of less than 0.1 kcal/mol). Our HF-r²SCAN-DC4 essentially recovers this high accuracy of HF-SCAN. Similar patterns observed for binding energies of water clusters are also seen here.

By studying the various plots, one can assess the importance of the relative contributions to HF-r²SCAN-DC4. First, the purple points give HF-r²SCAN, to be contrasted with HF-SCAN. We see that HF-r²SCAN significantly (on this scale) underestimates the interaction energy. Even though r²SCAN was designed to

reproduce the results of SCAN, these differences are so small as to be negligible for most purposes. However, they are clearly significant here, showing HF-r²SCAN is noticeably less accurate for these dimers. The addition of the D4 correction, however, makes their errors comparable.

On the other hand, we may also consider the importance of density correction. We see that SC-r²SCAN-D4 considerably overestimates interaction energies. In fact, SC-r²SCAN does rather well, as the errors due to poor density and missing dispersion cancel.

We can also observe from Figure 2(b) that the improvement of HF-r²SCAN-DC4 over SC-r²SCAN-D4 decreases with the distance between the two oxygen atoms in water dimers. This can be understood in terms of underlying density sensitivity which also decreases with the O-O distance (see Figure S1).

ii. Many-body interactions in larger water clusters

In Figure 3 we compare errors of HF-r²SCAN-DC4 and HF-SCAN for the interaction energies of the eight standard water hexamers.[59, 60] In addition to total interaction energies, we also use the many-body expansion (MBE) to show the K -body contributions to these energies (with K in between 2 and 6). This is a standard methodology for understanding the origins of errors in water models.[3, 13, 67] The energetic importance of the K -body contributions decreases rapidly with K (Figure S6), making the 2-body contributions by far the most important, and these are where significant differences emerge when the density is corrected. But in order to reach chemical accuracy, a proper description of the higher-order contributions also matters. The 2-body plot shows that HF-SCAN has a rather systematic overestimate of about 0.5 kcal/mol, whereas HF-r²SCAN-DC4 is substantially less for about half the clusters. The 3-body plot shows them being almost identical. But in the total error, we see that HF-r²SCAN-DC4 is far more systematic, as HF-SCAN makes errors of opposite sign, while




Figure 3: K -body interaction energy errors with (a) $K = 2$, (b) $K = 3$, and (c) total, and (d) the interaction energy for 8 water hexamers. (For higher order K -body interaction energies, see Figure S5 and S6.) Geometries and CCSD(T)/CBS reference interaction energies are from Ref. [13]. The MAEs of HF- r^2 SCAN-DC4 and HF-SCAN are 0.19 kcal/mol and 0.22 kcal/mol, respectively.

HF- r^2 SCAN-DC4 is always an overestimate of about 0.2 kcal/mol.

This consistency is important on the plot (d), showing the interaction energy of the 8 hexamers. Because HF- r^2 SCAN-DC4 is so consistent, it gets the ordering in interaction energies of all clusters correct, whereas HF-SCAN incorrectly predicts that the interaction energy in the bag is higher than that of the chair. The MAE of HF- r^2 SCAN-DC4 is 0.19 kcal/mol, lower than 0.22 kcal/mol for HF-SCAN. On average, HF- r^2 SCAN-DC4 also improves individual K -body contributions to the interaction energies, except for $K = 4$, where both are marginally small (Figure S5). This MBE test shows us that the improvement of HF- r^2 SCAN-DC4 over HF-SCAN for the water hexamer interaction energies (seen also for the relative isomer energies (Figure 1(b)) is systematic and does not result from the error cancellations between different K -body contributions (for the detailed information of water hexamer isomerization energy in Figure 1, see Figure S9).

iii. Water...cytosine interaction energies

In Figure 4, we study the performance of different variations for microhydration of cytosine, by specifically focusing on the interaction energies in water...cytosine complexes. We generate these complexes as described in Section S7, and in all of them, water interacts with cytosine through the hydrogen bond formed between the hydrogen atom in water and the oxygen atom in cytosine. For each complex, the errors of HF- r^2 SCAN-DC4 are small, and with the MAE of 0.09 kcal/mol, it is the best performer in Figure 4.

The errors of HF-SCAN are much smaller here than for cytosine dimers (Figure 1(a)), in which the role of dispersion is more important. Nevertheless, HF- r^2 SCAN-DC4 provides here a significant improvement over HF-SCAN. It is also interesting to observe what happens after we add the dispersion correction to HF- r^2 SCAN and its SC counterpart. In the case of HF- r^2 SCAN, the errors in the interaction energies are greatly reduced (roughly by a factor of 6




Figure 4: Errors in interaction energies of water...cytosine complexes sorted by the distance between the oxygen atom in cytosine and the oxygen atom in water. Reference interaction energies have been computed at the DLPNO-CCSD(T)-F12/AVQZ level of theory.

on average). In stark contrast, adding D4 to SC- r^2 SCAN significantly deteriorates its accuracy, as SC- r^2 SCAN already overbinds water...cytosine complexes and D4 makes the overbinding stronger.

iv. Wide applicability of HF- r^2 SCAN-DC4

A functional that works extremely well for pure water but nothing else is not widely applicable. Recently, GMTKN55 of 55 databases has become a popular benchmark for testing the accuracy of density functionals for main-group chemistry. Figure 5 has been designed to illustrate performance of functionals for both pure water and on the GMTKN55 database simultaneously. The water metric (y -axis on the left) combines most of the reactions with water used in this paper, and is carefully defined in the supporting information Section S8.

Figure 5(a) shows errors on GMTKN55 on the x -axis and errors on the water metric on the y -axis, each in kcal/mol. The x -axis ranges from about 3-10 kcal/mol, spanning the performance of modern approximations for main group chemistry, such as atomization energies. The y -axis range is much smaller, running less than 4.0 kcal/mol, reflecting the much smaller magnitude of NCIs in water, and how high accuracy needs to be in order to have an accurate model for water. Here, HF-SCAN sets a high standard, with a water error near 1.0 kcal/mol (the chemical accuracy claimed in Ref. [3]), while most standard-use functionals cannot compete. On the other hand, SCAN is designed mainly to improve materials calculations without the cost of a hybrid functional, and HF-SCAN has a high error on GMTKN55 (about 9 kcal/mol). Popular functionals have much smaller GMTKN55 errors, but perform worse on water. We also show the many combinations of HF- r^2 SCAN-DC4 that do not include all the right ingredients, showing they all perform less well on water than HF-SCAN. We finally include ω B97M-V functional [69], which might be considered the DFT gold-standard here, with the smallest errors for both water and main-group chemistry. But this range-separated functional with nonlocal correlation functional is far more expensive to compute than most functionals, including its own D4 variant,[70] and is less practical for DFT-MD simulations

than e.g., SCAN. We have included it here only to show what is possible in principle with DFT.

But the performance of HF- r^2 SCAN-DC4 is remarkable. Its errors on both water *and* the GMTKN55 dataset are almost half of those of HF-SCAN. No other functional in our collection comes close for water. Clearly, all the chemically-inclined approximations which are comparable for main-group chemistry do much worse.

In Figure 5(b), we show the hexagon plots comparing the MAEs of several density functionals, where the position of five vertices denote the MAEs for individual water-based datasets, while the sixth vertex denotes the overall performance of the functionals for the whole GMTKN55 databases, as measured by the weighted-mean-absolute-deviation-2 (WTMAD-2). It is the MAE for all the reactions from these five water-based datasets that we use as the quantity on the y -axis in Figure 5(a). The size of the hexagon of HF- r^2 SCAN-DC4 is the closest to that of more costly ω B97M-V. We can also see that the performance of HF- r^2 SCAN-DC4 is far superior to that of HF-SCAN. M062X-D3(0), a meta-hybrid that is very accurate for small organic molecules,[40] and yields WTMAD-2 which is slightly lower than that of HF- r^2 SCAN-DC4. But, for water simulations, M062X-D3(0) is nowhere close to HF- r^2 SCAN-DC4, as can be seen from the position of the remaining five vertices.

III. METHODS

The basic principles of DC-DFT are covered elsewhere in the literature[19, 22], and reviewed in the supporting information. In most KS-DFT calculations, the error in the density has a negligible effect on the energy errors. But sometimes the error in a SC density leads to a noticeable contribution, which can be reduced if a more accurate density is used instead. For many semi-local exchange-correlation approximations in molecular calculations, when a calculation is density sensitive, often the HF density then yields significantly smaller energy errors. These principles have led to improved energetics in reaction barrier heights, electron affinities, and also for the ground state geometries of noncovalent interaction systems, etc.[20, 71, 72, 73]

Application of the principles of DC-DFT is subtle in the case of r^2 SCAN-D4, because of the need to separate out the error due to density correction from the fitting of the D4 corrections. For example, for halogen bonds, the density-driven errors are far larger than dispersion corrections, so all fitting must be done on density-corrected energetics. Moreover, when empirical functionals contain parameters, such parameters should be fit only on density-insensitive calculations, so that the parameters optimize the 'true' functional error.

With these principles in mind, we find the parameters for HF- r^2 SCAN-DC4 using the density-insensitive calculations in the GMTKN55 dataset as a training set while using water...water pair interaction energy as a validation set. We find their optimum values by minimizing MAE values over all such cases. This is detailed in the supporting information section S4. This is why we use the acronym DC4 instead of D4, meaning that we use the principles of DC-DFT to find the underlying D4 parameters.

In Refs. [31] and [66], we proposed DC(HF)-DFT, a DC-DFT procedure that discriminately uses HF densities based on the density sensitivity criterion. The main idea of DC(HF)-DFT is to use HF-DFT for density-sensitive (DS) reactions and SC-DFT for




Figure 5: (a) The mean absolute error (MAE) for the water-based reactions appear in this work (hexamer isomer energies, water 20-mers isomer energies, WATER27 binding energies, water-small organic molecule interaction energies, and water dimer interaction energies) versus the weighted-mean-absolute-deviation-2 (WTMAD-2) for the GMTKN55 database for selected functionals. For a further description of the reactions used in the y-axis, see Section S8 in the supporting information. HF-SCAN-D4 functional used here is from Ref. [68]. (b) The hexagon plot with MAEs for selected water-based datasets and WTMAD-2 values for the whole GMTKN55 databases (for WTMAD-2 values for other GMTKN55 database, see Figure S10). Abbreviations of isomerization (I), binding (B), and interaction (T) energy are noted in the vertex caption. MAEs of HF-r²SCAN-DC4 for individual GMTKN55 datasets are shown in Table S1. In Figure S11, we give further details about the interaction energies used in the "water-small organic molecule" dataset.

density-insensitive (DI) reactions (possible spin-contaminations of the HF results are also taken into account as detailed in Section S3). While we consider DC(HF)-DFT a state-of-the-art DC-DFT procedure, for our HF-r²SCAN-DC4 we use HF-DFT, meaning that the functional is always evaluated on the HF density regardless of the sensitivity criterion. To use DC(HF)-DFT, we need to compute density sensitivity for each reaction of interest and possibly make adjustments to its cutoff value which is used to declare whether a given reaction is DS or DI.[31, 66] This would also require having two sets of D4 parameters, one for DS and the other for DI reactions. All these efforts would undermine the ease of use of r²SCAN, which is a general-purpose functional. For this reason and encouraged by the very good performance of HF-DFT with SCAN-like functionals[32, 42], we employ HF-DFT[74] as a DC-DFT procedure for HF-r²SCAN-DC4. While our HF-r²SCAN-DC4 can be routinely used by applying it to HF orbitals without ever needing to calculate density sensitivity of a given reaction, the use of DC-DFT principles and density sensitivity is vital for our training of HF-r²SCAN-DC4 as explained above.

To illustrate what can happen when these principles are not applied, we show results from Ref. [65]. This is a version of HF-r²SCAN-D4, but where all reactions in GMTKN55 were used, and the WTMAD-2[40] was used as the cost function instead. Figure 6 illustrates the results for the larger water clusters. In every case, they are noticeably worse than ours. Moreover, (d) shows that, apart from matching on WTMAD-2 measure, HF-r²SCAN-DC4 yields more accurate results in every other case.

IV. CONCLUSIONS

The work of Ref. [3] was a breakthrough in models for water, showing that, by using the principles of DC-DFT, a moderate-cost density functional approximation approached chemical accuracy for many relevant properties of small water clusters. However that functional is lacking in dispersion corrections, yielding large errors for energetics between organic and biological molecules. It also inherits some of the numerical issues of the original SCAN functional, which have been eliminated by using r²SCAN instead in most other applications. However, the small differences between these two wreak havoc on the much smaller scale of subtle energy differences of water clusters.

The present work shows that, by a very careful application of the principles of DC-DFT, all these difficulties can be overcome, and even greater accuracy achieved for pure water, while still including dispersion for other molecules where it can be vital. Finding the correct parameters depends crucially on training on only density-insensitive chemical reactions, as inclusion of density-sensitive reactions yields suboptimal values for the parameters.

Even if HF-r²SCAN-DC4 could be run at close to meta-GGA cost, KS-DFT MD simulations are typically far more costly than MD with machine learning (ML) interatomic potentials. But accurate force-field generation requires highly accurate reference energetics data as a training set, and CCSD(T) or Quantum Monte Carlo (QMC) methods are frequently used as reference methods these days.[12] Due to the large computational cost for such *ab initio* calculation, a more practical yet accurate method is in demand, and HF-r²SCAN-DC4 can replace them for




Figure 6: Comparison between HF-r²SCAN-DC4 and SM21 for (a) water hexamer isomerization energy, (b) water hexamer interaction energy, (c) water 20-mer isomerization energy, and (d) hexagonal plot same as Figure 5(b). SM21 (green) is HF-r²SCAN but with different D4 parameters obtained in Ref. [65].

calculating moderately large biomolecular systems. We suggest HF-r²SCAN-DC4 be tested and applied in solution wherever practical.

ACKNOWLEDGEMENT

ES, SS, YK, and HY are grateful for support from the National Research Foundation of Korea (NRF-2020R1A2C2007468 and NRF-2020R1A4A1017737) and Samsung Electronics (IO211126-09176-01). KB acknowledges funding from NSF (CHE-2154371). SV acknowledges funding from the Marie Skłodowska-Curie grant 101033630 (EU’s Horizon 2020 programme). We thank John Perdew and Francesco Paesani and his group for many useful discussions.

REFERENCES

- [1] Linfeng Zhang, Han Wang, Roberto Car, and E Weinan. Phase diagram of a deep potential water model. *Physical Review Letters*, 126(23):236001, 2021.
- [2] Kamal Sharkas, Kamal Wagle, Biswajit Santra, Sharmin Akter, Rajendra R Zope, Tunna Baruah, Koblar A Jackson, John P Perdew, and Juan E Peralta. Self-interaction error overbinds water clusters but cancels in structural energy differences. *Proceedings of the National Academy of Sciences*, 117(21):11283–11288, 2020.
- [3] Saswata Dasgupta, Eleftherios Lambros, John P Perdew, and Francesco Paesani. Elevating density functional theory to chemical accuracy for water simu-

- lations through a density-corrected many-body formalism. *Nature Communications*, 12(1):1–12, 2021.
- [4] Saswata Dasgupta, Chandra Shahi, Pradeep Bhetwal, John P. Perdew, and Francesco Paesani. How good is the density-corrected scan functional for neutral and ionic aqueous systems, and what is so right about the Hartree-Fock density? *Journal of Chemical Theory and Computation*, 18(8):4745–4761, 2022.
- [5] Chunyi Zhang, Fujie Tang, Mohan Chen, Jianhang Xu, Linfeng Zhang, Diana Y Qiu, John P Perdew, Michael L Klein, and Xifan Wu. Modeling liquid water by climbing up jacob’s ladder in density functional theory facilitated by using deep neural network potentials. *Journal of Physical Chemistry B*, 125(41):11444–11456, 2021.
- [6] Alberto Torres, Luana S Pedroza, Marivi Fernandez-Serra, and Alexandre R Rocha. Using neural network force fields to ascertain the quality of ab initio simulations of liquid water. *Journal of Physical Chemistry B*, 125(38):10772–10778, 2021.
- [7] Eleftherios Lambros, Jie Hu, and Francesco Paesani. Assessing the accuracy of the scan functional for water through a many-body analysis of the adiabatic connection formula. *Journal of Chemical Theory and Computation*, 17(6):3739–3749, 2021.
- [8] Etienne Palos, Eleftherios Lambros, Steven Swee, Jie Hu, Saswata Dasgupta, and Francesco Paesani. Assessing the interplay between functional-driven and density-driven errors in dft models of water. *Journal of Chemical Theory and Computation*, 18(6):3410–3426, 2022.
- [9] Etienne Palos, Eleftherios Lambros, Saswata Dasgupta, and Francesco Paesani. Density functional theory of water with the machine-learned dm21 functional. *Journal of Chemical Physics*, 156(16):161103, 2022.
- [10] Volodymyr Babin, Claude Leforestier, and Francesco Paesani. Development of a “first principles” water potential with flexible monomers: Dimer potential energy surface, vrt spectrum, and second virial coefficient. *Journal of Chemical Theory and Computation*, 9(12):5395–5403, 2013.
- [11] Volodymyr Babin, Gregory R Medders, and Francesco Paesani. Development of a “first principles” water potential with flexible monomers. ii: Trimer potential energy surface, third virial coefficient, and small clusters. *Journal of Chemical Theory and Computation*, 10(4):1599–1607, 2014.
- [12] Gregory R Medders, Volodymyr Babin, and Francesco Paesani. Development of a “first principles” water potential with flexible monomers. iii. liquid phase properties. *Journal of Chemical Theory and Computation*, 10(8):2906–2910, 2014.
- [13] Sandeep K Reddy, Shelby C Straight, Pushp Bajaj, C. Huy Pham, Marc Riera, Daniel R Moberg, Miguel A Morales, Chris Knight, Andreas W Götz, and Francesco Paesani. On the accuracy of the mb-pol many-body potential for water: Interaction energies, vibrational frequencies, and classical thermodynamic and dynamical properties from clusters to liquid water and ice. *Journal of Chemical Physics*, 145(19):194504, 2016.
- [14] Mohan Chen, Hsin-Yu Ko, Richard C Remsing, Marcos F Calegari Andrade, Biswajit Santra, Zhaoru Sun, Annabella Selloni, Roberto Car, Michael L Klein, John P Perdew, et al. Ab initio theory and modeling of water. *Proceedings of the National Academy of Sciences*, 114(41):10846–10851, 2017.
- [15] I-Feng W Kuo, Christopher J Mundy, Matthew J McGrath, J Ilja Siepmann, Joost VandeVondele, Michiel Sprik, Jürg Hutter, Bin Chen, Michael L Klein, Fawzi Mohamed, et al. Liquid water from first principles: Investigation of different sampling approaches. *Journal of Physical Chemistry B*, 108(34):12990–12998, 2004.
- [16] Thomas D Kühne, Matthias Krack, and Michele Parrinello. Static and dynamical properties of liquid water from first principles by a novel car-parrinello-like approach. *Journal of Chemical Theory and Computation*, 5(2):235–241, 2009.
- [17] Biswajit Santra, Jiří Klimeš, Dario Alfe, Alexandre Tkatchenko, Ben Slater, Angelos Michaelides, Roberto Car, and Matthias Scheffler. Hydrogen bonds and van der waals forces in ice at ambient and high pressures. *Physical Review Letters*, 107(18):185701, 2011.
- [18] Michael J Gillan, Dario Alfe, and Angelos Michaelides. Perspective: How good is dft for water? *Journal of Chemical Physics*, 144(13):130901, 2016.
- [19] Min-Cheol Kim, Eunji Sim, and Kieron Burke. Understanding and reducing errors in density functional calculations. *Physical Review Letters*, 111(7):073003, 2013.
- [20] Min-Cheol Kim, Eunji Sim, and Kieron Burke. Ions in solution: Density corrected density functional theory (dc-dft). *Journal of Chemical Physics*, 140(18):18A528, 2014.
- [21] Adam Wasserman, Jonathan Nafziger, Kaili Jiang, Min-Cheol Kim, Eunji Sim, and Kieron Burke. The importance of being self-consistent. *Annual Review of Physical Chemistry*, 68(1):555–581, 2017.

-
- [22] Stefan Vuckovic, Suhwan Song, John Kozłowski, Eunji Sim, and Kieron Burke. Density functional analysis: The theory of density-corrected dft. *Journal of Chemical Theory and Computation*, 15(12):6636–6646, 2019.
- [23] Eunji Sim, Suhwan Song, and Kieron Burke. Quantifying density errors in dft. *Journal of Physical Chemistry Letters*, 9(22):6385–6392, 2018.
- [24] Seungsoo Nam, Suhwan Song, Eunji Sim, and Kieron Burke. Measuring density-driven errors using kohn–sham inversion. *Journal of Chemical Theory and Computation*, 16(8):5014–5023, July 2020.
- [25] Subrata Jana, Abhilash Patra, Szymon Śmiga, Lucian A Constantin, and Prasanjit Samal. Insights from the density functional performance of water and water–solid interactions: Scan in relation to other meta-ggas. *Journal of Chemical Physics*, 153(21):214116, 2020.
- [26] Suhwan Song, Stefan Vuckovic, Eunji Sim, and Kieron Burke. Density sensitivity of empirical functionals. *Journal of Physical Chemistry Letters*, 12(2):800–807, 2021.
- [27] Min-Cheol Kim, Eunji Sim, and Kieron Burke. Communication: Avoiding unbound anions in density functional calculations. *Journal of Chemical Physics*, 134(17):171103, 2011.
- [28] Min-Cheol Kim, Hansol Park, Suyeon Son, Eunji Sim, and Kieron Burke. Improved dft potential energy surfaces via improved densities. *Journal of Physical Chemistry Letters*, 6(19):3802–3807, 2015.
- [29] Suhwan Song, Min-Cheol Kim, Eunji Sim, Anouar Benali, Olle Heinonen, and Kieron Burke. Benchmarks and reliable dft results for spin gaps of small ligand fe (ii) complexes. *Journal of Chemical Theory and Computation*, 14(5):2304–2311, 2018.
- [30] Yeil Kim, Suhwan Song, Eunji Sim, and Kieron Burke. Halogen and chalcogen binding dominated by density-driven errors. *Journal of Physical Chemistry Letters*, 10(2):295–301, 2018.
- [31] Suhwan Song, Stefan Vuckovic, Eunji Sim, and Kieron Burke. Density-corrected dft explained: Questions and answers. *Journal of Chemical Theory and Computation*, 18(2):817–827, 2022.
- [32] Jianwei Sun, Adrienn Ruzsinszky, and John P Perdew. Strongly constrained and appropriately normed semilocal density functional. *Physical Review Letters*, 115(3):036402, 2015.
- [33] Axel D Becke. Density-functional exchange-energy approximation with correct asymptotic behavior. *Physical Review A*, 38(6):3098–3100, 1988.
- [34] Chengteh Lee, Weitao Yang, and Robert G Parr. Development of the colle-salvetti correlation-energy formula into a functional of the electron density. *Physical Review B*, 37(2):785–789, 1988.
- [35] Axel D Becke. Density-functional thermochemistry. iii. the role of exact exchange. *Journal of Chemical Physics*, 98(7):5648–5652, 1993.
- [36] P. J. Stephens, F. J. Devlin, C. F. Chabalowski, and M. J. Frisch. Ab initio calculation of vibrational absorption and circular dichroism spectra using density functional force fields. *Journal of Physical Chemistry*, 98(45):11623–11627, 1994.
- [37] Jianwei Sun, Richard C Remsing, Yubo Zhang, Zhaoru Sun, Adrienn Ruzsinszky, Haowei Peng, Zenghui Yang, Arpita Paul, Umesh Waghmare, Xifan Wu, et al. Accurate first-principles structures and energies of diversely bonded systems from an efficient density functional. *Nature Chemistry*, 8(9):831–836, 2016.
- [38] Fabien Tran, Julia Stelzl, and Peter Blaha. Rungs 1 to 4 of dft jacob’s ladder: Extensive test on the lattice constant, bulk modulus, and cohesive energy of solids. *Journal of Chemical Physics*, 144(20):204120, 2016.
- [39] Narbe Mardirossian and Martin Head-Gordon. Thirty years of density functional theory in computational chemistry: an overview and extensive assessment of 200 density functionals. *Molecular Physics*, 115(19):2315–2372, 2017.
- [40] Lars Goerigk, Andreas Hansen, Christoph Bauer, Stephan Ehrlich, Asim Najibi, and Stefan Grimme. A look at the density functional theory zoo with the advanced gmtkn55 database for general main group thermochemistry, kinetics and noncovalent interactions. *Physical Chemistry Chemical Physics*, 19(48):32184–32215, 2017.
- [41] Yubo Zhang, Jianwei Sun, John P Perdew, and Xifan Wu. Comparative first-principles studies of prototypical ferroelectric materials by lda, gga, and scan meta-gga. *Physical Review B*, 96(3):035143, 2017.
- [42] James W Furness, Aaron D Kaplan, Jinliang Ning, John P Perdew, and Jianwei Sun. Accurate and numerically efficient r2scan meta-generalized gradient approximation. *Journal of Physical Chemistry Letters*, 11(19):8208–8215, 2020.
- [43] Stefan Grimme, Jens Antony, Stephan Ehrlich, and Helge Krieg. A consistent and accurate ab initio parametrization of density functional dispersion correction (dft-d) for the 94 elements h-pu. *Journal of Chemical Physics*, 132(15):154104, 2010.

-
- [44] Xueyuan Wu, MC Vargas, S Nayak, V Lotrich, and G Scoles. Towards extending the applicability of density functional theory to weakly bound systems. *Journal of Chemical Physics*, 115(19):8748–8757, 2001.
- [45] Evert Jan Meijer and Michiel Sprik. A density-functional study of the intermolecular interactions of benzene. *Journal of Chemical Physics*, 105(19):8684–8689, 1996.
- [46] Stefan Grimme. Accurate description of van der waals complexes by density functional theory including empirical corrections. *Journal of Computational Chemistry*, 25(12):1463–1473, 2004.
- [47] Axel D Becke and Erin R Johnson. Exchange-hole dipole moment and the dispersion interaction. *Journal of Chemical Physics*, 122(15):154104, 2005.
- [48] Stefan Grimme. Semiempirical gga-type density functional constructed with a long-range dispersion correction. *Journal of Computational Chemistry*, 27(15):1787–1799, 2006.
- [49] Axel D Becke and Erin R Johnson. Exchange-hole dipole moment and the dispersion interaction revisited. *Journal of Chemical Physics*, 127(15):154108, 2007.
- [50] Alexandre Tkatchenko and Matthias Scheffler. Accurate molecular van der waals interactions from ground-state electron density and free-atom reference data. *Physical Review Letters*, 102(7):073005, 2009.
- [51] Oleg A Vydrov and Troy Van Voorhis. Nonlocal van der waals density functional: The simpler the better. *Journal of Chemical Physics*, 133(24):244103, 2010.
- [52] Jan Rezac and Pavel Hobza. Benchmark calculations of interaction energies in noncovalent complexes and their applications. *Chemical Reviews*, 116(9):5038–5071, 2016.
- [53] Alastair JA Price, Kyle R Bryenton, and Erin R Johnson. Requirements for an accurate dispersion-corrected density functional. *Journal of Chemical Physics*, 154(23):230902, 2021.
- [54] Brian J Smith, David J Swanton, John A Pople, Henry F Schaefer III, and Leo Radom. Transition structures for the interchange of hydrogen atoms within the water dimer. *Journal of Chemical Physics*, 92(2):1240–1247, 1990.
- [55] Ilyas Yildirim and Douglas H Turner. Rna challenges for computational chemists. *Biochemistry*, 44(40):13225–13234, 2005.
- [56] Holger Kruse and Jiri Sponer. Revisiting the potential energy surface of the stacked cytosine dimer: Fno-ccsd (t) interaction energies, sapt decompositions, and benchmarking. *Journal of Physical Chemistry A*, 123(42):9209–9222, 2019.
- [57] K Nauta and RE Miller. Formation of cyclic water hexamer in liquid helium: The smallest piece of ice. *Science*, 287(5451):293–295, 2000.
- [58] Yimin Wang, Volodymyr Babin, Joel M Bowman, and Francesco Paesani. The water hexamer: cage, prism, or both. full dimensional quantum simulations say both. *Journal of the American Chemical Society*, 134(27):11116–11119, 2012.
- [59] Desiree M Bates and Gregory S Tschumper. Ccsd (t) complete basis set limit relative energies for low-lying water hexamer structures. *Journal of Physical Chemistry A*, 113(15):3555–3559, 2009.
- [60] Yiming Chen and Hui Li. Intermolecular interaction in water hexamer. *Journal of Physical Chemistry A*, 114(43):11719–11724, 2010.
- [61] A Otero-De-La-Roza and Erin R Johnson. Non-covalent interactions and thermochemistry using xdm-corrected hybrid and range-separated hybrid density functionals. *Journal of Chemical Physics*, 138(20):204109, 2013.
- [62] Biswajit Santra, Angelos Michaelides, Martin Fuchs, Alexandre Tkatchenko, Claudia Filippi, and Matthias Scheffler. On the accuracy of density-functional theory exchange-correlation functionals for h bonds in small water clusters. ii. the water hexamer and van der waals interactions. *Journal of Chemical Physics*, 129(19):194111, 2008.
- [63] Ryan M Olson, Jonathan L Bentz, Ricky A Kendall, Michael W Schmidt, and Mark S Gordon. A novel approach to parallel coupled cluster calculations: Combining distributed and shared memory techniques for modern cluster based systems. *Journal of Chemical Theory and Computation*, 3(4):1312–1328, 2007.
- [64] MJ Gillan, FR Manby, MD Towler, and D Alfè. Assessing the accuracy of quantum monte carlo and density functional theory for energetics of small water clusters. *Journal of Chemical Physics*, 136(24):244105, 2012.
- [65] Golokesh Santra and Jan ML Martin. Pure and hybrid scan, rscan, and r2scan: Which one is preferred in ks-and hf-dft calculations, and how does d4 dispersion correction affect this ranking? *Molecules*, 27(1):141, 2021.

- [66] Eunji Sim, Suhwan Song, Stefan Vuckovic, and Kieron Burke. Improving results by improving densities: Density-corrected density functional theory. *Journal of the American Chemical Society*, 144(15):6625–6639, 2022.
- [67] Urszula Góra, Rafał Podeszwa, Wojciech Cencek, and Krzysztof Szalewicz. Interaction energies of large clusters from many-body expansion. *Journal of Chemical Physics*, 135(22):224102, 2011.
- [68] Golokesh Santra and Jan ML Martin. What types of chemical problems benefit from density-corrected dft? a probe using an extensive and chemically diverse test suite. *Journal of Chemical Theory and Computation*, 17(3):1368–1379, 2021.
- [69] Narbe Mardirossian and Martin Head-Gordon. ω b97m-v: A combinatorially optimized, range-separated hybrid, meta-gga density functional with vv10 non-local correlation. *Journal of Chemical Physics*, 144(21):214110, 2016.
- [70] Narbe Mardirossian and Martin Head-Gordon. Survival of the most transferable at the top of jacob’s ladder: Defining and testing the ω b97m (2) double hybrid density functional. *Journal of Chemical Physics*, 148(24):241736, 2018.
- [71] Benjamin G Janesko and Gustavo E Scuseria. Hartree-fock orbitals significantly improve the reaction barrier heights predicted by semilocal density functionals. *Journal of Chemical Physics*, 128(24):244112, 2008.
- [72] Donghyung Lee, Filipp Furche, and Kieron Burke. Accuracy of electron affinities of atoms in approximate density functional theory. *Journal of Physical Chemistry Letters*, 1(14):2124–2129, 2010.
- [73] Donghyung Lee and Kieron Burke. Finding electron affinities with approximate density functionals. *Molecular Physics*, 108(19-20):2687–2701, 2010.
- [74] Yonsei University Theoretical and Computational Chemistry Laboratory. Density Corrected-Density Functional Theory. <http://tccl.yonsei.ac.kr/mediawiki/index.php/DC-DFT>, 2022. (accessed 2022-07-07).
- [75] Walter Kohn and Lu Jeu Sham. Self-consistent equations including exchange and correlation effects. *Physical Review*, 140(4A):A1133, 1965.
- [76] John P Perdew, Kieron Burke, and Matthias Ernzerhof. Generalized gradient approximation made simple. *Physical Review Letters*, 77(18):3865, 1996.
- [77] Enrico Clementi and Subhas J Chakravorty. A comparative study of density functional models to estimate molecular atomization energies. *Journal of Chemical Physics*, 93(4):2591–2602, 1990.
- [78] Peter MW Gill, Benny G Johnson, John A Pople, and Michael J Frisch. An investigation of the performance of a hybrid of hartree-fock and density functional theory. *International Journal of Quantum Chemistry*, 44(S26):319–331, 1992.
- [79] Peter MW Gill, Benny G Johnson, John A Pople, and Michael J Frisch. The performance of the Becke-Lee-Yang-Parr (B-LYP) density functional theory with various basis sets. *Chemical Physics Letters*, 197(4-5):499–505, 1992.
- [80] Gustavo E Scuseria. Comparison of coupled-cluster results with a hybrid of hartree-fock and density functional theory. *Journal of chemical physics*, 97(10):7528–7530, 1992.
- [81] Nevin Oliphant and Rodney J Bartlett. A systematic comparison of molecular properties obtained using hartree-fock, a hybrid hartree-fock density-functional-theory, and coupled-cluster methods. *Journal of Chemical Physics*, 100(9):6550–6561, 1994.
- [82] Adam Rettig, Diptarka Hait, Luke W Bertels, and Martin Head-Gordon. Third-order møller-plesset theory made more useful? the role of density functional theory orbitals. *Journal of Chemical Theory and Computation*, 16(12):7473–7489, 2020.
- [83] Eike Caldeweyher, Sebastian Ehlert, Andreas Hansen, Hagen Neugebauer, Sebastian Spicher, Christoph Bannwarth, and Stefan Grimme. A generally applicable atomic-charge dependent london dispersion correction. *Journal of Chemical Physics*, 150(15):154122, 2019.
- [84] Christoph Bannwarth, Sebastian Ehlert, and Stefan Grimme. Gfn2-xtb-an accurate and broadly parametrized self-consistent tight-binding quantum chemical method with multipole electrostatics and density-dependent dispersion contributions. *Journal of Chemical Theory and Computation*, 15(3):1652–1671, 2019.
- [85] Sebastian Spicher and Stefan Grimme. Robust atomistic modeling of materials, organometallic, and biochemical systems. *Angewandte Chemie International Edition*, 59(36):15665–15673, 2020.
- [86] Frank Neese, Frank Wennmohs, Ute Becker, and Christoph Riplinger. The orca quantum chemistry program package. *Journal of Chemical Physics*, 152(22):224108, 2020.

-
- [87] Sebastian Ehlert, Uwe Huniar, Jinliang Ning, James W Furness, Jianwei Sun, Aaron D Kaplan, John P Perdew, and Jan Gerit Brandenburg. r2scan-d4: Dispersion corrected meta-generalized gradient approximation for general chemical applications. *Journal of Chemical Physics*, 154(6):061101, 2021.
- [88] Jan Rezáč. Non-covalent interactions atlas benchmark data sets: Hydrogen bonding. *Journal of Chemical Theory and Computation*, 16(4):2355–2368, 2020.

Supporting Information

S1. DFT AND DC-DFT

In KS-DFT [75], the ground state energy is obtained by minimizing the following functional over densities,

$$E_v[n] = T_S[n] + U_H[n] + E_{XC}[n] + \int d^3r n(\mathbf{r})v(\mathbf{r}), \quad (1)$$

where the minimizing $n_v(\mathbf{r})$ is the ground-state density, $T_S[n]$ is the KS noninteracting kinetic energy functional, $U_H[n]$ the Hartree energy, and $E_{XC}[n]$ is the exchange-correlation (XC) functional. In practical KS calculations, the only term that of Eq. 1 that is approximated is $E_{XC}[n]$. Whatever approximation we use for XC, it yields an error in both energies and the densities, as the minimizing density in Eq. 1 is different from the exact when an approximate XC functional is used in place of its exact counterpart. We can write the error of any KS calculation as: $\Delta E = \tilde{E}[\tilde{n}] - E[n]$ as, where $E[n]$ is the exact functional (Eq. 1) and n is the exact density (we drop the v subscript for brevity), while tildes denote approximate quantities. The main idea of DC-DFT is to separate the errors in ΔE into a functional error, which is present even with the exact density [19]:

$$\Delta E_F = \tilde{E}[n] - E[n] = \tilde{E}_{XC}[n] - E_{XC}[n], \quad (2)$$

and the density-driven error is the remainder of ΔE :

$$\Delta E_D = \tilde{E}[\tilde{n}] - \tilde{E}[n]. \quad (3)$$

In most KS-DFT calculations, ΔE_F strongly dominates ΔE , implying that approximate KS densities are very good as measured by the impact on energies. A very simple example for such case is the total energy of the helium atom, whose energy computed from the PBE functional[76] barely changes after the PBE density is replaced with the exact one.[19] However, for a significant number of chemical domains (e.g., anions and barrier heights), ΔE_D can be much larger than ΔE_F [22, 24]. A good example for such a case is H^- with the PBE functional, which gives excellent energies when evaluated on the exact densities, whereas the self-consistent PBE density cannot even bind two electrons for this anion.

S2. SPOTTING AND CURING LARGE DENSITY-DRIVEN ERRORS

The following two questions are of key importance in DC-DFT: (i) How do we spot cases where ΔE_D is large?; (ii) What do we do in such cases to reduce large density-driven errors? With access to exact densities we can easily answer (i), as with those we can easily measure ΔE_D by Eq. 3. At the same time, we answered (ii), as ΔE_D entirely vanishes

with exact densities. However, exact densities are available only for small systems [19, 23], and if we always had access to exact energies and densities from highly accurate wavefunction theories, we would not even bother with KS-DFT. Thus one needs to find more practical ways to answer (i) and (ii). In relation to (i), the following quantity has been introduced:

$$\tilde{S} = \left| \tilde{E}[n^{\text{LDA}}] - \tilde{E}[n^{\text{HF}}] \right|, \quad (4)$$

and is called *density sensitivity*. \tilde{S} requires two nonempirical densities: the HF densities which are typically overlocalized and the local density approximation (LDA) densities which are typically delocalized. \tilde{S} serves as a practical measure of density sensitivity of a given reaction and approximate functional. For small molecules, \tilde{S} greater than the heuristic cutoff of 2 kcal/mol implies density sensitivity, indicating that the calculation may suffer from a large ΔE_D . What should we do then to reduce large ΔE_D ? In these cases, evaluating an approximate functional on the HF in place of self-consistent densities will likely reduce ΔE_D and likely improve the functional’s performance. This procedure, called HF-DFT, is the practice of evaluating an XC approximation on the HF density and orbitals. It had been used a long before DC-DFT was proposed [77, 78, 79, 80, 81] but only because HF densities were more convenient than self-consistent densities. KS-DFT does not always benefit from HF densities (e.g, cases where ΔE_D is small) and in Refs. [26, 31, 66] we discuss in more details formal and practical (dis)advantages of HF-DFT over SC-DFT.

S3. HF-DFT AND RELATED DC-DFT PROCEDURES

Following the idea that in some cases HF-DFT works better and SC-DFT in others, DC(HF)-DFT has been proposed [31, 66]. It is a procedure that discriminately uses HF densities, as DC(HF)-DFT becomes HF-DFT for cases that are both density-sensitive (\tilde{S} above a given cut-off value, 2 kcal/mol as discussed in Ref. [23].) and whose HF solution is not severely spin-contaminated. Otherwise, DC(HF)-DFT reverts to SC-DFT. Since HF-DFT uses the HF density as a proxy for the exact density, we only use it when there is little or no spin contamination. We calculate the expectation values of the spin-squared operator, S^2 , and only use the HF density if the $\langle S^2 \rangle$ from the HF calculation deviates less than 10% from the exact $\langle S^2 \rangle$ as discussed in Refs. [31] and [82]. Otherwise, we use the self-consistent density. As it combines the best of both of them, DC(HF)-DFT comes with a range of advantages over both HF-DFT and SC-DFT as further detailed in Ref. [31]. These advantages come at a small extra cost, as for DC(HF)-DFT we need to run up to three distinct self-consistent cycles to obtain the three densities (the SC density for a given functional and those from HF and LDA needed to calculate \tilde{S}). While

we consider DC(HF)-DFT the state-of-the-art DC-DFT-based procedure, in the present work we want a simple DC framework that can be applied easily and routinely. r²SCAN and SCAN are general-purpose functionals and the ease of their use would be undermined if tandem with DC(HF)-DFT, which would require always calculating \tilde{S} and possibly making adjustments to its cut-off value. For this reason and encouraged by the very good performance of HF-DFT with SCAN-like functionals[32, 42], we employ HF-DFT as a DC-DFT procedure throughout this work. As said, constructing the robust and accurate HF-r²SCAN-DC4 is the central objective of this work. While the resulting HF-r²SCAN-DC4 can be routinely used by applying it to HF orbitals without ever needing to calculate \tilde{S} of a given reaction, the use of \tilde{S} is vital for our training of HF-r²SCAN-DC4. Specifically, we use density-sensitivities of the training reactions to fit the D4 part of HF-r²SCAN. Further technical details of this fitting procedure will be given in the next section.

S4. OPTIMIZING DISPERSION PARAMETERS

D4 stands for the generally applicable atomic-charge dependent London dispersion correction term developed by Grimme and co-workers.[83]. It has 4 functional-dependent parameters s_6 , s_8 , a_1 , and a_2 . Following Refs. [43] and [83], we set s_6 to unity as is common for functionals that do not capture long-range dispersion interactions. We optimized the s_8 , a_1 , and a_2 parameters by minimizing the mean absolute error (MAE) for the density-insensitive GMTKN55 reactions by following the DC-DFT ideas of Ref. [26]. However, the density-insensitive reactions in GMTKN55 largely fall into two distinct parameter groups for HF-r²SCAN: s_8 has a negative value for noncovalent interactions, but is positive for the rest. The difference in MAE of density-insensitive cases between those two groups is miniscule (below 0.01 kcal/mol). For example, $(s_8, a_1, a_2) = (-0.20, 0.07, 6.50)$ gives 1.209 kcal/mol for the density-insensitive MAE while $(0.39, 0.09, 7.02)$ gives 1.210 kcal/mol. Such a difference is not meaningful. Small changes in computational details such as DFT grid information, two-electron operator fitting scheme, etc. changes the values of the parameters, since reaction energy errors and density sensitivity values can be changed by 0.01 kcal/mol with those changes. To eliminate this ambiguity while ensuring accuracy in water interactions, we include the density-insensitive water...water pair interaction energy as a validation set. The two most stable water hexamers, the prism and the cage, are used to calculate the water...water 2-body interaction energy error per dimer, relative to CCSD(T)/CBS in Ref. [13]. We multiply its weight by 7 in our loss function to produce a better defined minimum and regularize the result (if we used 1, it has no effect; if we used 1000, we simply fit to this data). We can rationalize this value by noting that the mean density

sensitivity of these pairs is 0.27 kcal/mol, which is about 1/7th of our density sensitivity cutoff. The resulting values for the three parameters are: -0.36, 0.23, 5.23 for s_8 , a_1 , and a_2 each.

S5. ADDITIONAL RESULTS FOR THE GMTKN55 DATABASE

In Table S1, we list MAE (kcal/mol) of SC-r²SCAN and HF-r²SCAN functional with and without the dispersion correction for the chemically diverse GMTKN55 database [40]. def2-QZVPPD basis set is used.

S6. ADDITIONAL RESULTS FOR WATER CLUSTERS

i. MD generated dimer structures

The structures used in Figures 1(e) and 2(b), have been obtained from the molecular dynamics (MD) simulation. The simulation is performed within the XTb package[84] and the GFN-FF force field[85], enabling us to generate the various dimer configurations. The total simulation time is 50 ps, while the integration time step is 4.0 fs using a Berendsen thermostat at 298K in the NVT ensemble. We use the SHAKE algorithm to constrain bonds, for all bonds with 4 amu for the hydrogen atom mass. Then, we randomly selected 110 different configurations for the water...water dimer and 80 for the water...Aspirin dimers. The reference interaction energies are then calculated with DLPNO-CCSD(T)-F12/TightPNO method with the aug-cc-pvqz basis set for water...water dimers and aug-cc-pvtz basis set for water...Aspirin dimers.

ii. Many-body expansion of the interaction energy

The interaction energy can be decomposed into 2-body, 3-body, etc. by using the many-body expansion.[67] For example, the interaction energy of the water hexamer can be divided into K -body contributions,

$$E_{int} = E_{int}^{2-body} + E_{int}^{3-body} + \dots + E_{int}^{6-body} \quad (5)$$

where E_{int}^{K-body} is the K -body interaction energy which can be calculated from the total energy of the subcluster of the N -mer cluster: [67]

$$E_{int}^{K-body} = \sum_{i=1}^K (-1)^{K-i} \binom{N-i}{K-i} S_{tot}^i \quad (6)$$

where S_{tot}^i stands for the total energy summation of the i -th monomer subcluster.

S7. ADDITIONAL RESULTS FOR THE COMPLEXES WITH CYTOSINE

In Figure 4, we plot the cytosine...water and cytosine...cytosine interaction energy error plots, respectively. The 14 different cytosine...cytosine configurations and reference interaction energies are from Ref. [56]. For the cytosine...water interaction, we place two water molecules around 14 different cytosine...cytosine structures and optimized the water molecular coordinates while fixing the cytosine...cytosine coordinates. B3LYP functional is used for the geometry optimization. DLPNO-CCSD(T)-F12/aug-cc-pvqz with TightPNO is used as a reference cytosine...water interaction energy with the ORCA package.[86]

S8. INTERACTIONS INCLUDING WATER

For the calculations shown in Figure 5(a), we combined energies of 145 reactions involving water. They include: the water hexamer isomerization in Figure 1(b), the water binding energy of WATER27 dataset in Figure 1(c), the water 20-mer isomerization in Figure 1(d), the water dimer stationary point geometry interaction energy in Figure 2(a), and water...small organic molecule interaction energy in Figure S11.

	HF-r ² SCAN	HF-r ² SCAN-DC4	SC-r ² SCAN	SC-r ² SCAN-D4
MAE (all)	2.82	2.42	2.98	2.91
MoM (all)	2.77	2.32	3.24	3.10
WTMAD-2 (all) [†]	8.54	5.18	8.65	7.10
basic [†]	4.19	4.09	4.92	4.89
react. [†]	8.10	5.85	9.48	8.11
barriers [†]	7.18	8.04	13.11	13.56
inter. NCI [†]	13.27	4.88	10.96	6.77
intra. NCI [†]	11.96	4.80	8.67	5.87
		basic		
W4	6.92	6.41	3.81	3.86
G21EA	4.59	4.55	3.88	3.86
G21IP	4.56	4.54	4.66	4.64
DIPCS10	4.42	4.34	5.18	5.14
PA26	1.85	1.80	2.44	2.40
SIE4x4	12.14	12.24	17.93	17.98
ALKBDE10	4.71	4.66	5.01	5.00
YBDE18	3.89	3.67	3.89	3.36
AL2X6	1.02	1.23	0.93	1.58
HEAVYSB11	5.60	4.32	3.91	3.15
NBPRC	1.24	1.24	1.60	1.52
ALK8	1.73	1.77	2.73	2.88
RC21	2.82	2.39	4.57	4.95
G2RC	4.24	4.56	5.38	5.55
BH76RC	2.57	2.56	2.97	2.97
FH51	1.71	1.72	2.19	2.16
TAUT15	1.11	1.10	1.58	1.57
DC13	8.96	7.97	8.63	7.72
		react.		
MB16-43	10.48	10.79	12.59	14.08
DARC	3.82	2.00	3.71	2.70
RSE43	0.96	0.96	1.55	1.51
BSR36	3.20	0.16	2.32	0.48
CDIE20	1.21	1.13	1.63	1.61
ISO34	1.52	1.36	1.36	1.29
ISOL24	4.34	3.01	4.96	4.10
C60ISO	3.52	3.88	5.35	5.57
PArel	1.14	1.15	1.55	1.54
		barriers		
BH76	2.85	2.90	6.87	6.98
BHPERI	4.14	5.95	3.86	4.65
BHDIV10	3.49	3.89	5.98	6.11
INV24	1.35	1.34	1.22	1.14
BHROT27	0.62	0.64	0.76	0.76
PX13	4.74	5.03	8.75	8.83
WCPT18	2.73	3.18	5.81	5.99
		inter. NCI		
RG18	0.26	0.10	0.23	0.16
ADIM6	2.65	0.45	1.98	0.34
s22	1.55	0.42	1.18	0.24
S66	1.42	0.30	1.02	0.26
HEAVY28	0.71	0.37	0.52	0.30
WATER27	3.94	1.01	4.24	6.30
CARBHB12	0.63	0.60	0.88	1.06
PNICO23	0.71	0.29	0.64	0.76
HAL59	1.01	0.41	0.99	0.80
AHB21	0.63	0.68	1.15	1.35
CHB6	0.46	0.58	0.49	0.52
IL16	1.88	0.43	0.33	0.64
		intra. NCI		
IDISP	6.83	1.58	10.67	7.21
ICONF	0.30	0.23	0.32	0.29
ACONF	0.54	0.16	0.38	0.18
Amino20x4	0.40	0.27	0.26	0.19
PCONF	1.22	0.44	1.05	0.41
MCONF	0.94	0.21	0.63	0.45
SCONF	0.57	0.19	0.37	0.51
UPU23	1.18	0.38	0.95	0.41
BUT14DIOL	0.40	0.14	0.14	0.23

Table S1: Mean-absolute-error (MAE) of GMTKN55 for selected functionals for the individual datasets in the GMTKN55 database. WTMAD-2 value from Ref. [40] and mean of means (MoM) values are also noted. WTMAD-2 values for individual GMTKN55 categories [†] (basic properties and reaction energies for small systems (basic), reaction energies for large systems and isomerisation reactions (react.), reaction barrier heights (barriers), intermolecular noncovalent interactions (inter. NCI), and intramolecular noncovalent interactions (intra. NCI)) are also added. All units are kcal/mol.




Figure S1: Density sensitivity plot of (left) Smith dimer configuration and (right) MD generated dimer configuration in Figure 2.




Figure S2: (left) Smith dimer interaction energy errors and (right) reference dimer interaction energies of MD generated dimer corresponding to Figure 2.




Figure S3: Interaction energy error plot of cytosine... cytosine compounds for $HF-r^2SCAN-DC4/aug-cc-pvqz$ and $HF-SCAN$. Reference $HF/7Z-CP+MP2/CBS(6,7)-CP+dCC(cc-pVTZ-F12)$ energies are from Ref. [56].




Figure S4: Binding energy error of WATER27 dataset for $SC-r^2SCAN$ and $SC-r^2SCAN-D4$. The x-axis indicates the reaction number in WATER27 and the detailed information of reaction including geometries can be found in the GMTKN55 database.[40] $D4$ parameters of $SC-r^2SCAN-D4$ are from Ref. [87].




Figure S5: The K -body energy plot corresponding to Figure 3. For comparison, SC -SCAN and HF -SCAN-DC4 is additionally plotted in the total subplot. $def2qzvppd$ basis set is used.




Figure S6: K -body percentage contributions to reference interaction energies of water hexamers (Figure 3).




Figure S7: Density sensitivity \bar{S} value for isomer energies for the water 20-mers corresponding to Figure 1(d).




Figure S8: *Aspirin*...*water* interaction including $SC-r^2SCAN-D_4$. The x -axis is the oxygen...oxygen distance between oxygen in the water and the specified oxygen in *Aspirin* (see inset of Fig. 1(e)). *aug-cc-pvtz* basis set is used.




Figure S9: (left) Water hexamer isomer relative energy compared to its global minimum geometry prism structure; (right) their \tilde{S} value calculated from the r^2 SCAN functional. Geometries are from Ref. [13]. MAEs of each functional are 0.61, 0.04, 0.13, 0.36, and 0.43 kcal/mol compared to the CCSD(T)/CBS from Ref. [61]. The ordering is the same as the legend ordering. For HF- r^2 SCAN-DC4, we used the dispersion parameters from Section S4, whilst for SC- r^2 SCAN-D4, we took the D4 parameters from Ref. [87]. The aug-cc-pvqz basis set was used for the calculations.




Figure S10: The hexagon plot with WTMAD-2 (kcal/mol) for all GMTKN55 and its categories for selected functionals. (See Ref. [40] for the detailed description of the categories).




Figure S11: Interaction energy curve for two non-hydrogen bonded systems. Geometries are from the HB375 dataset described in Ref. [88]. The x-axis show the scaled distance factor, f , which is used to make a translated vector t , $t = \frac{v}{|v|}(f-1)r_{ref}$ where v is the bond direction vector and r_{ref} is the distance between the hydrogen and the electron-donor atom. Detailed information about the x-axis can be found in Ref. [88].