

Bergman theory for the inhomogeneous Cimmino system

José Oscar González-Cervantes⁽¹⁾, Dante Arroyo-Sánchez⁽¹⁾
and
Juan Bory-Reyes^{(2)*}

⁽¹⁾ Departamento de Matemáticas, ESFM-Instituto Politécnico Nacional. 07338,
Ciudad México, México

Emails: jogc200678@gmail.com, tlacaelela.s68@gmail.com

⁽²⁾ SEPI, ESIME-Zacatenco-Instituto Politécnico Nacional. 07338, Ciudad
México, México

Email: juanboryreyes@yahoo.com

Abstract

We first prove a Cauchy's integral theorem and Cauchy type formula for certain inhomogeneous Cimmino system from quaternionic analysis perspective. The second part of the paper directs the attention towards some applications of the mentioned results, dealing in particular with four kinds of weighted Bergman spaces, reproducing kernels, projection and conformal invariant properties.

Keywords: Cimmino system; Bergman spaces; Reproducing kernel; Conformally covariant and invariant properties

AMS Subject Classification (2020): 30E20, 30G35, 32A25, 32A36, 47B32.

1 Introduction

Gianfranco Cimmino (12 March, 1908 - 30 May, 1989), was the first to consider the homogeneous first order linear partial differential equations (named in his honour

*corresponding author

Cimmino system):

$$\begin{cases} \frac{\partial f_0}{\partial x_0} + \frac{\partial f_2}{\partial x_2} - \frac{\partial f_1}{\partial x_1} - \frac{\partial f_3}{\partial x_3} = 0 \\ \frac{\partial f_0}{\partial f_0} + \frac{\partial f_1}{\partial f_1} - \frac{\partial f_2}{\partial f_2} + \frac{\partial f_3}{\partial f_3} = 0 \\ \frac{\partial x_1}{\partial f_0} + \frac{\partial x_0}{\partial f_3} - \frac{\partial x_3}{\partial f_1} + \frac{\partial x_2}{\partial f_2} = 0 \\ \frac{\partial x_2}{\partial f_0} + \frac{\partial x_1}{\partial f_1} - \frac{\partial x_3}{\partial f_2} - \frac{\partial x_0}{\partial f_3} = 0 \\ \frac{\partial f_0}{\partial x_3} + \frac{\partial f_1}{\partial x_2} + \frac{\partial f_2}{\partial x_1} + \frac{\partial f_3}{\partial x_0} = 0, \end{cases} \quad (1)$$

for $f_m, m = 1, 2, 3, 4$ continuously differentiable \mathbb{R} -valued unknown functions in a domain $\Omega \subset \mathbb{R}^4$, all of whose solutions are harmonic functions, see [10].

The solution theory of the Cimmino system can be seen as a generalization of the complex analysis of two complex variables. In [11] Dragomir and Lanconelli made a detailed investigation of Cimmino system, succeeded (among many other results) in extend most of the highlights from the holomorphic function theory to the Cimmino system theory, in particular they obtain conditions for the resolvability of the Dirichlet problem for Cimmino system.

Although nowadays the Cimmino system theory has been attacked with greater hope of success, see [1, 2, 4, 7, 9, 29], the investigation of inhomogeneous Cimmino system is limited for right-hand sides depending only on the real variables $x_m, m = 1, 2, 3, 4$, see [27, 28].

The history of Bergman theory goes back to the work of Stefan Bergman (5 May, 1895 – 6 June, 1977) in the early twentieth century, where the first systematic treatment of the subject of reproducing kernel for holomorphic functions of one or several complex variables was establish, see [5] for the classical bibliography.

Since then, efforts were made to develop a similar theory of functions for different linear differential equations in a fixed domain. Some standard works here are [6, 12, 21] and the references therein, which contain a broad summary and historical notes of the subject.

At present, systematic investigation of Bergman theory, mainly applied quaternionic methods, is conducted for suitably chosen classes of systems of partial differential equations in a series of papers, see [14–19] and references therein.

In this work we prove a Cauchy integral theorem and a Cauchy type formula for the following inhomogeneous Cimmino system:

$$\begin{cases} \frac{\partial f_0}{\partial x_0} + \frac{\partial f_2}{\partial x_2} - \frac{\partial f_1}{\partial x_1} - \frac{\partial f_3}{\partial x_3} = q_0 f_0 + q_2 f_2 - q_1 f_1 - q_3 f_3 \\ \frac{\partial f_0}{\partial f_0} + \frac{\partial f_1}{\partial f_1} - \frac{\partial f_2}{\partial f_2} + \frac{\partial f_3}{\partial f_3} = q_1 f_0 + q_0 f_1 - q_3 f_2 + q_2 f_3 \\ \frac{\partial x_1}{\partial f_0} + \frac{\partial x_0}{\partial f_3} - \frac{\partial x_3}{\partial f_1} + \frac{\partial x_2}{\partial f_2} = q_2 f_0 + q_1 f_3 - q_3 f_1 - q_0 f_3 \\ \frac{\partial x_2}{\partial f_0} + \frac{\partial x_1}{\partial f_1} - \frac{\partial x_3}{\partial f_2} - \frac{\partial x_0}{\partial f_3} = q_3 f_0 + q_2 f_1 + q_1 f_2 + q_0 f_3. \end{cases} \quad (2)$$

for four desired continuously differentiable \mathbb{R} -valued functions f_m depending on four real variables $x_m, m = 1, 2, 3, 4$ and a fix four-tuple $(q_0, q_1, q_2, q_3) \in \mathbb{R}^4$.

Together with the Cauchy integral theorem and Cauchy type formula versions alluded to earlier, we shall concern ourselves with some applications, in particular with four kinds of weighted Bergman spaces, reproducing functions, projections and conformal invariant properties.

2 Preliminaries

The skew-field of quaternions, customarily denoted by \mathbb{H} ((in honor of William Rowan Hamilton (August 4, 1805- September 2, 1865)) [20]), is a four dimensional associative algebra over \mathbb{R} with generators 1 and the quaternionic imaginary units e_1, e_2, e_3 under the multiplication rules:

$$e_1^2 = e_2^2 = e_3^2 = -1, e_1e_2 = e_3, e_2e_3 = e_1, e_3e_1 = e_2.$$

For any $q \in \mathbb{H}$ it can be written as these linear combinations of 1, e_1, e_2 and e_3 , namely $q = q_0 + \vec{q} = q_0 + q_1e_1 + q_2e_2 + q_3e_3$, where $q_k \in \mathbb{R}$ for $k = 0, 1, 2, 3$. Note that a quaternion q may also be viewed as a vector $(q_0, q_1, q_2, q_3) \in \mathbb{R}^4$.

The rules for multiplication among e_1, e_2 and e_3 are enough, with the distributive law, to multiply all quaternions.

Quaternionic conjugation and the module of q are defined respectively by $\bar{q} := q_0 - \vec{q}$ and $|q| := \sqrt{q\bar{q}} = \sqrt{q_0^2 + q_1^2 + q_2^2 + q_3^2}$.

The dot product of two quaternions $q, r \in \mathbb{H}$ is given by

$$\langle q, r \rangle := \frac{1}{2}(\bar{q}r + \bar{r}q) = \frac{1}{2}(q\bar{r} + r\bar{q}).$$

A four-tuple $\psi = \{\psi_0, \psi_1, \psi_2, \psi_3\} \in \mathbb{H}^4$ is called structural set if $\langle \psi_k, \psi_m \rangle = \delta_{k,m}$, for $k, m = 0, 1, 2, 3$ and any quaternion q can be rewritten as $q_\psi := \sum_{k=0}^3 q_k \psi_k$, where $q_k \in \mathbb{R}$ for all k . Note that any structural set can be thought of as an orthonormal (in the usual Euclidean sense) basis in \mathbb{R}^4 . The notion of structural set goes back as far as [22].

Given $q, x \in \mathbb{H}$ we follow the notation used in [25] to write

$$\langle q, x \rangle_\psi = \sum_{k=0}^3 q_k x_k,$$

where $q_\psi = \sum_{k=0}^3 q_k \psi_k$ and $x_\psi = \sum_{k=0}^3 x_k \psi_k$ with $q_k, x_k \in \mathbb{R}$ for all k .

Functions f defined in a bounded domain $\Omega \subset \mathbb{H} \cong \mathbb{R}^4$ with value in \mathbb{H} are considered. They may be written as: $f = \sum_{k=0}^3 f_k \psi_k$, where $f_k, k = 0, 1, 2, 3$, are \mathbb{R} -valued functions in Ω . Properties as continuity, differentiability, integrability and so on, which as ascribed to f have to be posed by all components f_k . We will follow standard notation, for example $C^1(\Omega, \mathbb{H})$ denotes the set of continuously differentiable \mathbb{H} -valued functions defined in Ω .

Furthermore, the conformal mappings in \mathbb{R}^4 (also referred to as the Möbius transformations), are represented by

$$T(x) = (ax + b)(cx + d)^{-1}, \quad (3)$$

where $a \neq 0$, if $c = 0$ and $b - ac^{-1}d \neq 0$, if $c \neq 0$, see [3] for more details.

In 1930's quaternionic functions of one quaternionic variable were investigated mainly by Karl Rudolf Fueter ((30 June, 1880 – 9 August, 1950) [13]), who was able to develop the appropriate generalization of the Cauchy-Riemann equations to the quaternionic case. The Cauchy-Fueter equations (as they are now called in honor of their inventor) were not developed until 1935, almost a century after Hamilton's discovery of quaternions. This function theory (called *Quaternionic analysis*) has become popular again since 1970's, see e.g. the very well known paper [26] of Anthony Sudbery (29 July, 1943 - Unknown).

Nowadays, quaternionic analysis is regarded as a broadly accepted branch of classical analysis offering a successful generalization of complex analysis. It relies heavily around the concept of ψ -hyperholomorphic functions, i.e., null solutions of the so-called ψ -Fueter operator related to the structural set ψ .

$${}^\psi\mathcal{D} = \frac{\partial}{\partial x_0} + \psi_1 \frac{\partial}{\partial x_1} + \psi_2 \frac{\partial}{\partial x_2} + \psi_3 \frac{\partial}{\partial x_3},$$

which represents a natural generalization of the complex Cauchy-Riemann operator, see [23–25] for more details.

In [1, 2, 7], motivated by [27, Subsection 6.4-6.5] it is shown how Cimmino system theory can be successfully embedded into that of ψ -hyperholomorphic functions for the simplest and therefore fundamental case of structural set

$$\psi = \{1, e_1, -e_2, e_3\}.$$

The theory of ψ -hyperholomorphic functions has recently rekindled by the study of a quaternionic valued function theory associated to a domain $\Omega \subset \mathbb{H}$ and induced by the left- and the right- q - ψ -Fueter type operators:

$${}^\psi\mathcal{D}[f] := {}^\psi\mathcal{D}[f] + qf \quad \text{and} \quad \mathcal{D}_q^\psi[f] := \mathcal{D}^\psi[f] + fq, \quad f \in C^1(\Omega, \mathbb{H}),$$

which the first author of the present paper has studied in some detail [14].

We will end this section by recalling relevant material from [14] and provide some additional notations and terminology that will be used in the sequel, thus making our exposition self-contained. To this end, let us introduce the following notations:

- $d\lambda_q^\psi := d\lambda_q^\psi(x) = e^{2\langle q, x \rangle_\psi} d\mu_x$,
- $d\mu_x$ denotes the oriented volume element on Ω
- $\nu_q^\psi(x) := e^{2\langle q, x \rangle_\psi} \sigma_x^\psi$,

- $\sigma_x^\psi := -\text{sgn}\psi \left(\sum_{k=0}^3 (-1)^k \psi_k d\hat{x}_k \right)$ stands for the quaternionic differential form of the 3 dimensional volume in \mathbb{R}^4 according to ψ , with $d\hat{x}_i = dx_0 \wedge dx_1 \wedge dx_2 \wedge dx_3$ omitting factor dx_i ,
- $\text{sgn}\psi$ is equal to 1, or -1 , provided that ψ and $(1, e_1, e_2, e_3)$ have the same orientation or not.

In addition, we write $K_q^\psi(y-x) := e^{\langle q, y-x \rangle_\psi} K_\psi(y-x)$, where

$$K_\psi(\tau-x) := \frac{1}{2\pi^2} \frac{\overline{\tau_\psi - x_\psi}}{|\tau_\psi - x_\psi|^4},$$

for the ℓ - q - ψ -hyperholomorphic Cauchy Kernel.

In particular, when $\psi = \{1, e_1, -e_2, e_3\}$ we denote it briefly by

$$\begin{aligned} K_q(y-x) &= e^{\langle q, y-x \rangle} K(y-x) \\ &= K_0^q(y-x) + K_1^q(y-x)e_1 + K_2^q(y-x)e_2 + K_3^q(y-x)e_3 \\ &= (K_0^q(y-x), K_1^q(y-x), K_2^q(y-x), K_3^q(y-x)). \end{aligned} \quad (4)$$

Proposition 2.1. *Let $\Omega \subset \mathbb{H}$ be a domain with a 3-dimensional compact smooth boundary, denoted by $\partial\Omega$ and fix $q \in \mathbb{H}$. Then*

$$\begin{aligned} d(f\nu_q^\psi(x)g) &= (\mathcal{D}_q^\psi[f]g + f_q^\psi \mathcal{D}[g])e^{2\langle q, x \rangle_\psi}, \\ \int_{\partial\Omega} f(x)\nu_q^\psi(x)g(x) &= \int_{\Omega} (\mathcal{D}_q^\psi[f](y)g(y) + f(y)_q^\psi \mathcal{D}[g](y))d\lambda_q^\psi(y) \end{aligned} \quad (5)$$

and

$$\begin{aligned} & \int_{\partial\Omega} (K_q^\psi(y-x)\sigma_y^\psi f(y) + g(y)\sigma_y^\psi K_q^\psi(y-x)) \\ & - \int_{\Omega} (K_q^\psi(y-x)_q^\psi \mathcal{D}[f](y) + \mathcal{D}_q^\psi[g](y)K_q^\psi(y-x))d\mu_y \\ & = \begin{cases} f(x) + g(x), & x \in \Omega, \\ 0, & x \in \mathbb{H} \setminus \overline{\Omega}, \end{cases} \end{aligned} \quad (6)$$

for all $f, g \in C^1(\overline{\Omega}, \mathbb{H})$.

Proposition 2.2. *Let $q, r \in \mathbb{H}$ and ψ be a structural set. Suppose $\Omega, \Xi \subset \mathbb{H}$ be two conformally equivalent domains such that $T(\Xi) = \Omega$ and set $y = T(x)$. Then*

$$\psi \mathcal{D}_x \circ [e^{\langle r-q, x \rangle_\psi} A_T f \circ T] = e^{\langle r-q, x \rangle_\psi} (B_T \circ T)_{(\delta_{T,r}^\psi)} \mathcal{D}_y [f] \circ T, \quad (7)$$

where

$$A_T(x) := \begin{cases} \bar{d}, & \text{if } c = 0; \\ \frac{\overline{cxV^{-1} + dV^{-1}}}{|cxV^{-1} + dV^{-1}|^4}, & \text{if } c \neq 0, \end{cases} \quad x \in \Omega$$

$$B_T(y) := \begin{cases} \bar{a}, & \text{if } c = 0; \\ -\bar{c}(y - ac^{-1})|y - ac^{-1}|^4, & \text{if } c \neq 0, \end{cases} \quad y \in \Xi,$$

$$\delta_{T,r}(y) := \begin{cases} (\bar{a})^{-1}r\bar{d}, & \text{if } c = 0; \\ -\frac{(y - ac^{-1})cr\bar{V}(y - ac^{-1})}{|c|^2|y - ac^{-1}|^4}, & \text{if } c \neq 0, \end{cases} \quad \forall y \in \Xi,$$

where $V = b - ac^{-1}d$ and $y = T(x)$ for $x \in \Omega$.

Under the conditions stated above let us introduce two more piece of notation:

$$C_T(x) := \begin{cases} \frac{|a|^2}{|d|}d^{-1}, & \text{if } c = 0, \\ \frac{|c|^2}{|V|}V^{-1}\frac{cxV^{-1} + dV^{-1}}{|cxV^{-1} + dV^{-1}|^4}, & \text{if } c \neq 0, \end{cases} \quad (8)$$

$$\rho_T(x) := \begin{cases} 1, & \text{if } c = 0, \\ \frac{1}{|cxV^{-1} + dV^{-1}|^2}, & \text{if } c \neq 0. \end{cases} \quad (9)$$

Definition 2.3. Let $\Omega \subset \mathbb{H}$ be a domain. Let ψ be a structural set and $q \in \mathbb{H}$.

1. The quaternionic right-linear space ${}^\psi_q\mathcal{A}(\Omega) = Ker_q{}^\psi\mathcal{D} \cap C^1(\Omega, \mathbb{H}) \cap \mathcal{L}_2(\Omega, \mathbb{H})$ is called ℓ - q - ψ -hyperholomorphic Bergman space associated to Ω and denote

$$\|f\|_{{}^\psi_q\mathcal{A}(\Omega)} = \left(\int_{\Omega} |f|^2 d\mu \right)^{\frac{1}{2}},$$

for all $f \in {}^\psi_q\mathcal{A}(\Omega)$. By $\mathcal{A}_{\bar{q}}^{\bar{\psi}}(\Omega) = \mathfrak{M}_{\bar{q}}^{\bar{\psi}}(\Omega) \cap \mathcal{L}_2(\Omega, \mathbb{H})$ we mean the right- \bar{q} - $\bar{\psi}$ -hyperholomorphic Bergman space associated to Ω .

2. The set ${}_{\lambda_q}{}^\psi\mathcal{A}(\Omega)$ consists of $f \in {}^\psi_q\mathfrak{M}(\Omega)$ such that

$$\|f\|_{{}_{\lambda_q}{}^\psi\mathcal{A}(\Omega)}^2 := \int_{\Omega} |f(x)|^2 d\lambda_q^\psi(x) < \infty.$$

Define

$$\langle f, g \rangle_{{}_{\lambda_q}{}^\psi\mathcal{A}(\Omega)} := \int_{\Omega} \bar{f}g d\lambda_q^\psi, \quad {}^\psi_q\mathcal{S}[f](x) := e^{\langle q, x \rangle_\psi} f(x), \quad x \in \Omega,$$

for all $f, g \in {}_{\lambda_q}{}^\psi\mathcal{A}(\Omega)$.

Clearly, ${}_{\lambda_0}{}^\psi\mathcal{A}(\Omega) = {}^\psi_0\mathcal{A}(\Omega) = {}^\psi\mathcal{A}(\Omega)$, as quaternionic right-Hilbert spaces.

Proposition 2.4. *Let $\Omega \subset \mathbb{H}$ be a domain and set $q, r \in \mathbb{H}$.*

1. *The quaternionic right-linear space $({}^\psi_q \mathcal{A}(\Omega), \langle \cdot, \cdot \rangle_{\psi_q \mathcal{A}(\Omega)})$ is a quaternionic right-Hilbert space and*

$$f(x) = \int_{\Omega} {}^\psi_q \mathcal{B}_{\Omega}(x, \zeta) f(\zeta) d\mu_{\zeta}, \quad f \in {}^\psi_q \mathcal{A}(\Omega), \quad (10)$$

where ${}^\psi_q \mathcal{B}_{\Omega}(x, \cdot)$ is the Bergman kernel of ${}^\psi_q \mathcal{A}(\Omega)$. Its Bergman projection is given by

$${}^\psi_q \mathfrak{B}_{\Omega}[f](x) := \int_{\Omega} {}^\psi_q \mathcal{B}_{\Omega}(x, \zeta) f(\zeta) d\mu_{\zeta}, \quad f \in \mathcal{L}_2(\Omega, \mathbb{H}). \quad (11)$$

2. ${}^\psi_{q-r} S : {}_{\lambda_q} \mathcal{A}(\Omega) \rightarrow {}_{\lambda_r} \mathcal{A}(\Omega)$ is an isometric isomorphisms of quaternionic right-Hilbert spaces given by

$${}^\psi_{q-r} S[f](x) = e^{\langle q-r, x \rangle} f(x), \quad x \in \Omega$$

and

$${}_{\lambda_r} \mathcal{B}_{\Omega}(x, \xi) = e^{\langle q-r, x+\xi \rangle} {}_{\lambda_q} \mathcal{B}_{\Omega}(x, \xi),$$

where ${}_{\lambda_r} \mathcal{B}_{\Omega}$ is the kernel of ${}_{\lambda_r} \mathcal{A}(\Omega)$.

Proposition 2.5. *Given $q, r \in \mathbb{H}$ and let $\psi \subset \mathbb{H}$ be a structural set. Assume Ω and Ξ be two conformally equivalent domains with $T(\Xi) = \Omega$.*

Let us define $\gamma_{T,r,q}(y) = e^{2(r-q, T^{-1}(y))}$ for all $y \in \Omega$ and the function space

$${}_{\delta_{T,r}} \mathcal{A}_{\gamma_{T,r,q}}(\Omega) := Ker {}_{\delta_{T,r}} \mathcal{D} \cap C^1(\Omega, \mathbb{H}) \cap \mathcal{L}_{2, \gamma_{T,r,q}}(\Omega, \mathbb{H}),$$

where

$${}_{\delta_{T,r}} \mathcal{D}[f] := {}^\psi \mathcal{D}[f] + \delta_{T,r} f, \quad f \in C^1(\Omega, \mathbb{H}).$$

The function space ${}_{\delta_{T,r}} \mathcal{A}_{\gamma_{T,r,q}}(\Omega)$ is equipped with the norm and the inner product inherited from the weighted space $\mathcal{L}_{2, \gamma_{T,r,q}}(\Omega, \mathbb{H})$:

$$\|f\|_{{}_{\delta_{T,r}} \mathcal{A}_{\gamma_{T,r,q}}(\Omega)} := \left[\int_{\Omega} \|f\|^2 \gamma_{T,r,q} d\mu \right]^{\frac{1}{2}}, \quad \langle f, g \rangle_{{}_{\delta_{T,r}} \mathcal{A}_{\gamma_{T,r,q}}(\Omega)} := \int_{\Omega} \bar{f} g \gamma_{T,r,q} d\mu,$$

for all $f, g \in {}_{\delta_{T,r}} \mathcal{A}_{\gamma_{T,r,q}}(\Omega)$.

On the other hand, set ${}^\psi_q \mathcal{A}_{\rho_T}(\Xi) := Ker {}^\psi_q \mathcal{D} \cap C^1(\Xi, \mathbb{H}) \cap \mathcal{L}_{2, \rho_T}(\Xi, \mathbb{H})$. Then

$$\|f\|_{{}^\psi_q \mathcal{A}_{\rho_T}(\Xi)} = \left[\int_{\Xi} \|f\|^2 \rho_T d\mu \right]^{\frac{1}{2}}, \quad \langle f, g \rangle_{{}^\psi_q \mathcal{A}_{\rho_T}(\Xi)} = \int_{\Xi} \bar{f} g \rho_T d\mu,$$

for all $f, g \in {}_q^\psi \mathcal{A}_{\rho_T}(\Xi)$.

The quaternionic right-linear operator $\mathcal{J} : {}_{\delta_{T,r}}^\psi \mathcal{A}_{\gamma_{T,r,q}}(\Omega) \rightarrow {}_q^\psi \mathcal{A}_{\rho_T}(\Xi)$ given by

$$\mathcal{J}[f](x) = e^{\langle r-q, x \rangle_\psi} C_T(x) f \circ T(x), \quad x \in \Xi, \quad (12)$$

for all $f \in {}_{\delta_{T,r}}^\psi \mathcal{A}_{\gamma_{T,r,q}}(\Omega)$ is an isometric isomorphism of quaternionic right-Hilbert spaces. What is more,

$${}_q^\psi \mathcal{B}_{\Xi, \rho_T}(x, \xi) = e^{\langle r-q, x+\xi \rangle_\psi} C_T(x) {}_{\delta_{T,r}}^\psi \mathcal{B}_{\Omega, \gamma_{T,r,q}}(T(x), T(\xi)) \overline{C_T(\xi)},$$

where ${}_q^\psi \mathcal{B}_{\Xi, \rho_T}$ and ${}_{\delta_{T,r}}^\psi \mathcal{B}_{\Omega, \gamma_{T,r,q}}$ stand for the Bergman kernels of ${}_q^\psi \mathcal{A}_{\rho_T}(\Xi)$ and ${}_{\delta_{T,r}}^\psi \mathcal{A}_{\gamma_{T,r,q}}(\Omega)$, respectively.

Corollary 2.6. *Let $\Omega \subset \mathbb{H}$ be a domain.*

1. *If $\Omega \subset \{x \in \mathbb{H} \mid \langle q, x \rangle_\psi < 0\}$ then ${}_q^\psi \mathcal{A}(\Omega) \subset {}_{\lambda_q^\psi} \mathcal{A}(\Omega)$ as function sets.*
2. *If $\Omega \subset \{x \in \mathbb{H} \mid \langle q, x \rangle_\psi > 0\}$ then ${}_{\lambda_q^\psi} \mathcal{A}(\Omega) \subset {}_q^\psi \mathcal{A}(\Omega)$ as function sets.*
3. *If Ω is a bounded domain then ${}_{\lambda_q^\psi} \mathcal{A}(\Omega) = {}_q^\psi \mathcal{A}(\Omega)$ as function sets.*

3 Main results

It is illuminating worth noting that while the equation ${}^\psi \mathcal{D}[f] = 0$, with $\psi = \{1, e_1, -e_2, e_3\}$, is equivalent to the Cimmino system (1), the inhomogeneous Cimmino system (2) is obtained from ${}^\psi \mathcal{D}[f] = 0$ with $q = q_0 + q_1 e_1 - q_2 e_2 + q_3 e_3 \in \mathbb{H}$ for the same structural set, so they are of great importance for the purposes of this article.

Throughout what follows, the main attention is given to the singular and important case of structural set $\psi = \{1, e_1, -e_2, e_3\}$.

The main results of the presented paper are formulated in the following theorems.

Theorem 3.1. *(Cauchy's integral theorem) Let $\Omega \subset \mathbb{R}^4$ be a domain with a 3-dimensional compact smooth boundary, denoted by $\partial\Omega$. If $f = (f_0, f_1, f_2, f_3)$ satisfies (2) then*

$$\begin{aligned} \int_{\partial\Omega} e^{2\langle q, x \rangle} [d\hat{x}_0 f_0(x) + d\hat{x}_1 f_1(x) + d\hat{x}_2 f_2(x) + d\hat{x}_3 f_3(x)] &= 0, \\ \int_{\partial\Omega} e^{2\langle q, x \rangle} [-d\hat{x}_1 f_0(x) + d\hat{x}_0 f_1(x) + d\hat{x}_3 f_2(x) - d\hat{x}_2 f_3(x)] &= 0, \\ \int_{\partial\Omega} e^{2\langle q, x \rangle} [-d\hat{x}_2 f_0(x) + d\hat{x}_1 f_3(x) + d\hat{x}_3 f_1(x) + d\hat{x}_0 f_2(x)] &= 0, \\ \int_{\partial\Omega} e^{2\langle q, x \rangle} [-d\hat{x}_3 f_0(x) + d\hat{x}_2 f_1(x) - d\hat{x}_1 f_2(x) + d\hat{x}_0 f_3(x)] &= 0. \end{aligned}$$

Proof. For $\psi = \{1, e_1, -e_2, e_3\}$ we see that $\text{sig}\psi = -1$ and $f = (f_0, f_1, f_2, f_3)$ satisfy (2) if and only if $f = f_0 + f_1 e_1 + f_2 e_2 + f_3 e_3$ belongs to $\text{Ker } {}_q^\psi \mathcal{D} \cap C^1(\Omega, \mathbb{H})$.

By (5) and taking $g = 1$ we obtain

$$\int_{\partial\Omega} \nu_q^\psi(x) f(x) = 0. \quad (13)$$

The real components of the previous identity establishes the formulas □

According to (4) with the notation

$$\begin{aligned} K_{\sigma,0}^q(y-x) &= K_0^q(y-x)d\hat{y}_0 + K_1^q(y-x)d\hat{y}_1 + K_2^q(y-x)d\hat{y}_2 + K_3^q(y-x)d\hat{y}_3, \\ K_{\sigma,1}^q(y-x) &= -K_0^q(y-x)d\hat{y}_1 + K_1^q(y-x)d\hat{y}_0 - K_2^q(y-x)d\hat{y}_3 + K_3^q(y-x)d\hat{y}_2, \\ K_{\sigma,2}^q(y-x) &= -K_0^q(y-x)d\hat{y}_2 + K_1^q(y-x)d\hat{y}_3 + K_2^q(y-x)d\hat{y}_0 - K_3^q(y-x)d\hat{y}_1, \\ K_{\sigma,3}^q(y-x) &= -K_0^q(y-x)d\hat{y}_3 - K_1^q(y-x)d\hat{y}_2 + K_2^q(y-x)d\hat{y}_1 + K_3^q(y-x)d\hat{y}_0. \end{aligned}$$

we have

Theorem 3.2. (Cauchy type formula) *Let $\Omega \subset \mathbb{R}^4$ be a domain with a 3-dimensional compact smooth boundary denoted by $\partial\Omega$. If $f = (f_0, f_1, f_2, f_3)$ satisfies (2) then*

$$\begin{aligned} & \int_{\partial\Omega} K_{\sigma,0}^q(y-x)f_0(y) - K_{\sigma,1}^q(y-x)f_1(y) - K_{\sigma,2}^q(y-x)f_2(y) - K_{\sigma,3}^q(y-x)f_3(y) \\ &= \begin{cases} f_0(x), & x \in \Omega, \\ 0, & x \in \mathbb{H} \setminus \bar{\Omega}, \end{cases} \\ & \int_{\partial\Omega} K_{\sigma,0}^q(y-x)f_1(y) + K_{\sigma,1}^q(y-x)f_0(y) + K_{\sigma,2}^q(y-x)f_3(y) - K_{\sigma,3}^q(y-x)f_2(y) \\ &= \begin{cases} f_1(x), & x \in \Omega, \\ 0, & x \in \mathbb{H} \setminus \bar{\Omega}, \end{cases} \\ & \int_{\partial\Omega} K_{\sigma,0}^q(y-x)f_2(y) - K_{\sigma,1}^q(y-x)f_3(y) + K_{\sigma,2}^q(y-x)f_0(y) - K_{\sigma,3}^q(y-x)f_1(y) \\ &= \begin{cases} f_2(x), & x \in \Omega, \\ 0, & x \in \mathbb{H} \setminus \bar{\Omega}, \end{cases} \\ & \int_{\partial\Omega} K_{\sigma,0}^q(y-x)f_3(y) + K_{\sigma,1}^q(y-x)f_2(y) - K_{\sigma,2}^q(y-x)f_1(y) + K_{\sigma,3}^q(y-x)f_0(y) \\ &= \begin{cases} f_3(x), & x \in \Omega, \\ 0, & x \in \mathbb{H} \setminus \bar{\Omega}. \end{cases} \end{aligned}$$

Proof. Set $\psi = \{1, e_1, -e_2, e_3\}$. Then

$$\begin{aligned} K_q(y-x)\sigma_y^\psi &= [K_0^q(y-x) + K_1^q(y-x)e_1 + K_2^q(y-x)e_2 + K_3^q(y-x)e_3] \\ & \quad (d\hat{y}_0 - d\hat{y}_1 e_1 - d\hat{y}_2 e_2 - d\hat{y}_3 e_3) \\ &= K_{\sigma,0}^q(y-x) + K_{\sigma,1}^q(y-x)e_1 + K_{\sigma,2}^q(y-x)e_2 + K_{\sigma,3}^q(y-x)e_3, \end{aligned}$$

where $\sigma_y^\psi := d\hat{y}_0 - e_1 d\hat{y}_1 - e_2 d\hat{y}_2 - e_3 d\hat{y}_3$.

Due to $f = (f_0, f_1, f_2, f_3)$ satisfy (2) if and only if $f = f_0 + f_1 e_1 + f_2 e_2 + f_3 e_3 \in \text{Ker } \overset{\psi}{\mathcal{D}} \cap C^1(\Omega, \mathbb{H})$ we can use (6) and $g \equiv 0$ to obtain

$$\int_{\partial\Omega} K_q^\psi(y-x)\sigma_y^\psi f(y) = \begin{cases} f(x), & x \in \Omega, \\ 0, & x \in \mathbb{H} \setminus \bar{\Omega}, \end{cases}$$

and finally, we find its real components and the proposition follows. \square

Proposition 3.3. *Given $q = (q_0, q_1, q_2, q_3), r = (r_0, r_1, r_2, r_3) \in \mathbb{R}^4$ and let $\Omega, \Xi \subset \mathbb{H}$ be two conformally equivalent domains such that $T(\Xi) = \Omega$. Then (f_0, f_1, f_2, f_3) defined on Ω satisfies the inhomogeneous Cimmino system*

$$\begin{aligned} \frac{\partial f_0}{\partial x_0} + \frac{\partial f_2}{\partial x_2} - \frac{\partial f_1}{\partial x_1} - \frac{\partial f_3}{\partial x_3} &= \delta_{T,r}^0 f_0 + \delta_{T,r}^2 f_2 - \delta_{T,r}^1 f_1 - \delta_{T,r}^3 f_3 \\ \frac{\partial f_0}{\partial x_1} + \frac{\partial f_1}{\partial x_0} - \frac{\partial f_2}{\partial x_3} + \frac{\partial f_3}{\partial x_2} &= \delta_{T,r}^1 f_0 + \delta_{T,r}^0 f_1 - \delta_{T,r}^3 f_2 + \delta_{T,r}^2 f_3 \\ \frac{\partial f_0}{\partial x_2} + \frac{\partial f_3}{\partial x_1} - \frac{\partial f_1}{\partial x_3} - \frac{\partial f_2}{\partial x_0} &= \delta_{T,r}^2 f_0 + \delta_{T,r}^1 f_3 - \delta_{T,r}^3 f_1 - \delta_{T,r}^0 f_2 \\ \frac{\partial f_0}{\partial x_3} + \frac{\partial f_1}{\partial x_2} + \frac{\partial f_2}{\partial x_1} + \frac{\partial f_3}{\partial x_0} &= \delta_{T,r}^3 f_0 + \delta_{T,r}^2 f_1 + \delta_{T,r}^1 f_2 + \delta_{T,r}^0 f_3, \end{aligned} \quad (14)$$

where $\delta_{T,r} = \delta_{T,r}^0 + \delta_{T,r}^1 e_1 - \delta_{T,r}^2 e_2 + \delta_{T,r}^3 e_3$ and $\delta_{T,r}^n$ is real-valued for $n = 0, 1, 2, 3$, if and only if the four-tuple of functions

$$\begin{aligned} g_0(x) &= e^{\langle r-q, x \rangle} (A_T^0(x) f_0(x) - A_T^1(x) f_1(x) - A_T^2(x) f_2(x) - A_T^3(x) f_3(x)), \\ g_1(x) &= e^{\langle r-q, x \rangle} (A_T^1(x) f_0(x) + A_T^0(x) f_1(x) - A_T^3(x) f_2(x) + A_T^2(x) f_3(x)), \\ g_2(x) &= e^{\langle r-q, x \rangle} (A_T^2(x) f_0(x) - A_T^1(x) f_3(x) + A_T^3(x) f_1(x) + A_T^0(x) f_2(x)), \\ g_3(x) &= e^{\langle r-q, x \rangle} (A_T^3(x) f_0(x) - A_T^2(x) f_1(x) + A_T^1(x) f_2(x) + A_T^0(x) f_3(x)), \quad x \in \Omega, \end{aligned}$$

where $A_T = A_T^0 + A_T^1 e_1 + A_T^2 e_2 + A_T^3 e_3$ and A_T^n is a real-valued for $n = 0, 1, 2, 3$, satisfies (2).

Proof. Consider $\psi = \{1, e_1, -e_2, e_3\}$. Due to $f = f_0 + f_1 e_1 + f_2 e_2 + f_3 e_3$ belongs to $\text{Ker } \overset{\psi}{\delta_{T,r}} \mathcal{D} \cap C^1(\Omega, \mathbb{H})$ we can use (7) and find its real components. \square

Definition 3.4. Let $\Omega \subset \mathbb{R}^4$ be a domain and fix $(q_0, q_1, q_2, q_3) \in \mathbb{R}^4$.

1. The Bergman space associated to Ω induced by (2), denoted by $CB_q(\Omega)$, consists of the four-tuples (f_0, f_1, f_2, f_3) of elements of $C^1(\Omega, \mathbb{R})$ that satisfy (2) and $\int_{\Omega} |f_k|^2 d\mu < \infty, \quad k = 0, 1, 2, 3$.

Define the real-bilinear form

$$\langle \cdot, \cdot \rangle_{CB_q(\Omega)} : CB_q(\Omega) \times CB_q(\Omega) \rightarrow \mathbb{R}^4$$

as follows:

$$\begin{aligned} & \langle f, g \rangle_{CB_q(\Omega)} \\ &= \left(\int_{\Omega} (f_0 g_0 + f_1 g_1 + f_2 g_2 + f_3 g_3) d\mu, \int_{\Omega} (f_0 g_1 - f_1 g_0 + f_2 g_3 - f_3 g_2) d\mu, \right. \\ & \quad \left. \int_{\Omega} (f_0 g_2 + f_3 g_1 - f_1 g_3 - f_2 g_0) d\mu, \int_{\Omega} (f_0 g_3 - f_1 g_2 + f_2 g_1 - f_3 g_0) d\mu \right), \end{aligned}$$

and set

$$\|f\|_{CB_q(\Omega)} = \left[\int_{\Omega} (f_0^2 + f_1^2 + f_2^2 + f_3^2) d\mu \right]^{\frac{1}{2}},$$

for all $f = (f_0, f_1, f_2, f_3)$, $g = (g_0, g_1, g_2, g_3) \in CB_q(\Omega)$.

Note that $\|f\|_{CB_q(\Omega)}^2 = \Pi_0(\langle f, f \rangle_{CB(\Omega)})$ where $\Pi_0 : \mathbb{R}^4 \rightarrow \mathbb{R}$ denotes the projection $\Pi_0(x_0, x_1, x_2, x_3) = x_0$ for all $(x_0, x_1, x_2, x_3) \in \mathbb{R}^4$.

2. The weighted Bergman space with wight function $e^{2\langle q, x \rangle}$ for all $x \in \Omega$ and induced by the inhomogeneous Cimmino system (2) is denoted by $CB_{\lambda_q}(\Omega)$ and it is formed by (f_0, f_1, f_2, f_3) , solutions (2) such that

$$\int_{\Omega} |f_k|^2 d\lambda_q = \int_{\Omega} |f_k(x)|^2 e^{2\langle q, x \rangle} d\mu_x < \infty, \quad k = 0, 1, 2, 3.$$

Given $f = (f_0, f_1, f_2, f_3)$, $g = (g_0, g_1, g_2, g_3) \in CB_{\lambda_q}(\Omega)$ define the real-bilinear form $\langle \cdot, \cdot \rangle_{CB_{\lambda_q}(\Omega)} : CB_{\lambda_q}(\Omega) \times CB_{\lambda_q}(\Omega) \rightarrow \mathbb{R}^4$ given by

$$\begin{aligned} & \langle f, g \rangle_{CB_{\lambda_q}(\Omega)} = \\ & \left(\int_{\Omega} (f_0 g_0 + f_1 g_1 + f_2 g_2 + f_3 g_3) d\lambda_q, \int_{\Omega} (f_0 g_1 - f_1 g_0 + f_2 g_3 - f_3 g_2) d\lambda_q, \right. \\ & \quad \left. \int_{\Omega} (f_0 g_2 + f_3 g_1 - f_1 g_3 - f_2 g_0) d\lambda_q, \int_{\Omega} (f_0 g_3 - f_1 g_2 + f_2 g_1 - f_3 g_0) d\lambda_q \right) \end{aligned}$$

and consider

$$\|f\|_{CB_{\lambda_q}(\Omega)} = \|(f_0, f_1, f_2, f_3)\|_{CB_{\lambda_q}(\Omega)} := \left[\int_{\Omega} (f_0^2 + f_1^2 + f_2^2 + f_3^2) d\lambda_q \right]^{\frac{1}{2}}.$$

Note that

$$\|f\|_{CB_{\lambda_q}(\Omega)}^2 = \Pi_0(\langle f, f \rangle_{CB_{\lambda_q}(\Omega)}),$$

for all $f \in CB_{\lambda_q}(\Omega)$.

3. Let $q, r \in \mathbb{R}^4$ and let Ω and Ξ be two conformally equivalent domains such that $T(\Xi) = \Omega$. According to Proposition 3.3 set $\gamma_{T, r, q}(y) = e^{2\langle r - q, T^{-1}(y) \rangle}$

for all $y \in \Omega$. Let $CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega)$ denote the set of four-tuples of functions in $C^1(\Omega, \mathbb{R})$ and let (f_0, f_1, f_2, f_3) defined on Ω satisfying (14) and such that

$$\int_{\Omega} |f_k|^2 \gamma_{T,r,q} d\mu < +\infty,$$

for $k = 0, 1, 2, 3$.

For $f = (f_0, f_1, f_2, f_3), g = (g_0, g_1, g_2, g_3) \in CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega)$ define the real-bilinear form

$$\langle \cdot, \cdot \rangle_{CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega)} : CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega) \times CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega) \rightarrow \mathbb{R}^4$$

as follows:

$$\begin{aligned} & \langle f, g \rangle_{CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega)} \\ &= \left(\int_{\Omega} (f_0 g_0 + f_1 g_1 + f_2 g_2 + f_3 g_3) \gamma_{T,r,q} d\mu, \int_{\Omega} (f_0 g_1 - f_1 g_0 + f_2 g_3 - f_3 g_2) \gamma_{T,r,q} d\mu, \right. \\ & \quad \left. \int_{\Omega} (f_0 g_2 + f_3 g_1 - f_1 g_3 - f_2 g_0) \gamma_{T,r,q} d\mu, \int_{\Omega} (f_0 g_3 - f_1 g_2 + f_2 g_1 - f_3 g_0) \gamma_{T,r,q} d\mu \right), \end{aligned}$$

and set $\|(f_0, f_1, f_2, f_3)\|_{CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega)}^2 = \Pi_0(\langle f, f \rangle_{CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega)})$.

At the same time, we will denote by $CB_{\rho_T}^q(\Xi)$, the set of four-tuples of elements of $C^1(\Omega, \mathbb{R})$: (f_0, f_1, f_2, f_3) such that satisfy (2) and

$$\int_{\Xi} |f_k|^2 \rho_T d\mu < +\infty,$$

for $k = 0, 1, 2, 3$. Recall that ρ_T is given by (9).

Given $f = (f_0, f_1, f_2, f_3), g = (g_0, g_1, g_2, g_3) \in CB_{\rho_T}^q(\Xi)$ define the real-form

$$\langle \cdot, \cdot \rangle_{CB_{\rho_T}^q(\Xi)} : CB_{\rho_T}^q(\Xi) \times CB_{\rho_T}^q(\Xi) \rightarrow \mathbb{R}^4$$

as follows:

$$\begin{aligned} & \langle f, g \rangle_{CB_{\rho_T}^q(\Xi)} = \\ & \left(\int_{\Xi} (f_0 g_0 + f_1 g_1 + f_2 g_2 + f_3 g_3) \rho_T d\mu, \int_{\Xi} (f_0 g_1 - f_1 g_0 + f_2 g_3 - f_3 g_2) \rho_T d\mu, \right. \\ & \quad \left. \int_{\Xi} (f_0 g_2 + f_3 g_1 - f_1 g_3 - f_2 g_0) \rho_T d\mu, \int_{\Xi} (f_0 g_3 - f_1 g_2 + f_2 g_1 - f_3 g_0) \rho_T d\mu \right), \end{aligned}$$

and

$$\|(f_0, f_1, f_2, f_3)\|_{CB_{\rho_T}^q(\Xi)}^2 = \Pi_0(\langle f, f \rangle_{CB_{\rho_T}^q(\Xi)}).$$

Proposition 3.5. Let $\Omega \subset \mathbb{H}$ be a domain and set $(q_0, q_1, q_2, q_3) \in \mathbb{R}^4$. Then $CB_q(\Omega)$ is a real-Banach space and there exist a four-tuple of real-valued reproducing functions, denoted by $(B_0(x, \cdot), B_1(x, \cdot), B_2(x, \cdot), B_3(x, \cdot)) \in CB_q(\Omega)$, such that

$$\langle (B_0(x, \cdot), -B_1(x, \cdot), -B_2(x, \cdot), -B_3(x, \cdot)), (f_0, f_1, f_2, f_3) \rangle_{CB_q(\Omega)} = (f_0(x), f_1(x), f_2(x), f_3(x)) \quad (15)$$

or equivalently

$$\begin{aligned} \int_{\Omega} (B_0(x, y)f_0(y) - B_1(x, y)f_1(y) - B_2(x, y)f_2(y) - B_3(x, y)f_3(y)) d\mu_y &= f_0(x), \\ \int_{\Omega} (B_0(x, y)f_1(y) + B_1(x, y)f_0(y) + B_2(x, y)f_3(y) - B_3(x, y)f_2(y)) d\mu_y &= f_1(x), \\ \int_{\Omega} (B_0(x, y)f_2(y) - B_1(x, y)f_3(y) + B_2(x, y)f_0(y) + B_3(x, y)f_1(y)) d\mu_y &= f_2(x), \\ \int_{\Omega} (B_0(x, y)f_3(y) + B_1(x, y)f_2(y) - B_2(x, y)f_1(y) + B_3(x, y)f_0(y)) d\mu_y &= f_3(x), \end{aligned}$$

for all $f = (f_0, f_1, f_2, f_3) \in CB_q(\Omega)$.

Additionally, if $g_k : \Omega \rightarrow \mathbb{R}$ is a measurable function and

$$\int_{\Omega} |g_k|^2 d\mu < \infty, \quad k = 0, 1, 2, 3$$

then the four-tuple of real-valued functions:

$$\langle (h_0(x), h_1(x), h_2(x), h_3(x)), (g_0, g_1, g_2, g_3) \rangle_{CB_q(\Omega)} = \quad (16)$$

for all $x \in \Omega$, belong to $CB_q(\Omega)$.

Proof. Let $\psi = \{1, e_1, -e_2, e_3\}$, we use the Fact 1. of Proposition 2.4 and denoting

$$\psi_q \mathcal{B}_{\Omega}(x, \cdot) = B_0(x, \cdot) + B_1(x, \cdot)e_1 + B_2(x, \cdot)e_2 + B_3(x, \cdot)e_3,$$

where $B_n(x, \cdot)$ are the real-valued functions for $n = 0, 1, 2, 3$ we can compute the real components of (10) and (11), which complete the proof. \square

Proposition 3.6. Let $\Omega \subset \mathbb{H}$ be a domain and set $q, r \in \mathbb{H}$. Then $(f_0, f_1, f_2, f_3) \in CB_q(\Omega)$ if and only if $(g_0, g_1, g_2, g_3) \in CB_r(\Omega)$, where $g_n(x) = e^{(q-r, x)} f_n(x)$, for all $x \in \Omega$ for $n = 0, 1, 2, 3$.

Proof. The proof is a direct computation and will be omitted. \square

Proposition 3.7. *Given $q, r \in \mathbb{R}^4$ and let Ω and Ξ be two conformally equivalent domains such that $T(\Xi) = \Omega$. Then the real normed function spaces*

$$\left(CB_{\lambda_q}(\Omega), \|\cdot\|_{CB_{\lambda_q}(\Omega)} \right), \left(CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega), \|\cdot\|_{CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega)} \right), \left(CB_{\rho_T}^q(\Xi), \|\cdot\|_{CB_{\rho_T}^q(\Xi)} \right)$$

are real-Banach function spaces that have four-tuple of reproducing functions, similar property to (15), that satisfies the projection property (16).

Proof. The proof follows via Propositions 2.4 and 2.5. \square

Proposition 3.8. *Given $q, r \in \mathbb{R}^4$ and let Ω and Ξ be two conformally equivalent domains such that $T(\Xi) = \Omega$. Then $(f_0, f_1, f_2, f_3) \in CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega)$ if and only if $(g_0, g_1, g_2, g_3) \in CB_{\rho_T}^q(\Xi)$, with*

$$\begin{aligned} g_0(x) &= e^{\langle r-q, x \rangle} [C_T^0(x)f_0 \circ T(x) - C_T^1(x)f_1 \circ T(x) - C_T^2(x)f_2 \circ T(x) - C_T^3(x)f_3 \circ T(x)], \\ g_1(x) &= e^{\langle r-q, x \rangle} [C_T^1(x)f_0 \circ T(x) + C_T^0(x)f_1 \circ T(x) - C_T^3(x)f_2 \circ T(x) + C_T^2(x)f_3 \circ T(x)], \\ g_2(x) &= e^{\langle r-q, x \rangle} [C_T^2(x)f_0 \circ T(x) - C_T^1(x)f_3 \circ T(x) + C_T^3(x)f_1 \circ T(x) + C_T^0(x)f_2 \circ T(x)], \\ g_3(x) &= e^{\langle r-q, x \rangle} [C_T^3(x)f_0 \circ T(x) - C_T^2(x)f_1 \circ T(x) + C_T^1(x)f_2 \circ T(x) + C_T^0(x)f_3 \circ T(x)], \end{aligned}$$

for all $x \in \Omega$, where C_T is expressed by $C_T = C_T^0 + C_T^1 e_1 + C_T^2 e_2 + C_T^3 e_3$ being C_T^n a real-valued function for $n = 0, 1, 2, 3$. What is more, the mapping

$$(f_0, f_1, f_2, f_3) \mapsto (g_0, g_1, g_2, g_3)$$

is a bijective real-linear operator from $CB_{\gamma_{T,r,q}}^{\delta_{T,r}}(\Omega)$ to $CB_{\rho_T}^q(\Xi)$.

Proof. Use (12) of Proposition 2.5 and compute its real components, which is our claim. \square

Corollary 3.9. *Let $\Omega \subset \mathbb{H}$ be a domain. Then*

1. *If $\Omega \subset \{x \in \mathbb{H} \mid \langle q, x \rangle < 0\}$ then $CB_q(\Omega) \subset CB_{\lambda_q}(\Omega)$ as function sets.*
2. *If $\Omega \subset \{x \in \mathbb{H} \mid \langle q, x \rangle > 0\}$ then $CB_{\lambda_q}(\Omega) \subset CB_q(\Omega)$ as function sets.*
3. *If Ω is a bounded domain then $CB_{\lambda_q}(\Omega) = CB_q(\Omega)$ as function sets.*

Proof. It follows from Corollary 2.6 choosing $\psi = \{1, e_1, -e_2, e_3\}$. \square

Declarations

Funding

Instituto Politécnico Nacional (grant numbers SIP20220017, SIP20221274) and CONACYT (grant number 1108687).

Conflict of interest

The authors declare that they have no conflict of interest regarding the publication of this paper.

Author contributions

The authors contributed equally to the manuscript and typed, read, and approved the final form of the manuscript, which is the result of an intensive collaboration.

Availability of data and material

Not applicable

Code availability

Not applicable

References

- [1] Abreu-Blaya, R., Bory-Reyes, J., Schneider, B. *On Cauchy type integrals related to the Cimmino system of partial differential equations*. Operator Theory: Advances and Applications, 242, 2014, 81–92.
- [2] Abreu-Blaya, R., Bory-Reyes, J., Guzmán-Adán, A., Schneider, B. *Boundary value problems for the Cimmino system via quaternionic analysis*. Appl. Math. Comput., 219, 2012, No. 8, 3872-3881.
- [3] Ahlfors, L. V. *Clifford numbers and Möbius transformations in \mathbb{R}^n* . Clifford Algebras and their Applications in Mathematical Physics, J. S. R. Chrisholm and A. K. Common (eds.), NATO Adv. Study Inst. Ser., Ser. C: Math. Phys. Sci., 1986, 183, Reidel, 167-175.
- [4] Avetisyan, K. *Subharmonicity and a version of Riesz theorem on harmonic conjugates*. Adv. Appl. Clifford Algebras 24 (2014), 909–919.
- [5] Bergman S. *The Kernel Function and Conformal Mapping*. Mathematical Surveys, No. 5. American Mathematical Society, New York, N. Y., 1950
- [6] Borichev A., Hedenmalm H., Zhu K. *Bergman spaces and related topics in complex analysis*. Contemporary Mathematics, 404. Israel Mathematical Conference Proceedings. American Mathematical Society, Providence, RI; Bar-Ilan University, Ramat Gan, 2006.

- [7] Bory Reyes, J., Abreu Blaya, R., Pérez-de la Rosa, M. A., Schneider, B. *Poincaré-Bertrand and Hilbert formulas for the Cauchy-Cimmino singular integrals*. Adv. Appl. Clifford Algebr. 27 (2017), no. 4, 2933–2960.
- [8] Brackx, F., Delanghe, R. *Hypercomplex function theory and Hilbert modules with reproducing kernel*. Proc. London Math. Soc. 1978, s3-37 3, 545–576.
- [9] Caramuta, P., Cialdea, A. *An application of self-conjugate differential forms to the Dirichlet problem for Cimmino system*. Complex Var. Elliptic Equ. 62, 2017, no. 7, 919–937.
- [10] Cimmino, G. *Su alcuni sistemi lineari omogenei di equazioni alle derivate parziali del primo ordine*. Rend. Sem. Mat. Univ. Padova 12, 1941, 89–113.
- [11] Dragomir, S., Lanconelli, E. *On first order linear PDF systems all of whose solutions are harmonic functions*. Tsukuba J. Math. 30(1), 2006, 149–170.
- [12] Duren P., Schuster A. *Bergman spaces*. Mathematical Surveys and Monographs, 100. American Mathematical Society, Providence, RI, 2004.
- [13] Fueter, R. *Die Funktionentheorie der Differentialgleichungen $\Delta u = 0$ und $\Delta\Delta u = 0$ mit vier reellen Variablen*. Comment. Math. Helv. 7 (1935) 307–330.
- [14] González-Cervantes J.O. *On a left- $\alpha - \psi$ -hyperholomorphic Bergman space*. Complex Var. Elliptic Equ. (2021), doi:10.1080/17476933.2021.1986033.
- [15] González-Cervantes, J. O., Bory-Reyes, J. *On Bergman spaces induced by a \mathbf{v} -Laplacian vector fields theory*. J. Math. Anal. Appl. 505 (2022), no. 2, Paper No. 125523, 18 pp.
- [16] González-Cervantes, J. O. Shapiro, M. *Hyperholomorphic Bergman spaces and Bergman operators associated with domains in \mathbb{C}^2* . Complex Anal. Oper. Theory 2 (2008), no. 2, 361–382.
- [17] González-Cervantes, J. O., Luna-Elizarrarás, M. E., Shapiro, M. *On some categories and functors in the theory of quaternionic Bergman spaces*. Adv. Appl. Clifford Algebr. 19 (2009), no. 2, 325–338.
- [18] González-Cervantes, J. O. Luna-Elizarrarás, M. E., Shapiro, M. *On the Bergman theory for solenoidal and irrotational vector fields, I: General theory*. Recent trends in Toeplitz and pseudodifferential operators, 79–106, Oper. Theory Adv. Appl., 210, Birkhäuser Verlag, Basel, 2010.
- [19] González-Cervantes, J. O., Luna-Elizarrarás, M. E., Shapiro, M. *On the Bergman theory for solenoidal and irrotational vector fields. II. Conformal covariance and invariance of the main objects*. Complex Anal. Oper. Theory 5 (2011), no. 1, 237–251.

- [20] Hamilton W. R. *Elements of quaternions*, 2 Vols (1899-1901); reprinted Chelsea, New York, 1969.
- [21] Hedenmalm H., Korenblum B., Zhu, K. *Theory of Bergman spaces*. Graduate Texts in Mathematics, 199. Springer-Verlag, New York, 2000.
- [22] Nôno, K. *Hyperholomorphic functions of a quaternion variable*. Bull. Fukuoka Univ. Ed. III 32, 21-37, 1983.
- [23] Mitelman, I., Shapiro, M. *Differentiation of the Martinelli-Bochner integrals and the notion of the hyperderivability*. Math. Nachr., 1995, 172, 211-238.
- [24] Shapiro, M. *Quaternionic analysis and some conventional theories*. In: Alpay, D. (ed.) *Operator Theory*, 1423-1446. Springer, Basel, 2015.
- [25] Shapiro M., Vasilevski, N. L. *Quaternionic ψ -monogenic functions, singular operators and boundary value problems. I. ψ -Hyperholomorphy function theory*. Compl. Var. Theory Appl. 1995, 27, 17–46.
- [26] Sudbery, A. *Quaternionic analysis*. Math. Proc. Cambridge Phil. Soc. 85 (1979) 199-225.
- [27] Tutschke, W. *Generalized analytic functions in higher dimensions*. Georgian Math. J. 14, 2007, no. 3, 581–595.
- [28] Wang, L., Xu, Z., Qiao, Y. *The mixed boundary value problem for the inhomogeneous Cimmino system*. Bound. Value Probl. 2015, 2015:13, 16 pp.
- [29] Xingya, F. *Conjugate harmonic functions of Fueter type*. Adv. Appl. Clifford Algebr. 30, 2020, no. 3, Paper No. 32, 13 pp.