

On the Origin of Coulomb Pseudopotential: Two Wrongs Make a “Right”

Tao Wang,¹ Xiansheng Cai,¹ Kun Chen,^{2,*} Boris V. Svistunov,^{1,3,†} and Nikolay V. Prokof'ev^{1,‡}

¹*Department of Physics, University of Massachusetts, Amherst, MA 01003, USA*

²*Center for Computational Quantum Physics, Flatiron Institute, 162 5th Avenue, New York, New York 10010*

³*Wilczek Quantum Center, School of Physics and Astronomy,
Shanghai Jiao Tong University, Shanghai 200240, China*

(Dated: July 13, 2022)

We explore the effect of Coulomb repulsion on the phonon-mediated BCS-type Cooper instability in the electron gas using a numeric approach that is controllably accurate at small to moderate values of the Coulomb parameter, $r_s \lesssim 2$. We demonstrate the necessity of breaking the net result into two distinctively different effects: the reduction of the Fermi-liquid renormalization factor and the change of the effective low-energy coupling. The latter quantity is shown to behave non-monotonically with an extremum at $r_s \approx 0.75$. Within the random-phase approximation, Coulomb interaction eventually starts to *enhance* the effective pairing coupling at $r_s > 2$ and the suppression of the critical temperature is entirely due to the renormalized Fermi-liquid properties. The onset of the enhancement is, however, sensitive to the approximation used for incorporating vertex corrections. Our results call for radical reconsideration of the widely accepted picture that the effect of Coulomb interactions reduces to a (weak) repulsive pseudopotential.

Introduction.—In the vast majority of low-temperature superconductors, the scenario of Cooper instability is of the emergent BCS type, implying quantitatively accurate low-energy effective description in terms of the two (partially related) parameters: the energy/frequency cutoff $\omega_0 \ll E_F$ (E_F is the Fermi energy; $\hbar = 1$) and the dimensionless effective coupling constant $g \ll 1$. Within this effective BCS theory, the expression for the critical temperature reads

$$T_c = \omega_0 e^{-1/g}. \quad (1)$$

For the phonon-mediated Cooper instability, one has $\omega_0 \sim \omega_{\text{ph}}$, where ω_{ph} is a typical phonon frequency. (The exact choice of ω_0 is a matter of convention, because it is always possible to change ω_0 by an order-unity factor in combination with small change in g .)

The emergent BCS regime also implies that the constant g can be decomposed into a product of two distinctively different factors—the pseudopotential U and the Fermi liquid factor f_{FL} :

$$g \propto U f_{\text{FL}}. \quad (2)$$

The pseudopotential is understood as an amplitude of the dimensionless (weak) attractive coupling between *bare* electrons near the Fermi surface (FS) and f_{FL} is defined by

$$f_{\text{FL}} = z^2 (m_*/m_0), \quad (3)$$

where z is the quasiparticle residue, m_* is the FS effective mass, and m_0 is the bare electron mass. It accounts for the fact that we are dealing with the correlated Fermi liquid rather than an ideal gas. Exponential sensitivity of the critical temperature to the small parameter g implies that the positive-definite factor f_{FL} —if noticeably smaller than unity—can dramatically suppress the value of T_c .

The strength of Coulomb interaction is characterized by the dimensionless coupling parameter (the Wigner-Seitz radius)

$$r_s = \left(\frac{3}{4\pi a_0^3 n} \right)^{\frac{1}{3}},$$

where n is the number density, and a_0 is the Bohr radius. Typical experimental values of $r_s \gtrsim 2$ correspond to a moderately strong interaction. Therefore *a priori* one might expect that repulsive Coulomb potential will simply eliminate the possibility of phonon-mediated pairing in materials. Meanwhile, the experiment tells us otherwise leading to the following chain of assumptions and arguments aimed at understanding this outcome. *Taking for granted* that effective (low-energy) Coulomb interaction is repulsive and *assuming* that f_{FL} can be ignored, one appeals to the effect of static screening to explain why the suppression of U is often moderate.

This reasoning lead to a widely accepted scenario, originally proposed by Tolmachev [1] and employed by Morel and Anderson [2], of logarithmic suppression of Coulomb repulsion near the FS. The net result is the so-called Morel–Anderson (MA) repulsive Coulomb pseudopotential

$$\mu^* = \frac{\mu}{1 + \mu \ln(E_F/\omega_0)}.$$

It is often used to estimate T_c with the help of McMillan’s formula [3], which was developed as a semi-phenomenological fit reproducing results of large number of experiments [4–8]. The main effect of the MA pseudopotential is to reduce the magnitude of the phonon-mediated U as $U \rightarrow U - \mu^*$, with most experiments suggesting that $\mu^* \in (0.1 \div 0.15)$. It should be noted that $\mu > 0$ was introduced by Morel and Anderson in a rather uncontrolled fashion as some coupling constant

characterizing screened Coulomb repulsion at frequencies $\sim E_F$. If μ is computed from $\rho_0 4\pi e^2 / \kappa^2$ where ρ_0 is the ideal gas FS density of states per spin component, and κ is the Thomas-Fermi screening momentum, then $\mu = 0.5$. The MA construction explains that the values of μ^* are small because the E_F/ω_0 ratio is large, but neither the values of μ nor its sign were derived from first principles, not to mention that Coulomb interaction cannot be fully screened at finite frequencies.

Recent breakthrough in precise computation of Fermi liquid properties of the uniform electron gas [9] points to yet another flaw in the MA scenario: the Fermi liquid factor f_{FL} is significantly smaller than unity at $r_s > 2$; see Fig. 1. In combination with the above-mentioned experimental fact that corrections to g remain small this result forces one to reconsider the effect of Coulomb potential on U at $r_s > 2$ —it has to be far smaller than predicted by μ^* and even opposite in sign, i.e. Coulomb interactions in the s -wave channel might actually *increase* the amplitude of attractive U .




FIG. 1. Fermi liquid factors f_{FL} of the uniform electron gas computed using data reported in Ref. [9].

In this Letter, we employ an implicit renormalization protocol for extracting effective coupling constant and critical temperature from the gap function equation [10] to study the effect of Coulomb repulsion on U and T_c . The four-point vertex function in the Cooper channel is based on approximations that guarantee exact results in the $r_s \rightarrow 0$ limit and their quantitative accuracy up to $r_s \lesssim 2$. We reveal that the suppression of U is maximal at $r_s \approx 0.75$, and the effect diminishes for larger values of r_s . Within the RPA, Coulomb interactions start to *enhance* attractive coupling at $r_s > 2$, but this result is sensitive to inclusion of vertex corrections. We discuss our findings in the context of earlier work suggesting/pointing to a possibility of pairing instability in the absence of electron-phonon coupling, i.e. exclusively




FIG. 2. Effective couplings, Fermi liquid factors, and critical temperatures for phonon-mediated superconductivity with and without the Coulomb vertex function approximated by either RPA or K0 interaction. Both approximation lead to qualitatively similar results for U : as r_s increases, the effective coupling goes through a minimum and starts to increase. The critical temperature follows a similar trend.

on the basis of dynamically screened Coulomb repulsion [1, 11–13].

Model.—The Hamiltonian of the uniform electron gas (UEG) on a neutralizing background is defined as:

$$H = \sum_{\vec{k}\sigma} \epsilon_{\vec{k}} a_{\vec{k}\sigma}^\dagger a_{\vec{k}\sigma} + \frac{1}{2} \sum_{\vec{q} \neq 0} \sum_{\vec{k}\sigma} \sum_{\vec{k}'\sigma'} V_{\vec{q}} a_{\vec{k}+\vec{q}\sigma}^\dagger a_{\vec{k}'-\vec{q}\sigma'}^\dagger a_{\vec{k}'\sigma'} a_{\vec{k}\sigma}, \quad (4)$$

Here $a_{\vec{k}\sigma}^\dagger$ is the creation operator of an electron with momentum \vec{k} and spin $\sigma = \uparrow, \downarrow$, $\epsilon_{\vec{k}} = \frac{\hbar^2 k^2}{2m_0} - \mu$, and $V_{\vec{q}} = \frac{4\pi e^2}{q^2}$ is the bare Coulomb interaction.

The gap function equation in the singlet channel reads:

$$\lambda(T) \Delta_{\omega_n, \mathbf{k}} = -T \sum_m \int \frac{d\mathbf{p}}{(2\pi)^d} \Gamma_{\omega_m, \mathbf{p}}^{\omega_n, \mathbf{k}} G_{\omega_m, \mathbf{p}} G_{-\omega_m, -\mathbf{p}} \Delta_{\omega_m, \mathbf{p}}. \quad (5)$$

Here Γ is the particle-particle irreducible four-point vertex, G is the single particle Green's function, Δ is the

gap function, and λ is its eigenvalue. The critical temperature T_c corresponds to the point where $\lambda_{\max}(T) = 1$.

We consider two approximations for Γ based on the screened Coulomb interaction, both depending only on the momentum and energy transfer, $\Gamma_{\omega_m, \mathbf{p}}^{\omega_n, \mathbf{k}} = \Gamma(\omega_m - \omega_n, \mathbf{p} - \mathbf{k})$. The RPA form is standard:

$$\Gamma_{\text{RPA}}(\omega, q) = \frac{V_q}{1 + V_q \Pi_0(\omega, q)}, \quad (6)$$

where Π_0 is the polarization function computed from the convolution of bare Green's function. For simplicity we take the functional form of Π_0 to be that at $T = 0$, which is justified by the smallness of the critical temperature. To account for vertex corrections and estimate their role as a function of r_s we employ the Kukkonen-Overhauser ansatz [14] when

$$\Gamma_{\text{KO}}(\omega, q) = V_q + V_+(q)^2 Q_+(\omega, q) - 3V_-(q)^2 Q_-(\omega, q), \quad (7)$$

with

$$Q_{\pm}(\omega, q) = -\frac{\Pi_0(\omega, q)}{1 + V_{\pm}(\omega, q)\Pi_0(\omega, q)}, \quad (8)$$

$$V_+ = (1 - G_+)V, \quad V_- = -G_-V. \quad (9)$$

Here Γ_{KO} is already projected on spin-singlet state as required by the fermionic parity. The higher-order vertex corrections neglected in RPA are encoded in the local field factors $G_{\pm}(q)$ for which we adopt the ansatz proposed by Takada [15].

Finally, we introduce phonon-mediated interactions taken to have the same functional form as considered by Ashcroft and Richardson to study the very same problem of superconductivity in the UEG with electron-phonon coupling [13].

$$\Gamma_{\text{ph}}(\omega, q) = -\frac{a\rho_0}{1 + (q/2k_F)^2} \frac{\omega_q^2}{\omega^2 + \omega_q^2}, \quad (10)$$

with the phonon dispersion $\omega_q = \frac{\omega_{\text{ph}}^2 (q/k_F)^2}{1 + (q/k_F)^2}$ and dimensionless coupling strength a . For every choice of the vertex function considered in this work the single particle self-energy was computed self-consistently from the convolution of G and Γ .

Implicit renormalization approach.—For the simplest case when $\Gamma = \Gamma_{\text{ph}}$, the eigenvalue $\lambda(T)$ is a linear function of $\ln T$ at low temperature $T \ll \omega_{\text{ph}}$ that can be written as:

$$\lambda(T) = -g \ln\left(\frac{T}{\omega_0}\right). \quad (11)$$

As expected, the condition $\lambda_{\max}(T_c) = 1$ leads to equation (1), and T_c can be determined accurately by fitting the data even if calculations need to be stopped

at $T \gg T_c$. When Coulomb interactions are included, screening and renormalization effects taking place in a broad frequency range above the phonon frequency ensure that $\lambda_{\max}(T)$ is an unknown non-linear function of $\ln T$ that can be used neither for reliable extrapolation towards lower temperature nor evaluation of the effective low energy coupling U . The implicit renormalization (IR) approach of Ref. [10] provides a solution to both problems by formulating an alternative eigenvalue problem. The gap function is decomposed into two complementary (low-frequency and high-frequency) parts, $\Delta = \Delta^{(1)} + \Delta^{(2)}$, with $\Delta_n^{(1)} = 0$ for $|\omega_n| > \Omega_c$, and $\Delta_n^{(2)} = 0$ for $|\omega_n| < \Omega_c$, and the eigenvalue problem is solved for $\Delta_n^{(1)}$ only. The condition $\bar{\lambda}(T_c) = 1$ for the largest eigenvalue of the new problem remains exact.

As shown in Fig. 3, the IR formulation brings back nearly perfect linear dependence of $\bar{\lambda}$ on $\ln T$ for a properly chosen frequency scale separation Ω_c . The slope of the curve is the emergent low-frequency coupling strength g while the vertical axis intercept determines the characteristic low-frequency scale ω_0 . Linear dependence is also crucial for accurate determination of T_c from simulations performed at $T \gg T_c$ when the critical temperature is extremely low and the number of Matsubara frequency points required to solve the gap equation is large.




FIG. 3. Temperature dependence of the largest eigenvalue $\bar{\lambda}$ for $\Gamma = \Gamma_{\text{ph}} + \Gamma_{\text{RPA}}$ at $r_s = 2$. The emergent BCS linear flow with effective coupling constant g and energy scale ω_0 is represented by the dotted line.

Results.—In Fig. 2 we show the breakdown of the coupling constant into U and f_{FL} and the resulting values of T_c for three different choices of Γ . We first consider the case when the Coulomb repulsion is omitted and Γ is based on the electron-phonon interaction with $a = 0.8$ and $\omega_{\text{ph}} = 0.01E_F$ in (10). According to Migdal's theorem [16], the phonon-mediated vertex correction is pro-




FIG. 4. Gap functions dependence on frequency at $k = k_F$ (left panel) and momentum at $\omega_n = \pi T$ (right panel) when for Γ equals to either Γ_{ph} , or Γ_{RPA} or $\Gamma_{\text{ph}} + \Gamma_{\text{RPA}}$ at $r_s = 3$ (with $\Delta(k_F, \pi T)$ normalized to unity).

portional to ω_{ph}/E_F and can be safely neglected. This calculation serves as a “baseline” for examining effects induced by the Coulomb repulsion. For $\Gamma = \Gamma_{\text{ph}} + \Gamma_{\text{RPA}}$, the pseudopotential U is first reduced to a minimum value at $r_s \approx 0.75$ but then starts to increase and eventually surpasses the electron-phonon interaction value at $r_s \approx 2$ (cf. Ref. [12]). However, the Fermi liquid factor f_{FL} is getting progressively smaller with increasing r_s . The net effect on the critical temperature is also non-monotonic, but $T_c(r_s)$ behavior is not as dramatic because increases in U at $r_s > 1$ are partially compensated by the suppression of f_{FL} .

When vertex corrections are accounted for Coulomb interaction and $\Gamma = \Gamma_{\text{ph}} + \Gamma_{\text{KO}}$, the Fermi liquid factor f_{FL} remains essentially the same for all values of r_s . However, changes in U are relatively small (less than 20%) only for $r_s < 1$. The most significant difference is the shift of the point of the onset of the Coulomb enhancement of U : from $r_s \approx 2$ to $r_s \approx 3$. This result underlines the importance of approximation-free high-order diagrammatic calculations. Nevertheless, the non-monotonic behavior of T_c and U is a robust effect based on the dynamic screening mechanism that tends to make effective Coulomb interactions attractive at large r_s . It is completely overlooked in the MA treatment. If we take our value of T_c at $r_s = 2$ and try to reproduce it with the help of McMillan’s formula, the phenomenological parameter μ^* ends up to be close to 0.08.

There exists yet another fundamental reason for non-additive effects when two pairing mechanisms are combined. If $\lambda_{i=1,2}$ and Δ_i are the largest eigenvalue and its eigenvector for matrix Γ_i and $\Delta_1 \neq \Delta_2$ then the largest eigenvalue of $\Gamma = \Gamma_1 + \Gamma_2$ is *always smaller* than $\lambda_1 + \lambda_2$. In Fig. 4 we show gap function solutions for $\Gamma_1 = \Gamma_{\text{ph}}$, $\Gamma_2 = \Gamma_{\text{RPA}}$ and $\Gamma = \Gamma_1 + \Gamma_2$ at $r_s = 3$. One can see that the eigenvector “mismatch” between these solutions is significant: while Δ_{ph} is sign-positive and monotonic, Δ_{RPA} changes sign both in the momen-

tum and frequency domains and features a pronounced singularity at $k = k_F$.

Discussion and conclusion.—It is instructive to put our findings in the context of historic developments. That Coulomb interaction can induce Cooper instability through dynamic screening mechanism is known for decades. Early work by Tolmachev [1] demonstrated that even if the Cooper channel coupling is repulsive at all frequencies, after its high-frequency part is renormalized to a smaller value the effective low-frequency potential might end up attractive. Later, Takada and others calculated critical temperatures of the UEG numerically using various approximate forms of the screened potential [12, 15, 17] featuring singular frequency/momentum dependence (ignored without justification by Morel and Anderson). These results raise an obvious question: Why are phenomenological values of μ^* used in material science always repulsive if Coulomb interaction alone can be the pairing glue?

Several studies attempted to account for Coulomb effects on superconductivity beyond the MA pseudopotential [13, 18, 19]. Most relevant to our study is the work by Richardson and Ashcroft [13] who calculated T_c for several metals by treating the electron-phonon, Eq. (10), and Coulomb interactions on equal footing. They reported that in Lithium (with $r_s = 3.25$) the inclusion of Coulomb interaction leads to smaller T_c . Our results explain, that for large values of r_s the suppression of f_{FL} is significant and cannot be dismissed as prescribed by the McMillan’s formula. However, this fact was not well established at the time, and Ashcroft and Richardson incorrectly attributed all effect to the repulsive MA pseudopotential μ^* .

By separating Coulomb suppression of the Fermi liquid factor f_{FL} from its contribution to the low-frequency pseudopotential U , we shed new light on the origin of small critical temperatures observed experimentally when compared to predictions of the Migdal-Eliashberg

theory. We reveal that Coulomb contribution to U changes from repulsive to attractive and conclude that the original interpretation of μ^* by Morel and Anderson is incorrect and misleading in two ways: (i) The scenario of enhancement of attractive U due to dynamic nature of screening is ignored, leading to the false impression that μ^* is always positive; (ii) strong renormalization of Fermi liquid properties is ignored, while it can easily reduce effective coupling by a factor of two. These two mistakes partially compensate each other in the phenomenological treatment, yielding reasonable effective coupling constants g within the freedom of choosing μ^* . However, the actual microscopic picture behind the procedure is missed.

Failure to appreciate non-additivity of the phonon and Coulomb contributions to the effective coupling constant g —implied by the structure of Eq. (2)—can lead to qualitatively wrong conclusions. For example, Ref. [12] stated that RPA is a deficient approximation at $r_s > 2$ because it predicts an attractive pseudopotential in contradiction with the “experimentally established” $\mu^* > 0$. Proper account of all the aspects of the interplay between dynamically screened Coulomb repulsion and (alternative) pairing mechanisms may bring new insights in the search for new superconducting materials, especially in cases when McMillan’s equation fails qualitatively.

N. Prokof’ev, B. Svistunov, and T. Wang acknowledge support by the National Science Foundation under Grant No. DMR-2032077. X. Cai acknowledges support by the Simons Collaboration on the Many Electron Problem.

-
- * kunchen@flatironinstitute.org
† svistunov@physics.umass.edu
‡ prokofev@physics.umass.edu
- [1] V. Tolmachev and S. Tiablikov, *Sov. Phys. JETP*. **7** (1958).
 - [2] P. Morel and P. W. Anderson, *Phys. Rev.* **125**, 1263 (1962).
 - [3] W. L. McMillan, *Phys. Rev.* **167**, 331 (1968).
 - [4] M. Calandra and F. Mauri, *Phys. Rev. Lett.* **95**, 237002 (2005).
 - [5] H. T. Hirose, T. Terashima, D. Hirai, Y. Matsubayashi, N. Kikugawa, D. Graf, K. Sugii, S. Sugiura, Z. Hiroi, and S. Uji, *Phys. Rev. B* **105**, 035116 (2022).
 - [6] B. Lian, Z. Wang, and B. A. Bernevig, *Phys. Rev. Lett.* **122**, 257002 (2019).
 - [7] M. Naskar, S. Ash, D. P. Panda, C. K. Vishwakarma, B. K. Mani, A. Sundaresan, and A. K. Ganguli, *Phys. Rev. B* **105**, 014513 (2022).
 - [8] H. Tan, Y. Liu, Z. Wang, and B. Yan, *Phys. Rev. Lett.* **127**, 046401 (2021).
 - [9] K. Haule and K. Chen, *Sci Rep* **12**, 2294 (2022).
 - [10] A. Chubukov, N. V. Prokof’ev, and B. V. Svistunov, *Phys. Rev. B* **100**, 064513 (2019).
 - [11] Y. Takada, *J. Phys. Soc. Jpn.* **45**, 786 (1978).
 - [12] H. Rietschel and L. J. Sham, *Phys. Rev. B* **28**, 5100 (1983).
 - [13] C. F. Richardson and N. W. Ashcroft, *Phys. Rev. B* **55**, 15130 (1997).
 - [14] C. A. Kukkonen and A. W. Overhauser, *Phys. Rev. B* **20**, 550 (1979).
 - [15] Y. Takada, *Phys. Rev. B* **39**, 11575 (1989).
 - [16] A. B. Migdal, *Sov. Phys. JETP*, **7** (1958).
 - [17] Y. Takada, *Phys. Rev. B* **47**, 5202 (1993).
 - [18] M. Lüders, M. A. L. Marques, N. N. Lathiotakis, A. Floris, G. Profeta, L. Fast, A. Continenza, S. Massidda, and E. K. U. Gross, *Phys. Rev. B* **72**, 024545 (2005).
 - [19] T. Wei and Z. Zhang, *Phys. Rev. B* **104**, 184503 (2021).