

# INFINITESIMAL AUTOMORPHISMS OF QUADRICS AND SECOND JET DETERMINATION FOR CR MAPPINGS

ALEXANDER TUMANOV

ABSTRACT. We consider a problem whether a CR mapping of a generic manifold in complex space is uniquely determined by its finite jet at a point, which is referred to as finite jet determination. We derive the finite jet determination for CR mappings of smooth Levi non-degenerate manifolds of arbitrary codimension from the finite dimensionality of the algebra of infinitesimal automorphisms of the corresponding quadrics. Previously, this implication was known for real analytic manifolds. We prove a new 2-jet determination result that covers most affirmative results on this matter obtained so far.

Key words: Infinitesimal automorphism, Second jet determination, CR mapping.

## 1. INTRODUCTION

Let  $M \subset \mathbb{C}^n$  be a generic real submanifold of real codimension  $k$ . Then  $M$  has CR dimension  $\dim_{CR} M = m = n - k$ . We introduce coordinates  $(z, w) \in \mathbb{C}^n$ ,  $z \in \mathbb{C}^m$ ,  $w = u + iv \in \mathbb{C}^k$ , so that  $M$  has a local equation of the form

$$v = h(z, u),$$

here  $h = (h_1, \dots, h_k)$  is a smooth real vector function with  $h(0) = 0$ ,  $dh(0) = 0$ . Furthermore, we can choose the coordinates so that the equation of  $M$  takes the form

$$v = h(z, u) = F(z, z) + O(|z|^3 + |u|^3).$$

Here

$$F = (F_1, \dots, F_k), \quad F_j(z, z) = \langle A_j z, z \rangle, \quad \langle a, b \rangle = \sum a_l \bar{b}_l,$$

$A_j$ -s are Hermitian matrices. The vector valued Hermitian form  $F$  can be regarded as the Levi form of  $M$  at 0.

- We say  $M$  is (Levi) *nondegenerate* at 0 if
  - (a) the matrices  $A_j$  are linearly independent and
  - (b)  $F(z, \zeta) = 0$  for all  $z \in \mathbb{C}^m$  implies  $\zeta = 0$ .
- We say  $M$  is *strongly nondegenerate* at 0 if  $M$  is nondegenerate and there is  $c \in \mathbb{R}^k$  such that  $\det(\sum c_j A_j) \neq 0$ .
- We say  $M$  is *strongly pseudoconvex* at 0 if  $M$  is nondegenerate and there is  $c \in \mathbb{R}^k$  such that  $\sum c_j A_j > 0$ .

---

2020 *Mathematics Subject Classification.* 32H12, 32V40.

Partially supported by Simons Foundation grant.

We will also call the form  $F$  itself respectively nondegenerate, strongly nondegenerate, or strongly pseudoconvex in these cases.

We recall that a *CR mapping* or a *CR diffeomorphism* between two generic submanifolds is a diffeomorphism that induces a complex isomorphism between their complex tangent bundles.

We are concerned with the problem whether a CR diffeomorphism between two manifolds is uniquely determined by its finite jet at a point, which is referred to as *finite jet determination*. The problem has been popular since 1970-s, and the number of publications on the matter has been very large, see surveys [1, 8, 17]. Nevertheless, there are fundamental open questions even in the Levi nondegenerate case, to which we restrict here.

Tanaka [14] gave a solution to the CR equivalence problem for nondegenerate CR manifolds of codimensions  $k = 1, m^2 - 1, m^2$ . His result implies 2-jet determination for real analytic CR mappings of real analytic nondegenerate manifolds of said codimensions. Tanaka's result for a real hypersurface ( $k = 1$ ) was rediscovered by Chern and Moser [10].

Beloshapka [2] proved finite jet determination for real analytic CR mappings of real analytic nondegenerate manifolds.

Bertrand and Meylan [5] prove 2-jet determination for  $C^3$ -smooth CR mappings of  $C^4$ -smooth generic nondegenerate manifold  $M$  with additional condition that the authors call *D-nondegeneracy*. In particular, it implies that there is  $z \in \mathbb{C}^m$  such that the vectors  $\{A_j z : 1 \leq j \leq k\}$  are  $\mathbb{R}$ -linearly independent. This condition is quite restrictive, in particular, it implies that  $k \leq 2m$ , whereas the dimension of the space of all Hermitian forms on  $\mathbb{C}^m$  is equal to  $m^2$ , and the nondegeneracy condition, part (a), imposes only the restriction  $k \leq m^2$ .

In [16], we prove 2-jet determination for smooth CR automorphisms of smooth *strongly pseudoconvex* manifolds.

Beloshapka [3] proves 2-jet determination for nondegenerate codimension  $\leq 3$  manifolds in the real analytic case.

On the other hand, Meylan [12] has constructed a surprising counterexample of a quadric for which 2-jet determination fails. For arbitrary large integer  $p$ , Gregerovič and Meylan [9] have constructed counterexamples for which  $p$ -jet determination fails.

In this paper, we prove 2-jet determination for smooth CR mappings under a new more general condition that we call *T-nondegeneracy*, see Definition 5.1. Our result implies all affirmative results on 2-jet determination mentioned above, in particular, for strictly pseudoconvex, D-nondegenerate and codimension  $\leq 3$  CR manifolds.

We note that Bertrand, Blanc-Centi, and Meylan [4, 5, 6] and the author [16] obtained their results by using stationary discs, a family of analytic discs invariant under CR mappings. This method works so far only for *strongly* nondegenerate manifolds because the existence of stationary discs is proven under this assumption.

In this paper, we follow the classical approach based on the infinitesimal automorphisms of quadrics. The Lie algebra  $\mathfrak{g}$  of infinitesimal automorphisms of the quadric  $v = F(z, z)$  has

a natural grading  $\mathfrak{g} = \mathfrak{g}_{-2} + \mathfrak{g}_{-1} + \mathfrak{g}_0 + \dots$ . It is long known [2, 15] that for a nondegenerate quadric,  $\mathfrak{g}$  has finite dimension, which implies (Beloshapka [2]) finite jet determination in the real analytic case. We prove (Theorem 4.1) this fact in the smooth case. In particular, we prove (Theorem 5.2) that for T-nondegenerate CR manifolds we have  $\mathfrak{g}_3 = 0$ , which implies (Theorem 4.2) 2-jet determination in the smooth case.

The paper is structured as follows. In Section 2, we introduce infinitesimal automorphisms of quadrics and state basic facts about them. In Section 3, we develop a rudimentary theory of weighted jets of mappings. In Section 4, we derive finite jet determination results for smooth CR mappings from the finite dimensionality of  $\mathfrak{g}$ . In Section 5, we prove the vanishing of  $\mathfrak{g}_3$  for T-nondegenerate quadrics. In Sections 6 and 7, we show that the previous affirmative results on 2-jet determination follow from our Theorems 4.2 and 5.2. Finally, in Section 8, we give several examples illustrating various nondegeneracy conditions and their effect on 2-jet determination.

The author wishes to thank Francine Meylan for letting him know about her counterexample [12] and Alexander Sukhov for useful discussions.

## 2. INFINITESIMAL AUTOMORPHISMS OF QUADRICS

An infinitesimal CR-automorphism of a CR-manifold  $M$  is a vector field on  $M$  that generates a local 1-parameter group of CR-mappings (CR-automorphisms)  $M \rightarrow M$ .

Let  $M_0$  be a nondegenerate quadric defined as before by the equations

$$v = F(z, z), \quad z \in \mathbb{C}^m, \quad w = u + iv \in \mathbb{C}^k.$$

Here

$$F = (F_1, \dots, F_k), \quad F_j(z, z) = \langle A_j z, z \rangle, \quad \langle a, b \rangle = \sum a_l \bar{b}_l.$$

Let  $G$  be the group of all CR-automorphisms  $M_0 \rightarrow M_0$ . Then  $G$  is a finite dimensional Lie group and its Lie algebra  $\mathfrak{g}$  is the set of all infinitesimal automorphisms of  $M_0$ . The dimension of  $G$  has an estimate depending on  $m$  and  $k$  (see Beloshapka [2], Isaev and Kaup [11], and the author [15]).

It turns out that all elements of  $G$  and  $\mathfrak{g}$  are respectively rational and polynomial. In particular, every vector field  $X \in \mathfrak{g}$  has the form

$$X = \sum f_j \frac{\partial}{\partial z_j} + \sum g_\ell \frac{\partial}{\partial w_\ell} = f \frac{\partial}{\partial z} + g \frac{\partial}{\partial w} =: (f, g),$$

where  $f$  and  $g$  are polynomial vector functions of  $z$  and  $w$  that satisfy the equation (see [2, 8])

$$(1) \quad \operatorname{Im}(g - 2iF(f, z)) = 0, \quad (z, w) \in M.$$

This equation implies (see [2, 8])

$$(2) \quad \deg_z f \leq 2, \quad \deg_z g \leq 1.$$

We will use the equation (1) rather than the infinitesimal automorphisms themselves.

We give the variables and differentiations  $z_j, w_j, \partial/\partial z_j, \partial/\partial w_j$  the weights  $1, 2, -1, -2$  respectively. Let  $\mathfrak{g}_p$  be the set of vector fields  $X \in \mathfrak{g}$  with weighted homogeneous degree  $p \in \mathbb{Z}$ . Then the elements  $X = (f_{p+1}, g_{p+2}) \in \mathfrak{g}_p$  are solutions of the equation

$$(3) \quad \text{Im} (g_{p+2} - 2iF(f_{p+1}, z))|_{w=u+iF(z,z)} = 0,$$

which is a linear homogeneous equation on the coefficients on the weighted homogeneous polynomials  $f_{p+1}$  and  $g_{p+2}$  of weighted degrees respectively  $p+1$  and  $p+2$ . Then

$$\mathfrak{g} = \sum_{p=-2}^{\infty} \mathfrak{g}_p$$

is a graded Lie algebra, that is,  $[\mathfrak{g}_p, \mathfrak{g}_q] \subset \mathfrak{g}_{p+q}$ . Since  $\mathfrak{g}$  has finite dimension,  $\mathfrak{g}_p = 0$  for big  $p$ . Since  $F$  is nondegenerate, it follows that each vector  $\xi \in \mathfrak{g}_p$  is uniquely determined by the map  $\text{ad } \xi : \mathfrak{g}_{-1} \rightarrow \mathfrak{g}_{p-1}$ , here  $(\text{ad } \xi)(\eta) = [\xi, \eta]$ . In particular, if  $\mathfrak{g}_p = 0$ , then  $\mathfrak{g}_q = 0$  for all  $q > p$ .

### 3. WEIGHTED JETS OF MAPPINGS

A CR manifold  $M$  has a complex tangent subbundle  $T^c(M) \subset T(M)$ . In this section, we consider a more general structure. Let  $M$  be a smooth real manifold of dimension  $n$  with a real vector subbundle  $H(M) \subset T(M)$  with fiber dimension  $m$ . (We will specify the smoothness class later.) For brevity we will call such manifolds *H-manifolds*. We will consider *morphisms* of H-manifolds, that is, mappings  $M \rightarrow M'$  whose tangent mappings take  $H(M)$  to  $H(M')$ . We introduce a rudimentary theory of weighted jet bundles of morphisms of H-manifolds.

Let  $M$  be a H-manifold. Let  $a \in M$ . Let  $(x, y) \in \mathbb{R}^m \times \mathbb{R}^k$ ,  $k = n - m$ , be local coordinates such that  $a$  has coordinates  $(0, 0)$ , and  $H_a(M)$  is represented by the equation  $y = 0$ . We will call such a coordinate system *admissible* with origin at  $a \in M$ . One should think of  $x$  and  $y$  variables as having weights respectively 1 and 2. We will use weighted distance to the origin  $\sigma = |x| + |y|^{1/2}$ .

Let  $M$  and  $M'$  be H-manifolds, and let  $a \in M$ ,  $a' \in M'$ . Let  $(x, y)$  and  $(x', y')$  be corresponding admissible coordinates. Let  $\Phi : M \rightarrow M'$  be a morphism,  $\Phi(a) = a'$ . Then  $\Phi$  can be represented by equations

$$x' = \phi(x, y), \quad y' = \psi(x, y).$$

Then we will write  $\Phi \cong (\phi, \psi)$ . Since  $\Phi(a) = a'$ , we have  $\phi(0, 0) = 0$ ,  $\psi(0, 0) = 0$ .

Let  $\Phi \cong (\phi, \psi)$  and  $\tilde{\Phi} \cong (\tilde{\phi}, \tilde{\psi})$  be morphisms sending  $a$  to  $a'$ . Let  $p, q$  be nonnegative integers with  $p \leq q \leq 2p$ . We say that  $\Phi$  and  $\tilde{\Phi}$  are  $(p, q)$ -equivalent and write  $\Phi \sim_{(p,q)} \tilde{\Phi}$  if

$$\phi - \tilde{\phi} = o(\sigma^p), \quad \psi - \tilde{\psi} = o(\sigma^q).$$

We claim that the equivalence relation  $\sim_{(p,q)}$  is well defined, that is, independent of coordinates  $(x, y)$  and  $(x', y')$ . Without loss of generality we restrict to the case  $\tilde{\Phi} \equiv a'$ , that is,  $\tilde{\phi} \equiv 0$ ,  $\tilde{\psi} \equiv 0$ .

We first show the independence of  $(x, y)$ . Indeed, let  $(\hat{x}, \hat{y})$  be another admissible system. Then  $\partial\hat{y}/\partial x = 0$ ,  $\hat{x} = O(|x| + |y|)$ ,  $\hat{y} = O(|x|^2 + |y|)$ . Then  $\hat{\sigma} = |\hat{x}| + |\hat{y}|^{1/2} = O(|x| + |y|) + O(|x| + |y|^{1/2}) = O(\sigma)$ . Similarly,  $\sigma = O(\hat{\sigma})$ . Hence, the relation is independent of  $(x, y)$ .

We now consider the effect of another coordinate system  $(\hat{x}', \hat{y}')$  in the target. Let  $\phi = o(\sigma^p)$ ,  $\psi = o(\sigma^q)$ . Then in the new coordinates we will have  $\hat{\phi} = O(|\phi| + |\psi|) = o(\sigma^p + \sigma^q) = o(\sigma^p)$  because  $p \leq q$ . We will also have  $\hat{\psi} = O(|\phi|^2 + |\psi|) = o(\sigma^{2p} + \sigma^q) = o(\sigma^q)$  because  $q \leq 2p$ . Hence, the relation is independent of  $(x', y')$ .

We denote the set of equivalence classes of morphisms sending  $a$  to  $a'$  by  $J_{a,a'}^{p,q}(M, M')$  and the set of all classes for all pairs  $(a, a')$  by  $J^{p,q}(M, M')$ . Then one can see that  $J^{p,q}(M, M') \rightarrow M \times M'$  is a fiber bundle whose fiber is a Euclidean space. We call the members of  $J^{p,q}(M, M')$  weighted jets of morphisms  $M \rightarrow M'$ .

Let  $\Phi : M \rightarrow M'$  be a morphism with  $\Phi(a) = a'$ . We denote by  $j_a^{p,q}(\Phi)$  the equivalence class of  $\Phi$  in  $J_{a,a'}^{p,q}(M, M')$  and call it the  $(p, q)$ -jet of  $\Phi$  at  $a$ . Given coordinates  $(x, y)$  and  $(x', y')$ , for  $\Phi \cong (\phi, \psi)$ , this jet is represented by the set

$$\{a, a', D^\alpha \phi(0), D^\beta \psi(0) : 0 < \|\alpha\| \leq p, 0 < \|\beta\| \leq q\},$$

here  $\alpha$  and  $\beta$  are multi-indices corresponding to the variables  $(x, y)$ , and for a multi-index  $\alpha = (\alpha_x, \alpha_y)$ , the weighted length  $\|\alpha\| = |\alpha_x| + 2|\alpha_y|$ .

For a morphism  $\Phi : M \rightarrow M'$ , we also introduce the map  $j^{p,q}(\Phi) : M \rightarrow J^{p,q}(M, M')$ ,  $a \mapsto j_a^{p,q}(\Phi)$ , and call it the  $(p, q)$ -jet of  $\Phi$ .

We now restrict to the case  $q = p + 1$  and in all notations introduced so far, we replace the pair  $(p, q)$  by the single number  $p$ . In our applications to CR manifolds, we will understand weighted jets of mappings this way. The main point of this section is the following.

**Proposition 3.1.** *Let  $p \geq 1$  be an integer. Let  $M$  and  $M'$  be  $C^{p+2}$ -smooth  $H$ -manifolds. Let  $\Phi : M \rightarrow M'$  be a morphism. Let  $a \in M$ ,  $a' = \Phi(a)$ , and let  $X \in H_a(M)$ . Then  $Xj^p(\Phi)$  (the derivative of the mapping along the tangent vector) can be expressed in terms of  $j_a^{p+1}(\Phi)$ .*

The result is quite intuitive because the derivative with respect to the  $x$ -variable, whose weight is 1, is supposed to add 1 to the weighted order of the jet. If the bundles  $H(M)$  and  $H(M')$  were integrable (involutive) then we could represent all jets in the same coordinate systems. Then the conclusion would be obvious. However, in our application to CR manifolds, they are not integrable, so the result needs some attention.

*Proof.* We fix admissible coordinates  $(x, y)$  and  $(x', y')$  in  $M$  and  $M'$ , and we call them main coordinates identifying  $M$  and  $M'$  with open sets in the Euclidean space. In order to represent  $j^p(\Phi)$ , we introduce admissible coordinates  $(\xi, \eta)$  and  $(\xi', \eta')$  with origins at  $(x, y)$  and  $(x', y')$  respectively, and we call them variable coordinates. We define  $(\xi, \eta)$  by the mapping

$$(\xi, \eta) \mapsto (\tilde{x}, \tilde{y}) = (x + \xi, y + \eta + A(x, y)\xi),$$

here  $A(x, y)$  is a matrix, so  $H_{(x,y)}(M)$  in the main coordinates  $(x, y)$  is defined as  $\{(\xi, A(x, y)\xi) : \xi \in \mathbb{R}^m\}$ . Then  $A(0, 0) = 0$ . We define  $(\xi', \eta')$  similarly.

Let  $\Phi \cong (\phi, \psi)$  in the main coordinates. We need the representation of  $\Phi$  in the variable coordinates, denote it  $\Phi \cong (f, g)$ . Then we have

$$\begin{aligned} f &= \phi(x + \xi, y + \eta + A(x, y)\xi) - \phi(x, y), \\ g &= \psi(x + \xi, y + \eta + A(x, y)\xi) - \psi(x, y) - A'(\phi(x, y), \psi(x, y))f. \end{aligned}$$

The desired claim means exactly that the derivatives

$$\begin{aligned} \partial_x|_{(x,y)=(0,0)} D_{\xi,\eta}^\alpha|_{(\xi,\eta)=(0,0)} f, \quad \|\alpha\| \leq p, \\ \partial_x|_{(x,y)=(0,0)} D_{\xi,\eta}^\beta|_{(\xi,\eta)=(0,0)} g, \quad \|\beta\| \leq p + 1, \end{aligned}$$

can be expressed in terms of  $D^\alpha \phi(0, 0)$  and  $D^\beta \psi(0, 0)$  with  $\|\alpha\| \leq p + 1$ ,  $\|\beta\| \leq p + 2$ . The verification is an exercise on the Chain Rule. Indeed, an increase in the weighted order of derivative by more than one unit can happen when we differentiate  $\phi$  or  $\psi$  with respect to the variable  $\xi$  that occurs in the second argument of the function, because it adds 2 to the weighed order. Then the factor  $A(x, y)$  comes out. Then in the subsequent differentiation with respect to  $x$ , we only have to differentiate  $A(x, y)$  because  $A(0, 0) = 0$ . Thus the final differentiation with respect to  $x$  does not increase the weighted order of the derivatives of  $\phi$  or  $\psi$ , so the resulting weighted order of the derivative increases by no more than one unit.

We point out that the choice  $q = p + 1$  is dictated by the above requirement on the derivatives of  $g$ . Since the equation of  $g$  involves  $f$ , the derivatives of  $g$  would include the derivatives of  $\phi$ , whose order has to be bounded by  $p + 1$ . We leave the details to the reader.  $\square$

#### 4. FINITE JET DETERMINATION

Let  $M$  be a nondegenerate CR manifold with equation

$$v = h(z, u) = F(z, z) + O(|z|^3 + |u|^3),$$

and let  $M_0$  be the corresponding quadric with equation

$$v = F(z, z).$$

Let  $\mathfrak{g}$  be the graded Lie algebra of infinitesimal automorphisms of  $M_0$ . It turns out that the finite dimensionality of  $\mathfrak{g}$  implies finite jet determination for smooth CR mappings of  $M$ .

A CR manifold  $M$  is an H-manifold as defined in Section 3 with  $H(M) = T^c(M)$ . Then we can use weighted jets of CR mappings as we defined there.

**Theorem 4.1.** *Let  $p \geq 1$  be an integer. Let  $M, M'$  be  $C^{p+2}$ -smooth nondegenerate CR manifolds defined as above. Suppose  $\mathfrak{g}_p = 0$ . Then every germ at 0 of a  $C^{p+2}$ -smooth CR diffeomorphism  $\Phi : M \rightarrow M'$  is uniquely determined by the weighted jet  $j_0^p(\Phi)$ .*

Beloshapka [2] obtained the real analytic version of this result. As a consequence we obtain the following.

**Theorem 4.2.** *Let  $M, M'$  be  $C^5$ -smooth non-degenerate CR manifolds defined as above. Suppose  $\mathfrak{g}_3 = 0$ . Then every germ at 0 of a  $C^5$ -smooth CR diffeomorphism  $\Phi : M \rightarrow M'$  is uniquely determined by the (standard) 2-jet of  $\Phi$  at 0.*

*Proof of Theorem 4.1.* We first show that  $j_0^{p+1}(\Phi)$  is uniquely determined by  $j_0^p(\Phi)$ . The argument is well known (see [2, 8]), and we describe it briefly.

We recall that a  $C^r$ -smooth CR function  $f$  on a  $C^r$ -smooth generic manifold in  $\mathbb{C}^n$  defined as above can be developed into a Taylor series with remainder of the form  $f(z, w) = P(z, w) + o(|z|^r + |w|^r)$ , where  $P(z, w)$  is a holomorphic polynomial of degree  $\leq r$ . We can represent  $P$  as a sum of weighted homogeneous holomorphic polynomials, in which variables  $z$  and  $w$  have weights respectively 1 and 2, and the index means the weighted degree of the polynomial. Then we obtain

$$f = f_0 + f_1 + \dots + f_r + o(|z|^r + |w|^{r/2}).$$

Following Chern and Moser [10] and Beloshapka [2], we expand the equations of  $M$  and  $M'$  and the CR mapping  $\Phi = (f, g)$  into Taylor series with remainders and represent them as sums of weighed homogeneous components, where  $z$ ,  $w$ , and  $u$  have weights respectively 1, 2, and 2. Without loss of generality,  $\Phi(0) = 0$ . Then we have

$$\begin{aligned} v &= h(z, u) = F + h_3 + \dots + h_{p+2} + o(|z|^{p+2} + |u|^{(p+2)/2}) \\ v' &= h'(z', u') = F' + h'_3 + \dots + h'_{p+2} + o(|z'|^{p+2} + |u'|^{(p+2)/2}) \\ z' &= f(z, w) = f_1 + f_2 + \dots + f_{p+2} + o(|z|^{p+2} + |w|^{(p+2)/2}) \\ w' &= g(z, w) = g_2 + g_3 + \dots + g_{p+2} + o(|z|^{p+2} + |w|^{(p+2)/2}). \end{aligned}$$

Since the derivative of  $\Phi$  maps the complex tangent plane  $w = 0$  to the plane  $w' = 0$ , we have  $g_1 = 0$ . By linear transformations of  $z$  and  $w$ , without loss of generality, we can put  $f_1 = z$ ,  $g_2 = w + Q(z)$ , where  $Q$  is a quadratic polynomial, but one can see that  $Q = 0$ . Also, one can see that  $F' = F$ .

By plugging  $z'$  and  $w'$  in terms of  $z$  and  $w = u + ih(z, u)$  in the equation of  $M'$  and collecting the terms of weight  $p + 2$ , we obtain:

$$(4) \quad \text{Im}(g_{p+2} - 2iF(f_{p+1}, z))|_{w=u+iF(z,z)} = \dots + o(|z|^{p+2} + |u|^{(p+2)/2}),$$

here the dots mean terms that include only  $f_{q+1}$  and  $g_{q+2}$  with  $q < p$ .

Note that the corresponding homogeneous equation (3) has only the trivial solution because  $\mathfrak{g}_p = 0$ . Then the component  $(f_{p+1}, g_{p+2})$  is uniquely determined by the components of  $\Phi$  of lower weighted degree. That is,  $j_0^{p+1}(\Phi)$  is uniquely determined by  $j_0^p(\Phi)$ .

We note that since  $\mathfrak{g}_p$  is obtained by solving a linear homogeneous system (3), the hypothesis that  $\mathfrak{g}_p = 0$  holds at zero implies that  $\mathfrak{g}_p = 0$  holds at nearby points even though the Levi form of  $M$  at those points is not necessarily equivalent to  $F$  as a vector valued Hermitian form, but it will be close to  $F$ . Without loss of generality,  $\mathfrak{g}_p = 0$  for every point of  $M$ .

Let  $\gamma : (-1, 1) \rightarrow M$  be a CR curve in  $M$ , that is,  $d\gamma(t)/dt \in T_{\gamma(t)}^c(M)$ , and let  $\gamma(0) = 0$ . By the above arguments,  $j_{\gamma(t)}^{p+1}(\Phi)$  is a function of  $j_{\gamma(t)}^p(\Phi)$ . By Proposition 3.1, this gives rise to a first order ordinary differential equation on the jet  $\Psi(t) = j_{\gamma(t)}^p(\Phi)$ . This equation has a unique solution with given initial condition  $\Psi(0) = j_0^p(\Phi)$ . In particular,  $\Phi(\gamma(t))$  is uniquely determined for all  $t$ .

Since  $M$  is nondegenerate, CR curves through 0 cover an open set in  $M$ . Then  $\Phi$  is uniquely determined by  $j_0^p(\Phi)$ , as desired.  $\square$

*Proof of Theorem 4.2.* By Theorem 4.1, the mapping  $\Phi$  is uniquely determined by the weighted jet  $j_0^3(\Phi)$ . We note that  $j_0^3(\Phi)$  is uniquely determined by the standard 2-jet of  $\Phi$  at 0. This follows by (4) with  $p = 2$  and the bounds (2) on the  $z$ -degrees of  $f_3$  and  $g_4$ . (See [7, 8] for details.)  $\square$

**Remark 4.3.** The regularity requirements in Theorems 4.1 and 4.2 are set so that all occurring jets make sense. We note that by Stein [13], a  $C^r$ -smooth CR function  $f$  on a nondegenerate generic CR manifold has roughly  $2r$  derivatives along complex tangential directions. Then a weighted jet  $j^p f$  makes sense for some  $p > r$ . Thus the required smoothness of class  $C^{p+2}$  in Theorem 4.1 in fact can be lowered. We leave the details to the reader.

## 5. VANISHING OF $\mathfrak{g}_3$

Let  $F$  be as above. Let  $S \subset \mathbb{C}^m$  be a set. We define the orthogonal complement

$$S^F = \{x \in \mathbb{C}^m : \forall y \in S, F(x, y) = 0\}.$$

Then  $S^F$  is a complex subspace of  $\mathbb{C}^m$ . Similarly, we can define  $S^{\text{Re}F}$ . We note that  $S^F \subset S^{\text{Re}F}$ , and if  $S$  is a complex subspace, then  $S^F = S^{\text{Re}F}$ . For  $a, b \in \mathbb{C}^m$ , we define complex subspaces

$$(5) \quad T(a, b) = \{y \in \mathbb{C}^m : \exists x \in \mathbb{C}^m : F(x, b) + F(a, y) = 0\}, \quad T(a) = T(a, a).$$

We observe

$$(6) \quad T(a, b) \supset a^F + \mathbb{C}b, \quad T(a, b)^F \subset (a^F)^F \cap b^F, \quad T(a)^F \subset (a^F)^F \cap a^F$$

If a certain property holds for every element  $x$  in an open dense set of a space  $X$ , then we will say that the property holds for a *generic* element of  $X$ .

**Definition 5.1.** We say that a nondegenerate form  $F$  is  $T$ -nondegenerate if for generic  $a \in \mathbb{C}^m$ , we have  $T(a)^F = 0$ .

**Theorem 5.2.** Let  $F$  be a  $T$ -nondegenerate form. Then  $\mathfrak{g}_3 = 0$ .

In the proof, we will apply the following lemma three times.

**Lemma 5.3.** Let  $\phi : \mathbb{C}^k \rightarrow \mathbb{C}^m$  be a  $C^1$  mapping. Suppose for every  $a, b \in \mathbb{C}^m$ , we have  $F(\phi(F(a, b)), b) = 0$ . Then for every  $a, b \in \mathbb{C}^m$ , we have  $\phi(F(a, b)) \in T(a, b)^F$ . In particular, for every  $a \in \mathbb{C}^m$ , we have  $\phi(F(a, a)) \in T(a)^F$ . Hence, if  $F$  is  $T$ -nondegenerate, then for every  $a \in \mathbb{C}^m$ , we have  $\phi(F(a, a)) = 0$ .



*Proof.* Let  $a, b, x, y \in \mathbb{C}^m$ , let  $F(x, b) + F(a, y) = 0$ , so  $y \in T(a, b)$ . For  $t \in \mathbb{R}$ , we put

$$a' = a + tx, \quad b' = b + ty.$$

Then we have

$$\begin{aligned} F(a', b') &= F(a, b) + t(F(x, b) + F(a, y)) + t^2 F(x, y) = F(a, b) + O(t^2), \\ \phi(F(a', b')) &= \phi(F(a, b)) + O(t^2). \end{aligned}$$

By the hypothesis we have

$$\begin{aligned} 0 &= F(\phi(F(a', b')), b') \\ &= F(\phi(F(a, b)) + O(t^2), b + ty) \\ &= F(\phi(F(a, b), b) + tF(\phi(F(a, b)), y) + O(t^2)). \end{aligned}$$

Then  $F(\phi(F(a, b)), y) = 0$ . Since it holds for every  $y \in T(a, b)$ , we have  $\phi(F(a, b)) \in T(a, b)^F$ , as desired.  $\square$

*Proof of Theorem 5.2.* An element  $(f, g) \in \mathfrak{g}_3$  has the following form (see [2, 8])

$$\begin{aligned} f(z, w) &= A(z, z, w) + B(w, w), \\ g(z, w) &= 2iF(z, B(\bar{w}, \bar{w})). \end{aligned}$$

Here  $A$  and  $B$  are complex multilinear forms such that  $A$  is symmetric in the first two arguments and  $B$  is symmetric. They are characterized by the following equations:

$$\begin{aligned} (7) \quad & F(A(z, z, F(z, b)), b) = 0, \\ (8) \quad & F(A(z, z, w), b) = 4iF(z, B(\bar{w}, F(b, z))). \end{aligned}$$

Here  $b, z \in \mathbb{C}^m$ , and  $w \in \mathbb{C}^k$  are arbitrary. In fact,  $A(z, z, F(z, b)) = 0$ , but we do not need this fact here. We would like to show that if  $F$  is T-nondegenerate, then  $A = 0$  and  $B = 0$ .

We first plug  $w = F(z, b)$  in (8) and using (7) we obtain

$$(9) \quad F(z, B(F(b, z), F(b, z))) = 0.$$

Put  $\phi(w) = B(w, w)$ . Then by (9) we have  $F(\phi(F(b, z)), z) = 0$  for all  $b, z \in \mathbb{C}^m$ . Since  $F$  is T-nondegenerate, by Lemma 5.3, we have  $\phi(F(b, b)) = 0$ , that is,

$$B(F(b, b), F(b, b)) = 0.$$

By polarization

$$B(F(b, z), F(b, z)) = 0.$$

Since  $B$  is symmetric, we have

$$(10) \quad B(F(b, a), F(b, z)) = 0$$

for all  $a, b, z \in \mathbb{C}^m$ . Plug  $w = F(a, b)$  in (8). Then by (10) we have

$$(11) \quad F(A(z, z, F(a, b)), b) = 4iF(z, B(F(b, a), F(b, z))) = 0.$$

Put  $\phi(w) = A(z, z, w)$ , where  $z \in \mathbb{C}^m$  is constant. Then by (11) we have  $F(\phi(F(a, b)), b) = 0$ . By Lemma 5.3, we have  $\phi(F(a, a)) = 0$ , that is,

$$A(z, z, F(a, a)) = 0.$$

Since  $F$  is nondegenerate, the values  $F(a, a)$  span all of  $\mathbb{C}^k$ , hence  $A = 0$ . Then by (8) we have

$$(12) \quad F(z, B(u, F(b, z))) = 0$$

for all  $b, z \in \mathbb{C}^m$  and  $u \in \mathbb{C}^k$ . Put  $\phi(w) = B(u, w)$ , where  $u \in \mathbb{C}^k$  is constant. Then by (12), we have  $F(\phi(F(b, z)), z) = 0$ . By Lemma 5.3, we have  $\phi(F(b, b)) = 0$ , that is,

$$B(u, F(b, b)) = 0.$$

Hence  $B = 0$ . Now  $A = 0$ ,  $B = 0$ ,  $f = 0$ ,  $g = 0$ , and  $\mathfrak{g}_3 = 0$ , as desired.  $\square$

## 6. OTHER CONDITIONS

Let  $F$  and  $A_j$  be as above. Let  $F$  be nondegenerate. Recall that  $F$  is strongly pseudoconvex if there is  $c \in \mathbb{R}^k$  such that  $A = \sum_{j=1}^k c_j A_j > 0$ , positive definite. In [16] we prove 2-jet determination for strongly pseudoconvex CR manifolds. We recover this result using the following.

**Proposition 6.1.** *Let the form  $F$  be strongly pseudoconvex. Then  $F$  is  $T$ -nondegenerate, hence  $\mathfrak{g}_3 = 0$ .*

*Proof.* If  $F$  is strongly pseudoconvex, then for  $z \neq 0$  we have  $F(z, z) \neq 0$ . Then we have  $(a^F)^F \cap a^F = 0$ , because if  $z \in (a^F)^F \cap a^F$ , then  $F(z, z) = 0$ . Now the result reduces to the following result.  $\square$

**Proposition 6.2.** *Suppose for generic  $a \in \mathbb{C}^m$ , we have  $(a^F)^F \cap a^F = 0$ . Then  $F$  is  $T$ -nondegenerate.*

*Proof.* By (6), we have  $T(a)^F \subset (a^F)^F \cap a^F = 0$ . Hence  $F$  is  $T$ -nondegenerate, as desired.  $\square$

Bertrand, Blanc-Centi, and Meylan [4], prove 2-jet determination for so called *fully nondegenerate* CR manifolds in the smooth case. Their condition, in particular, implies that there is  $a \in \mathbb{C}^m$  such that the vectors  $(A_j a)_{j=1}^k$  are  $\mathbb{C}$ -linear independent. Bertrand and Meylan [6] show that this weaker condition alone implies 2-jet determination in the analytic case. We recover these results here by proving the following.

**Proposition 6.3.** *Suppose there is  $a \in \mathbb{C}^m$  such that the vectors  $(A_j a)_{j=1}^k$  are  $\mathbb{C}$ -linear independent. Then  $F$  is  $T$ -nondegenerate, hence  $\mathfrak{g}_3 = 0$ .*

*Proof.* By the hypotheses, for generic  $a \in \mathbb{C}^m$ , we have  $F(\mathbb{C}^m, a) = \mathbb{C}^k$ . Then  $T(a) = \mathbb{C}^m$ , and  $T(a)^F = 0$ , hence  $F$  is  $T$ -nondegenerate.  $\square$

Beloshapka [3] proves a sufficient condition for  $\mathfrak{g}_3 = 0$  that in particular includes the hypothesis that there is  $a \in \mathbb{C}$  such that the vectors  $(A_j a)_{j=1}^k$  span  $\mathbb{C}^m$  over  $\mathbb{C}$ . We observe that this hypothesis alone suffices for  $\mathfrak{g}_3 = 0$ .

**Proposition 6.4.** *Suppose that there is  $a \in \mathbb{C}^m$  such that the vectors  $(A_j a)_{j=1}^k$  span  $\mathbb{C}^m$  over  $\mathbb{C}$ . Then  $F$  is  $T$ -nondegenerate, hence  $\mathfrak{g}_3 = 0$ .*

*Proof.* The hypothesis is equivalent to the fact that for generic  $a \in \mathbb{C}^m$ , we have  $a^F = 0$ . Then this proposition is a special case of Proposition 6.2.  $\square$

Suppose there is  $c \in \mathbb{R}^k$  such that  $A = \sum_{j=1}^k c_j A_j$  is nonsingular, that is,  $F$  is strongly nondegenerate. Let  $a \in \mathbb{C}^m$ . Following Bertrand and Meylan [5], we introduce a  $m \times k$  matrix

$$D = (A_1 a, \dots, A_k a)$$

and a  $k \times k$  matrix

$$(13) \quad B = D^* A^{-1} D, \quad B_{lj} = \langle A_l A^{-1} A_j a, a \rangle.$$

**Definition 6.5** ([5]).  *$F$  is called D-nondegenerate if there exist  $c \in \mathbb{R}^k$  and  $a \in \mathbb{C}^m$  such that the matrix  $\operatorname{Re} B := \frac{1}{2}(B + \overline{B})$  is nonsingular.*

Bertrand and Meylan [5] prove 2-jet determination for D-nondegenerate CR manifolds. We recover their result by proving the following.

**Proposition 6.6.** *Suppose  $F$  is D-nondegenerate. Then  $F$  is  $T$ -nondegenerate, therefore,  $\mathfrak{g}_3 = 0$ .*

**Lemma 6.7.**  $T(a)^F \subset (a^{\operatorname{Re} F})^{\operatorname{Re} F} \cap a^{\operatorname{Re} F}$ .

To compare with (6), we observe that  $(a^{\operatorname{Re} F})^{\operatorname{Re} F} \subset (a^F)^F$ . Indeed,  $a^F \subset a^{\operatorname{Re} F}$  implies  $(a^{\operatorname{Re} F})^{\operatorname{Re} F} \subset (a^F)^{\operatorname{Re} F}$ . Since  $a^F$  is a complex subspace, we have  $(a^F)^{\operatorname{Re} F} = (a^F)^F$ , hence  $(a^{\operatorname{Re} F})^{\operatorname{Re} F} \subset (a^F)^F$ .

*Proof.* By putting  $a = b$  and  $x = y$  in (5), we obtain  $a^{\operatorname{Re} F} \subset T(a)$ . Then

$$T(a)^F \subset (a^{\operatorname{Re} F})^F \subset (a^{\operatorname{Re} F})^{\operatorname{Re} F}.$$

Since  $a \in T(a)$ , we have  $T(a)^F \subset a^F \subset a^{\operatorname{Re} F}$ , and the lemma follows.  $\square$

**Lemma 6.8.** *Suppose  $\operatorname{Re} B$  is nonsingular. Then  $(a^{\operatorname{Re} F})^{\operatorname{Re} F} \cap a^{\operatorname{Re} F} = 0$ .*

*Proof.* Define  $V = \operatorname{Span}_{\mathbb{R}}\{A_j a : 1 \leq j \leq k\}$ . We claim that  $a^{\operatorname{Re} F} = V^{\perp_{\mathbb{R}}}$ , the orthogonal complement of  $V$  with respect to the standard *real* inner product  $\operatorname{Re} \langle \cdot, \cdot \rangle$ . Indeed,  $z \in a^{\operatorname{Re} F}$  means exactly that  $\operatorname{Re} \langle A_j a, z \rangle = 0$  for every  $j$ , that is,  $z \in V^{\perp_{\mathbb{R}}}$ .

We now claim  $z \in (a^{\operatorname{Re} F})^{\operatorname{Re} F}$  iff  $A_j z \in V$  for every  $j$ . Indeed, it means that  $\operatorname{Re} \langle A_j z, x \rangle = 0$  for every  $1 \leq j \leq k$  and  $x \in a^{\operatorname{Re} F} = V^{\perp_{\mathbb{R}}}$ . That is,  $A_j z \in (V^{\perp_{\mathbb{R}}})^{\perp_{\mathbb{R}}} = V$  for every  $j$ .

Let  $z \in (a^{\operatorname{Re} F})^{\operatorname{Re} F} \cap a^{\operatorname{Re} F}$ . Since  $z \in (a^{\operatorname{Re} F})^{\operatorname{Re} F}$ , we have  $A_j z \in V$  for all  $j$ . Recall  $A = \sum c_j A_j$ ,  $c \in \mathbb{R}^k$ . Then  $Az \in V$ . Then there is  $\lambda \in \mathbb{R}^k$  such that  $Az = \sum \lambda_j A_j a$ . Then  $z = \sum \lambda_j A^{-1} A_j a$ .

Since  $z \in a^{\operatorname{Re} F}$ , we have  $\operatorname{Re} \langle A_l z, a \rangle = 0$  for every  $l$ . Using the above expression for  $z$ , we obtain  $\operatorname{Re} \langle \sum_j \lambda_j A_l A^{-1} A_j a, a \rangle = 0$ . By (13) we rewrite  $\operatorname{Re} \sum_j B_{lj} \lambda_j = (\operatorname{Re} B) \lambda = 0$ .

Since  $\operatorname{Re} B$  is nonsingular,  $\lambda = 0$ . Then  $z = 0$ , hence the conclusion.  $\square$

*Proof of Proposition 6.6.* We observe that the condition  $\det \operatorname{Re} B \neq 0$  is stable under small perturbations of  $a \in \mathbb{C}^m$ , therefore, by Lemmas 6.7 and 6.8, we have  $T(a)^F = 0$  for generic  $a \in \mathbb{C}^m$ , that is,  $F$  is T-nondegenerate, as desired.  $\square$

We finally observe that in Tanaka's [14] cases  $k = m^2 - 1$  and  $k = m^2$ , nondegenerate CR manifolds are T-nondegenerate. Indeed, for  $k = m^2 - 1$ , one can see that for generic  $a \in \mathbb{C}^m$  we have  $a^F = 0$ , so by Proposition 6.2 the manifold is T-nondegenerate. Then the case  $k = m^2$  also falls into this category, but it is also strongly pseudoconvex.

## 7. CODIMENSION $k \leq 3$

By Tanaka [14] and Chern and Moser [10], 2-jet determination holds for codimension  $k = 1$ . Blanc-Centi and Meylan [7] prove 2-jet determination for holomorphic CR mappings for  $k = 2$ . Beloshapka [3] proves that  $\mathfrak{g}_3 = 0$  for  $k = 3$ . We recover these results here by proving the following.

**Proposition 7.1.** *Let  $F$  be a nondegenerate quadric with  $k \leq 3$ . Then  $F$  is T-nondegenerate, hence  $\mathfrak{g}_3 = 0$ .*

For  $k \leq 2$  we have a stronger result.

**Proposition 7.2.** *Let  $F$  be a nondegenerate quadric with  $k \leq 2$ . Then for generic  $a \in \mathbb{C}^m$ , we have  $(a^F)^F \cap a^F = 0$ .*

For  $k = 1$  the result is obvious. Indeed, for  $k = 1$  we have  $(a^F)^F = \mathbb{C}a$ . If  $F(a, a) \neq 0$ , then  $(a^F)^F \cap a^F = 0$ .

**Lemma 7.3.** *Let  $L$  be a  $m \times m$  matrix such that for all  $a, b \in \mathbb{C}^m$ ,*

$$(14) \quad \langle La, b \rangle^2 = \langle a, b \rangle \langle L^2 a, b \rangle.$$

*Then  $L = \lambda I$ , a scalar matrix.*

*Proof.* The equation (14) is invariant under similarity  $L \rightarrow PLP^{-1}$ ,  $a \rightarrow Pa$ ,  $b \rightarrow P^{-1}b$ . Then without loss of generality we can assume that  $L$  is in Jordan normal form.

We first show that there are no Jordan cells of order greater than 1, that is,  $L$  is diagonal. Suppose, otherwise, that there is a Jordan cell of order greater than 1. Without loss of generality we assume that  $L$  consists of a single Jordan cell  $L = \lambda I + J$ , here  $J$  is the standard Jordan cell with zero eigenvalue and units above the main diagonal.

Then after some cancellations, (14) reduces to  $\langle Ja, b \rangle^2 = \langle a, b \rangle \langle J^2 a, b \rangle$ . For  $b = e_1$ , the first element of the standard basis, this equation turns into  $a_2^2 = a_1 a_3$  ( $a_2 = 0$  if  $m = 2$ ), which is absurd because  $a$  is arbitrary. Hence  $L$  is diagonal.

Let  $L$  be diagonal with eigenvalues  $\lambda_1, \dots, \lambda_m$ . Then we have

$$\langle a, b \rangle = \sum a_j \bar{b}_j, \quad \langle La, b \rangle = \sum \lambda_j a_j \bar{b}_j, \quad \langle L^2 a, b \rangle = \sum \lambda_j^2 a_j \bar{b}_j.$$

Put  $c_j = a_j \bar{b}_j$ . Then (14) turns into

$$\left( \sum \lambda_j c_j \right)^2 = \left( \sum c_j \right) \left( \sum \lambda_j^2 c_j \right).$$

Put  $c_1 = c_2 = 1$ ,  $c_j = 0$  for  $j > 2$ . Then  $(\lambda_1 + \lambda_2)^2 = 2(\lambda_1^2 + \lambda_2^2)$ , which implies  $\lambda_1 = \lambda_2$ . Similarly, all  $\lambda_j$ -s are equal, hence  $L$  is scalar, as desired.  $\square$

The following lemma must be well known.

**Lemma 7.4.** *Let  $A$  and  $B$  be linearly independent  $m \times m$  Hermitian matrices. Then for generic  $a \in \mathbb{C}^m$ , the vectors  $Aa$  and  $Ba$  are  $\mathbb{C}$ -linearly independent.*

For completeness, we include a proof.

*Proof.* Arguing by contradiction, we assume that for all  $z \in \mathbb{C}^m$ , the vectors  $Az$  and  $Bz$  are linearly dependent.

Let  $r = \max\{\text{rank}(\alpha A + \beta B) : \alpha, \beta \in \mathbb{R}\}$ .

We first consider the case  $r = m$ . For definiteness,  $A$  is nonsingular. Then for all  $z \in \mathbb{C}^m$ , there is  $\lambda \in \mathbb{C}$  such that  $Bz = \lambda Az$ , that is,  $A^{-1}Bz = \lambda z$ . Then every  $z \in \mathbb{C}^m$  is an eigenvector of the matrix  $A^{-1}B$ . It can occur only if  $A^{-1}B$  is a scalar matrix. Then  $A^{-1}B = \lambda I$ , and  $B = \lambda A$ , which is absurd. Thus for  $r = m$  the lemma holds.

We now consider the case  $r < m$ . Without loss of generality assume  $A$  is diagonal and  $\text{rank } A = r$ . We represent  $A$  and  $B$  as  $2 \times 2$  block matrices  $A = (A_{pq})$ ,  $B = (B_{pq})$ ,  $p, q = 1, 2$  according to splitting the coordinates into two groups of  $r$  and  $m - r$  coordinates. We number the coordinates so that the  $r \times r$  matrix  $A_{11}$  is nonsingular and the rest  $A_{pq} = 0$ . Since  $\text{rank}(tA + B) \leq r$  for big  $t \in \mathbb{R}$ , we must have  $B_{22} = 0$ .

By the hypothesis, for every  $z = (z_1, z_2) \in \mathbb{C}^m$ , we have  $Bz = \lambda Az$  or  $Az = 0$ . The equation  $Az = 0$  means that  $z_1 = 0$ . The equation  $Bz = \lambda Az$ , in particular, implies  $B_{21}z_1 = 0$ . Then for every  $z_1 \neq 0$  we have  $B_{21}z_1 = 0$ . Then  $B_{21} = 0$ . Since  $B$  is Hermitian, we have also  $B_{12} = 0$ . Then the result reduces to the case  $r = m$  for the matrices  $A_{11}$  and  $B_{11}$  that we considered earlier.  $\square$

*Proof of Proposition 7.2.* Let  $k = 2$ . We change notation from  $A_1$  and  $A_2$  to  $A$  and  $B$ .

As in the proof of Lemma 7.4, let  $r = \max\{\text{rank}(\alpha A + \beta B) : \alpha, \beta \in \mathbb{R}\}$ .

We first consider the strongly nondegenerate case  $r = m$ . For definiteness,  $A$  is nonsingular. Arguing by contradiction, we assume that for generic  $a \in \mathbb{C}^m$  we have  $(a^F)^F \cap a^F \neq 0$ .

Put  $U = \text{Span}_{\mathbb{C}}(Aa, Ba)$ . We claim that  $a^F = U^\perp$ , here  $U^\perp$  stands for the orthogonal complement of  $U$  with respect to the standard *complex* inner product. We claim that  $z \in (a^F)^F$  iff  $Az, Bz \in U$ . The proofs of these claims are similar to the proofs of the corresponding claims in the proof of Lemma 6.8.

Let  $z \in (a^F)^F \cap a^F$ ,  $z \neq 0$ . Since  $z \in (a^F)^F$ , we have  $Az = \alpha Aa + \beta Ba$  with  $\alpha, \beta \in \mathbb{C}$ . Since  $z \in a^F$ , we have  $0 = \langle Az, a \rangle = \alpha \langle Aa, a \rangle + \beta \langle Ba, a \rangle$ . For generic  $a$  we can assume  $\langle Aa, a \rangle \neq 0$ . Then  $\alpha = -\beta \langle Aa, a \rangle^{-1} \langle Ba, a \rangle$ . Then  $Az = -\beta (\langle Aa, a \rangle^{-1} \langle Ba, a \rangle Aa - Ba)$ . Since  $A$  is nonsingular, we have  $Az \neq 0$ , and  $\beta \neq 0$ . Without loss of generality  $\beta = -\langle Aa, a \rangle$ . Then  $Az = \langle Ba, a \rangle Aa - \langle Aa, a \rangle Ba$ . Then  $z = \langle Ba, a \rangle a - \langle Aa, a \rangle A^{-1} Ba$ . Then  $Bz = \langle Ba, a \rangle Ba - \langle Aa, a \rangle BA^{-1} Ba$ .

Since  $z \in a^F$ , we have  $0 = \langle Bz, a \rangle = \langle Ba, a \rangle^2 - \langle Aa, a \rangle \langle BA^{-1} Ba, a \rangle$ .

Since the equation  $\langle Ba, a \rangle^2 = \langle Aa, a \rangle \langle BA^{-1} Ba, a \rangle$  holds for generic  $a$ , it holds for all  $a \in \mathbb{C}^m$ . By polarization, we have  $\langle Ba, b \rangle^2 = \langle Aa, b \rangle \langle BA^{-1} Ba, b \rangle$ . By replacing  $a$  by  $A^{-1}a$ , we have  $\langle BA^{-1}a, b \rangle^2 = \langle a, b \rangle \langle BA^{-1} BA^{-1}a, b \rangle$ .

Put  $L = BA^{-1}$ . Then we have  $\langle La, b \rangle^2 = \langle a, b \rangle \langle L^2a, b \rangle$  for all  $a, b \in \mathbb{C}^m$ . By Lemma 7.3  $L = \lambda I$  then  $B = \lambda A$ , which is absurd. Thus the result holds in the strongly nondegenerate case  $r = m$ .

Now consider the case  $r < m$ , that is, every linear combination of  $A$  and  $B$  is singular. Without loss of generality assume  $A$  is diagonal and  $\text{rank } A = r$ .

As in the proof of Lemma 7.4, we represent  $A$  and  $B$  as  $2 \times 2$  block matrices  $A = (A_{pq})$ ,  $B = (B_{pq})$ ,  $p, q = 1, 2$ . We assume that the  $r \times r$  matrix  $A_{11}$  is nonsingular and the rest  $A_{pq} = 0$ . Then we again have  $B_{22} = 0$ . Since  $F$  is nondegenerate,  $B_{21} \neq 0$ .

We take  $a = (a_1, a_2) \in \mathbb{C}^m$  such that  $\langle Aa, a \rangle = \langle A_{11}a_1, a_1 \rangle \neq 0$  and  $B_{21}a_1 \neq 0$ . We prove  $(a^F)^F \cap a^F = 0$ .

Let  $z = (z_1, z_2) \in (a^F)^F \cap a^F$ . As above, we have  $Az = \alpha Aa + \beta Ba$  with  $\alpha, \beta \in \mathbb{C}$  and  $0 = \langle Az, a \rangle = \alpha \langle Aa, a \rangle + \beta \langle Ba, a \rangle$ . In more detail,

$$Az = \begin{pmatrix} A_{11}z_1 \\ 0 \end{pmatrix} = \alpha \begin{pmatrix} A_{11}a_1 \\ 0 \end{pmatrix} + \beta \begin{pmatrix} \cdots \\ B_{21}a_1 \end{pmatrix}.$$

Since  $B_{21}a_1 \neq 0$ , we have  $\beta = 0$ . Then  $\alpha \langle Aa, a \rangle = 0$ . Since  $\langle Aa, a \rangle \neq 0$ , we have  $\alpha = 0$ . Then  $Az = 0$ , that is,  $A_{11}z_1 = 0$ , hence  $z_1 = 0$ .

Similarly, we have  $Bz = \gamma Aa + \delta Ba$  with  $\gamma, \delta \in \mathbb{C}$  and  $0 = \langle Bz, a \rangle = \gamma \langle Aa, a \rangle + \delta \langle Ba, a \rangle$ . Since  $z_1 = 0$ , we have

$$Bz = \begin{pmatrix} B_{12}z_2 \\ 0 \end{pmatrix} = \gamma \begin{pmatrix} A_{11}a_1 \\ 0 \end{pmatrix} + \delta \begin{pmatrix} \cdots \\ B_{21}a_1 \end{pmatrix}.$$

By the same argument as above, we have  $\delta = \gamma = 0$ , and  $Bz = 0$ . Since  $F$  is nondegenerate,  $Az = Bz = 0$  implies  $z = 0$ , and  $(a^F)^F \cap a^F = 0$ , as desired.  $\square$

*Proof of Proposition 7.1.* By Proposition 7.2, it only remains to consider the case  $k = 3$ . Let  $r = \max\{\text{rank}_{\mathbb{C}}(A_1a, A_2a, A_3a) : a \in \mathbb{C}^m\}$ . By Lemma 7.4, the case  $r = 1$  can not occur.

If  $r = 3$ , then for generic  $a$ , the vectors  $A_1a$ ,  $A_2a$ , and  $A_3a$  are  $\mathbb{C}$ -linearly independent. Then by Proposition 6.3,  $F$  is T-nondegenerate.

Finally, consider the case  $r = 2$ . For definiteness, assume that for generic  $a \in \mathbb{C}^m$ , the vectors  $A_1a$  and  $A_2a$  are  $\mathbb{C}$ -linearly independent, then  $A_3a$  is a  $\mathbb{C}$ -linear combination of  $A_1a$  and  $A_2a$ .

We denote by  $F'$  the form defined by the matrices  $A_1$  and  $A_2$ . Then  $F'$  is nondegenerate. Indeed, let  $a \in \mathbb{C}^m$  be a vector such that for all  $z \in \mathbb{C}^m$ , we have  $F'(z, a) = 0$ . For generic  $z$ , the vector  $A_3z$  is a  $\mathbb{C}$ -linear combination of  $A_1z$  and  $A_2z$ . Then for generic  $z$ , we have  $F(z, a) = 0$ . Then  $F(z, a) = 0$  for all  $z$ . Since  $F$  is nondegenerate,  $a = 0$ . Thus  $F'$  is nondegenerate.

By Proposition 7.2, for generic  $a$ , we have  $(a^{F'})^{F'} \cap a^{F'} = 0$ . Since  $A_3a$  is a  $\mathbb{C}$ -linear combination of  $A_1a$  and  $A_2a$  for generic  $a$ , we have  $a^F = a^{F'}$  for generic  $a$ . Then for generic  $a$ , we have  $(a^F)^F = (a^{F'})^F \subset (a^{F'})^{F'}$ , then  $(a^F)^F \cap a^F \subset (a^{F'})^{F'} \cap a^{F'} = 0$ . Now by Proposition 6.2,  $F$  is T-nondegenerate, as desired.  $\square$

## 8. EXAMPLES

**Example 8.1.** Meylan [12] has found an example of a strongly nondegenerate quadric for which  $\mathfrak{g}_4 \neq 0$ , so 2-jet determination fails. Here  $m = 4, k = 5, F = (F_1, \dots, F_5)$ .

$$\begin{aligned} F_1(z, z) &= \operatorname{Re}(z_1\bar{z}_3 + z_2\bar{z}_4), \\ F_2(z, z) &= |z_1|^2, \\ F_3(z, z) &= |z_2|^2, \\ F_4(z, z) &= \operatorname{Re}(z_1\bar{z}_2), \\ F_5(z, z) &= \operatorname{Im}(z_1\bar{z}_2). \end{aligned}$$

In this example,  $F$  is *not* T-nondegenerate, and  $a^F \cap (a^F)^F \neq 0$  for generic  $a$ .

**Example 8.2.** Let  $m = 4, k = 4$ , and  $F = (F_1, F_2, F_3, F_4)$  from Example 8.1. Then  $F$  is *not* T-nondegenerate, but one can see that  $\mathfrak{g}_3 = 0$ , so 2-jet determination takes place. This example shows that T-nondegeneracy is not necessary for  $\mathfrak{g}_3 = 0$  to hold.

**Example 8.3.** Let  $m = 4, k = 3$ , and  $F = (F_1, F_2, F_3)$  from Example 8.1. Since  $k = 3$ , the form  $F$  is T-nondegenerate,  $\mathfrak{g}_3 = 0$ , so 2-jet determination takes place. However, one can see that  $F$  is *not* D-nondegenerate and  $(a^F)^F \cap a^F \neq 0$  for generic  $a$ . This example shows that Proposition 7.2 does not necessarily hold for  $k = 3$ .

**Example 8.4.** Monomial quadric. We call  $F$  monomial if all components of  $F$  have the form  $\operatorname{Re}(z_p\bar{z}_q)$  (in particular,  $|z_p|^2$ ) or  $\operatorname{Im}(z_p\bar{z}_q)$  ( $p \neq q$ ), and there are no repeated components. A monomial quadric  $F$  is nondegenerate iff each variable  $z_p$  occurs in at least one component of  $F$ . For such a quadric, the codimension  $k$  can be any integer  $\frac{m}{2} \leq k \leq m^2$ . One can see that if  $a \in \mathbb{C}^m$  has no zero components, then  $(a^F)^F \cap a^F = 0$ . Hence  $F$  is T-nondegenerate. However,  $F$  can not be D-nondegenerate if  $k > 2m$ .

**Example 8.5.** Monomial antisymmetric quadric of an odd dimension. This is a special case of the previous example in which  $m \geq 3$  is an odd integer, and all components of  $F$  have the form  $\text{Im}(z_p \bar{z}_q)$ . Then  $k \leq \frac{m(m-1)}{2}$ . Note that  $F$  is not *strongly* nondegenerate, hence not D-nondegenerate. Indeed, the matrices of the components of  $F$  are antisymmetric, and every linear combination of them is an antisymmetric matrix of an odd order, hence singular. As a special case of the previous example, if  $F$  is nondegenerate, then  $F$  is T-nondegenerate, and 2-jet determination holds. However, this result would be difficult to obtain by means of stationary discs because  $F$  is not strongly nondegenerate.

**Conflict of interest statement:** There is no conflict of interest.

**Data availability statement:** This manuscript has no associated data.

#### REFERENCES

- [1] M. S. Baouendi, Strong unique continuation and finite jet determination for Cauchy-Riemann mappings. Phase space analysis of partial differential equations, 17–28, Progr. Nonlinear Differential Equations Appl., 69, Birkhäuser Boston, Boston, MA, 2006.
- [2] V. K. Beloshapka, Finite-dimensionality of the group of automorphisms of a real analytic surface. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 52 (1988), 437–442; translation in Math. USSR-Izv. 32 (1989), 443–448.
- [3] V. K. Beloshapka, On special quadrics. Russ. J. Math. Phys. 29 (2022), 11–27.
- [4] F. Bertrand, L. Blanc-Centi, and F. Meylan, Stationary discs and finite jet determination for nondegenerate generic real submanifolds. Adv. Math. 343 (2019), 910–934. Addendum: Adv. Math. 383 (2021), 107701, 3 pp.
- [5] F. Bertrand and F. Meylan, Nondefective stationary discs and 2-jet determination in higher codimension. J. Geom. Anal. 31 (2021), 6292–6306.
- [6] F. Bertrand and F. Meylan, The 1-jet determination of stationary discs attached to generic CR submanifolds. Preprint, arXiv:2112.12176
- [7] L. Blanc-Centi and F. Meylan, Chern-Moser operators and weighted jet determination problems in higher codimension. Internat. J. Math. 33 (2022), Paper No. 2250045, 15 pp.
- [8] L. Blanc-Centi and F. Meylan, On nondegeneracy conditions for the Levi map in higher codimension: a survey. Preprint, arXiv:1711.11481
- [9] J. Gregorovič and F. Meylan, Construction of counterexamples to the 2-jet determination Chern-Moser Theorem in higher codimension. Preprint, arXiv:2010.10220
- [10] S. S. Chern and J. K. Moser, Real hypersurfaces in complex manifolds. Acta Math. 133 (1974), 219–271.
- [11] A. Isaev and W. Kaup, Regularization of local CR-automorphisms of real-analytic CR-manifolds. J. Geom. Anal. 22 (2012), 244–260.
- [12] F. Meylan, A counterexample to the 2-jet determination Chern-Moser theorem in higher codimension. Preprint, arXiv: 2003.11783.
- [13] E. M. Stein, Singular integrals and estimates for the Cauchy-Riemann equations. Bull. Amer. Math. Soc. 79 (1973), 440–445.
- [14] N. Tanaka, On generalized graded Lie algebras and geometric structures I. J. Math. Soc. Japan, 19 (1967), 215–254.



- [15] A. Tumanov, Finite dimensionality of the group of CR automorphisms of a standard CR manifold, and proper holomorphic mappings of Siegel domains. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 52 (1988), 651–659; translation in *Math. USSR-Izv.* 32 (1989), 655–662.
- [16] A. Tumanov, Stationary discs and finite jet determination for CR mappings in higher codimension. *Adv. Math.* 371 (2020), 107254, 11 pp.
- [17] D. Zaitsev, Unique determination of local CR-maps by their jets: a survey. Harmonic analysis on complex homogeneous domains and Lie groups (Rome, 2001). *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.* 13 (2002), 295–305.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 WEST GREEN ST., URBANA, IL 61801

*Email address:* `tumanov@illinois.edu`