### Generalized Hilbert Operator Acting on Weighted Bergman Spaces and on Dirichlet Spaces\*

Shanli Ye<sup>†</sup> Guanghao Feng<sup>‡</sup>

(School of Sciences, Zhejiang University of Science and Technology, Hangzhou 310023, China)

#### Abstract

Let  $\mu$  be a positive Borel measure on the interval [0,1). For  $\beta > 0$ , The generalized Hankel matrix  $\mathcal{H}_{\mu,\beta} = (\mu_{n,k,\beta})_{n,k\geq 0}$  with entries  $\mu_{n,k,\beta} = \int_{[0,1)} \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} t^{n+k} d\mu(t)$ , induces formally the operator

$$\mathcal{H}_{\mu,\beta}(f)(z) = \sum_{n=0}^{\infty} \left( \sum_{k=0}^{\infty} \mu_{n,k,\beta} a_k \right) z^n$$

on the space of all analytic function  $f(z) = \sum_{k=0}^{\infty} a_k z^n$  in the unit disc  $\mathbb{D}$ . In this paper, we characterize those positive Borel measures on [0,1) such that  $\mathcal{H}_{\mu,\beta}(f)(z) = \int_{[0,1)} \frac{f(t)}{(1-tz)^{\beta}} d\mu(t)$  for all in weighted Bergman Spaces  $A^p_{\alpha}(0 -1)$ , and among them we describe those for which  $\mathcal{H}_{\mu,\beta}(\beta > 0)$  is a bounded(resp.,compact) operator on weighted Bergman spaces and Dirichlet spaces.

**Keywords** Generalized Hilbert operator, Weighted Bergman Spaces, Dirichlet spaces, Carleson measure

**2020** MR Subject Classification 47B35, 30H20, 30H30

### 1 Introduction

Let  $\mu$  be a positive Borel measure on the interval [0,1). for  $\beta > 0$ , we define  $\mathcal{H}_{\mu,\beta} = (\mu_{n,k,\beta})_{n,k\geq 0}$  to be generalized Hankel matrix with entries  $\mu_{n,k,\beta} = \int_{[0,1]} \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} t^{n+k} d\mu(t)$ .

For analytic functions  $f(z) = \sum_{k=0}^{\infty} a_k z^k$ , the generalized Hilbert operator define as

$$\mathcal{H}_{\mu,\beta}(f)(z) = \sum_{n=0}^{\infty} \left( \sum_{k=0}^{\infty} \mu_{n,k,\beta} a_k \right) z^n, \tag{1.1}$$

whenever the right hand side makes sense and defines an analytic function in D.

In recent decades, the generalized Hilbert operator  $\mathcal{H}_{\mu,1}$  which induced by the Hankel matrix  $\mathcal{H}_{\mu}$  have been studied extensively. For example, Galanopoulos and Peláez[8] characterized the Borel measure  $\mu$  for which the Hankel operator is a bounded (resp.,compact) operator on  $H^1$ . Then Chatzifountas, Girela and Peláez [4] extended this work the all Hardy spaces  $H^p$  with  $0 . In [9], Girela and Merchán studied the operator acting on certain conformally invariant spaces of analytic functions on the disk. For the case <math>\beta = 2$ , it had been studied the

<sup>\*</sup>The research was supported by the National Natural Science Foundation of China (Grant No. 11671357)

<sup>&</sup>lt;sup>†</sup>Corresponding author. E-mail address: slye@zust.edu.cn

<sup>&</sup>lt;sup>‡</sup>E-mail address:gh945917454@foxmail.com

operator in [18, 19] and the  $\mathcal{H}_{\mu,2}$  was called the Derivative-Hilbert operator. In [20], the  $\mathcal{H}_{\mu,\beta}$  was called the generalized Hilbert operator.

Another generalized Hilbert-integral operator related to  $\mathcal{H}_{\mu,\beta}$  denoted by  $\mathcal{I}_{\mu,\beta}(\beta \in \mathbb{R}^+)$  is defined by

$$\mathcal{I}_{\mu,\beta}(f)(z) = \int_{[0,1)} \frac{f(t)}{(1-tz)^{\beta}} d\mu(t),$$

whenever the right hand side makes sense and defines an analytic function in  $\mathbb{D}$ .

In this paper, we consider the operators

$$\mathcal{H}_{\mu,\beta}: A^p_{\alpha} \to A^q_{\beta-2}, \quad 0 -1 \text{ and } \beta > 1.$$
  
 $\mathcal{H}_{\mu,\beta}: \mathcal{D}^p_{\alpha} \to \mathcal{D}^q_{\beta-1}, \quad 1 -1 \text{ and } \beta > 0.$ 

The aim is to study the boundedness (resp.,compactness) of  $\mathcal{H}_{\mu,\beta}$ .

In this article we characterize the positive Borel measures  $\mu$  for which the operator which  $\mathcal{I}_{\mu,\beta}$  and  $\mathcal{H}_{\mu,\beta}$  is well defined in the weighted Bergman space  $A^p_{\alpha}$ . Then we give the necessary and sufficient conditions such that operator  $\mathcal{H}_{\mu,\beta}$  is bounded (resp.,compact) from the weighted Bergman spaces  $A^p_{\alpha}(0 -1)$  into the space  $A^q_{\beta-2}(\beta > 1, q \ge 1 \text{ and } p \le q < \infty)$ , and we also give necessary and sufficient conditions such that operator  $\mathcal{H}_{\mu,\beta}$  is bounded (resp.,compact) from the Dirichlet type space  $\mathcal{D}^p_{\alpha}(\alpha > p - 1, 0 into the space <math>\mathcal{D}^q_{\beta-1}(\beta > q, p \le q < \infty \text{ and } q > 1)$ , and get a series of corollaries.

### 2 Notation and Preliminaries

Let  $\mathbb{D}$  denote the open unit disk of the complex plane, and let  $H(\mathbb{D})$  denote the set of all analytic functions in  $\mathbb{D}$ .

For  $0 and <math>\alpha > -1$  the weighted Bergman space  $A^p_\alpha$  consists of those  $f \in H(\mathbb{D})$  such that

$$||f||_{A^p_{\alpha}} \stackrel{def}{=} \left( (\alpha+1) \int_{\mathbb{D}} |f(z)|^p (1-|z|^2)^{\alpha} dA(z) \right)^{1/p} < \infty.$$

Here, dA stands for the area measure on  $\mathbb{D}$ , normalized so that the total area of  $\mathbb{D}$  is 1. Thus  $dA(z) = \frac{1}{\pi} dx dy = \frac{1}{\pi} r dr d\theta$ . The unweighted Bergman space  $A_0^p$  is simply denoted by  $A^p$ . We refer to [2, 3] for the notation and results about Bergman spaces.

The space of Dirichlet type  $\mathcal{D}^p_{\alpha}$  (0 -1) consists of those  $f \in H(\mathbb{D})$  such that  $f' \in A^p_{\alpha}$ . In other words, a function  $f \in H(\mathbb{D})$  belongs to  $\mathcal{D}^p_{\alpha}$  if and only if

$$||f||_{\mathcal{D}^p_{\alpha}} \stackrel{def}{=} |f(0)| + \left( (\alpha + 1) \int_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^{\alpha} dA(z) \right)^{1/p} < \infty.$$

Finally, we recall that a function  $f \in H(\mathbb{D})$  is said to be a Bloch function if

$$||f||_{\mathcal{B}} \stackrel{def}{=} |f(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| < \infty.$$

The space of all Bloch functions is denoted by  $\mathcal{B}$ . A classical reference for the theory of Bloch functions is [11, 16]. The little Bloch space  $\mathcal{B}_0$  is the subspace of the Bloch space consisting of those functions  $f \in H(\mathbb{D})$  such that.

$$\lim_{|z|\to 0} (1-|z|^2)|f'(z)| = 0.$$

Let us recall the knowledge of Carleson measure, which is a very useful tool in the study of Banach spaces of analytic functions. For  $0 < s < \infty$ , a positive Borel measure  $\mu$  on  $\mathbb D$  will be called a s-Carleson measure, if there exists a positive constant C such that

$$\mu(S(I)) \le C|I|^s$$
.

The Carleson square S(I) is defined as

$$S(I) = \{z = re^{i\theta} : e^{i\theta} \in I; 1 - \frac{|I|}{2\pi} \le r \le 1\}.$$

where I is an interval of  $\partial \mathbb{D}$ , |I| denotes the length of I, if  $\mu$  satisfies  $\lim_{|I|\to 0} \frac{\mu(S(I))}{|I|^s} = 0$ , we call  $\mu$  is a vanishing s-Carleson measure.

Let  $\mu$  be a positive Borel measure on  $\mathbb{D}$ . For  $0 \le \alpha < \infty$  and  $0 < s < \infty$  we say that  $\mu$  is  $\alpha$ -logarithmic s-Carleson measure, if there exists a positive constant C such that

$$\frac{\mu(S(I))(\log \frac{2\pi}{|I|})^{\alpha}}{|I|^{s}} \le C \qquad I \subset \partial \mathbb{D}.$$

If  $\mu(S(I))(\log \frac{2\pi}{|I|})^{\alpha} = o(|I|^s)$ , as  $|I| \to 0$ , we say that  $\mu$  is vanishing  $\alpha$ -logarithmic s-Carleson measure [15, 21].

Suppose that  $\mu$  is a s-Carleson measure on  $\mathbb{D}$ , we see that the identity mapping i is well defined from  $A^p_\alpha$  into  $L^q(\mathbb{D},\mu)$ . Let  $\mathcal{N}(\mu)$  be the norm of i. For 0 < r < 1, let

$$d\mu_r(z) = \chi_{r < |z| < 1}(t)d\mu(t).$$

Then  $\mu$  is a vanishing s-Carleson measure if and only if

$$\mathcal{N}(\mu_r) \to 0 \quad as \quad r \to 1^-.$$
 (2.1)

A positive Borel measure on [0,1) also can be seen as a Borel measure on  $\mathbb D$  by identifying it with the measure  $\mu$  defined by

$$\tilde{\mu}(E) = \mu(E \bigcap [0,1)).$$

for any Borel subset E of  $\mathbb{D}$ . Then a positive Borel measure  $\mu$  on [0,1) can be seen as an s-Carleson measure on  $\mathbb{D}$ , if

$$\mu([t,1)) \le C(1-t)^s, \quad 0 \le t < 1.$$

Also, we have similar statements for vanishing s-Carleson measures,  $\alpha$ -logarithmic s-Carleson and vanishing  $\alpha$ -logarithmic s-Carleson measures.

As usual, throughout this paper, C denotes a positive constant which depends only on the displayed parameters but not necessarily the same from one occurrence to the next. For any given p > 1, p' will denote the conjugate index of p, that is, 1/p + 1/p' = 1.

# 3 Conditions such that $\mathcal{H}_{\mu,\beta}$ is well defined on Weighted Bergman spaces

In this section, we find the sufficient condition such that  $\mathcal{H}_{\mu,\beta}(\beta > 0)$  are well defined in  $A^p_{\alpha}(0 -1)$ , and obtain that  $\mathcal{H}_{\mu,\beta}(f) = \mathcal{I}_{\mu,\beta}(f)$ , for all  $f \in A^p_{\alpha}$ , with the certain condition. We first recall a result about the coefficients of functions in weighted Bergman spaces.

**Lemma 3.1** [12, p. 191] Let  $f \in H(\mathbb{D})$ ,  $f(z) = \sum_{n=1}^{\infty} a_n z^n$  and let  $-1 < \alpha < \infty$ .

- (i) If  $0 , and <math>f \in A_{\alpha}^{p}$ , then  $a_n = o(n^{(\alpha+2)/p-1})$ .
- (ii) If  $1 \le p < \infty$  and  $f \in A_{\alpha}^p$ , then  $a_n = o(n^{(\alpha+1)/p})$ .
- (iii) If 0 , then

$$\sum_{n=1}^{\infty} n^{-\alpha-1} |a_n|^p < \infty \Rightarrow f \in A^p_{\alpha} \Rightarrow \sum_{n=1}^{\infty} n^{p-\alpha-3} |a_n|^p < \infty.$$

(iv) If  $2 \le p < \infty$ , then

$$\sum_{n=1}^{\infty} n^{p-\alpha-3} |a_n|^p < \infty \Rightarrow f \in A_{\alpha}^p \Rightarrow \sum_{n=1}^{\infty} n^{-\alpha-1} |a_n|^p < \infty.$$

**Lemma 3.2** For  $0 and <math>\alpha > -1$ ,  $\mu$  is a  $\left(\frac{(2+\alpha)q}{p}\right)$ -Carleson measure if and only if there exists a positive constant C such that

$$\left\{ \int_{\mathbb{D}} |f(z)|^q d\mu(z) \right\}^{1/q} \le C \|f\|_{A^p_\alpha}, \quad \text{for all } f \in A^p_\alpha. \tag{3.1}$$

The result was proved by several authors which can be found in [13] and [14],

**Theorem 3.1** Suppose that  $0 and <math>-1 < \alpha < \infty$  and let  $\mu$  be a positive Borel measure on [0,1). Then the power series in (1.1) defines a well defined analytic function in  $\mathbb{D}$  for every  $f \in A^p_\alpha$  in any of the three following cases.

- (i)  $\mu$  is a  $(\frac{\alpha+2}{p})$ -Carleson measure if 0 .
- (ii)  $\mu$  is a  $\left(\frac{\alpha+2-(p-1)^2}{p}\right)$ -Carleson measure if  $1 \leq p \leq 2$ .
- $(iii) \ \mu \ is \ a \ (\tfrac{\alpha+1}{p}) \text{-} Carleson \ measure \ if} \ 2 \leq p < \infty.$

Furthermore, in such cases we have that

$$\mathcal{H}_{\mu,\beta}(f)(z) = \int_{[0,1)} \frac{f(t)}{(1-tz)^{\beta}} d\mu(t) = \mathcal{I}_{\mu,\beta}(f)(z). \tag{3.2}$$

**Proof** (i) Suppose that  $0 and <math>\mu$  is a  $(\frac{\alpha+2}{p})$ -Carleson measure. Then (3.1) gives

$$\int_{[0,1)} |f(t)| d\mu(t) \le C ||f||_{A^p_\alpha}, \quad for \ all \ f \in A^p_\alpha.$$

Fix  $f(z) = \sum_{k=0}^{\infty} a_k z^k \in A^p_{\alpha}$  and z with |z| < r, 0 < r < 1. It follows that

$$\begin{split} \int_{[0,1)} \frac{|f(t)|}{|1-tz|^{\beta}} d\mu(t) &\leq \frac{1}{(1-r)^{\beta}} \int_{[0,1)} |f(t)| d\mu(t) \\ &\leq C \frac{1}{(1-r)^{\beta}} ||f||_{A^p_{\alpha}}. \end{split}$$

This implies that the integral  $\int_{[0,1)} \frac{f(t)}{(1-tz)^{\beta}} d\mu(t)$  uniformly converges, and that

$$\mathcal{I}_{\mu,\beta}(f)(z) = \int_{[0,1)} \frac{f(t)}{(1-tz)^{\beta}} d\mu(t) 
= \sum_{n=0}^{\infty} \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} \left( \int_{[0,1)} t^n f(t) d\mu(t) \right) z^n.$$
(3.3)

Since  $\mu$  is  $(\frac{\alpha+2}{p})$ -Carleson measure. Take  $f(z)=\sum_{k=0}^{\infty}a_kz^k\in A^p_{\alpha}$ , By [4, Proposition 1], and Lemma 3.1, we have that there exists C>0 such that

$$|\mu_{n,k,\beta}| = \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} |\mu_{n+k}| \le \frac{C}{(k+1)^{\frac{\alpha+2}{p}}} \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)},$$

$$|a_k| \le C(k+1)^{(\alpha+2)/p-1}$$
 for all  $n, k$ .

Then it follows that, for every n,

$$\sum_{k=0}^{\infty} |\mu_{n,k,\beta}| |a_k| \le C \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} \sum_{k=0}^{\infty} \frac{|a_k|}{(k+1)^{(\alpha+2)/p}} = C \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} \sum_{k=0}^{\infty} \frac{|a_k|^p |a_k|^{(1-p)}}{(k+1)^{(\alpha+2)/p}}$$

$$\le C \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} \sum_{k=0}^{\infty} \frac{|a_k|^p (k+1)^{(\frac{\alpha+2}{p}-1)(1-p)}}{(k+1)^{(\alpha+2)/p}}$$

$$= C \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} \sum_{k=0}^{\infty} (k+1)^{p-\alpha-3} |a_k|^p.$$

This, Lemma 3.1 and (3.3) readily yield that the series in (1.1) is well defined for all  $z \in \mathbb{D}$  and that

$$\mathcal{H}_{\mu,\beta}(f)(z) = \int_{[0,1)} \frac{f(t)}{(1-tz)^{\beta}} d\mu(t) \quad z \in \mathbb{D},$$

this give that  $\mathcal{H}_{\mu,\beta}(f) = \mathcal{I}_{\mu,\beta}(f)$ .

(ii) When  $1 \le p \le 2$ , since  $\mu$  is a  $\left(\frac{\alpha+2-(p-1)^2}{p}\right)$ -Carleson, Arguing as in the preceding one, we obtain that there exists a constant C > 0 such that, for every n,

$$\sum_{k=0}^{\infty} |\mu_{n,k,\beta}| |a_k| \le C \frac{\Gamma(n+\beta)}{n! \Gamma(\beta)} \sum_{k=0}^{\infty} (k+1)^{p-\alpha-3} |a_k|^p.$$

Then Lemma 3.1 again implies that the power series in (1.1) is well defined and

$$\sum_{k=0}^{\infty} \mu_{n,k,\beta} a_k = \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} \int_{[0,1)} t^n f(t) d\mu(t).$$

It follows that for each  $z \in \mathbb{D}$ ,

$$\begin{split} &\sum_{n=0}^{\infty} \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} \left( \int_{[0,1)} t^n f(t) d\mu(t) \right) |z|^n \\ &= \sum_{n=0}^{\infty} \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} \left( \sum_{k=0}^{\infty} \mu_{n,k} |a_k| \right) |z|^n \\ &\leq C \sum_{n=0}^{\infty} \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} |z|^n \sum_{k=0}^{\infty} (k+1)^{p-\alpha-3} |a_k|^p \\ &\leq \frac{C}{(1-|z|)^{\beta}} \sum_{k=0}^{\infty} (k+1)^{p-\alpha-3} |a_k|^p < \infty, \end{split}$$

we deduce that

$$\mathcal{H}_{\mu,\beta}(f)(z) = \sum_{n=0}^{\infty} \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} \left( \int_{[0,1)} t^n f(t) d\mu(t) \right) z^n$$

$$= \int_{[0,1)} \sum_{n=0}^{\infty} \frac{\Gamma(n+\beta)}{n!\Gamma(\beta)} (tz)^n f(t) d\mu(t)$$

$$= \int_{[0,1)} \frac{f(t)}{(1-tz)^{\beta}} d\mu(t), \tag{3.4}$$

which together implies that  $\mathcal{H}_{\mu,\beta}$  is a well defined for all  $z \in \mathbb{D}$ , and  $\mathcal{H}_{\mu,\beta}(f) = \mathcal{I}_{\mu,\beta}(f)$ . (iii) The proof is similar to that of (ii). We omit the details.

**Remark 3.1** Theorem 3.1 is the promotion of Theorem 2.1 in [19], when  $\alpha = 0$ , Theorem 3.1 is same as Theorem 2.1 in [19].

# 4 Bounededness of $\mathcal{H}_{\mu,\beta}$ acting on Weighted Bergman spaces and Dirichlet spaces

In this section, we mainly characterize those measures  $\mu$  for which  $\mathcal{H}_{\mu,\beta}$  is a bounded operator on the weighted Bergman spaces  $A^p_{\alpha}$  and the Dirichlet spaces  $\mathcal{D}^p_{\alpha}$  for some different  $\beta$ , p and  $\alpha$ .

**Lemma 4.1** [10] If  $1 and <math>\alpha > -1$ , then the dual of  $A^p_{\alpha}$  can be identified with  $A^{p'}_{\gamma}$  and  $\gamma$  is any number with  $\gamma > -1$ , under the pairing

$$< h, f>_{A_{p,\alpha,\gamma}} = \int_{\mathbb{D}} h(z) \overline{f(z)} (1-|z|^2)^{\frac{\alpha}{p}+\frac{\gamma}{p'}} dA(z), \quad h \in A^p_{\alpha}, \quad f \in A^{p'}_{\gamma}.$$

If  $1 and <math>\alpha > -1$ , then the dual of  $\mathcal{D}^p_{\alpha}$  can be identified with  $\mathcal{D}^{p'}_{\gamma}$  where and  $\gamma$  is any number with  $\gamma > -1$ , under the pairing

$$< h, f>_{\mathcal{D}_{p,\alpha,\gamma}} = h(0)\overline{f(0)} + \int_{\mathbb{D}} h'(z)\overline{f'(z)}(1-|z|^2)^{\frac{\alpha}{p}+\frac{\gamma}{p'}}dA(z), \quad h \in \mathcal{D}^p_{\alpha}, \quad f \in \mathcal{D}^{p'}_{\gamma}.$$

**Theorem 4.1** Suppose that  $0 , <math>\alpha > -1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If  $q \ge p$  and q > 1, then  $\mathcal{H}_{\mu,\alpha+2}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^q_{\alpha}$  if and only if  $\mu$  is  $a\left(\frac{\alpha+2}{p}+\frac{\alpha+2}{q'}\right)$ -Carleson measure.
- (ii) If  $q \ge p$  and q = 1, then  $\mathcal{H}_{\mu,\alpha+2}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^1_{\alpha}$  if and only if  $\mu$  is a 1-logarithmic  $(\frac{\alpha+2}{p})$ -Carleson measure.

The Theorem 4.1 is also closely related to Theorem 4.2, it we suffices to prove Theorem 4.2 **Theorem 4.2** Suppose that  $0 , <math>\alpha > -1$  and  $\beta > 1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If  $q \ge p$  and q > 1, if  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^q_{\beta-2}$  then  $\mu$  is a  $\left(\frac{\alpha+2}{p} + \frac{\beta}{q'}\right)$ 
  Carleson measure, conversely if  $\mu$  is a  $\left(\max\{\alpha+2,\beta\}(\frac{1}{p} + \frac{1}{q'})\right)$ -Carleson measure then  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^q_{(\beta-2)}$ .
- (ii) If  $q \ge p$  and q = 1, then  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^1_{\beta-2}$  if and only if  $\mu$  is a 1-logarithmic  $(\frac{\alpha+2}{p})$ -Carleson measure.

**Proof** Suppose that  $0 , <math>\alpha > -1$  and  $\beta > 1$ . Since  $\mu$  satisfies the condition in Theorem 3.1, as in the proof of Theorem 3.1, we obtain that

$$\int_{[0,1)} |f(t)| d\mu(t) < \infty, \quad f \in A^p_{\alpha}.$$

Hence, it follows that

$$(\alpha+1) \int_{\mathbb{D}} \int_{[0,1)} \left| \frac{f(t)g(z)(1-|z|^{2})^{\alpha}}{(1-tz)^{\beta}} \right| d\mu(t) dA(z)$$

$$\leq \frac{1}{(1-r)^{\beta}} \int_{[0,1)} |f(t)| d\mu(t) \int_{\mathbb{D}} (\alpha+1)(1-|z|^{2})^{\alpha} |g(z)| dA(z)$$

$$\leq \frac{C}{(1-r)^{\beta}} ||g||_{A_{\alpha}^{1}}, \ 0 \leq r < 1, \ f \in A_{\alpha}^{p}, \ g \in A_{\alpha}^{1},$$

$$(4.1)$$

we can obtain that the reproducing kernel function of  $A^p_{\alpha}$  is

$$K_{\alpha}(z,w) = \frac{1}{(1-z\bar{w})^{2+\alpha}}.$$

Now the reproducing property

$$f(z) = \int_{\mathbb{D}} (\alpha + 1) f(w) (1 - |w^2|)^{\alpha} K_{\alpha}(z, w) dA(w), \quad \text{for all } f \in A^1_{\alpha}.$$

Thus by using Fubini's theorem imply that

$$\int_{\mathbb{D}} \overline{\mathcal{H}_{\mu,\beta}(f)(z)} g(z) (1 - |z|^{2})^{\beta - 2} dA(z) 
= \int_{\mathbb{D}} \int_{[0,1)} \frac{\overline{f(t)} d\mu(t)}{(1 - t\overline{z})^{\beta}} g(z) (1 - |z|^{2})^{\beta - 2} dA(z) 
= \int_{[0,1)} \int_{\mathbb{D}} \frac{(1 - |z|^{2})^{\beta - 2} g(z)}{(1 - t\overline{z})^{\beta}} dA(z) \overline{f(t)} d\mu(t) 
= \frac{1}{\beta - 1} \int_{[0,1)} \overline{f(t)} g(t) d\mu(t), \quad f \in A_{\alpha}^{p}, \ g \in A_{\beta - 2}^{1}.$$
(4.2)

(i) First consider the case  $0 , <math>q \ge 1$  and  $q \ge p$ . Using (4.2) and the duality theorem for Lemma 4.1, take  $(A_{\beta-2}^q)^* \cong A_{\beta-2}^{q'}$  and  $(A_{\beta-2}^{q'})^* \cong A_{\beta-2}^q (q > 1)$ . We obtain that  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A_{\alpha}^p$  into  $A_{\beta-2}^q$  if and only if there exists a positive constant C such that

$$\left| \int_{[0,1)} \overline{f(t)} g(t) d\mu(t) \right| \le C \|f\|_{A^p_\alpha} \|g\|_{A^{q'}_{\beta-2}}, \quad f \in A^p_\alpha, \ g \in A^{q'}_{\beta-2}. \tag{4.3}$$

Assume that  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^{q'}_{\beta-2}$ . Take the families of text functions

$$f_a(z) = \left(\frac{1 - a^2}{(1 - az)^2}\right)^{(\alpha + 2)/p}$$
 and  $g_a(z) = \left(\frac{1 - a^2}{(1 - az)^2}\right)^{\beta/q'}$ ,  $0 < a < 1$ .

A calculation shows that  $\{f_a\} \subset A^p_{\alpha}, \{g_a\} \subset A^{q'}_{\beta-2}$  and

$$\sup_{a\in[0,1)}||f||_{A^p_\alpha}<\infty\quad and\quad \sup_{a\in[0,1)}||g||_{A^{q'}_{\beta-2}}<\infty.$$

It follows that

$$\infty > C \sup_{a \in [0,1)} \|f\|_{A_{\alpha}^{p}} \sup_{a \in [0,1)} \|g\|_{A_{\beta-2}^{q'}} 
\ge \left| \int_{[0,1)} f_{a}(t) g_{a}(t) d\mu(t) \right| 
\ge \int_{[a,1)} \left( \frac{1 - a^{2}}{(1 - at)^{2}} \right)^{(\alpha + 2)/p} \left( \frac{1 - a^{2}}{(1 - at)^{2}} \right)^{\beta/q'} d\mu(t) 
\ge \int_{[a,1)} \left( \frac{1 - a^{2}}{(1 - at)^{2}} \right)^{(\alpha + 2)/p + \beta/q'} d\mu(t) 
\ge \frac{1}{(1 - a^{2})^{(\alpha + 2)/p + \beta/q'}} \mu([a, 1)).$$
(4.4)

This is equivalent to saying that  $\mu$  is a  $\left(\frac{\alpha+2}{p}+\frac{\beta}{q'}\right)$ -Carleson measure.

On the other hand, if  $\mu$  is a  $\left(\max\{\alpha+2,\beta\}(\frac{1}{p}+\frac{1}{q'})\right)$ -Carleson measure. Let s=1+p/q'. Then s'=1+q'/p. By Lemma 3.2, we obtain that there exists a constant C>0 such that

$$\left( \int_{[0,1)} |f(t)|^s d\mu(t) \right)^{1/s} \le C \|f\|_{A^p_\alpha}, \quad \text{for all } f \in A^p_\alpha, \tag{4.5}$$

and

$$\left(\int_{[0,1)} |g(t)|^{s'} d\mu(t)\right)^{1/s'} \le C \|g\|_{A^{q'}_{\beta-2}}, \quad \text{for all } g \in A^{q'}_{\beta-2}. \tag{4.6}$$

Then by Hölder's inequality,

$$\begin{split} \int_{[0,1)} |f(t)| \, |g(t)| \, d\mu(t) & \leq \left( \int_{[0,1)} |f(t)|^s \, d\mu(t) \right)^{1/s} \left( \int_{[0,1)} |g(t)|^{s'} \, d\mu(t) \right)^{1/s'} \\ & \leq C \|f\|_{A^p_\alpha} \|g\|_{A^{q'}_{\beta-2}}. \end{split}$$

Hence, (4.3) holds and then  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^q_{\beta-2}$ .

(ii) Let us first recall the duality theorem for  $A^1_{\alpha}$  [3], we see that  $A^1_{\alpha}$  can be identified as the dual of the little Bloch space under the pairing

$$\langle h, g \rangle = \int_{\mathbb{D}} h(z)\overline{g(z)}(1 - |z|^2)^{\alpha} dA(z), \quad h \in \mathcal{B}_0, \quad g \in A^1_{\alpha}.$$
 (4.7)

we obtain that  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^1_{\beta-2}$  if and only if there exists a positive constant C such that

$$\left| \int_{[0,1)} \overline{f(t)} g(t) d\mu(t) \right| \le C \|f\|_{A^p_\alpha} \|g\|_{\mathcal{B}}, \quad f \in A^p_\alpha, \ g \in \mathcal{B}_0.$$

$$(4.8)$$

Assume that  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^1_{\beta-2}$ . Take the families of text functions

$$f_a(z) = \left(\frac{1 - a^2}{(1 - az)^2}\right)^{(\alpha + 2)/p}$$
 and  $g_a(z) = \log \frac{2}{1 - az}$ ,  $0 < a < 1$ .

A calculation shows that  $\{f_a\} \subset A^p_\alpha$ ,  $\{g_a\} \subset \mathcal{B}_0$  and

$$\sup_{a \in [0,1)} ||f||_{A^p_\alpha} < \infty \quad and \quad \sup_{a \in [0,1)} ||g||_{\mathcal{B}} < \infty.$$

It follows that

$$\infty > C \sup_{a \in [0,1)} ||f||_{A_{\alpha}^{p}} \sup_{a \in [0,1)} ||g||_{\mathcal{B}} 
\ge \left| \int_{[0,1)} f_{a}(t) g_{a}(t) d\mu(t) \right| 
\ge \int_{[a,1)} \left( \frac{1 - a^{2}}{(1 - at)^{2}} \right)^{(\alpha + 2)/p} \log \frac{2}{1 - at} d\mu(t) 
\ge \frac{\log \frac{2}{1 - a}}{(1 - a)^{(\alpha + 2)/p}} \mu([a, 1)).$$
(4.9)

This is equivalent to saying that  $\mu$  is a 1-logarithmic  $(\frac{\alpha+2}{p})$ -Carleson measure. Conversely, we assume that  $\mu$  is a 1-logarithmic  $(\frac{\alpha+2}{p})$ -Carleson measure. It is well known that any function  $g \in \mathcal{B}$  [16] has the property

$$|g(z)| \le C||g||_{\mathcal{B}} \log \frac{2}{1 - |z|}.$$

Thus, if  $0 , for every <math>f \in A^p_\alpha$  and  $g \in \mathcal{B}_0 \subset \mathcal{B}$ , we have

$$\left| \int_{[0,1)} \overline{f(t)} g(t) d\mu(t) \right| \le C \|g\|_{\mathcal{B}} \int_{[0,1)} |f(t)| \log \frac{2}{1-t} d\mu(t).$$

Using [5, Proposition 2.5], we obtain that  $\log \frac{2}{1-t}d\mu(t)$  is a  $(\frac{\alpha+2}{p})$ -Carleson measure. Hence, Lemma 3.2 implies that

$$\left| \int_{[0,1)} \overline{f(t)} g(t) d\mu(t) \right| \le C \|f\|_{\alpha}^{p} \|g\|_{\mathcal{B}} \quad f \in A_{\alpha}^{p}, \ g \in \mathcal{B}_{0}.$$
 (4.10)

Therefore,  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^1_{\beta-2}$ .

**Corollary 4.1** Suppose that  $1 \le p < \infty$ ,  $\alpha > -1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If p > 1, then  $\mathcal{H}_{\mu,\alpha+2}$  is a bounded operator on  $A^p_{\alpha}$  if and only if  $\mu$  is a  $(\alpha + 2)$ -Carleson measure.
- (ii) If p = 1, then  $\mathcal{H}_{\mu,\alpha+2}$  is a bounded operator on  $A^1_{\alpha}$  if and only if  $\mu$  is a 1-logarithmic  $(\alpha + 2)$ -Carleson measure.

Now, it is well known that  $A^p_{\alpha} = D^p_{\alpha+p}$  (see [3, Theorem 4.28]). Hence, regarding Dirichlet spaces Theorem 4.2 says the following.

**Corollary 4.2** Suppose that  $0 , <math>\alpha > p-1$  and  $\beta > q+1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If  $q \geq p$  and q > 1, if  $\mathcal{H}_{\mu,\beta-q}$  is a bounded operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-2}$  then  $\mu$  is a  $\left(\frac{\alpha+2}{p} + \frac{\beta-q}{q'} 1\right)$ -Carleson measure. Conversely if  $\mu$  is a  $\left(\max\{\alpha + 2 p, \beta q\}(\frac{1}{p} + \frac{1}{q'})\right)$ -Carleson measure then  $\mathcal{H}_{\mu,\beta-q}$  is a bounded operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-2}$ .
- (ii) If  $q \ge p$  and q = 1, then  $\mathcal{H}_{\mu,\beta-1}$  is a bounded operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^1_{\beta-2}$  if and only if  $\mu$  is a 1-logarithmic  $(\frac{\alpha+2}{p}-1)$ -Carleson measure.

**Corollary 4.3** Suppose that  $1 \le p < \infty$ ,  $\alpha > p - 1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If p > 1, then  $\mathcal{H}_{\mu,\alpha+2-p}$  is a bounded operator on  $\mathcal{D}^p_{\alpha}$  if and only if  $\mu$  is a  $(\alpha + 2 p)$ Carleson measure.
- (ii) If p = 1, then  $\mathcal{H}_{\mu,\alpha+2-p}$  is a bounded operator on  $\mathcal{D}^1_{\alpha}$  if and only if  $\mu$  is a 1-logarithmic  $(\alpha + 1)$ -Carleson measure.

Then we discuss those measures  $\mu$  for which  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-1}$ .

**Theorem 4.3** Suppose that q > 1 and  $0 , <math>\alpha > p - 1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1. Then  $\mathcal{H}_{\mu,\alpha+1-p}$  is a bounded operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\alpha-p}$  if and only if  $\mu$  is a  $\left(\frac{\alpha+2-p}{p} + \frac{\alpha+2-p}{q'}\right)$ -Carleson measure.

The Theorem 4.3 is also closely related to Theorem 4.4, thus it suffices to prove Theorem 4.4

**Theorem 4.4** Suppose that q > 1 and  $0 , <math>\alpha > p-1$  and  $\beta > q$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-1}$  then  $\mu$  is a  $\left(\frac{\alpha+2}{p} + \frac{\beta+1}{q'} 1\right)$ -Carleson measure.
- (ii) If  $\mu$  is a  $\left(\max\{\alpha+2-p,\beta+1\}\left(\frac{1}{p}+\frac{1}{q'}\right)\right)$ -Carleson measure then  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-1}$ .

**Proof** By using Fubini's theorem and reproducing property imply that

$$\int_{\mathbb{D}} \overline{\mathcal{H}_{\mu,\beta}(f)'(z)} g'(z) (1 - |z|^{2})^{\beta - 1} dA(z) 
= \int_{\mathbb{D}} \int_{[0,1)} t\beta \frac{\overline{f(t)} d\mu(t)}{(1 - t\overline{z})^{\beta + 1}} g'(z) (1 - |z|^{2})^{\beta - 1} dA(z) 
= \int_{[0,1)} \int_{\mathbb{D}} t\beta \frac{(1 - |z|^{2})^{\beta - 1} g'(z)}{(1 - t\overline{z})^{\beta + 1}} dA(z) \overline{f(t)} d\mu(t) 
= \frac{\beta}{\beta - 1} \int_{[0,1)} t\overline{f(t)} g'(t) d\mu(t), \quad f \in \mathcal{D}_{\alpha}^{p}, \ g' \in A_{\beta - 1}^{1}.$$
(4.11)

(i) First consider the case  $0 , <math>q \ge 1$  and  $q \ge p$ . Using (4.11) and the duality theorem for Lemma 4.1, take  $(\mathcal{D}_{\beta-1}^q)^* \cong \mathcal{D}_{\beta-1}^{q'}$  and  $(\mathcal{D}_{\beta-1}^{q'})^* \cong \mathcal{D}_{\beta-1}^q$  (q > 1). we obtain that  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $\mathcal{D}_{\alpha}^p$  into  $\mathcal{D}_{\beta-1}^q$  if and only if there exists a positive constant C such that

$$\left| \int_{[0,1)} t \overline{f(t)} g'(t) d\mu(t) \right| \le C \|f\|_{\mathcal{D}^{p}_{\alpha}} \|g\|_{\mathcal{D}^{q'}_{\beta-1}}, \quad f \in \mathcal{D}^{p}_{\alpha}, \ g' \in A^{q'}_{\beta-1}.$$
 (4.12)

Assume that  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-1}$ . Take the families of text functions

$$f_a(z) = \frac{1}{a} \frac{(1-a^2)^{(\alpha+2)/p}}{(1-az)^{2(\alpha+2)/p-1}}$$
 and  $g'_a(z) = \left(\frac{1-a^2}{(1-az)^2}\right)^{(\beta+1)/q'}$ ,  $0 < a < 1$ .

A calculation shows that  $\{f_a\} \subset \mathcal{D}^p_{\alpha}, \{g_a\} \subset \mathcal{D}^{q'}_{\beta-1}$  and

$$\sup_{a \in [0,1)} ||f||_{\mathcal{D}^p_\alpha} < \infty \quad and \quad \sup_{a \in [0,1)} ||g||_{\mathcal{D}^{q'}_{\beta-1}} < \infty.$$

It follows that

$$\infty > C \sup_{a \in [0,1)} ||f||_{\mathcal{D}_{\alpha}^{p}} \sup_{a \in [0,1)} ||g||_{\mathcal{D}_{\beta-1}^{q'}} 
\ge \left| \int_{[0,1)} t f_{a}(t) g'_{a}(t) d\mu(t) \right| 
\ge \int_{[a,1)} \frac{t}{a} \frac{(1-a^{2})^{(\alpha+2)/p}}{(1-at)^{2(\alpha+2)/p-1}} \left( \frac{1-a^{2}}{(1-at)^{2}} \right)^{(\beta+1)/q'} d\mu(t) 
\ge \frac{1}{(1-a^{2})^{(\alpha+2)/p+(\beta+1)/q'-1}} \mu([a,1)).$$
(4.13)

This is equivalent to saying that  $\mu$  is a  $\left(\frac{\alpha+2}{p} + \frac{\beta+1}{q'} - 1\right)$ -Carleson measure.

(ii) On the other hand, if  $\mu$  is a  $\left(\max\{\alpha+2-p,\beta+1\}(\frac{1}{p}+\frac{1}{q'})\right)$ -Carleson measure. Let s=1+p/q'. Then s'=1+q'/p. We find  $A^p_{\alpha}=\mathcal{D}^p_{\alpha+p}$  which means  $f\in A^p_{\alpha-p}$ . By Lemma 3.2, we obtain that there exists a constant C>0 such that

$$\left( \int_{[0,1)} |f(t)|^s d\mu(t) \right)^{1/s} \le C \|f\|_{A^p_{(\alpha-p)}} = C \|f\|_{\mathcal{D}^p_{\alpha}}, \quad \text{for all } f \in \mathcal{D}^p_{\alpha}, \tag{4.14}$$

and

$$\left(\int_{[0,1)} |g'(t)|^{s'} d\mu(t)\right)^{1/s'} \le C\|g'\|_{A_{\beta-1}^{q'}} \le C\|g\|_{\mathcal{D}_{\beta-1}^{q'}}, \quad \text{for all } g' \in A_{\beta-1}^{q'}. \tag{4.15}$$

Then by Hölder's inequality,

$$\int_{[0,1)} |f(t)| |g'(t)| d\mu(t) \le \left( \int_{[0,1)} |f(t)|^s d\mu(t) \right)^{1/s} \left( \int_{[0,1)} |g'(t)|^{s'} d\mu(t) \right)^{1/s'}$$

$$\le C \|f\|_{\mathcal{D}^p_\alpha} \|g\|_{\mathcal{D}^{q'}_\alpha}.$$

Hence, (4.12) holds and then  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $\mathcal{D}_{\alpha}^{p}$  into  $\mathcal{D}_{\beta-1}^{q}$ .

Corollary 4.4 Suppose that  $1 , <math>\alpha > p - 1$ . Let  $\mu$  be a positive Borel measure on [0, 1) which satisfies the condition in Theorem 3.1. Then  $\mathcal{H}_{\mu,\alpha+1-p}$  is a bounded operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^p_{\alpha-p}$  if and only if  $\mu$  is a  $(\alpha+2-p)$ -Carleson measure.

**Corollary 4.5** Suppose that q > 1 and  $0 , <math>\alpha > -1$  and  $\beta > 0$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If  $\mathcal{H}_{\mu,\beta+q}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^q_{\beta-1}$  then  $\mu$  is a  $\left(\frac{\alpha+2}{p} + \frac{\beta+1+q}{q'}\right)$ -Carleson measure.
- (ii) If  $\mu$  is a  $\left(\max\{\alpha+2,\beta+1+q\}(\frac{1}{p}+\frac{1}{q'})\right)$ -Carleson measure then  $\mathcal{H}_{\mu,\beta+q}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^q_{\beta-1}$ .

Corollary 4.6 Suppose that  $1 , <math>\alpha > p - 1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1. Then  $\mathcal{H}_{\mu,\alpha+1}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^p_{\alpha-p}$  if and only if  $\mu$  is a  $(\alpha + 2)$ -Carleson measure.

# 5 Compactness of $\mathcal{H}_{\mu,\beta}$ acting on Weighted Bergman spaces and Dirichlet spaces

In this section we characterize the compactness of the Generalized Hilbert  $\mathcal{H}_{\mu,\beta}$ . We begin with the following Lemma 5.1 which is useful to deal with the compactness.

**Lemma 5.1** For  $0 and <math>0 < q < \infty$ ,  $\alpha > -1$  and  $\beta > 1$ . Suppose that  $\mathcal{H}_{\mu,\beta}$  is a bounded operator from  $A^p_{\alpha}$  into  $A^q_{\beta-2}$ . Then  $\mathcal{H}_{\mu,\beta}$  is a compact operator if and only if for any bounded sequence  $\{f_n\}$  in  $A^p_{\alpha}$  which converges uniformly to 0 on every compact subset of  $\mathbb{D}$ , we have  $\mathcal{H}_{\mu,\beta}(f_n) \to 0$  in  $A^q_{\beta-2}$ , for Dirichlet spaces also have an analogous conclusion.

The proof is similar to that of [5, Proposition 3.11]. We omit the details.

**Theorem 5.1** Suppose that  $0 , <math>\alpha > -1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

(i) If  $q \ge p$  and q > 1, then  $\mathcal{H}_{\mu,\alpha+2}$  is a compact operator from  $A^p_{\alpha}$  into  $A^q_{\alpha}$  if and only if  $\mu$  is a vanishing  $\left(\frac{\alpha+2}{p} + \frac{\alpha+2}{q'}\right)$ -Carleson measure.

(ii) If  $q \ge p$  and q = 1, then  $\mathcal{H}_{\mu,\alpha+2}$  is a compact operator from  $A^p_{\alpha}$  into  $A^1_{\alpha}$  if and only if  $\mu$  is a vanishing 1-logarithmic  $(\frac{\alpha+2}{p})$ -Carleson measure.

The Theorem 5.1 is closely related to Theorem 5.2, thus it suffices to prove Theorem 5.2

**Theorem 5.2** Suppose that  $0 , <math>\alpha > -1$  and  $\beta > 1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If  $q \ge p$  and q > 1, if  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $A^p_{\alpha}$  into  $A^q_{\beta-2}$  then  $\mu$  is a vanishing  $\left(\frac{\alpha+2}{p} + \frac{\beta}{q'}\right)$ -Carleson measure, conversely if  $\mu$  is a vanishing  $\left(\max\{\alpha+2,\beta\}(\frac{1}{p} + \frac{1}{q'})\right)$ -Carleson measure then  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $A^p_{\alpha}$  into  $A^q_{\beta-2}$ .
- (ii) If  $q \ge p$  and q = 1, then  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $A^p_{\alpha}$  into  $A^1_{\beta-2}$  if and only if  $\mu$  is a vanishing 1-logarithmic  $(\frac{\alpha+2}{p})$ -Carleson measure.

**Proof** (i) First consider  $q \geq p$  and q > 1. Suppose that  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $A^p_{\alpha}$  into  $A^q_{\beta-2}$ . Let  $\{a_n\} \subset (0,1)$  be any sequence with  $a_n \to 1$ . We set

$$f_{a_n}(z) = \left(\frac{1 - a_n^2}{(1 - a_n z)^2}\right)^{(\alpha + 2)/p}, \quad z \in \mathbb{D}.$$

Then  $f_{a_n}(z) \in A^p_{\alpha}$ ,  $\sup_{n\geq 1} ||f_{a_n}||_{H^p} < \infty$  and  $f_{a_n} \to 0$ , uniformly on any compact subset of  $\mathbb{D}$ . Using Lemma 5.1 and bearing in mind that  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $A^p_{\alpha}$  into  $A^q_{\beta-2}$ , we obtain that  $\{\mathcal{H}_{\mu,\beta}(f_{a_n})\}$  converges to 0 in  $A^q_{\beta-2}$ . This and (4.2) imply that

$$\lim_{n \to \infty} \int_{[0,1)} \overline{f_{a_n}(t)} g(t) d\mu(t)$$

$$= \lim_{n \to \infty} \int_{\mathbb{D}} \overline{\mathcal{H}_{\mu,\beta}(f_{a_n})(z)} g(z) d(1 - |z|^2)^{\beta - 2} A(z) = 0, \quad g \in A_{\beta - 2}^{q'}.$$
(5.1)

Now we set

$$g_{a_n}(z) = \left(\frac{1 - a_n^2}{(1 - a_n z)^2}\right)^{(\beta)/q'}, \quad z \in \mathbb{D}.$$

It is easy to check that  $g \in A_{\beta-2}^{q'}$ .

For  $r \in (a_n, 1)$ , we deduce that

$$\int_{[0,1)} \overline{f_{a_n}(t)} g_{a_n}(t) d\mu(t) 
\geq \int_{a_n}^1 \left( \frac{1 - a_n^2}{(1 - a_n t)^2} \right)^{(\alpha + 2)/p} \left( \frac{1 - a_n^2}{(1 - a_n t)^2} \right)^{\beta/q'} d\mu(t) 
\geq \frac{C}{(1 - a_n)^{(\alpha + 2)/p + \beta/q'}} \mu([a_n, 1)).$$

Then (5.1) and the fact that  $\{a_n\} \subset (0,1)$  is a sequence with  $a_n \to 1$  imply that

$$\lim_{a \to 1^{-}} \frac{1}{(1-a)^{(\alpha+2)/p+\beta/q'}} \mu([a,1)) = 0,$$

which shows that  $\mu$  is a vanishing  $\left(\frac{\alpha+2}{p}+\frac{\beta}{q'}\right)$ -Carleson measure.

On the other hand, suppose now that  $\mu$  is a vanishing  $\left(\max\{\alpha+2,\beta\}(\frac{1}{p}+\frac{1}{q'})\right)$ -Carleson measure. Let  $\{f_n\}_{n=1}^{\infty}$  be a sequence of  $A_{\alpha}^p$  functions with  $\sup_{n\geq 1}\|f_n\|_{A_{\alpha}^p}<\infty$  and such that  $\{f_n\}\to 0$ , uniformly on  $\mathbb D$ . Then by Lemma 5.1, it is enough to prove that  $\mathcal H_{\mu,\beta}(f_n)\to 0$  in  $A_{\beta-2}^q$ .

Taking  $g \in A_{\beta-2}^{q'}$  and  $r \in [0,1)$ , we have

$$\int_{[0,1)} |f_n(t)| \, |g(t)| d\mu(t) = \int_{[0,r)} |f_n(t)| \, |g(t)| d\mu(t) + \int_{[r,1)} |f_n(t)| \, |g(t)| d\mu(t).$$

For the first term on the right side tends to 0 since  $\{f_n\} \to 0$  uniformly on compact subsets of  $\mathbb{D}$ . For the second term, arguing as in proof of the boundedness (Theorem 4.2 (i)), we obtain that

$$\int_{[r,1)} |f_n(t)| |g(t)| d\mu(t) \leq \left( \int_{[0,1)} |f_n(t)|^s d\mu_r(t) \right)^{1/s} \left( \int_{[0,1)} |g(t)|^{s'} d\mu_r(t) \right)^{1/s'} \\
\leq (\mathcal{N}(\mu_r))^2 \|f\|_{A^p_{\alpha-2}} \|g\|_{A^{q'}_{\beta-2}}.$$

It tends to 0 by (2.1) and then

$$\lim_{n \to \infty} \int_{[0,1)} |f_n(t)| |g(t)| \, d\mu(t) = 0, \quad \text{ for all } g \in A_{\beta-2}^{q'}.$$

Thus

$$\lim_{n\to\infty} \left| \int_{\mathbb{D}} \overline{\mathcal{H}_{\mu,\beta}(f_n)(z)} g(z) \left(1 - |z|^{\beta-2}\right) dA(z) \right| = 0, \quad \text{ for all } g \in A_{\beta-2}^{q'}.$$

It means  $\mathcal{H}_{\mu,\beta}(f_n) \to 0$  in  $A^q_{\beta-2}$ , by Lemma 5.1, we obtain  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $A^p_{\alpha}$  into  $A^q_{\beta-2}$ .

(ii) Let  $0 . Suppose that <math>\mathcal{H}_{\mu,\beta}$  is a compact operator from  $A^p_{\alpha}$  into  $A^1_{\beta-2}$ . Let  $\{a_n\} \subset (0,1)$  be any sequence with  $a_n \to 1$  and  $f_{a_n}$  defines like in (i). Lemma 5.1 implies that  $\{\mathcal{H}_{\mu,\beta}(f_{a_n})\}$  converges to 0 in  $A^1_{\beta-1}$ . Then we have

$$\lim_{n \to \infty} \int_{[0,1)} \overline{f_{a_n}(t)} g(t) d\mu(t)$$

$$= \lim_{n \to \infty} \int_{\mathbb{D}} \overline{\mathcal{H}_{\mu,\beta}(f_{a_n})(z)} g(z) d(1 - |z|^2)^{\beta - 2} A(z) = 0, \quad g \in \mathcal{B}_0.$$
(5.2)

Now we set

$$g(z) = \log \frac{2}{1 - z}.$$

It is well know that  $g \in \mathcal{B}_0$ . For  $r \in (a_n, 1)$ , we deduce that

$$\int_{[0,1)} \overline{f_{a_n}(t)} g(t) d\mu(t) 
\geq C \int_{[a_n,1)} \left( \frac{1-a_n^2}{(1-a_nt)^2} \right)^{(\alpha+2)/p} \log \frac{2}{1-t} d\mu(t) 
\geq \frac{C \log \frac{2}{1-a_n}}{(1-a_n)^{(\alpha+2)/p}} \mu([a_n,1)).$$

Letting  $a_n \to 1^-$  as  $n \to \infty$ , we have

$$\lim_{a \to 1^{-}} \frac{\log \frac{2}{1 - a_n}}{(1 - a_n^2)^{(\alpha + 2)/p}} \mu([a_n, 1)) = 0.$$

We can obtain that  $\mu$  is a vanishing 1-logarithmic  $(\frac{\alpha+2}{p})$ -Carleson measure.

On the other hand, suppose that  $\mu$  is a vanishing 1-logarithmic  $(\frac{\alpha+2}{p})$ -Carleson measure. Let  $d\nu(t) = \log \frac{2}{1-t} d\mu(t)$ . By Proposition 2.5 of [9], we know that  $\nu$  is a vanishing  $(\frac{\alpha+2}{p})$ -Carleson. Let  $\{f_n\}_{n=1}^{\infty}$  be a sequence of  $A_{\alpha}^p$  functions with  $\sup_{n\geq 1}||f_n||_{A_{\alpha}^p}<\infty$ , and such that  $\{f_n\} \to 0$ , uniformly on any compact subset of  $\mathbb{D}$ . Then by Lemma 5.1, it is enough to prove that  $\{\mathcal{H}_{\mu,\beta}(f_n)\}\to 0$  in  $A^1_{\beta-2}$ . Then for every  $g\in\mathcal{B}_0,\ 0< r<1$ , using (2.1), we deduce that

$$\int_{[0,1)} |f_n(t)| |g(t)| d\mu(t) = \int_{[0,r)} |f_n(t)| |g(t)| d\mu(t) + \int_{[r,1)} |f_n(t)| |g(t)| d\mu(t) 
\leq \int_{[0,r)} |f_n(t)| |g(t)| d\mu(t) + C ||g||_{\mathcal{B}} \int_{[r,1)} |f_n(t)| \log \frac{2}{1-t} d\mu(t) 
\leq \int_{[0,r)} |f_n(t)| |g(t)| d\mu(t) + C ||g||_{\mathcal{B}} \int_{[0,1)} |f_n(t)| d\nu_r(t) 
\leq \int_{[0,r)} |f_n(t)| |g(t)| d\mu(t) + C \mathcal{N}(\nu_r) \sup_{n \geq 1} ||f_n||_{A^p} ||g||_{\mathcal{B}}.$$
(5.3)

Hence, (2.1), the conditions  $\{f_n\} \to 0$ , uniformly on compact subsets of  $\mathbb{D}$ , and  $\sup_{n \geq 1} \|f_n\|_{A^p}$ imply that

$$\lim_{n \to \infty} \int_{[0,1)} |f_n(t)| |g(t)| \, d\mu(t) = 0, \quad \text{ for all } g \in \mathcal{B}_0.$$

Using this and (5.2), we have

$$\lim_{n\to\infty} \left| \int_{\mathbb{D}} \overline{\mathcal{H}_{\mu,\beta}(f_n)(z)} g(z) (1-|z|^2)^{\beta-2} dA(z) \right| = 0, \quad \text{ for all } g \in \mathcal{B}_0.$$

It means  $\mathcal{H}_{\mu,\beta}(f_n) \to 0$  in  $A^1_{\beta-2}$ , by Lemma 5.1 we obtain  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $A^p_{\alpha}$ into  $A^1_{\beta-2}$ .

Corollary 5.1 Suppose that  $1 \le p < \infty$ ,  $\alpha > -1$ . Let  $\mu$  be a positive Borel measure on [0,1)which satisfies the condition in Theorem 3.1.

- (i) If p > 1, then  $\mathcal{H}_{\mu,\alpha+2}$  is a compact operator on  $A^p_{\alpha}$  if and only if  $\mu$  is a vanishing  $(\alpha+2)$ -
- (ii) If p=1, then  $\mathcal{H}_{\mu,\alpha+2}$  is a compact operator on  $A^1_{\alpha}$  if and only if  $\mu$  is a vanishing 1logarithmic ( $\alpha + 2$ )-Carleson measure.

It is well known that  $A_{\alpha}^{p} = D_{\alpha+p}^{p}$  (see [3, Theorem 4.28]). Hence, we can immediately obtain the following.

Corollary 5.2 Suppose that  $0 , <math>\alpha > p-1$  and  $\beta > q+1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If  $q \geq p$  and q > 1, if  $\mathcal{H}_{\mu,\beta-q}$  is a compact operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-2}$  then  $\mu$  is a vanishing  $\left(\frac{\alpha+2}{p} + \frac{\beta-q}{q'} 1\right)$ -Carleson measure. Conversely, if  $\mu$  is a vanishing  $\left(\max\{\alpha+2-p,\beta-q\}(\frac{1}{p}+\frac{1}{q'})\right)$ -Carleson measure then  $\mathcal{H}_{\mu,\beta-q}$  is a compact operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-2}$ .
- (ii) If  $q \geq p$  and q = 1, then  $\mathcal{H}_{\mu,\beta-1}$  is a compact operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^1_{\beta-2}$  if and only if  $\mu$  is a vanishing 1-logarithmic  $(\frac{\alpha+2}{p}-1)$ -Carleson measure.

**Corollary 5.3** Suppose that  $1 \le p < \infty$ ,  $\alpha > p - 1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If p > 1, then  $\mathcal{H}_{\mu,\alpha+2-p}$  is a compact operator on  $\mathcal{D}^p_{\alpha}$  if and only if  $\mu$  is a vanishing  $(\alpha+2-p)$ -Carleson measure.
- (ii) If p = 1, then  $\mathcal{H}_{\mu,\alpha+2-p}$  is a compact operator on  $\mathcal{D}^1_{\alpha}$  if and only if  $\mu$  is a vanishing 1-logarithmic  $(\alpha + 1)$ -Carleson measure.

Then we discuss those measures  $\mu$  for which  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $\mathcal{D}_{\alpha}^{p}$  into  $\mathcal{D}_{\beta-1}^{q}$ .

**Theorem 5.3** Suppose that q > 1 and  $0 , <math>\alpha > p - 1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1. Then  $\mathcal{H}_{\mu,\alpha+1-p}$  is a compact operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\alpha-p}$  if and only if  $\mu$  is a vanishing  $\left(\frac{\alpha+2-p}{p} + \frac{\alpha+2-p}{q'}\right)$ -Carleson measure.

The Theorem 5.3 is also closely related to Theorem 5.4, thus it suffices to prove Theorem 5.4

**Theorem 5.4** Suppose that q > 1 and  $0 , <math>\alpha > p-1$  and  $\beta > q$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-1}$  then  $\mu$  is a vanishing  $\left(\frac{\alpha+2}{p} + \frac{\beta+1}{q'} 1\right)$ -Carleson measure.
- (ii) If  $\mu$  is a vanishing  $\left(\max\{\alpha+2-p,\beta+1\}\left(\frac{1}{p}+\frac{1}{q'}\right)\right)$ -Carleson measure then  $\mathcal{H}_{\mu,\beta}$  is a compact operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^q_{\beta-1}$ .

**Proof** The proof is the same as that of Theorem 4.4(i) and Theorem 5.2(i). We omit the details here.

Corollary 5.4 Suppose that  $1 , <math>\alpha > p-1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1. Then  $\mathcal{H}_{\mu,\alpha+1-p}$  is a compact operator from  $\mathcal{D}^p_{\alpha}$  into  $\mathcal{D}^p_{\alpha-p}$  if and only if  $\mu$  is a vanishing  $(\alpha+2-p)$ -Carleson measure.

**Corollary 5.5** Suppose that q > 1 and  $0 , <math>\alpha > -1$  and  $\beta > 0$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1.

- (i) If  $\mathcal{H}_{\mu,\beta+q}$  is a compact operator from  $A^p_{\alpha}$  into  $A^q_{\beta-1}$  then  $\mu$  is a vanishing  $\left(\frac{\alpha+2}{p} + \frac{\beta+1+q}{q'}\right)$ -Carleson measure.
- (ii) If  $\mu$  is a vanishing  $\left(\max\{\alpha+2,\beta+1+q\}\left(\frac{1}{p}+\frac{1}{q'}\right)\right)$ -Carleson measure then  $\mathcal{H}_{\mu,\beta+q}$  is a compact operator from  $A^p_{\alpha}$  into  $A^q_{\beta-1}$ .

Corollary 5.6 Suppose that  $1 , <math>\alpha > p - 1$ . Let  $\mu$  be a positive Borel measure on [0,1) which satisfies the condition in Theorem 3.1. Then  $\mathcal{H}_{\mu,\alpha+1}$  is a compact operator from  $A^p_{\alpha}$  into  $A^p_{\alpha-p}$  if and only if  $\mu$  is a vanishing  $(\alpha+2)$ -Carleson measure.

### References

- [1] Bao G, Wulan H. Hankel matrices acting on Dirichlet spaces, J Math Anal Appl, 2014 **409**(1): 228-235.
- [2] Duren P L, Schuster A. Bergman spaces. American Mathematical Soc, 2004.
- [3] Zhu K. Operator theory in function spaces, American Mathematical Soc, 2007.
- [4] Chatzifountas C, Girela D, Peláez J Á. A generalized Hilbert matrix acting on Hardy spaces, J Math Anal Appl, 2014, **413**(1): 154-168.
- [5] Cowen Jr C C, MacCluer B I. Composition Operators on Spaces of Analytic Functions. CRC Press, 1995.
- [6] Diamantopoulos E , Siskakis A G. Composition operators and the Hilbert matrix, Studia Math, 2000, 140(2): 191-198.
- [7] Galanopoulos P , Girela D, Peláez, J Á , Siskakis A G. Generalized Hilbert operators, Ann Acad Sci Fenn Math, 2014, **39**: 231-258.
- [8] Galanopoulos P , Peláez J Á. A Hankel matrix acting on Hardy and Bergman spaces, Studia Math ,2010, **200**(3): 201-220.
- [9] Girela D, Merchán N. A generalized Hilbert operator acting on conformally invariant spaces, Banach J Math Anal ,2018, **12**(2): 374-398.
- [10] Girela D, Merchán N. Hankel matrices acting on the Hardy space  $H^1$  and on Dirichlet spaces, Revista Matemática Complutense, 2019, 32(3): 799-822.
- [11] Gnuschke-Hauschild D, Pommerenke C. On Bloch functions and gap series, J Reine Angew Math, 1986, 367: 172-186.
- [12] Jevtić M, Vukotić D, Arsenović M. Taylor coefficients and coefficient multipliers of Hardy and Bergman-type spaces, Springer International Publishing, 2016.
- [13] Hastings W W. A Carleson measure theorem for Bergman spaces, Proc Amer Math Soc, 1975, **52**(1): 237-241.
- [14] Luecking D H, Forward and reverse Carleson inequalities for functions in Bergman spaces and their derivatives, American Journal of Mathematics, 1985, **107**(1): 85-111.
- [15] MacCluer B, Zhao R. Vanishing logarithmic Carleson measures, Illinois J Math , 2002, 46(2):507-518.
- [16] Pommerenke C, Clunie J, Anderson J. On Bloch functions and normal functions, J Reine Angew Math, 1974, 270: 12-37.
- [17] Romberg B W, Duren P L, Shields A L. Linear functionals on Hp spaces with 0 , Reine Angew Math, 1969,**238**: 32-60.
- [18] Ye S, Zhou Z. A derivative-Hilbert operator acting on the Bloch space, Complex Anal Oper Theory, 2021, **15**(5): 88.

- [19] Ye S, Zhou Z. A derivative-Hilbert operator acting on Bergman spaces, J Math Anal Appl, 2022, **506**(1): 125553.
- [20] Ye S, Zhou, Z. Generalized Hilbert Operator Acting on Bloch Type Spaces, Acta Mathematic Sinica, Chinese Series (In Press), 2022.
- [21] Zhao R. On logarithmic Carleson measures, Acta Sci Math, 2003, 69: 605-618.
- [22] Zhu K. Bloch Type Spaces of Analytic Functions, Rocky Mountain J Math, 1993, **23**(3): 1143-1177.