

The emergence of interstellar molecular complexity explained by interacting networks, PNAS 119 (30) e2119734119 (2022)

<https://www.pnas.org/doi/10.1073/pnas.2119734119>

Miguel García-Sánchez,^{1,2,3} Izaskun Jiménez-Serra,¹ Fernando Puente-Sánchez,⁴ and Jacobo Aguirre^{1,3,*}

¹*Centro de Astrobiología (CSIC-INTA), ctra. de Ajalvir km 4, 28850 Torrejón de Ardoz, Madrid, Spain*

²*Instituto de Investigación Tecnológica (IIT), Universidad Pontificia Comillas, 28015 Madrid, Spain*

³*Grupo Interdisciplinar de Sistemas Complejos (GISC), Madrid, Spain*

⁴*Department of Aquatic Sciences and Assessment,
Swedish University of Agricultural Sciences, 76551 Uppsala, Sweden*

Abstract: Recent years have witnessed the detection of an increasing number of complex organic molecules in interstellar space, some of them being of prebiotic interest. Disentangling the origin of interstellar prebiotic chemistry and its connection to biochemistry and ultimately to biology is an enormously challenging scientific goal where the application of complexity theory and network science has not been fully exploited. Encouraged by this idea, we present a theoretical and computational framework to model the evolution of simple networked structures toward complexity. In our environment, complex networks represent simplified chemical compounds, and interact optimizing the dynamical importance of their nodes. We describe the emergence of a transition from simple networks toward complexity when the parameter representing the environment reaches a critical value. Notably, although our system does not attempt to model the rules of real chemistry, nor is dependent on external input data, the results describe the emergence of complexity in the evolution of chemical diversity in the interstellar medium. Furthermore, they reveal an as yet unknown relationship between the abundances of molecules in dark clouds and the potential number of chemical reactions that yield them as products, supporting the ability of the conceptual framework presented here to shed light on real scenarios. Our work reinforces the notion that some of the properties that condition the extremely complex journey from the chemistry in space to prebiotic chemistry and finally to life could show relatively simple and universal patterns.

Significance statement: *The road to life is punctuated by transitions toward complexity, from astrochemistry to biomolecules and eventually to living organisms. Disentangling the origin of such transitions is a challenge where the application of complexity and network theory has not been fully exploited. We introduce a computational framework in which simple networks simulate the most basic building bricks of life and interact to form complex structures, leading to an explosion of diversity when the parameter representing the environment reaches a critical value. While this model is abstract and unrelated to chemical theory, its predictions reliably mimic the molecular evolution in the interstellar medium during the transition toward chemical complexity, suggesting that the rules leading to the emergence of complexity may be universal.*

The origin of life on Earth is far away from being unveiled. Life could have appeared spontaneously in our early planet about 4 billion years ago, or exist previously in the outer space and be brought by dust and meteoroids, as panspermia states. An intermediate hypothesis is the *molecular panspermia*, which proposes that the original building blocks of life could have been produced in the interstellar medium (ISM) and be introduced in the early Earth by asteroids and meteorites during the Late Heavy Bombardment that took place between 3.8 and 4.1 billion years ago, importantly enriching prebiotic chemistry. Over 200 molecules have been detected in the ISM, with some of them being prebiotically relevant as e.g. glycolaldehyde, urea or ethanalamine [1–4]. Prebiotic molecules such as glycine or ribose have indeed been found in meteorites and comets [5, 6], which supports the idea that prebiotic species could initially form in inter-

stellar space, and be transferred later on to planetesimals and to Earth during the formation of the Solar system.

In parallel to the efforts to understand the origin of life from the biochemical optics, the development and advance of computation in the last fifty years propitiated the study of life modeled as a cellular automaton [7, 8]. Despite the simplicity of the rules, the different systems under study, in particular Conway’s Game of Life [9], presented an unexpected variety of spatio-temporal patterns and cast light on how complexity, emergence, and self-organization arise from a simple system. More complex systems were introduced in the 80’s to model Darwinian evolution with the use of a new type of artificial life where organisms described as computer programs could self-replicate, adapt and mutate by natural selection, mostly competing for the control of the memory of the computer (e.g. CoreWar [10] and Avida [11, 12]). The introduction of these digital organisms to address fundamental biological questions was supported on two main statements. First, they provide a way to generalize life beyond the organisms detected so-far in our

*jaguirre@cab.inta-csic.es

biosphere. Second, they allow to perform, enlarge and repeat experiments on a scale that is unachievable with real entities [11].

In order to assess whether artificial life and its connection with complex networks theory can bring light to the study of the origin of life, in this work we present a computational framework, NetWorld, where networks interact following very simple local rules inspired in network science and game theory leading to a *chemistry of networks*. Our objective is to test using real astrochemical data whether a basic digital framework can reproduce certain general properties of the difficult transition from chemistry to biology, and therefore describe in an abstract level the creation of the basic building blocks of life.

I. RESULTS

In NetWorld every chemical compound is represented by a network, and this allows us to apply the strength and tools of complex network theory. From simply isolated nodes that simulate an initial state of total lack of complexity, this new artificial chemistry shows the emergence of a transition beyond which the environment permits the appearance of a rich variety of networks with different spectral, topological and dynamical properties, mimicking in a very simplified manner the first steps of prebiotic chemistry and its natural evolution toward complexity. We will pay special attention to the descriptive and predictive ability of this framework, showing that the results throw light on the chemical evolution toward complexity of molecular clouds in the interstellar medium and reveal a so-far-unknown relationship between the abundances of the molecules present in dark clouds and the number of chemical reactions that have them as products.

A. Description and rules of the model

An artificial chemistry model is defined by three components: a set of all possible structures, a set of rules that govern the interaction among structures and an algorithm that describes the reaction domain [13]. Furthermore, depending on their level of abstraction, models can be classified in analogous, when they try to be faithful to natural chemistry, and abstract, otherwise. According to these definitions, NetWorld could be understood as an extremely abstract artificial chemistry model, where nodes stand for indistinguishable basic entities –they are unweighted, and have no different properties to represent the chemical valency or size of atoms, for example–, and the bonds between nodes are represented by undirected and unweighted links. In a simple case these entities could be molecules where nodes stand for atoms and links for their interactions. However, as its internal rules are so different to those that govern real chemistry, our approach differs drastically from the large myriad

of artificial chemistry models that quantitatively represent real processes through detailed descriptions of the physico-chemical interaction between atoms [13–15].

Figure 1 presents a visual description of the evolutionary dynamics described by NetWorld, and Fig. 1(a) shows a toy example of 4 initial nodes. A rigorous explanation of the algorithm is presented in Supporting Information Texts S1 to S3, an analysis of the dependence of the computation time with NetWorld’s parameters is introduced in Supporting Information Text S4 and information on the public availability of the code is found in Supporting Information Text S5. Each process is started with an initial number $n(0) = N$ of isolated nodes. Nodes are neither created nor destroyed during the process. In each time step t , the population consists of $n(t)$ networks that will be made of the N available nodes for the total ensemble. At the beginning of each time step t , two networks A and B of the total population are chosen randomly. They interact following a simple set of rules explained below and in Supporting Information Text S1.A. They either form a new network C , simulating the reaction $A + B \rightarrow C$, or fail to join and remain as A and B . In the latter case, a different pair of networks A' and B' is chosen until one pair succeeds in forming a new network. At the end of the time step, every network i in the population has a partition probability

$$P_i = \frac{2}{1 + \exp(\mu_i \beta)} \quad (1)$$

of being divided into smaller pieces, simulating the reaction $C \rightarrow \sum D_j$ (see Supporting Information Text S1.B for a full description of the partition algorithm). The stability parameter μ_i is the second smallest eigenvalue associated with the Laplacian matrix of network i . It is also known as the algebraic connectivity or the Fiedler eigenvalue, and represents the resistance of a network to being split into different communities [16]. The environment parameter β concentrates the whole physico-chemical properties of the environment, such as the temperature or the radiation. β is constant during the whole process and it is the unique relevant global parameter of our model. When a new time step starts, the same process is repeated with the new collection of $n(t + 1)$ networks. Note that $n(t + 1) \geq n(t) - 1$ and $n(t) \in [1, N]$ for all t . The process finishes when (i) the totality of nodes collapse in a unique structure (i.e. $n(t) = 1$), (ii) $n(t) > 1$ but no networks will accept any new connections, or (iii) t reaches a limit value of 10^4 steps. A typical realization of the total process for an environment parameter $\beta = 2.5$ and $N = 20$ initial nodes is plotted in Fig. 1(c).

The interaction between the two networks A and B chosen randomly at time t to give rise to network C requires special attention (see Fig. 1(b)). It is inspired by a connecting method already applied to socio-economic networks [17, 18], and it is grounded in the extensive work devoted to the description of the competition/cooperation between networks developed during the last decade for ordinary interactions [19–22], or more




FIG. 1: Description of the evolutionary dynamics created by NetWorld. (a) Sketch of the potential evolution of a simple system formed by 4 nodes. (b) In every time step of a simulation, two networks get in touch and interact until the system reaches a final network (defined by a Nash equilibrium) or a cycle of several alternating structures. The connector links (blue) join the original networks (red and green). (c) Example of a realization of the whole process with environment parameter $\beta = 2.5$ and $N = 20$ initial isolated nodes. The process reaches an end when a single network of size 20 is created.

recently for higher-order interactions [23–25]. We randomly choose one node a in A and one node b in B (connector nodes from now on) and connect them through an undirected link (connector link from now on). This new link is accepted only if both connector nodes a and b increase their *dynamical importance* in the network, measured as $I = \lambda_1 u_l$. λ_1 is the largest eigenvalue of the adjacency matrix of the new network formed by A , B and the connector link just added, and \vec{u} represents its associated eigenvector, L1-normalized such that $\sum u_k = 1$. u_l is the eigenvector centrality of connector node l , and measures the importance of a node based on how well connected it is and how important its neighbors are. Both λ_1 and \vec{u} are important measurements of the dynamical properties associated with a complex network. If there was a dynamical process of the type $\vec{m}(t+1) = \mathbf{G}\vec{m}(t)$ evolving on a network of adjacency matrix \mathbf{G} , it is known that $\vec{m}(t) \rightarrow \vec{u}$ when $t \rightarrow \infty$, and the population growth rate would be λ_1 [17]. The process of choosing randomly a new pair of nodes a' in A and b' in B –already connected through a connector link from a to b – and checking whether they accept a link or not is repeated, taking into account that, for simplicity and without loss of generality, only one connector link per node is allowed [18], and therefore any pre-existing connector links associated with nodes a' and b' are erased. In general, this algorithm leads to the connection of A and B through a cascade and rewiring of connector links until the total network

$C = A + B$ reaches a Nash equilibrium or a cycle between several final configurations, and in the latter case one of the final cyclic configurations is chosen randomly as C . If no links are accepted between A and B when all possible connections between both networks have been tried, we suppose that they do not react and A and B remain unchanged. In this case, the time is not increased and the interaction between two other networks starts.

In summary, our framework is grounded in the simulation of abstract networked entities that evolve following rules inherited from network science and game theory, and there is a total absence of machine learning techniques or any kind of fitting with real data in the different steps of the model.

B. Description of the artificial chemosphere created at NetWorld. Transition toward complexity

Every simulation of the whole process throughout this work started from $N = 40$ initial isolated nodes, lasted a maximum of 10^4 time steps and was repeated 25 times for each value of the environment parameter β in the range $\beta = [0, 2.7]$. The limit case $\beta \rightarrow \infty$ does not permit any network partition ($P = 0$ in Eq. 1) and is also studied. All structures detected at the end of any time step for all realizations of each β , even if they were destroyed later, represent its diversity. The relative abundance of




FIG. 2: Description of the diversity of structures created by NetWorld for different values of the environment parameter β and $N = 40$ initial nodes. (a) Histogram of network sizes (number of nodes in a network) for $\beta = 1.8$. A log-normal fitting is plotted in bold line (goodness of fit of $r = 0.993$). Dependence with β of (b) size of the networks, (c) average degree \bar{k} , (d) maximum eigenvalue λ_1 , (e) degree entropy H (calculated as $H = -\sum_{i=1}^K p_i \times \log_2(p_i)$, where p_i is the fraction of nodes of degree i , and K the maximum degree), and (f) stability parameter μ . (g) First (blue, 54% of variability) and second (yellow, 21% of variability) components of the PCA of the system. (h) Number of different configurations N_{conf} as a function of β . Exponential approximations are plotted in dashed lines ($N_{conf} = 2.56 \exp(1.43\beta)$, $r = 0.971$ for $\beta < \beta_{crit}$, and $N_{conf} = 0.03 \exp(4.25\beta)$, $r = 0.998$ for $\beta > \beta_{crit}$). The critical value of β ($\beta_{crit} \approx 1.55$) is remarked. In (b-f) every single network is represented by a blue dot, and the average value and the 10-90 percentile range for each β are plotted in red.

each configuration is given by the probability of finding it in the set of networks accumulated during all times and realizations (see Supporting Information Text S2 for a detailed explanation of how to compute the number of different configurations and the relative abundance in a simulation).

1. Diversity vs. environment

We start focusing on how the environment shapes the networks emerging in the system. Figure 2 shows the topological and structural description of the diversity created for a wide range of values of the environment parameter. For $\beta = 0$, every network of 2 nodes, i.e. the first to be created from isolated nodes, immediately breaks up and only isolated nodes are found for all times.

When β grows, the partition probability P of a network decreases (Eq. 1) and diverse structures emerge, but it is not until large values of β are reached ($\beta = 2.5$ for $N = 40$) that networks of the maximum possible size are created, as shown in Fig. 2(b). [50] The size, mean degree \bar{k} , largest eigenvalue of the adjacency matrix λ_1 and the degree entropy of the created structures H , plotted in Figs. 2(b-e), positively correlate with β , as shown in the first component of the Principal Components Analysis (PCA) calculated for the whole ensemble of existing structures and plotted in Fig. 2(g) [26]. On the contrary, the average stability of the networks $\bar{\mu}$ correlates negatively with β (Fig. 2(f) and second component of the PCA in (Fig. 2(g)), as more extensive and heterogeneous structures –i.e. with large entropy H – are in general easier to divide, but the low partition probability of networks for large β permits their appearance and survival.

Importantly, in Fig. 2(h) the growth of the number of different configurations with β suffers a transition around $\beta_{crit} = 1.55$ such that beyond that critical point the diversity explodes exponentially faster than before it. While the 10 – 90 percentile range plotted in red in Figs. 2(b-f) shows that the population is formed by a large number of networks with similar topological properties, for $\beta > \beta_{crit}$ there is also a *rare chemosphere* of structures that show very different topologies from the one of the majority, introducing a large amount of variability. Note that the existence of long tails of low-abundance entities is a typical property of real complex systems, notably ecological communities [27, 28]. In summary, for low values of β the main bricks of future complexity are formed, but only when a critical state of the environment is reached may these motifs be sufficiently abundant and interact successfully to enrich the system with a large number of new structures. This new population is made of very diverse entities, where (i) regular/robust structures but also (ii) heterogeneous/less stable networks can be detected. The former are made out of very similar blocks that, from the optics of information theory, would not code complex information but show redundancy, a fundamental property to prevent attacks and failures when basic tasks must be developed. The latter consist of complex networks where more information could be coded but are more sensitive to divisions, external perturbations, or losses of nodes.

2. Abundance vs. environment

Figure 3 describes the relative abundance of the networked structures obtained for the different environments, that is, the probability of finding them in the set of networks accumulated during all times and realizations. In Fig. 3(a) we plot the relative abundance of each configuration as a function of β , and it is clear that the simplest structures are created at low values of β and are especially abundant. A gradual appearance of larger and/or more complex configurations takes place for moderate β , and beyond $\beta = \beta_{crit}$ a cascade of new configurations leads to the exponential emergence of diversity already measured in Fig. 2(h). The sharpness of the transition toward complexity is specially visible in Fig. 3(b), where the normalized entropy of the total ensemble of networks H_{norm} [29] shows a transition from a linear growth with β to a constant value. For $\beta < \beta_{crit}$, the abundances of the few structures that exist tend to be more uniformly distributed when β grows (see Fig. 3(a)), increasing the entropy of the system (which reaches the maximum value $H_{norm} = 1$ when all abundances are equal, see the mathematical expression for H_{norm} in the caption of Fig. 3). However, once the critical environment is surpassed, the growth in complexity due to the tendency toward the relative abundance uniform distribution is balanced by the exponential emergence of diversity. Finally, in Fig. 3(c) we plot the abundance ranks of the population for differ-




FIG. 3: Analysis of the relative abundance of structures created by NetWorld for different values of the environment parameter β and $N = 40$ initial nodes. (a) Abundance of each structure as a function of β . Each color represents a different structure, but all structures emerged for $\beta > \beta_{crit}$ are plotted in dark blue for clarity. (b) Normalized entropy H_{norm} as a function of β . $H_{norm} = \sum_{i=1}^{N_{conf}} p_i \times \log_2 p_i / \log_2(N_{conf})$, where p_i is the relative abundance of configuration i and N_{conf} is the number of different configurations. Linear approximations are plotted in dashed lines: $H_{norm} = (0.006 \pm 0.004) + (0.121 \pm 0.005)\beta$ for $\beta < \beta_{crit}$, and $H_{norm} = (0.20 \pm 0.01) - (0.006 \pm 0.005)\beta$ for $\beta > \beta_{crit}$. The critical value of $\beta_{crit} \approx 1.55$ is remarked. (c) Abundance rank for different values of β (excluding isolated nodes). $\beta \in [0.1, 2.7]$, β grows from left to right in intervals of $\Delta\beta = 0.2$. The abundance rank for $\beta_{crit} = 1.55$ is plotted in brown line, and that for $\beta \rightarrow \infty$ (i.e., when the environment does not permit any network partition) is plotted in black circles.

ent values of β . For low β , the curves are very skewed and show exponential decays, typical behavior of ecological environments with little diversity [30]. When β grows and crosses the critical value (brown line), the curves gradually lose skewness and become power-laws, showing long tails consequence of a large diversity of rare configurations, and tend toward the limit case of the process in which $\beta = \infty$ and the structures cannot divide.

C. Application to a real scenario: Chemical complexity in the interstellar medium

Molecules are an important component of the interstellar medium since they regulate its ionization state and energy dissipation. Molecules are typically found




FIG. 4: Comparison between the results of the digital environment NetWorld (\times) and the evolution of chemical complexity of interstellar clouds, the astrochemical environment where the most basic bricks of life are created (\circ). All abundance sets shown in this figure were L1-normalized (the sum is 1). (a) Abundance rank of the molecular compounds detected in four interstellar clouds: the diffuse molecular cloud ζ Ophiuci [31], the translucent cloud located in the direction of the ultracompact HII region K4 in the SgrB2 molecular complex (a cloud in the transition from the diffuse regime to the dense regime, see [32, 33]), and the dense clouds L134N (Serpens) and TMC-1 (Taurus) [34]. (b) Abundance rank obtained in NetWorld for $N = 40$ initial nodes and environments that are qualitatively compatible with the four sets of real data plotted in (a). The abundance rank for $\beta_{crit} = 1.55$ is plotted in brown line. (c) and (d) Comparison between the abundances in TMC-1 and L134N dense clouds with NetWorld simulations for $\beta = 1.8$ and 1.7 respectively. (e) Dependence of abundances of the molecules detected in L134N and TMC-1 on the number of astrophysical reactions that have them as products. (f) Dependence of abundances of networks of size 10 created by NetWorld (in the limit $\beta = \infty$ for simplicity) on the number of paths used to create them. Curve fits are shown in (e) and (f) as dashed lines. Note that CO abundances are out of range in (e) for clarity, but were considered in the fits.

in interstellar clouds where the amount of interstellar dust, and thus of visual extinction A_v , is large enough to prevent the photo-dissociation of molecular species by the external interstellar UV radiation field, which enables their formation and survival. The level of chemical complexity in interstellar clouds is however very different depending on their level of extinction, and on the available amount of molecular hydrogen (H_2) and carbon monoxide (CO) within them [31]. In this way, inter-

stellar clouds can be classified as diffuse atomic (with a fraction of H_2 with respect to total amount of atomic H of $f(H_2) < 10\%$), diffuse molecular ($f(H_2) > 10\%$), translucent ($f(H_2) > 10\%$ and with a fraction of CO with respect to total amount of atomic C of $f(CO) < 90\%$), or dense clouds ($f(H_2) > 10\%$ and $f(CO) > 90\%$). The chemistry in diffuse atomic clouds is very limited [31] while the chemistry in dense clouds presents a very high level of chemical complexity (see e.g. [35, 36]). Therefore, we use here the molecular abundances measured toward diffuse molecular, translucent and dense clouds as test cases for the applicability of the digital environment NetWorld to real scenarios.

In Fig. 4(a) we show the abundances of the chemical compounds detected toward four interstellar clouds ranked in order of their decreasing magnitude: (i) the interstellar cloud ζ Ophiuci, a diffuse molecular cloud where only a few molecules have been found [31]; dust extinction is so weak ($A_v = 1.06$ mag) that UV radiation destroys most of the molecular material; (ii) the translucent cloud located in the direction of the ultracompact HII region K4 in the SgrB2 massive star-forming region [32, 33]; this cloud has an extinction of $A_v = 2.0$ mag, just enough to enable the formation of new molecular species and to protect the molecular content recently formed within the cloud, playing the role of the critical transition in the model; and (iii) L134N (Serpens) and (iv) TMC-1 (Taurus), two dense clouds with $A_v > 10$ mag where an extensive number of both simple and complex molecules have been synthesized thanks not only to the protection of the high A_v , but also to the high fraction of CO present [34]. We refer to Supplementary Information Text S6 for the explanation on how we have obtained the molecular abundances toward the different clouds used in our analysis. Fig. 4(b) shows the abundance rank obtained with NetWorld for 4 different environments that are qualitatively compatible with these four sets of real data: a low value of β representing a harsh environment where most created compounds are rapidly destroyed, a value close to the critical β_{crit} beyond which complexity expands, and two values of β slightly over this transition point. We include the critical environment $\beta_{crit} = 1.55$ for comparison. In Figs. 4(c) and (d) we focus on the potential quantitative agreement between real data and the results of the artificial framework. We compare the abundance ranks for TMC-1 and L134N with those of NetWorld's $\beta = 1.8$ and 1.7 respectively. The abundances are provided excluding the most frequent elements of each ensemble, H_2 in the molecular abundances and the isolated nodes in NetWorld. The real and numerical curves show a quantitative agreement between the number of molecules present in the cloud and the number of configurations in NetWorld, and also in the relative abundance for a large set of molecules and configurations. The framework does not reproduce, however, the truncation shown in the real curves for the 2-3 lowest abundances, but this behavior would disappear if new real data were introduced. Note that the astro-

chemical datasets of molecular species and their measured abundances remain largely incomplete even for the most observed clouds such as TMC-1, and especially for low abundance species or very large molecules that are more difficult to detect (see e.g. [35, 37]). It is also remarkable that the values of β that best fit the astrochemical data of L134N and TMC-1 ($\beta = 1.7$ and 1.8) are slightly beyond $\beta_{crit} = 1.55$, and thus belong to a regime in which expansion of their chemical diversity is expected. Indeed, recent observational works toward the TMC-1 dense cloud have revealed the presence of small polycyclic aromatic hydrocarbons (PAHs), demonstrating the ability of these environments to generate complex molecular structures [35–37].

Finally, a relevant pattern obtained in NetWorld is that the relative abundances of the different structures correlate with the number of paths identified to create them, following a very simple functional dependence of the type $y \propto x^\alpha$ (Fig. 4(f), $\alpha = 1.2 \pm 0.2$ for networks of size 10, $N = 40$ and $\beta = \infty$, see Supporting Information Text S3 for details on how to compute the number of paths to reach a configuration[51]). Note that this is a highly non-trivial result, as different paths have in general very different occurrence probabilities, and this expected diversity could in principle spoil the correlation. In order to check whether a similar relationship might also emerge in astrochemical environments, we plotted in Fig. 4(e) the dependence of the molecular abundances measured toward the dense clouds L134N and TMC-1 on the number of chemical reactions that have them as products, as a simple proxy for the chemical paths. The number of chemical reactions is extracted from the astrochemical reaction dataset KIDA [38]. Surprisingly, the abundance data correlate with the number of reactions following the same functional dependence as NetWorld’s simulations ($y \propto x^\alpha$, where $\alpha_{TMC-1} = 1.0 \pm 0.2$, $r = 0.57$, $p = 2 \cdot 10^{-6}$ and $\alpha_{L134N} = 1.2 \pm 0.3$, $r = 0.54$, $p = 6 \cdot 10^{-4}$, see Supporting Information Text S7). In addition, the α -coefficient of the power-law that relates these two magnitudes is close to 1, indicating that the abundance of a certain molecule would be expected to be approximately proportional to the number of reactions that create it.

We caution, however, that the latter analysis could suffer from important biases. Small molecules that do not have dipole moment, such as N₂ or CH₄, cannot be observed at radio wavelengths. Furthermore, large molecules not only show low abundances but also their partition function is so large that it spreads the emission across many energy levels, making the molecular line intensities very weak. Recent spectroscopic surveys carried out toward chemically rich sources such as SgrB2 N₂ [39], IRAS16293-2422 [40], TMC-1 [35, 37] or G+0.693 (see e.g. [3, 4]) with ALMA, the GBT or the Yebes 40m telescopes, have boosted the detection rate of new molecular species in the ISM in the past decade (see the review [41]), which is starting to alleviate this potential issue. Second, the astrochemical reaction databases are strongly biased

to small molecules. All these chemical networks are based on the pioneering work of [42], whose goal was to investigate the interstellar chemistry of small molecular species. In addition, experimental and theoretical works of gas-phase reactions are extremely challenging especially at the low temperatures typical of interstellar conditions, in particular of radical-radical reactions [43, 44], and not all reactions yielding the same product are equally efficient due to the existence of, for example, high energy barriers. Because of all these reasons, the number of detected molecules and the datasets of reactions that can have them as products are inaccurate.

To address these limitations, we have statistically verified that the correlations between the molecular abundances and the number of reactions shown in Fig. 4(e) for the interstellar clouds TMC-1 and L134N still hold when removing the effect of a controlling variable such as the molecular size, understood as the number of atoms contained within a molecule (partial correlation coefficient $r' = 0.49$ and $p' = 7 \cdot 10^{-5}$ for TMC-1 and $r' = 0.42$ and $p' = 0.007$ for L134N, see Supporting Information Text S7 for more details). This result points to the correlation detected being independent of potential biases due to the molecular size. A thorough statistical analysis, considering other astrochemical magnitudes related to the molecules found in different astrophysical environments, will be carried out in the future.

II. DISCUSSION

In this work we have introduced a conceptual and computational framework called NetWorld describing the evolution of networked structures, where nodes interact following exclusively the optimization of their own dynamical importance. Our results show that, although there is not a causal relationship between NetWorld’s framework and real astrochemical phenomena, a simple model grounded in network science and game theory captures key properties of the process toward chemical complexity and the creation of the building blocks of life. In particular, while our approach does not try to mimic real astrochemistry, it succeeds to explain the emergence of interstellar molecular complexity, and points out as yet unknown astrochemical relationships that could be of importance in our understanding of the formation of prebiotic species in interstellar space.

Multiple methods have been proposed to map real chemistry to artificial chemistry models. Graphs, binary strings, character strings or numbers, among others, have been used to represent molecules [45–48]. Our networks describe molecules where atoms are nodes and their interactions are links. Beyond this mapping, one could add properties and impose more complex rules of interaction in order to get closer to real chemistry. Node labels and properties might be introduced to represent the atom type, the hybridization type, charge, valency, or radicals. Moreover, an energy function could be used to choose

the structure resulting from a reaction on the basis of the change of energy [14]. Based on the current state of our framework, adding these features in NetWorld is possible and, a priori, computationally feasible. However, we stress that fundamental features of NetWorld are its simplicity and abstraction, as we aim to present a framework that transcends chemistry to describe the interaction between complex structures of different nature and in diverse environments, from nodes representing atoms to biomolecules or even species.

In the same line of thought, the natural transition toward complexity emerged from our computational environment has also been observed in complex ecosystems. In particular, Fisher & Mehta [30] reported a strikingly similar pattern in which the skewness of biodiversity rank-abundance curves decreased and the overall diversity increased with carrying capacity, highlighting the sharp transition between stochastic, neutral regimes and selection-dominated, niche regimes. The environment parameter β introduced here and the dust extinction A_v in interstellar chemistry –two quantities directly related since the rate constants of interstellar UV photo-destruction reactions depend exponentially of $-A_v$ [49]–, could be understood as loose proxies for the carrying capacity in the ecological context, in the sense that a low β /low A_v results in a “harsher” environment that limits network/molecules/species richness. Continuing the analogy, in the low β /low A_v regime our network communities and the interstellar molecular abundances show a highly skewed abundance distribution (see Figs. 4(a-b)) and are indeed dominated by stochasticity (the partition probability P introduced in Eq. 1 and the interaction with UV radiation respectively). On the other hand, in the high β /high A_v regime they present low skewness and high diversity, and are dominated by selection of structures with a higher number of paths/reactions leading to them (as shown in Figs. 4(e-f)). All in all, we believe that (i) the similarities between the results in [30], based on models that are firmly rooted in classical ecological theory and checked with real data, (ii) those obtained from molecular abundances in interstellar clouds, and (iii) the ones introduced by our computational environment, derived from a simple framework with no a priori ecological or chemical assumptions, are not coincidental. They instead hint that the long path from the creation of the basic prebiotic compounds in the interstellar medium to the origin of life and its evolution on the early Earth could show universal patterns and common phenomenologies at all scales and across all stages.

Finally, while we have exclusively used sets of isolated and indistinguishable nodes as initial conditions, and have focused on the description of the emerging diversity created, the framework here introduced could be of use in many other contexts. Advancing in the subject of the origin and evolution of early life, potentially fruitful lines of future work could be the analysis of the interaction of small motifs to simulate the polymerization of simple chemical compounds, resembling the phe-

nomenology present in the RNA World, or the search for autocatalytic reactions that could help us advance toward a proto-replication of networks where the concepts of mutation and fitness could be analysed as emerging properties of the system, instead of introducing them *ad-hoc* as has been done so far in the literature.

Acknowledgments

The authors acknowledge fruitful conversations with A. Aguirre-Tamaral, C. Briones, M. Castro, J. García de la Concepción, J.A. García-Martín, R. Guantes, D. Hochberg, J. Iranzo, A. Lucía-Sanz, S. Manrubia and M. Ruiz-Bermejo, and technical support from N. Aguirre and J. Aguirre. I.J.-S. and J.A. received support from Projects PID2019-105552RB-C41, PID2021-122936NB-I00 and MDM-2017-0737 Unidad de Excelencia “María de Maeztu”-Centro de Astrobiología (CSIC-INTA) by the Spanish Ministry of Science and Innovation/State Agency of Research MCIN/AEI/10.13039/501100011033 and by “Fondo Europeo de Desarrollo Regional (FEDER) Una manera de hacer Europa”, through Project ESP2017-86582-C4-1-R. F.P.-S. was supported by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 892961.

Author contributions

M.G.-S., I.J.-S., F.P.-S., and J.A. designed the project and wrote the paper; M.G.-S., F.P.-S., and J.A. conceived the computational environment; M.G.-S. and J.A. implemented the algorithm and performed the numerical experiments; and I.J.-S. and J.A. contributed to the applicability of the results to real astrochemical data.

Supporting Information

The Supporting Information of this paper consists of Supporting Figs. 1-5 and Supporting Information Texts S1-S7.

It is accessible at https://raw.githubusercontent.com/MiguelGarciaSanchez/NetWorld/main/garcia_sanchez_et_al_Supp_Info_2022.pdf

The titles of Supporting Information Texts S1-S7 are:

- S1. NetWorld’s algorithm.
- S2. Computation of the diversity and the relative abundance of the different configurations for a fixed environment in NetWorld.
- S3. Computation of the number of paths to reach a configuration in NetWorld.

- S4. Dependence of the computation time with Net-World's parameters.
- S5. Algorithm availability.
- S6. Molecular abundances measured toward diffuse

molecular, translucent and dense clouds.

- S7. Statistical analysis of the data associated with molecular clouds TMC-1 and L134N.

-
- [1] J. M. Hollis, P. R. Jewell, F. J. Lovas, and A. Remijan, *apjl* **613**, L45 (2004).
- [2] Belloche, A., Garrod, R. T., Müller, H. S. P., Menten, K. M., Medvedev, I., Thomas, J., and Kisiel, Z., *A&A* **628**, A10 (2019), URL <https://doi.org/10.1051/0004-6361/201935428>.
- [3] I. Jiménez-Serra, J. Martín-Pintado, V. M. Rivilla, L. Rodríguez-Almeida, E. R. Alonso Alonso, S. Zeng, E. J. Cocinero, S. Martín, M. Requena-Torres, R. Martín-Domenech, et al., *Astrobiology* **20**, 1048 (2020), pMID: 32283036, <https://doi.org/10.1089/ast.2019.2125>, URL <https://doi.org/10.1089/ast.2019.2125>.
- [4] V. M. Rivilla, I. Jiménez-Serra, J. Martín-Pintado, C. Briones, L. F. Rodríguez-Almeida, F. Rico-Villas, B. Tercero, S. Zeng, L. Colzi, P. de Vicente, et al., *Proceedings of the National Academy of Sciences* **118** (2021), ISSN 0027-8424, <https://www.pnas.org/content/118/22/e2101314118.full.pdf>, URL <https://www.pnas.org/content/118/22/e2101314118>.
- [5] K. Altwegg, H. Balsiger, J. Berthelier, A. Bieler, U. Calmonte, S. Fuselier, F. Goesmann, S. Gasc, T. I. Gombosi, L. Le Roy, et al., *Monthly Notices of the Royal Astronomical Society* **469**, S130 (2017), ISSN 0035-8711, <https://academic.oup.com/mnras/article-pdf/469/Suppl.2/S130/18563276/stx1415.pdf>, URL <https://doi.org/10.1093/mnras/stx1415>.
- [6] Y. Furukawa, Y. Chikaraishi, N. Ohkouchi, N. O. Ogawa, D. P. Glavin, J. P. Dworkin, C. Abe, and T. Nakamura, *Proceedings of the National Academy of Sciences* **116**, 24440 (2019), ISSN 0027-8424, <https://www.pnas.org/content/116/49/24440.full.pdf>, URL <https://www.pnas.org/content/116/49/24440>.
- [7] J. V. Neumann and A. W. Burks, *Theory of Self-Reproducing Automata* (University of Illinois Press, USA, 1966).
- [8] S. Wolfram, *Rev. Mod. Phys.* **55**, 601 (1983), URL <https://link.aps.org/doi/10.1103/RevModPhys.55.601>.
- [9] M. Gardner, *Scientific American* **223**, 120 (1970).
- [10] S. Rasmussen, C. Knudsen, R. Feldberg, and M. Hindsholm, *Physica D: Nonlinear Phenomena* **42**, 111–134 (1990).
- [11] R. E. Lenski, C. Ofria, T. C. Collier, and C. Adami, *Nature* **400**, 661–664 (1999).
- [12] C. Adami, C. Ofria, and T. C. Collier, *Proceedings of the National Academy of Sciences* **97**, 4463 (2000), ISSN 0027-8424, <https://www.pnas.org/content/97/9/4463.full.pdf>, URL <https://www.pnas.org/content/97/9/4463>.
- [13] P. Dittrich, J. Ziegler, and W. Banzhaf, *Artificial Life* **7**, 225 (2001).
- [14] G. Benkő, C. Flamm, and P. F. Stadler, *Journal of Chemical Information and Computer Sciences* **43**, 1085 (2003).
- [15] W. Banzhaf and L. Yamamoto, *Artificial Chemistries* (The MIT Press, Cambridge, MA, USA, 2010).
- [16] M. E. J. Newman, *Networks: An Introduction* (Oxford University Press, Inc., New York, NY, USA, 2010).
- [17] J. Iranzo, J. M. Buldú, and J. Aguirre, *Nature Communications* **7**, 13273 (2016).
- [18] J. Iranzo, F. Pablo-Martí, and J. Aguirre, *Phys. Rev. Research* **2**, 043352 (2020), URL <https://link.aps.org/doi/10.1103/PhysRevResearch.2.043352>.
- [19] J. Gómez-Gardeñes, I. Reinares, A. Arenas, and L. M. Floria, *Sci Rep* **2**, 620 (2012).
- [20] J. Aguirre, D. Papo, and J. M. Buldú, *Nature Physics* **9**, 230 (2013).
- [21] Z. Wang, A. Szolnoki, and M. Perc, *New Journal of Physics* **16**, 033041 (2014).
- [22] Z. Wang, L. Wang, A. Szolnoki, and M. Perc, *The European Physical Journal B* **88**, 124 (2015).
- [23] U. Alvarez-Rodriguez, F. Battiston, G. de Arruda, Y. Moreno, M. Perc, and V. Latora, *Nature Human Behavior* **5**, 586 (2021).
- [24] A. Kumar, S. Chowdhary, V. Capraro, and M. c. v. Perc, *Phys. Rev. E* **104**, 054308 (2021), URL <https://link.aps.org/doi/10.1103/PhysRevE.104.054308>.
- [25] G. Bianconi, *Higher-Order Networks* (Cambridge University Press, 2021).
- [26] I. T. Jolliffe and J. Cadima, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **374**, 20150202 (2016).
- [27] B. J. McGill, R. S. Etienne, J. S. Gray, D. Alonso, M. J. Anderson, H. K. Benecha, M. Dornelas, B. J. Enquist, J. L. Green, F. He, et al., *Ecology Letters* **10**, 995–1015 (2007).
- [28] C. Pedrós-Alió, *Annual Review of Marine Science* **4**, 449 (2012), pMID: 22457983, <https://doi.org/10.1146/annurev-marine-120710-100948>, URL <https://doi.org/10.1146/annurev-marine-120710-100948>.
- [29] J. Gregori, C. Perales, F. Rodriguez-Frias, J. I. Esteban, J. Quer, and E. Domingo, *Virology* **493**, 227 (2016), ISSN 0042-6822, URL <https://www.sciencedirect.com/science/article/pii/S004268221630037X>.
- [30] C. K. Fisher and P. Mehta, *Proceedings of the National Academy of Sciences* **111**, 13111 (2014), ISSN 0027-8424, <https://www.pnas.org/content/111/36/13111.full.pdf>, URL <https://www.pnas.org/content/111/36/13111>.
- [31] T. P. Snow and B. J. McCall, *Annual Review of Astronomy and Astrophysics* **44**, 367 (2006).
- [32] Thiel, V., Belloche, A., Menten, K. M., Garrod, R. T., and Müller, H. S. P., *A&A* **605**, L6 (2017), URL <https://doi.org/10.1051/0004-6361/201731495>.
- [33] Corby, J. F., McGuire, B. A., Herbst, E., and Remijan, A. J., *Astronomy & Astrophysics* **610**, A10 (2018), URL

- <https://doi.org/10.1051/0004-6361/201730988>.
- [34] M. Agúndez and V. Wakelam, *Chem. Rev.* **113**, 8710 (2013).
- [35] Cernicharo, J., Agúndez, M., Cabezas, C., Tercero, B., Marcelino, N., Pardo, J. R., and de Vicente, P., *A&A* **649**, L15 (2021), URL <https://doi.org/10.1051/0004-6361/202141156>.
- [36] B. A. McGuire, R. A. Loomis, A. M. Burkhardt, K. L. K. Lee, C. N. Shingledecker, S. B. Charnley, I. R. Cooke, M. A. Cordiner, E. Herbst, S. Kalenskii, et al., *Science* **371**, 1265 (2021).
- [37] B. A. McGuire, **239**, 17 (2018), URL <https://doi.org/10.3847/1538-4365/aae5d2>.
- [38] V. Wakelam, J.-C. Loison, E. Herbst, B. Pavone, A. Bergeat, K. Béroff, M. Chabot, A. Faure, D. Galli, W. D. Geppert, et al., *ApJS* **217**, 20 (2015), URL <https://doi.org/10.1088/0067-0049/217/2/20>.
- [39] Belloche, A., Müller, H. S. P., Garrod, R. T., and Menten, K. M., *Astronomy & Astrophysics* **587**, A91 (2016), URL <https://doi.org/10.1051/0004-6361/201527268>.
- [40] J. K. Jørgensen, M. H. D. van der Wiel, A. Coutens, J. M. Lykke, H. S. P. Müller, E. F. van Dishoeck, H. Calcutt, P. Bjerkeli, T. L. Bourke, M. N. Drozdovskaya, et al., *Astronomy & Astrophysics* **595**, A117 (2016), 1607.08733.
- [41] B. A. McGuire, *The Astrophysical Journal Supplement Series* **259**, 30 (2022), URL <https://doi.org/10.3847/1538-4365/ac2a48>.
- [42] S. S. Prasad and J. Huntress, W. T., *Astrophysical Journal* **43**, 1 (1980).
- [43] R. Shannon, M. Blitz, A. Goddard, and D. Heard, *Nature Chemistry* **5**, 745 (2013), © 2013 Macmillan Publishers Limited. All rights reserved. This is an author produced version of a paper published in *Nature Chemistry*. Uploaded in accordance with the publisher's self-archiving policy, URL <https://eprints.whiterose.ac.uk/87629/>.
- [44] J. G. de la Concepción, C. Puzzarini, V. Barone, I. Jiménez-Serra, and O. Roncero, *The Astrophysical Journal* **922**, 169 (2021), URL <https://doi.org/10.3847/1538-4357/ac1e94>.
- [45] J. Farmer, S. A. Kauffman, and N. H. Packard, *Physica D: Nonlinear Phenomena* **22**, 50 (1986), ISSN 01672789, URL <https://linkinghub.elsevier.com/retrieve/pii/0167278986902332>.
- [46] W. Banzhaf, *Biological Cybernetics* **69**, 275 (1993), ISSN 0340-1200, 1432-0770, URL <http://link.springer.com/10.1007/BF00203124>.
- [47] P. Dittrich and W. Banzhaf, *Artificial Life* **4**, 203 (1998), ISSN 1064-5462, 1530-9185, URL <https://direct.mit.edu/artl/article/4/2/203-220/2298>.
- [48] P. Dittrich, F. Liljeros, A. Soulier, and W. Banzhaf, *Physical Review Letters* **84**, 3205 (2000), ISSN 0031-9007, 1079-7114, URL <https://link.aps.org/doi/10.1103/PhysRevLett.84.3205>.
- [49] J. Holdship, S. Viti, I. Jiménez-Serra, A. Makrymallis, and F. Priestley, **154**, 38 (2017), URL <https://doi.org/10.3847/1538-3881/aa773f>.
- [50] Note that, in opposition to real chemistry, in our model all nodes are equal, and therefore there are very few potential configurations of small size (e.g. there is only 1 for size 2, while there is a large number of molecules of two atoms in nature).
- [51] We calculated the number of paths of each configuration for $\beta = \infty$ (with no loss of generality), as it is the case where there is no network partition and the computation is simpler.