

# CONNECTED REDUCED PRODUCTS

Miloš S. Kurilić<sup>1</sup>

## Abstract

A structure  $\mathbb{X} = \langle X, \rho \rangle$ , where  $\rho$  is a binary relation on the set  $X$ , is connected iff the minimal equivalence relation containing  $\rho$  is the full relation on  $X$ . Connectivity is not a first order property but it is expressible by a  $L_{\omega_1\omega}$ -sentence. We show that, for a set  $I$  the following conditions are equivalent

- (a)  $|I|$  is less than the first measurable cardinal,
- (b) For each filter  $\Phi \subset P(I)$  and each family  $\{\mathbb{X}_i : i \in I\}$  of binary structures, the reduced product  $\prod_{\Phi} \mathbb{X}_i$  is connected, iff there are a finite set  $K \subset I$  and  $n \in \omega$  such that  $\mathbb{X}_i$  is connected, for each  $i \in K$  and  $\{i \in I : \mathbb{X}_i \text{ is of diameter } \leq n\} \cup K \in \Phi$ ,
- (c) The ultraproduct  $\prod_{\mathcal{U}} \mathbb{G}_{\omega}$  is a disconnected graph for each non-principal ultrafilter  $\mathcal{U} \subset P(I)$ , where  $\mathbb{G}_{\omega}$  is the linear graph on  $\omega$ .

Moreover, the implication “ $\Leftarrow$ ” in (b) holds in ZFC. For a disconnected  $\prod_{\Phi} \mathbb{X}_i$  we describe the components of  $\prod_{\Phi} \mathbb{X}_i$ .

*2010 Mathematics Subject Classification:* 03C20

*Key words and phrases:* Reduced product, connected binary relation, components.

## 1 Preliminaries

**Reduced products** Let  $L = \{R\}$  be the language with one binary relational symbol. Let  $I$  be a non-empty set,  $\Phi \subset P(I)$  a filter and  $\mathbb{X}_i = \langle X_i, \rho_i \rangle$ , for  $i \in I$ ,  $L$ -structures. Let  $\sim_{\Phi}$  be the binary relation on the product  $\prod_{i \in I} X_i$  defined in the following way: for  $x = \langle x_i : i \in I \rangle, y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$

$$x \sim_{\Phi} y \Leftrightarrow \{i \in I : x_i = y_i\} \in \Phi.$$

It is evident that the relation  $\sim_{\Phi}$  is an equivalence relation on the set  $\prod_{i \in I} X_i$ . Let  $[x]_{\Phi} = \{y \in \prod_{i \in I} X_i : x \sim_{\Phi} y\} \in (\prod_{i \in I} X_i) / \sim_{\Phi}$  denote the equivalence class of  $x$ . The corresponding *reduced product*  $(\prod_{i \in I} \mathbb{X}_i) / \Phi$  (shortly  $\prod_{\Phi} \mathbb{X}_i$ ) is the structure

$$\left\langle (\prod_{i \in I} X_i) / \sim_{\Phi}, \rho \right\rangle,$$

where the binary relation  $\rho$  on the quotient  $(\prod_{i \in I} X_i) / \sim_{\Phi}$  is defined by

$$[x]_{\Phi} \rho [y]_{\Phi} \Leftrightarrow \{i \in I : x_i \rho_i y_i\} \in \Phi.$$

We will use the following evident fact (see [1], p. 177).

<sup>1</sup>Department of Mathematics and Informatics, Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 4, 21000 Novi Sad, Serbia. e-mail: milos@dmi.uns.ac.rs

**Fact 1.1** *If  $\Phi$  is a filter on  $I$  and  $J \in \Phi$ , then  $\Phi \upharpoonright J := \Phi \cap P(J)$  is a filter on  $J$  and  $f : \prod_{\Phi} \mathbb{X}_i \cong \prod_{\Phi \upharpoonright J} \mathbb{X}_j := (\prod_{i \in J} \mathbb{X}_i) / \sim_{\Phi \upharpoonright J}$ , where  $f([x]_{\Phi}) = [x \upharpoonright J]_{\Phi \upharpoonright J}$ .*

First-order properties of reduced products are well understood. If  $L$  is any language, an  $L$ -formula is a *basic Horn formula* iff it is of the form  $\eta_0 \vee \dots \vee \eta_{m-1}$ , where  $\eta_i$ 's are literals and at most one of them is atomic. The set of *Horn formulas* is the closure of the set of basic Horn formulas under  $\wedge$ ,  $\exists$  and  $\forall$ . It is known that each Horn formula  $\varphi(v_0, \dots, v_{n-1})$  is *preserved in reduced products*; i.e., for each set  $I$ , filter  $\Phi \subset P(I)$ , family of  $L$ -structures  $\{\mathbb{X}_i : i \in I\}$  and  $n$ -tuple  $\langle x^0, \dots, x^{n-1} \rangle \in (\prod_{i \in I} X_i)^n$  we have

$$\left\{ i \in I : \mathbb{X}_i \models \varphi[x_i^0, \dots, x_i^{n-1}] \right\} \in \Phi \Rightarrow \prod_{\Phi} \mathbb{X}_i \models \varphi \left[ [x^0]_{\Phi}, \dots, [x^{n-1}]_{\Phi} \right]. \quad (1)$$

By the well known result of Keisler (under CH) and Galvin (in ZFC) we have: an  $L$ -sentence  $\varphi$  is *preserved in reduced products*, i.e., for each  $I$ ,  $\Phi$  and  $\{\mathbb{X}_i : i \in I\}$  as above

$$\left\{ i \in I : \mathbb{X}_i \models \varphi \right\} \in \Phi \Rightarrow \prod_{\Phi} \mathbb{X}_i \models \varphi, \quad (2)$$

iff  $\varphi$  is logically equivalent to a Horn  $L$ -sentence (see [1], p. 365).

Putting the implication “ $\Leftarrow$ ” in (2) we obtain a definition of a sentence *preserved under reduced factors*. It is known that the positive sentences (without  $\neg$ ) are of that kind (see [1], p. 340).

**Connected binary structures** Let  $L = \{R\}$  be the language with one binary relational symbol. Let  $\mathbb{X} = \langle X, \rho \rangle$  be an  $L$ -structure, where  $\rho \subset X^2$  is a reflexive binary relation and let  $\rho^{-1}$  be the inverse of  $\rho$ . For  $\varepsilon \in 2 := \{0, 1\}$  let

$$\rho^{\varepsilon} = \begin{cases} \rho & \text{if } \varepsilon = 0, \\ \rho^{-1} & \text{if } \varepsilon = 1. \end{cases} \quad (3)$$

If  $x, y \in X$  and  $n \in \omega$ , a *path of length  $n + 1$  from  $x$  to  $y$*  is a pair  $\langle \bar{z}, \varepsilon \rangle$ , where  $\bar{z} = \langle z_0, \dots, z_{n-1} \rangle \in X^n$  and  $\varepsilon = \langle \varepsilon_0, \dots, \varepsilon_n \rangle \in 2^{n+1}$  and such that

$$x \rho^{\varepsilon_0} z_0 \rho^{\varepsilon_1} z_1 \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} z_{n-1} \rho^{\varepsilon_n} y.$$

(So, for  $n = 0$ ,  $x \rho^{\varepsilon_0} y$ .) Then we will write  $d(x, y) \leq n + 1$  and  $d(x, y) = \infty$  will denote that there is no (finite) path from  $x$  to  $y$ .

The binary relation  $\sim$  on the set  $X$  defined by  $x \sim y$  iff there is finite path from  $x$  to  $y$ , is an equivalence relation, the corresponding equivalence classes are the (*connected*) *components of  $\mathbb{X}$*  and the structure  $\mathbb{X}$  is *connected* iff it has only one component. (It is easy to see that  $\sim$  is the transitive closure  $(\rho_s)_t$  of the symmetrization  $\rho_s := \rho \cup \rho^{-1}$  of the relation  $\rho$ .)

We extend these definitions for arbitrary  $L$ -structure  $\mathbb{X} = \langle X, \rho \rangle$  in the following way: instead of  $\rho$  we take its reflexivization  $\rho_R := \rho \cup \Delta_X$  and call  $\mathbb{X}$  connected iff  $\mathbb{X}_R := \langle X, \rho_R \rangle$  is connected, components of  $\mathbb{X}$  are components of  $\mathbb{X}_R$ , etc.

The statement  $d(x, y) \leq n + 1$  is expressible by the first order formula

$$\varphi_{d \leq n+1}(x, y) := \exists z_0, \dots, z_{n-1} \bigvee_{\varepsilon \in 2^{n+1}} x \rho^{\varepsilon_0} z_0 \rho^{\varepsilon_1} \dots \rho^{\varepsilon_{n-1}} z_{n-1} \rho^{\varepsilon_n} y.$$

It is evident that  $\varphi_{d \leq n+1}$  is positive and  $\neg \varphi_{d \leq n+1}$  is Horn. The property of an  $L$ -structure that each two points are connected by a path of length  $\leq n + 1$  is expressible by the first order sentence

$$\varphi_n^{\text{conn}} := \forall x, y \varphi_{d \leq n+1}(x, y).$$

If such  $n \in \omega$  exists we will say that  $\mathbb{X}$  is of *finite diameter*. The connectedness of an  $L$ -structure is expressible by the  $L_{\omega_1 \omega}$ -sentence

$$\varphi^{\text{conn}} := \forall x, y \bigvee_{n \in \omega} \varphi_{d \leq n+1}(x, y).$$

## 2 Connected reduced products. Components

In the sequel we assume that

- (\*)  $I \neq \emptyset$  is a set,  $\Phi$  is a filter in  $P(I)$ ,  $\mathbb{X}_i = \langle X_i, \rho_i \rangle$ ,  $i \in I$ , are  $L$ -structures and  $\prod \mathbb{X}_i := \langle \prod_{i \in I} X_i, \sigma \rangle$  and  $\prod_{\Phi} \mathbb{X}_i := \langle \prod_{i \in I} X_i / \sim_{\Phi}, \rho \rangle$  are the corresponding direct and reduced product respectively.

**Fact 2.1** (\*) *If  $x, y \in \prod_{i \in I} X_i$ ,  $A \subset I$ ,  $x \upharpoonright (I \setminus A) = y \upharpoonright (I \setminus A)$ ,  $n \in \omega$ , and there is  $\varepsilon \in 2^{n+1}$  such that for each  $i \in A$  there is  $\bar{z} \in X_i^n$  such that  $\langle \bar{z}, \varepsilon \rangle$  is a path from  $x_i$  to  $y_i$ , then in  $\prod_{\Phi} \mathbb{X}_i$  there is a path from  $[x]_{\Phi}$  to  $[y]_{\Phi}$ .*

**Proof.** Let  $\varepsilon = \langle \varepsilon_0, \dots, \varepsilon_n \rangle$ , and for  $i \in A$  let  $\langle \langle z_i^0, \dots, z_i^{n-1} \rangle, \varepsilon \rangle$  be a path from  $x_i$  to  $y_i$

$$x_i \rho_i^{\varepsilon_0} z_i^0 \rho_i^{\varepsilon_1} z_i^1 \rho_i^{\varepsilon_2} \dots \rho_i^{\varepsilon_{n-1}} z_i^{n-1} \rho_i^{\varepsilon_n} y_i. \quad (4)$$

Let  $t^m \in \prod_{i \in I} X_i$ , for  $m < n$ , be defined by

$$t_i^m = \begin{cases} x_i & \text{if } i \in I \setminus A, \\ z_i^m & \text{if } i \in A. \end{cases} \quad (5)$$

By (4) in the direct product we have  $x \sigma^{\varepsilon_0} t^0 \sigma^{\varepsilon_1} t^1 \sigma^{\varepsilon_2} \dots \sigma^{\varepsilon_{n-1}} t^{n-1} \sigma^{\varepsilon_n} y$  and, hence, in the reduced product  $\prod_{\Phi} \mathbb{X}_i$  we have

$$[x]_{\Phi} \rho^{\varepsilon_0} [t^0]_{\Phi} \rho^{\varepsilon_1} [t^1]_{\Phi} \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [t^{n-1}]_{\Phi} \rho^{\varepsilon_n} [y]_{\Phi}. \quad (6)$$

So, this is a path from  $[x]_{\Phi}$  to  $[y]_{\Phi}$ .  $\square$

**Fact 2.2** *If  $\kappa \geq \omega$  is a non-measurable cardinal and  $\Phi \subset P(\kappa)$  a filter consisting of infinite sets, then there is a partition  $\kappa = \bigcup_{n \in \omega} I_n$  such that for each  $F \in \Phi$  we have  $F \cap I_n \neq \emptyset$ , for infinitely many  $n \in \omega$ .*

**Proof.** First let  $\bigcap \Phi = \emptyset$ . Let  $\mathcal{U} \subset P(\kappa)$  be an ultrafilter such that  $\Phi \subset \mathcal{U}$ . Then the ultrafilter  $\mathcal{U}$  is non-principal and, since the cardinal  $\kappa$  is non-measurable,  $\mathcal{U}$  is not  $\omega$ -complete. Thus there is a partition  $\kappa = \bigcup_{n \in \omega} I_n$  such that  $I_n \notin \mathcal{U}$  and, hence,  $\kappa \setminus I_n \in \mathcal{U}$ , for each  $n \in \omega$ . Assuming that there are  $F \in \Phi$  and  $m \in \omega$  such that  $F \subset \bigcup_{n \leq m} I_n$ , we would have  $F \cap \bigcap_{n \leq m} \kappa \setminus I_n = F \setminus \bigcup_{n \leq m} I_n = \emptyset \in \mathcal{U}$ , which is false. Thus,  $|\{n \in \omega : F \cap I_n \neq \emptyset\}| = \omega$ , for each  $F \in \Phi$ .

Second let  $A := \bigcap \Phi \neq \emptyset$ . If  $|A| < \omega$ , then, since  $\Phi \subset P(\kappa) \setminus [\kappa]^{<\omega}$ ,  $\Phi' := \Phi \upharpoonright (\kappa \setminus A)$  is a non-principal filter on  $\kappa \setminus A$  and we work as above.

If  $|A| \geq \omega$ ,  $I_0 = \kappa \setminus A$  and  $A = \bigcup_{n \in \mathbb{N}} I_n$  is any partition of  $A$ , then for each  $F \in \Phi$  we have  $F \cap I_n \neq \emptyset$ , for all  $n \in \mathbb{N}$ .  $\square$

**Theorem 2.3** (\*) *If the size of the index set  $I$  is less than the first measurable cardinal, then the following conditions are equivalent*

- (a) *The reduced product  $\prod_{\Phi} \mathbb{X}_i$  is connected,*
- (b) *There are a finite set  $K \subset I$  and  $n \in \omega$  such that  $\mathbb{X}_i$  is connected, for each  $i \in K$ , and  $\{i \in I : \mathbb{X}_i \models \varphi_n^{\text{conn}}\} \cup K \in \Phi$ .*

*The implication (b)  $\Rightarrow$  (a) is true in ZFC (without any assumptions on  $|I|$ ).*

**Proof.** W.l.o.g. we can assume that  $\rho_i \subset X_i$ ,  $i \in I$ , are reflexive relations.

(a)  $\Rightarrow$  (b). We prove the contrapositive: assuming that

$$\forall K \in [I]^{<\omega} \left( \forall i \in K (\mathbb{X}_i \models \varphi^{\text{conn}}) \Rightarrow \forall n \in \omega \{i \in I : \mathbb{X}_i \models \varphi_n^{\text{conn}}\} \cup K \notin \Phi \right) \quad (7)$$

we show that  $\prod_{\Phi} \mathbb{X}_i$  is disconnected.

If the filter  $\Phi$  contains a finite set, then  $\Phi$  is a principal filter generated by some  $K \in [I]^{<\omega}$  and, by (7), there is  $i_0 \in K$  such that  $\mathbb{X}_{i_0}$  is disconnected. Since  $\Phi \upharpoonright K = \{K\}$ , by Fact 1.1 we have  $\prod_{\Phi} \mathbb{X}_i \cong \prod_{\Phi \upharpoonright K} \mathbb{X}_i \cong \prod_{i \in K} \mathbb{X}_i$ . We

take  $x_{i_0}, y_{i_0} \in X_{i_0}$  such that there is no path from  $x_{i_0}$  to  $y_{i_0}$  in  $\mathbb{X}_{i_0}$  and choose arbitrary  $x_i = y_i \in X_i$ , for  $i \in K \setminus \{i_0\}$ . Assuming that there is a path from  $x := \langle x_i : i \in K \rangle$  to  $y := \langle y_i : i \in K \rangle$  in  $\prod_{i \in K} \mathbb{X}_i$  we would obtain a path from  $x_{i_0}$  to  $y_{i_0}$  in  $\mathbb{X}_{i_0}$ , which is impossible. Thus the reduced product  $\prod_{\Phi} \mathbb{X}_i$  is disconnected.

So, the remaining case is when  $\Phi$  is a filter consisting of infinite subsets of  $I$ . Let  $A_n := \{i \in I : \mathbb{X}_i \models \varphi_n^{\text{conn}}\}$ , for  $n \in \omega$ . Then  $A_0 \subset A_1 \subset A_2 \subset \dots$  and, defining  $I_0 := A_0$ ,  $I_n := A_n \setminus A_{n-1}$ , for  $n \in \mathbb{N}$ , and  $I_\infty = I \setminus \bigcup_{n \in \omega} A_n$ , we have

$$I = \bigcup_{n \in \omega} I_n \cup I_\infty.$$

We continue with a case analysis.

1.  $I_\infty \in \Phi$ . Then  $|I_\infty| \geq \omega$  is a non-measurable cardinal,  $\Phi \upharpoonright I_\infty$  is a filter on  $I_\infty$  consisting of infinite sets, and, by Fact 2.2, there is a partition  $I_\infty = \bigcup_{n \in \omega} I'_n$  such that for each  $F \in \Phi \upharpoonright I_\infty$  we have  $F \cap I'_n \neq \emptyset$ , for infinitely many  $n \in \omega$ . If  $n \in \omega$ , then for  $i \in I'_n$  we have  $i \notin A_n$  and we choose  $x_i, y_i \in X_i$  such that  $d(x_i, y_i) > n + 1$ . For  $i \in I \setminus I_\infty$  we choose arbitrary  $x_i = y_i \in X_i$ . Let  $x := \langle x_i : i \in I \rangle$ ,  $y := \langle y_i : i \in I \rangle$  and suppose that in  $\prod_{\Phi} \mathbb{X}_i$  there is a path from  $[x]_{\Phi}$  to  $[y]_{\Phi}$

$$[x]_{\Phi} \rho^{\varepsilon_0} [z^0]_{\Phi} \rho^{\varepsilon_1} [z^1]_{\Phi} \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [z^{n-1}]_{\Phi} \rho^{\varepsilon_n} [y]_{\Phi}. \quad (8)$$

Then there would be  $F \in \Phi$  which witnesses that; in other words,

$$\forall i \in F \quad x_i \rho_i^{\varepsilon_0} z_i^0 \rho_i^{\varepsilon_1} z_i^1 \rho_i^{\varepsilon_2} \dots \rho_i^{\varepsilon_{n-1}} z_i^{n-1} \rho_i^{\varepsilon_n} y_i. \quad (9)$$

Now let  $m > n$  be such that there is  $i \in F \cap I'_m$ . By (9) there is an  $(n + 1)$ -path from  $x_i$  to  $y_i$ , which is impossible by the choice of  $x_i$  and  $y_i$  ( $d(x_i, y_i) > n + 1$ ). So, there is no path from  $[x]_{\Phi}$  to  $[y]_{\Phi}$  in  $\prod_{\Phi} \mathbb{X}_i$  and this structure is disconnected.

2.  $I_\infty \notin \Phi$ . Then we have  $F \cap \bigcup_{n \in \omega} I_n \neq \emptyset$ , for all  $F \in \Phi$ .

2.1. Each  $F \in \Phi$  intersects infinitely many sets  $I_n$ . If  $n \in \mathbb{N}$ , then for  $i \in I_n$  we have  $i \notin A_{n-1}$  and we choose  $x_i, y_i \in X_i$  such that  $d(x_i, y_i) > n$ . For  $i \in I \setminus \bigcup_{n \in \mathbb{N}} I_n$  we choose arbitrary  $x_i = y_i \in X_i$ . Let  $x := \langle x_i : i \in I \rangle$ ,  $y := \langle y_i : i \in I \rangle$  and suppose that in  $\prod_{\Phi} \mathbb{X}_i$  there is a path from  $[x]_{\Phi}$  to  $[y]_{\Phi}$

$$[x]_{\Phi} \rho^{\varepsilon_0} [z^0]_{\Phi} \rho^{\varepsilon_1} [z^1]_{\Phi} \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [z^{n-1}]_{\Phi} \rho^{\varepsilon_n} [y]_{\Phi}. \quad (10)$$

Then there would be  $F \in \Phi$  which witnesses that; in other words,

$$\forall i \in F \quad x_i \rho_i^{\varepsilon_0} z_i^0 \rho_i^{\varepsilon_1} z_i^1 \rho_i^{\varepsilon_2} \dots \rho_i^{\varepsilon_{n-1}} z_i^{n-1} \rho_i^{\varepsilon_n} y_i. \quad (11)$$

By the assumption of 2.1 there is  $m > n + 1$  such that  $F \cap I_m \neq \emptyset$ . So, for  $i \in F \cap I_m$ , by (11) there is an  $(n + 1)$ -path from  $x_i$  to  $y_i$ , which is impossible by

the choice of  $x_i$  and  $y_i$  ( $d(x_i, y_i) > m$ ). So, there is no path from  $[x]_\Phi$  to  $[y]_\Phi$  in  $\prod_\Phi \mathbb{X}_i$  and this structure is disconnected.

2.2. There are  $n_0 \in \omega$  and  $F_0 \in \Phi$  such that  $F_0 \subset A_{n_0} \cup I_\infty$ . By (7) taking  $K = \emptyset$  we obtain  $A_n \notin \Phi$ , for all  $n \in \omega$ . So, assuming that  $F \cap I_\infty = \emptyset$ , for some  $F \in \Phi$ , we would have  $A_{n_0} \in \Phi$ , which is false. Thus

$$\forall F \in \Phi \quad F \cap I_\infty \neq \emptyset, \quad (12)$$

and  $\Phi \upharpoonright I_\infty$  is a proper filter on  $I_\infty$ .

2.2.1.  $|F \cap I_\infty| \geq \omega$ , for all  $F \in \Phi$ . Then  $|I_\infty| \geq \omega$  is a non-measurable cardinal,  $\Phi \upharpoonright I_\infty$  is a filter on  $I_\infty$  consisting of infinite sets, and, by Fact 2.2, there is a partition  $I_\infty = \bigcup_{n \in \omega} I'_n$  such that for each  $F \in \Phi \upharpoonright I_\infty$  we have  $F \cap I'_n \neq \emptyset$ , for infinitely many  $n \in \omega$ . Now we continue exactly as in Case 1.

2.2.2. The set  $F \cap I_\infty$  is finite, for some  $F \in \Phi$ . Then  $\Phi \upharpoonright I_\infty$  is a principal filter generated by some finite set  $J \subset I_\infty$  and, hence,  $A_{n_0} \cup J \in \Phi$ . So, by (7), there is  $i_0 \in J$  such that  $\mathbb{X}_{i_0}$  is disconnected. We take  $x_{i_0}, y_{i_0} \in X_{i_0}$  such that there is no path from  $x_{i_0}$  to  $y_{i_0}$  in  $\mathbb{X}_{i_0}$  and choose arbitrary  $x_i = y_i \in X_i$ , for  $i \in I \setminus \{i_0\}$ . Let  $x := \langle x_i : i \in I \rangle$ ,  $y := \langle y_i : i \in I \rangle$  and suppose that there is a path from  $[x]_\Phi$  to  $[y]_\Phi$  in  $\prod_\Phi \mathbb{X}_i$ ,

$$[x]_\Phi \rho^{\varepsilon_0} [z^0]_\Phi \rho^{\varepsilon_1} [z^1]_\Phi \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [z^{n-1}]_\Phi \rho^{\varepsilon_n} [y]_\Phi. \quad (13)$$

Then there would be  $F \in \Phi$  which witnesses that; in other words,

$$\forall i \in F \quad x_i \rho_i^{\varepsilon_0} z_i^0 \rho_i^{\varepsilon_1} z_i^1 \rho_i^{\varepsilon_2} \dots \rho_i^{\varepsilon_{n-1}} z_i^{n-1} \rho_i^{\varepsilon_n} y_i. \quad (14)$$

But  $i_0 \in J \subset F$  and we obtain a path from  $x_{i_0}$  to  $y_{i_0}$  in  $\mathbb{X}_{i_0}$ , which is impossible. Thus the reduced product  $\prod_\Phi \mathbb{X}_i$  is disconnected.

(b)  $\Rightarrow$  (a). Let  $K \in [I]^{<\omega}$ ,  $n \in \omega$  and, defining  $A := \{i \in I : \mathbb{X}_i \models \varphi_n^{\text{conn}}\}$ , suppose that

- (i)  $\mathbb{X}_i$  is connected, for each  $i \in K$ , and
- (ii)  $A \cup K \in \Phi$ .

Since  $\varphi_n^{\text{conn}} \Rightarrow \varphi^{\text{conn}}$  we can assume that  $A \cap K = \emptyset$ . For  $x = \langle x_i : i \in I \rangle$ ,  $y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$  we construct a path from  $[x]_\Phi$  to  $[y]_\Phi$  in  $\prod_\Phi \mathbb{X}_i$ .

If  $i \in A$ , then  $\mathbb{X}_i \models \varphi_n^{\text{conn}}$  and, since  $x_i, y_i \in X_i$ , there is a path  $\langle \bar{z}, \varepsilon \rangle$  of length  $n + 1$  from  $x_i$  to  $y_i$ ; thus we can choose one such  $\varepsilon$ . Let  $\varepsilon^k \in 2^{n+1}$ ,  $k < l$ , be an enumeration of the  $n + 1$ -tuples chosen in this way for  $i \in A$ . Then defining  $A_{\varepsilon^k} = \{i \in A : \varepsilon^k \text{ is chosen for } x_i \text{ and } y_i\}$ , for  $k < l$ , we have

$$A = \bigcup_{k < l} A_{\varepsilon^k}.$$

Let  $y' \in \prod_{i \in I} X_i$  be defined by

$$y'_i = \begin{cases} x_i & \text{if } i \in I \setminus A, \\ y_i & \text{if } i \in A. \end{cases} \quad (15)$$

We define sequences  $s^m = \langle s_i^m : i \in I \rangle \in \prod_{i \in I} X_i$ , for  $m \leq l-2$ , such that there are  $l$ -many paths of length  $(n+1)$  witnessing that  $[x]_\Phi \sim [s^0]_\Phi \sim [s^1]_\Phi \sim \dots \sim [s^{l-2}]_\Phi \sim [y']_\Phi$ , in the following way

$$s_i^m = \begin{cases} x_i & \text{if } i \in I \setminus \bigcup_{k \leq m} A_{\varepsilon^k}, \\ y_i & \text{if } i \in \bigcup_{k \leq m} A_{\varepsilon^k}. \end{cases} \quad (16)$$

Now,  $x$  and  $s^0$  satisfy the assumptions of Fact 2.1 over  $A_{\varepsilon^0}$ ; so,  $[x]_\Phi \sim [s^0]_\Phi$ .  $s^0$  and  $s^1$  satisfy the assumptions of Fact 2.1 over  $A_{\varepsilon^1}$ ; so,  $[s^0]_\Phi \sim [s^1]_\Phi$ . Continuing in this way we have that  $s^{l-2}$  and  $y'$  satisfy the assumptions of Fact 2.1 over  $A_{\varepsilon^{l-1}}$ ; so,  $[s^{l-2}]_\Phi \sim [y']_\Phi$ . Thus we have  $[x]_\Phi \sim [y']_\Phi$ .

Let  $K = \{i_j : j < k\}$  be an enumeration. For  $j < k$  the structure  $\mathbb{X}_{i_j}$  is connected so we can choose a path  $\langle \bar{z}_j, \varepsilon_j \rangle$  of length  $n_j$  from  $x_{i_j}$  to  $y_{i_j}$ . By (ii) we have  $F := A \cup K \in \Phi$ . Let  $y'' \in \prod_{i \in I} X_i$  be defined by

$$y''_i = \begin{cases} x_i & \text{if } i \in I \setminus F, \\ y_i & \text{if } i \in F. \end{cases} \quad (17)$$

We define sequences  $t^j = \langle t_i^j : i \in I \rangle \in \prod_{i \in I} X_i$ , for  $j \leq k-2$ , such that  $[y']_\Phi \sim [t^0]_\Phi \sim [t^1]_\Phi \sim \dots \sim [t^{k-2}]_\Phi \sim [y'']_\Phi$ , by

$$t_i^j = \begin{cases} x_i & \text{if } i \in I \setminus (A \cup \{i_0, \dots, i_j\}), \\ y_i & \text{if } i \in A \cup \{i_0, \dots, i_j\}. \end{cases} \quad (18)$$

Namely,  $y'$  and  $t^0$  satisfy the assumptions of Fact 2.1 over  $\{i_0\}$ ; so,  $[y']_\Phi \sim [t^0]_\Phi$ .  $t^0$  and  $t^1$  satisfy the assumptions of Fact 2.1 over  $\{i_1\}$ ; so,  $[t^0]_\Phi \sim [t^1]_\Phi$ . Continuing in this way we have that  $t^{k-2}$  and  $y''$  satisfy the assumptions of Fact 2.1 over  $\{i_{k-1}\}$ ; so  $[t^{k-2}]_\Phi \sim [y'']_\Phi$ . Thus we have proved that  $[y']_\Phi \sim [y'']_\Phi$  and, since  $[x]_\Phi \sim [y']_\Phi$  and  $[y'']_\Phi = [y]_\Phi$ , we finally have  $[x]_\Phi \sim [y]_\Phi$ .  $\square$

**Remark 2.4** Regarding Theorem 2.3, if  $|I| \geq \omega$  and the filter  $\Phi$  contains all cofinite sets, then condition (b) can be replaced by the following simpler condition

(b') There is  $n \in \omega$  such that  $\{i \in I : \mathbb{X}_i \models \varphi_n^{\text{conn}}\} \in \Phi$ .

In particular cases, when the filter  $\Phi$  is minimal (i.e.  $\Phi = \{I\}$ ) and when the filter  $\Phi$  is maximal (an ultrafilter in  $P(I)$ ) by Theorem 2.3 we have the following two corollaries.

**Corollary 2.5** (Direct products and powers) *If (\*) holds and  $\Phi = \{I\}$ , then*

- (a)  $\prod_{i \in I} \mathbb{X}_i$  is connected iff all the structures  $\mathbb{X}_i$  are connected and there is  $n \in \omega$  such that  $\mathbb{X}_i$  is of diameter  $\leq n$ , for all except finitely many  $i \in I$ ;
- (b) The direct power  $\prod_{i \in I} \mathbb{X}$  is connected iff  $\mathbb{X}$  is of finite diameter;
- (c) If  $|I| < \omega$ , then  $\prod_{i \in I} \mathbb{X}_i$  is connected iff all the structures  $\mathbb{X}_i$  are connected.

**Corollary 2.6** (Ultraproducts and ultrapowers) *If (\*) holds,  $\omega \leq |I|$  is less than the first measurable cardinal and  $\mathcal{U} \subset P(I)$  is a non-principal ultrafilter, then*

- (a)  $\prod_{\mathcal{U}} \mathbb{X}_i$  is connected, iff  $\{i \in I : \mathbb{X}_i \models \varphi_n^{\text{conn}}\} \in \mathcal{U}$ , for some  $n \in \omega$ ;
- (b) The ultrapower  $\prod_{\mathcal{U}} \mathbb{X}$  is connected iff  $\mathbb{X}$  is of finite diameter.

The implications  $\Leftarrow$  are true in ZFC (without any assumptions on  $|I|$ ).

**Theorem 2.7** *Let  $|I|$  be a measurable cardinal,  $\mathcal{U}$  a non-principal  $\omega$ -complete ultrafilter on  $I$  and  $\mathbb{X}_i = \langle X_i, \rho_i \rangle$ ,  $i \in I$ ,  $L$ -structures. Then*

$$\prod_{\mathcal{U}} \mathbb{X}_i \text{ is connected} \Leftrightarrow \{i \in I : \mathbb{X}_i \text{ is connected}\} \in \mathcal{U}.$$

**Proof.** ( $\Leftarrow$ ) Let  $U := \{i \in I : \mathbb{X}_i \text{ is connected}\} \in \mathcal{U}$  and let  $x = \langle x_i : i \in I \rangle, y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$ . We construct a path from  $[x]_{\mathcal{U}}$  to  $[y]_{\mathcal{U}}$  in  $\prod_{\mathcal{U}} \mathbb{X}_i$ . For  $i \in U$  there is a path  $\langle \bar{z}^i, \varepsilon^i \rangle$  of length  $n_i + 1$  from  $x_i$  to  $y_i$  and we choose such  $\varepsilon_i \in 2^{n_i+1}$ . Since  $|\bigcup_{n \in \omega} 2^{n+1}| = \omega$ , there are  $\leq \omega$  such choices; let  $\{\varepsilon^k : k < \lambda\}$  be an enumeration of them; so,  $\lambda \leq \omega$ . Then we have  $U = \bigcup_{k < \lambda} A_k$ , where  $A_k := \{i \in U : \varepsilon^i = \varepsilon^k\}$  and, since  $\mathcal{U}$  is  $\omega$ -complete, there is  $k < \lambda$  such that  $A_k \in \mathcal{U}$ . Now, using Fact 2.1 we construct a path from  $[x]_{\mathcal{U}}$  to  $[y]_{\mathcal{U}}$ .

( $\Rightarrow$ ) Let  $U := \{i \in I : \mathbb{X}_i \text{ is not connected}\} \in \mathcal{U}$ . For each  $i \in U$  we pick  $x_i, y_i \in X_i$  such that  $d(x_i, y_i) = \infty$ , for  $i \in I \setminus U$  we take arbitrary  $x_i = y_i \in X_i$  and define  $x = \langle x_i : i \in I \rangle, y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$ . Assuming that there is a path from  $[x]_{\mathcal{U}}$  to  $[y]_{\mathcal{U}}$  in  $\prod_{\mathcal{U}} \mathbb{X}_i$ ,

$$[x]_{\mathcal{U}} \rho^{\varepsilon_0} [z^0]_{\mathcal{U}} \rho^{\varepsilon_1} [z^1]_{\mathcal{U}} \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [z^{n-1}]_{\mathcal{U}} \rho^{\varepsilon_n} [y]_{\mathcal{U}}, \quad (19)$$

there would be  $U_1 \in \mathcal{U}$  which witnesses that and, hence,

$$\forall i \in U \cap U_1 \quad x_i \rho_i^{\varepsilon_0} z_i^0 \rho_i^{\varepsilon_1} z_i^1 \rho_i^{\varepsilon_2} \dots \rho_i^{\varepsilon_{n-1}} z_i^{n-1} \rho_i^{\varepsilon_n} y_i. \quad (20)$$

which is impossible because  $d(x_i, y_i) = \infty$ , for all  $i \in U$ . So, there is no path from  $[x]_{\mathcal{U}}$  to  $[y]_{\mathcal{U}}$  in  $\prod_{\mathcal{U}} \mathbb{X}_i$  and this structure is disconnected.  $\square$

A simple example of a connected  $L$ -structure of infinite diameter is the linear graph  $\mathbb{G}_{\omega} = \langle \omega, \rho \rangle$ , where  $\rho = \{\langle m, n \rangle : |m - n| = 1\}$ . In the absence of measurable cardinals Theorem 2.3 gives a characterization of connected reduced products of  $L$ -structures. In the presence of measurable cardinals we have the following equivalence.



**Theorem 2.8** For each infinite set  $I$  the following conditions are equivalent

- (i) The cardinal  $|I|$  is less than the first measurable cardinal,
- (ii) The implication (a)  $\Rightarrow$  (b) of Theorem 2.3 is true for each filter  $\Phi \subset P(I)$  and each family of  $L$ -structures  $\{\mathbb{X}_i : i \in I\}$ ,
- (iii) The ultraproduct  $\prod_{\mathcal{U}} \mathbb{G}_\omega$  is a disconnected graph for each non-principal ultrafilter  $\mathcal{U} \subset P(I)$ .

**Proof.** The implication (i)  $\Rightarrow$  (ii) is proved in Theorem 2.3.

(ii)  $\Rightarrow$  (iii). Let (ii) hold and let  $\mathcal{U} \subset P(I)$  be a non-principal ultrafilter. Assuming that the ultraproduct  $\prod_{\mathcal{U}} \mathbb{G}_\omega$  is connected we would have a finite set  $K \subset I$  and  $n \in \omega$  such that  $\{i \in I : \mathbb{G}_\omega \models \varphi_n^{\text{conn}}\} \cup K = K \in \Phi$ , which is false.

(iii)  $\Rightarrow$  (i). Let (iii) hold. Suppose that  $|I| \geq \kappa$ , where  $\kappa$  is the first measurable cardinal and w.l.o.g. suppose that  $\kappa \subset I$ . Let  $\mathcal{U}_\kappa \subset P(\kappa)$  be a non-principal  $\omega$ -complete ultrafilter and let  $\mathcal{U} \subset P(I)$  be its extension to  $I$ . Then by Fact 1.1 we have  $\prod_{\mathcal{U}} \mathbb{G}_\omega := (\prod_{i \in I} \mathbb{G}_\omega) / \mathcal{U} \cong (\prod_{i \in \kappa} \mathbb{G}_\omega) / \mathcal{U}_\kappa$  and by Theorem 2.7 the ultraproduct  $(\prod_{i \in \kappa} \mathbb{G}_\omega) / \mathcal{U}_\kappa$  is connected, which is false by (iii).  $\square$

The connectivity relation (and, hence, the components) of a reduced product are described by the following theorem.

**Theorem 2.9** (\*) If  $x = \langle x_i : i \in I \rangle, y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$ , then

$$[x]_\Phi \sim [y]_\Phi \Leftrightarrow \exists n \in \omega \{i \in I : \mathbb{X}_i \models \varphi_{d \leq n+1}[x_i, y_i]\} \in \Phi.$$

**Proof.** ( $\Rightarrow$ ) If  $[x]_\Phi \sim [y]_\Phi$ , then in  $\prod_{\Phi} \mathbb{X}_i$  there is a path from  $[x]_\Phi$  to  $[y]_\Phi$ ; namely, there are  $n \in \omega, \varepsilon \in 2^{n+1}$  and  $z^0, \dots, z^{n-1} \in \prod_{i \in I} X_i$  such that we have  $[x]_\Phi \rho^{\varepsilon_0} [z^0]_\Phi \rho^{\varepsilon_1} [z^1]_\Phi \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [z^{n-1}]_\Phi \rho^{\varepsilon_n} [y]_\Phi$ . Since  $\Phi$  is closed under finite intersections there is  $F \in \Phi$  such that for each  $i \in F$  we have  $x_i \rho_i^{\varepsilon_0} z_i^0 \rho_i^{\varepsilon_1} z_i^1 \rho_i^{\varepsilon_2} \dots \rho_i^{\varepsilon_{n-1}} z_i^{n-1} \rho_i^{\varepsilon_n} y_i$ . So  $F \subset \{i \in I : \mathbb{X}_i \models \varphi_{d \leq n+1}[x_i, y_i]\} \in \Phi$ .

( $\Leftarrow$ ) Let  $n \in \omega$  and  $A := \{i \in I : \mathbb{X}_i \models \varphi_{d \leq n+1}[x_i, y_i]\} \in \Phi$ . If  $i \in A$ , then there is a path  $\langle \bar{z}, \varepsilon \rangle$  of length  $n+1$  from  $x_i$  to  $y_i$ ; thus we can choose one such  $\varepsilon$ . Let  $\varepsilon^k \in 2^{n+1}, k < l$ , be an enumeration of the  $n+1$ -tuples chosen in this way for  $i \in A$ . Then defining  $A_{\varepsilon^k} = \{i \in A : \varepsilon^k \text{ is chosen for } x_i \text{ and } y_i\}$ , for  $k < l$ , we have  $A = \bigcup_{k < l} A_{\varepsilon^k}$ . Let  $y' \in \prod_{i \in I} X_i$ , where  $y' \upharpoonright A = y \upharpoonright A$  and  $y' \upharpoonright (I \setminus A) = x \upharpoonright (I \setminus A)$ . Then  $[y']_\Phi = [y]_\Phi$ . We define sequences  $s^m = \langle s_i^m : i \in I \rangle \in \prod_{i \in I} X_i$ , for  $m \leq l-2$ , such that there are  $l$ -many paths of length  $(n+1)$  witnessing that  $[x]_\Phi \sim [s^0]_\Phi \sim [s^1]_\Phi \sim \dots \sim [s^{l-2}]_\Phi \sim [y']_\Phi$ , by

$$s_i^m = \begin{cases} x_i & \text{if } i \in I \setminus \bigcup_{k \leq m} A_{\varepsilon^k}, \\ y_i & \text{if } i \in \bigcup_{k \leq m} A_{\varepsilon^k}. \end{cases} \quad (21)$$

Now,  $x$  and  $s^0$  satisfy the assumptions of Fact 2.1 over  $A_{\varepsilon^0}$ ; so,  $[x]_\Phi \sim [s^0]_\Phi$ .  $s^0$  and  $s^1$  satisfy the assumptions of Fact 2.1 over  $A_{\varepsilon^1}$ ; so,  $[s^0]_\Phi \sim [s^1]_\Phi$ . Continuing

in this way we have that  $s^{l-2}$  and  $y'$  satisfy the assumptions of Fact 2.1 over  $A_{\varepsilon^{l-1}}$ ; so,  $[s^{l-2}]_{\Phi} \sim [y']_{\Phi}$ . Thus we have  $[x]_{\Phi} \sim [y']_{\Phi} = [y]_{\Phi}$ .  $\square$

**Acknowledgement** This research was supported by the Science Fund of the Republic of Serbia, Program PROMIS, Grant No. 6062228, *Classification of Large Objects–Ultrafilters and Directed Sets–CLOUDS*.

## References

- [1] C. C. Chang, H. J. Keisler, *Model Theory*, North Holland, Amsterdam, 1973.