

Klein-Gordon particles in Gödel-type Som-Raychaudhuri cosmic string spacetime and the phenomenon of spacetime associated degeneracies

Omar Mustafa*

Department of Physics, Eastern Mediterranean University, G. Magusa, north Cyprus, Mersin 10 - Turkey.

Abstract: We argue that only exact, comprehensive, and explicit solutions for the fundamental models, the Klein-Gordon (KG) oscillators and the KG-Coulomb, would help to understand the effects of gravitational fields on the dynamics of such quantum mechanical systems. In the current methodical proposal, the effects of the gravitational fields generated by a Gödel-type Som-Raychaudhuri (SR) cosmic string spacetime on KG-oscillators (KG-particles in general) are studied and reported. In so doing, we revisit the KG-oscillators in a topologically trivial Gödel-type spacetime background and use textbook procedures to report its exact solution that covers even and odd parities. Next, we discuss the drawbacks associated with the power series expansion approach that implies the biconfluent Heun functions/polynomials solution. We, therefore, recollect the so called pseudo perturbative shifted ℓ expansion technique (PSLET) as an alternative and more sophisticated method/technique. Illustrative examples are used: (i) a KG-oscillator in a topologically trivial Gödel-type spacetime, (ii) a quasi-free KG-oscillator in Gödel SR-type cosmic string spacetime, (iii) a KG-Coulombic particle in Gödel SR-type cosmic string spacetime at zero vorticity, and (iv) a massless KG-particle in Gödel SR-type cosmic string spacetime in a Cornell-type Lorentz scalar potential. The corresponding exact energies are obtained from the zeroth (leading) order correction of PSLET, where all higher order correction identically vanish. The comprehensive exactness of the reported solutions manifestly suggest degeneracies associated with spacetime (STAD) phenomenon.

PACS numbers: 05.45.-a, 03.50.Kk, 03.65.-w

Keywords: Klein-Gordon (KG) particles, Gödel-type Som-Raychaudhuri cosmic string spacetime, Spacetime associated degeneracies.

I. INTRODUCTION

Topological defects in spacetime have stimulated intensive research studies in quantum gravity. Amongst are, domain walls [1, 2], cosmic string [3, 4], global monopole [5] and textures [6]. The energy levels of relativistic/non-relativistic quantum particles are shown to be affected by the gravitational fields generated by different spacetime backgrounds with such topological defects, not only in general relativity but also in the geometrical theory of defects in condensed matter physics. In general relativity, for example, the Gödel spacetime metric [7], with an embedded cosmic string, introduces itself as the first cosmological solution to Einstein's equation with rotating matter. Its compact form allowed analytical research studies of many physical and mathematical systems in gravitational backgrounds with rotation and causality violation. It has been shown that all spacetime homogeneous (ST-homogeneous) Gödel-type metrics characterized by the vorticity Ω of spacetime (a real number that represents uniform angular velocity or rotational material, i.e, $\Omega = \pm |\Omega|$ [8]) and a given value of the parameter $\tilde{\mu}$; $-\infty \leq \tilde{\mu}^2 \leq \infty$, can be transformed in cylindrical coordinates [8, 9, 13–21] to

$$ds^2 = - \left(dt + \alpha \Omega \frac{\sinh^2(\tilde{\mu} r)}{\tilde{\mu}^2} d\varphi \right)^2 + \alpha^2 \frac{\sinh^2(2\tilde{\mu} r)}{4\tilde{\mu}^2} d\varphi^2 + dr^2 + dz^2. \quad (1)$$

Where, for $\tilde{\mu}^2 < 0$ there is an infinite number of successive causal and noncausal regions, and for $0 \leq \tilde{\mu}^2 < \Omega^2$ there is one noncausal region for a given $r > r_c$. Here, r_c is the critical radius determined by $g_{\varphi\varphi} = 0$ (e.g., [8]) to imply

$$\tanh(\tilde{\mu} r_c) = \frac{\tilde{\mu}}{|\Omega|} \implies r_c = \frac{1}{\tilde{\mu}} \operatorname{arctanh} \left(\frac{\tilde{\mu}}{|\Omega|} \right).$$

For $\tilde{\mu}^2 = \Omega^2$ the Gödel spacetime metric (the first exact Gödel solution of the Einstein equation describing a complete causal, ST-homogeneous rotating universe [13]) is recovered, and for $\tilde{\mu} < |\Omega|$ there is a causal region (classified as

*Electronic address: omar.mustafa@emu.edu.tr

chronologically safe) with radius r_c surrounded by the outer noncausal space [8] Nevertheless, at the limit $\tilde{\mu} \rightarrow 0$ of the Gödel spacetime metric (1) we obtain the ST-homogeneous Som-Raychaudhuri (SR) solution

$$ds^2 = - (dt + \alpha \Omega r^2 d\varphi)^2 + \alpha^2 r^2 d\varphi^2 + dr^2 + dz^2 \quad (2)$$

of the Einstein field equations [9–12], in the presence of cosmic string. Here, the disclination parameter α (curvature parameter) admits the values $0 < \alpha < 1$ in general relativity (for cosmic strings with positive curvature), $\alpha > 1$ in the geometric theory of defects in condensed matter (for a negative curvature), and $\Omega = 0$ and $\alpha = 1$ corresponds to Minkowski flat spacetime metric. Moreover, the covariant and contravariant metric tensors associated with the Som-Raychaudhuri spacetime are, respectively, given by

$$g_{\mu\nu} = \begin{pmatrix} -1 & 0 & -\alpha\Omega r^2 & 0 \\ 0 & 1 & 0 & 0 \\ -\alpha\Omega r^2 & 0 & \alpha^2 r^2 (1 - \Omega^2 r^2) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \iff g^{\mu\nu} = \begin{pmatrix} (\Omega^2 r^2 - 1) & 0 & -\frac{\Omega}{\alpha} & 0 \\ 0 & 1 & 0 & 0 \\ -\frac{\Omega}{\alpha} & 0 & \frac{1}{\alpha^2 r^2} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}; \det(g) = -\alpha^2 r^2. \quad (3)$$

The objective of the current methodical proposal is to explore in a comprehensive manner the effect of the gravitational field generated by the Gödel-type cosmic string SR-spacetime (2) on the spectra of some Klein-Gordon (KG) particles of fundamental quantum gravity nature, like the KG-oscillators and KG-Coulomb models. Such fundamental models have been studied by different authors (e.g., [15–24]) where the vorticity parameter Ω is assumed to take the form $\Omega \geq 0$ and the energies are assumed to be positive $E = +|E|$ or $E = \pm|E|$ where all nodeless states are lost in the process (excluding Hassanabadi et al [20, 21] who have included all nodeless states with the radial quantum number $n_r = 0$). However, in Duarte de Oliveira et al [8], Ω is a real number that denotes uniform angular velocity or rotation of material (i.e., $\Omega = \pm|\Omega|$). In the current methodical proposal, we discuss KG-particles' spectral properties under the influence of the gravitational field generated by the Gödel-type cosmic string SR-spacetime (2) taking into account $E = \pm|E|$, $\Omega = \pm|\Omega|$, and including nodeless states. In so doing, nevertheless, it is unavoidably inviting to reconsider and report on the KG-oscillators in a topologically trivial Gödel-type spacetime metric, used by Ahmed [18],

$$ds^2 = -dt^2 + dx^2 + (1 - \alpha^2 x^2) dy^2 - 2\alpha x dt dy + dz^2, \quad (4)$$

where $H(x) = \alpha x$ and $D(x) = 1$ in the general Gödel metric [7, 13]

$$ds^2 = - (dt + H(x) dy)^2 + dx^2 + D(x)^2 dy^2 + dz^2,$$

using the coordinates $(x^0 = t, x^1 = x, x^2 = y, x^3 = z; x, y, z \in (-\infty, \infty))$. Where, the covariant and contravariant elements of the spacetime metric tensors are given by

$$\begin{aligned} g_{00} &= -1, g_{11} = g_{33} = 1, g_{02} = g^{20} = -\alpha x, g_{22} = 1 - \alpha^2 x^2, \\ g^{00} &= \alpha^2 x^2 - 1, g^{11} = g^{22} = g^{33} = 1, \det(g) = -1. \end{aligned} \quad (5)$$

The organization of our methodical proposal is in order. We start with the KG-oscillators in a topologically trivial Gödel-type spacetime (4) considered by Ahmed [18], who has reported only the positive energy states and missed not only the ground state but also the negative energy states. For the sake of scientific correctness, we give this problem a proper textbook quantum mechanical treatment [25] and report general forms of the spectra, in section 2, that include all positive and negative energy states. In section 3, we discuss KG-particles in Gödel SR-type cosmic string spacetime in a Cornell-type Lorentz scalar potentials and show the disadvantages of using a power series (Frobinous method) expansion method that leads to biconfluent Heun functions/polynomials. In section 4, we recollect, as a powerful alternative method, the so called pseudo-perturbative shifted- ℓ expansion technique (PSLET), to solve for Schrödinger equation (a common form that relativistic wave equations collapse into) in D -dimensions [26–33] to serves our current methodical proposal. PSLET has shown, with a brut force evidence through comparison with numerical integration techniques, to provide very highly accurate results (even with its non-Hermitian \mathcal{PT} -symmetric version by Mustafa and Znojil [34, 35]). We use some illustrative examples, in section 5, that include a KG-oscillator (of section 2) in a topologically trivial Gödel-type spacetime (4), a quasi-free KG-oscillator in Gödel SR-type cosmic string spacetime (2), a KG-Coulombic particle in Gödel SR-type cosmic string spacetime (2) at zero vorticity (i.e., $\Omega = 0$), and a massless KG-particles in Gödel SR-type cosmic string spacetime in a Cornell-type Lorentz scalar potential [36]. We conclude in section 6.

II. KG-OSCILLATORS IN A TOPOLOGICALLY TRIVIAL GÖDEL-TYPE SPACETIME BACKGROUND

A. KG-oscillator I as a topologically trivial Gödel-type spacetime byproduct:

In this section we recollect the topologically trivial Gödel-type spacetime metric (4), along with the metric tensor elements in (5) and $\det(g) = -1$, used by Ahmed [18]. Then, a KG-particle of rest mass energy m (denoting mc^2 in $c = \hbar = 1$ units) in such a Gödel-type spacetime is described by the KG-equation

$$\frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu \Psi) = m^2 \Psi. \quad (6)$$

Consequently, with

$$\Psi(t, x, y, z) = \exp(i[k_y y + k_z z - Et]) \phi(x), \quad (7)$$

it yields

$$\phi''(x) - (\gamma^2 x^2 + 2\gamma k_y x) \phi(x) = \tilde{\beta} \phi(x); \quad \tilde{\beta} = m^2 + k_y^2 + k_z^2 - E^2, \quad \gamma^2 = \alpha^2 E^2. \quad (8)$$

At this point, one should notice that $\gamma = \pm\alpha|E|$ as mandated by the very nature of the Gödel-type spacetime in (4). Obviously, this equation resembles the well known one-dimensional Schrödinger equation for the so called shifted-oscillator (but does not inherit its parametric characterizations) that can be treated, in a straightforward textbook manner, by rewriting (8) as

$$\phi''(x) - (\gamma x + k_y)^2 \phi(x) = \tilde{\beta} \phi(x); \quad \tilde{\beta} = \tilde{\beta} - k_y^2 = m^2 + k_z^2 - E^2. \quad (9)$$

One would then use the change of variable $u = \gamma x + k_y$ to obtain

$$\phi''(u) - \tilde{\gamma}^2 u^2 \phi(u) + \lambda \phi(u) = 0; \quad \lambda = -\frac{\tilde{\beta}}{\gamma^2}, \quad \tilde{\gamma}^2 = \frac{1}{\gamma^2}. \quad (10)$$

This is the one-dimensional Schrödinger-like harmonic oscillator form that admits exact textbook solution with the eigenvalues and eigenfunctions, respectively, given by

$$\lambda_n = |\tilde{\gamma}|(2n+1), \quad \text{and} \quad \phi(u) = \mathcal{N}_n e^{-\frac{|\tilde{\gamma}|u^2}{2}} H_n(\sqrt{|\tilde{\gamma}|}u); \quad n = 0, 1, 2, \dots, \quad (11)$$

where $H_n(\sqrt{|\tilde{\gamma}|}u)$ are the Hermite polynomials and \mathcal{N}_n are the normalization constants. The finiteness and square integrability of the wave function enforces the quantum mechanical condition that $|\tilde{\gamma}| = \pm\tilde{\gamma}_\pm \geq 0$ (i.e., $\tilde{\gamma} = \tilde{\gamma}_\pm$; $\tilde{\gamma}_+ = 1/\alpha E_+$, $\tilde{\gamma}_- = 1/\alpha E_- \implies |\tilde{\gamma}| = \pm\tilde{\gamma}_\pm$, where $E_\pm = \pm|E|$). Consequently, with the parametric substitutions above (8), one obtains

$$\frac{m^2 + k_z^2 - E_\pm^2}{\alpha^2 E_\pm^2} = -|\tilde{\gamma}|(2n+1) \implies E_\pm^2 \mp \alpha E_\pm(2n+1) - (m^2 + k_z^2) = 0; \quad n = 0, 1, 2, \dots \quad (12)$$

Hence,

$$E_n = E_\pm = \pm\alpha \left(n + \frac{1}{2}\right) \pm \sqrt{\alpha^2 \left(n + \frac{1}{2}\right)^2 + m^2 + k_z^2}. \quad (13)$$

Which in turn implies

$$E_+ = +\alpha \left(n + \frac{1}{2}\right) + \sqrt{\alpha^2 \left(n + \frac{1}{2}\right)^2 + m^2 + k_z^2} \quad (14)$$

as the positive energy solution, and

$$E_- = -\alpha \left(n + \frac{1}{2}\right) - \sqrt{\alpha^2 \left(n + \frac{1}{2}\right)^2 + m^2 + k_z^2} \quad (15)$$

for the negative energy solution. Notably, unlike the solution reported by Ahmed [18] (see equation (58) in [18]), the textbook solution above includes both even and odd parities, positive and negative energies, as well as the ground state [25].

B. KG-oscillator II as a topologically trivial Gödel-type spacetime and momentum operator deformation byproduct

Using similar recipe as that of Moshinsky and Szczepaniak [37] and Mirza and Mohadesi [38] for the Dirac and KG oscillators, we redefine the momentum operator as

$$p_\mu \longrightarrow p_\mu + i\eta\chi_\mu; \quad \chi_\mu = (0, x, 0, 0), \quad (16)$$

where the dimensions of the second term suits the dimensions of the momentum operator p_μ and here we consider η to have positive and/or negative values. Then the KG-equation reads

$$\frac{1}{\sqrt{-g}} (\partial_\mu + \eta\chi_\mu) \sqrt{-g}g^{\mu\nu} (\partial_\nu - \eta\chi_\nu) \Psi = m^2\Psi. \quad (17)$$

At this point, one may use $\eta = m\omega \geq 0$ (with m denoting rest mass of the KG-particle) to recover the traditionally used values as in (e.g., [18, 22–24, 37, 38] and other related references cited therein). However, we shall use a general parameter η and allow it to take positive and/or negative values. Yet, in this case we avoid eminent confusion and inconsistency between m^2 (denoting $m^2c^2 = mmc^2$, rest mass multiplied by rest mass energy) on the R.H.S. and the rest mass of the particle on the L.H.S. of the KG-equation (17) for $\eta = m\omega$. This point is made clear by Moshinsky and Szczepaniak [37] and Mirza and Mohadesi [38] while dealing with the Dirac and KG oscillators. We therefore stick with our assumption and use the spacetime metric tensor elements in (5), to recast (17) as

$$\phi''(x) - (\tilde{\omega}^2 x^2 + 2\gamma k_y x) \phi(x) = \zeta \phi(x); \quad \zeta = m^2 + k_y^2 + k_z^2 - E^2 + \eta, \quad \tilde{\omega}^2 = \gamma^2 + \eta^2, \quad \gamma = \alpha E. \quad (18)$$

Hereby, one should be aware that $\eta^2 x^2$ would take the form of $\eta^2 x^2 = m(m\omega^2 x^2)$ as in [37, 38] when the speed of light is kept intact in Dirac and/or KG equations (i.e., consistent with $m^2 c^2 = m(mc^2)$ on the R.H.S. of (18)). The source of confusion is therefore clear and should be avoided. Moreover, equation (18) admits solution in the form of hypergeometric functions [25] given by

$$\phi(x) = e^{-\frac{|\tilde{\omega}|x^2}{2} - \frac{\delta x}{|\tilde{\omega}|}} \left[A {}_1F_1 \left(\tilde{a}, \frac{1}{2}, \frac{(\tilde{\omega}^2 x + \delta)^2}{|\tilde{\omega}|^3} \right) + B (\tilde{\omega}^2 x + \delta) {}_1F_1 \left(\tilde{b}, \frac{3}{2}, \frac{(\tilde{\omega}^2 x + \delta)^2}{|\tilde{\omega}|^3} \right) \right], \quad (19)$$

where $\delta = \gamma k_y = \alpha E k_y$, $|\tilde{\omega}| \geq 0$ to secure finiteness and square integrability of the wave function, and

$$\tilde{b} = \frac{3}{4} - \frac{\delta^2}{4|\tilde{\omega}|^3} + \frac{\zeta}{4|\tilde{\omega}|} = \tilde{a} + \frac{1}{2}. \quad (20)$$

Again, the condition for the hypergeometric functions to become polynomials of degree $n \geq 0$ suggests that $a = -n$ and $b = a + 1/2 = -n$. This would imply that

$$\tilde{a} = \frac{1}{4} - \frac{\delta^2}{4|\tilde{\omega}|^3} + \frac{\zeta}{4|\tilde{\omega}|} = -n \implies \zeta_{2n} = -|\tilde{\omega}|(4n+1) + \frac{\delta^2}{\tilde{\omega}^2}, \quad (21)$$

for even parity, and

$$\tilde{b} = -n \implies \zeta_{2n+1} = -|\tilde{\omega}|(4n+3) + \frac{\delta^2}{\tilde{\omega}^2}. \quad (22)$$

for odd parity solutions. The two results (21) and (22) would reduce to a common form

$$\zeta_n = -|\tilde{\omega}|(2n+1) + \frac{\delta^2}{\tilde{\omega}^2}; \quad |\tilde{\omega}| = \pm\alpha E_\pm \sqrt{1 + \frac{\eta^2}{\alpha^2 E_\pm^2}} \geq 0, \quad (23)$$

to imply that

$$E_\pm^2 - (m^2 + k_y^2 + k_z^2 + \eta) = \pm\alpha E_\pm \sqrt{1 + \frac{\eta^2}{\alpha^2 E_\pm^2}} (2n+1) - \frac{\alpha^2 E_\pm^2 k_y^2}{\alpha^2 E_\pm^2 + \eta^2}. \quad (24)$$

Obviously, this result collapses into that in (13) for $\eta = 0$, where the effect of even and odd parities is reflected on the related spectrum.

III. KG-PARTICLES IN A 4-VECTOR AND SCALAR LORENTZ POTENTIALS IN GÖDEL SR-TYPE COSMIC STRING SPACETIME BACKGROUND

The KG-equation for a spin-0 particle in a 4-vector A_μ and a radial scalar $S(r)$ potentials in the Gödel SR-type spacetime background (2) is given by

$$\frac{1}{\sqrt{-g}} D_\mu (\sqrt{-g} g^{\mu\nu} D_\nu \Psi) = (m + S(r))^2 \Psi, \quad (25)$$

where the gauge-covariant derivative is given by $D_\mu = \partial_\mu - ieA_\mu$. This would in turn allow us to cast equation (25) as

$$\left\{ -D_t^2 + \frac{1}{r} D_r (r D_r) + \left(\Omega r D_t - \frac{1}{\alpha r} D_\varphi \right)^2 + D_z^2 - (m + S(r))^2 \right\} \Psi = 0. \quad (26)$$

At this point, we may now define the corresponding gauge-covariant derivatives so that

$$D_t = \partial_t - ieA_t = \partial_t - iV(r), \quad D_r = \partial_r, \quad D_\varphi = \partial_\varphi - ieA_\varphi, \quad D_z = \partial_z, \quad (27)$$

where, $V(r) = eA_t$ is the Lorentz 4-vector potential (i.e., transforms within the 4-vector potential A_μ), $S(r)$ is the Lorentz scalar potential (i.e., transforms like the rest mass energy $m \rightarrow m + S(r)$), and eA_φ may include both magnetic and Aharonov-Bohm flux fields effects [40, 41]. We now use the assumption that

$$\Psi(t, r, \varphi, z) = \exp(i[\ell\varphi + k_z z - Et]) \psi(r) = \exp(i[\ell\varphi + k_z z - Et]) \frac{R(r)}{\sqrt{r}} \quad (28)$$

in (26) to obtain

$$\left\{ \partial_r^2 + \frac{1}{4r^2} + (E + V(r))^2 - \left[\Omega r (E + V(r)) + \frac{(\ell - eA_\varphi)}{\alpha r} \right]^2 - k_z^2 - (m + S(r))^2 \right\} R(r) = 0, \quad (29)$$

This result would describe KG-particles in Gödel SR-type cosmic string spacetime background.

In what follows, we shall be interested in a set of KG-particles in Gödel SR-type cosmic string spacetime under the influence of $A_\varphi = 0$ (i.e., no magnetic or Aharonov-Bohm flux fields effects), and no Lorentz 4-vector potential (i.e., $V(r) = 0$). Under such settings, the KG-equation (29) reduces to

$$\left\{ \partial_r^2 - \frac{(\tilde{\ell}^2 - 1/4)}{r^2} - \Omega^2 E^2 r^2 - 2mS(r) - S(r)^2 + \tilde{\lambda} \right\} R(r) = 0; \quad \tilde{\ell} = \frac{\ell}{\alpha}, \quad \tilde{\lambda} = E^2 - 2\tilde{\ell}\Omega E - k_z^2 - m^2. \quad (30)$$

Obviously, this equation resembles the one-dimensional form of the two-dimensional radial Schrödinger equation. Which, with a Cornell-type Lorentz scalar potential $S(r) = ar + b/r$ would result

$$\left\{ \partial_r^2 - \frac{(\mathcal{L}^2 - 1/4)}{r^2} - \tilde{\Omega}^2 r^2 - 2mar - \frac{2mb}{r} + \Lambda \right\} R(r) = 0, \quad (31)$$

where

$$\tilde{\Omega}^2 = \Omega^2 E^2 + a^2, \quad \mathcal{L}^2 = \tilde{\ell}^2 + b^2, \quad \Lambda = E^2 - 2\tilde{\ell}\Omega E - (k_z^2 + m^2 + 2ab). \quad (32)$$

Equation (31) is known to admit, using power series method, a solution in the form of biconfluent Heun functions

$$R(r) = .C r^{|\mathcal{L}|+1/2} \exp\left(-\frac{|\tilde{\Omega}|}{2} r^2 - \frac{am}{|\tilde{\Omega}|} r\right) H_B\left(\alpha, \beta, \gamma, \delta, \sqrt{|\tilde{\Omega}|} r\right). \quad (33)$$

where $\alpha = 2|\mathcal{L}|$, $\beta = \frac{2am}{|\tilde{\Omega}|^{3/2}}$, $\gamma = \frac{a^2 m^2 + \Lambda \tilde{\Omega}^2}{|\tilde{\Omega}|^3}$, and $\delta = \frac{4mb}{\sqrt{|\tilde{\Omega}|}}$. For the case $a = 0 = b$, this solution implies

$$\begin{aligned} R(r) &= .C r^{|\tilde{\ell}|+1/2} \exp\left(-\frac{|\Omega E|}{2} r^2\right) H_B\left(2|\tilde{\ell}|, 0, \frac{\Lambda}{|\Omega E|}, 0, \sqrt{|\Omega E|} r\right) \\ &= .C r^{|\tilde{\ell}|+1/2} \exp\left(-\frac{|\Omega E|}{2} r^2\right) {}_1F_1\left(\left[\frac{1}{2} + \frac{\dot{\alpha}}{4} - \frac{\dot{\gamma}}{4}\right], \left[1 + \frac{\dot{\alpha}}{2}\right], |\Omega E| r^2\right), \end{aligned} \quad (34)$$

where the condition that

$$\frac{1}{2} + \frac{\dot{\alpha}}{4} - \frac{\dot{\gamma}}{4} = \frac{1}{2} + \frac{|\tilde{\ell}|}{2} - \frac{\Lambda}{4|\Omega E|} = -n_r \quad (35)$$

is required for the confluent hypergeometric function to become a polynomial of degree $n_r \geq 0$ and to secure the finiteness and square integrability of the wave function. This would immediately result, in terms of the associated Laguerre polynomials $L_{n_r}^{|\tilde{\ell}|}(z^2)$, that

$$\Lambda = 2|\Omega E| \left(2n_r + |\tilde{\ell}| + 1 \right), \text{ and } R(r) = .C r^{|\tilde{\ell}|+1/2} \exp\left(-\frac{|\Omega E|}{2} r^2\right) L_{n_r}^{|\tilde{\ell}|}(|\Omega E| r^2) \quad (36)$$

as the exact eigen energies and eigen functions for the two dimensional Schrödinger harmonic oscillator. Moreover, the same condition (35) should also be imposed on the biconfluent Heun function (33) so that a biconfluent Heun polynomial [39] of degree $n = 2n_r \geq 0$ is obtained (c.f., e.g., [23, 40, 42, 43]) to imply

$$\Lambda = 2|\Omega E| \sqrt{1 + \frac{a^2}{\Omega^2 E^2}} (2n_r + |\mathcal{L}| + 1), \quad (37)$$

and

$$E^2 - 2\tilde{\ell}\Omega E - (k_z^2 + m^2 + 2ab) = 2|\Omega E| \sqrt{1 + \frac{a^2}{\Omega^2 E^2}} (2n_r + |\mathcal{L}| + 1). \quad (38)$$

Where the detailed analysis of such a quadratic equation are given in section 5-B, taking into account that

$$|\Omega E| = \left\{ \begin{array}{l} +\Omega_{\pm} E_{\pm} \\ -\Omega_{\mp} E_{\pm} \end{array} \right\} \geq 0; \quad E_{\pm} = \pm |E|, \quad \Omega_{\pm} = \pm |\Omega|, \quad (39)$$

At this point, the truncation order n of a power series (biconfluent Heun function here) does not make it a valid quantum number, but rather it should be correlated to the well known quantum numbers as done above (i.e., $n = 2n_r \geq 0$). Hereby, one should give credentials for Neto and co-workers [23] as they resemble a group of the very few who have correctly used condition (35) on the biconfluent Heun polynomials. Only under such condition that the result of (38) would recover those of (36) for $a = 0 = b$. Obviously, however, for the case $a = 0 = \Omega$ the solution in (38) collapses into a free relativistic particle eigenvalues

$$E^2 - (k_z^2 + m^2) = 0, \quad (40)$$

although we have an effective two-dimensional Coulomb problem (i.e., a KG-Coulombic particle). This is a clear drawback of the Heun power series (Frobinous method) expansion approach. Moreover, if condition (35) is not implemented then the biconfluent Heun polynomial solution would fail to address the exact solution for the KG-oscillator for $a = 0 = b$. One should, therefore, search for an alternative method to deal with such a case in (31).

As a more reliable alternative method, to the power series (Frobinous method) expansion, we recollect the so called pseudo-perturbative shifted- ℓ expansion technique (PSLET) for the D -dimensional radial Schrödinger-type equation and solve for different KG-particles' settings in the Gödel SR-type cosmic string spacetime backgrounds of (31). In order to make our proposal self contained, we, in short, discuss PSLET in the following section. We shall also use Ahmed's model (10) [18] as one of the illustrative examples to be used in the current methodical proposal.

IV. PSLET FOR THE D -DIMENSIONAL SCHRÖDINGER-TYPE EQUATION

Let us recollect that the D -dimensional radial Schrödinger equation ($\hbar = 2m = 1$ units are used in this section) reads

$$\left\{ -\partial_r^2 + \frac{\ell_D (\ell_D + 1)}{r^2} + V(r) \right\} \Phi_{k,\ell}(r) = E_{k,\ell} \Phi_{k,\ell}(r), \quad (41)$$

where $\ell_D = \ell + (D - 3)/2$ to incorporate interdimensional degeneracies associated with the isomorphism between angular momentum and the dimensionality D . Hereby, one should notice that for $D = 1$, $\ell_1 = \ell - 1$ and hence

$\ell_D(\ell_D + 1) = \ell(\ell - 1) = 0$ for $\ell = 0$ and $\ell = 1$ to denote even and odd parity, respectively, for the one-dimensional case with $r = x \in (-\infty, \infty)$. In the two dimensional case with radially cylindrical symmetry, $r = \sqrt{x^2 + y^2} \in (0, \infty)$ and $\ell_2 = |\ell| - 1/2$ with ℓ denoting the magnetic quantum number. Moreover, in the three-dimensional radially spherical symmetry, $r = \sqrt{x^2 + y^2 + z^2} \in (0, \infty)$ and $\ell_3 = \ell$ where ℓ denotes angular momentum quantum number. Since we are going to use $1/\ell_D$ as a small perturbation expansion parameter at large- ℓ_D limit (i.e., pseudo-classical limit, e.g., [26–34]) we avoid the trivial case for $\ell_D = 0$ (for $D = 3$ and $\ell = 0$) and use a shift β so that $\check{\ell} = \ell_D - \beta$. Under such settings, our Schrödinger equation in (41) reads

$$\left\{ -\partial_r^2 + \frac{\check{\ell}^2 + \check{\ell}(2\beta + 1) + \beta(\beta + 1)}{r^2} + V(r) \right\} \Phi_{k,\ell}(r) = E_{k,\ell} \Phi_{k,\ell}(r). \quad (42)$$

For convenience, we now shift the origin of the coordinate system and use

$$x = \frac{\check{\ell}^{1/2}}{r_0} (r - r_0), \quad (43)$$

and expand about $x = 0$ to obtain

$$\left(\frac{x}{\check{\ell}^{1/2}} + 1 \right)^{-2} = 1 - \frac{2x}{\check{\ell}^{1/2}} + \frac{3x^2}{\check{\ell}} + \frac{4x^3}{\check{\ell}^{3/2}} + \dots, \quad (44)$$

$$V(x(r)) = \frac{\check{\ell}^2}{Q} \left[V(r_0) + V'(r_0) \frac{r_0 x}{\check{\ell}^{1/2}} + V''(r_0) \frac{r_0^2 x^2}{2\check{\ell}} + V'''(r_0) \frac{r_0^3 x^3}{6\check{\ell}^{3/2}} + \dots \right]. \quad (45)$$

Moreover, let us expand the energy so that

$$E_{k,\ell} = \frac{\check{\ell}^2}{Q} \left[E_0 + \frac{E_1}{\check{\ell}} + \frac{E_2}{\check{\ell}^2} + \frac{E_3}{\check{\ell}^3} + \dots \right], \quad (46)$$

where Q is a constant parameter that scales the potential at large- ℓ_D limit and is set equal to $\check{\ell}^2$ at the end of the calculation. Consequently, we may now re-write (42) as

$$\left\{ -\partial_x^2 + \left[\check{\ell} + (2\beta + 1) + \frac{\beta(\beta + 1)}{\check{\ell}} \right] \left(1 - \frac{2x}{\check{\ell}^{1/2}} + \frac{3x^2}{\check{\ell}} + \frac{4x^3}{\check{\ell}^{3/2}} + \dots \right) + \frac{r_0^2 \check{\ell}}{Q} \left[V(r_0) + V'(r_0) \frac{r_0 x}{\check{\ell}^{1/2}} + V''(r_0) \frac{r_0^2 x^2}{2\check{\ell}} + \dots \right] \right\} \Phi_k(x(r)) = \xi_k \Phi_k(x(r)) \quad (47)$$

with

$$\xi_k = \frac{r_0^2 \check{\ell}}{Q} \left[E_0 + \frac{E_1}{\check{\ell}} + \frac{E_2}{\check{\ell}^2} + \frac{E_3}{\check{\ell}^3} + \dots \right]. \quad (48)$$

This equation is to be compared with that of the one-dimensional anharmonic oscillator

$$\left[-\partial_x^2 + \frac{1}{4} \omega^2 x^2 + \varepsilon_0 + P(x) \right] \chi_k(x) = \lambda_k \chi_k(x) \quad (49)$$

with $P(x)$ is a perturbation given by

$$P(x) = \check{\ell}^{-1/2} [\varepsilon_1 x + \varepsilon_3 x^3] + \check{\ell}^{-1} [\varepsilon_2 x^2 + \varepsilon_4 x^4] + \check{\ell}^{-3/2} [\delta_1 x + \delta_3 x^3 + \delta_5 x^5] + \check{\ell}^{-2} [\delta_2 x^2 + \delta_4 x^4 + \delta_6 x^6]. \quad (50)$$

It is obvious that equation (49) represents the one-dimensional Schrödinger anharmonic oscillator that has been readily discussed by Imbo et al. [27, 28] and by Mustafa and Barakat [29] and reported to admit the eigenvalues

$$\lambda_k = \varepsilon_0 + \left(k + \frac{1}{2} \right) \omega + \frac{\alpha_1}{\check{\ell}} + \frac{\alpha_2}{\check{\ell}^2}. \quad (51)$$

where α_1 and α_2 are given in the Appendix, and $(k + \frac{1}{2}) \omega$ are the energies of the unperturbed one-dimensional Schrödinger oscillator with $k = 0, 1, 2, \dots$ that denotes the number of nodes in the wave function (for $D = 2, 3$ it denotes the radial quantum number or if you wish, the number of nodes in the wave function). A comparison between (47) and (49) would imply that

$$\lambda_k = \check{\ell} \left(1 + \frac{r_o^2}{Q} V(r_o) \right) + \left[2\beta + 1 + \left(k + \frac{1}{2} \right) \omega \right] + \frac{1}{\check{\ell}} [\beta(\beta + 1) + \alpha_1] + \frac{\alpha_2}{\check{\ell}^2} + \dots \quad (52)$$

We may now simply compare (48) with (52) to obtain

$$E_0 = \frac{Q}{r_o^2} + V(r_o), \quad E_1 = \frac{Q}{r_o^2} \left[2\beta + 1 + \left(k + \frac{1}{2} \right) \omega \right], \quad E_2 = \frac{Q}{r_o^2} [\beta(\beta + 1) + \alpha_1], \quad E_3 = \frac{Q}{r_o^2} \alpha_2, \quad (53)$$

where r_o is chosen to minimize the zeroth order term so that it satisfies the relations

$$\frac{dE_0}{dr_o} = 0, \quad \frac{d^2 E_0}{dr_o^2} > 0 \implies (\ell_D - \beta) = \sqrt{\frac{r_o^3 V'(r_o)}{2}}. \quad (54)$$

and the shifting parameter β is determined through the requirement that the first order term vanishes. This would result in

$$E_1 = 0 \implies \beta = -\frac{1}{2} \left(k + \frac{1}{2} \right) \omega - \frac{1}{2}, \quad (55)$$

with

$$\omega = 2 \sqrt{3 + \frac{r_o V''(r_o)}{V'(r_o)}}. \quad (56)$$

One would therefore recast our anharmonic oscillator energies of (46) as

$$E_k = E_0 + \frac{1}{r_o^2} [\beta(\beta + 1) + \alpha_1] + \frac{\alpha_2}{\check{\ell} r_o^2}. \quad (57)$$

Now we recall the corresponding Schrödinger potential for our KG-particles in the Gödel SR-type cosmic string spacetime with a Cornell-type Lorentz scalar interaction of (31). This would result, using (56), (55) and (54), that

$$\omega = 2 \sqrt{\frac{4\tilde{\Omega}^2 r_o^3 + 3amr_o^2 - bm}{\tilde{\Omega}^2 r_o^3 + amr_o^2 - bm}}, \quad \beta = -\frac{1}{2} \left(k + \frac{1}{2} \right) \omega - \frac{1}{2}, \quad \tilde{\Omega} = \Omega E \quad (58)$$

and

$$\check{\ell} = (\ell_D - \beta) = \sqrt{\tilde{\Omega}^2 r_o^4 + amr_o^4 - bmr_o^2}; \quad \ell_D = \ell + (D - 3)/2. \quad (59)$$

Various especial cases of such settings are given in the following illustrative examples.

V. ILLUSTRATIVE EXAMPLES

In this section, we use PSLET and consider the following illustrative examples.

A. KG-oscillator I as a topologically trivial Gödel-type spacetime byproduct of (10)

We hereby recall the KG-oscillator I as a trivial Gödel-type spacetime of section 2 described by (10) as

$$\phi''(u) - \tilde{\gamma}^2 u^2 \phi(u) = -\lambda \phi(u); \quad \lambda = -\frac{\tilde{\beta}}{\gamma^2}, \quad \tilde{\gamma}^2 = \frac{1}{\gamma^2} \cdot u = \gamma x + k_y. \quad (60)$$

Obviously, the central repulsive/attractive core $\ell_D (\ell_D + 1) / r^2$ in the D -dimensional Schrödinger equation (41) must vanish to resemble the one dimensional form of Schrödinger equation (60). In this case, $D = 1 \implies \ell_D (\ell_D + 1) = \ell (\ell - 1) = 0$ for $\ell = 0$ and $\ell = 1$ to denote even and odd parities $(-1)^\ell$, respectively. Moreover, $u \equiv r \in (-\infty, \infty)$ and u_o is determined by (54) to read $u_o = \sqrt{\check{\ell}/|\check{\gamma}|} \in \mathbb{R}$; $\check{\ell} = \ell_D - \beta = \ell - 1 - \beta$. This would in turn imply that $\omega = 4$ by (56) and $\beta = -2k - 3/2$ by (55). Consequently, equations (53) would yield the zeroth-order correction

$$E_0 = 2 |\check{\gamma}| \left(2k + \ell + \frac{1}{2} \right) = \left\{ \begin{array}{l} 2 |\check{\gamma}| \left(2k + \frac{1}{2} \right) ; \text{ for even parity } \ell = 0 \\ 2 |\check{\gamma}| \left(2k + \frac{3}{2} \right) ; \text{ for odd parity } \ell = 1 \end{array} \right\}, \quad (61)$$

which results in a common form

$$E_0 = 2 |\check{\gamma}| \left(n + \frac{1}{2} \right); \quad n = 0, 1, 2, \dots, \quad (62)$$

with E_1, E_2 , and E_3 identically vanish. Therefore, the energies associated with the shifted Schrödinger oscillator are

$$\lambda_n = 2 |\check{\gamma}| \left(n + \frac{1}{2} \right); \quad n = 0, 1, 2, \dots. \quad (63)$$

Consequently, the KG-oscillator I above admits the exact energies reported in (13) and goes through the same procedure discussed in section 2-A.

B. Quasi-free KG-oscillator in Gödel SR-type cosmic string spacetime (2)

Next, we consider a KG-particle in Gödel SR-type cosmic string spacetime background described by (29) with the assumption that $V(r) = S(r) = 0$ (no interaction potential is involved, hence the notion "quasi-free KG-particle" is introduced). In this case one would rewrite (29) as

$$\left\{ \partial_r^2 - \frac{(\check{\ell}^2 - 1/4)}{r^2} - \check{\Omega}^2 r^2 + \check{\lambda} \right\} R(r) = 0, \quad (64)$$

where

$$\check{\lambda} = E^2 - 2 \check{\ell} \Omega E - (k_z^2 + m^2); \quad \check{\ell} = \frac{\ell}{\alpha}, \quad \check{\Omega}^2 = \Omega^2 E^2. \quad (65)$$

Which resembles the two-dimensional radial Schrödinger-oscillator with an irrational angular frequency $\check{\Omega} = \pm |\Omega E|$ (now angular velocity). Under such settings, our PSLET implies that for $D = 2 \implies \ell_D = |\check{\ell}| - 1/2$ (with $\check{\ell}$ denoting irrational magnetic quantum number and $\check{\ell}$ replaces our ℓ of PSLET in (41)), $\mathbb{R} \ni r \in (0, \infty)$ and r_o is determined by (54) to read $r_o = \sqrt{\check{\ell}/|\check{\Omega}|} \in \mathbb{R}$ with $\omega = 4$, $\beta = -2k - 3/2$ by (58), and $\check{\ell} = \ell_D - \beta = 2k + |\check{\ell}| + 1$ by (59). Consequently, equations (53) would yield the zeroth-order correction

$$E_0 = 2 |\Omega E| \left(2k + |\check{\ell}| + 1 \right). \quad (66)$$

One should be aware that $r_o = \sqrt{\check{\ell}/|\check{\Omega}|} \in \mathbb{R}$ is manifestly a condition that secures the finiteness and square integrability wave function, and E_1, E_2 , and E_3 identically vanish.

Therefore, the energies associated with quasi-free KG-oscillator in Gödel SR-type cosmic string spacetime are given by (65) as

$$E^2 - 2 \check{\ell} \Omega E - (k_z^2 + m^2) = 2 |\Omega E| \left(2k + |\check{\ell}| + 1 \right); \quad |\Omega E| = \left\{ \begin{array}{l} +\Omega_\pm E_\pm \\ -\Omega_\mp E_\pm \end{array} \right\} \geq 0. \quad (67)$$

This would allow us to obtain

$$\begin{aligned} E_{\pm,1} &= \pm |\Omega| \dot{n}_+ \pm \sqrt{\Omega^2 \dot{n}_+^2 + k_z^2 + m^2}; \quad \dot{n}_+ = 2k + |\check{\ell}| + \tilde{\ell} + 1, \text{ for } |\Omega E| = +\Omega_\pm E_\pm \\ E_{\pm,2} &= \pm |\Omega| \dot{n}_- \pm \sqrt{\Omega^2 \dot{n}_-^2 + k_z^2 + m^2}; \quad \dot{n}_- = 2k + |\check{\ell}| - \tilde{\ell} + 1, \text{ for } |\Omega E| = -\Omega_\mp E_\pm \end{aligned} \quad (68)$$

However, we may rearrange such energies so that

$$E_{k,\ell,\pm}^{(+)} = \pm |\Omega| \dot{n}_{\pm} \pm \sqrt{\Omega^2 \dot{n}_{\pm}^2 + k_z^2 + m^2} \Rightarrow \left\{ \begin{array}{l} E_{k,\ell,+}^{(+)} = + |\Omega| \dot{n}_+ + \sqrt{\Omega^2 \dot{n}_+^2 + k_z^2 + m^2} \\ E_{k,\ell,-}^{(+)} = - |\Omega| \dot{n}_- - \sqrt{\Omega^2 \dot{n}_-^2 + k_z^2 + m^2} \end{array} \right\}; \text{ for } \Omega = +|\Omega|$$

$$E_{k,\ell,\pm}^{(-)} = \pm |\Omega| \dot{n}_{\mp} \pm \sqrt{\Omega^2 \dot{n}_{\mp}^2 + k_z^2 + m^2} \Rightarrow \left\{ \begin{array}{l} E_{k,\ell,+}^{(-)} = + |\Omega| \dot{n}_- + \sqrt{\Omega^2 \dot{n}_-^2 + k_z^2 + m^2} \\ E_{k,\ell,-}^{(-)} = - |\Omega| \dot{n}_+ - \sqrt{\Omega^2 \dot{n}_+^2 + k_z^2 + m^2} \end{array} \right\}; \text{ for } \Omega = -|\Omega|$$
(69)

where in $E_{k,\ell,\pm}^{(+)}$ and $E_{k,\ell,\pm}^{(-)}$ the superscripts (+) and (-) identify states with positive and negative vorticities, $\Omega = \Omega_+ = +|\Omega|$ and $\Omega = \Omega_- = -|\Omega|$, respectively. Moreover, one should notice that $\dot{n}_{\pm}(\tilde{\ell} = -|\tilde{\ell}|) = \dot{n}_{\mp}(\tilde{\ell} = |\tilde{\ell}|)$ and consequently $E_{k,\ell,\pm}^{(+)}(\tilde{\ell} = -|\tilde{\ell}|) = E_{k,\ell,\pm}^{(-)}(\tilde{\ell} = |\tilde{\ell}|)$ indicating spacetime-type associated degeneracies (STAD). Moreover, when this result is compared with that of Carvalho et al [15] (i.e., equation (14) in [15]) we observe that not only the negative energies are missed but also the STADs discussed above. Their results should, therefore, be generalized into those reported in (69). So should be the case with the results reported on the linear confinement of a scalar particle in Gödel-type space-time by Vitória [22] (their Eq. (23)) and in the related comment by Neto et al [23] (their Eq. (18)) for zero linear confinement. That would also include the results reported by Ahmed [44], where he also missed all nodeless (i.e., states with $n_r = 0$) states.

C. KG-Coulombic particle in Gödel SR-type cosmic string spacetime (2) at zero vorticity, $\Omega = 0$.

Now, we consider a KG-particle in a Gödel SR-type cosmic string spacetime (2) at zero vorticity, $\Omega = 0$, and subjected to a Lorentz scalar potential $S(r) = \beta_2/r$ in (29) to obtain

$$\left\{ \partial_r^2 - \frac{(\mathcal{L}^2 - 1/4)}{r^2} - \frac{\tilde{\beta}_2}{r} + \tilde{\lambda} \right\} R(r) = 0, \quad (70)$$

where

$$\tilde{\lambda} = E^2 - (k_z^2 + m^2); \quad \tilde{\ell} = \frac{\ell}{\alpha}, \quad \mathcal{L}^2 = \tilde{\ell}^2 + \beta_2^2, \quad \tilde{\beta}_2 = 2m\beta_2. \quad (71)$$

In this case, our PSLET recipe above implies that $D = 2 \implies \ell_D = |\mathcal{L}| - 1/2$ (with \mathcal{L} now denotes irrational magnetic quantum number and \mathcal{L} replaces our ℓ of PSLET in (41)), $r \in (0, \infty)$ and r_o is determined by (59) to read $r_o = 2\tilde{\ell}^2/\tilde{\beta}_2$ with $\omega = 2$, $\beta = -k - 1$ by (58), and $\check{\ell} = \ell_D - \beta = k + |\mathcal{L}| + 1/2$. Consequently, equations (53) would yield

$$E_0 = -\frac{\tilde{\beta}_2^2}{4(k + |\mathcal{L}| + 1/2)^2} \quad (72)$$

with E_1, E_2 , and E_3 identically vanish. Therefore, the energies associated with KG-Coulombic particle in Gödel SR-type cosmic string spacetime are given by

$$E_{k,\ell} = \pm \sqrt{k_z^2 + m^2 - \frac{m^2 \beta_2^2}{(k + |\mathcal{L}| + 1/2)^2}}. \quad (73)$$

One should notice that the result of the Heun polynomials solution for the KG-Coulomb in (40) for the case $a = 0 = \Omega$ and $b = \beta_2$ does not match the exact solution (73), where the effect of the Coulombic parameter $b = \beta_2$ have no contribution in (40).

D. Massless KG-particles in Gödel SR-type cosmic string spacetime in a Cornell-type Lorentz scalar potential

We hereby use the massless KG equation (30) with a Cornell-type Lorentz scalar potential $S(r) = \beta_1 r + \beta_2/r$ to obtain

$$\left\{ \partial_r^2 - \frac{(\mathcal{L}^2 - 1/4)}{r^2} - \tilde{\Omega}^2 r^2 + \tilde{\lambda} \right\} R(r) = 0, \quad (74)$$

$$\tilde{\Omega}^2 = \Omega^2 E^2 + \beta_1^2, \mathcal{L}^2 = \tilde{\ell}^2 + \beta_2^2, \tilde{\ell} = \frac{\ell}{\alpha}, \tilde{\lambda} = E^2 - 2\tilde{\ell}\Omega E - k_z^2 - 2\beta_1\beta_2. \quad (75)$$

Then, our PSLET recipe implies that $D = 2 \implies \ell_D = |\mathcal{L}| - 1/2$ (with \mathcal{L} denoting irrational magnetic quantum number and \mathcal{L} replaces our ℓ of PSLET in (41)), where r_o is determined by (59) to read $r_o = \sqrt{\check{\ell}/|\tilde{\Omega}|} \in \mathbb{R}$, $\omega = 4$ and $\beta = -(2k + 3/2)$ by (58) to imply $\check{\ell} = \ell_D - \beta = 2k + |\mathcal{L}| + 1$. Under such settings, one obtains

$$E_0 = 2 |\Omega E| \sqrt{1 + \frac{\beta_1^2}{\Omega^2 E^2}} (2k + |\mathcal{L}| + 1) \implies \tilde{\lambda} = 2 |\Omega E| \sqrt{1 + \frac{\beta_1^2}{\Omega^2 E^2}} (2k + |\mathcal{L}| + 1), \quad (76)$$

where $E_2 = 0$, $E_3 = 0$, and $|\tilde{\Omega}| = |\Omega E| \sqrt{1 + \beta_1^2/\Omega^2 E^2} \geq 0$ is used. Consequently, we get

$$\tilde{\lambda} = E^2 - 2\tilde{\ell}\Omega E - (k_z^2 + 2\beta_1\beta_2) = 2 |\Omega E| \sqrt{1 + \frac{\beta_1^2}{\Omega^2 E^2}} (2k + |\mathcal{L}| + 1) \quad (77)$$

to yield

$$E_{\pm}^2 - 2\tilde{\ell}\Omega_{\pm} E_{\pm} - (k_z^2 + 2\beta_1\beta_2) = 2\Omega_{\pm} E_{\pm} \sqrt{1 + \frac{\beta_1^2}{\Omega^2 E^2}} (2k + |\mathcal{L}| + 1) \quad (78)$$

for $|\Omega E| = \Omega_{\pm} E_{\pm}$, and

$$E_{\pm}^2 - 2\tilde{\ell}\Omega_{\mp} E_{\pm} - (k_z^2 + 2\beta_1\beta_2) = -2\Omega_{\mp} E_{\pm} \sqrt{1 + \frac{\beta_1^2}{\Omega^2 E^2}} (2k + |\mathcal{L}| + 1). \quad (79)$$

for $|\Omega E| = -\Omega_{\mp} E_{\pm}$. However, the two results above would eventually imply that

$$E_{\pm}^2 \mp 2 |\Omega| E_{\pm} N_{\pm} - (k_z^2 + 2\beta_1\beta_2) = 0 \Rightarrow E_{k,l,\pm} = \pm |\Omega| N_{\pm} \pm \sqrt{\Omega^2 N_{\pm}^2 + k_z^2 + 2\beta_1\beta_2}, \quad (80)$$

to read

$$E_{k,l,+}^{(+)} = + |\Omega| N_+ + \sqrt{\Omega^2 N_+^2 + k_z^2 + 2\beta_1\beta_2} \quad \text{and} \quad E_{k,l,-}^{(+)} = - |\Omega| N_- - \sqrt{\Omega^2 N_-^2 + k_z^2 + 2\beta_1\beta_2} \quad (81)$$

for $\Omega = \Omega_+ = +|\Omega|$, and

$$E_{\pm}^2 \mp 2 |\Omega| E_{\pm} N_{\mp} - (k_z^2 + 2\beta_1\beta_2) = 0 \Rightarrow E_{k,l,\pm} = \pm |\Omega| N_{\mp} \pm \sqrt{\Omega^2 N_{\mp}^2 + k_z^2 + 2\beta_1\beta_2}, \quad (82)$$

to read

$$E_{k,l,+}^{(-)} = + |\Omega| N_- + \sqrt{\Omega^2 N_-^2 + k_z^2 + 2\beta_1\beta_2} \quad \text{and} \quad E_{k,l,-}^{(-)} = - |\Omega| N_+ - \sqrt{\Omega^2 N_+^2 + k_z^2 + 2\beta_1\beta_2}, \quad (83)$$

for $\Omega = \Omega_- = -|\Omega|$, where

$$N_{\pm} = \sqrt{1 + \frac{\beta_1^2}{\Omega^2 E^2}} (2k + |\mathcal{L}| + 1) \pm \tilde{\ell}. \quad (84)$$

Again, one observes that $N_{\pm} (\tilde{\ell} = -|\tilde{\ell}|) = N_{\mp} (\tilde{\ell} = |\tilde{\ell}|)$ and consequently $E_{k,\ell,\pm}^{(+)} (\tilde{\ell} = -|\tilde{\ell}|) = E_{k,\ell,\pm}^{(-)} (\tilde{\ell} = |\tilde{\ell}|)$ to represent spacetime-type associated degeneracies (STAD).

VI. CONCLUDING REMARKS

In this paper, we started with KG-oscillators in a topologically trivial Gödel-type spacetime background. A KG-oscillator I as a topologically trivial Gödel-type spacetime byproduct, and a KG-oscillator II as a topologically trivial

Gödel-type spacetime plus momentum operator deformation byproduct. We have used simple and straightforward textbook procedures and reported, for the sake of scientific correctness, their exact relativistic energies in (14), (15), and (24). This would make the results reported by [18] on KG-oscillators in topologically trivial Gödel-type spacetime (4) only partially correct and consequently should be generalized into those reported in (24). Next, we have discussed the drawbacks associated with the power series (Frobinous method) expansion approach that implied the biconfluent Heun functions/polynomials solution (33). We have shown that for the effective interaction potential $V_{eff}(r) = \tilde{\Omega}^2 r^2 + 2mar + 2mb/r$ in the Schrödinger-like quantum model (31), the biconfluent Heun polynomial solution (33) cease to be able to address the especial case of KG-Coulombic particle when $a = 0 = \Omega$. Consequently, the biconfluent Heun polynomial solution (33) can only be classified as a conditionally exact solution that may address the KG-oscillator (36), when $a = 0 = b$, but tragically fails to address the KG-Coulomb case when $a = 0 = \Omega$.

As an alternative method/technique, we have recollected, in short, the so called pseudo perturbative shifted ℓ expansion technique PSLET (e.g., [27–35]) to solve for the D -dimensional Schrödinger-type equations (which is a common form the relativistic wave equations collapse into, like (60), (31), etc). Some illustrative examples on the straightforward/simplistic applicability of PSLET procedure are used. Amongst, a KG-oscillator as a topologically trivial Gödel-type spacetime byproduct of (10) that collapses into the one dimensional form of Schrödinger equation, and three examples that resemble the radial two-dimensional form of Schrödinger equation: a Quasi-free KG-oscillator in Gödel SR-type cosmic string spacetime (2), a KG-Coulombic particle in Gödel SR-type cosmic string spacetime (2) at zero vorticity $\Omega = 0$, and a massless KG-particle in Gödel SR-type cosmic string spacetime in a Cornell-type Lorentz scalar potential. In all of our illustrative examples above, the exact energies for the harmonic oscillator (i.e., (60), (64), and (74)) like and Coulomb like (70) interactions are obtained from the zeroth (leading) order correction, where all higher order correction identically vanish. In connection with the effective potential $V_{eff}(r) = \tilde{\Omega}^2 r^2 + 2mar + 2mb/r$ in the Schrödinger-like quantum model (31), nevertheless, one would need to numerically solve for ω and r_o in (58) and (59) for each parametric value involved in the Gödel SR-type cosmic string spacetime metric (2). In this case, although tedious, one would work out the corresponding eigenvalues for such a quantum mechanical problem. The accuracy of PSLET is shown to be very satisfactory and reliable for similar Schrödinger-like models [29–32, 34, 35]. This issue already lies far beyond the scope of the current proposal.

Finally, we believe that the effects of gravitational fields introduced by different spacetime structures can be clarified if only if the fundamental models like the KG-oscillators and KG-Coulomb are comprehensively and explicitly reported. To the best of our knowledge, the results of the current methodical proposal have never been published elsewhere.

VII. APPENDIX: EXPLICIT FORMS OF α_1 AND α_2

In this section we give the explicit forms of α_1 and α_2 as given by [28, 29],

$$\alpha_1 = [(1 + 2k) e_2 + 3(1 + 2k + 2k^2) e_4] - \frac{1}{\omega} [e_1^2 + 6(1 + 2k) e_1 e_3 + (11 + 30k + 30k^2) e_3^2], \quad (85)$$

$$\begin{aligned} \alpha_2 = & [(1 + 2k) d_2 + 3(1 + 2k + 2k^2) d_4 + 5(3 + 8k + 6k^2 + 4k^3) d_6] \\ & - \frac{1}{\omega} [(1 + 2k) e_2^2 + 12(1 + 2k + 2k^2) e_2 e_4 + 2e_1 d_1 + 2(21 + 59k + 51k^2 + 34k^3) e_4^2 \\ & + 6(1 + 2k) e_1 d_3 + 30(1 + 2k + 2k^2) e_1 d_5 + 6(1 + 2k) e_3 d_1 + 2(11 + 30k + 30k^2) e_3 d_3 \\ & + 10(13 + 40k + 42k^2 + 28k^3) e_3 d_5] + \frac{1}{\omega^2} [4e_1^2 e_2 + 36(1 + 2k) e_1 e_2 e_3 \\ & + 8(11 + 30k + 30k^2) e_2 e_3^2 + 24(1 + 2k) e_1^2 e_4 + 8(31 + 78k + 78k^2) e_1 e_2 e_4 \\ & + 12(57 + 189k + 225k^2 + 150k^3) e_3^2 e_4] - \frac{1}{\omega^3} [8e_1^3 e_3 + 108(1 + 2k) e_1^2 e_3^2 \\ & + 48(11 + 30k + 30k^2) e_1 e_3^3 + 30(31 + 109k + 141k^2 + 94k^3) e_3^4], \end{aligned} \quad (86)$$

with

$$e_j = \varepsilon_j / \omega^{j/2}; \quad d_i = \delta_i / \omega^{i/2}; \quad j = 1, 2, 3, 4, \quad i = 1, 2, 3, 4, 5, 6, \quad (87)$$

$$\varepsilon_1 = -4\beta, \quad \varepsilon_2 = 6\beta, \quad \varepsilon_3 = -4 + r_o^5 V'''(r_o) / 6Q, \quad \varepsilon_4 = 5 + r_o^6 V''''(r_o) / 24Q, \quad (88)$$

$$\begin{aligned} \delta_1 = -2(\beta^2 - 1/4), \quad \delta_2 = 3(\beta^2 - 1/4), \quad \delta_3 = -8\beta, \quad \delta_4 = 10\beta, \\ \delta_5 = -6 + r_o^7 V'''''(r_o) / 120Q, \quad \delta_6 = 7 + r_o^8 V''''''(r_o) / 120Q. \end{aligned} \quad (89)$$

-
- [1] A. Vilenkin, Phys. Let. B **133** (1983) 177.
- [2] A. Vilenkin, Phys. Rep. **121** (1985) 263.
- [3] T. W. Kibble, J. Phys. **A 19** (1976) 1387.
- [4] M. Hiscock, Phys. Rev. **D 31** (1985) 3288.
- [5] M. Barriola, A. Vilenki, Phys. Rev. Lett. **63** (1989) 341.
- [6] M. Cruz, N. Turok, P. Vielva, E. M. Gonzalez, M. Hobson, Science **318** (2007) 1612.
- [7] K. Gödel, Rev. Mod. Phys. **21** (1949) 447.
- [8] J. Duarte de Oliveira, A.F. Teixeira, J. Tiomno, Phys. Rev. D **34** (1986) 3661.
- [9] M.M. Som, A.K. Raychaudhuri, Proc. R. Soc. London **A304** (1968) 81.
- [10] R.J. Gleiser, M. Gürses, A. Karasu, O. Sarioğlu, Class. Quant. Grav. **23** (2006) 2653.
- [11] M. Gurses, Gen. Relativ. Gravit. **41** (2009) 31.
- [12] H. Sobhani, H. Hassanabadi, W. S. Chung, Int. J. Geom. Methods Mod. Phys. **15** (2018) 1850037.
- [13] M.J. Rebouças, J. Tiomno, Phys. Rev. D **28** (1983) 1251.
- [14] N. Drukker, B. Fiol, J. Simón, JCAP **0410** (2004) 012.
- [15] J. Carvalho, A. M. de M. Carvalho, C. Furtado, Eur. Phys. J. **C 74** (2014) 2935.
- [16] B. D. B. Figueiredo, I. D. Soares, J. Tiomno, Class. Quantum Gravit. **9** (1992) 1593.
- [17] Z. Wang, Z. Long, C. Long, M. Wu, Eur. Phys. J. Plus **130** (2015) 36.
- [18] F. Ahmed, Eur. Phys. J. C **78** (2018) 598.
- [19] L. Inzunza, M. S. Plyushchay, JHEP **01** (2022) 179.
- [20] H. Hassanabadi, S. Zare, M. de Montigny, Gen. Relativ. Gravit. **50** (2018) 47.
- [21] P. Sedaghatnia, H. Hassanabadi, F. Ahmed, Eur. Phys. J. C **79** (2019) 541.
- [22] R.L.L. Vitória, C. Furtado, K. Bakke, Eur. Phys. J. C **78** (2018) 44.
- [23] F.A.C. Neto, C.C. Soares, L.B. Castro, Eur. Phys. J. C **80** (2020) 53.
- [24] F. Ahmed, Eur. Phys. J. C **79** (2019) 682.
- [25] S. Flügge, *Practical quantum mechanics*, Springer-Verlag, New York, Heidelberg, Berlin 1974.
- [26] D.R. Herschbach et al. (1993) *Dimensional Scaling in Chemical Physics (Dordrecht, Kluwer)*.
- [27] U. Sukhatme, T. Imbo, Phys. Rev. D **28** ((1983) 418.
- [28] T. Imbo, A. Pagnamenta, U Sukhatme, Phys. Rev. D **29** ((1984) 1669.
- [29] O. Mustafa, T. Barakat, Commun. Theor. Phys. **28** (1997) 257.
- [30] O. Mustafa, M. Odeh, J. Phys. A: Math. Gen **33** (2000) 5207.
- [31] O. Mustafa, J. Phys.: Condens. Matter **5** (1993) 1327.
- [32] O. Mustafa, S.C. Chhajlany, Phys. Rev. A **50** (1994) 2926.
- [33] O. Mustafa, R. Sever, Phys. Rev. A **44** (1991) 4142.
- [34] O. Mustafa, M Znojil, J. Phys. A: Math. Gen **35** (2002) 8929.
- [35] M. Znojil, F. Gemperle, O. Mustafa, J. Phys. A: Math. Gen **35** (2002) 5781.
- [36] C. Quigg, J. J. Rosner, Phys. Rep. **56** (1979) 167.
- [37] M. Moshinsky, A. Szczepaniak, J. Phys. A: Math. Gen. **22** (1989) L817.
- [38] B. Mirza, M. Mohadesi, Commun. Theor. Phys. **42** (2004) 664.
- [39] E.R. Arriola, A. Zarzo, J.S. Dehesa, J. Comput. Appl. Math. **37** (1991) 161.
- [40] O. Mustafa, Ann. Phys. **440** (2022) 168857.
- [41] O. Mustafa, Z. Algadhi, Chin. J. Phys. **65** (2020) 554.
- [42] O. Mustafa, Eur. Phys. J. C **82** (2022) 82.
- [43] O. Mustafa, arXiv:2111.10077, " Klein-Gordon oscillator in a cosmic spacetime with a cosmic space-like dislocation: Conditional solvability and position-dependent mass settings" (2022).
- [44] F. Ahmed, Eur. Phys. J. Plus **135** (2020) 588.