# APPROACHING THE ISOPERIMETRIC PROBLEM IN  $H_{\mathbb{C}}^{m}$  VIA THE HYPERBOLIC LOG-CONVEX DENSITY CONJECTURE

### LAURO SILINI

Abstract. We prove that geodesic balls centered at some base point are isoperimetric in the real hyperbolic space  $H_{\mathbb{R}}^n$  endowed with a smooth, radial, strictly log-convex density on the volume and perimeter. This is an analogue of the result by G. R. Chambers for log-convex densities on  $\mathbb{R}^n$ . As an application we prove that in any rank one symmetric space of non-compact type, geodesic balls are isoperimetric in a class of sets enjoying a suitable notion of radial symmetry.

### 1. INTRODUCTION

We denote by  $(H_{\mathbb{R}}^n, g_H)$  the real hyperbolic space of dimension  $n \in \mathbb{N}$  with constant sectional curvature equal to  $-1$ . Call  $d_H$  the induced Riemannian distance. Choose an arbitrary base point  $o \in H^n_{\mathbb{R}}$ . We say that a function  $f: H^n_{\mathbb{R}} \to \mathbb{R}_{>0}$  is (strictly) radially log-convex if

$$
\ln(f(x)) = h(d_H(o, x)),
$$

for a smooth, (strictly) convex and even function  $h : \mathbb{R} \to \mathbb{R}$ . We define the weighted perimeter and volume of a set with finite perimeter  $E \subset H_{\mathbb{R}}^n$  as

<span id="page-0-0"></span>
$$
V_f(E) := \int_E f \, d\mathscr{H}^n, \quad \text{and} \quad P_f(E) = \int_{\partial^* E} f \, d\mathscr{H}^{n-1}.
$$

Here, following the notation in [\[9\]](#page-16-0),  $\partial^* E$  denotes the reduced boundary of E. A set of finite perimeter E with volume  $V_f(E) = v > 0$  is called isoperimetric if it solves the minimization problem

(1.1) 
$$
\inf \Big\{ P_f(F) : V_f(F) = v, F \subset H^n_{\mathbb{R}} \text{ of finite perimeter} \Big\}.
$$

The first goal of this paper is to show the following characterization of the isoperimetric sets, which will be developed in Section [3.](#page-5-0)

<span id="page-0-1"></span>Theorem 1.1. For any strictly radially log-convex density f, geodesic balls centered at  $o \in H_{\mathbb{R}}^n$  are isoperimetric sets with respect to the weighted volume and perimeter  $V_f$  and  $P_f$ .

This result is the hyperbolic twin of an analogous result in the Euclidean space, conjectured first by Kenneth Brakke (see [\[14,](#page-16-1) Conjecture 3.12]), stating that Euclidean balls centered at the origin are isoperimetric in  $\mathbb{R}^n$  endowed with a log-convex density. We invite the reader to consult F. Morgan [\[11\]](#page-16-2) and F. Morgan and A. Pratelli [\[13\]](#page-16-3) for existence and regularity properties, V. Bayle, A. Cañete, F. Morgan and C. Rosales [\[14](#page-16-1)] for the stability, A. Figalli and F. Maggi [\[5](#page-15-0)] for the small volume regime, A. Kolesnikov and R. Zhdanov [\[8\]](#page-15-1) for the large volume regime. The conjecture was then recently completely solved by the remarkable article by G. R. Chambers [\[2\]](#page-15-2). In fact, the first and main part of this paper is an adaptation of the method presented in the latter to the real hyperbolic space.

In the hyperbolic setting, the two dimensional case was solved by I. McGillivray in [\[10\]](#page-16-4). We refer as well to [\[1](#page-15-3), [3](#page-15-4), [7](#page-15-5), [12](#page-16-5)] for other works related to this problem.

#### 2 LAURO SILINI

Even if interesting by itself, our main motivation in proving such result is the tight relation of this problem with the (unweighted) isoperimetric problem in the complex hyperbolic spaces  $H_{\mathbb{C}}^m$ , the quaternionic spaces  $H_{\mathbb{H}}^m$  and the Cayley plane  $H_{\mathbb{O}}^2$  restricted to a family of sets sharing a particular symmetry that we define as follows.

<span id="page-1-0"></span>**Definition 1.2** (Hopf-symmetric sets). Let  $\mathbb{K} \in \{ \mathbb{C}, \mathbb{H}, \mathbb{O} \}, d = \dim(\mathbb{K}) \in \{2, 4, 8\}$  and  $(M, g) = (H_{\mathbb{K}}^{m}, g)$  be the associated rank one symmetric space of non-compact type of real dimension  $n = dm$ ,  $m = 2$  if  $K = \mathbb{O}$ . Fix an arbitrary point  $o \in M$  and let N be the unit length radial vector field emanating from o. Then, up to renormalization of the metric, the Jacobi operator  $R(\cdot, N)N$  arising from the Riemannian curvature tensor R is a self adjoint operator of  $TM$ , and has exactly three eigenvalues:  $\{0, -1, -4\}$ . The  $(-4)$ -eigenspace defines at every point  $x \neq o$  a distribution  $\mathcal{H}_x$  of real dimension  $d-1$ . A  $C^1$ -set  $E \subset M$  with normal vector field  $\nu$  is said to be *Hopf-symmetric* if  $\nu(x)$  is orthogonal to  $\mathcal{H}_x$  at each point  $x \in \partial E$ ,  $o \notin \partial E$ .

Remark 1.3. Let  $h: S^{n-1} \to \mathbb{K}P^{m-1}$  be the celebrated Hopf fibration. Then, for any  $C^1$ -profile  $\rho: S^{n-1} \to (0, +\infty)$  so that  $\rho$  is constant along the fibres of h, the set with boundary

$$
\partial E := \{ \exp_o(\rho(x)) : x \in S^{n-1} \subset T_o M \},\
$$

is Hopf-symmetric.

Remark 1.4. Being Hopf-symmetric has not to be confused with the standard notion of being  $Hopf$  in  $H_{\mathbb{C}}^m$ , that is a set with principal curvature along the characteristic directions  $J\nu$ , where  $J$  denotes the associated complex structure. It is worth saying that spheres are the only Hopf, compact, embedded mean curvature surfaces in  $H_{\mathbb{C}}^{m}$ , as it is proven by X. Wang in [\[15](#page-16-6)]. The natural generalization of this concept when  $\mathbb{K} \in \{\mathbb{H}, \mathbb{O}\}\$ is being a curvature-adapted hypersurface, that is the normal Jacobi operator  $R(\cdot,\nu)\nu$  commutes with the shape operator.

We adopt the notation of Definition [1.2](#page-1-0) for the rest of the paper. Let  $P$  and  $V$  be the perimeter and volume functionals induced by  $g$  in  $H_{\mathbb{K}}^m$ . Consider the (unweighted) isoperimetric problem

(1.2) 
$$
\inf \Big\{ P(F) : V(F) = v, F \subset H^m_{\mathbb{K}} \text{ Hopf-symmetric} \Big\}.
$$

We dedicate Section [4](#page-14-0) to the proof of this theorem.

<span id="page-1-2"></span>**Theorem 1.5.** If geodesic balls centered at  $o \in H_{\mathbb{R}}^n$  are isoperimetric with respect to Problem [\(1.1\)](#page-0-0) for the strictly radial log-convex density

<span id="page-1-1"></span>
$$
f(x) = \cosh(d_{H_{\mathbb{R}}^n}(o, x)))^{d-1}, \quad d = \dim(\mathbb{K}),
$$

then geodesic balls in  $H^m_{\mathbb{K}}$  are optimal with respect to the isoperimetic Problem [\(1.2\)](#page-1-1).

Combining this with Theorem [1.1](#page-0-1) we get immediately the following Corollary.

Corollary 1.6. In the class of Hopf-symmetric sets, geodesic balls are isoperimetric regions in  $H_{\mathbb{K}}^m$ .

Acknowledgments. The author would like to thank Urs Lang and Alessio Figalli for their precious guidance and constant support. A special thanks to Raphael Appenzeller for the very enjoyable and instructive exchanges about the hyperbolic plane. Lastly, the author would like to thank Miguel Domínguez Vázquez for suggesting an efficient way to include the octonionic case in the definition of Hopf-symmetric, and Frank Morgan, for the useful insight about the earlier contributions to the problem. The author has received funding from the European Research Council under the Grant Agreement No. 721675 "Regularity and Stability in Partial Differential Equations (RSPDE)".

## <span id="page-2-1"></span>2. Preliminaries

In what follows, we will always assume  $E \subset H_{\mathbb{R}}^n$  to be an isoperimetric set with respect to the weighted problem [\(1.1\)](#page-0-0).

2.1. Qualitative properties of the isoperimetric sets. Supposing  $E$  smooth, the volume preserving first variation of the perimeter gives us that

(2.1) 
$$
\mathbf{H}_f := H + \partial_\nu h = \text{constant},
$$

on  $\partial E$ , where H is the unaveraged inward mean curvature and  $\nu$  is the outward pointing unit normal. Existence and regularity properties of isoperimetric sets are summarized in the following theorem. We refer to the papers of F. Morgan [\[11\]](#page-16-2) and F. Morgan and A. Pratelli [\[13\]](#page-16-3). Their results on  $\mathbb{R}^n$  generalise directly to  $H^n_{\mathbb{R}}$ .

<span id="page-2-0"></span>**Theorem 2.1** (Existence and regularity). For any volume  $v > 0$  there exists a set  $E \subset$  $H_{\mathbb{R}}^n$  of finite perimeter and weighted volume  $V_f(E) = v$  solving the isoperimetric problem  $(1.1)$ . Moreover, E enjoys the following properties:

- $\partial E$  is a bounded embedded hypersurface with singular set of Hausdorff dimension at most  $n-8$ .
- There exists  $\lambda \in \mathbb{R}$  such that at any regular point  $x \in \partial E$ ,  $H_f(x) = \lambda$ . As a consequence,  $\partial E$  is mean-convex at each regular point  $y \in \partial E$ , that is  $H(y) \geq$  $(n-1)$ .
- If the tangent cone at  $x \in \partial E$  lies in an halfspace, then it is an hyperplane, and therefore  $\partial E$  is regular at x. In particular,  $\partial E$  is regular at points  $x^* \in \partial E$ satisfying  $d_H(x^*, o) = \sup_{x \in \partial E} d_H(x, o)$ .

2.2. The Poincaré model of  $H^n_{\mathbb{R}}$ . Adopting the Poincaré model,  $H^n_{\mathbb{R}}$  is conformal to the open Euclidean unit ball. At a point  $x \in H_{\mathbb{R}}^n$  the metric is

$$
g_H = \frac{4}{(1 - r^2)^2} g_{\text{flat}},
$$

where  $r = |x|$  will always denote the Euclidean distance of x from the origin, and  $g_{\text{flat}}$  the usual Euclidean metric of  $\mathbb{R}^n$ . The hyperbolic distance from the origin is then given by

$$
d_H(x,0) = 2\operatorname{artanh}(r).
$$

We define the boundary at infinity  $\partial_{\infty} H_{\mathbb{R}}^n$  of  $H_{\mathbb{R}}^n$  to be the Euclidean unit sphere  $\partial B(0,1) =$  $S^{n-1}$ . We will identify the base point  $o \in H_{\mathbb{R}}^{n}$  of the radial density f with the origin 0 in  $B(0, 1)$ .

2.3. Isometries and special frames in  $H^2_{\mathbb{R}}$ . Denote by  $e_1$  and  $e_2$  the horizontal and vertical Cartesian axis in the two dimensional Poincaré disk model. Also, let  $(H_{\mathbb{R}}^2)_+$  be the intersection of  $H^2_{\mathbb{R}}$  with the closed upper half-plane having  $e_1$  as boundary. The isometry group of  $(H^2_{\mathbb{R}}, g_H)$  is completely determined (up to orientation) by the Möbius transformations fixing the boundary  $\partial_{\infty} H_{\mathbb{R}}^2$ . Hence, geodesic spheres coincide with Euclidean circles completely contained in  $H^2_{\mathbb{R}}$ . Their curvature lies in  $(1, +\infty)$ . Circles touching  $\partial_{\infty}H_{\mathbb{R}}^2$  in a point are called horospheres, and have curvature equal to 1. Geodesics are arcs of (possibly degenerate) circles hitting  $\partial_{\infty} H_{\mathbb{R}}^2$  perpendicularly in two points. Arcs of (possibly degenerate) circles that are not geodesics are called hypercycles, and have constant curvature in  $(-1, 1) \setminus \{0\}$ . It will be convenient to work with a particular frame: define

$$
S:(H^2_{\mathbb{R}})_+\to\mathbb{R},
$$

to be the hyperbolic distance of a point in  $(H_{\mathbb{R}}^2)_+$  from the horizontal axis  $e_1$ . Set  $X =$  $\nabla S$ , where we naturally extend by continuity X at  $e_1$ . Then, denoting with  $X^{\perp}$  the

#### 4 LAURO SILINI

counterclockwise rotation of X by  $\frac{\pi}{2}$  radians, since the level sets of S are equidistant to each other,  $\{X, X^{\perp}\}\)$  forms an orthonormal frame of  $(H_{\mathbb{R}}^2)_+$ , see Figure [1.](#page-3-0) The integral curves of X are all geodesic rays hitting  $e_1$  perpendicularly. For each  $l \in [0,1)$ , let  $\delta_l$  be the integral curve of  $X^{\perp}$  so that  $\delta_l(0) = (0, l)$ . Then,  $(\delta_l)_{l \in [0,1)}$  is a family of equidistant hypercycles foliating  $(H_{\mathbb{R}}^2)_+$ , crossing  $e_2$  perpendicularly and with constant curvature which coincides with the Eucliden one:  $K_1 = \frac{2l}{1+l}$  $\frac{2l}{1+l^2} = \frac{1}{R(1)}$  $\frac{1}{R(l)}$ , where  $R(l) \in (0, +\infty]$  is the radius of the Euclidean circle representing the curve. Similarly, set  $\{N, N^{\perp}\}\$ to be the orthonormal frame on  $H^2_{\mathbb{R}} \setminus \{0\}$  where N is the radial unit length vector field emanating from the origin. Then, integral curves of N are rays of geodesics, and integral curves of  $N^{\perp}$ are concentric geodesic spheres. Notice that the frame  $\{X, X^{\perp}\}\$ is invariant under the oneparameter subgroup of hyperbolic isometries fixing  $e_1$  ( $X^{\perp}$  is the infinitesimal generator of the action by translations) and, up to orientation reversing, under the reflections with respect to any geodesic integral curve of X. Finally, notice that on  $e_1$  and  $e_2$ ,  $\{X, X^{\perp}\}\$ is a positive rescaling of  $\{(0, 1), (-1, 0)\}.$ 



FIGURE 1. The special frame  $\{X, X^{\perp}\}.$ 

For a regular curve parametrized by arc length  $\eta$  we denote with  $\kappa_n(t)$  the inward signed curvature of  $\eta$  at  $\eta(t)$ . We recall the identity

<span id="page-3-0"></span>
$$
\kappa_\eta \dot{\eta}^\perp = \nabla_{\dot{\eta}} \dot{\eta},
$$

where here  $\nabla$  denotes the standard Levi-Civita connection associated to  $q_H$ .

2.4. Reduction to  $H^2_{\mathbb{R}}$ . From now on, let E be an isoperimetric set with arbitrary weighted volume. Since both the density f and the conformal term of  $g_H$  are radial, the coarea formula implies that spherical symmetrization pointed at the origin preserves the weighted volume and does not increase the weighted perimeter of  $E$ . For this reason, we will assume E spherically symmetric with respect to the  $e_1$  axis. Now, intersecting E with the the Euclidean plane spanned by  $\{e_1, e_2\}$ , we obtain a spherically symmetric profile  $\Omega \subset H^2_{\mathbb{R}}$ . Let  $x^*$  be the furthest point of  $\Omega$  lying in the positive part of the  $e_1$  axis (this is always possible by reflecting  $\Omega$  with respect to the  $e_2$  geodesic). Let  $\gamma : [-a, a] \to H^2_{\mathbb{R}}$  be a counter-clockwise, arclength parametrization of the boundary of the connected component of  $\Omega$  containing  $x^*$ , so that  $\gamma(0) = x^*$ , see Figure [2.](#page-4-0) The curve  $\gamma$  enjoys the following properties:

- $-γ$  is smooth on  $(-a, a)$ . Indeed, if there exists  $a^* ∈ (-a, a)$  such that  $γ(a^*)$  is not regular, then  $\partial E$  contains a singular set of Hausdorff dimension  $n-2$ , but this cannot be because of Theorem [2.1.](#page-2-0)
- The curve  $\gamma$  forms a simple, closed curve.
- Writing  $\gamma = (\gamma_1, \gamma_2)$  in cartesian coordinates, one has that  $sgn(\gamma_2(t)) = sgn(t)$ . In particular,  $\gamma : [0, a) \to (H_{\mathbb{R}}^2)_+.$
- $-\dot{\gamma}(0) = X(\gamma(0)).$



<span id="page-4-0"></span>FIGURE 2. The spherical symmetrization.

To translate Equation [\(2.1\)](#page-2-1) as a property of the profile  $\gamma$ , we need the following definition.

<span id="page-4-2"></span>**Definition 2.2.** For any  $t \in [0, a)$ , denote by

- $C_t = C_t(s)$  the (possibly degenerated) oriented circle tangent to  $\gamma(t)$ , with center on  $e_1$ , parametrized by arclength and such that  $C_t(0) = \gamma(t)$ . Denote by  $\kappa(C_t)$  its constant curvature.
- Similarly, call  $c_t = c_t(s)$  the (possibly degenerated) oriented circle tangent to  $\gamma(t)$ , parametrized by arclength, such that  $c_t(0) = \gamma(t)$  and  $\kappa(c_t) = \kappa_{\gamma}(t)$ .
- Define  $x(C_t)$  and  $x(c_t)$  to be the hyperbolic center of  $C_t$  and  $c_t$  respectively. Similarly, let  $x_1(C_t)$  and  $x_1(c_t)$  be the first Euclidean coordinate of  $x(C_t)$  and  $x(c_t)$ respectively.

<span id="page-4-1"></span>Remark 2.3. Let  $F \subset B(0,1) \subset \mathbb{R}^n$ . Then, at every regular point  $x \in \partial F$ , the mean curvature  $H$  is related with the Euclidean mean curvature  $H^{\text{flat}}$  by

$$
H(x) = \frac{1 - r^2}{2} H^{\text{flat}}(x) + (n - 1) g_{\text{flat}}(x, \tilde{\nu}),
$$

where  $\tilde{\nu}$  is the outward normal vector to  $\partial F$  with Euclidean norm equal to one. In particular, when  $n = 2$ , denoting with  $\kappa^{\text{flat}}$  the usual Euclidean curvature, one has that

$$
\kappa_{\eta} = \frac{1 - |\gamma(t)|^2}{2} \kappa_{\eta}^{\text{flat}} + g_{\text{flat}}(\eta, \tilde{\nu}),
$$

Therefore,  $\kappa^{\text{flat}}(c_t) = \kappa^{\text{flat}}_{\gamma}$ <sup>flat</sup>, that is comparison circles  $c_t$  and  $C_t$  in the hyperbolic setting coincide with comparison circles with respect to the Euclidean metric. From this formula, we also deduce that for any Euclidean circle  $\mathscr C$ 

$$
\kappa_{\mathscr{C}} = \frac{1}{2} \left( \frac{1 - |x_0|^2}{\tau_0} + \tau_0 \right) = \coth(\tau),
$$

where  $x_0$  and  $\tau_0$  are the Euclidean center and radius, and  $\tau$  is the hyperbolic radius.

<span id="page-5-3"></span>**Lemma 2.4.** On  $t \in [0, a)$  it holds

$$
H = \kappa_{\gamma} + (n-2)\kappa(C_t).
$$

In particular,

$$
H_f = \kappa_\gamma + (n-2)\kappa(C_t) + h'g_H(\nu, N) = \lambda,
$$

where  $\nu = -\dot{\gamma}^{\perp}$ .

We call  $H_1 := h' g_H(\nu, N)$  the term coming from the log-convex density.

Proof. In [\[2,](#page-15-2) Proposition 3.1] it is shown that the Euclidean mean curvature of the spherically symmetric set  $E$  can be computed as

$$
H^{\text{flat}} = \kappa_{\gamma}^{\text{flat}} + (n-2)\kappa^{\text{flat}}(C_t).
$$

Thanks to Remark [2.3](#page-4-1) we have that

$$
H(\gamma(t)) = \frac{1 - |\gamma(t)|^2}{2} H^{\text{flat}}(\gamma(t)) + (n - 1)g_{\text{flat}}(\gamma(t), \nu)
$$
  
= 
$$
\frac{1 - |\gamma(t)|^2}{2} \left(\kappa_{\gamma}^{\text{flat}} + (n - 2)\kappa^{\text{flat}}(C_t)\right) + (n - 1)g_{\text{flat}}(\gamma(t), \nu)
$$
  
=  $\kappa_{\gamma} + (n - 1)\kappa(C_t).$ 

## 3. The proof

<span id="page-5-0"></span>The reduction to  $H_{\mathbb{R}}^2$  implies that Theorem [1.1](#page-0-1) is equivalent to showing that  $\gamma$  represents a circumference centered in the origin. Essentially, we prove that ruling out this possibility, implies that  $\gamma$  has to make a curl (see Figure [3\)](#page-6-0), contradicting the fact that  $\gamma$  parametrizes a spherically symmetric set. This is made rigorous by the combination of the next two lemmas.

<span id="page-5-2"></span>**Lemma 3.1.** For every  $t \in (0, a)$ 

$$
g_H(N,\dot{\gamma}) \leq 0.
$$

Proof. The set  $\Omega$  spherically symmetric implies that  $t \mapsto g_{\text{flat}}(\gamma(t), \gamma(t))$  is non increasing.<br>Differentiating in t gives the desired sign of the angle between N and  $\dot{\gamma}$ Differentiating in t gives the desired sign of the angle between N and  $\dot{\gamma}$ .

Section [3.1](#page-6-1) is devoted to the proof of the next lemma.

<span id="page-5-1"></span>**Lemma 3.2** (The tangent lemma). If  $\gamma$  is not a circle centered in the origin, there exist  $0 < a_0 < a_1 < a_2 < a$  such that  $\dot{\gamma}(a_0) = X^{\perp}(\gamma(a_0)), \dot{\gamma}(a_1) = -X(\gamma(a_1))$  and  $\dot{\gamma}(a_2) = X(\gamma(a_2)).$ 

Assuming now that Lemma [3.2](#page-5-1) holds true, the proof of the main result goes as follows.

*Proof of Theorem [1.1.](#page-0-1)* If  $\gamma$  is a circle centered at the origin we are done. Otherwise, Lemma [3.2](#page-5-1) ensures the existence of  $0 < a_2 < a$  such that  $\dot{\gamma}(a_2) = X(\gamma(a_2))$ . This violates the inequality of Lemma [3.1,](#page-5-2) because at  $a_2$ 

$$
g_H(N, \dot{\gamma}) = g_H(N, X) > 0.
$$

 $\Box$ 

 $\Box$ 



<span id="page-6-0"></span>Figure 3. The curl described in Lemma [3.2.](#page-5-1)

<span id="page-6-1"></span>3.1. Proof of the tangent lemma. The proof is made by following the behaviour of  $\gamma$  step by step: first we show that  $\gamma$  has to arch upwards with curvature strictly greater than one. The endpoint of this arc will be  $\gamma(a_0)$ , where  $\dot{\gamma}(a_0) = X^{\perp}(\gamma(a_0))$ . Then, it goes down curving strictly faster than before, and this result about curvature comparison is the tricky point to generalize in the hyperbolic setting. It turns out that the special frame given by the hypercyclical foliation  $(\delta_l)_{l\in[0,1)}$  is the good one. Then, arguing by contradiction, we will show that this behaviour must end at a point  $0 < a_0 < a_1 < a$ , where  $\dot{\gamma}(a_1) = -X(\gamma(a_1))$ . Finally, we prove the existence of  $a_2$  so that  $\dot{\gamma}(a_2) = X(\gamma(a_2))$ taking advantage of the mean-curvature convexity of  $\Omega$ . We start by looking at what happens at the starting point.

<span id="page-6-4"></span>**Lemma 3.3.** One has that  $\dot{\gamma}(0) = X(\gamma(0)), \dot{\kappa}_\gamma(0) = 0$  and  $\kappa_\gamma(0) \ge \kappa(C_0) > 1$ .

*Proof.* This is a consequence of the symmetry of  $\gamma$  with respect to the  $e_1$  axis, and that  $\gamma(0)$  represent the furthest point from the origin of  $\Omega$ .

<span id="page-6-3"></span>**Lemma 3.4.** If there exists  $t^* \in [0, a)$  such that  $x_1(C_{t^*}) = 0$  and  $\kappa_\gamma(t^*) = \kappa(C_{t^*})$ , then  $\gamma$ has to be a centered circle.

*Proof.* In this case  $\gamma(t)$  and  $C_{t^*}(s)$  solve the same ODE, with same initial data. Therefore, they have to coincide locally, and hence globally.  $\Box$ 

**Definition 3.5.** Call  $\alpha$  :  $[0, a) \rightarrow [\pi, -\pi]$  the oriented angle made by  $\dot{\gamma}$  with  $X^{\perp}$ . We say that  $\dot{\gamma}(t)$  is in the I, II, III and IV quadrant if  $\alpha(t)$  belongs to  $[\pi/2, \pi]$ ,  $[0, \pi/2]$ ,  $[0, -\pi/2]$ and  $[-\pi/2, -\pi]$  respectively. We add *strictly* if  $\gamma$  is not collinear to X and  $X^{\perp}$ .

<span id="page-6-5"></span>**Lemma 3.6.** If for  $t \in [0, a)$ ,  $\dot{\gamma}(t)$  belongs to the II quadrant, then  $x_1(C_t) > 0$ .

*Proof.* We first treat the case  $t \in (0, a)$ . Expressing  $N(\gamma(t))$  in the  $\{X, X^{\perp}\}\)$  frame, we have thanks to Lemma [3.1](#page-5-2) that

<span id="page-6-2"></span>(3.1) 
$$
0 \ge g_H(N, \dot{\gamma}) = g_H(X, N)g_H(X, \dot{\gamma}) + g_H(X^{\perp}, N)g_H(X^{\perp}, \dot{\gamma}) = g_H(X, N)\sin(\alpha) + g_H(X^{\perp}, N)\cos(\alpha).
$$

If  $\alpha = \pi/2$ , then

<span id="page-6-6"></span>
$$
0 \ge g_H(X, N),
$$

which is possible only when  $\gamma_2(t) = 0$ , that is  $t \in \{0, a\}$ . If  $\alpha \in [0, \pi/2)$  then  $\cos(\alpha) > 0$ , implying that  $g_H(X^{\perp}, N) \leq -g(X, N) \sin(\alpha) \leq 0$ . Notice that this is possible only if  $\gamma_1(t) \geq 0$ . Calling  $-\theta < 0$  the angle that N makes with  $X^{\perp}$ , we get by Equation [\(3.1\)](#page-6-2) that

(3.2) tan(α) ≤ tan(ϑ).

Now, observe that the two geodesic rays  $\sigma_{\gamma}$ ,  $\sigma_N$  stating at  $\gamma(t)$  with initial velocities  $\dot{\sigma}_{\gamma}(0) = \dot{\gamma}^{\perp}(t)$  and  $\dot{\sigma}_{N}(0) = N^{\perp}(\gamma(t))$ , together with the axis  $e_1$  and the geodesic orthogonal to  $e_1$  passing from  $\gamma(t)$  bound two geodesic triangles  $\Delta_{\gamma}$  and  $\Delta_N$ . Call d the distance between  $\gamma(t)$  and  $e_1$ . Then, the length of the sides  $\ell_{\gamma}$  and  $\ell_N$  of  $\Delta_{\gamma}$  and  $\Delta_N$ respectively, lying on  $e_1$  are given via hyperbolic trigonometric laws by

$$
\tanh(\ell_{\gamma}) = \tan(\alpha)\sinh(d) \leq \tan(\vartheta)\sinh(d) = \tanh(\ell_{N}).
$$

But this implies that  $x(C_t)$ , which is the intersection of  $\sigma_\gamma$  with  $e_1$ , has first coordinate positive, as claimed. If  $t = 0$ , then  $C_0 = c_0$  and approximates  $\gamma(0)$  up to the fourth order. Therefore, if  $x_1(C_t) < 0$ , then there exists  $\varepsilon > 0$  such that  $\gamma|_{(\varepsilon,2\varepsilon)}$  lies outside the ball of centered in the origin and with radius  $d_H(\gamma(0), o)$ . This is a contradiction because by construction  $\gamma(0)$  is the furthest point of  $\Omega$  from  $\sigma$ .

Our next goal is to show four important properties of the curve  $\gamma$ . The proof is made by comparison with the circles  $c_t$  and  $C_t$ , and the preservation of the weighted mean curvature  $H_f$ . For this reason, we need the following preliminary lemma.

<span id="page-7-3"></span>**Lemma 3.7.** Let  $\eta = \eta(s)$  be an arc-length, counter-clockwise parametrization of a circle centered at  $(0, y)$  such that  $\eta(0) = (\tau, y)$  and  $\eta(L) = (0, y + \tau)$ . Let  $O = (-\tilde{o}, 0)$  be an arbitrary point lying on  $e_1$  with  $\tilde{o} \in [0, 1)$ , and  $\nu(s)$  the outward pointing normal to  $\eta(s)$ . Then, setting

<span id="page-7-2"></span><span id="page-7-1"></span><span id="page-7-0"></span>
$$
\tilde{H}_1(s) := \partial_{\nu}(h(d_H(\eta(s), O))),
$$

if  $y = 0$  then

(3.3) 
$$
\tilde{H}'_1(s) \le 0, \text{ in } (0, L],
$$

and

(3.4) 
$$
\tilde{H}_1''(0) \le 0.
$$

Both inequalities are strict if  $\tilde{o} \neq 0$ . If  $y \in (0, 1)$  and  $\tilde{o} \neq 0$ , then

(3.5) 
$$
\tilde{H}'_1(L) < 0.
$$

*Proof.* Let  $T: H^2_{\mathbb{R}} \to H^2_{\mathbb{R}}$  be the unique isometry fixing  $e_1$  and sending the origin to O. Then,

$$
\tilde{H}'_1(s) = h''g_H(T_*N, \dot{\eta})g_H(\nu, T_*N) + h'\frac{d}{ds}g_H(\nu, T_*N) \n= h''g_H(T_*N, \dot{\eta})g_H(\nu, T_*N) - h'\Big(g_H(\nabla_{\dot{\eta}}\dot{\eta}^{\perp}, T_*N) + g_H(\dot{\eta}^{\perp}, \nabla_{\dot{\eta}}T_*N)\Big) \n= h''g_H(T_*N, \dot{\eta})g_H(\nu, T_*N) - h'\Big(-g_H(T_*N, \dot{\eta})\kappa_{\eta} + g_H(T_*N, \dot{\eta})g_H(T_*N^{\perp}, \dot{\eta})\kappa_1\Big),
$$

where  $\kappa_1$  is the curvature of the integral curve of  $T_*N^{\perp}$  passing through  $\eta(s)$ , which is a geodesic sphere centered at O. Suppose first that  $y = 0$  and  $\tilde{o} \neq 0$ . Then,  $\dot{\eta} = N^{\perp}$ , and

$$
\tilde{H}'_1(s) = h''g_H(T_*N, N^{\perp})g_H(T_*N, N) - h'g_H(T_*N, N^{\perp})(-\kappa_{\eta} + g_H(T_*N, N)\kappa_1) < 0,
$$

because  $h'' > 0$ ,  $h' > 0$ ,  $g_H(T_*N, N^{\perp}) < 0$ ,  $g_H(T_*N, N) > 0$  and  $\kappa_{\eta} > \kappa_1$  since  $\tilde{o} \neq 0$ . This proves Equation [\(3.3\)](#page-7-0) when  $\tilde{\sigma} \neq 0$ . The same holds in the context of Equation [\(3.5\)](#page-7-1) since  $\eta(L) = N^{\perp}$ . Up to relaxing the inequalities, the proof when  $\tilde{\sigma} = 0$  is exactly the same. To prove Equation [\(3.4\)](#page-7-2), we differentiate  $H_1$  one more time, obtaining

$$
\tilde{H}_{1}''(s) = h''' g_{H}(T_{*}N, \dot{\eta})^{2} g_{H}(\nu, T_{*}N) + h'' \frac{d}{ds} g_{H}(T_{*}N, \dot{\eta}) g_{H}(\nu, T_{*}N) \n+ h'' g_{H}(T_{*}N, \dot{\eta}) \frac{d}{ds} g_{H}(\nu, T_{*}N) + h'' g_{H}(T_{*}N, \dot{\eta}) \frac{d}{ds} g_{H}(\nu, T_{*}N) \n+ h' \frac{d^{2}}{ds^{2}} g_{H}(\nu, T_{*}N).
$$

Observe that in zero  $g_H(T_*N, \dot{\eta}) = 0$ , hence only the second and last term survive

$$
\tilde{H}_1''(0) = h'' \frac{d}{ds} \Big|_{s=0} g_H(T_*N, N^\perp) g_H(T_*N, N) + h' \frac{d^2}{ds^2} \Big|_{s=0} g_H(T_*N, N).
$$

Taking advantage of the explicit expression for  $\frac{d}{ds}g(T_*N, N)$  we obtained before, we get

$$
\frac{d^2}{ds^2}\Big|_{s=0}g_H(T_*N,N) = -\frac{d}{ds}\Big|_{s=0}\Big(g_H(T_*N,N^\perp)\Big(-\kappa_\eta + g_H(T_*N,N)\kappa_1\Big)\Big) \n= (\kappa_\eta - g_H(T_*N,N)\kappa_1)\frac{d}{ds}\Big|_{s=0}g_H(T_*N,N^\perp),
$$

which implies that

$$
\tilde{H}_1''(0) = \underbrace{(h''g_H(T_*N,N) + h'\kappa_\eta - h'g_H(T_*N,N)\kappa_1)}_{>0} \frac{d}{ds}\Big|_{s=0} g_H(T_*N,N^\perp).
$$

Hence, we are left to show that  $\frac{d}{ds}$  $\Big|_{s=0} g_H(T_*N, N^{\perp}) < 0.$  Developing again we get

$$
\frac{d}{ds}\Big|_{s=0}g_H(T_*N, N^{\perp}) = g_H(\nabla_{N^{\perp}}T_*N, N^{\perp}) + g_H(T_*N, \nabla_{N^{\perp}}N^{\perp})|_{s=0}
$$
  
=  $g_H(T_*N, N)^2 \kappa_1|_{s=0} - g_H(TN, N)\kappa_\eta|_{s=0}$   
=  $\kappa_1 - \kappa_\eta < 0$ .

We are now ready to prove the next result.

<span id="page-8-4"></span><span id="page-8-0"></span>Lemma 3.8. The following four points hold.

- i. If for  $t \in (0, a)$  one has that  $\kappa_{\gamma}(t) \geq \kappa(C_t) > 1$ , then  $t \mapsto x_1(C_t)$  is smooth and  $\frac{d}{dt}x_1(C_t) \geq 0.$
- <span id="page-8-2"></span><span id="page-8-1"></span>ii. If  $\gamma$  is not a centered circle, then  $\ddot{\kappa}_\gamma(0) > 0$ .
- iii. If for  $t \in (0, a)$ ,  $\dot{\gamma}(t)$  is in the II quadrant and  $\kappa_{\gamma}(t) = \kappa(C_t) > 1$ , then  $\dot{\kappa}_{\gamma}(t) \geq 0$ . Moreover, if  $\gamma(t) \neq X^{\perp}(\gamma(t))$  and  $C_t$  is not centered in the origin, then  $\kappa_{\gamma}(t) > 0$ .
- <span id="page-8-3"></span>iv. If for  $t \in (0, a)$  one has that  $\gamma(t) = X^{\perp}(\gamma(t)), \gamma_1(t) > 0$  and  $\kappa_{\gamma}(t) \geq \kappa(C_t) > 1$ , then  $\dot{\kappa}_{\gamma}(t) > 0.$

*Proof.* We start with point [i.](#page-8-0) Observe that since  $c_t$  approximates  $\gamma$  up to the third order around  $\gamma(t)$ , it suffices to prove  $\frac{d}{dt}x_1(C_t) \geq 0$  replacing  $\gamma$  with  $c_t$ . Also, we can suppose  $x(c_t)$  on  $e_2$  by composing with the unique hyperbolic isometry translating  $x(c_t)$  on  $e_2$  and fixing  $e_1$ . The curvature condition  $\kappa_{\gamma}(t) \geq \kappa(C_t) > 1$  ensures that  $x(c_t) \in (H_{\mathbb{R}}^2)_+$ . By monotonicity of the function  $tanh(\cdot/2)$ , it suffices to prove the claim for the Euclidean center of  $C_t$ . Thus, we have reduced the problem to an explicit computation in the Euclidean plane, that can be found in [\[2](#page-15-2), Lemma 5.3]. Thanks to Lemma [3.7](#page-7-3) the proofs of the other points go exactly as in [\[2,](#page-15-2) Lemma 3.4, Lemma 3.5 and Lemma 3.7]. We show point [ii.](#page-8-1) Differentiating  $H_f$  twice, we get that

$$
\ddot{\kappa}_\gamma(0) = -(n-2)\ddot{\kappa}(C_0) - H_1''.
$$

 $\Box$ 

By symmetry,  $c_0 = C_0$ . Moreover, since  $\dot{\kappa}_{\gamma}(0) = 0$ , we have that both  $c_0$  and  $C_0$  approximate  $\gamma$  up to the fourth order near zero. Hence,  $\ddot{\kappa}(C_0) = 0$ . Therefore, it suffices to determine the sign of  $H''_1$  replacing  $\gamma$  with  $C_0$ . Let  $T : H^2_{\mathbb{R}} \to H^2_{\mathbb{R}}$  be the unique isometry fixing  $e_1$  that moves  $x(C_0)$  to the origin. The result follows by Equation [\(3.4\)](#page-7-2) of Lemma [3.7](#page-7-3) setting  $O = T(0)$ , and noticing that  $T(0) \neq 0$  by Lemma [3.4.](#page-6-3) The proofs of points [iii.](#page-8-2) and [iv.](#page-8-3) are similar: in the first case the condition  $\kappa_{\gamma}(t) = \kappa(C_t)$  implies that  $c_t = C_t$  approximates  $\gamma$  near t up to the third order, the same holds if  $\dot{\gamma}(t) = X^{\perp}(\gamma(t))$ by symmetry. Hence, substituting  $\gamma$  with  $c_t$  and differentiating one time  $\mathbf{H}_f$ , we have to determine the sign of  $H'_1$  in the case of a circle, via Lemma [3.7.](#page-7-3)

We are now ready to analyse the first behaviour of  $\gamma$ .

Definition 3.9 (Upper curve). The upper curve is the (possibly empty) maximal connected interval  $I_U \subset [0, a)$  such that  $0 \in I_C$  and for all  $t \in I_U$ 

a.  $\dot{\gamma}(t)$  is in the II quadrant,

b.  $\kappa_{\gamma}(t) \geq \kappa(C_t) > 1$ , c.  $t \mapsto x_1(C_t)$  is smooth and  $\frac{d}{dt}x_1(C_t) \geq 0$ .

We set

$$
a_0 := \sup I_U.
$$

In the discussion, we will sometimes identify the upper curve with its image through  $\gamma$ .

**Definition 3.10.** We say that a curve  $\eta$  is graphical with respect to the hypercyclic foliation  $(\delta_l)_{l\in[0,1)}$  if  $\eta$  meets each  $\delta_l$  at most once.

Notice that the upper curve (if non empty) is graphical with respect to the hypercyclical foliation because  $\dot{\gamma}$  is in the II quadrant

<span id="page-9-0"></span>Proposition 3.11. The upper curve is non empty and enjoys the following properties

i.  $0 < a_0 < a$ , ii.  $a_0 \in I_U$ , iii.  $\gamma_1(a_0) > 0$ , *iv.*  $\dot{\gamma}(a_0) = X^{\perp}(\gamma(a_0)).$ 

Proof. Thanks to Lemma [3.8,](#page-8-4) the proof goes exactly as [\[2,](#page-15-2) Lemma 3.11 and Proposition 3.12]. We sketch for completeness the idea behind each point. The upper curve is non empty because by Lemma [3.3](#page-6-4) and Lemma [3.8](#page-8-4) points [i.](#page-8-0) and [ii.](#page-8-1) since  $c_0 = C_0$  approximates  $\gamma$  up to the fourth order near zero, we have that there exists  $\varepsilon > 0$  such that in  $[0, \varepsilon)$ ,  $\gamma$  belongs to the II quadrant,  $\kappa_{\gamma} \geq \kappa(C_t) > 1$  and  $\frac{d}{dt}x_1(C_t) \geq 0$ . Hence,  $[0, \epsilon) \subset I_U$ . Notice that  $0 < a_0$  cannot be equal to a since otherwise the curve  $\gamma$  does not close itself on  $e_1$ , simply because  $\dot{\gamma}$  belongs to the II quadrant by definition of  $I_U$ . By the regularity of  $\gamma$  and that  $I_U$  is defined by three close conditions, we have that  $a_0 \in I_U$ . By composing with the unique hyperbolic isometry sending  $\gamma(a_0)$  on  $e_2$  fixing  $e_1$ , we can see that  $x_1(C_{a_0}) \leq 0$  because  $\gamma(a_0)$  belongs to the II quadrant. Lemma [3.4](#page-6-3) and Lemma [3.6](#page-6-5) imply that  $x_1(C_0) > 0$  and since  $\frac{d}{dt}x_1(C_t) \ge 0$  in  $I_U$ , one must have that  $\gamma_1(a_0) > 0$ . The last point is proved by contradiction: if  $\dot{\gamma}(a_0) \neq X^{\perp}(\gamma(a_0))$ , then  $a_0 \in I_U$  implies that  $\dot{\gamma}(a_0)$  is strictly in the II quadrant. If  $\kappa_{\gamma}(a_0) = \kappa(C_{a_0}) > 1$ , then  $c_{a_0} = C_{a_0}$  approximates  $\gamma$  to the third order and Lemma [3.8](#page-8-4) point [iii.](#page-8-2) implies that there exists some  $\delta > 0$  such that  $\kappa_{\gamma}(t) \geq \kappa(C_t) > 1$  for  $t \in [a_0, a_0 + \delta)$ . The same holds if  $\kappa_{\gamma}(a_0) > \kappa(C_{a_0}) > 1$  by continuity. This means that  $[a_0, a_0 + \delta) \subset I_U$ , which is not possible by the very definition of  $a_0$ . Hence,  $\dot{\gamma}(a_0) = X^{\perp}(\gamma(a_0))$ .

**Definition 3.12** (Lower curve). The lower curve is the maximal connected interval  $I_L \subset$  $[a_0, a)$  such that for all  $t \in I_L$ 

 $\mathbb{C}$  11

a.  $a_0 \in I_L$ ,

b.  $\dot{\gamma}(t)$  is in the III quadrant,

c. calling  $\bar{t} \in I_U$  the unique time such that  $S(\gamma(t)) = S(\gamma(\bar{t}))$  we have that  $\kappa_{\gamma}(t) \geq \kappa_{\gamma}(\bar{t})$ . We set

$$
a_1 := \sup I_L.
$$

Notice that  $a_0$  truly belongs to  $I_L$ , so  $I_L \neq {\}$ . Also, the lower curve is graphical with respect to the hypercyclical foliation because  $\dot{\gamma}$  is in the III quadrant. Our next goal is to prove that  $a_1 < a$ . Again, we proceed by contradiction, and the intuition is the following: suppose that  $a_1 = a$ . If  $\kappa_{\gamma}(t) = \kappa_{\gamma}(t)$  for all  $t \in I_L$ , then the lower curve is nothing else than the upper curve reflected with respect to the geodesic orthogonal to  $e_1$  and passing through  $\gamma(a_0)$ . Hence,  $\lim_{t\to a^+} \alpha(t) = -\frac{\pi}{2}$  $\frac{\pi}{2}$ . Otherwise, if the  $\gamma|_{I_L}$  curves strictly faster than the upper curve at some point, then the angle of incidence  $\lim_{t\to a^+} \alpha(t) < -\frac{\pi}{2}$  $\frac{\pi}{2}$  (see Figure [4\)](#page-11-0). But this cannot be, because it contradicts the regularity of  $\partial E$  points pointed out in Theorem [2.1.](#page-2-0) To prove that the lower curve curves strictly faster than the upper curve we need first to express the curvature with respect to the  $\{X, X^{\perp}\}\$ frame, and next prove three comparison lemmas.

<span id="page-10-0"></span>**Lemma 3.13.** Let  $\eta$  any regular curve parametrized by arclength. Then,

$$
-\kappa_{\eta}(t) = \dot{\beta}(t) - K_1(\eta(t))\cos(\beta(t)),
$$

where  $\beta(t)$  denotes the angle between  $\eta$  and  $X^{\perp}$ , and  $K_1$  is the curvature of the leaf  $\delta_l$ passing through  $\eta(t)$ .

*Proof.* Decompose  $\dot{\eta} = AX + BX^{\perp}$ . Then, since  $\kappa_{\eta} \dot{\eta}^{\perp} = \nabla_{\dot{\eta}} \dot{\eta}$ , we get that

$$
-\cos(\beta)\kappa_{\eta} = g_H(\nabla_{\dot{\eta}}\dot{\eta}, X) = \partial_t(\sin(\beta)) - g_H(\dot{\eta}, \nabla_{\dot{\beta}}X)
$$

.

Now, keeping in mind that  $\nabla_X X = 0$  and  $g_{\mathbb{H}}(\nabla_{X^{\perp}} X^{\perp}, X) = -K_1(\eta(t))$ , we get

$$
g_H\Big(AX + BX^{\perp}, \nabla_{AX + BX^{\perp}}X\Big) = B^2 g_H\Big(X^{\perp}, \nabla_{X^{\perp}}X\Big) = \cos(\beta)^2 K_1(\eta(t)).
$$

Dividing both sides by  $cos(\beta)$  we get the desired identity.

We can prove our first curvature comparison lemma.

<span id="page-10-1"></span>**Lemma 3.14** ( $\kappa$  comparison lemma). Let  $\eta_1 : (0, A_1] \to H^2_{\mathbb{R}}$  and  $\eta_2 : (0, A_2] \to H^2_{\mathbb{R}}$  be two hypercyclical graphical curves parametrized by arclength and with velocity vectors in the II quadrant. Suppose that there exists  $l_0 \in [0, 1)$  such that

$$
\lim_{t \to 0^+} \eta_1(t)
$$
 and 
$$
\lim_{t \to 0^+} \eta_2(t)
$$
,

exist and belong to the same leaf  $\delta_{l_0}$ . Also, suppose that  $\eta_1(A_1) = \eta_2(A_2)$ ,  $\dot{\eta}_1(A_1) = \dot{\eta}_2(A_2)$ , and that if  $S(\eta_1(t)) = S(\eta_2(\tau))$  then  $\kappa_{\eta_1}(t) \geq \kappa_{\eta_2}(\tau)$ . Then, calling  $\alpha_1$  and  $\alpha_2$  the angle made by  $\eta_1$  and  $\eta_2$  with  $X^{\perp}$  we have that

$$
\lim_{t \to 0^+} \alpha_1(t) \ge \lim_{t \to 0^+} \alpha_2(t).
$$

Moreovoer, if for some t and  $\tau$  such that  $S(\eta_1(t)) = S(\eta_2(\tau))$ , one has that  $\kappa_1(t) > \kappa_2(\tau)$ , then

$$
\lim_{t \to 0^+} \alpha_1(t) > \lim_{t \to 0^+} \alpha_2(t).
$$

Proof. Since the curves are graphical with respect to the hypercyclical foliation we can operate a change of variable: we observe that the two height functions  $s_1(t) := S(\eta_1(t))$ and  $s_2(\tau) = S(\eta_2(\tau))$  are bijections with same image of the form  $(l_0, L]$ . By hypothesis  $\kappa_{\eta_1}(s_1^{-1}(l)) \geq \kappa_{\eta_2}(s_2^{-1}(l))$  for every  $l \in (l_0, L]$ . Comparing the two curves in the  $l \in (l_0, L]$ 

variable, since  $s_i' = g_H(\nabla S, \dot{\eta}_i) = g_H(X, \dot{\eta}_i) = \sin(\alpha_i)$ ,  $i = 1, 2$ , we get by Lemma [3.13](#page-10-0) that

$$
0 \leq \kappa_{\eta_1}(l) - \kappa_{\eta_2}(l) = \dot{\alpha}_2(l) \sin(\alpha_2(l)) - \dot{\alpha}_1(l) \sin(\alpha_1(l)) - \frac{2l}{1+l^2} (\cos(\alpha_2(l)) - \cos(\alpha_1(l))).
$$

Multiplying by  $(1 + l^2)$  and integrating we finally get that

$$
0 \leq \int_{l_0}^{L} (1 + l^2) (\cos(\alpha_1) \cos(\alpha_2))' + 2l(\cos(\alpha_1) - \cos(\alpha_2)) dl
$$
  
= 
$$
\int_{l_0}^{L} \frac{d}{dl} ((1 + l^2) (\cos(\alpha_1) - \cos(\alpha_2))) dl = \lim_{l \to l_0^+} (1 + l^2) (\cos(\alpha_2(l)) - \cos(\alpha_1(l))).
$$

If the two curvatures are different somewhere, then the inequality between the two angles is strict.



<span id="page-11-0"></span>FIGURE 4. The curvature comparison.

<span id="page-11-1"></span>**Lemma 3.15** ( $\kappa(C_t)$  comparison lemma). Let  $\eta_1$ ,  $\eta_2$  as in Lemma [3.14.](#page-10-1) Then, for any two points  $\eta_1(t_1)$  and  $\eta_2(t_2)$  on the same leaf  $\delta_l$ , calling  $C^1$  and  $C^2$  the comparison circles at  $\eta_1(t_1)$  and  $\eta_2(t_2)$  as in Definition [2.2,](#page-4-2) we have that

$$
\kappa(C^1) \le \kappa(C^2).
$$

*Proof.* For  $i = 1, 2$ , the hyperbolic radius of  $C<sup>i</sup>$  together with  $e<sub>1</sub>$  and the geodesic starting from  $\eta_i(t_i)$  and hitting  $e_1$  perpendicularly bound a geodesic triangle  $\Delta_i$ . Let  $d_1^i$  be the hyperbolic radius,  $d_2^i$  be the side touching  $e_1$  and  $d_3^i$  the the remaining side of  $\Delta_i$ . Similarly, for  $i = 1, 2$  and  $j = 1, 2, 3$ , call  $\beta_j^i$  the angle opposite to  $d_j^i$ , and  $\ell_i^j$  $i$ <sup>the</sup> length of  $d_j^i$ . We refer to Figure [5.](#page-12-0) By construction  $\beta_1^1 = \beta_1^2 = \frac{\pi}{2}$  $\frac{\pi}{2}$ ,  $\beta_2^i = \alpha_i$ , and since  $\eta_1(t_1)$  and  $\eta_2(t_2)$  are in the same hypercycle by hypothesis, we get  $\ell_3^1 = \ell_3^2$ . Then, by the hyperbolic law of cosines and by Lemma [3.14](#page-10-1) we get

$$
\kappa(C^1) = \coth(\ell_1^1) = \frac{\cos(\alpha_1)}{\tanh(\ell_3^1)} \le \frac{\cos(\alpha_2)}{\tanh(\ell_3^2)} = \coth(\ell_1^2) = \kappa(C^2).
$$

<span id="page-11-2"></span>**Lemma 3.16** ( $H_1$  comparison lemma). Let  $\eta_1$ ,  $\eta_2$  as in Lemma [3.14](#page-10-1) and let  $\tilde{\eta}_1$  be the reflection of  $\eta_1$  with respect to the geodesic passing through  $\eta_1(A_1)$  and crossing  $e_1$  perpendicularly. Reverse its parametrization, so that the angle that  $\tilde{\eta}_1$  makes with  $X^{\perp}$  is equal to  $\tilde{\alpha}_1 = -\alpha_1$ . Moreover, suppose that

$$
g_H(N, \dot{\eta}_2) \leq 0.
$$

 $\overline{I}$ 



<span id="page-12-0"></span>FIGURE 5. The curvature comparison for  $C_t$ .

Denote the unitary outward pointing normals to  $\eta_1$  and  $\eta_2$  by  $\tilde{\nu}_1$  and  $\nu_2$ . Then, for any two points  $\tilde{\eta}_1(t_1)$  and  $\eta_2(t_2)$  on the same leaf  $\delta_l$  we have that

$$
g_H(N(\tilde{\eta}_1(t_1)), \tilde{\nu}_1(t_1)) \le g_H(N(\eta_1(t_2)), \nu_2(t_2)),
$$

with equality if and only if  $\eta_2(t_2)$  and  $\dot{\eta}_1(t_1)$  are tangent to the same circle centered in the origin and  $\dot{\tilde{\eta}}_1(t_1) = -\dot{\eta}_2(t_2)$ .

*Proof.* Let  $\vartheta(s)$  be the angle that N makes with  $X^{\perp}$  at  $\delta_l(s)$  and  $s_2 < s_1$  be such that  $\delta_l(s_1) = \tilde{\eta}_1(t_1)$  and  $\delta_l(s_2) = \eta_1(t_2)$ . Setting  $\Theta_1 := \vartheta(s_1) - \tilde{\alpha}_1 - \frac{\pi}{2}$  $\frac{\pi}{2}$  and  $\Theta_2 := \vartheta(s_2) - \alpha_2 - \frac{\pi}{2}$  $\frac{\pi}{2}$ we get that

$$
g_H(N(\tilde{\eta}_1(t_1)), \tilde{\nu}_1(t_1)) = \cos(\Theta_1),
$$

and

$$
g_H(N(\eta_2(t_2)), \nu_2(t_2)) = \cos(\Theta_2).
$$

Let  $s_2^*$  $\frac{1}{2}$  such that the unit vector at  $\delta_l(s_2^*)$ <sup>\*</sup>/<sub>2</sub>) that forms an angle of  $\alpha_2$  with  $X^{\perp}$  is tangent to a circle centered in the origin. The value  $s_2^*$  must exist and it is greater than  $s_1$  because by Lemma [3.6,](#page-6-5) Equation [\(3.2\)](#page-6-6), we have that  $\vartheta(s_2) \geq \alpha_2$ . Now, on the intersection of  $\delta_l$  with  $e_1$  we have  $\vartheta = 0$ , hence by continuity there must be a point  $s_2^*$  between this intersection and  $\delta_l(s_1)$  such that  $\vartheta(s_2^*)$  $i_2^*$ ) =  $\alpha$ . Then

$$
\Theta_2 = \vartheta(s_2) - \vartheta(s_2^*) = \int_{s_2^*}^{s_2} \dot{\vartheta}(s) \, ds.
$$

Let  $s_1^* = -s_2^*$  $_2^*$ , and notice that  $\vartheta(-s) = \pi - \vartheta(s)$  for every  $s \geq 0$ . Then,

$$
\Theta_1 = \vartheta(s_1) - \tilde{\alpha}_1 - \frac{\pi}{2} = \vartheta(s_2) + \alpha_2 - (\tilde{\alpha}_1 + \alpha_2) - \frac{\pi}{2}
$$
  
=  $\vartheta(s_1) + \vartheta(s_2^*) - \pi - (\tilde{\alpha}_1 + \alpha_2)$   
=  $\vartheta(s_1) - \vartheta(-s_2^*) - (\tilde{\alpha}_1 + \alpha_2)$   
=  $-\int_{s_1}^{s_1^*} \dot{\vartheta}(s) ds - (\tilde{\alpha}_1 + \alpha_2).$ 

Hence,

$$
\Theta_2 - \Theta_1 = (\tilde{\alpha}_1 + \alpha_2) + \int_{s_2^*}^{s_2} \dot{\vartheta}(s) \, ds + \int_{s_1}^{s_1^*} \dot{\vartheta}(s) \, ds \le 0,
$$

since  $\dot{\vartheta}(s) < 0$  and  $\tilde{\alpha}_1 + \alpha_2 \le 0$  by Lemma [3.14.](#page-10-1) The equality is attained only when  $s_0^* = s_0$ ,  $s_1^* = s_1$  and  $\alpha_2 = \alpha_1$  as predicted  $s_2^* = s_2, s_1^* = s_1$  and  $\alpha_2 = \alpha_1$  as predicted.

We can prove the main result about the lower curve.

<span id="page-13-0"></span>**Proposition 3.17.** It  $\gamma$  is not a circle centered in the origin, the lower curve is contained in  $[a_0, a)$ , that is  $0 < a_1 < a$ . Furthermore,  $a_1 \in I_L$  and  $\dot{\gamma}(a_1) = -X(\gamma(a_1)).$ 

*Proof.* By property [iv.](#page-8-3) of Lemma [3.8,](#page-8-4)  $a_1 > a > 0$ . Suppose by contradiction that  $a_1 = a$ . Set  $\tilde{\eta}_1$  to be the (reparametrized) lower curve and  $\eta_2$  the upper curve. Choose any point  $t \in I_L$  with corresponding  $\bar{t} \in I_U$ . Applying Lemma [3.15](#page-11-1) and Lemma [3.16](#page-11-2) to  $\tilde{\eta}_1(t)$  and  $\eta_2(\bar{t})$ , and taking advantage of the expression for  $H_f$  given in Lemma [2.4,](#page-5-3) we get that

$$
\mathbf{H}_f(\gamma(\bar{t})) = \mathbf{H}_f(\gamma(t)) = \kappa_\gamma(t) + (n-2)\kappa(C_t) + h'(d_H(0, \gamma(t)))g_H(\nu(t), N)
$$
  
< 
$$
< \kappa_\gamma(t) + (n-2)\kappa(C_{\bar{t}}) + h'(d_H(0, \gamma(t)))g_H(\nu(t), N).
$$

We have that

$$
d_H(0,\gamma(t)) \le d_H(0,\gamma(\bar{t})).
$$

This can be verified again via the trigonometric rules for hyperbolic triangles: fix  $t \in I_U$ , and call  $\beta$  and  $\bar{\beta}$  the angle that N makes with  $X(\gamma(t))$  and  $X(\gamma(\bar{t}))$  respectively. Notice that  $0 \le \beta \le \bar{\beta}$ . Then, calling d the distance of  $\gamma(t)$  and  $\gamma(\bar{t})$  from  $e_1$ , we get that

$$
\tanh(d_H(0,\gamma(t))) = \frac{\tanh(d)}{\cos(\beta)} \le \frac{\tanh(d)}{\cos(\overline{\beta})} = \tanh(d_H(0,\gamma(\overline{t}))).
$$

Hence

$$
\mathbf{H}_f(\gamma(\bar{t})) < \kappa_\gamma(t) + (n-2)\kappa(C_{\bar{t}}) + h'(d_H(0,\gamma(\bar{t})))g_H(\nu(\bar{t}),N),
$$

implying

$$
\kappa_{\gamma}(\bar{t}) < \kappa_{\gamma}(t),
$$

since h is strictly convex. Again, Lemma [3.14](#page-10-1) tells us that the lower curve hits the  $e_1$ axis with an angle strictly smaller than  $-\frac{\pi}{2}$  $\frac{\pi}{2}$ . Contradiction. Therefore,  $a_1 < a$ . Since  $\gamma$  is smooth in  $a_1 < a$ , and the conditions on  $I_L$  are closed, we deduce that  $a_1 \in I_L$ . Suppose now that  $\alpha(a_1)$  is strictly in the III quadrant. Since  $a_1 \in I_L$ , we can apply again the comparison lemmas to  $\gamma(a_1)$  and  $\gamma(\bar{a}_1)$  to infer

$$
\kappa_{\gamma}(\bar{a}_1) < \kappa_{\gamma}(a_1).
$$

By continuity of  $\kappa_{\gamma}$  and  $\dot{\gamma}$  around  $a_1$ , we get that there exists a neighbourhood of  $a_1$  in which  $\dot{\gamma}$  is in the III quadrant and the above inequality holds in the not strict sense. But this implies that  $a_1$  is not the supremum of  $I_L$ . Therefore, the velocity vector of  $\gamma$  at  $a_1$ has to be equal to  $-X$ .

We prove the last part of the tangent lemma.

<span id="page-13-1"></span>**Proposition 3.18.** If  $\gamma$  is not a centered circle, then there exists  $0 < a_1 < a_2$  such that  $\dot{\gamma}(a_2) = X(\gamma(a_2)).$ 

*Proof.* If  $a_2$  exists we are done. Otherwise, we show that the non existence contradicts the mean-curvature convexity of  $\Omega$ . Let

 $I_c := \{t \in [a_1, a) : \dot{\gamma} \text{ is in the I or IV quadrant}\}.$ 

Here the index stands for *curl curve*. Set  $\tilde{a}_2 := \sup I_c$ . Since  $\kappa(a_1) > 1$  we have that  $a_1 < \tilde{a}_2$ . If  $\tilde{a}_2 < a$ , then the mean convexity of  $\Omega$  implies that

$$
\kappa_{\gamma}(\tilde{a}_2) \ge (n-1) - (n-2)\kappa(C_{\tilde{a}_2}) > 0.
$$

To see this, move  $\gamma(\tilde{a}_2)$  on  $e_2$  as in Lemma [3.8.](#page-8-4) Then,  $C_{\tilde{a}_2}$  is oriented clockwise, and hence has negative curvature. But this implies that we can extend  $I_c$  after  $\tilde{a}_2$ , contradicting the definition of  $\tilde{a}_2$ . So, we need to rule out the situation in which  $\tilde{a}_2 = a$ . If it is the case, then again for mean-convexity one has that in  $I_c$ 

$$
\kappa_{\gamma}(t) > 1.
$$

Moreover, for  $t \in I_c \setminus \{a_1\}$  we have that  $\dot{\gamma}$  lies in the IV quadrant, because otherwise this implies together with  $\kappa_{\gamma}(t) > 0$  that  $\gamma$  cannot close at  $e_1$ . Therefore  $\alpha(t)$  lies in the IV quadrant and it is strictly increasing, implying that

$$
\lim_{t \to a^+} \alpha(t) < -\frac{\pi}{2}.
$$

This cannot be because of the before mentioned regularity properties of isoperimetric  $sets.$ 

The proof of the tangent lemma is then the collection of the results we showed in this section.

<span id="page-14-0"></span>*Proof of Lemma [3.2.](#page-5-1)* The existence of the chain  $0 < a_0 < a_1 < a_2 < a$  is ensured by Proposition [3.11,](#page-9-0) Proposition [3.17](#page-13-0) and Proposition [3.18.](#page-13-1)

# 4. SYMMETRIC SETS IN  $H_{\mathbb{K}}^m$

Consider any rank one symmetric space of non-compact type  $(M^n, g) = (H^m_{\mathbb{K}}, g)$ ,  $\mathbb{K} \in$  $\{\mathbb C, \mathbb H, \mathbb O\}$ . Set  $d = \dim(\mathbb K) \in \{2, 4, 8\}$  so that the real dimension of M is  $n = md$ . Recall that if  $\mathbb{K} = \mathbb{O}$ , we only have the Cayley plane  $H_0^2$ . As classical references on symmetric spaces we cite the books of Eberlein [\[4](#page-15-6)] and Helgason [\[6\]](#page-15-7). Fix an arbitrary base point  $o \in M$ , and let N be the unit-length, radial vector field emanating from it. As in Definition [1.2,](#page-1-0) let H be the distribution on  $M \setminus \{o\}$  induced by the  $(-4)$ -eigenspace of the Jacobi operator  $R(\cdot, N)N$ . Denote with V the orthogonal complement of H with respect to g. For every  $x \in M \setminus \{o\}$ , we have the orthogonal splitting

$$
T_xM=\mathcal{H}_x\oplus \mathcal{V}_x,
$$

with orthogonal projections  $(\cdot)^{\mathcal{H}}$  and  $(\cdot)^{\mathcal{V}}$ . Let now  $(\bar{M}^n, g_H) = (H^n_{\mathbb{R}}, g_H)$ , and choose an arbitrary base point  $\bar{\sigma}$  in it. The isometric identification of  $T_oM$  with  $T_{\bar{o}}M$  according to the flat metrics  $(\exp_o^M)^* g|_0$  and  $(\exp_{\bar{o}}^{\bar{M}})^* g_H|_0$ , induces a well defined diffeomorphism

$$
\Psi = \exp_{\bar{o}}^{\bar{M}} \circ (\exp_o^M)^{-1} : M \to \bar{M}.
$$

With a slight abuse of notation, we still denote with  $g_H$  the metric  $\Psi^* g_H$ , that makes M isometric to  $\overline{M}$ . The following lemma allows us to compare q with  $q_H$ .

<span id="page-14-2"></span>**Lemma 4.1.** For every  $x \in M \setminus \{o\}$ , the splitting

$$
T_xM=\mathcal{H}_x\oplus \mathcal{V}_x,
$$

is orthogonal with respect to  $g_H$ . In particular, letting  $d_H$  be the Riemannian distance induced by  $g_H$  on M, one has that

<span id="page-14-1"></span>(4.1) 
$$
g(v, w) = \cosh^2(d_H(o, x))g_H(v^{\mathcal{H}}, w^{\mathcal{H}}) + g_H(v^{\mathcal{V}}, w^{\mathcal{V}}),
$$

for all  $v, w \in T_xM$ .

*Proof.* Fix an arbitrary unit direction  $N_o \in T_oM$ , and let  $V_o \in T_oM$  be any vector orthogonal to it with respect to  $g|_o = g_H|_o$ . Since the radial geodesics emanating from o are the same for g and  $g_H$ , the Jacobi field  $Y(t)$  along the geodesic  $\sigma : t \mapsto \exp_{o}^{M}(tN_o)$ , determined by the initial conditions  $Y(0) = 0$ ,  $\dot{Y}(0) = V_o$  is the same for both metrics. Let  $V(t)$  and  $V_H(t)$  be the parallel transport of  $V_o$  along  $\sigma$  with respect to g and  $g_H$ , respectively. By the very definition of symmetric spaces, the curvature tensor  $R$  is itself parallel along geodesics. This implies that

$$
\sinh(t)V_H(t) = Y(t) = \frac{\sinh(\sqrt{-\kappa}t)}{\sqrt{-\kappa}}V(t),
$$

#### 16 LAURO SILINI

provided  $V_o$  belongs to the *κ*-eigenspace of the Jacobi operator  $R(\cdot, N_o)N_o$ . Therefore, parallel vector fields in the eigenspaces are collinear for the two metrics. Hence, for  $t > 0$  the linear subspaces  $\mathcal{H}_{\sigma(t)}$  and  $\mathcal{V}_{\sigma(t)}$  are nothing else than the parallel transport of the corresponding eigenspaces of  $R(\cdot, N_o)N_o$  along  $\sigma$ . It follows that the splitting  $T_xM = H_x \oplus V_x$  is orthogonal not only with respect to g, but also with respect to the hyperbolic metric  $g_H$ . Equation [\(4.1\)](#page-14-1) is a direct consequence of this fact and the definition of the distribution  $\mathcal{H}$ .

We can now prove Theorem [1.5.](#page-1-2)

*Proof of Theorem [1.5.](#page-1-2)* Let  $E \subset M$  be an Hopf-symmetric set with outward pointing normal vector field  $\nu$  with respect to q. By the very definition of being Hopf-symmetric,  $\nu^{\mathcal{H}} \equiv 0$ . Therefore, thanks to Lemma [4.1,](#page-14-2)  $\nu$  is orthonormal to  $\partial E$  also with respect to  $g_H$ . Let vol and vol<sub>H</sub> the volume forms associated to g and  $g_H$ . We have that

$$
P(E) = \int_{\partial E} \iota_{\nu} \operatorname{vol} = \int_{\partial E} \cosh^{d-1}(d_H(o, x)) \iota_{\nu} \operatorname{vol}_H(x),
$$

where  $\iota : \Omega(M)^p \to \Omega(M)^{p-1}$  denotes the interior product  $\iota_X \alpha(\cdot) = \alpha(X, \cdot)$ . The volume of  $E$  is given by the formula

$$
V(E) = \int_E \text{vol} = \int_E \cosh^{d-1}(d_H(o, x)) \text{ vol}_H(x).
$$

Hence, the volume and perimeter of Hopf-symmetric sets in  $M$  correspond to the volume and perimeter of  $\Psi(E)$  in  $H_{\mathbb{R}}^n$  with density equal to  $h(r) = \cosh^{d-1}(r)$ , concluding the proof.  $\Box$ 

## **REFERENCES**

- <span id="page-15-3"></span>[1] Eliot Bongiovanni et al. "Isoperimetry in Surfaces of Revolution with Density". Missouri Journal of Mathematical Sciences 30.2 (Nov. 2018). DOI: [10.35834/mjms/1544151692](https://doi.org/10.35834/mjms/1544151692). url: <https://doi.org/10.35834%2Fmjms%2F1544151692>.
- <span id="page-15-2"></span>[2] Gregory R Chambers. "Proof of the log-convex density conjecture". Journal of the European Mathematical Society 21.8 (2019), pp. 2301–2332.
- <span id="page-15-4"></span>[3] Leonardo Di Giosia et al. "The Log Convex Density Conjecture in Hyperbolic Space" (2016). DOI: [10.48550/ARXIV.1610.07043](https://doi.org/10.48550/ARXIV.1610.07043). URL: <https://arxiv.org/abs/1610.07043>.
- <span id="page-15-6"></span>[4] Patrick Eberlein. Geometry of nonpositively curved manifolds. eng. Chicago lectures in mathematics series. Chicago: University of Chicago Press, 1996. ISBN: 0226181987.
- <span id="page-15-0"></span>[5] Alessio Figalli and Francesco Maggi. "On the isoperimetric problem for radial log-convex densities". Calculus of Variations and Partial Differential Equations 48.3 (2013), pp. 447–489.
- <span id="page-15-7"></span>[6] Sigurdur Helgason. Differential geometry, Lie groups, and symmetric spaces. eng. Graduate studies in mathematics, volume 34. Providence, Rhode Island: American Mathematical Society, 2001 - 1978. ISBN: 1-4704-1146-6.
- <span id="page-15-5"></span>[7] Sean Howe. "The Log-Convex Density Conjecture and vertical surface area in warped products". Advances in Geometry 15.4 (2015), pp. 455–468. DOI: [doi:10.1515/advgeom-2015-0026](https://doi.org/doi:10.1515/advgeom-2015-0026). url: <https://doi.org/10.1515/advgeom-2015-0026>.
- <span id="page-15-1"></span>[8] Alexander Kolesnikov and Roman Zhdanov. "On isoperimetric sets of radially symmetric measures". Concentration, Functional inequalities and Isoperimetry, Contemp. Math. 545 (Feb. 2010). DOI: [10.1090/conm/545/10769](https://doi.org/10.1090/conm/545/10769).
- <span id="page-16-0"></span>[9] Francesco Maggi. Sets of Finite Perimeter and Geometric Variational Problems: An Introduction to Geometric Measure Theory. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2012. DOI: [10.1017/CBO9781139108133](https://doi.org/10.1017/CBO9781139108133).
- <span id="page-16-4"></span>[10] Ivor E. McGillivray. "A weighted isoperimetric inequality on the hyperbolic plane" (2017). doi: [10.48550/ARXIV.1712.07690](https://doi.org/10.48550/ARXIV.1712.07690). url: <https://arxiv.org/abs/1712.07690>.
- <span id="page-16-2"></span>[11] Frank Morgan. "Regularity of isoperimetric hypersurfaces in Riemannian manifolds". Transactions of the American Mathematical Society 355.12 (2003), pp. 5041–5052.
- <span id="page-16-5"></span>[12] Frank Morgan, Michael Hutchings, and Hugh Howards. "The isoperimetric problem on surfaces of revolution of decreasing Gauss curvature". Transactions of the American Mathematical Society 352.11 (2000), pp. 4889–4909.
- <span id="page-16-3"></span>[13] Frank Morgan and Aldo Pratelli. "Existence of isoperimetric regions in  $\mathbb{R}^n$  with density". Annals of Global Analysis and Geometry 43.4 (2013), pp. 331–365.
- <span id="page-16-1"></span>[14] César Rosales et al. "On the isoperimetric problem in Euclidean space with density". Calculus of Variations and Partial Differential Equations 31.1 (2008), pp. 27–46.
- <span id="page-16-6"></span>[15] Xiaodong Wang. "An integral formula in Kähler geometry with applications". Communications in Contemporary Mathematics 19.05 (2017), p. 1650063.