How fast can quantum brachistochrone evolve?

Fu-Quan Dou^{*} and Min-Peng Han

College of Physics and Electronic Engineering, Northwest Normal University, Lanzhou, 730070, China

The quantum speed limit (QSL) represents the cornerstone of essential physical limit on the speed of evolution process in quantum mechanics. To derive the QSL for arbitrary quantum systems, here we present a geometric method, which is estiblished on the quantum brachistochrone characterized by the Riemannian metric. As an application to the complex quantum system, we consider the nonlinear Landau-Zener type two-level system, representing coherent atoms for BECs with mean-field interaction. The analytical expression of the quantum brachistochrone and population are obtained for the nonlinear system. We further derive the corresponding optimal procotol and demonstrate that the QSL is tight. We also generalize our formulation to open quantum systems.

Introduction. How fast can quantum brachistochrone (QB) evolve? This issue is of the essence in pratically all domains of quantum physics expecting to achieve quantum speed limit (QSL), such as quantum computation [1, 2] and quantum control [3, 4]. As the fundamental principle of quantum mechanics, the QSL displays the bound on the minimal evolution time of quantum systems between two given states on the quantum-state space [2, 4-7]. At present, the potential of approaching QSL by quantum control has also been demonstrated in experiments [8-11]. However, in different systems, the factors affecting the QSL are complicated [12-19]. In addition, accurate analytic solutions of optimal protocol are always subject to mathematical or numerical difficulties [20, 21]. The quantum brachistochrone equation (QBE) can provide the optimal protocol on the premise of satisfying the task fault tolerance, which is considered to be a promising method [20]. Based on the general framework [20, 22–24], QBE gradually plays an important role in quantum computation and quantum control of various models [25–28], especially in parity-time (PT) symmetric systems [29–31] and adiabatic evolution, i.e., the quantum adiabatic brachistochrone [32–35]. The corresponding numerical methods have also been proposed in succession [20, 21, 33, 34]. The relationship of quantum coherence [36] and entanglement [37, 38] to QB is also discussed.

In this Letter, we present a geometric method established on the QB characterized by the Riemannian metric to derive the QSL for arbitrary quantum systems with unitary dynamics. In contrast to the Mandelstam-Tamm and the Margolus-Levitin inequalities, we obtain a bound which is of geometric properties and expressed as the global and local geometric quantum speed estimation, i.e., Eq. (10). We apply this formulation to the nonlinear Landau-Zener (LZ) type two-level system to investigate the evolution speed of the complex system. Compared with the linear case, it is far from fully understanding the dynamical evolution of the nonlinear twolevel model. In order to derive the QB of this model, we find the analytical expression of population represented by the Bloch parameterization. As this basis, we obtain

the QB and analytical optimal protocol corresponding to the QSL via the QBE and further prove that the QSL is tight. These results are consistent with the result given by the experiment [8] and optimal control [39, 40] in the linear limit. Particularly, the application to this model reveals the relationship between the optimal control and the traditional geometric approach which focuses on estimating for the maximal quantum speed [2]: As common methods to calculate the minimal evolution time, they are actually different aspects of the QSL. Specifically, the optimal control and the traditional geometric approach contribute the local and global geometric quantum speed estimation to the QSL respectively. In fact, there is no obstacle to the application of this geometric framework to open system dynamics. Finally, we generalize our formulation to open quantum systems.

Evolution speed. The QB refers to the time-optimal path \mathscr{L}_{op} on the quantum-state space given by the functional $\tau[\bullet] = \int d\mathscr{L}/\mathcal{V}$, which satisfies $\delta\tau/\delta\mathscr{L} = 0$. Here, $d\mathscr{L}$ donates the differential of Bures angle [41, 42], which represents the Bures angle moving from ψ to $\psi + d\psi$ through unitary evolution and $\mathcal{V} = d\mathscr{L}/dt$ characterizes the geometric quantum speed [2, 43]. On the Hilbert space, $\mathscr{L} = \arccos(|\langle \psi_0 | \psi_\tau \rangle|)$ which can be written in the tensor form by the Fubini-Study metric [42, 44]

$$d\mathscr{L}^2 = \sum_{i=1}^n \frac{dp_i^2}{p_i} + 4\left[\sum_{i=1}^n p_i d\varphi_i^2 - \sum_{i,j=1}^n p_i p_j d\varphi_i d\varphi_j\right], \quad (1)$$

where $p_{i,j}$ and $\varphi_{i,j}$ correspond to the amplitude and phase of the eigenstates respectively. For general representation, considering a set of parameters $\{\lambda_i(t)\}, i =$ $1, \ldots, 2n$ to describe analytically quantum states such as $p_{i,j}$, $\varphi_{i,j}$ or the Bloch vector [44–46]. Over the evolutionary duration $0 \leq t \leq \tau$, the dynamic process of the quantum system draws geometrically a curve varying over the Riemannian manifold \mathscr{M} from the initial state $|\psi(\lambda_1(0), \ldots, \lambda_{2n}(0))\rangle$ to the final state $|\psi(\lambda_1(\tau), \ldots, \lambda_{2n}(\tau))\rangle$. First of all, reparameterization of functional $\tau[\bullet]$ is necessary. To unify the coordinates, we introduce

$$x_1 = p_1, \ldots, x_n = p_n, x_{n+1} = \varphi_1, \ldots, x_{2n} = \varphi_n.$$

Thus Eq. (1) becomes $d\mathscr{L}^2 = \sum_{i,j=1}^{2n} g_{ij} dx_i dx_j$, where

$$g_{ij} = \begin{cases} \delta_{ij}/x_i, & \text{if } max\{i,j\} \le n, \\ 4\delta_{ij}x_{i-n} - 4x_{i-n}x_{j-n}, & \text{if } min\{i,j\} > n, \\ 0, & \text{otherwise.} \end{cases}$$
(2)

Under the coordinate transformations of arbitrary parameters λ , the property of tensor produces the form

$$d\mathscr{L}^2 = \sum_{\mu,\nu=1}^{2n} g_{\mu\nu} d\lambda_\mu d\lambda_\nu, \qquad (3)$$

where $g_{\mu\nu} = \sum_{i,j=1}^{2n} \partial_{\lambda_{\mu}} x_i \ \partial_{\lambda_{\nu}} x_j$. $\tau[\bullet]$ is further reparameterized explicitly

$$\tau[\bullet] = \int \frac{\sqrt{\sum_{\mu,\nu=1}^{2n} g_{\mu\nu} \frac{d\lambda_{\mu}}{d\zeta} \frac{d\lambda_{\nu}}{d\zeta}}}{\frac{d}{dt} \mathscr{L}(|\psi_0\rangle, |\psi_{\tau}\rangle)} d\zeta, \qquad (4)$$

where \hbar is set to be one and ζ is the natural parameter selected. QBE requires the Euler-Lagrange (EL) equations $d\partial_{\dot{\lambda}_i} \mathscr{L}(\lambda_1, \ldots, \lambda_{2n})/d\zeta = \partial_{\lambda_i} \mathscr{L}(\lambda_1, \ldots, \lambda_{2n}), i =$ $1, \ldots, 2n$. Now we aim at deriving the bound on the functional τ for the QB on the manifold \mathscr{M} . Utilizing the Riemannian metric Eq. (3), geometric quantum speed reads $\mathcal{V} = d\mathscr{L}(|\psi_0\rangle, |\psi_\tau\rangle)/dt = \sqrt{\sum_{\mu,v=1}^{2n} g_{\mu v} v_{\mu} v_v}$, where $v_{\mu}(\lambda_1, \ldots, \lambda_{2n}) = d\lambda_{\mu}/dt$. \mathcal{V} and v_{μ} are the global evolution speed and local evolution speed respectively. Selecting arbitrary component λ_i as the natural parameter, Eq. (4) becomes

$$\tau[\bullet] = \int \frac{\sqrt{\sum_{\mu,\nu=1}^{2n} g_{\mu\nu} \frac{d\lambda_{\mu}}{d\lambda_{i}} \frac{d\lambda_{\nu}}{d\lambda_{i}}}}{\sqrt{\sum_{\mu,\nu=1}^{2n} g_{\mu\nu} \frac{d\lambda_{\mu}}{dt} \frac{d\lambda_{\nu}}{dt}}} d\lambda_{i} = \int \frac{d\lambda_{i}}{v_{i}}.$$
 (5)

For any parameter λ_i , the dynamics of the element on the manifold \mathcal{M} is limited by each velocity component v_i . These simultaneous 2n inequalities lead to

$$\tau[\bullet] \ge \max_{i=1,\dots,2n} \left\{ \frac{\mathfrak{L}_i(|\psi_0\rangle, |\psi_\tau\rangle)}{|v_i|_{max}} \right\} = \frac{\mathfrak{L}(|\psi_0\rangle, |\psi_\tau\rangle)}{\mathscr{V}}, \quad (6)$$

where $\mathfrak{L}_i(|\psi_0\rangle, |\psi_\tau\rangle) = |\lambda_i(0) - \lambda_i(\tau)|, \mathfrak{L}(|\psi_0\rangle, |\psi_\tau\rangle)$ and \mathscr{V} describe distance and speed respectively corresponding to the component which contribute the most. Likewise, there exists estimation for the global geometric quantum speed $|\mathcal{V}_{max}|$

$$\tau[\bullet] \ge \frac{\mathscr{L}(|\psi_0\rangle, |\psi_\tau\rangle)}{|\mathcal{V}_{max}|}.$$
(7)

Mathematically we consider a concise example to show that Eq. (6) and Eq. (7) are independent of each other. Assume that \mathcal{V} approaches maximum

when each component is optimal, i.e., $|\mathcal{V}_{max}| = \sqrt{\sum_{\mu,\nu=1}^{2n} g_{\mu\nu} |v_{\mu}|_{max} |v_{\nu}|_{max}}$. For arbitrary component λ_i and the corresponding speed v_i there holds the relation $\int d\lambda_i / v_i = \int d\mathscr{L} / \mathcal{V}$ given by Eq. (5). Then the bound follows the form

$$\tau[\bullet] \ge \max\left\{\int \frac{d\mathscr{L}}{\frac{|v_i|_{max}}{v_i}}, \int \frac{d\mathscr{L}}{\frac{|\mathcal{V}_{max}|}{\mathcal{V}}}\right\}.$$
 (8)

Eq. (8) indicates that the independence of Eqs. (6) and (7) depends on whether there exists strict unequal relationship between $|\mathcal{V}_{max}|/\mathcal{V}$ and $|v_i|_{max}/v_i$. The formulation $\max\left\{\frac{\alpha'}{\alpha} + \frac{\beta'}{\beta}\right\} \geq \frac{\alpha' + \beta'}{\alpha + \beta} \geq \min\left\{\frac{\alpha'}{\alpha} + \frac{\beta'}{\beta}\right\}$ provides the following answer:

$$\min\left\{\frac{|v_i|_{max}}{v_i}, \sqrt{\sum_{\mu\neq\nu}^{2n} g_{\mu\nu} |v_\mu|_{max} |v_\nu|_{max}} / \mathcal{V}\right\} \le \frac{|\mathcal{V}_{max}|}{\mathcal{V}}$$
$$\le \max\left\{\frac{|v_i|_{max}}{v_i}, \sqrt{\sum_{\mu\neq\nu}^{2n} g_{\mu\nu} |v_\mu|_{max} |v_\nu|_{max}} / \mathcal{V}\right\}, \quad (9)$$

which means that the unequal relationship between Eqs. (6) and (7) depends on the contribution of the nondiagonal elements of Riemannian metric to Eq. (9). As a consequence, the QSL should take the upper bound of Eqs. (6) and (7):

$$\tau_{QSL} = \max\left\{\frac{\mathscr{L}(|\psi_0\rangle, |\psi_\tau\rangle)}{|\mathcal{V}_{max}|}, \frac{\mathfrak{L}(|\psi_0\rangle, |\psi_\tau\rangle)}{\mathscr{V}}\right\}.$$
 (10)

Distinguished from the geometric approach of estimating QSL using maximum evolution velocity [2], the second term of Eq. (10) applies geometric estimation locally for each parameter. That demonstrates the duration of evolution process is always constrained by the global and local geometric quantum speed bound. Eq. (10) also presents a feasible perspective of estimating the QSL of quantum systems by measuring local geometric quantum speed experimentally.

Application. As an intuitive representation of two-level systems, the Bloch sphere has particular advantages: a) The geodesic between two states on the sphere is equivalent to the Bures angle by considering the conservation of probabilistic flow and ignoring the relative phase. b) The metric tensor is easy to calculate for sphere. To show the validity of Eq. (10), we consider the nonlinear LZ type two-level system, which is related to a model for BECs in the mean-field [47–54]. It is determined by the following Schrödinger equation:

$$i\frac{d}{dt}\begin{pmatrix}\psi_1\\\psi_2\end{pmatrix} = H\begin{pmatrix}\psi_1\\\psi_2\end{pmatrix},\tag{11}$$

where

$$H = \frac{\Gamma(t) - c(|\psi_2|^2 - |\psi_1|^2)}{2}\sigma_z + \frac{v}{2}\sigma_x.$$
 (12)

 ψ_1 and ψ_2 are the probability amplitudes and $\hbar = 1$. The Hamiltonian is characterized by the coupling strength vbetween the two states, the level bias Γ and the nonlinear strength c which describes the interaction between atoms. With the usual Bloch parametrization $\psi_1 = \cos(\chi/2)e^{i\varphi_1}, \psi_2 = \sin(\chi/2)e^{i\varphi_2}$ [55], we introduce $\chi(t)$ to donate the population difference $\cos \chi = |\psi_2|^2 - |\psi_1|^2$ and $\varphi(t)$ to donate the relative phase $\varphi = \varphi_2 - \varphi_1$. In the following calculation, f' defaults to the derivative of fwith respect to time and \dot{f} defaults to the derivative of fwith respect to χ . On the Bloch sphere, the Schrödinger equation reads (see Sec. I of the Supplemental Material [56] for calculation)

$$\begin{cases} \chi' = -v \sin \varphi, \\ \varphi' = \Gamma - \cos \chi \left(c + v \frac{\cos \varphi}{\sin \chi} \right). \end{cases}$$
(13)

In order to derive the QB of the nonlinear LZ type twolevel system, we calculate the analytical expressions of the time-dependent evolution of $\chi(t)$ and $\varphi(t)$. Under the situation $\Gamma = 0$, we find that $\sin \chi = -2v \cos \varphi/c$ is a particular solution of Eq. (13), where 2v/c corresponds exactly to the self-trapping in Josephson junction [57– 59].

For $\Gamma \neq 0$, our main idea is to solve the undetermined function f which is inserted into $\sin \chi = -2v \cos \varphi/c$. Different positions of f are associated with different types of ordinary differential equations. We find that the denominator is the correct position:

$$\sin \chi = -\frac{2v}{c+f}\cos\varphi.$$
 (14)

The derivative with respect to χ on both sides of Eq. (14) leads to

$$\cos \chi = \frac{2v[(c+f)\dot{\varphi}\sin\varphi + \dot{f}\cos\varphi]}{(c+f)^2}.$$
 (15)

With substituting Eq. (14) into Eq. (13), we obtain $\dot{\varphi} = [-2\Gamma + (c - f)\cos\chi]/2v\sin\varphi$. Substituting $\dot{\varphi}$ into (15) and replacing $\cos\varphi$ by Eq. (14) yields: $\dot{f} = -2f\tan\chi - 2\Gamma/\sin\chi$ and further obtains $f = (-2\int\Gamma\sin\chi d\chi + C_0)/\sin^2\chi$. The formulation between $\chi(t)$ and $\varphi(t)$ for arbitrary Γ yields:

$$\sin \chi = -\frac{2v}{c + \frac{-2\int \Gamma \sin \chi d\chi + C_0}{\sin^2 \chi}} \cos \varphi, \qquad (16)$$

where C_0 is the integration constant determined by the initial phase $\varphi(0)$ and initial angle $\chi(0)$. Here we introduce the approximation that the protocol Γ is the arbitrary function of χ . Now $\varphi(t)$ is fully expressed as a function of $\chi(t)$. Finally, by employing Eq. (13), the analytical expressions for the time-dependent evolution of $\chi(t)$ and $\varphi(t)$ can be further calculated by the following



FIG. 1. In the case that χ_i is less than χ_{τ} , the optimal path Eq. (21) is showed on the Bloch sphere. The upper star donates the initial state and the lower star donates the final state. For unconstrained Γ , the speed of φ approaches infinity. The main contribution to evolution time comes from the longitude .

integral (see Sec. II of the Supplemental Material [56] for more examples):

$$d\tau = \frac{d\eta}{v\sqrt{1 - \eta^2 - \frac{(C_0 - 2\Gamma\eta)^2}{4v^2}}}.$$
 (17)

The states of the two-level system are expressed as the points on the Bloch sphere represented by χ and φ . We select χ as the natural parameter and calculate the corresponding metric tensor on the sphere, and then the variational problem can be written as

$$\int_{\chi_0}^{\chi_\tau} \frac{\sqrt{1 + \dot{\varphi}^2 \sin^2 \chi}}{\frac{d}{dt} \mathscr{L}(|\psi_0\rangle, |\psi_\tau\rangle)} d\chi = \int_{\chi_0}^{\chi_\tau} \frac{\dot{\mathscr{L}}}{\mathcal{V}} d\chi = \int_{\chi_0}^{\chi_\tau} T d\chi.$$
(18)

The geometric quantum speed should be consistent with the evolution speed of the point on the Bloch sphere, thus $|d\mathscr{L}(|\psi_0\rangle, |\psi_\tau\rangle)/dt|^2 = (d\varphi/dt)^2 + (d\chi/dt)^2 \sin^2 \chi$. Further simplifying gives $\mathcal{V}^2 = v^2 + (\dot{\xi}^2 \sin^2 \chi - \xi^2)/4$ (see Sec. III of the Supplemental Material [56] for calculation). The solution of the variational problem corresponds to the EL equations $d\partial_{\dot{\varphi}}T/d\chi = \partial_{\varphi}T$, $d\partial_{\dot{\xi}}T/d\chi = \partial_{\xi}T$, where

$$\begin{cases} \frac{\partial T}{\partial \dot{\varphi}} = \frac{\dot{\varphi} \sin^2 \chi}{\dot{\mathscr{L}} \mathcal{V}}, \ \frac{\partial T}{\partial \varphi} = 0, \\ \frac{\partial T}{\partial \dot{\xi}} = \frac{\xi}{2} \frac{\mathscr{L}}{\mathcal{V}^3}, \ \frac{\partial T}{\partial \xi} = -\frac{\dot{\xi} \sin^2 \chi}{2} \frac{\dot{\mathscr{L}}}{\mathcal{V}^3}. \end{cases}$$
(19)

We obtain the QB of this nonlinear LZ type two-level system, which reads

$$\varphi = \pm \frac{\pi}{2}, \ \dot{\varphi} = \xi = \dot{\xi} = 0.$$
 (20)

When the upper limit of the integral is greater than the lower limit, $\varphi = -\pi/2$; otherwise $\varphi = \pi/2$. To sum up,

the minimal time should be $|\chi_0 - \chi_\tau|/v$. Without loss of generality, we consider the case $\chi_0 < \chi_\tau$. The QB we obtained is showed in Fig. 1 for two given states. Combining with the relative phase of the initial and target states, we rewrite the QB by the step function \mathscr{H} of the following form (see Sec. IV of the Supplemental Material [56] for details):

$$\varphi = \begin{cases} \varphi_0 - \left(\frac{\pi}{2} + \varphi_0\right) \mathscr{H} \left(\chi - \chi_0\right), & \text{if } \chi \le (\chi_0 + \chi_\tau)/2, \\ \varphi_\tau - \left(\frac{\pi}{2} + \varphi_\tau\right) \mathscr{H} \left(\chi_\tau - \chi\right), & \text{if } \chi > (\chi_0 + \chi_\tau)/2. \end{cases}$$
(21)

Moreover, the optimal protocol Γ is calculated in terms of Eq. (16):

$$\Gamma = c \cos \chi - \delta(\chi - \chi_0) + \delta(\chi - \chi_\tau), \qquad (22)$$

where $\delta(\chi)$ is the Dirac function. Eq. (22) is consistent with the results given by the experiment [8] and optimal control [39, 40]. For the variational method, it is necessary to further verify the protocol Γ is globally optimal. Eq. (21) shows that the contribution of φ to the evolution time tends to be 0 and the duration of evolution process only depends on χ . In fact, it is evident that the following inequality holds with Eq. (17) only if $\Gamma = c \cos \chi$, $C_0 = 0$:

$$\tau[\bullet] \ge \left| \int_{\eta_0}^{\eta_\tau} \frac{1}{v} \frac{d\eta}{\sqrt{1-\eta^2}} \right| = \frac{|\chi_0 - \chi_\tau|}{v}, \qquad (23)$$

which presents the QSL of the nonlinear LZ type twolevel system. Note that Eq. (23) indicates that the QB has the same formulation for repulsive and attractive interactions.

Discussion. By inserting parameters of this nonlinear model, the QSL can be expressed explicitly from Eq. (10):

$$\tau_{QSL} = \max\left\{\frac{|\chi_0 - \chi_\tau|}{|v_\chi|_{max}}, \frac{|\varphi_0 - \varphi_\tau|}{|v_\varphi|_{max}}, \frac{\mathscr{L}(|\psi_0\rangle, |\psi_\tau\rangle)}{\mathcal{V}_{max}}\right\}$$
$$= \max\left\{\frac{|\chi_0 - \chi_\tau|}{v}, 0, 0\right\}, \tag{24}$$

which shows that the bound we derived is tight and illustrates that the coupling strength v acts the maximum local geometric quantum speed. Optimal control gives the QSL which is the first term of Eq. (10) [39, 40]. The conventional geometric approach which focuses on the maximum geometric speed is corresponding to the global estimation. Coincidently, these two methods reflect two different terms of Eq. (10). The QB is essentially the transition of these two methods. Thus we reach that our framework provides the perspective of viewing the previous methods via the geometric quantum speed estimation.

Now we introduce the generalization of Eq. (10) to the open quantum systems. For the mixed states, the metric

of Bures angle can also be rewritten into the form of Eq. (3) [43] and the corresponding time functional is

$$\tau[\bullet] = \int \frac{\sqrt{\sum_{\mu,\nu=1}^{2n} g_{\mu\nu} \frac{d\lambda_{\mu}}{d\zeta} \frac{d\lambda_{\nu}}{d\zeta}}}{\frac{d}{dt} \mathscr{L}(\rho_0, \rho_{\tau})} d\zeta, \qquad (25)$$

where $\mathscr{L}(\rho_0, \rho_\tau) = \arccos(tr\sqrt{\sqrt{\rho_\tau}\rho_\sqrt{\rho_\tau}})$ [60, 61], ρ_0 donates the initial state and ρ_τ donates the final state. Therefore, following the steps above, there also exists estimation Eq. (10), which shows that it is possible to give effective bound on actual evolution time in practical applications only by measuring quantum geometric speed.

Conclusions. We have derived the QSL for arbitrary quantum systems with unitray dynamics via the framework of the QB. This bound is expressed as the global and local geometric quantum speed estimation. We have applied our method for the nonlinear LZ type two-level system to investgate the evolution speed of the complex system. We have found the analytical expression of evolution population with the usual Bloch parametrization to derive the corresponding optimal procotol Γ and QB of this nonlinear system. We further demonstrate that the QSL is tight, which is consistent with the result given by the optimal control and the experiment in the linear limit. We also generalize our formulation to open quantum systems. We hope that the present geometric method to the QSL opens up novel systematic investigations of this fundamental principle.

The work is supported by the National Natural Science Foundation of China (Grant No. 12075193).

* doufq@nwnu.edu.cn

- [1] S. Lloyd, Nature **406**, 1047 (2000).
- [2] S. Deffner and S. Campbell, J. Phys. A: Math. Theor. 50, 453001 (2017).
- [3] Y.-J. Zhang, W. Han, Y.-J. Xia, J.-P. Cao, and H. Fan, Phys. Rev. A 91, 032112 (2015).
- [4] M. R. Frey, Quantum Inf. Process. 15, 3919 (2016).
- [5] A. del Campo, I. L. Egusquiza, M. B. Plenio, and S. F. Huelga, Phys. Rev. Lett. **110**, 050403 (2013).
- [6] M. M. Taddei, B. M. Escher, L. Davidovich, and R. L. de Matos Filho, Phys. Rev. Lett. **110**, 050402 (2013).
- [7] E. O'Connor, G. Guarnieri, and S. Campbell, Phys. Rev. A 103, 022210 (2021).
- [8] M. G. Bason, M. Viteau, N. Malossi, P. Huillery, E. Arimondo, D. Ciampini, R. Fazio, V. Giovannetti, R. Mannella, and O. Morsch, Nat. Phys. 8, 147 (2012).
- [9] M. R. Lam, N. Peter, T. Groh, W. Alt, C. Robens, D. Meschede, A. Negretti, S. Montangero, T. Calarco, and A. Alberti, Phys. Rev. X 11, 011035 (2021).
- [10] S. van Frank, M. Bonneau, J. Schmiedmayer, S. Hild, C. Gross, M. Cheneau, I. Bloch, T. Pichler, A. Negretti, T. Calarco, *et al.*, Sci. Rep. **6**, 1 (2016).

- [11] Y. Dong, C. Feng, Y. Zheng, X.-D. Chen, G.-C. Guo, and F.-W. Sun, Phys. Rev. Research 3, 043177 (2021).
- [12] N. Il'in and O. Lychkovskiy, Phys. Rev. A 103, 062204 (2021).
- [13] P. M. Poggi, S. Campbell, and S. Deffner, PRX Quantum 2, 040349 (2021).
- [14] T.-N. Xu, J. Li, T. Busch, X. Chen, and T. Fogarty, Phys. Rev. Research 2, 023125 (2020).
- [15] J. Wang, Y. Wu, M. Mo, and H. Zhang, Sci. Rep. 6, 1 (2016).
- [16] S. Sun and Y. Zheng, Phys. Rev. Lett. **123**, 180403 (2019).
- [17] S.-X. Wu and C.-S. Yu, Sci. Rep. 10, 1 (2020).
- [18] K. Suzuki and K. Takahashi, Phys. Rev. Research 2, 032016(R) (2020).
- [19] X. Meng, C. Wu, and H. Guo, Sci. Rep. 5, 1 (2015).
- [20] X. Wang, M. Allegra, K. Jacobs, S. Lloyd, C. Lupo, and M. Mohseni, Phys. Rev. Lett. **114**, 170501 (2015).
- [21] F. Campaioli, W. Sloan, K. Modi, and F. A. Pollock, Phys. Rev. A 100, 062328 (2019).
- [22] A. Carlini, A. Hosoya, T. Koike, and Y. Okudaira, Phys. Rev. Lett. 96, 060503 (2006).
- [23] H. Wakamura and T. Koike, New J. Phys. 22, 073010 (2020).
- [24] B. Russell and S. Stepney, J. Phys. A: Math. Theor. 48, 115303 (2015).
- [25] T. Hatomura, Phys. Rev. A **100**, 043619 (2019).
- [26] A. R. Kuzmak and V. M. Tkachuk, Phys. Lett. A 379, 1233 (2015).
- [27] B. Amghar and M. Daoud, Phys. Lett. A 384, 126682 (2020).
- [28] A. R. Kuzmak and V. M. Tkachuk, J. Phys. A: Math. Theor. 46, 155305 (2013).
- [29] A. Mostafazadeh, Phys. Rev. Lett. 99, 130502 (2007).
- [30] U. Günther and B. F. Samsonov, Phys. Rev. Lett. 101, 230404 (2008).
- [31] A. Beygi and S. P. Klevansky, Phys. Rev. A 98, 022105 (2018).
- [32] T. Albash and D. A. Lidar, Rev. Mod. Phys. 90, 015002 (2018).
- [33] A. T. Rezakhani, W.-J. Kuo, A. Hamma, D. A. Lidar, and P. Zanardi, Phys. Rev. Lett. **103**, 080502 (2009).
- [34] A. C. Santos, C. J. Villas-Boas, and R. Bachelard, Phys. Rev. A 103, 012206 (2021).
- [35] K. Takahashi, Phys. Rev. A 91, 042115 (2015).
- [36] B. K. Zhao and J. B. Lu, Europhys. Lett. 100, 50002 (2012).
- [37] A. K. Pati, B. Pradhan, and P. Agrawal, Quantum Inf. Process. 11, 841 (2012).
- [38] A. Carlini and T. Koike, J. Phys. A: Math. Theor. 50, 105304 (2017).
- [39] G. C. Hegerfeldt, Phys. Rev. Lett. 111, 260501 (2013).
- [40] G. C. Hegerfeldt, Phys. Rev. A **90**, 032110 (2014).
- [41] W. K. Wootters, Phys. Rev. D 23, 357 (1981).
- [42] S. L. Braunstein and C. M. Caves, Phys. Rev. Lett. 72, 3439 (1994).
- [43] D. P. Pires, M. Cianciaruso, L. C. Céleri, G. Adesso, and D. O. Soares-Pinto, Phys. Rev. X 6, 021031 (2016).
- [44] I. Bengtsson and K. Życzkowski, Geometry of Quantum States: An Introduction to Quantum Entanglement, 2nd ed. (Cambridge University Press, 2017).
- [45] M. S. Byrd and N. Khaneja, Phys. Rev. A 68, 062322 (2003).

- [46] F. Campaioli, F. A. Pollock, F. C. Binder, and K. Modi, Phys. Rev. Lett. **120**, 060409 (2018).
- [47] M. Jona-Lasinio, O. Morsch, M. Cristiani, N. Malossi, J. H. Müller, E. Courtade, M. Anderlini, and E. Arimondo, Phys. Rev. Lett. **91**, 230406 (2003).
- [48] A. M. Ishkhanyan, Phys. Rev. A 81, 055601 (2010).
- [49] F.-Q. Dou, L.-B. Fu, and J. Liu, Phys. Rev. A 89, 012123 (2014).
- [50] X. Chen, Y. Ban, and G. C. Hegerfeldt, Phys. Rev. A 94, 023624 (2016).
- [51] F.-Q. Dou, J. Liu, and L.-B. Fu, Phys. Rev. A 98, 022102 (2018).
- [52] F.-Q. Dou, H. Cao, J. Liu, and L.-B. Fu, Phys. Rev. A 93, 043419 (2016).
- [53] J. Liu, L. Fu, B.-Y. Ou, S.-G. Chen, D.-I. Choi, B. Wu, and Q. Niu, Phys. Rev. A 66, 023404 (2002).
- [54] B. Wu and Q. Niu, Phys. Rev. A **61**, 023402 (2000).
- [55] M. V. Berry, J. Phys. A: Math. Theor. 42, 365303 (2009).
- [56] See supplemental material for more details.
- [57] G. J. Milburn, J. Corney, E. M. Wright, and D. F. Walls, Phys. Rev. A 55, 4318 (1997).
- [58] G.-F. Wang, L.-B. Fu, and J. Liu, Phys. Rev. A 73, 013619 (2006).
- [59] M. Albiez, R. Gati, J. Fölling, S. Hunsmann, M. Cristiani, and M. K. Oberthaler, Phys. Rev. Lett. 95, 010402 (2005).
- [60] A. Uhlmann, Rep. Math. Phys. 9, 273 (1976).
- [61] S. Kakutani, Ann. Math. **49**, 214 (1948).

Supplemental Material for "How fast can quantum brachistochrone evolve?"

Fu-Quan Dou and Min-Peng Han

College of Physics and Electronic Engineering, Northwest Normal University, Lanzhou, 730070, China

I. DERIVATION OF DYNAMICS EQUATIONS BY THE BLOCH PARAMETRIZATION

To begin, we calculate the derivative of χ with respect to time. The general form of the Hamilton which we consider reads

$$H = p\sigma_z + q\sigma_x. \tag{S1}$$

For $|\psi_1|^2$ we have $d|\psi_1|^2/dt = (\psi_1^*)'\psi_1 + \psi_1^*\psi_1'$, where ψ^* donates the complex conjugate of ψ . Submitting it into the Schrödinger equation and we obtain

$$\frac{d|\psi_1|^2}{dt} = i\psi_1 \left(p\psi_1^* + q\psi_2^*\right) - i\psi_1^* \left(p\psi_1 + q\psi_2\right)$$
$$= iq \left(\psi_1\psi_2^* - \psi_1^*\psi_2\right). \tag{S2}$$

Using the substitution $\psi_1 = \cos(\chi/2)e^{i\varphi_1}, \psi_2 = \sin(\chi/2)e^{i\varphi_2}, \varphi = \varphi_2 - \varphi_1$ yields

$$\frac{d|\psi_1|^2}{dt} = q\sin\chi\sin\varphi,\tag{S3}$$

which can be further written as

$$\frac{d|\psi_1|^2}{dt} = \frac{d\left[\cos^2\left(\frac{\chi}{2}\right)\right]}{dt} = \frac{-\chi'\sin\chi}{2}.$$
 (S4)

Combining with Eq. (S3) and Eq. (S4) and we get the derivative of χ with respect to time

$$\chi' = -2q\sin\varphi. \tag{S5}$$

For ψ_1 , the Bloch parametrization and dynamics equation lead to

$$\begin{cases} \psi_1' = -\frac{1}{2}\sin\left(\frac{\chi}{2}\right)e^{i\varphi_1} + i\varphi_1'\cos\left(\frac{\chi}{2}\right)e^{i\varphi_1}, \\ \psi_1' = -i(p\psi_1 + q\psi_2), \end{cases}$$
(S6)

with

$$\varphi_1' = -p - q \tan\left(\frac{\chi}{2}\right) e^{i\varphi} - \frac{i\chi'}{2} \tan\left(\frac{\chi}{2}\right).$$
 (S7)

In the same way for ψ_2 we have

$$\begin{cases} \psi_2' = -\frac{1}{2}\cos\left(\frac{\chi}{2}\right)e^{i\varphi_2} + i\varphi_2'\sin\left(\frac{\chi}{2}\right)e^{i\varphi_2}, \\ \psi_2' = -i(q\psi_1 - p\psi_2), \end{cases}$$
(S8)

and φ_2 satisfies

$$\varphi_2' = p - q \cot\left(\frac{\chi}{2}\right) e^{-i\varphi} + \frac{i\chi'}{2} \cot\left(\frac{\chi}{2}\right).$$
 (S9)

 $\varphi' = \varphi'_2 - \varphi'_1$ is further calculated. The difference between the first term of Eq. (S7) and Eq. (S9) equals to 2p. The difference between the second terms lead to

$$- q \cot\left(\frac{\chi}{2}\right) e^{-i\varphi} + q \tan\left(\frac{\chi}{2}\right) e^{i\varphi}$$
$$= q[-\cot\left(\frac{\chi}{2}\right)\cos\varphi + i\cot\left(\frac{\chi}{2}\right)\sin\varphi$$
$$+ \tan\left(\frac{\chi}{2}\right)\cos\varphi + i\tan\left(\frac{\chi}{2}\right)\sin\varphi].$$
(S10)

Note that

$$\begin{cases}
\tan\left(\frac{\chi}{2}\right) + \cot\left(\frac{\chi}{2}\right) = 2\csc\chi, \\
\tan\left(\frac{\chi}{2}\right) - \cot\left(\frac{\chi}{2}\right) = -2\cot\chi,
\end{cases}$$
(S11)

then Eq. (S10) becomes

$$- q \cot\left(\frac{\chi}{2}\right) e^{-i\varphi} + q \tan\left(\frac{\chi}{2}\right) e^{i\varphi}$$
$$= -2q \cos\varphi \cot\chi + 2iq \sin\varphi \csc\chi.$$
(S12)

The difference between the third terms yield

$$\frac{i\chi'}{2}\cot\left(\frac{\chi}{2}\right) + \frac{i\chi'}{2}\tan\left(\frac{\chi}{2}\right) = i\chi'\csc\chi$$
$$= -2iq\sin\varphi\csc\chi.$$
(S13)

Thus $\varphi'_2 - \varphi'_1$, namely the derivative of relative phase with respect to t equals to

$$\varphi' = \varphi'_2 - \varphi'_1 = 2p - 2q\cos\varphi\cot\chi.$$
 (S14)

Specifically, for the nonlinear LZ type two-level system, $p = (\Gamma - c \cos \chi)/2$, q = v/2. Inserting p and q, the Schrödinger equation representing by the Bloch parametrization reads

$$\begin{cases} \chi' = -v \sin \varphi, \\ \varphi' = \Gamma - \cos \chi \left(c + \frac{v \cos \varphi}{\sin \chi} \right). \end{cases}$$
(S15)

II. THE APPLICATION OF THE ANALYTICAL EXPRESSION OF THE POPULATION

In general, the evolution time of the nonlinear LZ type two-level system can be expressed as

$$d\tau = \frac{d\eta}{v\sqrt{1 - \eta^2 - \frac{\mathscr{F}^2}{4v^2}}},\tag{S16}$$

where $\mathscr{F} = c(1 - \eta^2) - 2 \int \Gamma d\eta + C_0$ and $\eta = \cos \chi$. To illustrate the application of this expression, here we take the nonlinear Josephson oscillation as an example, which describes the dynamics for BECs in a double-well system with nonlinear interaction. For this model, the level bias Γ is constant in Eq. (S16).

A. The linear case with c = 0

Under the linear case, the nonlinear Josephson oscillation model becomes the known Rabi oscillation and the Eq. (S16) reads

$$d\tau = \frac{d\eta}{v\sqrt{1 - \eta^2 - \frac{(C_0 - 2\Gamma\eta)^2}{4v^2}}}.$$
 (S17)

Considering the initial condition $\lim_{\tau \to 0} \varphi = \pi/2$, $\lim_{\tau \to 0} \chi = 0$, we obtain $C_0 = 2\Gamma$ and the evolution time can be written as

$$d\tau = \frac{d\eta}{v\sqrt{1 - \eta^2 - \frac{\Gamma^2}{v^2}(1 - \eta)^2}}.$$
 (S18)

For the Rabi oscillation, the analytical expression of the population is well known

$$\begin{aligned} |\psi_1(\tau)|^2 &= 1 - \frac{v^2}{v^2 + \Gamma^2} \sin^2\left(\frac{\sqrt{v^2 + \Gamma^2}}{2}\tau\right), \\ |\psi_2(\tau)|^2 &= \frac{v^2}{v^2 + \Gamma^2} \sin^2\left(\frac{\sqrt{v^2 + \Gamma^2}}{2}\tau\right), \end{aligned}$$
(S19)

which yields

$$\tau = \frac{2}{\sqrt{v^2 + \Gamma^2}} \arcsin\left[\sqrt{\frac{v^2 + \Gamma^2}{2v^2}(1 - \cos\chi)}\right].$$
 (S20)

Differentiating the evolution time with respect to χ gives Eq. (S17). Eq. (S20) indicates that under the linear case, Eq. (S16) is equivalent to the Rabi oscillation that we are familiar with.

B. Explanation of the self-trapping for the nonlinear Josephson oscillation

When $\Gamma = 0$ and $c \neq 0$, we have the relation

S

$$\sin \chi = -\frac{2v}{c}\cos\varphi. \tag{S21}$$

With the nonlinear interaction, the tunneling between two wells is influenced by the nonlinear strength like in Fig. S1. We use the fourth to fifth-order Runge-Kutta method to simulate the quantum state evolution on the Bloch sphere and illustrate the physical significance of Eq. (S21).

For $|\sin \chi| \leq 1$, when $2v/c \geq 1$, i.e., $|c/v| \leq 2$, the difference of the population will not be limited. This is reflected in the fact that the state can always evolve from the North pole to the South pole on the sphere. However, when |2v/c| < 1, i.e., |c/v| > 2, the difference of the population will be limited by the nonlinear strength and the maximum of the sin χ becomes |2v/c|. χ never approaches $\pi/2$, which means that the state never crosses the equator, namely self-trapping.



FIG. S1. Evolution process of the Josephson oscillation on the Bloch sphere. (a) c/v = 4, (b) c/v = 2, (c) c/v = 1.5, (d) c/v = 1.

III. SIMPLIFICATION OF \mathcal{V}^2

 \mathcal{V}^2 can be rewritten as the form

$$\mathcal{V}^2 = v^2 (1 - \cos^2 \varphi) + \mathscr{G}^2 \sin^2 \chi, \qquad (S22)$$

where $\mathscr{G} = \Gamma - \cos \chi (c + v \cos \varphi / \sin \chi)$. Combining with $\sin \chi = -2v \cos \varphi / (c + f)$, we obtain

$$\mathcal{V}^2 = v^2 - \frac{(c+f)^2 \sin^2 \chi}{4}$$
(S23)
+ $\left(\Gamma - \frac{c \cos \chi}{2} + \frac{f \cos \chi}{2}\right)^2 \sin^2 \chi,$
 $\mathscr{G} = \Gamma - \frac{c \cos \chi}{2} + \frac{f \cos \chi}{2},$

where

$$f = \frac{-2\int\Gamma\sin\chi d\chi + C_0}{\sin^2\chi}.$$
 (S24)

Multiplying both sides of Eq. (S24) by $\sin^2 \chi$ and taking the derivative of both sides with respect to χ give

$$\Gamma = -\frac{\dot{f}\sin\chi}{2} - f\cos\chi.$$
 (S25)

Submitting Eq. (S25) into Eq. (S24), we obtain

$$\mathcal{V}^{2} = v^{2} - \frac{(c+f)^{2} \sin^{2} \chi}{4} + \frac{\left(f \cos \chi + \dot{f} \sin \chi + c \cos \chi\right)^{2}}{4} \sin^{2} \chi. \quad (S26)$$

For the sake of simplification, we introduce $\xi = f \sin \chi + c \sin \chi$, which leads to the simplified form

$$\mathcal{V}^2 = v^2 + \frac{\dot{\xi}^2 \sin^2 \chi - \xi^2}{4}.$$
 (S27)

IV. TRANSFORMATION OF TRAJECTORY EQUATION

which is equivalent to

With the step function

$$\mathscr{H}(x) = \begin{cases} 0, & \text{if } x \le 0, \\ 1, & \text{if } x > 0, \end{cases}$$
(S28)

the relative phase φ can be written as:

$$\varphi = \begin{cases} \varphi_0 - \left(\frac{\pi}{2} + \varphi_0\right) \mathscr{H} \left(\chi - \chi_0\right), & \text{if } \chi_0 \le \chi \le \frac{\chi_0 + \chi_\tau}{2}, \\ \varphi_\tau - \left(\frac{\pi}{2} + \varphi_\tau\right) \mathscr{H} \left(\chi_\tau - \chi\right), & \text{if } \frac{\chi_0 + \chi_\tau}{2} < \chi \le \chi_\tau, \end{cases}$$
(S29)

$$\varphi = \begin{cases} \varphi_0, & \text{if } \chi = \chi_0, \\ -\frac{\pi}{2}, & \text{if } \chi_0 < \chi < \chi_\tau, \\ \varphi_\tau, & \text{if } \chi = \chi_\tau. \end{cases}$$
(S30)

For Eq. (S29), the position of the subsection should be within $[\chi_0, \chi_\tau]$. Here, $(\chi_0 + \chi_\tau)/2$ is selected.