

Quantum-inspired permanent identities

Ulysse Chabaud¹, Abhinav Deshpande¹, and Saeed Mehraban²

¹Institute for Quantum Information and Matter, California Institute of Technology, Pasadena, CA 91125, USA

²Computer Science, Tufts University, Boston, MA 02155, USA

The permanent is pivotal to both complexity theory and combinatorics. In quantum computing, the permanent appears in the expression of output amplitudes of linear optical computations, such as in the Boson Sampling model. Taking advantage of this connection, we give quantum-inspired proofs of many existing as well as new remarkable permanent identities. Beyond their purely combinatorial applications, our results demonstrate the classical hardness of exact and approximate sampling of linear optical quantum computations with input cat states.

Contents

1	Introduction	2
2	Background	3
2.1	Notations and preliminary material	3
2.2	Boson Sampling	5
2.3	The MacMahon master theorem	5
3	Main results	6
4	Discussion	8
5	Quantum-inspired proofs of permanent identities	8
5.1	Glynn’s formula	8
5.2	The Glynn–Kan formula	10
5.3	The Cauchy–Binet theorem	11
5.4	Generating functions	12
6	New quantum-inspired permanent identities	14
6.1	Generalizations of the MacMahon master theorem	15
6.2	New generating functions	20
7	Boson Sampling with input cat states	23

uchabaud@caltech.edu

saeed.mehraban@tufts.edu

1 Introduction

The *permanent* of an $m \times m$ matrix $A = (a_{ij})_{1 \leq i, j \leq m}$ is a combinatorial function defined as [1, 2]:

$$\text{Per}(A) = \sum_{\sigma \in \mathcal{S}_m} \prod_{k=1}^m a_{k\sigma(k)}, \quad (1)$$

where \mathcal{S}_m is the symmetric group over m symbols. While the closely related determinant can be computed efficiently using many methods such as Gaussian elimination, Valiant famously proved the $\#P$ -hardness of computing the permanent [3].

Interestingly, the permanent appears in the expression of the output amplitudes of linear optical quantum computations with noninteracting bosons [4, 5], as in the Boson Sampling model of quantum computation [6]. This connection has led to several linear optical proofs of existing and new classical complexity results: computation of the permanent is $\#P$ -hard [7], multiplicative estimation of the permanent of positive semidefinite matrices is in BPP^{NP} [8], multiplicative estimation of the permanent of orthogonal matrices is $\#P$ -hard [9], and computation of a class of multidimensional integrals is $\#P$ -hard [10]. It also led to the introduction of a quantum-inspired classical algorithm for additive estimation of the permanent of positive semidefinite matrices [11].

Beyond its importance for complexity theory, the permanent has numerous applications for solving combinatorial problems [1, 2] and identities for the permanent have been instrumental in these applications. For example, the MacMahon master theorem [12], which relates the permanent to a coefficient of the Taylor series of a determinant, is an invaluable tool for proving combinatorial identities [13–15]. Similarly, Ryser’s formula [16] and Glynn’s formula [17–19] are routinely used to compute the permanent more efficiently than the naive brute-force approach.

While linear optics has been used as a tool to explore the classical complexity of the permanent, previous work suggests that it can also be a useful way to obtain simple proofs of theorems about the permanent: for instance, [5] shows that the permanent of a unitary matrix U lies in the (closed) complex unit disk by expressing $|\text{Per}(U)|^2$ as an output probability of a linear optical computation, while [6] derives simple permanent identities using various representations of the same linear optical computation. This begs the following questions:

1. Can we use linear optics to prove existing remarkable permanent identities, such as the MacMahon master theorem?
2. Can we use linear optics to derive *new* remarkable permanent identities?

In this work, we show that the answer to both questions is *yes*.

We give quantum-inspired proofs of several important permanent identities in section 5. In particular, we show that the MacMahon master theorem can be understood as two different ways of computing an inner product between two Gaussian quantum states.

We also derive new quantum-inspired identities for the permanent. Our main results are summarized in section 3 and proven in section 6. These include generalizations of the MacMahon master theorem (Theorems 1 and 2), new generating functions for the permanent (Theorem 3) and a formula for the sum of two permanents (Theorem 4).

As a bonus, our findings also have consequences for the classical complexity of exact and approximate sampling of linear optical computations with input cat states (Theorem 5), which we discuss in section 7. We rigorously prove that the corresponding quantum probability distributions are as hard to sample as the original Boson Sampling distribution [6],

for all cat state amplitudes in the exact case and for small enough amplitudes in the approximate case. Until now, a formal proof was available only in the exact case [20], based on post-selection.

2 Background

2.1 Notations and preliminary material

$\mathbf{0} = (0, \dots, 0)$ $\mathbf{1} = (1, \dots, 1)$ $ \mathbf{p}\rangle = p_1 + \dots + p_m$ $\mathbf{p}! = p_1! \dots p_m!$ $ \mathbf{p}\rangle = p_1\rangle \otimes \dots \otimes p_m\rangle$ $\mathbf{p} + \mathbf{q} = (p_1 + q_1, \dots, p_m + q_m)$ $\mathbf{p} \oplus \mathbf{q} = (p_1, \dots, p_m, q_1, \dots, q_m)$	$\mathbf{p} \leq \mathbf{q} \Leftrightarrow \forall k \in \{1, \dots, m\}, p_k \leq q_k$ $-\mathbf{z} = (-z_1, \dots, -z_m)$ $\mathbf{z}^* = (z_1^*, \dots, z_m^*)$ $\ \mathbf{z}\ ^2 = z_1 ^2 + \dots + z_m ^2$ $\mathbf{z}^{\mathbf{p}} = z_1^{p_1} \dots z_m^{p_m}$ $\partial_{\mathbf{z}}^{\mathbf{p}} = \partial_{z_1}^{p_1} \dots \partial_{z_m}^{p_m}$ $d^m \mathbf{z} d^m \mathbf{z}^* = d \operatorname{Re}\{z_1\} d \operatorname{Im}\{z_1\} \dots d \operatorname{Re}\{z_m\} d \operatorname{Im}\{z_m\}$
---	---

Table 1: Multi-index notations used in this paper, for $m \in \mathbb{N}^*$, $\mathbf{p} = (p_1, \dots, p_m) \in \mathbb{N}^m$, $\mathbf{q} = (q_1, \dots, q_m) \in \mathbb{N}^m$ and $\mathbf{z} = (z_1, \dots, z_m) \in \mathbb{C}^m$.

We use bold math for multi-index expressions (see Table 1 above). We denote by $\mathbb{T} = \{z \in \mathbb{C}, |z| = 1\}$ the complex unit circle. For all $m \in \mathbb{N}^*$, all $\mathbf{z} = (z_1, \dots, z_m) \in \mathbb{C}^m$ and all $\mathbf{p} = (p_1, \dots, p_m) \in \mathbb{N}^m$, we use the notation $[\mathbf{z}^{\mathbf{p}}]$ to denote the coefficient of $\mathbf{z}^{\mathbf{p}} = z_1^{p_1} \dots z_m^{p_m}$ in an analytic expression. For all $m \in \mathbb{N}^*$, all $\mathbf{p} = (p_1, \dots, p_m) \in \mathbb{N}^m$ and all $\mathbf{q} = (q_1, \dots, q_m) \in \mathbb{N}^m$, we denote by $A_{\mathbf{p}, \mathbf{q}}$ the matrix obtained from an $m \times m$ matrix A by repeating its i^{th} row p_i times (deleting the row if $p_i = 0$) and repeating its j^{th} column q_j times (deleting the column if $q_j = 0$) for all $i, j \in \{1, \dots, m\}$. By convention, we set the permanent of non-square matrices to 0 and $\operatorname{Per}(A_{\mathbf{0}, \mathbf{0}}) = 1$.

Let $m \in \mathbb{N}^*$ denote the number of modes. We denote (unnormalized) quantum states using the Dirac ket notation. These are vectors in an infinite-dimensional Hilbert space spanned by the orthonormal Fock basis

$$\{|\mathbf{p}\rangle = |p_1\rangle \otimes \dots \otimes |p_m\rangle\}_{\mathbf{p}=(p_1, \dots, p_m) \in \mathbb{N}^m}. \quad (2)$$

Hereafter, we label Fock states using p , \mathbf{p} , \mathbf{q} , and \mathbf{k} , coherent states using α , β , $\boldsymbol{\alpha}$, and $\boldsymbol{\beta}$, and two-mode squeezed states using λ , μ , $\boldsymbol{\lambda}$, and $\boldsymbol{\mu}$. In particular, coherent states are defined as

$$|\alpha\rangle = e^{-\frac{1}{2}|\alpha|^2} \sum_{p \geq 0} \frac{\alpha^p}{\sqrt{p!}} |p\rangle, \quad (3)$$

for all $\alpha \in \mathbb{C}$, and (unnormalized) two-mode squeezed states as

$$|\lambda\rangle = \sum_{p \geq 0} \lambda^p |pp\rangle, \quad (4)$$

for all $\lambda \in \mathbb{C}$ with $|\lambda| < 1$. Moreover, cat states are defined as

$$|\operatorname{cat}_{\alpha}\rangle = \frac{e^{-\frac{1}{2}|\alpha|^2}}{2\sqrt{\sinh(|\alpha|^2)}} (|\alpha\rangle - |-\alpha\rangle), \quad (5)$$

for all $\alpha \in \mathbb{C}$.

We make use of the following inner products involving these states: for all $p \in \mathbb{N}$, $\alpha \in \mathbb{C}$, and $\lambda \in \mathbb{C}$ with $|\lambda| < 1$,

$$\langle p|\alpha\rangle = e^{-\frac{1}{2}|\alpha|^2} \frac{\alpha^p}{\sqrt{p!}} \quad (6)$$

$$\langle \alpha|\beta\rangle = e^{-\frac{1}{2}|\alpha|^2 - \frac{1}{2}|\beta|^2 + \alpha^*\beta}, \quad (7)$$

$$\langle pp|\lambda\rangle = \lambda^p, \quad (8)$$

which can be readily checked in the Fock basis. Moreover, for all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ [5, 6],

$$\langle \mathbf{p}|\hat{U}|\mathbf{q}\rangle = \frac{\text{Per}(U_{\mathbf{p},\mathbf{q}})}{\sqrt{\mathbf{p}!\mathbf{q}!}}, \quad (9)$$

where \hat{U} is a passive linear operation over m modes whose action on the creation operators of the modes is described by the unitary matrix $U = (u_{ij})_{1 \leq i, j \leq m}$ as

$$\hat{U}\hat{a}_j^\dagger\hat{U}^\dagger = \sum_{i=1}^m u_{ij}\hat{a}_i^\dagger, \quad (10)$$

for all $j \in \{1, \dots, m\}$. As their name indicates, passive linear operations do not change the total number of photons [21]: for all $n \in \mathbb{N}$,

$$\hat{U}\Pi_n = \Pi_n\hat{U}, \quad (11)$$

where $\Pi_n := \sum_{|\mathbf{p}|=n} |\mathbf{p}\rangle\langle \mathbf{p}|$ is the m -mode projector onto states with total photon number equal to n . Passive linear operations map tensor products of coherent states to tensor products of coherent states [21]: for all $\boldsymbol{\alpha} \in \mathbb{C}^m$,

$$\hat{U}|\boldsymbol{\alpha}\rangle = |U\boldsymbol{\alpha}\rangle. \quad (12)$$

We will also make use of the following Gaussian inner product [22]: for all $\boldsymbol{\lambda}, \boldsymbol{\mu} \in \mathbb{C}^m$ with $|\lambda_k| < 1$ and $|\mu_k| < 1$ for all $k \in \{1, \dots, m\}$, and for all passive linear operation \hat{U} over $2m$ modes,

$$\begin{aligned} \langle \boldsymbol{\lambda}^*|\hat{U}|\boldsymbol{\mu}\rangle &= \int_{\mathbf{z} \in \mathbb{C}^{2m}} e^{\sum_{k=1}^m \lambda_k z_k^* z_{m+k}^*} e^{\sum_{k=1}^m \mu_k (U^T \mathbf{z})_k (U^T \mathbf{z})_{m+k}} e^{-\sum_{j=1}^{2m} z_j^* z_j} \frac{d^{2m} \mathbf{z} d^{2m} \mathbf{z}^*}{\pi^{2m}} \\ &= \int_{\mathbf{z} \in \mathbb{C}^{2m}} \exp \left[-\frac{1}{2} \begin{pmatrix} \mathbf{z} \\ \mathbf{z}^* \end{pmatrix}^T V_U(\boldsymbol{\lambda}, \boldsymbol{\mu}) \begin{pmatrix} \mathbf{z} \\ \mathbf{z}^* \end{pmatrix} \right] \frac{d^{2m} \mathbf{z} d^{2m} \mathbf{z}^*}{\pi^{2m}} \\ &= \frac{1}{\sqrt{\text{Det}(V_U(\boldsymbol{\lambda}, \boldsymbol{\mu}))}}, \end{aligned} \quad (13)$$

where we have introduced the $(4m) \times (4m)$ symmetric matrix

$$V_U(\boldsymbol{\lambda}, \boldsymbol{\mu}) := \begin{pmatrix} -UV_{\boldsymbol{\mu}}U^T & I_{2m} \\ I_{2m} & -V_{\boldsymbol{\lambda}} \end{pmatrix}, \quad (14)$$

where for all $\mathbf{w} \in \mathbb{C}^n$,

$$V_{\mathbf{w}} := \begin{pmatrix} 0_m & \text{Diag}(\mathbf{w}) \\ \text{Diag}(\mathbf{w}) & 0_m \end{pmatrix}. \quad (15)$$

Note that we associate mode 1 with mode $m + 1$, mode 2 with mode $m + 2$, and so on, i.e. $|\boldsymbol{\lambda}\rangle = \bigotimes_{k=1}^m |\lambda_k\rangle$, where $|\lambda_k\rangle$ is an unnormalized two-mode squeezed state over modes k and $m + k$.

Finally, we note that any $m \times m$ matrix A with $\|A\| \leq 1$ may be embedded as the top-left submatrix of a $(2m) \times (2m)$ unitary matrix U [6, Lemma 29]. We will use this fact multiple times throughout the paper to extend identities proven for unitary matrices to the case of general matrices.

2.2 Boson Sampling

Boson Sampling is a sub-universal model of quantum computation introduced by Aaronson and Arkhipov (AA) in [6], which takes as input a Fock state $|\mathbf{1} \oplus \mathbf{0}\rangle = |1\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}$, evolves it according to a passive linear operation \hat{U} over m modes with unitary matrix U , and measures the photon number of each output mode.

With Eq. (9), the probability of detecting $\mathbf{p} = (p_1, \dots, p_m) \in \mathbb{N}^m$ output photons is given by:

$$\begin{aligned} P_{\text{BS}}(\mathbf{p}|n) &:= |\langle p_1 \dots p_m | \hat{U}(|1\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n})|^2 \\ &= \frac{|\text{Per}(U_{\mathbf{p}, \mathbf{1} \oplus \mathbf{0}})|^2}{\mathbf{p}!}. \end{aligned} \quad (16)$$

This model of quantum computation, while not being universal, is already capable of outperforming its classical counterparts: AA showed that the output probability distribution P_{BS} is hard to sample exactly classically for $m \geq 2n$, or the polynomial hierarchy of complexity classes collapses to its third level [6]. Moreover, under additional plausible conjectures, AA showed that this collapse holds for $n = o(m^2)$ even if only an efficient classical algorithm for *approximate sampling* exists (i.e. a classical algorithm which samples efficiently from a probability distribution that has a small total variation distance with the ideal Boson Sampling output probability distribution P_{BS}).

2.3 The MacMahon master theorem

The MacMahon master theorem is an important result in combinatorics which relates the permanent to the determinant:

Theorem (MacMahon master theorem [12]). *For any $m \times m$ matrix A ,*

$$\sum_{\mathbf{p} \in \mathbb{N}^m} \frac{z^{\mathbf{p}}}{\mathbf{p}!} \text{Per}(A_{\mathbf{p}, \mathbf{p}}) = \frac{1}{\text{Det}(I - ZA)}, \quad (17)$$

where $Z = \text{Diag}(\mathbf{z})$, with $\mathbf{z} \in \mathbb{C}^m$.

This theorem is particularly useful to derive short proofs of combinatorial identities: it expresses the permanent of an $m \times m$ matrix A with rows and columns repeated in the same way as the coefficient

$$\text{Per}(A_{\mathbf{p}, \mathbf{p}}) = \mathbf{p}! [\mathbf{z}^{\mathbf{p}}] \left(\frac{1}{\text{Det}(I - ZA)} \right), \quad (18)$$

where $Z = \text{Diag}(\mathbf{z})$ and $\mathbf{p} \in \mathbb{N}^m$, while the same permanent may also be expressed as the coefficient

$$\text{Per}(A_{\mathbf{p}, \mathbf{p}}) = \mathbf{p}! [\mathbf{z}^{\mathbf{p}}] (A\mathbf{z})^{\mathbf{p}}. \quad (19)$$

Picking a specific matrix A and a pattern \mathbf{p} and computing the above expressions yields combinatorial identities, a famous example being the short proof of Dixon's identity [13, 23], $\sum_{k=0}^{2n} (-1)^k \binom{2n}{k}^3 = (-1)^n \binom{3n}{n, n, n}$, by taking

$$A = \begin{pmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{pmatrix}, \quad (20)$$

and $\mathbf{p} = (2n, 2n, 2n)$ for $n \in \mathbb{N}^*$.

Various generalizations of the MacMahon master theorem have been introduced over the years [24–28]. In physics, this theorem plays an important role in the quantum theory of angular momentum [29] and is also oftentimes interpreted as an instance of the boson-fermion correspondence [25].

3 Main results

In this section, we summarize our main findings, which we prove in section 6. We obtain the following generalization of the MacMahon master theorem:

Theorem 1. *For all $m \times m$ matrices A and B ,*

$$\sum_{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m} \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}) \text{Per}(B_{\mathbf{q}, \mathbf{p}}) = \frac{1}{\text{Det}(I - XAYB)}, \quad (21)$$

where $X = \text{Diag}(\mathbf{x})$ and $Y = \text{Diag}(\mathbf{y})$, with $\mathbf{x}, \mathbf{y} \in \mathbb{C}^m$.

We further show that this generalization extends to N matrices:

Theorem 2. *Let $N \geq 2$, for all $m \times m$ matrices $A^{(1)}, \dots, A^{(N)}$,*

$$\sum_{\mathbf{p}_1, \dots, \mathbf{p}_N \in \mathbb{N}^m} \prod_{k=1}^N \frac{\mathbf{z}_k^{\mathbf{p}_k}}{\mathbf{p}_k!} \text{Per}(A_{\mathbf{p}_1, \mathbf{p}_2}^{(1)}) \text{Per}(A_{\mathbf{p}_2, \mathbf{p}_3}^{(2)}) \dots \text{Per}(A_{\mathbf{p}_N, \mathbf{p}_1}^{(N)}) = \frac{1}{\text{Det}(I - Z_1 A^{(1)} \dots Z_N A^{(N)}), \quad (22)$$

where $\mathbf{z}_k = (z_{k1}, \dots, z_{km}) \in \mathbb{C}^m$ and $Z_k = \text{Diag}(\mathbf{z}_k)$ for all $k \in \{1, \dots, N\}$.

As a corollary of Theorem 1, we obtain:

Corollary 1. *For any $m \times m$ matrix A and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ with $|\mathbf{p}| = |\mathbf{q}| = n \in \mathbb{N}$,*

$$\text{Per}(A_{\mathbf{p}, \mathbf{q}}) = \frac{\mathbf{p}! \mathbf{q}!}{n!} [\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}] (\mathbf{x}^T A \mathbf{y})^n. \quad (23)$$

As a consequence of Corollary 1, we obtain the following family of generating functions for the permanent:

Theorem 3. *Let $f(z) = \sum_{n=0}^{+\infty} f_n z^n$ be a series. For any $m \times m$ matrix A ,*

$$f(\mathbf{x}^T A \mathbf{y}) = \sum_{\substack{n \in \mathbb{N}, \mathbf{p}, \mathbf{q} \in \mathbb{N}^m \\ |\mathbf{p}| = |\mathbf{q}| = n}} f_n n! \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}), \quad (24)$$

where $\mathbf{x}, \mathbf{y} \in \mathbb{C}^m$. Equivalently,

$$\partial_{\mathbf{x}}^{\mathbf{p}} \partial_{\mathbf{y}}^{\mathbf{q}} f(\mathbf{x}^T A \mathbf{y}) \Big|_{\mathbf{x}=\mathbf{y}=\mathbf{0}} = \text{Per}(A_{\mathbf{p}, \mathbf{q}}) \partial_z^n f(z) \Big|_{z=0}, \quad (25)$$

for all $n \in \mathbb{N}$ and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ such that $|\mathbf{p}| = |\mathbf{q}| = n$. As a result, when $f_n \neq 0$,

$$\text{Per}(A_{\mathbf{p}, \mathbf{q}}) = \mathbb{E}_{\mathbf{x}, \mathbf{y} \in \mathbb{T}^m} \left[\frac{\mathbf{p}! \mathbf{q}!}{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}} \frac{f(\mathbf{x}^T A \mathbf{y})}{\partial_z^n f(z) \Big|_{z=0}} \right], \quad (26)$$

where the average is over random vectors with complex coefficients of modulus 1.

From this theorem we derive various permanent identities, the most notable one being a formula for the sum of two permanents:

Theorem 4. For all $m \times m$ matrices A and B , all $n \in \mathbb{N}$ and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ such that $|\mathbf{p}| = |\mathbf{q}| = n$,

$$\begin{aligned} & \text{Per}(A_{\mathbf{p}, \mathbf{q}}) + \text{Per}(B_{\mathbf{p}, \mathbf{q}}) \\ &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k}{\binom{n-1}{k}} \sum_{\substack{\mathbf{a}+\mathbf{b}+\mathbf{c}=\mathbf{p} \\ \mathbf{a}'+\mathbf{b}'+\mathbf{c}'=\mathbf{q} \\ |\mathbf{a}|=|\mathbf{b}|=|\mathbf{a}'|=|\mathbf{b}'|=k}} \frac{\mathbf{p}! \mathbf{q}!}{\mathbf{a}! \mathbf{b}! \mathbf{c}! \mathbf{a}'! \mathbf{b}'! \mathbf{c}'!} \text{Per}(A_{\mathbf{a}, \mathbf{a}'}) \text{Per}(B_{\mathbf{b}, \mathbf{b}'}) \text{Per}((A+B)_{\mathbf{c}, \mathbf{c}'}). \end{aligned} \quad (27)$$

In particular, when $\mathbf{p} = \mathbf{q} = \mathbf{1}$,

$$\text{Per}(A) + \text{Per}(B) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k}{\binom{n-1}{k}} \sum_{\substack{\mathbf{a}+\mathbf{b}+\mathbf{c}=\mathbf{1} \\ \mathbf{a}'+\mathbf{b}'+\mathbf{c}'=\mathbf{1} \\ |\mathbf{a}|=|\mathbf{b}|=|\mathbf{a}'|=|\mathbf{b}'|=k}} \text{Per}(A_{\mathbf{a}, \mathbf{a}'}) \text{Per}(B_{\mathbf{b}, \mathbf{b}'}) \text{Per}((A+B)_{\mathbf{c}, \mathbf{c}'}). \quad (28)$$

Section 5 is primarily devoted to quantum-inspired proofs of existing permanent identities, including a new proof of the MacMahon master theorem. Along the way, we obtain the following inner product formula:

Lemma 1. For all $\alpha \in \mathbb{C}$, all $\mathbf{p} \in \mathbb{N}^m$, all $n \leq m$, and any passive linear operation \hat{U} over m modes with unitary matrix U ,

$$\begin{aligned} \langle \mathbf{p} | \hat{U} (|\text{cat}_{\alpha}\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}) &= \frac{\alpha^n}{\sqrt{\sinh^n(|\alpha|^2)}} \langle \mathbf{p} | \hat{U} (|1\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}) \\ &= \frac{\alpha^n}{\sqrt{\sinh^n(|\alpha|^2)}} \frac{\text{Per}(U_{\mathbf{p}, \mathbf{1} \oplus \mathbf{0}})}{\sqrt{\mathbf{p}!}}. \end{aligned} \quad (29)$$

Lemma 1 has the following direct implications for the hardness of Boson Sampling with input cat states [20], which we discuss in section 7:

Theorem 5. Let $m \geq 2n$. Boson Sampling with input $|\text{cat}_{\alpha}\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}$ with $\alpha \in \mathbb{C}$ is hard to sample exactly classically unless the polynomial hierarchy of complexity classes collapses to its third level.

Moreover, assuming $|\alpha| = \mathcal{O}(n^{-1/4} \log^{1/4} n)$ and $n = o(m^2)$, Boson Sampling with input cat states is as hard to sample approximately as Boson Sampling with input single photons, i.e. Boson Sampling with input cat states is hard to sample approximately classically unless the polynomial hierarchy of complexity classes collapses to its third level modulo the complexity conjectures introduced by AA [6].

This result extends the arguments of [20]—which gave formal proof of classical hardness in the exact sampling case—to the approximate sampling case.

4 Discussion

In the next sections, we introduce quantum-inspired proofs of permanent identities. This approach allows us to give quantum-operational interpretations of seminal results, such as the MacMahon master theorem [12]. In particular, we show that this theorem can be seen as two facets of the same bosonic Gaussian amplitude.

This quantum-inspired approach also yields a breadth of new permanent identities. We give some examples of combinatorial applications of these identities in section 6 and we anticipate that they have many more. Beyond these purely combinatorial applications, it would be interesting to investigate whether our new identities may be used to obtain more efficient classical algorithms for computing or estimating the permanent. In particular, our Theorem 3 provides new estimators for the permanent which could be of interest, by minimizing the variance of these estimators over the choice of the analytic function f .

We use the formalism of linear optics with noninteracting bosons, but our approach can be applied more generally to linear optics with other types of particles. For instance, we expect our proof techniques to lead to determinant identities in the fermionic case [30], immanant identities in the case of partially distinguishable particles [31, 32], and additional permanent identities in the case of generalized bosons [33]. Moreover, graphical languages are currently being developed for linear optical quantum computations [34, 35], which could lead to graphical proofs of remarkable identities in combination with our approach.

Finally, our Theorem 5 gives solid complexity-theoretic foundations for the hardness of Boson Sampling with input cat states. Generation of such states has progressed tremendously in the recent years, thanks to circuit QED in particular [36–38]. We hope that these foundations will motivate an experimental demonstration of quantum speedup based on Schrödinger’s cat states.

5 Quantum-inspired proofs of permanent identities

In this section, we derive quantum-inspired proofs of existing permanent identities. Along the way, we obtain a generalization of Glynn’s formula [19] for the permanent of matrices with repeated rows and columns in Eq. (40), as well as a generalization of the Glynn–Kan formula [39] for the permanent of matrices with repeated rows and columns in Eq. (46).

5.1 Glynn’s formula

Glynn’s formula for the permanent of an $m \times m$ matrix $A = (a_{ij})_{1 \leq i, j \leq m}$ is [19]:

$$\text{Per}(A) = \frac{1}{2^{m-1}} \sum_{\substack{\mathbf{x} \in \{-1, 1\}^m \\ x_1 = 1}} x_1 \dots x_m \prod_{i=1}^m \left(\sum_{j=1}^m a_{ij} x_j \right). \quad (30)$$

By symmetry, it is equivalent to the identity

$$\text{Per}(A) = \frac{1}{2^m} \sum_{\mathbf{x} \in \{-1, 1\}^m} x_1 \dots x_m \prod_{i=1}^m \left(\sum_{j=1}^m a_{ij} x_j \right). \quad (31)$$

Proof of Glynn’s formula. To prove this identity using quantum-mechanical tools, let us introduce the unnormalized cat state $|\tilde{\text{cat}}_\alpha\rangle := \frac{1}{2\alpha}(|\alpha\rangle - |-\alpha\rangle)$, for $\alpha \in \mathbb{C}$. With Eq. (6), $\lim_{\alpha \rightarrow 0} |\tilde{\text{cat}}_\alpha\rangle = |1\rangle$ in trace distance. Hence, with Eq. (9),

$$\text{Per}(U) = \lim_{\alpha \rightarrow 0} \langle 1 \dots 1 | \hat{U} | \tilde{\text{cat}}_\alpha \dots \tilde{\text{cat}}_\alpha \rangle, \quad (32)$$

where \hat{U} is a passive linear operation over m modes with unitary matrix $U = (u_{ij})_{1 \leq i, j \leq m}$. We compute the right hand side of this equation:

$$\begin{aligned}
\langle 1 \dots 1 | \hat{U} | \tilde{\text{c\hat{a}}}_\alpha \dots \tilde{\text{c\hat{a}}}_\alpha \rangle &= \frac{1}{(2\alpha)^m} \sum_{\mathbf{x} \in \{-1, 1\}^m} x_1 \dots x_m \langle 1 \dots 1 | \hat{U} | x_1 \alpha \dots x_m \alpha \rangle \\
&= \frac{1}{(2\alpha)^m} \sum_{\mathbf{x} \in \{-1, 1\}^m} x_1 \dots x_m \langle 1 \dots 1 | (\alpha U \mathbf{x})_1 \dots (\alpha U \mathbf{x})_m \rangle \\
&= \frac{1}{(2\alpha)^m} \sum_{\mathbf{x} \in \{-1, 1\}^m} x_1 \dots x_m e^{-\frac{1}{2}|\alpha|^2 \|\mathbf{U}\mathbf{x}\|^2} \prod_{i=1}^m (\alpha U \mathbf{x})_i \\
&= \frac{e^{-\frac{m}{2}|\alpha|^2}}{2^m} \sum_{\mathbf{x} \in \{-1, 1\}^m} x_1 \dots x_m \prod_{i=1}^m \left(\sum_{j=1}^m u_{ij} x_j \right),
\end{aligned} \tag{33}$$

where we used Eq. (12) in the second line, Eq. (6) in the third line, and the fact that U is unitary in the last line. Taking the limit when $\alpha \rightarrow 0$ yields Glynn's formula for unitary matrices. \square

With the same calculations, we may obtain a more general version of Eq. (33): for all $\alpha \in \mathbb{C}$ and $\mathbf{p} \in \mathbb{N}^m$,

$$\begin{aligned}
\langle p_1 \dots p_m | \hat{U} | \tilde{\text{c\hat{a}}}_\alpha \dots \tilde{\text{c\hat{a}}}_\alpha \rangle &= \frac{1}{(2\alpha)^m} \sum_{\mathbf{x} \in \{-1, 1\}^m} x_1 \dots x_m e^{-\frac{1}{2}|\alpha|^2 \|\mathbf{U}\mathbf{x}\|^2} \prod_{i=1}^m \frac{[(\alpha U \mathbf{x})_i]^{p_i}}{\sqrt{p_i!}} \\
&= \frac{\alpha^{|\mathbf{p}|-m} e^{-\frac{m}{2}|\alpha|^2}}{2^m \sqrt{\mathbf{p}!}} \sum_{\mathbf{x} \in \{-1, 1\}^m} x_1 \dots x_m \prod_{i=1}^m \left(\sum_{j=1}^m u_{ij} x_j \right)^{p_i}.
\end{aligned} \tag{34}$$

Note that when $|\mathbf{p}| < m$ all products in the sum have at least one x_i missing. Hence, by symmetry, $\langle p_1 \dots p_m | \hat{U} | \tilde{\text{c\hat{a}}}_\alpha \dots \tilde{\text{c\hat{a}}}_\alpha \rangle = 0$ when $|\mathbf{p}| < m$. With Eq. (9), taking the limit when $\alpha \rightarrow 0$ yields a version of Glynn's formula for unitary matrices with repeated rows:

$$\text{Per}(U_{\mathbf{p}, \mathbf{1}}) = \frac{\delta_{|\mathbf{p}|, m}}{2^m} \sum_{\mathbf{x} \in \{-1, 1\}^m} x_1 \dots x_m \prod_{i=1}^m \left(\sum_{j=1}^m u_{ij} x_j \right)^{p_i}, \tag{35}$$

where δ is the Kronecker symbol.

So far, all identities are derived for unitary matrices U . In order to retrieve the same identity for a general (nonzero) matrix $A = (a_{ij})_{1 \leq i, j \leq n}$ of size n , we can embed $\frac{1}{\|A\|} A$ as a submatrix of a unitary matrix U of size $m = 2n$ and compute:

$$\frac{\text{Per}(A_{\mathbf{q}, \mathbf{1}})}{\sqrt{\mathbf{q}!}} = \|A\|^{|\mathbf{q}|} \lim_{\alpha \rightarrow 0} (\langle q_1 \dots q_n | \otimes \langle 0 |^{\otimes n}) \hat{U} (| \tilde{\text{c\hat{a}}}_\alpha \rangle^{\otimes n} \otimes | 0 \rangle^{\otimes n}). \tag{36}$$

To do so, we compute a slightly more general inner product: for $\alpha \in \mathbb{C}$, $\mathbf{p} \in \mathbb{N}^m$ and $n \leq m$,

$$\begin{aligned}
&\langle p_1 \dots p_m | \hat{U} (| \tilde{\text{c\hat{a}}}_\alpha \rangle^{\otimes n} \otimes | 0 \rangle^{\otimes m-n}) \\
&= \frac{1}{(2\alpha)^n} \sum_{\mathbf{x} \in \{-1, 1\}^n} x_1 \dots x_n \langle p_1 \dots p_m | \hat{U} | x_1 \alpha \dots x_n \alpha 0 \dots 0 \rangle \\
&= \frac{1}{(2\alpha)^n} \sum_{\mathbf{x} \in \{-1, 1\}^n} x_1 \dots x_n e^{-\frac{1}{2}|\alpha|^2 \|\mathbf{U}(\mathbf{x} \oplus \mathbf{0})\|^2} \prod_{i=1}^m \frac{[(\alpha U(\mathbf{x} \oplus \mathbf{0}))_i]^{p_i}}{\sqrt{p_i!}} \\
&= \frac{\alpha^{|\mathbf{p}|-n} e^{-\frac{n}{2}|\alpha|^2}}{2^n \sqrt{\mathbf{p}!}} \sum_{\mathbf{x} \in \{-1, 1\}^n} x_1 \dots x_n \prod_{i=1}^m \left(\sum_{j=1}^n u_{ij} x_j \right)^{p_i}.
\end{aligned} \tag{37}$$

In particular, setting $m = 2n$ and $\mathbf{p} = \mathbf{q} \oplus \mathbf{0}$ for $\mathbf{q} \in \mathbb{N}^n$,

$$(\langle q_1 \dots q_n | \otimes \langle 0 |^{\otimes n}) \hat{U} (|\tilde{\text{cat}}_\alpha\rangle^{\otimes n} \otimes |0\rangle^{\otimes n}) = \frac{\alpha^{|\mathbf{q}|-n} e^{-\frac{n}{2}|\alpha|^2}}{2^n \sqrt{\mathbf{q}!}} \sum_{\mathbf{x} \in \{-1,1\}^n} x_1 \dots x_n \prod_{i=1}^n \left(\sum_{j=1}^n u_{ij} x_j \right)^{q_i}. \quad (38)$$

Choosing $u_{ij} = \frac{1}{\|A\|} a_{ij}$ for $1 \leq i, j \leq n$ and letting α go to 0 proves the claim: the left hand side converges to $\frac{1}{\sqrt{\mathbf{q}!}} \text{Per}(U_{\mathbf{q} \oplus \mathbf{0}, \mathbf{1} \oplus \mathbf{0}}) = \frac{1}{\|A\|^{|\mathbf{q}|} \sqrt{\mathbf{q}!}} \text{Per}(A_{\mathbf{q}, \mathbf{1}})$, while the right hand side converges to $\frac{\delta_{|\mathbf{q}|, n}}{\|A\|^{|\mathbf{q}|} 2^n \sqrt{\mathbf{q}!}} \sum_{\mathbf{x} \in \{-1,1\}^n} x_1 \dots x_n \prod_{i=1}^n \left(\sum_{j=1}^n a_{ij} x_j \right)^{q_i}$. Hence,

$$\text{Per}(A_{\mathbf{q}, \mathbf{1}}) = \frac{\delta_{|\mathbf{q}|, n}}{2^n} \sum_{\mathbf{x} \in \{-1,1\}^n} x_1 \dots x_n \prod_{i=1}^n \left(\sum_{j=1}^n a_{ij} x_j \right)^{q_i}. \quad (39)$$

A similar generalization of Glynn's formula for matrices with repeated rows (or columns) based on roots of unity has previously appeared in [40]. Combining this generalization with the above reasoning yields, for any $m \times m$ matrix $A = (a_{ij})_{1 \leq i, j \leq m}$ and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ with $|\mathbf{p}| = |\mathbf{q}| = n$,

$$\text{Per}(A_{\mathbf{p}, \mathbf{q}}) = \frac{\mathbf{q}!}{n^m} \sum_{\mathbf{x} \in \{1, e^{\frac{2i\pi}{n}}, \dots, e^{\frac{2i(n-1)\pi}{n}}\}^m} \mathbf{x}^{-\mathbf{q}} (A\mathbf{x})^{\mathbf{p}}, \quad (40)$$

which is a version of Glynn's formula for matrices with repeated rows *and* columns. This formula, which appears to be new, can be easily proven by expanding the product $(A\mathbf{x})^{\mathbf{p}} = \prod_{i=1}^m \left(\sum_{j=1}^m a_{ij} x_j \right)^{p_i}$ and using $\sum_{l=0}^{n-1} e^{\frac{2ikl\pi}{n}} = \delta_{k,n}$ for all $k \in \{1, \dots, n\}$.

5.2 The Glynn–Kan formula

A symmetrized version of Glynn's formula for the permanent of an $m \times m$ matrix A has been recently derived [39] under the name Glynn–Kan formula:

$$\text{Per}(A) = \frac{1}{4^m m!} \sum_{\mathbf{x}, \mathbf{y} \in \{-1,1\}^m} x_1 \dots x_m y_1 \dots y_m (\mathbf{x}^T A \mathbf{y})^m. \quad (41)$$

Proof of the Glynn–Kan formula. To prove this identity using quantum-mechanical tools, we again use the unnormalized cat state $|\tilde{\text{cat}}_\alpha\rangle := \frac{1}{2\alpha}(|\alpha\rangle - |-\alpha\rangle)$, as in the previous section. With Eq. (6), $\lim_{\alpha \rightarrow 0} |\tilde{\text{cat}}_\alpha\rangle = |1\rangle$ in trace distance. Hence, with Eq. (9),

$$\text{Per}(U) = \lim_{\alpha \rightarrow 0} \langle \tilde{\text{cat}}_\alpha \dots \tilde{\text{cat}}_\alpha | \hat{U} | \tilde{\text{cat}}_\alpha \dots \tilde{\text{cat}}_\alpha \rangle, \quad (42)$$

where \hat{U} is a passive linear operation over m modes with unitary matrix U . We compute the right hand side of this equation:

$$\begin{aligned} & \langle \tilde{\text{cat}}_\alpha \dots \tilde{\text{cat}}_\alpha | \hat{U} | \tilde{\text{cat}}_\alpha \dots \tilde{\text{cat}}_\alpha \rangle \\ &= \frac{1}{4^m |\alpha|^{2m}} \sum_{\mathbf{x}, \mathbf{y} \in \{-1,1\}^m} x_1 \dots x_m y_1 \dots y_m \langle x_1 \alpha \dots x_m \alpha | \hat{U} | y_1 \alpha \dots y_m \alpha \rangle \\ &= \frac{1}{4^m |\alpha|^{2m}} \sum_{\mathbf{x}, \mathbf{y} \in \{-1,1\}^m} x_1 \dots x_m y_1 \dots y_m \langle x_1 \alpha \dots x_m \alpha | (\alpha U \mathbf{y})_1 \dots (\alpha U \mathbf{y})_m \rangle \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{4^m |\alpha|^{2m}} \sum_{\mathbf{x}, \mathbf{y} \in \{-1, 1\}^m} x_1 \dots x_m y_1 \dots y_m e^{-\frac{1}{2} |\alpha|^2 \|\mathbf{x}\|^2} e^{-\frac{1}{2} |\alpha|^2 \|U\mathbf{y}\|^2} \prod_{i=1}^m e^{x_i \alpha^* (\alpha U \mathbf{y})_i} \quad (43) \\
&= \frac{e^{-m|\alpha|^2}}{4^m |\alpha|^{2m}} \sum_{\mathbf{x}, \mathbf{y} \in \{-1, 1\}^m} x_1 \dots x_m y_1 \dots y_m e^{|\alpha|^2 \mathbf{x}^T U \mathbf{y}} \\
&= \frac{e^{-m|\alpha|^2}}{4^m} \sum_{\mathbf{x}, \mathbf{y} \in \{-1, 1\}^m} x_1 \dots x_m y_1 \dots y_m \sum_{k=0}^{+\infty} \frac{|\alpha|^{2k-2n}}{k!} (\mathbf{x}^T U \mathbf{y})^k,
\end{aligned}$$

where we used Eq. (12) in the third line, Eq. (7) in the fourth line, and the fact that U is unitary in the fifth line. For $k < m$, all products in the expansion of $(\mathbf{x}^T U \mathbf{y})^k$ have at least one x_i missing. Hence, by symmetry, the terms for $k < m$ vanish and we obtain

$$\langle \tilde{\text{cat}}_\alpha \dots \tilde{\text{cat}}_\alpha | \hat{U} | \tilde{\text{cat}}_\alpha \dots \tilde{\text{cat}}_\alpha \rangle = \frac{e^{-m|\alpha|^2}}{4^m} \sum_{\mathbf{x}, \mathbf{y} \in \{-1, 1\}^m} x_1 \dots x_m y_1 \dots y_m \sum_{l=0}^{\infty} \frac{|\alpha|^{2l}}{(m+l)!} (\mathbf{x}^T U \mathbf{y})^{m+l}. \quad (44)$$

Finally, taking the limit when α goes to 0 yields

$$\text{Per}(U) = \frac{1}{4^m m!} \sum_{\mathbf{x}, \mathbf{y} \in \{-1, 1\}^m} x_1 \dots x_m y_1 \dots y_m (\mathbf{x}^T U \mathbf{y})^m. \quad (45)$$

This proves the Glynn–Kan formula for a unitary matrix U . Once again, the proof extends straightforwardly to any matrix A of size n by embedding $\frac{1}{\|A\|} A$ as a submatrix of a unitary matrix U of size $2n$ and computing $\lim_{\alpha \rightarrow 0} (\langle \tilde{\text{cat}}_\alpha |^{\otimes n} \otimes \langle 0 |^{\otimes n}) \hat{U} (| \tilde{\text{cat}}_\alpha \rangle^{\otimes n} \otimes | 0 \rangle^{\otimes n})$. \square

Similar to Eq. (40), we obtain a version of the Glynn–Kan formula for matrices with repeated rows and columns using roots of unity as

$$\text{Per}(A_{\mathbf{p}, \mathbf{q}}) = \frac{\mathbf{p}! \mathbf{q}!}{(2n)^m m!} \sum_{\mathbf{x}, \mathbf{y} \in \{1, e^{\frac{2i\pi}{n}}, \dots, e^{\frac{2i(n-1)\pi}{n}}\}^m} \mathbf{x}^{-\mathbf{p}} \mathbf{y}^{-\mathbf{q}} (\mathbf{x}^T A \mathbf{y})^n, \quad (46)$$

for any $m \times m$ matrix $A = (a_{ij})_{1 \leq i, j \leq m}$ and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ with $|\mathbf{p}| = |\mathbf{q}| = n$. This formula, which also appears to be new, can be easily proven using our Corollary 1 by expanding the product $(\mathbf{x}^T A \mathbf{y})^n$ and using $\sum_{l=0}^{n-1} e^{\frac{2ikl\pi}{n}} = \delta_{k,n}$ for all $k \in \{1, \dots, n\}$.

5.3 The Cauchy–Binet theorem

The Cauchy–Binet theorem for the permanent expresses the permanent of the product of two $m \times m$ matrices A and B as a sum of products involving permanents of submatrices of these two matrices [1, 2]: for all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$,

$$\text{Per}((AB)_{\mathbf{p}, \mathbf{q}}) = \sum_{\mathbf{k} \in \mathbb{N}^m} \frac{1}{\mathbf{k}!} \text{Per}(A_{\mathbf{p}, \mathbf{k}}) \text{Per}(B_{\mathbf{k}, \mathbf{q}}). \quad (47)$$

Proof of the Cauchy–Binet theorem. Using Eq. (9) gives a quick quantum-inspired proof of this identity: for U and V two $m \times m$ unitary matrices and for all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$,

$$\begin{aligned}
\text{Per}((UV)_{\mathbf{p}, \mathbf{q}}) &= \sqrt{\mathbf{p}! \mathbf{q}!} \langle \mathbf{p} | \widehat{UV} | \mathbf{q} \rangle \\
&= \sqrt{\mathbf{p}! \mathbf{q}!} \langle \mathbf{p} | \widehat{U} \widehat{V} | \mathbf{q} \rangle
\end{aligned}$$

$$\begin{aligned}
&= \sqrt{\mathbf{p}! \mathbf{q}!} \langle \mathbf{p} | \hat{U} \left(\sum_{\mathbf{k} \in \mathbb{N}^m} |\mathbf{k}\rangle \langle \mathbf{k}| \right) \hat{V} | \mathbf{q} \rangle \quad (48) \\
&= \sum_{\mathbf{k} \in \mathbb{N}^m} \sqrt{\mathbf{p}! \mathbf{q}!} \langle \mathbf{p} | \hat{U} |\mathbf{k}\rangle \langle \mathbf{k}| \hat{V} | \mathbf{q} \rangle \\
&= \sum_{\mathbf{k} \in \mathbb{N}^m} \frac{1}{\mathbf{k}!} \text{Per}(U_{\mathbf{p}, \mathbf{k}}) \text{Per}(V_{\mathbf{k}, \mathbf{q}}),
\end{aligned}$$

where we used the fact that Fock states form a basis in the third line and where we used Eq. (9) once in the first line and twice in the last line.

In order to retrieve the formula for general matrices A and B , we can embed $\frac{1}{\|A\|}A$ as a submatrix of a unitary matrix U and $\frac{1}{\|B\|}B$ as a submatrix of a unitary matrix V and compute $\text{Per}((UV)_{\mathbf{p} \oplus \mathbf{0}, \mathbf{q} \oplus \mathbf{0}})$. \square

5.4 Generating functions

The permanent may be seen as a monomial coefficient in the Taylor expansion of various functions [1, 2]. For example, the permanent of an $m \times m$ matrix $A = (a_{ij})_{1 \leq i, j \leq m}$ with rows repeated according to $\mathbf{p} \in \mathbb{N}^m$ and columns repeated according to $\mathbf{q} \in \mathbb{N}^m$ is given by the coefficient of $\mathbf{z}^{\mathbf{q}} = z_1^{q_1} \dots z_m^{q_m}$ in

$$\mathbf{q}!(A\mathbf{z})^{\mathbf{p}} = \prod_{i=1}^m q_i! \left(\sum_{j=1}^m a_{ij} z_j \right)^{p_i}. \quad (49)$$

This may be thought of as the ‘monomial version’ of Glynn’s formula in Eq. (39). Formally:

$$\sum_{\mathbf{q} \in \mathbb{N}^m} \frac{\mathbf{z}^{\mathbf{q}}}{\mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}) = (A\mathbf{z})^{\mathbf{p}}, \quad (50)$$

for all $\mathbf{z} \in \mathbb{C}^m$.

Proof of ‘Glynn’s monomial formula’. To prove this relation with quantum mechanical tools, we fix $\mathbf{p} \in \mathbb{N}^m$, $\boldsymbol{\alpha} \in \mathbb{C}^m$, a unitary matrix U of size m , and we compute:

$$\begin{aligned}
\sum_{\mathbf{q} \in \mathbb{N}^m} \frac{\boldsymbol{\alpha}^{\mathbf{q}}}{\mathbf{q}!} \text{Per}(U_{\mathbf{p}, \mathbf{q}}) &= \sqrt{\mathbf{p}!} e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2} \langle \mathbf{p} | \hat{U} \left(\sum_{\mathbf{q} \in \mathbb{N}^m} |\mathbf{q}\rangle \langle \mathbf{q}| \right) | \boldsymbol{\alpha} \rangle \\
&= \sqrt{\mathbf{p}!} e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2} \langle \mathbf{p} | \hat{U} | \boldsymbol{\alpha} \rangle \\
&= \sqrt{\mathbf{p}!} e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2} \langle \mathbf{p} | (U \boldsymbol{\alpha}) \rangle \\
&= e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2 - \frac{1}{2} \|U \boldsymbol{\alpha}\|^2} (U \boldsymbol{\alpha})^{\mathbf{p}} \\
&= (U \boldsymbol{\alpha})^{\mathbf{p}},
\end{aligned} \quad (51)$$

where we used Eqs. (6) and (9) in the first line, the fact that Fock states form a basis in the second line, Eq. (12) in the third line, Eq. (6) again in the fourth line, and the fact that U is unitary in the last line. Once again, the relation for a generic matrix A is obtained by embedding it as a submatrix of a unitary matrix U . \square

Another generating function for the permanent is due to Jackson [41]: for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^m$,

$$\sum_{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m} \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}) = e^{\mathbf{x}^T A \mathbf{y}}. \quad (52)$$

Note that this relation implies Corollary 1, by expanding the Taylor series of the exponential and considering the $\mathbf{x}^p \mathbf{y}^q$ coefficient when $|\mathbf{p}| = |\mathbf{q}| = n \in \mathbb{N}^*$ (an alternative proof of this result is given in the next section):

$$\text{Per}(A_{\mathbf{p},\mathbf{q}}) = \frac{\mathbf{p}! \mathbf{q}!}{n!} [\mathbf{x}^p \mathbf{y}^q] (\mathbf{x}^T A \mathbf{y})^n. \quad (53)$$

Proof of Jackson's formula. Jackson's formula may be derived using quantum mechanical tools in a similar way: for $\boldsymbol{\alpha}, \boldsymbol{\beta} \in \mathbb{C}^m$ and a unitary matrix U ,

$$\begin{aligned} \sum_{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m} \frac{\boldsymbol{\alpha}^p \boldsymbol{\beta}^q}{\mathbf{p}! \mathbf{q}!} \text{Per}(U_{\mathbf{p},\mathbf{q}}) &= e^{\frac{1}{2}\|\boldsymbol{\alpha}\|^2 + \frac{1}{2}\|\boldsymbol{\beta}\|^2} \langle \boldsymbol{\alpha}^* | \left(\sum_{\mathbf{p} \in \mathbb{N}^m} |\mathbf{p}\rangle \langle \mathbf{p}| \right) \hat{U} \left(\sum_{\mathbf{q} \in \mathbb{N}^m} |\mathbf{q}\rangle \langle \mathbf{q}| \right) | \boldsymbol{\beta} \rangle \\ &= e^{\frac{1}{2}\|\boldsymbol{\alpha}\|^2 + \frac{1}{2}\|\boldsymbol{\beta}\|^2} \langle \boldsymbol{\alpha}^* | \hat{U} | \boldsymbol{\beta} \rangle \\ &= e^{\frac{1}{2}\|\boldsymbol{\alpha}\|^2 + \frac{1}{2}\|\boldsymbol{\beta}\|^2} \langle \boldsymbol{\alpha}^* | U \boldsymbol{\beta} \rangle \\ &= e^{\frac{1}{2}\|\boldsymbol{\beta}\|^2 - \frac{1}{2}\|U\boldsymbol{\beta}\|^2} e^{\boldsymbol{\alpha}^T U \boldsymbol{\beta}} \\ &= e^{\boldsymbol{\alpha}^T U \boldsymbol{\beta}}, \end{aligned} \quad (54)$$

where we used Eqs. (6) and (9) in the first line, the fact that Fock states form a basis in the second line, Eq. (12) in the third line, Eq. (7) in the fourth line, and the fact that U is unitary in the last line. Once again, the relation for a generic matrix A is obtained by embedding it as a submatrix of a unitary matrix U . \square

We conclude this section with arguably one of the most remarkable permanent identities, the MacMahon master theorem [12], which relates the permanent and the determinant through a generating function (see section 2.3):

$$\sum_{\mathbf{p} \in \mathbb{N}^m} \frac{\mathbf{z}^{\mathbf{p}}}{\mathbf{p}!} \text{Per}(A_{\mathbf{p},\mathbf{p}}) = \frac{1}{\text{Det}(I - ZA)}, \quad (55)$$

where $Z = \text{Diag}(\mathbf{z})$, with $\mathbf{z} \in \mathbb{C}^m$.

Proof of the MacMahon master theorem. To prove this relation with quantum mechanical tools, we show that the MacMahon master theorem describes two different ways of computing an inner product between two Gaussian states.

For $\lambda \in \mathbb{C}$ and $|\lambda| < 1$, we make use of (unnormalized) two-mode squeezed states of the form $|\lambda\rangle = \sum_{p \geq 0} \lambda^p |pp\rangle$. We write $|\boldsymbol{\lambda}\rangle$ with $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_m) \in \mathbb{C}^m$ a tensor product of m two-mode squeezed states (note that we are associating mode 1 with mode $m+1$, mode 2 with mode $m+2$, and so on). For U a unitary matrix and $\boldsymbol{\lambda}, \boldsymbol{\mu} \in \mathbb{C}^m$, with $|\lambda_k| < 1$ and $|\mu_k| < 1$ for all $k \in \{1, \dots, m\}$, we compute, using Eq. (9):

$$\sum_{\mathbf{p} \in \mathbb{N}^m} \frac{\boldsymbol{\lambda}^p \boldsymbol{\mu}^p}{\mathbf{p}!} \text{Per}(U_{\mathbf{p},\mathbf{p}}) = \sum_{\mathbf{p} \in \mathbb{N}^m} \boldsymbol{\lambda}^p \boldsymbol{\mu}^p \langle \mathbf{p} | \hat{U} | \mathbf{p} \rangle. \quad (56)$$

By Eq. (8), we have

$$\boldsymbol{\lambda}^p = \langle \boldsymbol{\lambda}^* | \mathbf{p} \mathbf{p} \rangle. \quad (57)$$

Moreover, a quick computation in Fock basis shows that

$$\boldsymbol{\mu}^p \langle \mathbf{p} | \hat{U} | \mathbf{p} \rangle = \langle \mathbf{p} \mathbf{p} | \hat{I} \otimes \hat{U} | \boldsymbol{\mu} \rangle. \quad (58)$$

Hence, Eq. (56) rewrites

$$\begin{aligned}
\sum_{\mathbf{p} \in \mathbb{N}^m} \frac{\lambda^{\mathbf{p}} \mu^{\mathbf{p}}}{\mathbf{p}!} \text{Per}(U_{\mathbf{p}, \mathbf{p}}) &= \sum_{\mathbf{p} \in \mathbb{N}^m} \langle \boldsymbol{\lambda}^* | \mathbf{p}\mathbf{p} \rangle \langle \mathbf{p}\mathbf{p} | \hat{I} \otimes \hat{U} | \boldsymbol{\mu} \rangle \\
&= \langle \boldsymbol{\lambda}^* | \left(\sum_{\mathbf{p} \in \mathbb{N}^m} |\mathbf{p}\mathbf{p}\rangle \langle \mathbf{p}\mathbf{p}| \right) \hat{I} \otimes \hat{U} | \boldsymbol{\mu} \rangle \\
&= \langle \boldsymbol{\lambda}^* | \hat{I} \otimes \hat{U} | \boldsymbol{\mu} \rangle,
\end{aligned} \tag{59}$$

where in the last line we used $\langle \boldsymbol{\lambda}^* | \left(\sum_{\mathbf{p} \in \mathbb{N}^m} |\mathbf{p}\mathbf{p}\rangle \langle \mathbf{p}\mathbf{p}| \right) = \langle \boldsymbol{\lambda}^* |$, which can be checked in Fock basis.

We now compute the Gaussian inner product $\langle \boldsymbol{\lambda}^* | \hat{I} \otimes \hat{U} | \boldsymbol{\mu} \rangle$. From Eq. (10), the unitary matrix associated to the passive linear operation $\hat{I} \otimes \hat{U}$ is $I \oplus U$. With Eqs. (13-15), we thus obtain:

$$\langle \boldsymbol{\lambda}^* | \hat{I} \otimes \hat{U} | \boldsymbol{\mu} \rangle = \frac{1}{\sqrt{\text{Det}(V_{I \oplus U}(\boldsymbol{\lambda}, \boldsymbol{\mu}))}}, \tag{60}$$

where

$$V_{I \oplus U}(\boldsymbol{\lambda}, \boldsymbol{\mu}) = \begin{pmatrix} -(I \oplus U)V_{\boldsymbol{\mu}}(I \oplus U)^T & I_{2m} \\ I_{2m} & -V_{\boldsymbol{\lambda}} \end{pmatrix}, \tag{61}$$

where for all $\mathbf{w} \in \mathbb{C}^m$,

$$V_{\mathbf{w}} = \begin{pmatrix} 0_m & \text{Diag}(\mathbf{w}) \\ \text{Diag}(\mathbf{w}) & 0_m \end{pmatrix}. \tag{62}$$

We have

$$\begin{aligned}
\text{Det}(V_{I \oplus U}(\boldsymbol{\lambda}, \boldsymbol{\mu})) &= \text{Det}(I - (I \oplus U)V_{\boldsymbol{\mu}}(I \oplus U)^T V_{\boldsymbol{\lambda}}) \\
&= \text{Det} \left(I - (I \oplus U) \begin{pmatrix} 0_m & \text{Diag}(\boldsymbol{\mu}) \\ \text{Diag}(\boldsymbol{\mu}) & 0_m \end{pmatrix} (I \oplus U)^T \begin{pmatrix} 0_m & \text{Diag}(\boldsymbol{\lambda}) \\ \text{Diag}(\boldsymbol{\lambda}) & 0_m \end{pmatrix} \right) \\
&= \text{Det} \begin{pmatrix} I - \text{Diag}(\boldsymbol{\mu})U^T \text{Diag}(\boldsymbol{\lambda}) & 0_m \\ 0_m & I - U \text{Diag}(\boldsymbol{\mu}) \text{Diag}(\boldsymbol{\lambda}) \end{pmatrix} \\
&= \text{Det}(I - \text{Diag}(\boldsymbol{\mu})U^T \text{Diag}(\boldsymbol{\lambda})) \text{Det}(I - U \text{Diag}(\boldsymbol{\mu}) \text{Diag}(\boldsymbol{\lambda})) \\
&= \text{Det}(I - \text{Diag}(\boldsymbol{\lambda}) \text{Diag}(\boldsymbol{\mu}) U)^2,
\end{aligned} \tag{63}$$

where the last line is obtained by assuming that $\text{Diag}(\boldsymbol{\lambda})$ is invertible (wlog by density) and taking the transpose for the first determinant, and using $\text{Diag}(\boldsymbol{\mu}) \text{Diag}(\boldsymbol{\lambda}) = \text{Diag}(\boldsymbol{\lambda}) \text{Diag}(\boldsymbol{\mu})$ for the second determinant. Combining Eqs. (59), (60) and (63), we obtain

$$\sum_{\mathbf{p} \in \mathbb{N}^m} \frac{\lambda^{\mathbf{p}} \mu^{\mathbf{p}}}{\mathbf{p}!} \text{Per}(U_{\mathbf{p}, \mathbf{p}}) = \langle \boldsymbol{\lambda}^* | \hat{I} \otimes \hat{U} | \boldsymbol{\mu} \rangle = \frac{1}{\text{Det}(I - \text{Diag}(\boldsymbol{\lambda}) \text{Diag}(\boldsymbol{\mu}) U)}. \tag{64}$$

Setting $\mathbf{z} = (\lambda_1 \mu_1, \dots, \lambda_p \mu_p)$ concludes the proof. The relation for a generic matrix A of size n is obtained by embedding it as a submatrix of a unitary matrix U of size $2n$ and taking $\mathbf{z} = (z_1, \dots, z_n, 0, \dots, 0) \in \mathbb{C}^{2n}$. \square

6 New quantum-inspired permanent identities

In this section, we derive new quantum-inspired identities for the permanent and we give some combinatorial applications of these identities.

6.1 Generalizations of the MacMahon master theorem

In this section, we introduce new generalizations of the MacMahon master theorem (see section 2.3).

We first derive new quantum-inspired identities involving the permanent of even-sized matrices: for M a $(2m) \times (2m)$ matrix,

$$\text{Per}(M) = [z^m] \left(\frac{1}{4^m} \sum_{\mathbf{x}, \mathbf{y} \in \{-1, 1\}^m} \frac{x_1 \dots x_m y_1 \dots y_m}{\sqrt{\text{Det}(I - zV_{\mathbf{x}} M V_{\mathbf{y}} M^T)}} \right), \quad (65)$$

and, for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^m$,

$$\sum_{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m} \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(M_{\mathbf{p} \oplus \mathbf{p}, \mathbf{q} \oplus \mathbf{q}}) = \frac{1}{\sqrt{\text{Det}(I - V_{\mathbf{x}} M V_{\mathbf{y}} M^T)}}, \quad (66)$$

where for all $\mathbf{w} \in \mathbb{C}^m$,

$$V_{\mathbf{w}} = \begin{pmatrix} 0_m & \text{Diag}(\mathbf{w}) \\ \text{Diag}(\mathbf{w}) & 0_m \end{pmatrix}. \quad (67)$$

As a direct consequence, when $M = A \oplus B$, with A and B two $m \times m$ matrices, we obtain Theorem 1:

$$\sum_{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m} \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}) \text{Per}(B_{\mathbf{p}, \mathbf{q}}) = \frac{1}{\text{Det}(I - X A Y B^T)}, \quad (68)$$

where $X = \text{Diag}(x_1, \dots, x_m)$ and $Y = \text{Diag}(y_1, \dots, y_m)$. In particular, $\text{Per}(A)\text{Per}(B)$ is given by the $x_1 \dots x_m y_1 \dots y_m$ coefficient of $1/\text{Det}(I - X A Y B^T)$. As a consequence, we obtain Corollary 1: for all $n \in \mathbb{N}^*$ and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ with $|\mathbf{p}| = |\mathbf{q}| = n$,

$$\text{Per}(A_{\mathbf{p}, \mathbf{q}}) = \frac{\mathbf{p}! \mathbf{q}!}{n!} [\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}] (\mathbf{x}^T A \mathbf{y})^n. \quad (69)$$

We also generalize Eq. (68) to the case of N matrices to obtain Theorem 2: for all $n \times n$ matrices $A^{(1)}, \dots, A^{(N)}$,

$$\begin{aligned} \sum_{\mathbf{p}_1, \dots, \mathbf{p}_N \in \mathbb{N}^m} \prod_{k=1}^N \frac{z_k^{\mathbf{p}_k}}{\mathbf{p}_k!} \text{Per}(A_{\mathbf{p}_1, \mathbf{p}_2}^{(1)}) \text{Per}(A_{\mathbf{p}_2, \mathbf{p}_3}^{(2)}) \dots \text{Per}(A_{\mathbf{p}_N, \mathbf{p}_1}^{(N)}) \\ = \frac{1}{\text{Det}(I - Z_1 A^{(1)} \dots Z_N A^{(N)})}, \end{aligned} \quad (70)$$

where $\mathbf{z}_k = (z_{k1}, \dots, z_{km}) \in \mathbb{C}^m$ and $Z_k = \text{Diag}(\mathbf{z}_k)$ for all $k \in \{1, \dots, N\}$.

Proofs of Theorem 1, Corollary 1, and Theorem 2. In section 5.1, we have obtained a linear optical proof of Glynn's formula for the permanent using the fact that a cat state $|\text{cat}_\alpha\rangle$ of small amplitude α approximates a single-photon Fock state $|1\rangle$. Hereafter, we consider another approximation of Fock states using superpositions of two-mode squeezed states. To prove the identity in Eq. (65), we introduce $|\lambda^-\rangle := \frac{1}{2\lambda} (|\lambda\rangle - |-\lambda\rangle)$, for $|\lambda| < 1$, where $|\lambda\rangle = \sum_{p \geq 0} \lambda^p |pp\rangle$ is an unnormalized two-mode squeezed state. We have $\lim_{\lambda \rightarrow 0} |\lambda^-\rangle = |11\rangle$ in trace distance. Hence, for U a $(2m) \times (2m)$ unitary matrix we have with Eq. (9):

$$\text{Per}(U) = \lim_{\lambda \rightarrow 0} \langle \lambda^- | \hat{U} | \lambda^- \rangle^{\otimes m}. \quad (71)$$

For all $\lambda \in \mathbb{R}$, with $|\lambda| < 1$, let us compute

$$\otimes^m \langle \lambda^- | \hat{U} | \lambda^- \rangle^{\otimes m} = \frac{1}{4^m \lambda^{2m}} \sum_{\mathbf{x}, \mathbf{y} \in \{-1, 1\}^m} x_1 \dots x_m y_1 \dots y_m g_U(\lambda \mathbf{x}, \lambda \mathbf{y}), \quad (72)$$

where we have defined for all $\boldsymbol{\lambda}, \boldsymbol{\mu} \in \mathbb{C}^m$ (note that we associate mode 1 with mode $m+1$, mode 2 with mode $m+2$, and so on)

$$g_U(\boldsymbol{\lambda}, \boldsymbol{\mu}) := \langle \boldsymbol{\lambda}^* | \hat{U} | \boldsymbol{\mu} \rangle. \quad (73)$$

With Eq. (13), we have

$$g_U(\boldsymbol{\lambda}, \boldsymbol{\mu}) = \frac{1}{\sqrt{\det(V_U(\boldsymbol{\lambda}, \boldsymbol{\mu}))}}, \quad (74)$$

where

$$V_U(\boldsymbol{\lambda}, \boldsymbol{\mu}) = \begin{pmatrix} -UV_\mu U^T & I_{2m} \\ I_{2m} & -V_\lambda \end{pmatrix}, \quad (75)$$

where for all $\mathbf{w} \in \mathbb{C}^m$,

$$V_{\mathbf{w}} = \begin{pmatrix} 0_m & \text{Diag}(\mathbf{w}) \\ \text{Diag}(\mathbf{w}) & 0_m \end{pmatrix}. \quad (76)$$

Moreover,

$$\text{Det}(V_U(\boldsymbol{\lambda}, \boldsymbol{\mu})) = \text{Det}(I - V_\lambda UV_\mu U^T), \quad (77)$$

so with Eq. (74)

$$g_U(\boldsymbol{\lambda}, \boldsymbol{\mu}) = \frac{1}{\sqrt{\text{Det}(I - V_\lambda UV_\mu U^T)}}. \quad (78)$$

Setting $\boldsymbol{\lambda} = \boldsymbol{\mu} = \lambda \mathbf{1}$ for $\lambda \in \mathbb{R}$ and letting λ go to 0 in Eq. (72), we obtain that $\text{Per}(U)$ is given by

$$\frac{1}{4^m} \sum_{\mathbf{x}, \mathbf{y} \in \{-1, 1\}^m} x_1 \dots x_m y_1 \dots y_m [\lambda^{2m}] g_U(\lambda \mathbf{x}, \lambda \mathbf{y}), \quad (79)$$

where $[\lambda^{2m}] g_U(\lambda \mathbf{x}, \lambda \mathbf{y})$ is the λ^{2m} coefficient in the Taylor expansion (in λ) of

$$g_U(\lambda \mathbf{x}, \lambda \mathbf{y}) = \frac{1}{\sqrt{\text{Det}(I - \lambda^2 V_{\mathbf{x}} UV_{\mathbf{y}} U^T)}}. \quad (80)$$

This concludes the proof of Eq. (65). Once again, the relation for a generic matrix A is obtained by embedding it as a submatrix of a unitary matrix U .

Now with Eq. (73), for all $\boldsymbol{\lambda}, \boldsymbol{\mu} \in \mathbb{C}^n$,

$$\begin{aligned} g_U(\boldsymbol{\lambda}, \boldsymbol{\mu}) &= \sum_{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m} \lambda^{\mathbf{p}} \mu^{\mathbf{q}} \langle \mathbf{p} \mathbf{p} | \hat{U} | \mathbf{q} \mathbf{q} \rangle \\ &= \sum_{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m} \frac{\lambda^{\mathbf{p}} \mu^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(U_{\mathbf{p} \oplus \mathbf{p}, \mathbf{q} \oplus \mathbf{q}}), \end{aligned} \quad (81)$$

where we used Eq. (9) in the second line. Combining Eqs. (73), (78) and (81) we obtain

$$\sum_{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m} \frac{\lambda^{\mathbf{p}} \mu^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(U_{\mathbf{p} \oplus \mathbf{p}, \mathbf{q} \oplus \mathbf{q}}) = \langle \boldsymbol{\lambda}^* | \hat{U} | \boldsymbol{\mu} \rangle = \frac{1}{\sqrt{\text{Det}(I - V_\lambda UV_\mu U^T)}}, \quad (82)$$

where for all $\mathbf{w} \in \mathbb{C}^n$,

$$V_{\mathbf{w}} = \begin{pmatrix} 0_m & \text{Diag}(\mathbf{w}) \\ \text{Diag}(\mathbf{w}) & 0_m \end{pmatrix}. \quad (83)$$

This concludes the proof of Eq. (66). Once again, the relation for a generic matrix A is obtained by embedding it as a submatrix of a unitary matrix U .

When $U = A \oplus B$, with A and B two $m \times m$ matrices, we have $U_{\mathbf{p} \oplus \mathbf{p}, \mathbf{q} \oplus \mathbf{q}} = A_{\mathbf{p}, \mathbf{q}} \oplus B_{\mathbf{p}, \mathbf{q}}$ and the permanent of a block-diagonal matrix is the product of the permanents of the blocks, so Eq. (82) gives

$$\sum_{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m} \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}) \text{Per}(B_{\mathbf{p}, \mathbf{q}}) = \frac{1}{\sqrt{\text{Det}(I - V_{\mathbf{x}}(A \oplus B)V_{\mathbf{y}}(A^T \oplus B^T))}}. \quad (84)$$

Writing $X = \text{Diag}(x_1, \dots, x_m)$ and $Y = \text{Diag}(y_1, \dots, y_m)$, we have

$$\begin{aligned} \text{Det}(I - V_{\mathbf{x}}(A \oplus B)V_{\mathbf{y}}(A^T \oplus B^T)) &= \text{Det}\left(I - \begin{pmatrix} 0_m & X \\ X & 0_m \end{pmatrix} \begin{pmatrix} A & 0_m \\ 0_m & B \end{pmatrix} \begin{pmatrix} 0_m & Y \\ Y & 0_m \end{pmatrix} \begin{pmatrix} A^T & 0_m \\ 0_m & B^T \end{pmatrix}\right) \\ &= \text{Det}\begin{pmatrix} I - XBYA^T & 0_m \\ 0_m & I - XAYB^T \end{pmatrix} \\ &= \text{Det}(I - XBYA^T) \text{Det}(I - XAYB^T) \\ &= \text{Det}(I - XAYB^T)^2, \end{aligned} \quad (85)$$

where the last line is obtained by assuming that A is invertible (wlog by density) and taking the transpose in the first determinant. Combining Eqs. (84) and (85) concludes the proof of Theorem 1.

This theorem reduces to the MacMahon master theorem when $A = I$ or $B = I$, since $\text{Per}(I_{\mathbf{p}, \mathbf{q}}) = \mathbf{p}! \delta_{\mathbf{p}, \mathbf{q}}$. Another particular case of interest is when $B = J_m$, where J_m is the all-1 matrix of size $m \times m$, which satisfies $\text{Per}(J_m) = m!$. In this case, we obtain, for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^m$,

$$\begin{aligned} \sum_{\substack{n \in \mathbb{N}, \mathbf{p}, \mathbf{q} \in \mathbb{N}^m \\ |\mathbf{p}| = |\mathbf{q}| = n}} \frac{n! \mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}) &= \frac{1}{\text{Det}(I - AYJ_mX)} \\ &= \frac{1}{\text{Det}(I - A\mathbf{y}\mathbf{x}^T)} \\ &= \frac{1}{1 - \mathbf{x}^T A\mathbf{y}}, \end{aligned} \quad (86)$$

where we used the fact that $A\mathbf{y}\mathbf{x}^T$ is a rank-one matrix to compute the determinant. This implies Corollary 1: for all $n \in \mathbb{N}^*$ and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ with $|\mathbf{p}| = |\mathbf{q}| = n$,

$$\begin{aligned} \text{Per}(A_{\mathbf{p}, \mathbf{q}}) &= \frac{\mathbf{p}! \mathbf{q}!}{n!} [\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}] \left(\frac{1}{1 - \mathbf{x}^T A\mathbf{y}} \right) \\ &= \frac{\mathbf{p}! \mathbf{q}!}{n!} [\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}] (\mathbf{x}^T A\mathbf{y})^n, \end{aligned} \quad (87)$$

where we used the Taylor series $\frac{1}{1-z} = \sum_{k=0}^{+\infty} z^k$.

We note that Theorem 1—which is a generalization of the MacMahon master theorem to two matrices—may also be obtained by combining the MacMahon master theorem in Eq. (55) with the Cauchy–Binet theorem in Eq. (47), applied to the matrix AYB^T . As it turns out, we may apply the same proof technique inductively in order to generalize

the MacMahon master theorem to N matrices $A^{(1)}, \dots, A^{(N)}$, for $N \geq 1$ and obtain Theorem 2: assuming that Eq. (70) holds for some $N \geq 2$, we have, for all $m \times m$ matrices $B^{(1)}, \dots, B^{(N)}$,

$$\begin{aligned} & \sum_{\mathbf{p}_1, \dots, \mathbf{p}_N \in \mathbb{N}^m} \prod_{k=1}^N \frac{z_k^{\mathbf{p}_k}}{\mathbf{p}_k!} \text{Per}(B_{\mathbf{p}_1, \mathbf{p}_2}^{(1)}) \text{Per}(B_{\mathbf{p}_2, \mathbf{p}_3}^{(2)}) \dots \text{Per}(B_{\mathbf{p}_N, \mathbf{p}_1}^{(N)}) \\ &= \frac{1}{\text{Det}(I - Z_1 B^{(1)} \dots Z_N B^{(N)})}, \end{aligned} \quad (88)$$

where $\mathbf{z}_k = (z_{k1}, \dots, z_{km}) \in \mathbb{C}^m$ and $Z_k = \text{Diag}(\mathbf{z}_k)$ for all $k \in \{1, \dots, N\}$.

Let $A^{(1)}, \dots, A^{(N+1)}$ be $m \times m$ matrices, $\mathbf{z}_{N+1} = (z_{N+1,1}, \dots, z_{N+1,m}) \in \mathbb{C}^m$ and $Z_{N+1} = \text{Diag}(\mathbf{z}_{N+1})$. Setting $B^{(N)} = A^{(N)} Z_N A^{(N+1)}$ and $B^{(k)} = A^{(k)}$ for all $k \in \{1, \dots, N\}$, we obtain with Eq. (88):

$$\begin{aligned} & \frac{1}{\text{Det}(I - Z_1 A^{(1)} \dots Z_N A^{(N)} Z_{N+1} A^{(N+1)})} \\ &= \sum_{\mathbf{p}_1, \dots, \mathbf{p}_N \in \mathbb{N}^m} \prod_{k=1}^N \frac{z_k^{\mathbf{p}_k}}{\mathbf{p}_k!} \text{Per}(A_{\mathbf{p}_1, \mathbf{p}_2}^{(1)}) \dots \text{Per}(A_{\mathbf{p}_{N-1}, \mathbf{p}_N}^{(N-1)}) \text{Per}((A^{(N)} Z_N A^{(N+1)})_{\mathbf{p}_N, \mathbf{p}_1}). \end{aligned} \quad (89)$$

On the other hand, with the Cauchy–Binet theorem from Eq. (47),

$$\begin{aligned} \text{Per}((A^{(N)} Z_{N+1} A^{(N+1)})_{\mathbf{p}_N, \mathbf{p}_1}) &= \sum_{\mathbf{p}_{N+1} \in \mathbb{N}^m} \frac{1}{\mathbf{p}_{N+1}!} \text{Per}(A_{\mathbf{p}_N, \mathbf{p}_{N+1}}^{(N)}) \text{Per}((Z_{N+1} A^{(N+1)})_{\mathbf{p}_{N+1}, \mathbf{p}_1}) \\ &= \sum_{\mathbf{p}_{N+1} \in \mathbb{N}^m} \frac{z_{N+1}^{\mathbf{p}_{N+1}}}{\mathbf{p}_{N+1}!} \text{Per}(A_{\mathbf{p}_N, \mathbf{p}_{N+1}}^{(N)}) \text{Per}(A_{\mathbf{p}_{N+1}, \mathbf{p}_1}^{(N+1)}), \end{aligned} \quad (90)$$

where we used the fact that multiplying any single row of M by a scalar z changes $\text{Per}(M)$ to $z \text{Per}(M)$. Combining Eqs. (89) and (90) completes the induction step and the proof of Theorem 2. \square

Applications. Recall that the MacMahon master theorem is a useful tool for deriving combinatorial identities (see section 2.3): it expresses the permanent of an $m \times m$ matrix A with rows and columns repeated in the same way as the coefficient

$$\text{Per}(A_{\mathbf{p}, \mathbf{p}}) = \mathbf{p}! [z^{\mathbf{p}}] \left(\frac{1}{\text{Det}(I - ZA)} \right), \quad (91)$$

where $Z = \text{Diag}(\mathbf{z})$ and $\mathbf{p} \in \mathbb{N}^m$, while the same permanent may also be expressed as the coefficient

$$\text{Per}(A_{\mathbf{p}, \mathbf{p}}) = \mathbf{p}! [z^{\mathbf{p}}] (AZ)^{\mathbf{p}}. \quad (92)$$

Our generalizations of the MacMahon master theorem may be used in a similar fashion to obtain simple proofs of combinatorial identities. Let us illustrate this with an example: for $n \in \mathbb{N}$ and $a, b \in \mathbb{C}$, setting

$$S_n(a, b) := \sum_{k=0}^n \binom{n}{k}^2 a^k b^{n-k}, \quad (93)$$

we aim to prove the relation

$$S_n(a, b)^2 = \sum_{l=0}^n \binom{2l}{l} \binom{n+l}{2l} (-1)^{n-l} (a-b)^{2n-2l} S_l(a^2, b^2), \quad (94)$$

which for $a = b = 1$ directly implies the well-known $\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n}$.

Proof. By Theorem 1 we have

$$\text{Per}(A_{\mathbf{p},\mathbf{q}})\text{Per}(B_{\mathbf{p},\mathbf{q}}) = \mathbf{p}!\mathbf{q}![\mathbf{x}^{\mathbf{p}}\mathbf{y}^{\mathbf{q}}] \left(\frac{1}{\text{Det}(I - XAYB^T)} \right), \quad (95)$$

where $X = \text{Diag}(x_1, \dots, x_m)$ and $Y = \text{Diag}(y_1, \dots, y_m)$. Setting

$$A = B = \begin{pmatrix} 1 & a \\ 1 & b \end{pmatrix}, \quad (96)$$

for $a, b \in \mathbb{C}$, $n \in \mathbb{N}^*$, and $\mathbf{p} = \mathbf{q} = (n, n)$, we obtain with Eq. (92):

$$\begin{aligned} \text{Per}(A_{\mathbf{p},\mathbf{q}}) &= (n!)^2 [z_1^n z_2^n] (z_1 + az_2)^n (z_1 + bz_2)^n \\ &= (n!)^2 \sum_{k+l=n} \binom{n}{k} \binom{n}{l} a^k b^l \\ &= (n!)^2 S_n(a, b). \end{aligned} \quad (97)$$

On the other hand, with Eq. (95) we have

$$\begin{aligned} \text{Per}(A_{\mathbf{p},\mathbf{q}})^2 &= (n!)^4 [x_1^n x_2^n y_1^n y_2^n] \left(\frac{1}{\text{Det}(I - \text{Diag}(x_1, x_2)A\text{Diag}(y_1, y_2)A^T)} \right) \\ &= (n!)^4 [x_1^n x_2^n y_1^n y_2^n] \left(\frac{1}{1 - x_1 y_1 - a^2 x_1 y_2 - x_2 y_1 - b^2 x_2 y_2 + (a-b)^2 x_1 x_2 y_1 y_2} \right) \\ &= (n!)^4 \sum_{k=n}^{2n} [x_1^n x_2^n y_1^n y_2^n] \left(x_1 y_1 + a^2 x_1 y_2 + x_2 y_1 + b^2 x_2 y_2 - (a-b)^2 x_1 x_2 y_1 y_2 \right)^k \\ &= (n!)^4 \sum_{k=n}^{2n} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} (a-b)^{2k-2j} \sum_{n_{11}+n_{12}+n_{21}+n_{22}=j} \frac{j! a^{2n_{12}} b^{2n_{22}}}{n_{11}! n_{12}! n_{21}! n_{22}!} \\ &\quad \times [x_1^n x_2^n y_1^n y_2^n] \left((x_1 x_2 y_1 y_2)^{k-j} x_1^{n_{11}+n_{12}} x_2^{n_{21}+n_{22}} y_1^{n_{11}+n_{21}} y_2^{n_{12}+n_{22}} \right), \end{aligned} \quad (98)$$

where we used the Taylor expansion of $\frac{1}{1-z}$ in the third line and the multinomial theorem in the last line. The indices in the above expression must satisfy $n_{11} = n_{22}$, $n_{12} = n_{21}$, which implies $j = 2n_{11} + 2n_{12}$, i.e. j is even. Moreover, $n = k - j + n_{11} + n_{12} = k - j/2$. Relabeling $j = 2l$ and $n_{11} = p$ we have $k = l + n$ and we obtain

$$\begin{aligned} \text{Per}(A_{\mathbf{p},\mathbf{q}})^2 &= (n!)^4 \sum_{l=0}^n \binom{n+l}{2l} (-1)^{n-l} (a-b)^{2n-2l} \sum_{p=0}^l \frac{(2l)! a^{2l-2p} b^{2p}}{p!^2 (l-p)!^2} \\ &= (n!)^4 \sum_{l=0}^n \binom{2l}{l} \binom{n+l}{2l} (-1)^{n-l} (a-b)^{2n-2l} S_l(a^2, b^2). \end{aligned} \quad (99)$$

With Eq. (97) this proves

$$S_n(a, b)^2 = \sum_{l=0}^n \binom{2l}{l} \binom{n+l}{2l} (-1)^{n-l} (a-b)^{2n-2l} S_l(a^2, b^2). \quad (100)$$

□

6.2 New generating functions

We have encountered several generating functions for the permanent of an $m \times m$ matrix A with differently repeated rows and columns. Jackson's formula gives [41]:

$$e^{\mathbf{x}^T A \mathbf{y}} = \sum_{\substack{n \in \mathbb{N}, \mathbf{p}, \mathbf{q} \in \mathbb{N}^m \\ |\mathbf{p}| = |\mathbf{q}| = n}} \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}), \quad (101)$$

for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^m$. Moreover, we have obtained in Eq. (86), as a corollary of our generalization of the MacMahon master theorem:

$$\frac{1}{1 - \mathbf{x}^T A \mathbf{y}} = \sum_{\substack{n \in \mathbb{N}, \mathbf{p}, \mathbf{q} \in \mathbb{N}^m \\ |\mathbf{p}| = |\mathbf{q}| = n}} n! \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}). \quad (102)$$

Furthermore, Corollary 1 gives:

$$(\mathbf{x}^T A \mathbf{y})^n = \sum_{\substack{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m \\ |\mathbf{p}| = |\mathbf{q}| = n}} n! \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}). \quad (103)$$

In this section we prove Theorem 3, which generalizes all three statements: for any series $f(z) = \sum_{n=0}^{+\infty} f_n z^n$,

$$f(\mathbf{x}^T A \mathbf{y}) = \sum_{\substack{n \in \mathbb{N}, \mathbf{p}, \mathbf{q} \in \mathbb{N}^m \\ |\mathbf{p}| = |\mathbf{q}| = n}} f_n n! \frac{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(A_{\mathbf{p}, \mathbf{q}}). \quad (104)$$

One may prove this statement by linearity, using Eq. (103). In what follows, we give a direct quantum-inspired proof which is similar to that of Jackson's formula from section 5.4.

Proof of Theorem 3. For $\boldsymbol{\alpha}, \boldsymbol{\beta} \in \mathbb{C}^m$ and a unitary matrix U ,

$$\begin{aligned} \sum_{\substack{n \in \mathbb{N}, \mathbf{p}, \mathbf{q} \in \mathbb{N}^m \\ |\mathbf{p}| = |\mathbf{q}| = n}} f_n n! \frac{\boldsymbol{\alpha}^{\mathbf{p}} \boldsymbol{\beta}^{\mathbf{q}}}{\mathbf{p}! \mathbf{q}!} \text{Per}(U_{\mathbf{p}, \mathbf{q}}) &= e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2 + \frac{1}{2} \|\boldsymbol{\beta}\|^2} \sum_{n=0}^{+\infty} f_n n! \sum_{\substack{\mathbf{p}, \mathbf{q} \in \mathbb{N}^m \\ |\mathbf{p}| = |\mathbf{q}| = n}} \langle \boldsymbol{\alpha}^* | \mathbf{p} \rangle \langle \mathbf{p} | \hat{U} | \mathbf{q} \rangle \langle \mathbf{q} | \boldsymbol{\beta} \rangle \\ &= e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2 + \frac{1}{2} \|\boldsymbol{\beta}\|^2} \sum_{n=0}^{+\infty} f_n n! \langle \boldsymbol{\alpha}^* | \Pi_n \hat{U} \Pi_n | \boldsymbol{\beta} \rangle \\ &= e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2 + \frac{1}{2} \|\boldsymbol{\beta}\|^2} \sum_{n=0}^{+\infty} f_n n! \langle \boldsymbol{\alpha}^* | \Pi_n \hat{U} | \boldsymbol{\beta} \rangle \\ &= e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2 + \frac{1}{2} \|\boldsymbol{\beta}\|^2} \sum_{n=0}^{+\infty} f_n n! \langle \boldsymbol{\alpha}^* | \Pi_n | U \boldsymbol{\beta} \rangle, \end{aligned} \quad (105)$$

where we used Eqs. (6) and (9) in the first line, the definition of the projector $\Pi_n = \sum_{|\mathbf{p}|=n} |\mathbf{p}\rangle \langle \mathbf{p}|$ in the second line, Eq. (11) in the third line, and Eq. (12) in the fourth line. Now,

$$\begin{aligned} e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2 + \frac{1}{2} \|\boldsymbol{\beta}\|^2} \langle \boldsymbol{\alpha}^* | \Pi_n | U \boldsymbol{\beta} \rangle &= e^{\frac{1}{2} \|\boldsymbol{\alpha}\|^2 + \frac{1}{2} \|\boldsymbol{\beta}\|^2} \sum_{|\mathbf{p}|=n} \langle \boldsymbol{\alpha}^* | \mathbf{p} \rangle \langle \mathbf{p} | U \boldsymbol{\beta} \rangle \\ &= \sum_{|\mathbf{p}|=n} e^{\frac{1}{2} \|\boldsymbol{\beta}\|^2 - \frac{1}{2} \|U \boldsymbol{\beta}\|^2} \frac{\boldsymbol{\alpha}^{\mathbf{p}} (U \boldsymbol{\beta})^{\mathbf{p}}}{\mathbf{p}!} \\ &= \sum_{|\mathbf{p}|=n} \frac{\boldsymbol{\alpha}^{\mathbf{p}} (U \boldsymbol{\beta})^{\mathbf{p}}}{\mathbf{p}!} \\ &= \frac{1}{n!} (\boldsymbol{\alpha}^T U \boldsymbol{\beta})^n, \end{aligned} \quad (106)$$

where we used Eq. (6) in the second line, the fact that U is unitary in the third line, and the multinomial theorem in the last line. Combining Eqs. (105) and (106) completes the proof for unitary matrices. Once again, the relation for a generic matrix A is obtained by embedding it as a submatrix of a unitary matrix U .

As a result, when $f_n \neq 0$ we have, for any $m \times m$ matrix A , all $n \in \mathbb{N}$ and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ such that $|\mathbf{p}| = |\mathbf{q}| = n$,

$$\begin{aligned} \mathbb{E}_{\mathbf{x}, \mathbf{y} \in \mathbb{T}^m} \left[\frac{\mathbf{p}! \mathbf{q}!}{\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}} \frac{f(\mathbf{x}^T A \mathbf{y})}{\partial_z^n f(z) \Big|_{z=0}} \right] &= \frac{\mathbf{p}! \mathbf{q}!}{\partial_z^n f(z) \Big|_{z=0}} \sum_{\substack{n \in \mathbb{N}, \mathbf{s}, \mathbf{t} \in \mathbb{N}^m \\ |\mathbf{s}| = |\mathbf{t}| = k}} f_k k! \frac{\text{Per}(A_{\mathbf{s}, \mathbf{t}})}{\mathbf{s}! \mathbf{t}!} \\ &\quad \times \prod_{j=1}^m \left(\int_{\theta_j, \varphi_j \in [0, 2\pi]} e^{i\theta_j (s_j - p_j)} e^{i\varphi_j (t_j - q_j)} \frac{d\theta_j d\varphi_j}{4\pi^2} \right) \quad (107) \\ &= \frac{1}{\partial_z^n f(z) \Big|_{z=0}} f_n n! \text{Per}(A_{\mathbf{p}, \mathbf{q}}) \\ &= \text{Per}(A_{\mathbf{p}, \mathbf{q}}), \end{aligned}$$

where we have set $\mathbf{x} = (e^{i\theta_1}, \dots, e^{i\theta_m}) \in \mathbb{T}^m$ and $\mathbf{y} = (e^{i\varphi_1}, \dots, e^{i\varphi_m}) \in \mathbb{T}^m$ in the first line, and where we have used $\int_0^{2\pi} e^{i(k-l)\theta} \frac{d\theta}{2\pi} = \delta_{kl}$ in the third line. \square

Applications. These generating functions may be used to derive remarkable identities for the permanent: for instance, for A and B two $m \times m$ matrices, taking $f(z) = e^z$ and equating the $\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}$ coefficients of $e^{\mathbf{x}^T (A+B) \mathbf{y}}$ and $e^{\mathbf{x}^T A \mathbf{y}} e^{\mathbf{x}^T B \mathbf{y}}$ yields the sum formula [1, 2]:

$$\text{Per}((A+B)_{\mathbf{p}, \mathbf{q}}) = \sum_{\substack{\mathbf{s} + \mathbf{t} = \mathbf{p} \\ \mathbf{u} + \mathbf{v} = \mathbf{q}}} \frac{\mathbf{p}! \mathbf{q}!}{\mathbf{s}! \mathbf{t}! \mathbf{u}! \mathbf{v}!} \text{Per}(A_{\mathbf{s}, \mathbf{u}}) \text{Per}(B_{\mathbf{t}, \mathbf{v}}), \quad (108)$$

for all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$.

Similarly, for A an $m \times m$ matrix, taking $f(z) = z^p$ for $p = k, l, k+l$ and equating the $\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}$ coefficients of $(\mathbf{x}^T A \mathbf{y})^{k+l}$ and $(\mathbf{x}^T A \mathbf{y})^k (\mathbf{x}^T A \mathbf{y})^l$ yields the Laplace expansion formula [1, 2]:

$$\text{Per}(A_{\mathbf{p}, \mathbf{q}}) = \frac{k! l!}{(k+l)!} \sum_{\substack{\mathbf{s} + \mathbf{t} = \mathbf{p} \\ \mathbf{u} + \mathbf{v} = \mathbf{q} \\ |\mathbf{s}| = |\mathbf{u}| = k \\ |\mathbf{t}| = |\mathbf{v}| = l}} \frac{\mathbf{p}! \mathbf{q}!}{\mathbf{s}! \mathbf{t}! \mathbf{u}! \mathbf{v}!} \text{Per}(A_{\mathbf{s}, \mathbf{u}}) \text{Per}(A_{\mathbf{t}, \mathbf{v}}), \quad (109)$$

for all $k, l \in \mathbb{N}$ and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}^m$ with $|\mathbf{p}| = |\mathbf{q}| = k+l$.

Finally, for A and B two $m \times m$ matrices, taking $f(z) = -\log(1-z)$ and equating the $\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}$ coefficients of $-\log(1 - \mathbf{x}^T A \mathbf{y}) - \log(1 - \mathbf{x}^T B \mathbf{y})$ and $-\log[(1 - \mathbf{x}^T B \mathbf{y})(1 - \mathbf{x}^T A \mathbf{y})]$ yields Theorem 4 after a derivation which we detail below.

Proof of Theorem 4. Let us write

$$-\log[(1 - \mathbf{x}^T B \mathbf{y})(1 - \mathbf{x}^T A \mathbf{y})] = -\log[1 - \mathbf{x}^T (A + B - B \mathbf{y} \mathbf{x}^T A) \mathbf{y}]. \quad (110)$$

Applying Theorem 3 for $f(z) = -\log(1-z)$ with $z = \mathbf{x}^T B \mathbf{y}, \mathbf{x}^T A \mathbf{y}, \mathbf{x}^T (A+B-B \mathbf{y} \mathbf{x}^T A) \mathbf{y}$ and considering the $\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}$ coefficient we obtain for all $n \in \mathbb{N}^*$ and all $\mathbf{p}, \mathbf{q} \in \mathbb{N}$ with

$$|\mathbf{p}| = |\mathbf{q}| = n:$$

$$\begin{aligned}
& \text{Per}(A_{\mathbf{p},\mathbf{q}}) + \text{Per}(B_{\mathbf{p},\mathbf{q}}) \\
&= \frac{\mathbf{p}! \mathbf{q}!}{(n-1)!} [\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}] \left(\sum_{\substack{l \in \mathbb{N}^*, \mathbf{s}, \mathbf{t} \in \mathbb{N}^m \\ |\mathbf{s}| = |\mathbf{t}| = l}} (l-1)! \frac{\mathbf{x}^{\mathbf{s}} \mathbf{y}^{\mathbf{t}}}{\mathbf{s}! \mathbf{t}!} \text{Per}((A+B - B\mathbf{y}\mathbf{x}^T A)_{\mathbf{s},\mathbf{t}}) \right) \\
&= \sum_{\substack{l \in \mathbb{N}^*, \mathbf{s}, \mathbf{t} \in \mathbb{N}^m \\ |\mathbf{s}| = |\mathbf{t}| = l}} \frac{(l-1)! \mathbf{p}! \mathbf{q}!}{(n-1)! \mathbf{s}! \mathbf{t}!} [\mathbf{x}^{\mathbf{p}} \mathbf{y}^{\mathbf{q}}] \left(\mathbf{x}^{\mathbf{s}} \mathbf{y}^{\mathbf{t}} \text{Per}((A+B - B\mathbf{y}\mathbf{x}^T A)_{\mathbf{s},\mathbf{t}}) \right) \\
&= \sum_{\substack{l \in \mathbb{N}^*, \mathbf{s}, \mathbf{t} \in \mathbb{N}^m \\ |\mathbf{s}| = |\mathbf{t}| = l \\ \mathbf{s} \leq \mathbf{p}, \mathbf{t} \leq \mathbf{q}}} \frac{(l-1)! \mathbf{p}! \mathbf{q}!}{(n-1)! \mathbf{s}! \mathbf{t}!} [\mathbf{x}^{\mathbf{p}-\mathbf{s}} \mathbf{y}^{\mathbf{q}-\mathbf{t}}] \left(\text{Per}((A+B - B\mathbf{y}\mathbf{x}^T A)_{\mathbf{s},\mathbf{t}}) \right) \\
&= \sum_{\substack{l \in \mathbb{N}^*, \mathbf{s}, \mathbf{t}, \mathbf{u}, \mathbf{v} \in \mathbb{N}^m \\ |\mathbf{s}| = |\mathbf{t}| = l \\ \mathbf{s} + \mathbf{u} = \mathbf{p} \\ \mathbf{t} + \mathbf{v} = \mathbf{q}}} \frac{(l-1)! \mathbf{p}! \mathbf{q}!}{(n-1)! \mathbf{s}! \mathbf{t}!} [\mathbf{x}^{\mathbf{u}} \mathbf{y}^{\mathbf{v}}] \left(\text{Per}((A+B - B\mathbf{y}\mathbf{x}^T A)_{\mathbf{s},\mathbf{t}}) \right).
\end{aligned} \tag{111}$$

Applying the sum formula from Eq. (108) to the matrices $(A+B)$ and $(-B\mathbf{y}\mathbf{x}^T A)$ we obtain, for all $l \in \mathbb{N}^*$ and all $\mathbf{s}, \mathbf{t} \in \mathbb{N}$ with $|\mathbf{s}| = |\mathbf{t}| = l$,

$$\begin{aligned}
\text{Per}((A+B - B\mathbf{y}\mathbf{x}^T A)_{\mathbf{s},\mathbf{t}}) &= \sum_{\substack{\mathbf{c} + \mathbf{k} = \mathbf{s} \\ \mathbf{c}' + \mathbf{k}' = \mathbf{t} \\ |\mathbf{c}| = |\mathbf{c}'|, |\mathbf{k}| = |\mathbf{k}'|}} \frac{\mathbf{s}! \mathbf{t}!}{\mathbf{c}! \mathbf{k}! \mathbf{c}'! \mathbf{k}'!} \text{Per}((A+B)_{\mathbf{c},\mathbf{c}'}) \text{Per}((-B\mathbf{y}\mathbf{x}^T A)_{\mathbf{k},\mathbf{k}'}) \\
&= \sum_{\substack{\mathbf{c} + \mathbf{k} = \mathbf{s} \\ \mathbf{c}' + \mathbf{k}' = \mathbf{t} \\ |\mathbf{c}| = |\mathbf{c}'|, |\mathbf{k}| = |\mathbf{k}'|}} \frac{(-1)^{|\mathbf{k}|} \mathbf{s}! \mathbf{t}!}{\mathbf{c}! \mathbf{k}! \mathbf{c}'! \mathbf{k}'!} \text{Per}((A+B)_{\mathbf{c},\mathbf{c}'}) \text{Per}((B\mathbf{y}\mathbf{x}^T A)_{\mathbf{k},\mathbf{k}'}).
\end{aligned} \tag{112}$$

The permanent of the outer product $\mathbf{v}\mathbf{w}^T$ of two vectors \mathbf{v} and \mathbf{w} of the same size k is easily computed as

$$\begin{aligned}
\text{Per}(\mathbf{v}\mathbf{w}^T) &= \sum_{\sigma \in \mathcal{S}_k} \prod_{j=1}^k v_j w_{\sigma(j)} \\
&= \sum_{\sigma \in \mathcal{S}_k} v_1 \dots v_k w_1 \dots w_k \\
&= k! v_1 \dots v_k w_1 \dots w_k.
\end{aligned} \tag{113}$$

The matrix $(B\mathbf{y}\mathbf{x}^T A)_{\mathbf{k},\mathbf{k}'}$ is the outer product between the vector $(B\mathbf{y})_{\mathbf{k}}$ (obtained from the vector $B\mathbf{y}$ by repeating k_i times its i^{th} entry) and the vector $(A^T \mathbf{x})_{\mathbf{k}'}$ (obtained from the vector $A^T \mathbf{x}$ by repeating k'_i times its i^{th} entry) of size $|\mathbf{k}|$, so its permanent is given by

$$\text{Per}((B\mathbf{y}\mathbf{x}^T A)_{\mathbf{k},\mathbf{k}'}) = (|\mathbf{k}|)! (B\mathbf{y})^{\mathbf{k}} (A^T \mathbf{x})^{\mathbf{k}'}. \tag{114}$$

With Eq. (112) we obtain

$$\text{Per}((A+B - B\mathbf{y}\mathbf{x}^T A)_{\mathbf{s},\mathbf{t}}) = \sum_{\substack{\mathbf{c} + \mathbf{k} = \mathbf{s} \\ \mathbf{c}' + \mathbf{k}' = \mathbf{t} \\ |\mathbf{c}| = |\mathbf{c}'|, |\mathbf{k}| = |\mathbf{k}'|}} \frac{(-1)^{|\mathbf{k}|} (|\mathbf{k}|)! \mathbf{s}! \mathbf{t}!}{\mathbf{c}! \mathbf{k}! \mathbf{c}'! \mathbf{k}'!} \text{Per}((A+B)_{\mathbf{c},\mathbf{c}'}) (B\mathbf{y})^{\mathbf{k}} (A^T \mathbf{x})^{\mathbf{k}'}. \tag{115}$$

Plugging this expression into Eq. (111) yields

$$\begin{aligned}
& \text{Per}(A_{\mathbf{p},\mathbf{q}}) + \text{Per}(B_{\mathbf{p},\mathbf{q}}) \\
&= \sum_{\substack{l \in \mathbb{N}^*, \mathbf{s}, \mathbf{t}, \mathbf{u}, \mathbf{v} \in \mathbb{N}^m \\ |\mathbf{s}|=|\mathbf{t}|=l \\ \mathbf{s}+\mathbf{u}=\mathbf{p} \\ \mathbf{t}+\mathbf{v}=\mathbf{q}}} \frac{(l-1)! \mathbf{p}! \mathbf{q}!}{(n-1)! \mathbf{s}! \mathbf{t}!} \sum_{\substack{\mathbf{c}+\mathbf{k}=\mathbf{s} \\ \mathbf{c}'+\mathbf{k}'=\mathbf{t} \\ |\mathbf{c}|=|\mathbf{c}'|, |\mathbf{k}|=|\mathbf{k}'|}} \frac{(-1)^{|\mathbf{k}|} (|\mathbf{k}|)! \mathbf{s}! \mathbf{t}!}{\mathbf{c}! \mathbf{k}! \mathbf{c}'! \mathbf{k}'!} \text{Per}((A+B)_{\mathbf{c},\mathbf{c}'})[\mathbf{x}^{\mathbf{u}} \mathbf{y}^{\mathbf{v}}](B\mathbf{y})^{\mathbf{k}} (A^T \mathbf{x})^{\mathbf{k}'} \\
&= \sum_{\substack{l \in \mathbb{N}^*, \mathbf{s}, \mathbf{t}, \mathbf{u}, \mathbf{v} \in \mathbb{N}^m \\ |\mathbf{s}|=|\mathbf{t}|=l \\ \mathbf{s}+\mathbf{u}=\mathbf{p} \\ \mathbf{t}+\mathbf{v}=\mathbf{q}}} \frac{(l-1)! \mathbf{p}! \mathbf{q}!}{(n-1)!} \sum_{\substack{\mathbf{c}+\mathbf{k}=\mathbf{s} \\ \mathbf{c}'+\mathbf{k}'=\mathbf{t} \\ |\mathbf{c}|=|\mathbf{c}'|, |\mathbf{k}|=|\mathbf{k}'|}} \frac{(-1)^{|\mathbf{k}|} (|\mathbf{k}|)!}{\mathbf{c}! \mathbf{k}! \mathbf{c}'! \mathbf{k}'!} \text{Per}((A+B)_{\mathbf{c},\mathbf{c}'})[\mathbf{x}^{\mathbf{u}}] (A^T \mathbf{x})^{\mathbf{k}'} [\mathbf{y}^{\mathbf{v}}] (B\mathbf{y})^{\mathbf{k}} \\
&= \sum_{\substack{l \in \mathbb{N}^*, \mathbf{s}, \mathbf{t}, \mathbf{u}, \mathbf{v} \in \mathbb{N}^m \\ |\mathbf{s}|=|\mathbf{t}|=l \\ \mathbf{s}+\mathbf{u}=\mathbf{p} \\ \mathbf{t}+\mathbf{v}=\mathbf{q}}} \frac{(l-1)! \mathbf{p}! \mathbf{q}!}{(n-1)!} \sum_{\substack{\mathbf{c}+\mathbf{k}=\mathbf{s} \\ \mathbf{c}'+\mathbf{k}'=\mathbf{t} \\ |\mathbf{c}|=|\mathbf{c}'| \\ |\mathbf{k}|=|\mathbf{k}'|=|\mathbf{u}|=|\mathbf{v}|}} \frac{(-1)^{|\mathbf{k}|} (|\mathbf{k}|)!}{\mathbf{c}! \mathbf{k}! \mathbf{c}'! \mathbf{k}'!} \text{Per}((A+B)_{\mathbf{c},\mathbf{c}'}) \frac{\text{Per}(A_{\mathbf{k}',\mathbf{u}}^T)}{\mathbf{u}!} \frac{\text{Per}(B_{\mathbf{k},\mathbf{v}})}{\mathbf{v}!} \\
&= \sum_{\substack{l \in \mathbb{N}^*, \mathbf{s}, \mathbf{t}, \mathbf{u}, \mathbf{v} \in \mathbb{N}^m \\ |\mathbf{s}|=|\mathbf{t}|=l \\ \mathbf{s}+\mathbf{u}=\mathbf{p} \\ \mathbf{t}+\mathbf{v}=\mathbf{q}}} \frac{(l-1)! \mathbf{p}! \mathbf{q}!}{(n-1)! \mathbf{u}! \mathbf{v}!} \sum_{\substack{\mathbf{c}+\mathbf{k}=\mathbf{s} \\ \mathbf{c}'+\mathbf{k}'=\mathbf{t} \\ |\mathbf{c}|=|\mathbf{c}'| \\ |\mathbf{k}|=|\mathbf{k}'|=|\mathbf{u}|=|\mathbf{v}|}} \frac{(-1)^{|\mathbf{k}|} (|\mathbf{k}|)!}{\mathbf{c}! \mathbf{k}! \mathbf{c}'! \mathbf{k}'!} \text{Per}(A_{\mathbf{u},\mathbf{k}'}) \text{Per}(B_{\mathbf{k},\mathbf{v}}) \text{Per}((A+B)_{\mathbf{c},\mathbf{c}'}) \\
&= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k k! (n-k-1)!}{(n-1)!} \sum_{\substack{\mathbf{a}+\mathbf{b}+\mathbf{c}=\mathbf{p} \\ \mathbf{a}'+\mathbf{b}'+\mathbf{c}'=\mathbf{q} \\ |\mathbf{a}|=|\mathbf{a}'|=|\mathbf{b}|=|\mathbf{b}'|=k}} \frac{\mathbf{p}! \mathbf{q}!}{\mathbf{a}! \mathbf{b}! \mathbf{c}! \mathbf{a}'! \mathbf{b}'! \mathbf{c}'!} \text{Per}(A_{\mathbf{a},\mathbf{a}'}) \text{Per}(B_{\mathbf{b},\mathbf{b}'}) \text{Per}((A+B)_{\mathbf{c},\mathbf{c}'}), \tag{116}
\end{aligned}$$

where we used Eq. (50) in the third line, $\text{Per}(M^T) = \text{Per}(M)$ in the fourth line, and where we relabeled \mathbf{u} as \mathbf{a} , \mathbf{k}' as \mathbf{a}' , \mathbf{k} as \mathbf{b} , \mathbf{v} as \mathbf{b}' , and $|\mathbf{k}| = |\mathbf{k}'| = |\mathbf{u}| = |\mathbf{v}|$ as $k = n - l$ in the last line. The cutoff of the sum at $\lfloor \frac{n}{2} \rfloor$ comes from the fact that $n = |\mathbf{p}| = |\mathbf{a}| + |\mathbf{b}| + |\mathbf{c}| = 2k + |\mathbf{c}| \geq 2k$. Writing $\frac{(-1)^k k! (n-k-1)!}{(n-1)!} = \frac{(-1)^k}{\binom{n-1}{k}}$ completes the proof of Theorem 4. \square

7 Boson Sampling with input cat states

In this section, we discuss the classical complexity of sampling from the output distribution of linear optical computations with input cat states, which we refer to as Boson Sampling with input cat states.

Since the introduction of Boson Sampling by Aaronson and Arkhipov [6] for the demonstration of quantum computational advantage using noninteracting bosons, several variants of this model have been analyzed [42–49]. These variants were introduced to address two different challenges: on the one hand, to reduce the experimental burden associated with the demonstration of quantum advantage; on the other hand, to understand the resources for this quantum advantage and which computational models are able to reproduce it. Boson Sampling with input cat states is another such variant and has been first considered in [20], where its classical hardness was shown for exact sampling and argued for approximate sampling. We strengthen these results in the following.

We first give a proof of Lemma 1, which provides a closed form expression for output amplitudes of Boson Sampling with input cat states.

Proof of Lemma 1. In the quantum-inspired proof of Glynn's formula in section 5.1 we have obtained the following identity:

$$\langle p_1 \dots p_m | \hat{U}(|\tilde{\text{cat}}_\alpha\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}) = \frac{\alpha^{|\mathbf{p}|-n} e^{-\frac{n}{2}|\alpha|^2}}{2^n \sqrt{\mathbf{p}!}} \sum_{\mathbf{x} \in \{-1,1\}^n} x_1 \dots x_n \prod_{i=1}^m \left(\sum_{j=1}^n u_{ij} x_j \right)^{p_i}, \quad (117)$$

for all $\alpha \in \mathbb{C}$, all $\mathbf{p} = (p_1, \dots, p_m) \in \mathbb{N}^m$, all $n \leq m$ and any passive linear operation \hat{U} over m modes with unitary matrix $U = (u_{ij})_{1 \leq i, j \leq m}$, where $|\text{cat}_\alpha\rangle = \frac{1}{2\alpha}(|\alpha\rangle - |-\alpha\rangle)$ is an unnormalized cat state. By Eq. (5), these states are related to normalized cat states $|\text{cat}_\alpha\rangle$ by

$$|\text{cat}_\alpha\rangle = \frac{\alpha e^{\frac{1}{2}|\alpha|^2}}{\sqrt{\sinh(|\alpha|^2)}} |\tilde{\text{cat}}_\alpha\rangle. \quad (118)$$

Setting $|\mathbf{p}| = n$ and replacing unnormalized cat states by normalized ones in Eq. (117), we obtain Lemma 1:

$$\begin{aligned} \langle p_1 \dots p_m | \hat{U}(|\text{cat}_\alpha\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}) &= \frac{\alpha^n}{\sqrt{\sinh^n(|\alpha|^2)} 2^n \sqrt{\mathbf{p}!}} \sum_{\mathbf{x} \in \{-1,1\}^n} x_1 \dots x_n \prod_{i=1}^m \left(\sum_{j=1}^n u_{ij} x_j \right)^{p_i} \\ &= \frac{\alpha^n}{\sqrt{\sinh^n(|\alpha|^2)}} \frac{\text{Per}(U_{\mathbf{p}, \mathbf{1} \oplus \mathbf{0}})}{\sqrt{\mathbf{p}!}} \\ &= \frac{\alpha^n}{\sqrt{\sinh^n(|\alpha|^2)}} \langle p_1 \dots p_m | \hat{U}(|1\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}), \end{aligned} \quad (119)$$

where we used Eq. (39) in the second line and Eq. (9) in the last line. \square

Interestingly, Lemma 1 implies that, up to a global factor, n cat states of amplitude α reproduce exactly the Boson Sampling statistics of n single-photons (however, detection events \mathbf{p} with $|\mathbf{p}| > n$ can occur for input cat states, contrary to single-photons).

This result has two immediate consequences, summarized by Theorem 5. Firstly, it implies that Boson Sampling with input cat states is hard to sample exactly for $m \geq 2n$ and for all choices of cat state amplitude unless the polynomial hierarchy collapses to its third level, since some of its outcome probabilities are $\#\text{P}$ -hard to estimate multiplicatively [6]. An alternative proof of this statement based on universality under post-selection was given in [20]. Secondly, as we show hereafter, it implies that Boson Sampling with input cat states of small enough amplitudes is also hard to sample approximately.

Proof of Theorem 5. Our proof uses arguments similar to the ones used in [20], extended to the case of approximate sampling. Let us prove that for $|\alpha| = \mathcal{O}(n^{-1/4} \log^{1/4} n)$, the detection events satisfying $|\mathbf{p}| = n$ represent an inverse-polynomial fraction of the outcomes for Boson Sampling with input cat states of amplitude α .

Let $P_{\text{BS}}(\mathbf{p}|n) := |\langle p_1 \dots p_m | \hat{U}(|1\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n})|^2$ be the probability of detecting $\mathbf{p} = (p_1, \dots, p_m)$ output photons in a Boson Sampling experiment with interferometer \hat{U} and input single-photons $|1\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}$. We have $P_{\text{BS}}(\mathbf{p}|n) = 0$ whenever $|\mathbf{p}| \neq n$ since \hat{U} conserves the total number of photons. Hence,

$$\begin{aligned} \sum_{|\mathbf{p}|=n} |\langle p_1 \dots p_m | \hat{U}(|1\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n})|^2 &= \sum_{|\mathbf{p}|=n} P_{\text{BS}}(\mathbf{p}|n) \\ &= \sum_{\mathbf{p}} P_{\text{BS}}(\mathbf{p}|n) \\ &= 1. \end{aligned} \quad (120)$$

With Eq. (119), the probability of detecting $\mathbf{p} = (p_1, \dots, p_m)$ output photons such that $|\mathbf{p}| = n$ in a Boson Sampling experiment with interferometer \hat{U} and input cat states $|\text{cat}_\alpha\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}$ is thus given by:

$$\begin{aligned} \sum_{|\mathbf{p}|=n} \left| \langle p_1 \dots p_m | \hat{U} (|\text{cat}_\alpha\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}) \right|^2 &= \frac{|\alpha|^{2n}}{\sinh^n(|\alpha|^2)} \sum_{|\mathbf{p}|=n} \left| \langle p_1 \dots p_m | \hat{U} (|1\rangle^{\otimes n} \otimes |0\rangle^{\otimes m-n}) \right|^2 \\ &= \frac{|\alpha|^{2n}}{\sinh^n(|\alpha|^2)}. \end{aligned} \tag{121}$$

Assuming $|\alpha| = \mathcal{O}(n^{-1/4} \log^{1/4} n)$, this last quantity satisfies $\frac{|\alpha|^{2n}}{\sinh^n(|\alpha|^2)} \geq \frac{1}{\text{poly}(n)}$.

This implies that any efficient classical algorithm for approximate sampling from the output probability distribution of a Boson Sampling instance with input cat states with $|\alpha| = \mathcal{O}(n^{-1/4} \log^{1/4} n)$ may also efficiently sample approximately from the output probability distribution of the corresponding Boson Sampling instance with input single-photons efficiently, by keeping only the samples \mathbf{p} satisfying $|\mathbf{p}| = n$. This proves that Boson Sampling with input cat states is hard to sample approximately, assuming the same conjectures as in [6]. \square

Acknowledgments

U. C. thanks Atul Singh Arora and Pierre-Emmanuel Emeriau for interesting discussions. U. C, A. D, and S. M. acknowledge funding provided by the Institute for Quantum Information and Matter, an NSF Physics Frontiers Center (NSF Grant PHY-1733907). A. D. also acknowledges support from the National Science Foundation RAISE-TAQS 1839204 and Amazon Web Services, AWS Quantum Program.

References

- [1] H. Minc, “Permanents,” vol. 6. [Cambridge University Press](#), 1984.
- [2] J. K. Percus, “Combinatorial methods,” vol. 4. [Springer Science & Business Media](#), 2012.
- [3] L. G. Valiant, “The complexity of computing the permanent,” *Theoretical computer science* **8**, 189–201 (1979).
- [4] E. R. Caianiello, “On quantum field theory—I: explicit solution of Dyson’s equation in electrodynamics without use of Feynman graphs,” *Il Nuovo Cimento (1943-1954)* **10**, 1634–1652 (1953).
- [5] S. Scheel, “Permanents in linear optical networks,” [quant-ph/0406127](#).
- [6] S. Aaronson and A. Arkhipov, “The computational Complexity of Linear Optics,” *Theory of Computing* **9**, 143 (2013).
- [7] S. Aaronson, “A linear-optical proof that the permanent is #P-hard,” *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* **467**, 3393–3405 (2011).
- [8] S. Rahimi-Keshari, A. P. Lund, and T. C. Ralph, “What can quantum optics say about computational complexity theory?,” *Physical review letters* **114**, 060501 (2015).
- [9] D. Grier and L. Schaeffer, “New hardness results for the permanent using linear optics,” [arXiv:1610.04670](#).
- [10] P. P. Rohde, D. W. Berry, K. R. Motes, and J. P. Dowling, “A Quantum Optics Argument for the #P-hardness of a Class of Multidimensional Integrals,” [arXiv:1607.04960](#).

- [11] L. Chakhmakhchyan, N. J. Cerf, and R. Garcia-Patron, “Quantum-inspired algorithm for estimating the permanent of positive semidefinite matrices,” *Physical Review A* **96**, 022329 (2017).
- [12] P. A. MacMahon, “Combinatory Analysis, Volumes I and II,” vol. 137. American Mathematical Soc., 2001.
- [13] I. Good, “Proofs of some ‘binomial’ identities by means of MacMahon’s ‘Master Theorem’,” in *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 58, pp. 161–162, Cambridge University Press. 1962.
- [14] L. Carlitz, “An application of MacMahon’s master theorem,” *SIAM Journal on Applied Mathematics* **26**, 431–436 (1974).
- [15] L. Carlitz, “Some expansions and convolution formulas related to MacMahon’s master theorem,” *SIAM Journal on Mathematical Analysis* **8**, 320–336 (1977).
- [16] H. J. Ryser, “Combinatorial mathematics,” vol. 14. American Mathematical Soc., 1963.
- [17] K. Balasubramanian, *Combinatorics and diagonals of matrices*. PhD thesis, Indian Statistical Institute-Kolkata, 1980.
- [18] E. T. Bax, *Finite-difference algorithms for counting problems*. PhD thesis, California Institute of Technology, 1998.
- [19] D. G. Glynn, “The permanent of a square matrix,” *European Journal of Combinatorics* **31**, 1887–1891 (2010).
- [20] P. P. Rohde, K. R. Motes, P. A. Knott, J. Fitzsimons, W. J. Munro, and J. P. Dowling, “Evidence for the conjecture that sampling generalized cat states with linear optics is hard,” *Physical Review A* **91**, 012342 (2015).
- [21] C. Weedbrook, S. Pirandola, R. García-Patrón, N. J. Cerf, T. C. Ralph, J. H. Shapiro, and S. Lloyd, “Gaussian quantum information,” *Reviews of Modern Physics* **84**, 621 (2012).
- [22] U. Chabaud and S. Mehraban, “Holomorphic Quantum Computing,” [arXiv:2111.00117](https://arxiv.org/abs/2111.00117).
- [23] A. C. Dixon, “On the sum of the cubes of the coefficients in a certain expansion by the binomial theorem,” *Messenger of mathematics* **20**, 79–80 (1891).
- [24] I. Good, “A short proof of MacMahon’s ‘Master Theorem’,” in *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 58, pp. 160–160, Cambridge University Press. 1962.
- [25] S. Garoufalidis, T. T. Lê, and D. Zeilberger, “The quantum MacMahon master theorem,” *Proceedings of the National Academy of Sciences* **103**, 13928–13931 (2006).
- [26] M. Konvalinka and I. Pak, “Non-commutative extensions of the MacMahon Master Theorem,” *Advances in Mathematics* **216**, 29–61 (2007).
- [27] M. P. Tuite, “Some generalizations of the MacMahon Master Theorem,” *Journal of Combinatorial Theory, Series A* **120**, 92–101 (2013).
- [28] V. V. Kocharovskiy, V. V. Kocharovskiy, and S. V. Tarasov, “The Hafnian Master Theorem,” *Linear Algebra and its Applications* 144–161 (2022).
- [29] W. Y. Chen, H. Galbraith, and J. Louck, “Angular momentum theory, umbral calculus, and combinatorics,” *Computers & Mathematics with Applications* **41**, 1199–1214 (2001).
- [30] B. M. Terhal and D. P. DiVincenzo, “Classical simulation of noninteracting-fermion quantum circuits,” *Physical Review A* **65**, 032325 (2002).
- [31] V. Shchesnovich, “Partial indistinguishability theory for multiphoton experiments in multipoint devices,” *Physical Review A* **91**, 013844 (2015).

- [32] D. Spivak, M. Y. Niu, B. C. Sanders, and H. de Guise, “Generalized interference of fermions and bosons,” *Physical Review Research* **4**, 023013 (2022).
- [33] E.-J. Kuo, Y. Xu, D. Hangleiter, A. Grankin, and M. Hafezi, “Boson Sampling for Generalized Bosons,” [arXiv:2204.08389](#).
- [34] A. Clément, N. Heurtel, S. Mansfield, S. Perdrix, and B. Valiron, “LO_v-Calculus: A Graphical Language for Linear Optical Quantum Circuits,” [arXiv:2204.11787](#).
- [35] G. De Felice and B. Coecke, “Quantum Linear Optics via String Diagrams,” [arXiv:2204.12985](#).
- [36] B. Peropadre, G. G. Guerreschi, J. Huh, and A. Aspuru-Guzik, “Proposal for microwave boson sampling,” *Physical review letters* **117**, 140505 (2016).
- [37] S. Girvin, “Schrödinger cat states in circuit qed,” [arXiv:1710.03179](#).
- [38] X. Gu, A. F. Kockum, A. Miranowicz, Y.-x. Liu, and F. Nori, “Microwave photonics with superconducting quantum circuits,” *Physics Reports* **718**, 1–102 (2017).
- [39] J. Huh, “A fast quantum algorithm for computing matrix permanent,” [arXiv:2205.01328](#).
- [40] S. Aaronson and T. Hance, “Generalizing and Derandomizing Gurvits’s Approximation Algorithm for the Permanent,” *Quantum Info. Comput.* **14**, 541–559 (2014).
- [41] D. M. Jackson, “The unification of certain enumeration problems for sequences,” *Journal of Combinatorial Theory, Series A* **22**, 92–96 (1977).
- [42] A. P. Lund, A. Laing, S. Rahimi-Keshari, T. Rudolph, J. L. O’Brien, and T. C. Ralph, “Boson sampling from a Gaussian state,” *Physical review letters* **113**, 100502 (2014).
- [43] J. P. Olson, K. P. Seshadreesan, K. R. Motes, P. P. Rohde, and J. P. Dowling, “Sampling arbitrary photon-added or photon-subtracted squeezed states is in the same complexity class as boson sampling,” *Physical Review A* **91**, 022317 (2015).
- [44] C. S. Hamilton, R. Kruse, L. Sansoni, S. Barkhofen, C. Silberhorn, and I. Jex, “Gaussian boson sampling,” *Physical review letters* **119**, 170501 (2017).
- [45] A. Lund, S. Rahimi-Keshari, and T. Ralph, “Exact boson sampling using Gaussian continuous-variable measurements,” *Physical Review A* **96**, 022301 (2017).
- [46] L. Chakhmakhchyan and N. J. Cerf, “Boson sampling with Gaussian measurements,” *Physical Review A* **96**, 032326 (2017).
- [47] U. Chabaud, T. Douce, D. Markham, P. van Loock, E. Kashefi, and G. Ferrini, “Continuous-variable sampling from photon-added or photon-subtracted squeezed states,” *Physical Review A* **96**, 062307 (2017).
- [48] N. Quesada, J. M. Arrazola, and N. Killoran, “Gaussian boson sampling using threshold detectors,” *Physical Review A* **98**, 062322 (2018).
- [49] A. Deshpande, A. Mehta, T. Vincent, N. Quesada, M. Hinsche, M. Ioannou, L. Madsen, J. Lavoie, H. Qi, J. Eisert, *et al.*, “Quantum computational advantage via high-dimensional Gaussian boson sampling,” *Science advances* **8**, 7894 (2022).