# Hear $\pi$ from Quantum Galperin Billiards

Yin Cai<sup>1</sup> and Fu-Lin Zhang<sup>1,\*</sup>

<sup>1</sup>Department of Physics, School of Science, Tianjin University, Tianjin 300072, China (Dated: August 2, 2022)

Galperin [Regul. Chaotic. Dyn. 8, 375 (2003)] introduced an interesting method to learn the digits of  $\pi$  by counting the collisions of two billiard balls and a hard wall. This paper studies two quantum versions of the Galperin billiards. It is shown that the digits of  $\pi$  can be observed in the phase shifts of the quantum models.

## I. INTRODUCTION

The number  $\pi$  has attracted interest in human history. There are various elegant ways to determine  $\pi$  with a good precision; for instance, the inscribed and circumscribed polygons on a circle [1–3] and the Monte Carlo Method baseed on computers. Twenty years ago, Galperin [4, 5] invented an extraordinary method to learn the digits of  $\pi$ , known as Galperin billiards. In such a scenario, as shown in Fig. 1, a big ball is hurled towards a small ball that has its back to a wall. An observer counts the ensuing elastic collisions until the big ball's momentum is redirected enough to fully turn around. When the ratio of masses for the two billiard balls  $\frac{M}{m}=100^N$ , the total number of hits in the system is  $[\pi 10^N]$  where  $[\ ]$  is the greatest integer function.

The interesting results of Galperin billiards have been discussed in different aspects, such as their relationships with geometrical optics and Grover algorithm for quantum search [4, 6] In the present paper, we study two quantum extensions of the Galperin billiards. In our first (semiclassical) model, the motion of the small ball is assumed to be described by quantum mechanics, and the big one is classical. In addition, the classical part is assumed to move slowly, and hence the quantum part is treated under the adiabatic approximation. In the second (fully quantum) version, both balls are quantum, which is equivalent to a scattering problem in a two-dimensional unbounded sector domain. The phase shifts in the two models are shown to correspond to the number of collisions in classical mechanics. As phase shifts in scattering problems are slosely related with state densities and energy spectrums [7, 8], this work connects the Galperin




FIG. 1: Schematic picture of Galperin billiards, consisting of a big ball M, a small ball m and a wall.

model with the classic problem  $Can\ you\ hear\ the\ shape$  of a drum? formulated by Mark Kac [9]. That is, the shape of two-dimensional unbounded sector domain, corresponding to the quantum Galperin model, determines the spectrum of a quantum particle in it and the digits of  $\pi$  in collisions.

#### II. SEMICLASSICAL MODEL

We begin by replacing the small ball in the original model with a quantum particle. As shown in Fig. 2, the classical big ball and the wall constitute a one-dimensional infinite potential well, as the quantum small ball is situated between them. The motions of the two balls are governed by their interaction. Let x and y denote the positions of the big and small balls respectively. The small ball satisfies the Schrödinger equation

$$i\hbar\frac{\partial}{\partial t}\Psi\left(y,t\right) = \left[-\frac{\hbar^{2}}{2m}\frac{\partial^{2}}{\partial y^{2}} + V\left(y\right)\right]\Psi\left(y,t\right),$$
 (1)

where m is its mass and

$$V(y) = \begin{cases} 0, & 0 \leq y \leq x, \\ +\infty, & \text{otherwise.} \end{cases}$$
 (2)




FIG. 2: Semiclassical Galperin billiards.

Here, we assume that the big ball moves very gradually and therefore the wave function of the quantum particle changes adiabatically. Under the adiabatic approximation, the wave function of a small ball is given by

$$\Psi = \sum_{n=1}^{+\infty} C_n \Psi_n e^{-i\frac{1}{\hbar} \int_0^t E_n dt + i\gamma_n},\tag{3}$$

where 
$$E_n = \frac{n^2 \pi^2 \hbar^2}{2mx^2}, \Psi_n = \sqrt{\frac{2}{x}} \sin \frac{n\pi y}{x}$$
, and  $\gamma_n$  is the

<sup>\*</sup>Corresponding author: flzhang@tju.edu.cn

Berry phase gained by each instantaneous eigenenstate [10]. Here, both the initial and final values of a are  $+\infty$ , and the final time  $t \to +\infty$ . The the Berry phase can be directly calculated as

$$\gamma_n = i \int_0^t \langle \Psi_n | \frac{d}{dt} | \Psi_n \rangle dt = 0, \tag{4}$$

as  $\langle \Psi_n | d | \Psi_n \rangle = 0$ .

To observe the motion of the quantum particle, we consider a superposition of two levels adjacent levels as

$$\Psi = \frac{1}{\sqrt{2}} (\Psi_n e^{-i\frac{1}{\hbar} \int_0^t E_n dt} + \Psi_{n+1} e^{-i\frac{1}{\hbar} \int_0^t E_{n+1} dt}).$$
 (5)

Its average position is given by

$$\overline{y} = \langle \Psi | y | \Psi \rangle$$

$$= \frac{x}{2} - \frac{x}{\pi^2} \frac{8n(n+1)}{(2n+1)^2} \cos \int_0^t \frac{E_{n+1} - E_n}{\hbar} dt.$$
 (6)

The average position vibrates around the midpoint  $\frac{x}{2}$  at the Bohr frequency  $\frac{E_{n+1}-E_n}{\hbar}$ . Hence, the phase difference, divided by  $\pi$ , between the two eigenenstates can be regarded as the quantum correspondence of the number of collisions in classical mechanics.

The energy of the quantum ball is

$$E = \frac{1}{2}(E_n + E_{n+1}) = \frac{2n^2 + 2n + 1}{2} \frac{\pi^2 \hbar^2}{mx^2}.$$
 (7)

We assume that the speed change of the classical ball is determined by its energy exchange with the quantum particle. One can obtain

$$\frac{2n^2 + 2n + 1}{2} \frac{\pi^2 \hbar^2}{m} \left(\frac{1}{x_{\min}^2} - \frac{1}{x^2}\right) = \frac{1}{2} M v^2, \tag{8}$$

where  $v = \frac{dx}{dt}$ , M is the mass of the classical ball and  $x_{\min}$  is its retracing point. Then, the definite integral in (6) can be solved by variable transformation from t to x. It is obtained as

$$\int_{0}^{+\infty} \frac{E_{n+1} - E_n}{\hbar} dt = \sqrt{\frac{4n^2 + 4n + 1}{4n^2 + 4n + 2}} \pi^2 R, \quad (9)$$

where  $R = \sqrt{\frac{M}{m}}$  is the square root of the two masses ratios. The number of the cosine function in (6) reaching its extreme values is  $\sqrt{\frac{4n^2+4n+1}{4n^2+4n+2}}\pi R$ , which does not depend on  $x_{\min}$ , or the initial speed of the classical ball.

In the classical model, the full change of phase of action-angle variables is  $\pi^2 R$  [11] and the collision times is  $\left[\frac{\pi}{\beta}\right]$  with  $\beta = \operatorname{arccot} R$  [4]. When  $n \to +\infty$  and  $R \to +\infty$ , the phase difference (9) in the semiclassical model is equal to the full change of phase in the classical model. At the same time, the number of extremums in the semiclassical model is equal to the collision times in




FIG. 3: A comparison between the average position  $\bar{y}$  and the corresponding results of a classical model when  $\beta = \pi/10$ . The three curves are obtained for the classical model (solid line), the semiclassical model with quantum number n=1 (dashed line), and n=10 (dot-dashed line).




FIG. 4: The fully quantum Galperin model is equivalent to a free particle in a two-dimensional unbounded sector domain in the first quadrant where  $\frac{y}{x} \in [0, \sqrt{\frac{m}{M}}]$ .

the classical model. We define a normalized time as

$$\alpha = \operatorname{sgn}(v) \arccos \frac{\rho_{\min}}{\rho},$$
 (10)

where sgn is the sign function and  $\rho = \sqrt{Mx^2 + my^2}$ . As t increases from 0 to  $+\infty$ , the value of  $\alpha$  changes continuously from  $-\frac{\pi}{2}$  to  $\frac{\pi}{2}$ , and  $\rho$  decreases from  $+\infty$  to  $\rho_{\min}$  and then goes back to  $+\infty$ . Using  $\alpha$  as the variable, in Fig. 3, we show x comparison between the average position  $\bar{y}$  and the result of a corresponding classical model. In general, the oscillation of  $\bar{y}$  correspond to the collisions in classical mechanics. The period of the quantum oscillation is greater than the one of the classical collisions, while the difference between them is reduced as the quantum number n increases.

### III. FULLY QUANTUM MODEL

We now consider a fully quantum model, in which both balls are described by quantum mechanics. The stationary Schrödinger equation of the fully quantum model can be written as

$$\[ -\frac{\hbar^2}{2M} \frac{\partial^2}{\partial x^2} - \frac{\hbar^2}{2m} \frac{\partial^2}{\partial y^2} + V(x, y) \] \Psi(x, y) = E\Psi(x, y).$$
(11)

where x and y denote respectively the positions of the big ball of mass M and the small ball of mass m, and

$$V(x,y) = \begin{cases} 0, & 0 \leqslant y \leqslant x, \\ +\infty, & \text{otherwise.} \end{cases}$$
 (12)

Such a two-body problem in one-dimensional space can be converted into a free particle in a two-dimensional unbounded sector domain. Namely, introducing the polar coordinates

$$\sqrt{M}x = \rho\cos\theta, \quad \sqrt{m}y = \rho\sin\theta,$$
(13)

and the Schr odinger equation turns to

$$\[ -\frac{\hbar^2}{2} \left( \frac{\partial^2}{\partial \rho^2} + \frac{1}{\rho} \frac{\partial}{\partial \rho} + \frac{1}{\rho^2} \frac{\partial^2}{\partial \theta^2} \right) + V(\theta) \] \Psi(\rho, \theta) = E \Psi(\rho, \theta), \tag{14}$$

where

$$V(\theta) = \begin{cases} 0, & 0 \leq \theta \leq \beta, \\ +\infty, & \text{otherwise.} \end{cases}$$
 (15)

When  $M \gg m$ ,  $\rho \approx \sqrt{M}x$  and  $\theta/\beta \approx y/x$ . The former approaches to the position of the big ball, and the latter becomes the (normalized) position of the small one corresponding to the curves in Fig. 3.

As the potential  $V(\theta)$  does not depend on the radial coordinate  $\rho$ , the stationary Schrödinger equation (14) can be solved by using the separation of variables method. We assume that  $\Psi(\rho,\theta)=R(\rho)\Phi(\theta)$  and  $\Phi(\theta)=0$  when  $\theta\notin[0,\beta]$ . The Schrödinger equation turns to

$$-\frac{\partial^2}{\partial\theta^2}\Phi(\theta)=l^2\Phi(\theta), \qquad (16)$$

$$-\frac{\hbar^2}{2} \left( \frac{\partial^2}{\partial \rho^2} + \frac{1}{\rho} \frac{\partial}{\partial \rho} - \frac{l^2}{\rho^2} \right) R(\rho) = ER(\rho). \tag{17}$$

The eigenfuntions and eigenvalues for the angular part can be directly found to be  $\Phi(\theta) = \sin l\theta$  when  $\theta \in [0, \beta]$  and  $l = n\pi/\beta$  with n = 1, 2, 3... The physical solution to the radial equation (17) is a linear combination of the first kind of Bessel function, as the second kind of Bessel function tends to  $-\infty$  when  $\rho \to 0$  [12]. Here, we write the common eigenfunctions in terms of the Hankel functions as

$$\Psi_{k,l}(\rho,\theta) = [H_l^{(1)}(k\rho) + H_l^{(2)}(k\rho)] \sin l\theta, \quad (18)$$

with  $k = \sqrt{2E}/\hbar$ .

The probability current desity of the stationary wave function (18) can be divided into two equal and opposite parts without cross item, which are along the radial direction and correspond to the two terms  $H_l^{(1)}(k\rho) \sin l\theta$  and

 $H_l^{(2)}(k\rho)\sin l\theta$  respectively. In addition, when  $\rho\to\infty$ ,

$$H_l^{(1)}(k\rho) \to \sqrt{\frac{2}{\pi k \rho}} e^{i[(k\rho - (l + \frac{1}{2})\frac{\pi}{2})]},$$
 (19)

$$H_l^{(2)}(k\rho) \to \sqrt{\frac{2}{\pi k \rho}} e^{-i[k\rho - (l + \frac{1}{2})\frac{\pi}{2}]},$$
 (20)

which become spherical waves in the two-dimensional sector domain. Therefore, the two terms,  $H_l^{(1)}(k\rho)\sin l\theta$  and  $H_l^{(2)}(k\rho)\sin l\theta$ , are incident and outgoing waves respectively. They can be regarded as the quantum correspondences of the two movement processes in the classical model divided by the retracing point of the big ball. Here, an observation is that, although a more reasonable incidence wave should promulgate along x-axis, the incident direction makes a difference of at most 1 on the number of classical reflections in the unbounded sector domain in Fig. 4 [4]. The phase shift between the incident and outgoing waves can be directly derived from the asymptotic formulations (19) and (20) as

$$\delta(n) = (l + \frac{1}{2})\pi = (\frac{n\pi}{\beta} + \frac{1}{2})\pi.$$
 (21)

It does not depend on the wavenumber k, or energy E equivalently, and can be regarded as the phase gained by the standing wave  $\sin l\theta$ . The difference between two adjacent phase shifts is

$$\Delta \delta = \delta(n+1) - \delta(n) = \frac{\pi^2}{\beta}.$$
 (22)

When  $M \gg m$ , it becomes  $\Delta \delta = \pi^2 R$ , which corresponds to the phase difference (9) in the semiclassical model and the change of phase in classical model.

To observe the reciprocal motion of y between 0 and x, *i.e.*  $\theta$  between 0 and  $\beta$ , we consider a superposition of two eigenfunctions with the same energy and adjacent angular quantum number l as

$$\Psi = \Psi_{k l} + e^{ic\pi} \Psi_{k l'}, \tag{23}$$

where  $l=n\pi/\beta$ ,  $l'=(n+1)\pi/\beta$  and  $c=\frac{\pi}{2\beta}$ . Here, the relative phase  $e^{ic\pi}$  ensures the phase consistency between the average position in quantum model and classical position. We also divide it into incident and outgoing parts according to the superscripts of Hankel functions. The incident wave reads

$$\Psi^{\rm in} = H_l^{(1)}(k\rho) \sin l\theta + e^{ic\pi} H_{l'}^{(1)}(k\rho) \sin l'\theta, \qquad (24)$$

and the outgoing term has the same form with the Hankel functions of the first kind replaced with the second kind. For a fixed  $\rho$ , the average values of  $\theta$  in the incident wave




FIG. 5: A comparison between the average position  $\bar{\theta}$  and the corresponding results of a classical model when  $\beta=\pi/10$ . The three curves are obtained for the classical model (solid line) and the fully quantum model with quantum number  $l=\frac{\pi}{\beta}$  (dashed line) and  $l=\frac{10\pi}{\beta}$  (dot-dashed line),

is

$$\begin{split} \overline{\theta}(\rho) &= \frac{\int \Psi^{\text{in}*} \theta \Psi^{\text{in}} d\theta}{\int \Psi^{\text{in}*} \Psi^{\text{in}} d\theta} \\ &= \frac{\beta}{2} - \frac{\beta}{\pi^2} \frac{8n(n+1)}{(2n^2+1)^2} \frac{e^{ic\pi} H_l^{(2)} H_{l'}^{(1)} + e^{-ic\pi} H_l^{(1)} H_{l'}^{(2)}}{H_l^{(1)} H_l^{(2)} + H_{l'}^{(1)} H_{l'}^{(2)}}, (25) \end{split}$$

where we omit the argument  $k\rho$  to the Hankel functions and the result in outgoing wave for simplicity. The expression (25) is similar to the average position (6) of the quantum particle in the semiclassical model. In addition, the vibration of the average value  $\overline{\theta}(\rho)$  with  $\rho$  depends on the relative phase between  $H_l^{(1)}$  and  $H_{l'}^{(2)}$ , since  $H_{\gamma}^{(1)*} = H_{\gamma}^{(2)}$  and  $|H_{\gamma}^{(1)}(k\rho)|$  is a monotonous decrease function of  $k\rho$ .

In Fig. 5, we show a comparison between the average value  $\overline{\theta}(\rho)$  and the classical motion of  $\theta$ , as functions of

$$\eta = \arccos \frac{l}{k\rho}.$$
 (26)

As  $\eta$  increases from 0 to  $\frac{\pi}{2}$ , the value of  $\rho$  changes continuously from  $\frac{l}{k}$  to  $+\infty$ . Here,  $\frac{l}{k}$  is the quantum correspondence of the minimum of  $\rho$  in classical model, *i.e.* its retracing point  $\rho = \rho_{\min}$ . Thus, the parameter  $\eta$  in classical model is defined by replacing  $\frac{l}{k}$  with  $\rho_{\min}$  in Eq. (26). Although the value of  $\rho$  is allowed to be less than  $\rho_{\min}$  in the fully quantum model, there is no vibration of  $\theta(\rho)$  in the region of  $\rho < \rho_{\min}$ . There is a obvious stationary region of the fully quantum model where  $\eta$  is near zero. The region becomes smaller as the quantum number l increases. When  $l \to +\infty$ , this region almost disappear, and the period of the quantum oscillation is consistent with the one of the classical collisions.

#### IV. SUMMARY

In summary, we study two quantum extensions of the Galperin billiards, one of which is semiclassical and the other is fully quantum. In addition, we assume the small quantum ball in the semiclassical model, can be treated under the adiabatic approximation as the big classical ball moves adiabatically. The phase shifts in the two models are shown to correspond to the number of collisions in classical mechanics.

As the quantum model connects the digits of  $\pi$  in the classical Galperin model with the spectrum of a quantum particle in a two-dimensional unbounded sector domain. It would be intriguing to study the relationship among the digits of  $\pi$  and other physical problems in the same domain, such as statistical mechanics and phonon eigenfunctions. Actually, these physical problems in similar domains xare attracting attention [13, 14] .

<sup>[1]</sup> P. Beckmann, A History of  $\pi$  (St. Martin's Press, New York, 1971).

<sup>[2]</sup> L. Berggren, J. Borwein, and P. Borwein, Pi: A Source Book, 3<sup>rd</sup> edition (Springer-Verlag, New York, 2004).

<sup>[3]</sup> J. P. Delahaye, Le fascinant nombre  $\pi$  (Pour La Science, Berlin, 1977).

<sup>[4]</sup> G. A. Galperin, Regular & Chaotic Dynamics 8, 375 (2003).

<sup>[5]</sup> G. A. Galperin, Matematicheskoye Prosveschenye 3, 137 (2001).

<sup>[6]</sup> A. R. Brown, Quantum 4, 357 (2020).

<sup>[7]</sup> W.-S. Dai and M. Xie, Journal of High Energy Physics 2009, 033 (2009).

<sup>[8]</sup> W.-D. Li and W.-S. Dai, The European Physical Journal C 75, 1 (2015).

<sup>[9]</sup> M. Kac, The American Mathematical Monthly 73, 1 (1966).

<sup>[10]</sup> D. J. Griffiths and D. F. Schroeter, Introduction to Quantum Mechanics (Cambridge university press, Cambridge, 2018).

<sup>[11]</sup> I. V. Gorelyshev, Regular & Chaotic Dynamics 11, 61 (2006).

<sup>[12]</sup> R. Courant and D. Hilbert, Methods of mathematical physics (John Wiley & Sons, New York, 1989).

<sup>[13]</sup> G. H. Zhang and D. R. Nelson, Physical Review Letters 125, 215503 (2020).

<sup>[14]</sup> G. H. Zhang and D. R. Nelson, Physical Review E 104, 065005 (2021).