# Force balance in thermal quantum many-body systems from Noether's theorem

Sophie Hermann<sup>1</sup> and Matthias Schmidt<sup>1</sup>

<sup>1</sup> Theoretische Physik II, Physikalisches Institut, Universität Bayreuth, D-95447 Bayreuth, Germany (Dated: 31 July 2022)

We address the consequences of invariance properties of the free energy of spatially inhomogeneous quantum many-body systems. We consider a specific position-dependent transformation of the system that consists of a spatial deformation and a corresponding locally resolved change of momenta. This operator transformation is canonical and hence equivalent to a unitary transformation on the underlying Hilbert space of the system. As a consequence, the free energy is an invariant under the transformation. Noether's theorem for invariant variations then allows to derive an exact sum rule, which we show to be the locally resolved equilibrium one-body force balance. For the special case of homogeneous shifting, the sum rule states that the average global external force vanishes in thermal equilibrium.

### I. INTRODUCTION

When investigating global equilibrium properties such as the equation of state for a given many-body Hamiltonian, the strategies in classical and quantum statistical mechanical treatments differ markedly from each other. Obtaining the partition sum in the classical case requires, in principle, to carry out the high-dimensional phase space integral over the Boltzmann factor of the Hamiltonian [1]. The quantum mechanical analog thereof is the trace over the Boltzmann factor of the Hamiltonian, where the latter is viewed as an infinite-dimensional matrix expressed in a suitable basis of e.g. energy eigenfunctions [2]. In both cases, quantum and classical, the leap from the dynamics of the particle-based many-body description to the thermal average is both powerful and abstract. As a result physically meaningful quantities, such as the pressure, chemical and thermal susceptibilities etc. become systematically available, at least in principle, through derivatives of the free energy, which is readily available from the partition sum. On a higher level of detail, locally resolved correlation functions are available as statistical averages and they characterize the microscopic structure of the system and allow to obtain global properties via suitable integration.

On the other hand the concept of forces, while being at the very heart of mechanics, often receives less attention in both statistical and quantal contexts. Nevertheless in the realm of quantum many-body systems several recent publications [3–5] addressed in detail the force balance relationship on the one-body level of correlation functions. Here the forces are resolved in position and also in time in the dynamic case. Tarantino and Ullrich [3] reformulated time-dependent Kohn-Sham density functional theory (DFT) in terms of the second time derivative of the density. In their approach forces feature prominently. They argue that the causal structure of their formulation is more transparent than that of the standard Kohn-Sham formalism of DFT. Rubio and his coworkers [4] have addressed the force balance in several advanced approximations in DFT. They state that their approach avoids differentiability and causality issues and having to carry out the optimized-effectivepotential procedure of orbital-dependent functionals.

Locally resolved force fields play a prominent role in the recent power functional framework for many-body dynamics [5]. Besides the time-dependent density profile, this variational approach includes the locally resolved current and acceleration distributions as its fundamental physical variables. The theory has been formulated for classical [6, 7] and quantum [8, 9] systems; reference [5] reviews these approaches and gives much background. The respective variational equation has the clear physical interpretation of a nonequilibrium force balance relationship and it allows to categorize flow and structural forces [10, 11] and acceleration viscous forces [12], all of which go beyond the adiabatic forces that are captured by the dynamical classical DFT [13–17]. Standard DFT is recovered as the equilibrium limit of the power functional theory [5].

Furthermore, in the classical context, forces were recently put to the fore in methods to obtain statistically averaged quantities, such as the density profile of a spatially inhomogeneous system, via computer simulation of the many-body problem. In his recent review [18], Rotenberg gives a clear account of such force-sampling techniques; see e.g. references [19–21] for original work. On the theoretical side, classical DFT [13-15] offers access to forces via building the gradient of the Euler-Lagrange minimization equation [5]. An alternative that applies to pairwise interparticle interactions is the force integral over the two-body density correlation function [33, 34]. The two-body density is explicitly available within stateof-the-art classical density functionals, such as fundamental measure theory, see e.g. Ref. [22]. Two-body density correlation functions are also central to the recently developed conditional probability DFT for quantum systems [23, 24].

In prior work we have applied Noether's theorem of invariant variations [25, 26] to the classical statistical mechanics of particle-based many-body systems [27–30]. Rather than starting with the invariance properties of an action functional, the approach is based on considering the symmetry properties of appropriate statistical

functionals, such as the partition sum, in order to derive exact identities. These "sum rules" carry clear physical interpretation as interrelations between forces when starting with spatial displacement, and between torques when starting with spatial rotations. Different types of identities result, depending on whether the elementary free energy is displaced (leading to external force sum rules), the excess free energy density functional (internal force identities) or the power functional (memory identities [27] that connect time direct correlation functions [31, 32]).

These invariant variational techniques have aided the development of a force-based approach to classical DFT [30]. Here the fundamental starting equation is the locally resolved equilibrium force balance relationship, which (for pairwise interparticle forces) is a classical result [1] that dates back to Yvon [33], and to Born and Green [34]. The derivation of this fundamental equation is performed by considering an inhomogeneous spatial displacement of the entire system, as described by a vector field  $\epsilon(\mathbf{r})$  in three-dimensional space. Together with a corresponding change of momenta (described in detail below) the change of variables constitutes a canonical transformation on classical phase space, and hence it preserves the phase space volume element [35]. The specific form of the transformation (in particular it being independent of time) also preserves the Hamiltonian. Hence the partition sum itself is unchanged under the transformation and so is the free energy. (The Hamiltonian, via its associated Boltzmann factor, and the phase space integral are the only nontrivial ingredients in the partition sum.) One is hence faced with an invariant variational problem, as addressed succinctly by Emmy Noether in her classical work [25]; see reference [26] for a historical account.

In the present contribution, we demonstrate that Noether's theorem is applicable to the equilibrium statistical properties of quantum many-body systems. We present a quantal shifting transformation of the position and momentum operators that reduces to the transformation of reference [30] in the classical case. Quantum mechanically, the transformation is canonical [36], i.e. it preserves the fundamental commutator relation between position and momentum. Such transformations represent unitary transformations on the Hilbert space of the considered system. The partition sum is hence invariant under the transformation, as it is given as the trace of the Hamiltonian's Boltzmann factor, with both the trace and the Hamiltonian being invariants, as is the case classically. The result, to first order in the displacement field, is the locally resolved equilibrium force balance relationship [3–5]. While one could expect on general grounds that the Noether line of thought would indeed apply to quantum systems, the details of the derivation differ markedly from the classical case, and we spell out the details in the following.

### II. THEORY

We consider Hamiltonians of the form

$$H = \sum_{i} \frac{\mathbf{p}_{i}^{2}}{2m} + u(\mathbf{r}^{N}) + \sum_{i} V_{\text{ext}}(\mathbf{r}_{i}), \tag{1}$$

where the sums run over all particles  $i=1,\ldots,N$ , with the total number of particles N. All particles possess identical mass m and each particle i is characterized by its position  $(\mathbf{r}_i)$  and momentum  $(\mathbf{p}_i)$  operator. The interparticle interaction potential  $u(\mathbf{r}^N)$  depends on all particle positions and we use the compact notation  $\mathbf{r}^N \equiv \mathbf{r}_1, \ldots, \mathbf{r}_N$ . The system is under the influence of an external one-body potential  $V_{\text{ext}}(\mathbf{r})$ , where  $\mathbf{r}$  is a generic position variable.

Position and momentum satisfy the fundamental commutator relations

$$[\mathbf{r}_i, \mathbf{p}_j] = i\hbar \delta_{ij} \mathbb{1}, \tag{2}$$

where i is the imaginary unit,  $\delta_{ij}$  is the Kronecker symbol and 1 indicates the  $3 \times 3$ -unit matrix. The commutator of two vectors involves transposition according to  $[\mathbf{r}_i, \mathbf{p}_j] = \mathbf{r}_i \mathbf{p}_j - \mathbf{p}_j \mathbf{r}_i^\mathsf{T}$ , where the superscript T denoting the transpose of a  $3 \times 3$ -matrix; the multiplication of two vectors is dyadic. Using Greek superscripts  $\alpha, \gamma$  to denote Cartesian components of position and momentum, Eq. (2) reads as  $[r_i^{\alpha}, p_j^{\gamma}] = i\hbar \delta_{ij} \delta_{\alpha\gamma}$ . We work in position representation, such that the momentum operator of particle i is given by  $\mathbf{p}_i = -i\hbar \nabla_i$ , where  $\nabla_i$  indicates the derivative with respect to  $\mathbf{r}_i$ .

We consider the following transformation of position and momenta

$$\mathbf{r}_{i} \to \mathbf{r}_{i} + \boldsymbol{\epsilon}(\mathbf{r}_{i}), \tag{3}$$

$$\mathbf{p}_{i} \to \left\{ (\mathbb{1} + (\nabla_{i}\boldsymbol{\epsilon}_{i}))^{-1} \cdot \mathbf{p}_{i} + \mathbf{p}_{i} \cdot (\mathbb{1} + (\nabla_{i}\boldsymbol{\epsilon}_{i})^{\mathsf{T}})^{-1} \right\} / 2, \tag{4}$$

where  $\epsilon(\mathbf{r}_i)$  is a given real-valued three-dimensional vector field,  $\epsilon_i = \epsilon(\mathbf{r}_i)$  is a shorthand notation, and the superscript -1 indicates matrix inversion. In Eq. (4) the gradient operator  $\nabla_i$  acts only on  $\epsilon_i$ , as is indicated by the surrounding parentheses; hence in position representation  $(\nabla_i \epsilon_i)$  acts only as a multiplication operator on the wave function.

We expand the inverse matrix in Eq. (4) to linear order in the gradient of the displacement field according to:  $(\mathbb{1} + (\nabla_i \epsilon_i))^{-1} = \mathbb{1} - (\nabla_i \epsilon_i)$ , where terms of the order  $(\nabla_i \epsilon_i)^2$  and higher have been omitted. Using this expansion, Eq. (4) in component notation is:  $p_i^{\alpha} \to p_i^{\alpha} - \sum_{\gamma} \{(\nabla_i^{\alpha} \epsilon_i^{\gamma}) p_i^{\gamma} + p_i^{\gamma} (\nabla_i^{\alpha} \epsilon_i^{\gamma})\}/2$ , which when resorting back to vector notation is:

$$\mathbf{p}_i \to \mathbf{p}_i - \{(\nabla_i \boldsymbol{\epsilon}_i) \cdot \mathbf{p}_i + \mathbf{p}_i \cdot (\nabla_i \boldsymbol{\epsilon}_i)^\mathsf{T}\}/2.$$
 (5)

We first ascertain that the new coordinates  $\tilde{\mathbf{r}}_i$  and new momenta  $\tilde{\mathbf{p}}_i$ , as defined by the right hand sides of the

transformation (3) and (4), also satisfy canonical commutation relations. We start with the prominent case:

$$[\tilde{\mathbf{r}}_{i}, \tilde{\mathbf{p}}_{j}] = [\mathbf{r}_{i} + \boldsymbol{\epsilon}_{i}, \\ \mathbf{p}_{j} - \{(\nabla_{j}\boldsymbol{\epsilon}_{j}) \cdot \mathbf{p}_{j} + \mathbf{p}_{j} \cdot (\nabla_{j}\boldsymbol{\epsilon}_{j})^{\mathsf{T}}\}/2]$$
(6)  
$$= [\mathbf{r}_{i}, \mathbf{p}_{j}] + [\boldsymbol{\epsilon}_{i}, \mathbf{p}_{j}]$$
$$- [\mathbf{r}_{i}, (\nabla_{j}\boldsymbol{\epsilon}_{j}) \cdot \mathbf{p}_{j}]/2 - [\mathbf{r}_{i}, \mathbf{p}_{j} \cdot (\nabla_{j}\boldsymbol{\epsilon}_{j})^{\mathsf{T}}]/2$$
(7)  
$$= i\hbar\delta_{ij}\mathbb{1},$$
(8)

where we have truncated in (7) at linear order in the displacement field and its gradient. As the left hand side of (6) involves no coupling between different particles, it is straightforward to see that for distinct particles,  $i \neq j$ , the result vanishes, as is indeed the case in Eq. (8). For i = j we use the explicit form of the momentum operator  $\mathbf{p}_i = -i\hbar\nabla_i$  to find that the second term in (7) is  $[\boldsymbol{\epsilon}_i, \mathbf{p}_i] = i\hbar(\nabla_i \boldsymbol{\epsilon}_i)$ . This contribution is precisely cancelled by the sum of the third and the fourth term in (7), which can be shown to have the form  $-[\mathbf{r}_i, (\nabla_i \boldsymbol{\epsilon}_i) \cdot \mathbf{p}_i] = -(\nabla_i \boldsymbol{\epsilon}_i) \cdot (i\hbar \mathbf{1}) = -i\hbar(\nabla_i \boldsymbol{\epsilon}_i)$ . Hence the first term in (7) alone gives the result (8) upon using the fundamental commutator (2).

For completeness, the new variables also satisfy  $[\tilde{\mathbf{r}}_i, \tilde{\mathbf{r}}_j] = 0$  and  $[\tilde{\mathbf{p}}_i, \tilde{\mathbf{p}}_j] = 0$ . The former relationship is trivial, as in position representation only coordinates are involved according to the transformation (3). The momentum identity can be worked out straightforwardly as is shown in appendix A. Furthermore the new degrees of freedom are self-adjoint operators. For the positions this is trivial, as we we have  $\tilde{\mathbf{r}}_i^{\dagger} = \mathbf{r}_i^* + \epsilon(\mathbf{r}_i)^* = \mathbf{r}_i + \epsilon(\mathbf{r}_i) \equiv \tilde{\mathbf{r}}_i$ , because  $\epsilon(\mathbf{r})$  is real-valued. For the momenta:  $\tilde{\mathbf{p}}_i^{\dagger} = \mathbf{p}_i^{\dagger} - \{(\nabla_i \epsilon_i) \cdot \mathbf{p}_i + \mathbf{p}_i \cdot (\nabla_i \epsilon_i)^{\mathsf{T}})\}^{\dagger}/2 = \mathbf{p}_i - \{\mathbf{p}_i \cdot (\nabla_i \epsilon_i)^{\mathsf{T}} + (\nabla_i \epsilon_i) \cdot \mathbf{p}_i\}/2 \equiv \tilde{\mathbf{p}}_i$ .

Having ascertained that the new variables form a sound basis for the description of the many-body system, we wish to illustrate the effect of the transformation on the system. The following considerations will be an essential ingredient in the thermal physics addressed further below. We wish to investigate the effect on the Hamiltonian  $H[\epsilon]$ , which is obtained by applying the operator replacements (3) and (4) in the form (1). We consider the functional derivative of the transformed Hamiltonian with respect to the displacement field:

$$\frac{\delta H[\boldsymbol{\epsilon}]}{\delta \boldsymbol{\epsilon}(\mathbf{r})} = \frac{\delta}{\delta \boldsymbol{\epsilon}(\mathbf{r})} \Big( \sum_{i} \frac{\tilde{\mathbf{p}}_{i}^{2}}{2m} + u(\tilde{\mathbf{r}}^{N}) + \sum_{i} V_{\text{ext}}(\tilde{\mathbf{r}}_{i}) \Big). \quad (9)$$

To make progress, we first require the fundamental derivatives of the new position and new momentum with respect to the displacement field. These are easily obtained as follows:

$$\frac{\delta \tilde{\mathbf{r}}_i}{\delta \epsilon(\mathbf{r})}\Big|_{\epsilon=0} = \delta(\mathbf{r} - \mathbf{r}_i)\mathbb{1},\tag{10}$$

$$\frac{\delta \tilde{\mathbf{p}}_i}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0} = \nabla \{ \delta(\mathbf{r} - \mathbf{r}_i) \mathbf{p}_i + \mathbf{p}_i \delta(\mathbf{r} - \mathbf{r}_i) \} / 2, \quad (11)$$

where  $\delta(\cdot)$  denotes the (three-dimensional) Dirac distribution and the derivatives are taken at vanishing displacement field  $\epsilon(\mathbf{r})=0$ , as is indicated in the notation on both left hand sides. The right hand side of Eq. (10) constitutes the density operator of particle i times the unit matrix. The right hand side of Eq. (11) is the spatial gradient of the momentum density operator of particle i. That both correctly localized operators appear naturally as functional derivatives is an initial indication that the considered transformation indeed can be used as a successful probe for the spatially resolved behaviour of the system.

For completeness, in index notation Eq. (11) reads as

$$\delta \tilde{p}_i^{\alpha} / \delta \epsilon^{\gamma} = \nabla^{\alpha} (\delta_i p_i^{\gamma} + p_i^{\gamma} \delta_i) / 2, \tag{12}$$

where we have introduced the shorthand notations  $\delta_i = \delta(\mathbf{r} - \mathbf{r}_i)$  and  $\epsilon^{\gamma} = \epsilon^{\gamma}(\mathbf{r})$  and the derivative is again evaluated at vanishing displacement field (such that higher than linear powers in the displacement gradient as they occur in the *finite* momentum transformation (4) vanish).

In order to obtain the functional derivative (9) of the Hamiltonian we proceed by first differentiating the kinetic energy, which reads as

$$\frac{\delta}{\delta \epsilon(\mathbf{r})} \sum_{i} \frac{\tilde{\mathbf{p}}_{i}^{2}}{2m} \Big|_{\epsilon=0} = \nabla \cdot \sum_{i} \frac{\mathbf{p}_{i} \delta_{i} \mathbf{p}_{i} + \mathbf{p}_{i} \delta_{i} \mathbf{p}_{i}^{\mathsf{T}}}{2m} - \frac{\hbar^{2}}{4m} \nabla \nabla^{2} \sum_{i} \delta(\mathbf{r} - \mathbf{r}_{i}), \quad (13)$$

where  $\nabla^2$  is the Laplace operator. That Eq. (13) holds is perhaps most easily seen in component notation. We use Einstein summation convention over pairs of Greek indices and after taking the functional derivative set  $\epsilon(\mathbf{r}) =$ 0 throughout. We consider particle i only, for which we have  $\delta(\tilde{p}_i^{\alpha}\tilde{p}_i^{\alpha})/\delta\epsilon^{\gamma} = p_i^{\alpha}\delta\tilde{p}_i^{\alpha}/\delta\epsilon^{\gamma} + \delta\tilde{p}_i^{\alpha}/\delta\epsilon^{\gamma}p_i^{\alpha}$ . The first term on the right hand side is  $p_i^{\alpha} \delta \tilde{p}_i^{\alpha} / \delta \epsilon^{\gamma} = \tilde{\nabla}^{\alpha} (p_i^{\alpha} \delta_i p_i^{\gamma} +$  $p_i^{\alpha} p_i^{\gamma} \delta_i)/2 = \nabla^{\alpha} \{ p_i^{\alpha} \delta_i p_i^{\gamma} + p_i^{\alpha} (p_i^{\gamma} \delta_i) + p_i^{\alpha} \delta_i p_i^{\gamma} \}/2, \text{ where}$ we have used Eq. (12) in the first equality and the product rule of differentiation for the application of  $p_i^{\gamma}$ in the second equality. The remaining second term is  $\frac{\delta \tilde{p}_{i}^{\alpha}/\delta \epsilon^{\gamma} p_{i}^{\alpha}}{\delta \tilde{p}_{i}^{\alpha}/\delta \epsilon^{\gamma} p_{i}^{\alpha}} = \nabla^{\alpha}(\delta_{i} p_{i}^{\gamma} p_{i}^{\alpha} + p_{i}^{\gamma} \delta_{i} p_{i}^{\alpha})/2 = \nabla^{\alpha}\{p_{i}^{\gamma} \delta_{i} p_{i}^{\alpha} - (p_{i}^{\gamma} \delta_{i}) p_{i}^{\alpha} + p_{i}^{\gamma} \delta_{i} p_{i}^{\alpha}\}/2.$  The appearance of the minus sign allows to carry out the following cancellation of the respective "middle" terms:  $p_i^{\alpha}(p_i^{\gamma}\delta_i) - (p_i^{\gamma}\delta_i)p_i^{\alpha} = (p_i^{\alpha}p_i^{\gamma}\delta_i) = -\hbar^2(\nabla_i^{\alpha}\nabla_i^{\gamma}\delta_i) = -\hbar^2\nabla^{\alpha}\nabla^{\gamma}\delta_i$ , where in the second step we have rewritten the action of the momentum operator on the delta function by the (negative) position gradient  $(p_i^{\gamma}\delta_i) = -i\hbar(\nabla_i^{\gamma}\delta_i) = i\hbar(\nabla^{\gamma}\delta_i)$ .

Collecting all terms we obtain the overall result for kinetic energy:

$$\frac{\delta}{\delta\epsilon^{\gamma}} \sum_{i} \frac{\tilde{p}_{i}^{\alpha} \tilde{p}_{i}^{\alpha}}{2m} \Big|_{\epsilon=0} = \nabla^{\alpha} \Big\{ \sum_{i} \frac{p_{i}^{\alpha} \delta_{i} p_{i}^{\gamma} + p_{i}^{\gamma} \delta_{i} p_{i}^{\alpha}}{2m} - \frac{\hbar^{2}}{4m} \nabla^{\gamma} \nabla^{\alpha} \sum_{i} \delta_{i} \Big\}$$
(14)

$$= -\nabla^{\alpha} \hat{\tau}^{\alpha \gamma}, \tag{15}$$

where, as desired, Eq. (14) is the  $\gamma$ th Cartesian component of Eq. (13), expressed with tensor contractions and transpositions in index notation. In Eq. (15) we have identified the position-resolved one-body kinematic stress operator  $\hat{\tau}(\mathbf{r})$  (see e.g. Ref. [5]). We have adopted the convention to include the wave-like contribution  $\hbar^2 \nabla \nabla \hat{\rho}(\mathbf{r})/(4m)$  into the kinematic stress, where  $\hat{\rho}(\mathbf{r}) = \sum_i \delta(\mathbf{r} - \mathbf{r}_i)$  is the standard form of the one-body density operator. As an aside, the classical case is recovered by letting  $\hbar \to 0$ , such that  $\hat{\tau}(\mathbf{r})$  reduces to  $-\sum_i \delta(\mathbf{r} - \mathbf{r}_i) \mathbf{p}_i \mathbf{p}_i / m$ , where here  $\mathbf{p}_i$  denotes the classical phase space variable, which trivially commutes with the spatial delta distribution, and the transpose becomes irrelevant as  $\mathbf{p}_i \mathbf{p}_i = \mathbf{p}_i \mathbf{p}_i^{\mathsf{T}}$ .

We have so far shown that the considered quantum canonical transformation the functional derivative of kinetic energy with respect to the displacement field creates the following simple result:

$$\frac{\delta H_{\text{kin}}[\epsilon]}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0} = -\nabla \cdot \hat{\boldsymbol{\tau}}(\mathbf{r}). \tag{16}$$

Here we have split the Hamiltonian (1) according to  $H = H_{\rm kin} + H_{\rm pot}$ , where the potential energy contains the interparticle and external contributions,  $H_{\rm pot} = u(\mathbf{r}^N) + \sum_i V_{\rm ext}(\mathbf{r}_i)$ . As already laid out above, the functional dependence on  $\epsilon(\mathbf{r})$  that is indicated on the left hand side of (16) arises from expressing the original positions and momenta in the Hamiltonian (1) via the transformation (3) and (4).

Treating the effects of the local displacement transformation on the potential energy is comparatively easier than the above kinetic energy consideration, as here only position coordinates are involved and hence the commutator structure is trivial. The calculation is very closely analogous to the classical case [28]. We obtain

$$\frac{\delta H_{\text{pot}}[\epsilon]}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0} = \frac{\delta u(\tilde{\mathbf{r}}^N)}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0} + \sum_{i} \frac{\delta V_{\text{ext}}(\tilde{\mathbf{r}}_i)}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0}$$

$$= \sum_{i} (\nabla_i u(\mathbf{r}_N)) \delta_i + \sum_{i} (\nabla_i V_{\text{ext}}(\mathbf{r}_i)) \delta_i$$
(18)

$$= -\hat{\mathbf{F}}_{\text{int}}(\mathbf{r}) + \hat{\rho}(\mathbf{r})\nabla V_{\text{ext}}(\mathbf{r}), \tag{19}$$

where we have defined the one-body interparticle force density operator  $\hat{\mathbf{F}}_{\mathrm{int}}(\mathbf{r}) = -\sum_i (\nabla_i u(\mathbf{r}^N)) \delta_i$ . The rewriting that involves the external force field  $-\nabla V_{\mathrm{ext}}(\mathbf{r})$  in (19) is possible as the derivatives  $\nabla_i$  and  $\nabla$ , as well as the positions  $\mathbf{r}$  and  $\mathbf{r}_i$ , can be identified with each other due to the presence of the delta function. The negative external force field  $\nabla V_{\mathrm{ext}}(\mathbf{r})$  can then be taken as a common factor outside of the second sum in Eq. (18) and the density operator remains.

Summing up the kinetic energy derivative (16) and the potential energy identity (19) we obtain

$$-\frac{\delta H[\boldsymbol{\epsilon}]}{\delta \boldsymbol{\epsilon}(\mathbf{r})}\Big|_{\boldsymbol{\epsilon}=0} = \nabla \cdot \hat{\boldsymbol{\tau}}(\mathbf{r}) + \hat{\mathbf{F}}_{int}(\mathbf{r}) - \hat{\rho}(\mathbf{r})\nabla V_{ext}(\mathbf{r}), \quad (20)$$

which makes explicit that the Hamiltonian generates, via its negative functional derivative with respect to the displacement field, the sum of all one-body force density distributions that act in the system.

That the transformation (3) and (4) has an effect on the Hamiltonian could have been expected from the outset, as the transformation has a nontrivial spatial structure via its dependence on the vector field  $\epsilon(\mathbf{r})$ . Hence Noether's theorem seemingly does not apply, due to the absence of a direct corresponding invariance. In contrast to this standard application, here we proceed differently and search for an invariance that applies in thermal equilibrium. This requires an average to be an invariant rather than the corresponding operator itself being an invariant, as we lay out in the following.

We hence turn to a statistical mechanical description which we base on the free energy in the canonical ensemble, expressed as

$$F = -k_B T \ln Z,\tag{21}$$

$$Z = \operatorname{Tr} e^{-\beta H} \tag{22}$$

$$= \sum_{n} \langle n|e^{-\beta H}|n\rangle, \tag{23}$$

where  $k_B$  denotes the Boltzmann constant and T is absolute temperature. The trace in Hilbert space is denoted by Tr and it is made explicit in (23) with  $|n\rangle$  denoting the complete set of orthonormal eigenstates of H labelled by index n. (Possibly degenerate energy eigenstates occur multiple times in the sum.) In more explicit notation, using position representation,  $|n\rangle = \phi_n(\mathbf{r}^N)$  such that  $\langle n|\cdot|n\rangle = \int d\mathbf{r}^N\phi_n^*(\mathbf{r}^N)\cdot\phi_n(\mathbf{r}^N)$ , where the integral is over all position coordinates,  $\int d\mathbf{r}^N \cdot = \int d\mathbf{r}_1 \int d\mathbf{r}_2 \dots \int d\mathbf{r}_N \cdot$  and the asterisk denotes complex conjugation.

We expand the free energy (21) in the transformation parameter according to:

$$\frac{\delta F[\epsilon]}{\delta \epsilon(\mathbf{r})}\Big|_{\epsilon=0} = -\frac{k_B T}{Z} \frac{\delta Z[\epsilon]}{\delta \epsilon(\mathbf{r})}\Big|_{\epsilon=0}$$
 (24)

$$= -\frac{k_B T}{Z} \operatorname{Tr} \frac{\delta e^{-\beta H[\epsilon]}}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0}$$
 (25)

$$= \sum_{n} \frac{e^{-\beta E_{n}}}{Z} \langle n | \frac{\delta H[\epsilon]}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0} |n\rangle, \quad (26)$$

where in Eq. (24) we have used the definition (21) of the free energy via the partition sum  $Z[\epsilon]$  of the transformed system. In Eq. (25) we have used the form (22) of the partition sum and have exchanged the order of the functional derivative and building the trace. Equation (26) constitutes a thermal equilibrium average, where  $E_n$  denotes the energy eigenvalue corresponding to the energy eigenstate  $|n\rangle$ . We have hence obtained

$$\frac{\delta F[\epsilon]}{\delta \epsilon(\mathbf{r})}\Big|_{\epsilon=0} = \left\langle \frac{\delta H[\epsilon]}{\delta \epsilon(\mathbf{r})}\Big|_{\epsilon=0} \right\rangle_{\text{eq}},\tag{27}$$

where on the right hand side we have used the notation  $\langle \cdot \rangle_{eq}$  to indicate the average over the canonical ensemble as it occurs in Eq. (26); explicitly this is  $\langle \cdot \rangle_{eq}$  =

 $\sum_{n} Z^{-1} e^{-\beta E_n} \langle n| \cdot |n\rangle$ . The identity (27) is remarkable as it indicates that the local transformation (3) and (4) to lowest order in the displacement field generates a well-defined and physically meaningful thermal average, that of the functional derivative of the Hamiltonian. This mathematical structure mirrors closely that of standard derivatives of the free energy.

Before exploiting the specific form of the right hand side of Eq. (27) further, we first proceed with the general invariance argument. We expand the free energy of the transformed system to linear order in the displacement field according to:

$$F[\epsilon] = F + \int d\mathbf{r} \frac{\delta F[\epsilon]}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0} \cdot \epsilon(\mathbf{r})$$
 (28)

$$= F + \int d\mathbf{r} \left\langle \frac{\delta H[\boldsymbol{\epsilon}]}{\delta \boldsymbol{\epsilon}(\mathbf{r})} \Big|_{\boldsymbol{\epsilon}=0} \right\rangle_{\text{eq}} \cdot \boldsymbol{\epsilon}(\mathbf{r}), \qquad (29)$$

where Eq. (28) is the functional Taylor expansion to linear order and the form (29) follows from using Eq. (27).

On the other hand, the free energy is an invariant under the quantum canonical transformation, and hence:

$$F[\epsilon] = F,\tag{30}$$

where F is the free energy (21) of the original representation of the system. Equation (30) holds due to the fact that canonical transformations are analogous to unitary transformations on the underlying Hilbert space of the considered system; see e.g. the account given by Anderson [36]. We will return to this point below.

From comparison of the Taylor expansion (29) with the free energy invariance (30) we can conclude that the linear term in the expansions vanishes identically and it has to do so irrespective of the form of  $\epsilon(\mathbf{r})$ . This can only hold provided that the prefactor vansishes:

$$\left\langle \frac{\delta H[\epsilon]}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0} \right\rangle_{\text{eq}} = 0.$$
 (31)

Equation (31) is a bare consequence of the invariance of the free energy under the displacement operation. The identity holds irrespective of the precise form of the interparticle interaction potential  $u(\mathbf{r}^N)$  and of the external potential  $V_{\text{ext}}(\mathbf{r})$  as they appear in the Hamiltonian (1).

In order to reveal the physical significance of the Noether sum rule (31) we proceed by inserting the explicit force form (20) of the functional derivative of the Hamiltonian, which yields

$$\nabla \cdot \boldsymbol{\tau}(\mathbf{r}) + \mathbf{F}_{\text{int}}(\mathbf{r}) - \rho(\mathbf{r}) \nabla V_{\text{ext}}(\mathbf{r}) = 0.$$
 (32)

Here we have introduced the equilibrium averages for the locally resolved kinetic stress:  $\mathbf{r}(\mathbf{r}) = \langle \hat{\mathbf{r}}(\mathbf{r}) \rangle_{\rm eq}$ , for the interparticle force density:  $\mathbf{F}_{\rm int}(\mathbf{r}) = \langle \hat{\mathbf{F}}_{\rm int}(\mathbf{r}) \rangle_{\rm eq}$ , and for the one-body density distribution:  $\rho(\mathbf{r}) = \langle \hat{\rho}(\mathbf{r}) \rangle_{\rm eq}$ . The force density balance relationship (32) is a known exact equilibrium sum rule, see e.g. Refs. [3–5]. Our derivation demonstrates its origin in the invariance of the free

energy under the quantum canonical transformation (3) and (4).

As a special case we consider a uniform displacement such that  $\epsilon(\mathbf{r}) = \epsilon_0 = \text{const.}$  For classical systems the invariance of the free energy under such homogeneous displacement leads to the sum rule of vanishing global external force in thermal equilibrium [27, 28]. This result readily translates to the quantum case as follows.

First we obtain the global identity by starting with the locally resolved force balance relationship (32) and integrating over all positions. Two of the resulting integrals vanish,  $\int d\mathbf{r} \nabla \cdot \boldsymbol{\tau}(\mathbf{r}) = 0$  and  $\int d\mathbf{r} \mathbf{F}_{int}(\mathbf{r}) = 0$ , where the former identity can be shown via integration by parts and the latter identity is a consequence of the translational invariance of the interparticle interaction potential:  $\int d\mathbf{r} \mathbf{F}_{int}(\mathbf{r}) = \int d\mathbf{r} \langle \hat{\mathbf{F}}_{int}(\mathbf{r}) \rangle_{eq} = \langle \int d\mathbf{r} \hat{\mathbf{F}}_{int}(\mathbf{r}) \rangle_{eq} =$  $\langle \hat{\mathbf{F}}_{\text{int}}^{\text{o}} \rangle_{\text{eq}} = 0$ . This holds due to the global force operator vanishing identically:  $\hat{\mathbf{F}}_{\mathrm{int}}^{0} = -\sum_{i} (\nabla_{i} u(\mathbf{r}_{N})) \equiv 0$ , which can be seen straightforwardly by displacing all positions arguments in  $u(\mathbf{r}^N)$  and observing that this leaves its value invariant. Explicitly the invariance is  $u(\mathbf{r}_1,\ldots,\mathbf{r}_N)=u(\mathbf{r}_1+\boldsymbol{\epsilon}_0,\ldots,\mathbf{r}_N+\boldsymbol{\epsilon}_0),$  as the global shift leaves all distance vectors  $\mathbf{r}_i - \mathbf{r}_j$  unchanged. The Taylor expansion of the right hand side is to first order  $u(\mathbf{r}_1, \dots, \mathbf{r}_N) + \epsilon_0 \cdot \partial u(\mathbf{r}_1 + \epsilon_0, \dots, \mathbf{r}_N + \epsilon_0) / \partial \epsilon_0 |_{\epsilon_0 = 0}$ . The linear term can be rewritten as  $\epsilon_0 \cdot \sum_i \nabla_i u(\mathbf{r}^N)$ . As this vanishes for any  $\epsilon_0$ , the prefactor vanishes identically which provides the anticipated vanishing of the global interparticle force. This reasoning is analogous to Newton's third law, actio equals reactio, which holds due to the interparticle forces being conservative.

The only term that remains of Eq. (32) after carrying out the position integral is the external force contribution, which reads as:

$$-\int d\mathbf{r}\rho(\mathbf{r})\nabla V_{\text{ext}}(\mathbf{r}) = 0. \tag{33}$$

Equation (33) expresses the vanishing of the average global external force in thermal equilibrium.

Briefly, in our second route to Eq. (33) we start from the free energy (21) and directly perform the transformation for the special case of a homogeneous displacement  $\epsilon_0$ . In this case the momenta are unchanged, as the gradient of the (constant) displacement field vanishes identically. Hence kinetic energy is trivially invariant. As laid out above, the coordinate change does not affect the interparticle potential energy, as the difference vectors are unaffected. Hence the only change in the Hamiltonian occurs in the external contribution and we obtain, following the same argumentation as in the case of position-dependent shifting, the result

$$-\left\langle \sum_{i} \nabla_{i} V_{\text{ext}}(\mathbf{r}_{i}) \right\rangle_{\text{eq}} = 0, \tag{34}$$

which is analogous to the previous form (33) upon prepending  $1 = \int d\mathbf{r} \delta(\mathbf{r} - \mathbf{r}_i)$  to Eq. (34), moving the

delta function into the thermal average, and identifying the density profile  $\rho(\mathbf{r}) = \langle \sum_i \delta(\mathbf{r} - \mathbf{r}_i) \rangle_{\text{eq}}$ .

For completeness and as a final step, we make explicit that the quantum canonical transformation corresponds indeed to a unitary transformation on Hilbert space, as is relevant for the invariance (30) of the free energy under the transformation. For the present transformation, to linear order in  $\epsilon(\mathbf{r})$ , the transformed Hamiltonian is obtained via functional Taylor expansion in the following form:

$$H[\epsilon] = H + \int d\mathbf{r} \frac{\delta H[\epsilon]}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0} \cdot \epsilon(\mathbf{r}),$$
 (35)

and we recall the explicit one-body force density form (20) of the functional derivative of the Hamiltonian. We treat the second term in Eq. (35) as a perturbation to the original Hamiltonian H. Then the transformed (perturbed) energy eigenstates  $|\tilde{n}\rangle$  are given by

$$|\tilde{n}\rangle = \sum_{k} U_{nk} |k\rangle, \tag{36}$$

$$U_{nk} = \delta_{nk} + \frac{1 - \delta_{nk}}{E_k - E_n} \int d\mathbf{r} \langle n | \frac{\delta H[\epsilon]}{\delta \epsilon(\mathbf{r})} \Big|_{\epsilon=0} |k\rangle \cdot \epsilon(\mathbf{r}).$$
 (37)

The form (37) of the matrix that performs the change of basis follows from applying time-independent first order perturbation theory, as is appropriate to capture the effects to linear order in  $\epsilon(\mathbf{r})$  that we consider. (We imply that the prefactor of the integral in Eq. (37) vanishes for k = n.) The matrix elements of the Hermitian conjugate to Eq. (37) can be obtained via exchanging indices n and k as

$$U_{nk}^{\dagger} = \delta_{kn} + \frac{1 - \delta_{kn}}{E_n - E_k} \int d\mathbf{r} \langle k | \frac{\delta H[\boldsymbol{\epsilon}]}{\delta \boldsymbol{\epsilon}(\mathbf{r})} \Big|_{\boldsymbol{\epsilon} = 0}^{\dagger} |n\rangle \cdot \boldsymbol{\epsilon}(\mathbf{r}) \quad (38)$$
$$= \delta_{nk} - \frac{1 - \delta_{nk}}{E_k - E_n} \int d\mathbf{r} \langle n | \frac{\delta H[\boldsymbol{\epsilon}]}{\delta \boldsymbol{\epsilon}(\mathbf{r})} \Big|_{\boldsymbol{\epsilon} = 0} |k\rangle \cdot \boldsymbol{\epsilon}(\mathbf{r}), \quad (39)$$

where to obtain the matrix elements (39) we have exploited that the functional derivative of the Hamiltonian is self-adjoint. We observe that the sole difference between Eqs. (37) and (39) is the minus sign. Hence we can see explicitly that unitarity holds,  $\sum_k U_{nk}^{\dagger} U_{km} = \delta_{nm}$  to linear order in  $\epsilon(\mathbf{r})$ , as was expected on general grounds [36].

## III. OUTLOOK AND CONCLUSIONS

In conclusion we have investigated the consequences of a specific local displacement operation for the free energy of a quantum mechanical many-body system. The transformation consists of position-dependent shifting, as parameterized by a real-valued strain field, and a corresponding transformation of the quantum mechanical momentum operator of each particle. The entirety of the transformation can be viewed as the self-adjoint version of the corresponding local shifting transformation of the classical phase space variables [30]. We have explicitly shown that the new position and momentum operators are self-adjoint and that they satisfy the fundamental commutator relations and hence form a valid and complete set of degrees of freedom of the considered system. The transformation can be viewed as a basis change of the underlying Hilbert space of the quantal system and we have spelled out explicitly the corresponding unitary transformation between the original and the new basis.

The resulting invariance of the free energy under changes in the displacement field then leads, following Noether's theorem for invariant variations, to an exact local identity ("sum rule") which we have shown to be the thermal equilibrium force balance. The present derivation of this known and fundamental result from Noether's theorem sheds new light on the very nature of the identity. Existing derivations are based e.g. on the second time derivative of the one-body density profile [3] or, equivalently, on the first time derivative of the one-body current distribution [5] and then taking the equilibrium limit.

Our results hold for the ground state state of the quantum system, as it is obtained in the limit  $T \to 0$  of the free energy of the thermal system. We have used the canonical ensemble throughout as it captures the essence of the required thermal physics. We expect the reasoning to carry over straightforwardly to the grand ensemble with fluctuating particle number, as the classical canonical [28] and grand canonical cases lead to analogous results upon identifying the respective statistical averages.

Future work could be addressed at investigating how further functional differentiation can be used to obtain quantum sum rules for higher-body correlation functions is an interesting topic, as previously shown for classical systems [27]. It would be interesting to address the effects beyond linear order in the displacement field; classically the variance of the global external force was shown to be constrained by the external potential energy curvature [29]. Last not least it would be worthwhile to find possible relationships of our displacement field and the strain field that is central to elasticity theory, see e.g. Refs. [37, 38] for recent work again in classical systems.

### Appendix A: Momentum commutator

To derive the commutator of the new momenta,  $[\tilde{\mathbf{p}}_i, \tilde{\mathbf{p}}_j]$ , we insert the definition of the transformation (5) and consider terms up to linear order in the displacement gradient. In index notation this reads as follows:

$$\begin{split} 2[\tilde{p}_{i}^{\alpha}, \tilde{p}_{j}^{\gamma}] &= 2[p_{i}^{\alpha}, p_{j}^{\gamma}] \\ &- [p_{i}^{\alpha}, (\nabla_{j}^{\gamma} \epsilon_{j}^{\delta}) p_{j}^{\delta}] - [p_{i}^{\alpha}, p_{j}^{\delta} (\nabla_{j}^{\gamma} \epsilon_{j}^{\delta})] \\ &- [(\nabla_{i}^{\alpha} \epsilon_{i}^{\delta}) p_{i}^{\delta}, p_{j}^{\gamma}] - [p_{i}^{\delta} (\nabla_{i}^{\alpha} \epsilon_{i}^{\delta}), p_{j}^{\gamma}]. \end{split} \tag{A1}$$

The correlator of the original momenta, as it appears in the first term on the right hand side, vanishes trivially,  $[p_i^{\alpha}, p_j^{\gamma}] = 0$ . This identity also allows to take the operators  $p_i^{\delta}$  and  $p_j^{\delta}$  out of the commutator in the remaining terms on the right hand side of Eq. (A1). We obtain

$$\begin{split} 2[\tilde{p}_{i}^{\alpha}, \tilde{p}_{j}^{\gamma}] &= -[p_{i}^{\alpha}, (\nabla_{j}^{\gamma} \epsilon_{j}^{\delta})] p_{j}^{\delta} - p_{j}^{\delta} [p_{i}^{\alpha}, (\nabla_{j}^{\gamma} \epsilon_{j}^{\delta})] \\ &+ [p_{i}^{\gamma}, (\nabla_{i}^{\alpha} \epsilon_{i}^{\delta})] p_{i}^{\delta} + p_{i}^{\delta} [p_{i}^{\gamma}, (\nabla_{i}^{\alpha} \epsilon_{i}^{\delta})], \end{split} \tag{A2}$$

where we have exploited the anti-symmetry of the commutator, [A, B] = -[B, A], for rewriting the third and the fourth term on the right hand side of Eq. (A2).

Writing out explicitly the commutator in the first contribution in Eq. (A2) yields  $[p_i^{\alpha}, (\nabla_j^{\gamma} \epsilon_j^{\delta})] = p_i^{\alpha} (\nabla_j^{\gamma} \epsilon_j^{\delta}) - (\nabla_j^{\gamma} \epsilon_j^{\delta}) p_i^{\alpha}$ . Hence the momentum operator only acts on the gradient of the displacement field,  $(p_i^{\alpha} \nabla_j^{\gamma} \epsilon_j^{\delta}) = -i\hbar(\nabla_i^{\alpha} \nabla_j^{\gamma} \epsilon_j^{\delta})$ , where as before the parentheses indicate that the derivative(s) only act on the displacement field and we have expressed the momentum operator in position representation. Analog manipulation of all remaining commutators in equation (A2) then yields

$$\frac{2i}{\hbar} [\tilde{p}_{i}^{\alpha}, \tilde{p}_{j}^{\gamma}] = -(\nabla_{i}^{\alpha} \nabla_{j}^{\gamma} \epsilon_{j}^{\delta}) p_{j}^{\delta} - p_{j}^{\delta} (\nabla_{i}^{\alpha} \nabla_{j}^{\gamma} \epsilon_{j}^{\delta}) 
+ (\nabla_{j}^{\gamma} \nabla_{i}^{\alpha} \epsilon_{i}^{\delta}) p_{i}^{\delta} + p_{i}^{\delta} (\nabla_{j}^{\gamma} \nabla_{i}^{\alpha} \epsilon_{i}^{\delta}).$$
(A3)

For  $i \neq j$  it is now straightforward to see that each term

on the right hand side of equation (A3) vanishes individually: As the displacement field  $\epsilon_i$  only depends on positions  $\mathbf{r}_i$ , derivatives with respect to  $\mathbf{r}_j$  vanish for  $i \neq j$ . For i = j the first and the third term, as well as the second and the fourth term, on the right hand side of Eq. (A3) cancel each other pairwise, as the derivatives  $\nabla_i^{\alpha}$  and  $\nabla_j^{\gamma}$  commute. Collecting the cases i = j and  $i \neq j$  the commutator of the new momentum operators hence vanishes,

$$[\tilde{p}_i^{\alpha}, \tilde{p}_i^{\gamma}] = 0, \tag{A4}$$

which ascertains that the position-dependent momentum transformation does not generate any spurious terms.

#### ACKNOWLEDGMENTS

This paper is dedicated to Sir Michael Berry on the occasion of his 80th birthday. MS is grateful for the manifold inspirations and reliable judgements, in scientific and other matters, that Michael provided over many years. SH and MS acknowledge useful discussions with Bob Evans and Daniel de las Heras on the topic of the present paper. This work is supported by the German Research Foundation (DFG) via Project No. 436306241.

- J. P. Hansen and I. R. McDonald, Theory of Simple Liquids, 4th ed. (Academic Press, London, 2013).
- [2] F. Reif, Statistical and Thermal Physics, 18th ed. (McGraw-Hill, New York, 2007).
- [3] W. Tarantino and C. A. Ullrich, A reformulation of time-dependent Kohn-Sham theory in terms of the second time derivative of the density, J. Chem. Phys. **154**, 204112 (2021).
- [4] M.-L. M. Tchenkoue, M. Penz, I. Theophilou, M. Ruggenthaler, and A. Rubio, Force balance approach for advanced approximations in density functional theories, J. Chem. Phys. 151, 154107 (2019).
- [5] M. Schmidt, Power functional theory for many-body dynamics, Rev. Mod. Phys. 94, 015007 (2022).
- [6] M. Schmidt and J. M. Brader, Power functional theory for Brownian dynamics, J. Chem. Phys. 138, 214101 (2013).
- [7] M. Schmidt, Power functional theory for Newtonian many-body dynamics,
   J. Chem. Phys. 148, 044502 (2018).
- [8] M. Schmidt, Quantum power functional theory for manybody dynamics, J. Chem. Phys. 143, 174108 (2015).
- [9] M. Brütting, M., T. Trepl, D. de las Heras, and M. Schmidt, Superadiabatic forces via the acceleration gradient in quantum many-body dynamics, Molecules 24, 3660 (2019).
- [10] D. de las Heras and M. Schmidt, Velocity gradient power functional for Brownian dynamics, Phys. Rev. Lett. 120, 028001 (2018).
- [11] D. de las Heras and M. Schmidt, Flow and structure in nonequilibrium Brownian many-body systems,

- Phys. Rev. Lett. **125**, 018001 (2020).
- [12] J. Renner, M. Schmidt, and D. de las Heras, Shear and bulk acceleration viscosities in simple fluids, Phys. Rev. Lett. 128, 094502 (2022).
- [13] R. Evans, The nature of the liquid-vapour interface and other topics in the statistical mechanics of non-uniform, classical fluids, Adv. Phys. 28, 143 (1979).
- [14] R. Evans, 'Density functionals in the theory nonuniform fluids', in *Fundamentals of inhomogeneous fluids*, edited by D. Henderson (Dekker, New York, 1992).
- [15] For an overview of new developments in classical density functional theory, see: R. Evans, M. Oettel, R. Roth, and G. Kahl, New developments in classical density functional theory, J. Phys.: Condens. Matter 28, 240401 (2016).
- [16] U. M. B. Marconi and P. Tarazona, Dynamic density functional theory of fluids, J. Chem. Phys. 110, 8032 (1999).
- [17] A. J. Archer and R. Evans, Dynamical density functional theory and its application to spinodal decomposition, J. Chem. Phys. 121, 4246 (2004).
- [18] B. Rotenberg, Use the force! Reduced variance estimators for densities, radial distribution functions, and local mobilities in molecular simulations, J. Chem. Phys. 153, 150902 (2020).
- [19] D. de las Heras and M. Schmidt, Better than counting: Density profiles from force sampling, Phys. Rev. Lett. 120, 218001 (2018).
- [20] D. Borgis, R. Assaraf, B. Rotenberg, and R. Vuilleumier, Computation of pair distribution functions and threedimensional densities with a reduced variance principle, Mol. Phys. 111, 3486 (2013).

- [21] A. Purohit, A. J. Schultz, and D. A. Kofke, Forcesampling methods for density distributions as instances of mapped averaging, Mol. Phys. 117, 2822 (2019).
- [22] S. M. Tschopp and J. M. Brader, Fundamental measure theory of inhomogeneous two-body correlation functions, Phys. Rev. E 103, 042103 (2021).
- [23] R. J. McCarty, D. Perchak, R. Pederson, R. Evans, Y. Qiu, S. R. White, and K. Burke, Bypassing the energy functional in density functional theory: direct calculation of electronic energies from conditional probability densities, Phys. Rev. Lett. 125, 266401 (2020).
- [24] R. Pederson, J. Chen, S. R. White, and K. Burke, Conditional probability density functional theory, arxiv:2203.09647 (2022).
- [25] E. Noether, Invariante Variationsprobleme, Nachr. d. König. Gesellsch. d. Wiss. zu Göttingen, Math. English translation by M. A. Tavel: Invariant variation problems, Transp. Theo. Stat. Phys. 1, 186 (1971); for a version in modern typesetting see: Frank Y. Wang, arXiv:physics/0503066v3 (2018).
- [26] N. Byers, E. Noether's discovery of the deep connection between symmetries and conservation laws, arXiv:physics/9807044 (1998).
- [27] S. Hermann and M. Schmidt, Noether's theorem in statistical mechanics, Commun. Phys. 4, 176 (2021).
- [28] S. Hermann and M. Schmidt, Why Noether's face be theorem applies to statistical mechanics, J. Chen J. Phys.: Condens. Matter (invited Topical Review) 34, 213001 (2022).

- [29] S. Hermann and M. Schmidt, Variance of fluctuations from Noether invariance, Comm. Phys. (submitted); arXiv:2203.15654 (2022).
- [30] S. M. Tschopp, F. Sammüller, S. Hermann, M. Schmidt, and J. M. Brader, Force density functional theory in- and out-of-equilibrium, Phys. Rev. E 106, 014115 (2022).
- [31] J. M. Brader and M. Schmidt, Nonequilibrium Ornstein-Zernike relation for Brownian many-body dynamics, J. Chem. Phys. 139, 104108 (2013).
- [32] J. M. Brader, and M. Schmidt, Dynamic correlations in Brownian many-body systems, J. Chem. Phys. 140, 034104 (2014).
- [33] J. Yvon, Actualités Scientifiques et Industrielles, (Hermann & Cie., 1935).
- E. Noether, Invariante Variationsprobleme, [34] M. Born and H. S. Green, A general kinetic the-Nachr. d. König. Gesellsch. d. Wiss. zu Göttingen, Math.-Phys. Khassef 235u/d/918). The molecular distribution functions, English translation by M. A. Tavel: Invariant variation

  Proc. R. Soc. London, Ser. A 188, 10 (1946).
  - [35] H. Goldstein, C. Poole, and J. Safko, Classical Mechanics (Addison-Wesley, New York, 2002).
  - [36] A. Anderson, Canonical transformations in quantum mechanics, Ann. Phys. 232, 292 (1994).
  - [37] M. Sprik, Continuum model of the simple dielectric fluid: consistency between density based and continuum mechanics methods, Mol. Phys. 119, e1887950 (2021).
  - [38] M. Sprik, Chemomechanical equilibrium at the interface between a simple elastic solid and its liquid phase, J. Chem. Phys. 155, 244701 (2021).