

# A Scotogenic Model with Two Inert Doublets

---

**Amine Ahriche** 

*Department of Applied Physics and Astronomy, University of Sharjah, P.O. Box 27272 Sharjah, UAE.*

*The Abdus Salam International Centre for Theoretical Physics, Strada Costiera 11, I-34014, Trieste, Italy.*

*Laboratoire de Physique des Particules et Physique Statistique, Ecole Normale Supérieure, BP 92 Vieux Kouba, DZ-16050 Algiers, Algeria.*

*E-mail:* [ahriche@sharjah.ac.ae](mailto:ahriche@sharjah.ac.ae)

**ABSTRACT:** In this work, we present a scotogenic model, where the neutrino mass is generated at one-loop diagrams. The standard model (SM) is extended by three singlet Majorana fermions and two inert scalar doublets instead of one doublet as in the minimal scotogenic model. The model scalar sector includes two CP-even, two CP-odd and two charged scalars in addition to the Higgs. The dark matter (DM) candidate could be either the light Majorana fermion (Majorana DM), or the lightest among the CP-even and the CP-odd scalars (scalar DM). We show that the model accommodates both Majorana and scalar DM within a significant viable parameter space, while considering all the relevant theoretical and experimental constraints such as perturbativity, vacuum stability, unitarity, the di-photon Higgs decay, electroweak precision tests and lepton flavor violating constraints. In addition to the collider signatures predicted by the minimal scotogenic model, our model predicts some novel signatures that can be probed through some final states such as  $8 \text{ jets} + \cancel{E}_T$ ,  $1\ell + 4 \text{ jets} + \cancel{E}_T$  and  $4b + \cancel{E}_T$ .

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The Model &amp; Neutrino Mass</b>	<b>2</b>
<b>3</b>	<b>Theoretical &amp; Experimental Constraints</b>	<b>5</b>
<b>4</b>	<b>Viable Parameter Space</b>	<b>8</b>
<b>5</b>	<b>Dark Matter: Majorana or Scalar?</b>	<b>10</b>
<b>6</b>	<b>Conclusion</b>	<b>12</b>
<b>A</b>	<b>Generalized Casas-Ibarra parameterization</b>	<b>13</b>
<b>B</b>	<b>Scotogenic Model with <math>n</math> inert Doublets</b>	<b>14</b>
<b>C</b>	<b>The Unitarity Amplitude Matrices</b>	<b>14</b>

---

## 1 Introduction

The standard model (SM) of particle physics was successful in describing the properties of elementary particles and their interactions around the electroweak (EW) scale. However, there are still unanswered questions such as the neutrino oscillation data and the dark matter (DM) nature. Neutrino oscillation data established that at least two of the three SM neutrinos have mass and nonzero mixing. However, their properties, such as their nature and the origin of the smallness of their mass, have no explanation within the SM, which demands for new physics. One of the most popular mechanisms to explain why neutrino mass is tiny; is the so-called seesaw mechanism [1], which assumes the existence of right-handed (RH) neutrinos with masses that are heavier by several orders of magnitude than the EW scale. Such heavy particles decouple from the low-energy spectrum at the EW scale, and hence, cannot be directly probed at high-energy physics experiments. Another approach that requires lower new physics scale, where small neutrino mass is generated naturally; is via quantum corrections, where loop suppression factor(s) ensures the neutrino mass smallness. This can be realized at one loop [2], two loops [3], three loops [4, 5], or four loops [6]. Interesting features for this TeV scale new physics class of models can be tested at high energy colliders [7] (For a review, see [8]).

The simplest and most popular realization of a radiative neutrino mass mechanism is the so-called scotogenic model [9], where the SM is extended by an inert scalar doublet and three singlet Majorana fermions. In addition, this model could accommodate two possible DM candidates, i.e., the lightest

among the Majorana fermions and the light neutral scalar in the inert dark doublet. The scalar DM scenario is identical to the inert doublet Higgs model (IDM), that is highly constrained by the DM direct detection experiments. In the scenario of Majorana DM, the right amount of the DM relic density requires relatively large values for the new Yukawa couplings that couple the inert doublet with the Majorana fermions [10, 11]. This can be achieved only by imposing a strong degeneracy between the masses of the CP-even and CP-odd scalars, making the quartic coupling suppressed  $\lambda_5 \sim 10^{-10} - 10^{-9}$ . Such fine-tuning can be avoided by extending the minimal scotogenic model (MSM) by two real singlets and impose a global  $Z_4/Z_2$  symmetry [12]. Indeed, there have many models that have been proposed and studied beyond the MSM [13].

Based on the current/future results of both ATLAS and CMS, the IDM (and therefore, the MSM with scalar DM model) parameter space will become more constrained as the precision measurements of the EW sector gets improved. In addition, the recent results from the direct detection experiments such as LUX-ZEPLIN experiment [14], would place more stringent bounds on the model. This makes the extension of the IDM interesting, that might evade some of the constraints and predict additional interesting signatures. On top of that, among the important motivations to extend the IDM; is that it does not allow CP violation in the scalar sector, where an additional  $SU(2)_L$  Higgs doublet is useful to do this task [15]. According to the previously mentioned motivations, it is natural to consider the realisation of a scotogenic model where the SM is extended by three singlet Majorana fermions  $N_{1,2,3}$  and two inert doublets  $\Phi_{1,2}$  that couple to the leptonic left-handed doublets and the singlet Majoranas. Here, the new additional fields transform as odd fields under a global  $Z_2$  symmetry  $\{N_{1,2,3}, \Phi_{1,2}\} \rightarrow \{-N_{1,2,3}, -\Phi_{1,2}\}$  to ensure the DM stability.

This work is organized as follow. In section 2, we present the model and show the neutrino mass generation at loop. In section 3, different theoretical and experimental constraints are discussed. Sections 4 and 5 are devoted to show the viable parameter space and the DM phenomenology, respectively, in both scenarios of Majorana and scalar DM candidates. We conclude in section 6. In addition, the paper includes three appendices: in Appendix A, we discuss how to estimate the new Yukawa couplings where a generalized parameterization like the Casas-Ibarra one is introduced. In Appendix B, we generalize our results for the case of a scotogenic model with  $n$  inert scalar doublets instead of two. In Appendix C, we give the amplitude matrices that are relevant to the perturbative unitarity conditions.

## 2 The Model & Neutrino Mass

Here, we extend the SM by two inert doublets denoted by  $\Phi_{1,2} \sim (1, 2, 1)$  and three singlet Majorana fermions  $N_i \sim (1, 1, 0)$ . The Lagrangian that involves the Majorana fermions can be written as

$$\mathcal{L} \supset \bar{L}_\alpha (h_{\alpha,k} \epsilon \cdot \Phi_1 + h_{\alpha,k+3} \epsilon \cdot \Phi_2) N_k + \frac{1}{2} \bar{N}_k^C M_k N_k + h.c., \quad (2.1)$$

where  $L_\alpha$  are the left-handed lepton doublets, and  $\epsilon = i\sigma_2$  is an anti-symmetric tensor. Here,  $h_{\alpha,k}$  is a  $3 \times 6$  new Yukawa matrix, the mass matrix  $M_k$  can be considered diagonal.

The most general  $Z_2$ -symmetric, renormalizable, and gauge invariant potential reads

$$\begin{aligned}
V(H, \Phi, S, \chi) = & -\mu_H^2 |\mathcal{H}|^2 + \mu_i^2 |\Phi_i|^2 + \frac{\lambda_H}{6} |\mathcal{H}|^4 + \frac{\lambda_i}{6} |\Phi_i|^4 + \omega_i |\mathcal{H}|^2 |\Phi_i|^2 + \kappa_i |\mathcal{H}^\dagger \Phi_i|^2 \\
& + \varrho_1 |\Phi_1|^2 |\Phi_2|^2 + \varrho_2 |\Phi_1^\dagger \Phi_2|^2 + \left\{ \mu_3^2 \Phi_1^\dagger \Phi_2 + \frac{1}{2} \zeta_i (\mathcal{H}^\dagger \Phi_i)^2 + \zeta_3 (\mathcal{H}^\dagger \Phi_1) (\mathcal{H}^\dagger \Phi_2) \right. \\
& \left. + \zeta_4 (\Phi_1^\dagger \mathcal{H}) (\mathcal{H}^\dagger \Phi_2) + h.c. \right\}, \tag{2.2}
\end{aligned}$$

with  $\mathcal{H}$ , and  $\Phi$  can be parameterized as follows

$$H = \begin{pmatrix} \chi^+ \\ \frac{1}{\sqrt{2}}(v + h + i\chi^0) \end{pmatrix}, \quad \Phi_i = \begin{pmatrix} S_i^+ \\ \frac{1}{\sqrt{2}}(S_i^0 + iQ_i^0) \end{pmatrix}. \tag{2.3}$$

The terms of the Lagrangian in (2.1) and (2.2) are invariant under a global  $Z_2$  symmetry according to the charges

$$\Phi_{1,2} \rightarrow -\Phi_{1,2}, \quad N_{1,2,3} \rightarrow -N_{1,2,3}, \tag{2.4}$$

where all other fields are even. This  $Z_2$  symmetry forbids such terms like  $(\mathcal{H}^\dagger \Phi_i)(\Phi_j^\dagger \Phi_k)$ . In case of complex values for  $\mu_3^2$  and/or  $\zeta_i$ , CP is spontaneously broken, where we obtain CP-violating interactions relevant to the four neutral inert eigenstates in the basis  $\{S_1^0, S_2^0, Q_1^0, Q_2^0\}$ . This case would be very interesting since the additional CP-violating sources are required for matter antimatter asymmetry generation [16]. In what follows, we consider real values for the parameters  $\mu_3^2$  and  $\zeta_{i=1,2,3,4}$  to avoid spontaneous CP violation. The case of spontaneous CP violation (complex values for  $\mu_3^2$  and/or  $\zeta_i$ ) requires an independent investigation since it may have important cosmological consequences.

After the electroweak symmetry breaking (EWSB), we are left with three CP-even scalars ( $h, H_{1,2}^0$ ), two CP-odd scalar  $A_{1,2}^0$  and two pair of charged scalars  $H_{1,2}^\pm$ . The Higgs mass mass is given  $m_h^2 = 2\mu_H^2 = \frac{\lambda_H}{3}v^2$ , and the inert eigenstates are defined as

$$\begin{pmatrix} H_1^0 \\ H_2^0 \end{pmatrix} = \begin{pmatrix} c_H & s_H \\ -s_H & c_H \end{pmatrix} \begin{pmatrix} S_1^0 \\ S_2^0 \end{pmatrix}, \quad \begin{pmatrix} A_1^0 \\ A_2^0 \end{pmatrix} = \begin{pmatrix} c_A & s_A \\ -s_A & c_A \end{pmatrix} \begin{pmatrix} Q_1^0 \\ Q_2^0 \end{pmatrix}, \quad \begin{pmatrix} H_1^\pm \\ H_2^\pm \end{pmatrix} = \begin{pmatrix} c_C & s_C \\ -s_C & c_C \end{pmatrix} \begin{pmatrix} S_1^\pm \\ S_2^\pm \end{pmatrix}, \tag{2.5}$$

with  $c_X = \cos \theta_X$ ,  $s_X = \sin \theta_X$  and  $\theta_H$ ,  $\theta_A$  and  $\theta_C$  are the mixing angles that diagonalise the CP-even, CP-odd and charged scalars mass matrices, respectively. The charged, neutral CP-even and CP-odd scalar squared mass matrices in the basis  $\{S_1^\pm, S_2^\pm\}$ ,  $\{S_1^0, S_2^0\}$  and  $\{Q_1^0, Q_2^0\}$ , respectively, are given by

$$M_C^2 = \begin{bmatrix} \mu_1^2 & \mu_3^2 \\ \mu_3^2 & \mu_2^2 \end{bmatrix} + \frac{v^2}{2} \begin{bmatrix} \omega_1 & 0 \\ 0 & \omega_2 \end{bmatrix}, \quad M_{H,A}^2 = M_C^2 + \frac{v^2}{2} \begin{bmatrix} \kappa_1 & \zeta_4 \\ \zeta_4 & \kappa_2 \end{bmatrix} \pm \frac{v^2}{2} \begin{bmatrix} \zeta_1 & \zeta_3 \\ \zeta_3 & \zeta_2 \end{bmatrix}. \tag{2.6}$$

The mass eigenvalues and the mixing are given by

$$\begin{aligned}
m_{X_{1,2}}^2 = & \frac{1}{2} \left( [M_X^2]_{11} + [M_X^2]_{22} \mp \sqrt{([M_X^2]_{22} - [M_X^2]_{11})^2 + 4[M_X^2]_{12}^2} \right), \\
\tan(2\theta_X) = & 2[M_X^2]_{12} / ([M_X^2]_{22} - [M_X^2]_{11}), \tag{2.7}
\end{aligned}$$

for  $([X_{1,2}, \theta_X] = [H_{1,2}^0, \theta_H], [A_{1,2}^0, \theta_A], [H_{1,2}^\pm, \theta_C])$ , respectively.

Then, the independent free parameters are

$$M_{1,2,3}, m_{H_{1,2}^\pm}, m_{H_{1,2}^0}, m_{A_{1,2}^0}, s_{H,A,C}, \omega_{1,2}, h_{\alpha i}, \quad (2.8)$$

and the of parameters are given by

$$\begin{aligned} \mu_1^2 &= m_{H_1^\pm}^2 c_C^2 + m_{H_2^\pm}^2 s_C^2 - \frac{1}{2} \omega_1 v^2, \quad \mu_2^2 = m_{H_1^\pm}^2 s_C^2 + m_{H_2^\pm}^2 c_C^2 - \frac{1}{2} \omega_2 v^2, \quad \mu_3^2 = c_C s_C (m_{H_2^\pm}^2 - m_{H_1^\pm}^2), \\ \kappa_1 v^2 &= m_{H_1^0}^2 c_H^2 + m_{H_2^0}^2 s_H^2 + m_{A_1^0}^2 c_A^2 + m_{A_2^0}^2 s_A^2 - 2(m_{H_1^\pm}^2 c_C^2 + m_{H_2^\pm}^2 s_C^2), \\ \kappa_2 v^2 &= m_{H_2^0}^2 c_H^2 + m_{H_1^0}^2 s_H^2 + m_{A_2^0}^2 c_A^2 + m_{A_1^0}^2 s_A^2 - 2(m_{H_2^\pm}^2 c_C^2 + m_{H_1^\pm}^2 s_C^2), \\ \xi_1 v^2 &= (m_{H_1^0}^2 c_H^2 + m_{H_2^0}^2 s_H^2) - (m_{A_2^0}^2 s_A^2 + m_{A_1^0}^2 c_A^2), \quad \xi_2 v^2 = (m_{H_1^0}^2 s_H^2 + m_{H_2^0}^2 c_H^2) - (m_{A_1^0}^2 s_A^2 + m_{A_2^0}^2 c_A^2), \\ \xi_3 v^2 &= s_H c_H (m_{H_2^0}^2 - m_{H_1^0}^2) - s_A c_A (m_{A_2^0}^2 - m_{A_1^0}^2), \\ \xi_4 v^2 &= s_H c_H (m_{H_2^0}^2 - m_{H_1^0}^2) + s_A c_A (m_{A_2^0}^2 - m_{A_1^0}^2) - 2c_C s_C (m_{H_2^\pm}^2 - m_{H_1^\pm}^2). \end{aligned} \quad (2.9)$$


The first term in (2.1), can be written as

$$\mathcal{L} \supset \sum_{\alpha,j,k} \frac{1}{\sqrt{2}} (g_{\alpha k}^{(j)} H_j^0 + i f_{\alpha k}^{(j)} A_j^0) \bar{\nu}_{\alpha L} N_k - \zeta_{\alpha k}^{(j)} H_j^\pm \bar{\ell}_{\alpha L} N_k + h.c., \quad (2.10)$$

with  $\alpha = e, \mu, \tau, j = 1, 2$  and  $k = 1, 3$ ; and

$$\begin{aligned} g_{\alpha k}^{(1)} &= c_H h_{\alpha,k} + s_H h_{\alpha,k+3}, \quad g_{\alpha k}^{(2)} = -s_H h_{\alpha,k} + c_H h_{\alpha,k+3}, \\ f_{\alpha k}^{(1)} &= c_A h_{\alpha,k} + s_A h_{\alpha,k+3}, \quad f_{\alpha k}^{(2)} = -s_A h_{\alpha,k} + c_A h_{\alpha,k+3}, \\ \zeta_{\alpha k}^{(1)} &= c_C h_{\alpha,k} + s_C h_{\alpha,k+3}, \quad \zeta_{\alpha k}^{(2)} = -s_C h_{\alpha,k} + c_C h_{\alpha,k+3}. \end{aligned} \quad (2.11)$$

The first two interaction terms in (2.10) give rise to the neutrino mass a la scotogenic way; and the last term leads to the LFV processes  $\ell_\beta \rightarrow \ell_\alpha + \gamma$  and  $\ell_\beta \rightarrow 3\ell_\alpha$ . Then, the neutrino mass  $[\alpha, \beta]$  matrix element comes from the contribution 12 diagrams as shown in Fig. 1.



**Figure 1:** The 12 Feynman diagrams responsible for neutrino mass generation.

It can be written as

$$m_{\alpha\beta}^{(v)} = \sum_{k=1}^3 \sum_{j=1}^2 M_k \left\{ g_{\alpha k}^{(j)} g_{\beta k}^{(j)} \mathcal{F}(m_{H_j^0}/M_k) - f_{\alpha k}^{(j)} f_{\beta k}^{(j)} \mathcal{F}(m_{A_j^0}/M_k) \right\}, \quad (2.12)$$

with  $\mathcal{F}(x) = \frac{x^2 \log(x)}{8\pi^2(x^2-1)}$ . The divergent part of the diagrams in Fig. 1 cancels due to the identity

$$\sum_{j=1}^2 \left\{ g_{\alpha k}^{(j)} g_{\beta k}^{(j)} - f_{\alpha k}^{(j)} f_{\beta k}^{(j)} \right\} = 0. \quad (2.13)$$

The neutrino mass matrix (2.12) can be parameterized as

$$(m^{(\nu)})_{3 \times 3} = (h)_{3 \times 6} \cdot (\Lambda)_{6 \times 6} \cdot (h^T)_{6 \times 3} , \quad (2.14)$$

$$\Lambda = \begin{pmatrix} \omega_1 & \omega_3 \\ \omega_3 & \omega_2 \end{pmatrix}, \quad (2.15)$$

where  $\omega_{1,2,3}$  are  $3 \times 3$  diagonal matrices whose elements are given by

$$\begin{aligned} (\omega_1)_{ik} &= \delta_{ik} M_k [(c_H^2 \mathcal{F}(m_{H_1^0}/M_k) + s_H^2 \mathcal{F}(m_{H_2^0}/M_k) - c_A^2 \mathcal{F}(m_{A_1^0}/M_k) - s_A^2 \mathcal{F}(m_{A_2^0}/M_k))], \\ (\omega_2)_{ik} &= \delta_{ik} M_k [s_H^2 \mathcal{F}(m_{H_1^0}/M_k) + c_H^2 \mathcal{F}(m_{H_2^0}/M_k) - s_A^2 \mathcal{F}(m_{A_1^0}/M_k) - c_A^2 \mathcal{F}(m_{A_2^0}/M_k)], \\ (\omega_3)_{ik} &= \delta_{ik} M_k [s_{HC} \mathcal{F}(m_{H_1^0}/M_k) - \mathcal{F}(m_{H_2^0}/M_k) - s_{AC} (\mathcal{F}(m_{A_1^0}/M_k) - \mathcal{F}(m_{A_2^0}/M_k))], \end{aligned} \quad (2.16)$$

for  $i, k = 1, 3$ . Here, the matrix  $\Lambda$  in (2.14) is not diagonal, and therefore cannot use the Casas-Ibarra parameterization [17] to estimate the Yukawa couplings  $h$ . However, we can derive a similar formula in such case ( $\Lambda$  is not diagonal). We give a general solution for the Yukawa coupling  $3 \times M$  matrix<sup>1</sup> as

$$h_{3 \times M} = (U_\nu)_{3 \times 3} \cdot (D_{\sqrt{m_\nu}})_{3 \times 3} \cdot (T)_{3 \times M} \cdot (D_{(\Lambda'_i)^{-1/2}})_{M \times M} \cdot (Q)_{M \times M} , \quad (2.17)$$

with  $U_\nu$  is the Pontecorvo-Maki-Nakawaga-Sakata (PMNS) mixing matrix,  $D_{\sqrt{m_\nu}} = \text{diag}\{m_1^{1/2}, m_2^{1/2}, m_3^{1/2}\}$  with  $m_i$  are the neutrino mass eigenvalues; and  $T$  is an arbitrary  $3 \times M$  matrix that fulfills the identity  $T_{3 \times M} \cdot T_{M \times 3}^T = \mathbf{1}_{3 \times 3}$ . The matrix  $(D_{(\Lambda'_i)^{-1/2}})_{M \times M} = \text{diag}\{(\Lambda'_i)^{-1/2}\}$ , with  $\Lambda'_i$  to be the eigenvalues of the matrix  $\Lambda$  in (2.14), that diagonalized using the matrix  $Q$ , i.e.,  $\text{diag}\{\Lambda'_i\} = Q \cdot \Lambda \cdot Q^T$ . In Appendix A, we discuss the derivation of (2.17) and the possible representations for this matrix. In addition, we discuss in Appendix B the generalization of this model by considering  $\mathbf{n}$  inert scalar doublets instead of two doublets.

### 3 Theoretical & Experimental Constraints

Here, we discuss the theoretical and the experimental constraints relevant to our model. These constraints are briefly discussed below:

- **Perturbativity:** all the quartic vertices of the scalar fields should be satisfy the perturbativity bounds, i.e.,

$$\lambda_H, \lambda_{1,2}, |\omega_{1,2}|, |\kappa_{1,2} + \omega_{1,2}|, |\omega_{1,2} + \kappa_{1,2} \pm \xi_{1,2}|, |\varrho_1|, |\varrho_1 + \varrho_2|, |\xi_{1,2}|, \frac{1}{2} |\kappa_{1,2} \pm \xi_{1,2}|, \frac{1}{2} |\varrho_2| \leq 4\pi. \quad (3.1)$$

- **Perturbative unitarity:** the perturbative unitarity has to be preserved in all processes involving scalars or gauge bosons. In the high-energy limit, the gauge bosons should be replaced by their

<sup>1</sup>Here,  $M = 6$  in our case. However, the solutions (2.17) are valid for any value  $M \geq 3$ .

Goldstones. In our model, the scattering amplitudes in the high-energy limit contain six submatrices that decouple from each other due to the conservation of electric-charge,  $Z_2$  symmetry and/or  $CP$ -quantum numbers.

These six submatrices are defined in the basis of initial/final states that are: (1)  $CP$ -even,  $Z_2$  even, &  $Q_{em} = 0$ ; (2)  $CP$ -even,  $Z_2$  odd, &  $Q_{em} = 0$ , (3)  $CP$ -odd,  $Z_2$  even, &  $Q_{em} = 0$ , (4)  $CP$ -odd,  $Z_2$  odd, &  $Q_{em} = 0$ , (5)  $Z_2$  even, &  $Q_{em} = \pm 1$ ; and (6)  $Z_2$  odd, &  $Q_{em} = \pm 1$ . We require that the eigenvalues of these matrices, that are given in Appendix C, to be smaller than  $8\pi$ .

- **Vacuum Stability:** the scalar potential is required to be bounded from below in all the directions of the field space. Therefore, the following conditions need to be fulfilled:

$$\lambda_H, \lambda_1, \lambda_2, \det \begin{bmatrix} \lambda_H & \overline{\omega_1 + \kappa_1 + \zeta_1} & \overline{\omega_2 + \kappa_2 + \zeta_2} \\ \frac{\omega_1 + \kappa_1 + \zeta_1}{\omega_2 + \kappa_2 + \zeta_2} & \lambda_1 & \frac{\overline{\omega_1 + \kappa_1 + \zeta_1}}{\overline{\omega_2 + \kappa_2 + \zeta_2}} \\ \frac{\omega_1 + \kappa_1 + \zeta_1}{\omega_2 + \kappa_2 + \zeta_2} & \frac{\overline{\omega_1 + \kappa_1 + \zeta_1}}{\overline{\omega_2 + \kappa_2 + \zeta_2}} & \lambda_2 \end{bmatrix} > 0, \quad (3.2)$$

with  $\overline{X} = \min(X, 0)$ . In addition, we require that the vacuum ( $\langle h \rangle = v, \langle S_i^0 \rangle = 0$ ) would be the deepest one. As a conservative choice in this work, we consider the conditions that the potential (2.2) should not develop a vev in the inert directions, i.e., we should have  $\langle S_i^0 \rangle = 0$  either at  $\langle h \rangle = 0$  or  $\langle h \rangle = v$ . These conditions can be translated into

$$\mu_1^2, \mu_2^2, \mu_1^2 + \mu_2^2 - \sqrt{(\mu_2^2 - \mu_1^2)^2 + 4(\mu_3^2)^2} > 0, \quad (3.3)$$

and obviously,  $m_{H_1^0}^2, m_{A_1^0}^2, m_{H_1^\pm}^2 > 0$ .

- **Gauge bosons decay widths:** The decay widths of the  $W/Z$ -bosons were measured with high precision at LEP. Therefore, we require that the decays of  $W/Z$ -bosons to  $Z_2$ -odd scalars is closed. This is fulfilled if one assumes that

$$\min(m_{H_1^0} + m_{A_1^0}, 2m_{H_1^\pm}) > m_Z, \min(m_{H_1^\pm} + m_{A_1^0}, m_{H_1^\pm} + m_{H_1^0}) > m_W. \quad (3.4)$$

- **Lepton flavor violating (LFV) decays:** in this model, LFV decay processes arise via one-loop diagrams mediated by the  $H_{i=1,2}^\pm$  and  $N_k$  particles. The branching ratio of the decay  $\mathcal{B}(\ell_\alpha \rightarrow \ell_\beta + \gamma)$  due to the contribution of the interactions (2.1) is [18]:

$$\mathcal{B}(\ell_\alpha \rightarrow \ell_\beta + \gamma) = \frac{3\alpha v^4}{32\pi} \left| \sum_{k=1}^3 \sum_{j=1}^2 \frac{(\zeta_{\alpha k}^{(j)}) (\zeta_{\beta k}^{(j)})^*}{m_{H_j^\pm}^2} F\left(\frac{M_k^2}{m_{H_j^\pm}^2}\right) \right|^2, \quad (3.5)$$

where  $\alpha = e^2/4\pi$  is the electromagnetic fine structure constant and  $F(x) = (1 - 6x + 3x^2 + 2x^3 - 6x^2 \log x)/6(1 - x)^4$ . Since the bounds on  $\ell_\alpha \rightarrow \ell_\beta + \gamma$  are more severe, especially  $\mu \rightarrow e + \gamma$ , we will consider only the LFV bounds on  $\ell_\alpha \rightarrow \ell_\beta + \gamma$  and not  $\ell_\alpha \rightarrow \ell_\beta \ell_\beta \ell_\beta$ , since the latter's would implicitly fulfilled [19].

- **Direct searches of charginos and neutralinos at the LEP-II experiment:** we use the null results of neutralinos and charginos at LEP [20] to put lower bounds on the masses of charged Higgs  $H_1^\pm$  and the light neutral scalars of the inert doublets ( $H_1^0, A_1^0$ ). The bound obtained from a re-interpretation of neutralino searches [21] cannot apply to our model since the decays  $A_1^0 \rightarrow H_1^\pm Z (\rightarrow \ell^+ \ell^-)$  are kinematically forbidden. On the other hand, in most regions of the parameter space, the charged Higgs decays exclusively into a Majorana fermion and a charged lepton. For Yukawa couplings of order one  $h_{ei} \simeq \mathcal{O}(1)$ , the following bounds are derived  $m_{H^\pm} > 100$  GeV [11].
- **The electroweak precision tests:** in this model, the oblique parameters acquire contributions from the existence of inert scalars. We take  $\Delta U = 0$  in our analysis, the oblique parameters are given by [22]

$$\begin{aligned}
\Delta T &= \frac{1}{16\pi s_W^2 M_W^2} \left\{ c_H^2 c_C^2 F(m_{H_1^0}^2, m_{H_1^\pm}^2) + c_H^2 s_C^2 F(m_{H_1^0}^2, m_{H_2^\pm}^2) + s_H^2 c_C^2 F(m_{H_2^0}^2, m_{H_1^\pm}^2) \right. \\
&\quad + s_H^2 s_C^2 F(m_{H_2^0}^2, m_{H_2^\pm}^2) + c_A^2 c_C^2 F(m_{A_1^0}^2, m_{H_1^\pm}^2) + c_A^2 s_C^2 F(m_{A_1^0}^2, m_{H_2^\pm}^2) + s_A^2 c_C^2 F(m_{A_1^0}^2, m_{H_1^\pm}^2) \\
&\quad + s_A^2 s_C^2 F(m_{A_1^0}^2, m_{H_2^\pm}^2) - c_H^2 c_A^2 F(m_{H_1^0}^2, m_{A_1^0}^2) - c_H^2 s_A^2 F(m_{H_1^0}^2, m_{A_2^0}^2) - s_H^2 c_A^2 F(m_{H_2^0}^2, m_{A_2^0}^2) \\
&\quad \left. - s_H^2 s_A^2 F(m_{H_2^0}^2, m_{A_2^0}^2) \right\}, \\
\Delta S &= \frac{1}{24\pi} \left\{ (2s_W^2 - 1)^2 \left[ c_C^4 G(m_{H_1^\pm}^2, m_{H_1^\pm}^2, m_Z^2) + s_C^4 G(m_{H_2^\pm}^2, m_{H_2^\pm}^2, m_Z^2) + 2c_C^2 s_C^2 G(m_{H_1^\pm}^2, m_{H_2^\pm}^2, m_Z^2) \right] \right. \\
&\quad + c_H^2 c_A^2 G(m_{H_1^0}^2, m_{A_1^0}^2, m_Z^2) + c_H^2 s_A^2 G(m_{H_1^0}^2, m_{A_2^0}^2, m_Z^2) + s_H^2 c_A^2 G(m_{H_2^0}^2, m_{A_1^0}^2, m_Z^2) \\
&\quad + s_H^2 s_A^2 G(m_{H_2^0}^2, m_{A_2^0}^2, m_Z^2) + c_H^2 c_C^2 \log\left(\frac{m_{H_1^0}^2}{m_{H_1^\pm}^2}\right) + c_H^2 s_C^2 \log\left(\frac{m_{H_1^0}^2}{m_{H_2^\pm}^2}\right) + s_H^2 c_C^2 \log\left(\frac{m_{H_2^0}^2}{m_{H_1^\pm}^2}\right) \\
&\quad + s_H^2 s_C^2 \log\left(\frac{m_{H_2^0}^2}{m_{H_2^\pm}^2}\right) + c_H^2 c_A^2 \log\left(\frac{m_{A_1^0}^2}{m_{H_1^\pm}^2}\right) + c_H^2 s_A^2 \log\left(\frac{m_{A_2^0}^2}{m_{H_1^\pm}^2}\right) + s_H^2 c_A^2 \log\left(\frac{m_{A_1^0}^2}{m_{H_2^\pm}^2}\right) \\
&\quad \left. + s_H^2 s_A^2 \log\left(\frac{m_{A_2^0}^2}{m_{H_2^\pm}^2}\right) \right\}, \tag{3.6}
\end{aligned}$$

where  $s_W \equiv \sin \theta_W$ , with  $\theta_W$  is the Weinberg mixing angle, and  $F(x, y)$  and  $G(x, y, z)$  are one-loop functions that can be found in [22].

- **The di-photon Higgs decay:** the Higgs couplings to the two charged scalar can change drastically the value of the Higgs boson loop-induced decay into two photons. These loop interactions depends on the signs and the values of  $\omega_{1,2}$ , as well on the mixing and  $\theta_C$ . The ratio  $R_{\gamma\gamma} = \mathcal{B}(h \rightarrow \gamma\gamma) / \mathcal{B}(h \rightarrow \gamma\gamma)^{\text{SM}} = 1.02_{-0.12}^{+0.09}$  [23], that characterizes the di-photon Higgs decay modification can be written as

$$R_{\gamma\gamma} = (1 - \mathcal{B}_{\text{BSM}}) \left| 1 + \frac{v^2}{2} \frac{\sum_{j=1}^2 \frac{r_j}{m_{H_j^+}^2} A_0^{\gamma\gamma}\left(\frac{m_h^2}{4m_{H_j^+}^2}\right)}{A_1^{\gamma\gamma}\left(\frac{m_h^2}{4m_W^2}\right) + \frac{4}{3} A_{1/2}^{\gamma\gamma}\left(\frac{m_h^2}{4m_t^2}\right)} \right|^2, \tag{3.7}$$



with  $r_1 = c_C^2 \omega_1 + s_C^2 \omega_2$ ,  $r_2 = s_C^2 \omega_1 + c_C^2 \omega_2$ ; and  $\mathcal{B}_{BSM}$  represents any non-SM decay for the Higgs like  $h \rightarrow H_i^0 H_k^0, A_i^0 A_k^0$ . The loop functions  $A_{0,1/2,1}^{\gamma\gamma}$  are given in the literature [24]. The  $R_{\gamma\gamma}$  ratio estimation in (3.7) is based on the assumption of the SM production rates for the SM Higgs boson [25]. Similarly, the ratio  $R_{\gamma Z} = \mathcal{B}(h \rightarrow \gamma Z) / \mathcal{B}(h \rightarrow \gamma Z)^{\text{SM}}$  can be written as

$$R_{\gamma Z} = (1 - \mathcal{B}_{BSM}) \left| 1 - \frac{v^2}{2} \frac{1 - 2s_W^2}{c_W} \frac{\sum_{j=1}^2 \frac{r_j}{m_{H_j^+}^2} A_0^{\gamma Z} \left( \frac{m_h^2}{4m_{H_j^+}^2}, \frac{m_Z^2}{4m_{H_j^+}^2} \right)}{c_W A_1^{\gamma Z} \left( \frac{m_h^2}{4m_W^2}, \frac{m_Z^2}{4m_W^2} \right) + \frac{6 - 16s_W^2}{3c_W} A_{1/2}^{\gamma Z} \left( \frac{m_h^2}{4m_t^2}, \frac{m_Z^2}{4m_t^2} \right)} \right|^2, \quad (3.8)$$

with  $c_W = \cos \theta_W$ ,  $s_W = \sin \theta_W$  and  $\theta_W$  is the Weinberg mixing angle; and the loop functions  $A_{0,1/2,1}^{\gamma Z}$  are given in the literature [24].

In Appendix B, we discuss the generalization of these results and constraints in the case of a  $n$  inert scalar doublets model.

## 4 Viable Parameter Space

In our analysis, we consider both cases where the DM candidate could be a Majorana fermion  $N_1$  and the light inert scalar  $H_1^0$ . By taking into account all the above mentioned theoretical and experimental constraints, we perform a full numerical scan over the model free parameters (2.8), where we will consider 4 k benchmark points (BPs) for each case. In our scan, we consider the following ranges for the free parameters


$$|s_{H,A,C}| \leq 1, \quad |\omega_{1,2}| < 4\pi, \quad m_{H_1^0, A_1^0} < 3 \text{ TeV}, \quad 78 \text{ GeV} < m_{H_1^\pm} < 3 \text{ TeV}, \quad (4.1)$$

while focusing on the parameter space regions that makes this model different than a two inert doublets model SM extension, i.e., the case with non negligible new Yukawa couplings  $\min |h_{\alpha i}| \geq 10^{-3}$ .

In Fig. 2, we show the mixing angles sine (left-up), the the strength of the new Yukawa couplings  $h_{\alpha i}$  (right-up) the masses; and the inert masses (bottom).

Clearly from Fig. 2, the mixing angles of the CP-even, CP-odd and charged sectors can take most of the possible ranges, however, there some preferred values around  $s_H = 0$  and  $s_A = s_H$ . All the above mentioned constraints together with the neutrino oscillation data that are taken into account via (2.17); lead to the new Yukawa couplings  $h_{\alpha i}$  strength as shown in Fig. 2-top-right. It is clear that all the couplings can suppressed which makes the LFV constraints easily fulfilled and the model indistinguishable from the SM extended by two inert singlets. As mentioned previously, here we are interested in the opposite case where  $|h_{\alpha i}| \geq 10^{-3}$ , then, the couplings can be large as  $|h_{\alpha i}| < \sqrt{4\pi}$  and a significant hierarchy between the largest and smallest couplings strength does exist as  $\max(|h_{\alpha i}|) > 1.8 \min(|h_{\alpha i}|)$ . According to Fig. 2-bottom, the masses of the inert eigenstates lie within large values intervals.


In Fig. 3, we present the Higgs decay modifiers of  $h \rightarrow \gamma\gamma, \gamma Z$  (left), the oblique parameters (middle); and the LFV branching ratios scaled by their experimental bounds.



**Figure 2:** Left-top: the CP-even mixing angle sine vs the corresponding CP-odd one, where the palette shows the mixing of the charged scalar sector. Right-top: the strength of the new Yukawa couplings  $h_{\alpha i}$  where the palette shows the ratio  $\delta^0 = (m_{A_1^0}^2 - m_{H_1^0}^2)/m_{H_1^0}^2$ . Left-bottom: the mass ratio  $\delta^0 = (m_{A_1^0}^2 - m_{H_1^0}^2)/m_{H_1^0}^2$  versus the light CP-even scalar mass  $m_{H_1^0}^2$ , where the palette shows the light charged scalar mass  $m_{H_1^\pm}^2$ . Right-bottom: the ratios  $\delta_{\text{even}} = (m_{H_2^0}^2 - m_{H_1^0}^2)/m_{H_1^0}^2$ ,  $\delta_{\text{even}} = (m_{A_2^0}^2 - m_{A_1^0}^2)/m_{A_1^0}^2$  and  $\delta_{\text{charged}} = (m_{H_2^\pm}^2 - m_{H_1^\pm}^2)/m_{H_1^\pm}^2$  that represent the mass relative difference in each sector.

One notices that for allowed values for the Higgs branching ratio  $h \rightarrow \gamma\gamma$ , the branching ratio  $h \rightarrow \gamma Z$  can be modified with respect to the SM within the range  $[-5.2\%, 5, 1\%]$ . Concerning the electroweak precision tests, as it is expected the  $T$  parameter is more sensitive to the quantum corrections due to the interactions of the gauge bosons to the two extra doublets, while the  $S$  parameter is sensitive to the corrections only in case of light charged scalars. Like in many models, the constraint from the branching ratio  $\mu \rightarrow e\gamma$  is the most severe one and the bounds from the other LFV decays ( $\tau \rightarrow e\gamma$  and  $\tau \rightarrow \mu\gamma$ ) are at least three orders of magnitude smaller.

At colliders, the model may predict many interesting signatures. For instance, it predicts all the signatures relevant to the MSM [10]. At the LHC, charged scalars can be pair produced as  $pp \rightarrow H_{1,2}^+ H_{1,2}^-$  or associated with neutral inert scalars  $pp \rightarrow H_{1,2}^\pm H_{1,2}^0, H_{1,2}^\pm A_{1,2}^0$ . These channels can be probed via the final states mono-lepton  $\ell + \cancel{E}_T$  and di-lepton  $2\ell + \cancel{E}_T$  in case if the channels  $H_{1,2}^\pm \rightarrow W^\pm H_{1,2}^0, W^\pm A_{1,2}^0$



**Figure 3:** Left: The Higgs decay modifiers of  $h \rightarrow \gamma\gamma, \gamma Z$ , where the colored band represents the allowed experimental values. Middle: the oblique parameters  $\Delta T$  vs  $\Delta S$ , where the palette shows the light charged scalar mass. Right: the LFV branching ratios scaled by their experimental bounds versus the light Majorana mass.

are closed. If these channels are open, then the final states  $4 jets + \cancel{E}_T$  and  $1\ell + 2 jets + \cancel{E}_T$  could be useful to probe this model as it is useful to probe the MSM [10]. In order to look for beyond MSM signatures, one has to consider the heavy inert scalar production at the LHC:  $pp \rightarrow H_2^\pm H_{1,2}^\mp, H_2^\pm H_{1,2}^0, H_2^\pm A_{1,2}^0, H_2^0 H_{1,2}^\pm, A_2^0 H_{1,2}^\pm, H_2^0 H_{1,2}^0, A_2^0 H_{1,2}^0, A_2^0 A_{1,2}^0$ , where the heavy scalars should decay as  $H_2^+ \rightarrow W^+ H_1^0, W^+ A_1^0, h H_1^+, H_2^0 \rightarrow Z A_1^0, h H_1^0$  and  $A_2^0 \rightarrow Z H_1^0, h A_1^0$ . These novel signatures can be probed through the final states  $8 jets + \cancel{E}_T, 1\ell + 4 jets + \cancel{E}_T, 1\ell + 2 jets + \cancel{E}_T, 4b + \cancel{E}_T, 2 jets + 2b + \cancel{E}_T, 2b + \cancel{E}_T$ , that can not be seen in the MSM or in other neutrino mass and DM motivated SM extensions. The investigation of these novel signatures requires a full and precise numerical scan to define the relevant regions of the parameter space, that should be confronted with the existing analyses.

## 5 Dark Matter: Majorana or Scalar?

In this model, DM candidate could be either the light Majorana fermion ( $N_1$ ), the light  $CP$ -even ( $CP$ -odd) scalar  $H_1^0$  ( $A_1^0$ ), or a mixture of all these components if they are degenerate in mass. In the case of scalar DM, the possible annihilation channels are  $W^\pm W^\mp, ZZ, q_i \bar{q}_i, hh, \bar{\ell}_\alpha \ell_\beta$  and  $\bar{\nu}_\alpha \nu_\beta$ . In this case, very small couplings  $h_{\alpha i}$  could be favored by the LFV constraints, and therefore the contributions of the channels  $\nu_\alpha \bar{\nu}_\beta, \ell_\alpha \ell_\beta$  to the annihilation cross section would be negligible. Then, this case is almost identical to the SM extended by two inert doublets with scalar DM. The co-annihilation effect along the channels  $H_1^0 A_1^0 (A_1^0 A_1^0) \rightarrow X_{SM} X'_{SM}$  could be important if the mass difference  $(m_{A_1^0}^2 - m_{H_1^0}^2) / (m_{A_1^0}^2 + m_{H_1^0}^2)$  is small enough. This makes the couplings  $h_{\alpha i}$  non-suppressed and therefore the annihilation into  $\nu_\alpha \bar{\nu}_\beta, \ell_\alpha \ell_\beta$  is more important<sup>2</sup>. Such effect could make this setup (spin-0 DM) better than the minimal IDM. In case of Majorana DM scenario, the DM (co-)annihilation could occur into charged leptons  $\ell_\alpha^- \ell_\beta^+$  and light neutrinos  $\nu_\alpha \bar{\nu}_\beta$ ; via  $t$ -channel diagrams mediated by the charged scalar  $H_{1,2}^\pm$ ; and the


<sup>2</sup>To be more precise, the mass degeneracy between  $H_1^0$  and  $A_1^0$  is not enough to get non-suppressed values for the couplings  $h_{\alpha i}$ , a similar degeneracy between both  $H_2^0$  and  $A_2^0$  and between the  $CP$ -even and  $CP$ -odd mixing angles; are also required.

neutral scalars  $H_{1,2}^0, A_{1,2}^0$ , respectively. Here, the annihilation cross section is fully triggered by the non-suppressed values of the couplings  $h_{ai}$ .

In case of scalar DM, negative searches from DM direct detection (DD) at underground detectors impose more constraints on the parameters space. This is translated into upper bounds on the DM-nucleon cross section. In our model, the DM interaction with nucleons occurs via a single t-channel diagram mediated by the Higgs, and the scattering cross-section is given by:

$$\sigma_{det} = \frac{(\lambda_{hH_1^0 H_1^0})^2 m_{\mathcal{N}}^2 (m_{\mathcal{N}} - \frac{7}{9}m_{\mathcal{B}})^2}{4\pi v^2 m_h^4 (m_{\mathcal{N}} + m_{H_1^0})^2}, \quad (5.1)$$

where  $m_{\mathcal{N}}$  and  $m_{\mathcal{B}}$  are, respectively, the nucleon and baryon masses in the chiral limit [26]; and  $\lambda_{hH_1^0 H_1^0} = \frac{v}{2}[c_H^2(\kappa_1 + \omega_1 + \xi_1) + s_H^2(\kappa_2 + \omega_2 + \xi_2) + 2s_H c_H(\xi_3 + \xi_4)]$  is the DM Higgs triple coupling. In Fig. 4, we show the DD cross section (5.1) versus the DM mass for the BPs used in Fig. 2 that correspond to scalar DM cases, compared with recent experimental bounds from the PandaX-4T experiment [27] and LUX-ZEPLIN [14].



**Figure 4:** The DD cross section vs the DM mass for the scalar DM BPs used in Fig. 2. The violet, blue and orange lines correspond to the bounds from LUX-ZEPLIN [14], Xenon-1T bound [28] and PandaX-4T [27], respectively, while the yellow one represents the neutrino floor [29].


In order to show that both scalar and Majorana DM scenarios are viable in this model, we consider the two PBs among the BPs used in Fig. 2, where their free parameters values are given Table ??, where BP1 and BP2 correspond to Majorana and scalar DM cases, respectively.

Then, we use the package MadDM [30] to estimate the relic density and the direct detection cross section for the scalar DM case by varying both the DM mass and the Yukawa couplings  $h_{ai}$  strength as shown in Fig. 5. In order to do so, we used the package FeynRules [31] to implement the model and produce the UFO files.

From Fig. 5-left, the dependence of the relic density on the DM mass in case of Majorana DM is quadratic  $\Omega_{DM} h^2 \sim m_{DM}^2$ . This can be understood due to the fact that the Majorana DM (co-

BP1	$s_{H,A,C} = \{-0.053734, 0.72187, 0.82912\}, \omega_{1,2} = \{-0.15715, -0.92515\},$ $m_{H_1^0} = \{2755.6, 2820.6\}, m_{A_1^0} = \{2766.7, 2823\},$ $m_{H_{1\pm}^\pm} = \{2318.7, 2327.8\}, M_i = \{2317.1, 2318.7, 2327.8\},$ $h_{\alpha i} = \begin{pmatrix} -1.3664 - i0.066927, & -0.0063156 - i1.1746, & -0.020184 + i0.28501, & -0.0063503 - i1.1811, & -1.3739 - i0.067294, & -0.020295 + i0.28657 \\ 1.8625 + i0.046833, & -0.09083 - i2.2339, & -0.0063982 - i0.33045, & -0.091329 - i2.2461, & 1.8727 + i0.04709, & -0.0064333 - i0.33226 \\ -2.4404 - i0.065694, & -0.667 - i1.8413, & 0.037294 - i0.36281, & -0.67066 - i1.8514, & -2.4538 - i0.066055, & 0.037499 - i0.36481 \end{pmatrix}$
BP2	$s_{H,A,C} = \{-0.38546, -0.27092, -0.019603\}, \omega_{1,2} = \{-1.0362, 5.1668\},$ $m_{H_1^0} = \{1228.9, 1246\}, m_{A_1^0} = \{1229.3, 1266\},$ $m_{H_{1\pm}^\pm} = \{1230.4, 1270.1\}, M_i = \{1633.4, 1850.7, 1853.5\},$ $h_{\alpha i} = 10^{-3} \times \begin{pmatrix} 2.5475 - i51.865, & 15.027 + i1.4322, & 19.457 + i0.69002, & 0.44292 - i9.0175, & 2.6139 + i0.24914, & 3.3845 + i0.12003 \\ -1.7657 + i70.885, & -31.844 + i0.82879, & 4.9703 - i0.46673, & -0.307 + i12.325, & -5.5394 + i0.14417, & 0.86461 - i0.08119 \\ 2.5076 - i92.163, & -33.608 + i0.041734, & 1.0967 - i6.4609, & 0.43599 - i16.024, & -5.8463 + i0.0072598, & 0.19077 - i1.1239 \end{pmatrix}$

**Table 1:** The values of the free parameters for BP1 and BP2. Here, all the masses are given in GeV.



**Figure 5:** The DM relic density versus the DM mass (left) and versus the couplings combination  $\sum_{i,\alpha} |h_{\alpha,i}|^2$  (right). The yellow band represents the observed the DM relic density value. One has to mention that in the left panel, we make all masses changing simultaneously, i.e., all the mass fulfil the condition  $m_i/m_{DM} = \text{constant}$  with all the DM mass interval.

)annihilation is simple, i.e.,  $N_1 N_1 (N_{2,3}) \rightarrow \nu_\alpha \bar{\nu}_\beta, \ell_\alpha \ell_\beta$ . However, in the scalar DM case, its dependence on the DM mass is not trivial due to the existence of many annihilation channels ( $W^\pm W^\mp, ZZ, q_i \bar{q}_i, hh, \bar{\ell}_\alpha \ell_\beta$  and  $\bar{\nu}_\alpha \nu_\beta$ ), where each channel could be dominant within a specific DM mass range. From Fig. 5-right, one learns that in the scalar DM case the DM annihilation into  $\nu_\alpha \bar{\nu}_\beta, \ell_\alpha \ell_\beta$  could be efficient only if  $\sum_{i,\alpha} |h_{\alpha,i}|^2 \sim 0.1$ . Clearly, the right amount of the DM relic density can be easily achieved in this model whether the DM is a Majorana or a scalar particle.

## 6 Conclusion

In this paper, we proposed a SM extension by three singlet Majorana fermions and two scalar inert doublets. After diagonalizing the scalar squared mass sub-matrices, we found that the model includes two CP-even ( $H_{1,2}^0$ ), two CP-odd ( $A_{1,2}^0$ ) and two charged scalar ( $H_{1,2}^\pm$ ) eigenstates, with different mixing angles. Then, the values of many physical observables got modified with respect to the MSM. For

instance, the neutrino mass is generated at one-loop level via 12 diagrams instead of 6 in the MSM. Similar remarks hold for the observables relevant to the electroweak precision tests, LFV and the di-photon Higgs decay, where many masses and mixing angles are included. This made the neutrino oscillation data easily explained without being in conflict with the different theoretical and experimental constraints for large parameter space regions.

The dark matter (DM) in this model could be either fermionic ( $N_1$ ) or scalar (the lightest among  $H_1^0$  and  $A_1^0$ ). We have shown that the right amount of the relic density can be easily achieved in both scenarios for different DM mass ranges. In addition, for the scalar DM scenario the model can accommodate the constraints for the direct detection cross section of the DM-nucleon scattering.

The model predicts all the signatures predicted by the MSM, however, novel signature relevant to the production and decay of the heavy scalars ( $H_2^\pm, H_2^0, A_2^0$ ) can be probed through some final states like:  $8 \text{ jets} + \cancel{E}_T, 1\ell + 4 \text{ jets} + \cancel{E}_T, 1\ell + 2 \text{ jets} + \cancel{E}_T, 4b + \cancel{E}_T, 2 \text{ jets} + 2b + \cancel{E}_T$  and  $2b + \cancel{E}_T$ . This point requires an independent investigation.

## Acknowledgments

This work is funded by the University of Sharjah under the research projects No 21021430100 “*Extended Higgs Sectors at Colliders: Constraints & Predictions*” and No 21021430107 “*Hunting for New Physics at Colliders*”.

## A Generalized Casas-Ibarra parameterization

Following the Casas-Ibarra approach [17], one assumes that the new Yukawa couplings matrix is a product of many matrices as  $h = S_1.S_2.S_3.S_4.S_5$ , then replace this in (2.14); and match the result with the known neutrino mass formula  $(m^{(\nu)})_{3\times 3} = (U_\nu)_{3\times 3}.(D_{m_\nu})_{3\times 3}.(U_\nu^T)_{3\times 3}$ , with  $U_\nu$  is the PMNS mixing matrix.

The most general solution for the matrix  $h$  implies  $S_1 = (U_\nu)_{3\times 3}, S_2 = (D_{\sqrt{m_\nu}})_{3\times 3}, S_3 = (T)_{3\times M}, S_4 = (D_{(\Lambda'_i)^{-1/2}})_{M\times M}$  and  $S_5 = (Q)_{M\times M}$ , with the conditions  $(Q)_{M\times M}.\Lambda_{M\times M}.(Q^T)_{M\times M} = D_{(\Lambda'_i)}$  and  $(T)_{3\times M}.(T^T)_{M\times 3} = \mathbf{1}_{3\times 3}$  where  $\Lambda'_i$  are the eigenvalues of the matrix  $\Lambda$ .

Here,  $T$  is a  $3 \times M$  matrix that includes the identity matrix  $\mathbf{1}_{3\times 3}$  with the remaining elements is vanishing, so it fulfils the condition  $(T)_{3\times M}.(T^T)_{M\times 3} = \mathbf{1}_{3\times 3}$ . In case of SM extended by  $n$  inert doublet, we have  $M = 3 \times n$ , and therefore the matrix  $T$  can be presented as

$$T_3 = \left( \alpha^{(1)} R_{3\times 3}^{(1)}, \alpha^{(2)} R_{3\times 3}^{(2)}, \dots \alpha^{(n)} R_{3\times 3}^{(n)} \right), \quad (\text{A.1})$$

with  $R^{(i)}$  are  $3 \times 3$  orthogonal matrices  $R^{(i)}.R^{(i)T} = R^{(i)T}.R^{(i)} = \mathbf{1}$ , and  $\alpha^{(i)}$  are real numbers that fulfil the identity  $\sum_i (\alpha^{(i)})^2 = 1$ . For collider purpose or Majorana DM scenario, the Yukawa couplings should be relatively large, and in order to ensure this, it is enough to choose a diagonalization matrix  $(Q)_{M\times M}$  that leads to the order  $|\Lambda'_1| < |\Lambda'_2| < \dots \ll |\Lambda'_M|$ ; and  $\alpha^{(1)} = 1, \alpha^{(i\neq 1)} = 0$ .

## B Scotogenic Model with $n$ inert Doublets

In case of a scotogenic model with  $N$  inert doublets ( $\Phi_{i=1,n}$ ) with the global  $Z_2$  symmetry  $N_i \rightarrow -N_i$ ,  $\Phi_i \rightarrow -\Phi_i$ , the Lagrangian (2.1) can be generalized as

$$\mathcal{L} \supset \sum_{i=1}^n \sum_{k=1}^3 \bar{L}_\alpha ((h_{\alpha,k+3(i-1)}) \epsilon \Phi_i) N_k + \frac{1}{2} \bar{N}_k^C M_k N_k + h.c., \quad (\text{B.1})$$

where the new Yukawa couplings  $h_{\alpha,i}$  here are a  $3 \times (3n)$  matrix. The scalar potential (2.2) can be generalized as

$$\begin{aligned} V(H, \Phi_i, S, \chi) = & -\mu_H^2 |\mathcal{H}|^2 + \mu_i^2 |\Phi_i|^2 + \frac{\lambda_H}{6} |\mathcal{H}|^4 + \frac{\lambda_i}{6} |\Phi_i|^4 + \omega_i |\mathcal{H}|^2 |\Phi_i|^2 + \kappa_i |\mathcal{H}^\dagger \Phi_i|^2 \\ & + \varrho_{i,j}^{(1)} |\Phi_i|^2 |\Phi_j|^2 + \varrho_{i,j}^{(2)} |\Phi_i^\dagger \Phi_j|^2 + \left\{ \mu_{3,ij}^2 \Phi_i^\dagger \Phi_j + \frac{1}{2} \tilde{\zeta}_i^{(1)} (\mathcal{H}^\dagger \Phi_i)^2 \right. \\ & \left. + \tilde{\zeta}_{i,j}^{(2)} (\mathcal{H}^\dagger \Phi_i) (\mathcal{H}^\dagger \Phi_j) + \tilde{\zeta}_{i,j}^{(3)} (\Phi_i^\dagger \mathcal{H}) (\mathcal{H}^\dagger \Phi_j) + h.c. \right\}. \end{aligned} \quad (\text{B.2})$$

In case of spontaneous CP-conservation (real values of all potential parameters), there are  $n$  charged eigenstates ( $H_{j=1,n}^\pm$ ),  $n$  neutral CP-even eigenstates ( $H_{j=1,n}^0$ ) and  $n$  neutral CP-odd eigenstates ( $A_{j=1,n}^0$ ) with similar couplings to (2.11). In (2.11), the mixing ( $c_{H,A,C}$  and  $s_{H,A,C}$  should be replaced by the elements of the mixing matrices  $\mathcal{U}^{(H,A,C)}$  that diagonalize the CP-even, CP-odd and charged scalars squared  $n \times n$  mass matrices, respectively, where the summation should be performed over  $j = 1, n$ . Consequently, the neutrino mass is generated via  $4 \times n$  diagrams mediated by  $N_{1,2,3}$  and  $H_{j=1,n}^0 / A_{j=1,n}^0$ . Therefore, the formulas of the mass matrix elements (2.12) and the values of the new Yukawa couplings in (2.17) hold for  $j = 1, n$  and  $M = 3 \times n$ . Similar conclusions can be achieved for the LFV branching ratios (3.5) and the di-photon Higgs decay ratio (3.7). For the oblique parameters (3.6), the summation should be performed over  $j = 1, n$ ; and the mixing ( $c_{H,A,C}$  and  $s_{H,A,C}$  should be replaced by the elements of the mixing matrix  $\mathcal{U}^{(H,A,C)}$ .

## C The Unitarity Amplitude Matrices

Here, we give the relevant amplitude matrices for the unitarity. We have the following basis that characterize the initial/final states:

**CP-even,  $Z_2$  even, &  $Q_{em} = 0$ :** the basis is  $\{hh, \chi^0 \chi^0, S_1^0 S_1^0, S_2^0 S_2^0, Q_1^0 Q_1^0, Q_2^0 Q_2^0, \chi^+ \chi^-, S_1^+ S_1^-, S_2^+ S_2^-, S_1^+ S_2^-\}$ ; and the matrix is

$$\begin{bmatrix} \lambda_H, & \frac{1}{3}\lambda_H, & \omega_1 + \kappa_1 + \xi_1, \omega_2 + \kappa_2 + \xi_2, \omega_1 + \kappa_1 - \xi_1, \omega_2 + \kappa_2 - \xi_2, & \frac{1}{3}\lambda_H, & \omega_1, & \omega_2, & 0 \\ \frac{1}{3}\lambda_H, & \lambda_H, & \omega_1 + \kappa_1 - \xi_1, \omega_2 + \kappa_2 - \xi_2, \omega_1 + \kappa_1 + \xi_1, \omega_2 + \kappa_2 + \xi_2, & \frac{1}{3}\lambda_H, & \omega_1, & \omega_2, & 0 \\ \omega_1 + \kappa_1 + \xi_1, \omega_1 + \kappa_1 - \xi_1, & \lambda_1, & \varrho_1 + \varrho_2, & \frac{1}{3}\lambda_1, & \varrho_1 + \varrho_2, & \omega_1, & \frac{1}{3}\lambda_1, & \varrho_1, & 0 \\ \omega_2 + \kappa_2 + \xi_2, \omega_2 + \kappa_2 - \xi_2, & \varrho_1 + \varrho_2, & \lambda_2, & \varrho_1 + \varrho_2, & \frac{1}{3}\lambda_2, & \omega_2, & \varrho_1, & \frac{1}{3}\lambda_2, & 0 \\ \omega_1 + \kappa_1 - \xi_1, \omega_1 + \kappa_1 + \xi_1, & \frac{1}{3}\lambda_1, & \varrho_1 + \varrho_2, & \lambda_1, & \varrho_1 + \varrho_2, & \omega_1, & \frac{1}{3}\lambda_1, & \varrho_1, & 0 \\ \omega_2 + \kappa_2 - \xi_2, \omega_2 + \kappa_2 + \xi_2, & \varrho_1 + \varrho_2, & \frac{1}{3}\lambda_2, & \varrho_1 + \varrho_2, & \lambda_2, & \omega_2, & \varrho_1, & \frac{1}{3}\lambda_2, & 0 \\ \frac{1}{3}\lambda_H, & \frac{1}{3}\lambda_H, & \omega_1, & \omega_2, & \omega_1, & \omega_2, & \frac{2}{3}\lambda_H, & \kappa_1 + \omega_1, \kappa_2 + \omega_2, & \xi_4 \\ \omega_1, & \omega_1, & \frac{1}{3}\lambda_1, & \varrho_1, & \frac{1}{3}\lambda_1, & \varrho_1, & \kappa_1 + \omega_1, & \frac{2}{3}\lambda_1, & \varrho_1 + \varrho_2, & 0 \\ \omega_2, & \omega_2, & \varrho_1, & \frac{1}{3}\lambda_2, & \varrho_1, & \frac{1}{3}\lambda_2, & \kappa_2 + \omega_2, & \varrho_1 + \varrho_2, & \frac{2}{3}\lambda_2, & 0 \\ 0, & 0, & 0, & 0, & 0, & 0, & \xi_4, & 0, & 0, & \varrho_1 + \varrho_2 \end{bmatrix} \quad (\text{C.1})$$

**CP-even,  $Z_2$  even, &  $Q_{em} = 0$ :** the basis is  $\{hS_1^0, hS_2^0, \chi^0 Q_1^0, \chi^0 Q_2^0, \chi^+ S_1^-, \chi^+ S_2^-\}$ ; and the matrix is

$$\begin{bmatrix} \omega_1 + \kappa_1 + \zeta_1, & \zeta_3 + \zeta_4, & \zeta_1, & \zeta_3, & \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4) \\ \zeta_3 + \zeta_4, & \omega_2 + \kappa_2 + \zeta_2, & \zeta_3, & \zeta_2, & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2) \\ \zeta_1, & \zeta_3, & \omega_1 + \kappa_1 + \zeta_1, & \zeta_3 + \zeta_4, & \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4) \\ \zeta_3, & \zeta_2, & \zeta_3 + \zeta_4, & \omega_2 + \kappa_2 + \zeta_2, & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2) \\ \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4), & \kappa_1 + \omega_1, & \zeta_4 \\ \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2), & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2), & \zeta_4, & \kappa_2 + \omega_2 \end{bmatrix} \quad (\text{C.2})$$

**CP-even,  $Z_2$  even, &  $Q_{em} = 0$ :** the basis is  $\{h\chi^0, S_1^0 Q_1^0, S_1^0 Q_2^0, S_2^0 Q_1^0, S_2^0 Q_2^0, \chi^+ \chi^-, S_1^+ S_1^-, S_2^+ S_2^-, S_1^+ S_2^-\}$ ; and the matrix is

$$\begin{bmatrix} \frac{1}{3}\lambda_H, & \zeta_1, & \zeta_3, & \zeta_3, & \zeta_2, & 0, & 0, & 0, & 0 \\ \zeta_1, & \frac{1}{3}\lambda_1, & 0, & 0, & 0, & 0, & 0, & 0, & 0 \\ \zeta_3, & 0, & \varrho_1 + \varrho_2, & 0, & 0, & 0, & 0, & 0, & -\frac{i}{2}\varrho_2 \\ \zeta_3, & 0, & 0, & \varrho_1 + \varrho_2, & 0, & 0, & 0, & 0, & \frac{i}{2}\varrho_2 \\ \zeta_2, & 0, & 0, & 0, & \frac{1}{3}\lambda_2, & 0, & 0, & 0, & 0 \\ 0, & 0, & 0, & 0, & 0, & \frac{2}{3}\lambda_H, & \kappa_1 + \omega_1, & \kappa_2 + \omega_2, & \zeta_4 \\ 0, & 0, & 0, & 0, & 0, & \kappa_1 + \omega_1, & \frac{2}{3}\lambda_1, & \varrho_1 + \varrho_2, & 0 \\ 0, & 0, & 0, & 0, & 0, & \kappa_2 + \omega_2, & \varrho_1 + \varrho_2, & \frac{2}{3}\lambda_2, & 0 \\ 0, & 0, & \frac{i}{2}\varrho_2, & -\frac{i}{2}\varrho_2, & 0, & \zeta_4, & 0, & 0, & \varrho_1 + \varrho_2 \end{bmatrix} \quad (\text{C.3})$$

**CP-even,  $Z_2$  even, &  $Q_{em} = 0$ :** the basis is  $\{hQ_1^0, hQ_2^0, S_1^0 \chi^0, S_2^0 \chi^0, \chi^+ S_1^-, \chi^+ S_2^-\}$ ; and the matrix is

$$\begin{bmatrix} \omega_1 + \kappa_1 - \zeta_1, & -\zeta_3 + \zeta_4, & \zeta_1, & \zeta_3, & -\frac{i}{2}(\kappa_1 - \zeta_1), & \frac{i}{2}(\zeta_3 - \zeta_4) \\ -\zeta_3 + \zeta_4, & \omega_2 + \kappa_2 - \zeta_2, & \zeta_3, & \zeta_2, & \frac{i}{2}(\zeta_3 - \zeta_4), & -\frac{i}{2}(\kappa_2 - \zeta_2) \\ \zeta_1, & \zeta_3, & \omega_1 + \kappa_1 - \zeta_1, & -\zeta_3 + \zeta_4, & \frac{i}{2}(\kappa_1 - \zeta_1), & -\frac{i}{2}(\zeta_3 - \zeta_4) \\ \zeta_3, & \zeta_2, & -\zeta_3 + \zeta_4, & \omega_2 + \kappa_2 - \zeta_2, & -\frac{i}{2}(\zeta_3 - \zeta_4), & \frac{i}{2}(\kappa_2 - \zeta_2) \\ \frac{i}{2}(\kappa_1 - \zeta_1), & -\frac{i}{2}(\zeta_3 - \zeta_4), & -\frac{i}{2}(\kappa_1 - \zeta_1), & \frac{i}{2}(\zeta_3 - \zeta_4), & \kappa_1 + \omega_1, & \zeta_4 \\ -\frac{i}{2}(\zeta_3 - \zeta_4), & \frac{i}{2}(\kappa_2 - \zeta_2), & \frac{i}{2}(\zeta_3 - \zeta_4), & -\frac{i}{2}(\kappa_2 - \zeta_2), & \zeta_4, & \kappa_2 + \omega_2 \end{bmatrix} \quad (\text{C.4})$$

**CP-even,  $Z_2$  even, &  $Q_{em} = 0$ :** the basis is  $\{h\chi^+, \chi^0 \chi^+, S_1^0 S_1^+, S_1^0 S_2^+, S_2^0 S_1^+, S_2^0 S_2^+, Q_1^0 S_1^+, Q_1^0 S_2^+, Q_2^0 S_1^+, Q_2^0 S_2^+\}$ ; and the matrix is

$$\begin{bmatrix} \frac{1}{3}\lambda_H, & 0, & \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2), & \frac{i}{2}(\kappa_1 - \zeta_1), & -\frac{i}{2}(\zeta_3 - \zeta_4), & -\frac{i}{2}(\zeta_3 - \zeta_4), & \frac{i}{2}(\kappa_2 - \zeta_2) \\ 0, & \frac{1}{3}\lambda_H, & -\frac{i}{2}(\kappa_1 - \zeta_1), & \frac{i}{2}(\zeta_3 - \zeta_4), & \frac{i}{2}(\zeta_3 - \zeta_4), & -\frac{i}{2}(\kappa_2 - \zeta_2), & \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2) \\ \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{i}{2}(\kappa_1 - \zeta_1), & \frac{1}{3}\lambda_1, & 0, & 0, & \frac{1}{2}\varrho_2, & 0, & 0, & 0, & \frac{i}{2}\varrho_2 \\ \frac{1}{2}(\zeta_3 + \zeta_4), & -\frac{i}{2}(\zeta_3 - \zeta_4), & 0, & \varrho_1, & \frac{1}{2}\varrho_2, & 0, & 0, & 0, & -\frac{i}{2}\varrho_2, & 0 \\ \frac{1}{2}(\zeta_3 + \zeta_4), & -\frac{i}{2}(\zeta_3 - \zeta_4), & 0, & \frac{1}{2}\varrho_2, & \varrho_1, & 0, & -\frac{i}{2}\varrho_2, & 0, & 0, & 0 \\ \frac{1}{2}(\kappa_2 + \zeta_2), & \frac{i}{2}(\kappa_2 - \zeta_2), & \frac{1}{2}\varrho_2, & 0, & 0, & \frac{1}{3}\lambda_2, & \frac{i}{2}\varrho_2, & 0, & 0, & 0 \\ -\frac{i}{2}(\kappa_1 - \zeta_1), & \frac{1}{2}(\kappa_1 + \zeta_1), & 0, & 0, & 0, & -\frac{i}{2}\varrho_2, & \frac{1}{3}\lambda_1, & 0, & 0, & \frac{1}{2}\varrho_2 \\ \frac{i}{2}(\zeta_3 - \zeta_4), & \frac{1}{2}(\zeta_3 + \zeta_4), & 0, & 0, & \frac{i}{2}\varrho_2, & 0, & 0, & \varrho_1, & \frac{1}{2}\varrho_2, & 0 \\ \frac{i}{2}(\zeta_3 - \zeta_4), & \frac{1}{2}(\zeta_3 + \zeta_4), & 0, & \frac{i}{2}\varrho_2, & 0, & 0, & 0, & \frac{1}{2}\varrho_2, & \varrho_1, & 0 \\ -\frac{i}{2}(\kappa_2 - \zeta_2), & \frac{1}{2}(\kappa_2 + \zeta_2), & -\frac{i}{2}\varrho_2, & 0, & 0, & 0, & \frac{1}{2}\varrho_2, & 0, & 0, & \frac{1}{3}\lambda_2 \end{bmatrix} \quad (\text{C.5})$$



**CP-even,  $Z_2$  even, &  $Q_{em} = 0$ :** the basis is  $\{hS_1^+, hS_2^+, \chi^0 S_1^+, \chi^0 S_2^+, S_1^0 \chi^+, S_2^0 \chi^+, Q_1^0 \chi^+, Q_2^0 \chi^+\}$ ; and the matrix is

$$\begin{bmatrix} \omega_1, & 0, & 0, & 0, & \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4), & -\frac{i}{2}(\kappa_1 - \zeta_1), & \frac{i}{2}(\zeta_3 - \zeta_4) \\ 0, & \omega_2, & 0, & 0, & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2), & \frac{i}{2}(\zeta_3 - \zeta_4), & -\frac{i}{2}(\kappa_2 - \zeta_2) \\ 0, & 0, & \omega_1, & 0, & \frac{i}{2}(\kappa_1 - \zeta_1), & -\frac{i}{2}(\zeta_3 - \zeta_4), & \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4) \\ 0, & 0, & 0, & \omega_2, & -\frac{i}{2}(\zeta_3 - \zeta_4), & \frac{i}{2}(\kappa_2 - \zeta_2), & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2) \\ \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4), & -\frac{i}{2}(\kappa_1 - \zeta_1), & \frac{i}{2}(\zeta_3 - \zeta_4), & \omega_1, & 0, & 0, & 0 \\ \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2), & \frac{i}{2}(\zeta_3 - \zeta_4), & -\frac{i}{2}(\kappa_2 - \zeta_2), & 0, & \omega_2, & 0, & 0 \\ \frac{i}{2}(\kappa_1 - \zeta_1), & -\frac{i}{2}(\zeta_3 - \zeta_4), & \frac{1}{2}(\kappa_1 + \zeta_1), & \frac{1}{2}(\zeta_3 + \zeta_4), & 0, & 0, & \omega_1, & 0 \\ -\frac{i}{2}(\zeta_3 - \zeta_4), & \frac{i}{2}(\kappa_2 - \zeta_2), & \frac{1}{2}(\zeta_3 + \zeta_4), & \frac{1}{2}(\kappa_2 + \zeta_2), & 0, & 0, & 0, & \omega_2 \end{bmatrix} \quad (\text{C.6})$$

## References

- [1] R. N. Mohapatra, Phys. Rev. Lett. **56** (1986), 561-563. R. N. Mohapatra and J. W. F. Valle, Phys. Rev. D **34** (1986), 1642
- [2] A. Zee, Phys. Lett. **161B**, 141 (1985). E. Ma, Phys. Rev. Lett. **81**, 1171 (1998) [hep-ph/9805219].
- [3] A. Zee, Nucl. Phys. **B 264**, 99 (1986); K. S. Babu, Phys. Lett. B **203**, 132 (1988); M. Aoki, S. Kanemura, T. Shindou, and K. Yagyu, J. High Energy Phys. **10 1007** (2010) 084; J. High Energy Phys. **11** (2010) 049; G. Guo, X. G. He, and G. N. Li, J. High Energy Phys. **10** (2012) 044; Y. Kajiyama, H. Okada, and K. Yagyu, Nucl. Phys. **B874**, 198 (2013).
- [4] M. Aoki, S. Kanemura, and O. Seto, Phys. Rev. Lett. **102**, 051805 (2009); M. Aoki, S. Kanemura, and O. Seto, Phys. Rev. D **80**, 033007 (2009).
- [5] L. M. Krauss, S. Nasri, and M. Trodden, Phys. Rev. D **67**, 085002 (2003). A. Ahriche and S. Nasri, JCAP **07** (2013), 035 [arXiv:1304.2055 [hep-ph]]. A. Ahriche, K. L. McDonald and S. Nasri, Phys. Rev. D **92** (2015) no.9, 095020 [arXiv:1508.05881 [hep-ph]].
- [6] T. Nomura and H. Okada, Phys. Lett. B **755**, 306 (2016) [arXiv:1601.00386 [hep-ph]].
- [7] K. Cheung and O. Seto, Phys. Rev. D **69**, 113009 (2004). A. Ahriche, S. Nasri, and R. Soualah, Phys. Rev. D **89**, 095010 (2014), C. Guella, D. Cherigui, A. Ahriche, S. Nasri and R. Soualah, Phys. Rev. D **93**, no. 9, 095022 (2016) [arXiv:1601.04342 [hep-ph]], D. Cherigui, C. Guella, A. Ahriche and S. Nasri, Phys. Lett. B **762**, 225 (2016) [arXiv:1605.03640 [hep-ph]],
- [8] Y. Cai, J. Herrero-Garcia, M. A. Schmidt, A. Vicente and R. R. Volkas, Front. in Phys. **5**, 63 (2017) [arXiv:1706.08524 [hep-ph]].
- [9] E. Ma, Phys. Rev. D **73** (2006), 077301 [arXiv:hep-ph/0601225 [hep-ph]].
- [10] A. Ahriche, A. Jueid and S. Nasri, Phys. Rev. D **97** (2018) no.9, 095012 [arXiv:1710.03824 [hep-ph]].
- [11] A. Ahriche, A. Arhrib, A. Jueid, S. Nasri and A. de La Puente, Phys. Rev. D **101** (2020) no.3, 035038 [arXiv:1811.00490 [hep-ph]].
- [12] A. Ahriche, A. Jueid and S. Nasri, Phys. Lett. B **814** (2021), 136077 [arXiv:2007.05845 [hep-ph]].
- [13] A. Ahriche, K. L. McDonald and S. Nasri, JHEP **06** (2016), 182 [arXiv:1604.05569 [hep-ph]]. A. Beniwal, J. Herrero-García, N. Leerdam, M. White and A. G. Williams, JHEP **21** (2020), 136 [arXiv:2010.05937 [hep-ph]]. D. M. Barreiros, F. R. Joaquim, R. Srivastava and J. W. F. Valle, JHEP **04** (2021), 249

- [arXiv:2012.05189 [hep-ph]]. P. Escribano, M. Reig and A. Vicente, JHEP **07** (2020), 097 [arXiv:2004.05172 [hep-ph]]. Z. L. Han, R. Ding, S. J. Lin and B. Zhu, Eur. Phys. J. C **79** (2019) no.12, 1007 [arXiv:1908.07192 [hep-ph]]. C. H. Chen and T. Nomura, JHEP **10** (2019), 005 [arXiv:1906.10516 [hep-ph]]. Z. L. Han and W. Wang, Eur. Phys. J. C **79** (2019) no.6, 522 [arXiv:1901.07798 [hep-ph]]. W. Wang, R. Wang, Z. L. Han and J. Z. Han, Eur. Phys. J. C **77** (2017) no.12, 889 [arXiv:1705.00414 [hep-ph]]. D. M. Barreiros, H. B. Camara and F. R. Joaquim, [arXiv:2204.13605 [hep-ph]].
- [14] J. Aalbers *et al.* [LUX-ZEPLIN], [arXiv:2207.03764 [hep-ex]].
- [15] B. Grzadkowski, O. M. Ogreid and P. Osland, Phys. Rev. D **80** (2009), 055013 [arXiv:0904.2173 [hep-ph]].
- [16] L. Sarma, P. Das and M. K. Das, Nucl. Phys. B **963** (2021), 115300 [arXiv:2004.13762 [hep-ph]]. L. Sarma, B. B. Boruah and M. K. Das, Eur. Phys. J. C **82** (2022) no.5, 488 [arXiv:2106.04124 [hep-ph]].
- [17] J. A. Casas and A. Ibarra, Nucl. Phys. B **618** (2001), 171-204 [arXiv:hep-ph/0103065 [hep-ph]].
- [18] T. Toma and A. Vicente, JHEP **01** (2014), 160 [arXiv:1312.2840 [hep-ph]].
- [19] M. Chekkal, A. Ahriche, A. B. Hammou and S. Nasri, Phys. Rev. D **95**, no. 9, 095025 (2017). [arXiv:1702.04399 [hep-ph]].
- [20] J. Abdallah *et al.* [DELPHI], Eur. Phys. J. C **31** (2003), 421-479 [arXiv:hep-ex/0311019 [hep-ex]].
- [21] E. Lundstrom, M. Gustafsson and J. Edsjo, Phys. Rev. D **79** (2009), 035013 [arXiv:0810.3924 [hep-ph]].
- [22] W. Grimus, L. Lavoura, O. M. Ogreid and P. Osland, Nucl. Phys. B **801** (2008), 81-96 [arXiv:0802.4353 [hep-ph]].
- [23] [ATLAS], ATLAS-CONF-2017-047.
- [24] A. Djouadi, Phys. Rept. **457** (2008), 1-216 [arXiv:hep-ph/0503172 [hep-ph]].
- [25] [ATLAS], ATLAS-CONF-2018-031.
- [26] X. G. He, T. Li, X. Q. Li, J. Tandean and H. C. Tsai, Phys. Rev. D **79** (2009), 023521 [arXiv:0811.0658 [hep-ph]].
- [27] Y. Meng *et al.* [PandaX-4T], Phys. Rev. Lett. **127** (2021) no.26, 261802 [arXiv:2107.13438 [hep-ex]].
- [28] E. Aprile *et al.* [XENON], JCAP **04** (2016), 027 [arXiv:1512.07501 [physics.ins-det]].
- [29] J. Billard, L. Strigari and E. Figueroa-Feliciano, Phys. Rev. D **89** (2014) no.2, 023524 [arXiv:1307.5458 [hep-ph]].
- [30] F. Ambrogio, C. Arina, M. Backovic, J. Heisig, F. Maltoni, L. Mantani, O. Mattelaer and G. Mohlabeng, Phys. Dark Univ. **24** (2019), 100249 [arXiv:1804.00044 [hep-ph]].
- [31] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr and B. Fuks, Comput. Phys. Commun. **185** (2014), 2250-2300 [arXiv:1310.1921 [hep-ph]].