

EXPLICIT CONDITIONS FOR THE CLT AND RELATED RESULTS FOR NON-UNIFORMLY PARTIALLY EXPANDING RANDOM DYNAMICAL SYSTEMS VIA EFFECTIVE RPF RATES

YEOR HAFOUTA

DEPARTMENT OF MATHEMATICS
THE OHIO STATE UNIVERSITY AND THE UNIVERSITY OF MARYLAND

ABSTRACT. The purpose of this paper is to provide a first class of explicit sufficient conditions for the central limit theorem and related results in the setup of non-uniformly (partially) expanding non iid random transformations, considered as stochastic processes together with some random Gibbs measure. More precisely, we prove a central limit theorem (CLT), an almost sure invariance principle, a moderate deviations principle, Berry-Esseen type estimates and a moderate local central limit theorem for random Birkhoff sums generated by a non-uniformly partially expanding dynamical systems T_ω and a random Gibbs measure μ_ω corresponding to a random potential ϕ_ω with a sufficiently regular variation. In the partially expanding case the maps we consider are similar to the ones in [44], with the exception that the amount of expansion dominates the amount of contraction fiberwise and not only on the average and with an additional regularity condition on a certain type of local variation of ϕ_ω along inverse branches of T_ω . A notable example when the maps are truly partially-expanding is the case when $\phi_\omega \equiv 0$ which corresponds to random measures of maximal entropy μ_ω , but any potential with a sufficiently small (fiberwise) variation can be considered. Our results in the partially expanding case are new even in the uniformly random case, where all the random variables describing the maps are uniformly controlled. For properly expanding maps (as in [42, 33]), the above local regularity condition allows applications also in the smooth case where the Gibbs measure is absolutely continuous with respect to the underlying volume measure and $\phi_\omega = -\ln J_{T_\omega}$. For instance, we can consider certain fiberwise piecewise C^2 -perturbations of piecewise linear or affine maps. All of the above is achieved by first proving random, real and complex, Ruelle-Perron-Frobenius (RPF) theorems with rates that can be expressed analytically by means of certain random parameters that describe the maps (such rates will be referred to as “effective”). Using these effective rates, our conditions for the limit theorems involve some weak type of upper mixing conditions on the driving system (base map) and some integrability conditions on the norm of the random function generating the Birkhoff sums. A big part of the proof of the moderate deviations, the Berry-Esseen type estimates and the local CLT is to show how Rugh’s theory [43] of complex cones contractions applies to the cones considered in [16] (and their random versions in [44]), which is new even for deterministic dynamical systems T and that case it yields explicit estimates on the spectral gap of appropriate deterministic complex perturbations of the transfer operator of T , as well as explicit constants in the corresponding Berry-Esseen theorem for deterministic partially expanding dynamical systems.

1. INTRODUCTION AND AN OVERVIEW OF THE MAIN RESULTS

1.1. Statistical properties of random Birkhoff sums and a preview. Probabilistic limit theorems for expanding random dynamical systems have been studied extensively in the past decades. This setup includes a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and a family locally expanding maps $T_\omega, \omega \in \Omega$ which are composed along an orbit of a probability preserving invertible and ergodic map $\theta : \Omega \rightarrow \Omega$ together with a family of equivariant¹ probability measures μ_ω (i.e. $(T_\omega)_* \mu_\omega = \mu_{\theta\omega}$ for \mathbb{P} -a.a. ω) on the domain

Date: May 11, 2023.

¹As will be discussed below, in applications $\{\mu_\omega\}$ is not just any equivariant family, but it is generated by an appropriate random potential (i.e. random Gibbs measures). In other situations μ_ω can be the appropriate volume measure.

\mathcal{E}_ω of T_ω . When considering a random point x_0 distributed according to μ_ω we get random orbits

$$T_\omega^n x_0 = T_{\theta^{n-1}\omega} \circ \dots \circ T_{\theta\omega} \circ T_\omega x_0$$

and the question is whether for \mathbb{P} -almost all $\omega \in \Omega$ random Birkhoff sums of the form $S_n^\omega = S_n^\omega u(x_0) = \sum_{j=0}^{n-1} u_{\theta^j\omega} \circ T_\omega^j(x_0)$ obey limit theorems like the quenched central limit theorem² (CLT) and its variety of stronger versions. Here u_ω is random function on the domain of T_ω satisfying some regularity conditions like Hölder continuity (not necessarily uniformly in ω).

1.1.1. *An illustrating example.* In this section we will give an example of a random map $T_\omega : [0, 1) \rightarrow [0, 1)$ which already captures the essence of the problem addressed in this manuscript. Let a_ω be a random variable taking values in $[\frac{1}{2}, 1)$. Let us consider the piecewise linear map T_ω on $[0, 1)$ so that on $[0, a_\omega)$ the map T_ω coincides with the linear function connecting the points $(0, 0)$ and $(a_\omega, 1)$, while on $[a_\omega, 1)$ it coincides with the linear function connecting the points $(a_\omega, 0)$ and $(1, 1)$. Then $T_\omega([0, 1)) = [0, 1)$ and $\gamma_\omega := a_\omega^{-1}$ is the minimal amount of expansion of the map T_ω . When a_ω is bounded away from 1 then the maps T_ω are uniformly expanding, and statistical properties (i.e. limit theorems) for random Birkhoff sums were extensively studied³ for random functions u_ω with uniformly bounded Hölder norms $\|u_\omega\|$ (here $\mu_\omega = \text{Lebesgue}$). However, when $\text{esssup}_{\omega \in \Omega}(a_\omega) = 1$ the maps T_ω are not uniformly expanding. Focusing for the moment on this example, in this paper we will prove limit theorems in the case when a_ω can take arbitrarily close to 1 values (i.e. we can have $\text{esssup}_{\omega \in \Omega}(a_\omega) = 1$), and the random variable $\omega \rightarrow \|u_\omega\|$ belongs to L^p for some $p > 2$ (how small can p be depends on the result, for the CLT we have sufficient conditions for every $p > 2$). Our results will be obtained under some (upper) mixing related assumptions on the sequence of random variables $\{a_{\theta^j\omega}, j \geq 0\}$. For instance, the quenched CLT holds when $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ is the Markov shift corresponding to a sufficiently fast mixing Markov chain X_n (e.g. geometrically ergodic) and $\beta(r) =: \|a_\omega - \mathbb{E}[a_\omega | X_{-r}, \dots, X_r]\|_{L^1}$, $\omega = (X_j)_{j \in \mathbb{Z}}$ decays polynomially fast as $r \rightarrow \infty$. Note that when a_ω depends only on finitely many coordinates then $\beta(r) = 0$ for r large enough, while when it is a Hölder continuous function of ω (in the case when the chain takes values on some metric space) then $\beta(r)$ decays exponentially fast (in this case we can take any $p > 2$ above). We refer to Examples 22, 25 and 26 for more details.

1.1.2. *Back to the general setup.* In general, the maps T_ω can very often be described by means of random parameters $a_{i,\omega}, i \leq d$ such as minimal amount of local expansion, degree, “ratio” between contraction and expansion, local variation of the logarithm of the Jacobian, etc (in the example in Section 1.1.1 we can take $d = 1$ and $a_{1,\omega} = a_\omega$). We call the maps uniformly random when the random variables $a_{i,\omega}$ take values on appropriate domains (e.g. minimal local expansion is bounded away from 1, bounded degree, bounded variation, etc.). In the example in Section 1.1.1 the uniform case corresponds to the case when $a_\omega \leq 1 - \epsilon$ for some $\epsilon \in (0, \frac{1}{2})$ and \mathbb{P} -a.a. ω . Then most of the statistical properties in literature were obtained in the uniformly random case⁴, with the exception⁵ of certain types of maps so that $\{T_{\theta^j\omega} : 0 \leq j < \infty\}$ and $\{u_{\theta^j\omega} : 0 \leq j < \infty\}$ are iid processes on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Beyond the iid case we are not aware of even a single explicit example with a true non-uniform behavior for which the quenched CLT holds true.

The purpose of this manuscript is to provide explicit sufficient conditions for several limit theorems like the CLT for non-uniformly random (partially) expanding maps (which will provide a variety of examples beyond the iid case). Let us note that in [39, Theorem 2.3] an inducing strategy was developed in order to prove the CLT and related results in the non-uniformly random case. The conditions in [39, Theorem 2.3] require certain type of regularity of the behavior of the first visiting time to a measurable set $A \subset \Omega$ with positive probability so that $\{T_\omega, \omega \in A\}$ are uniformly expanding in an appropriate sense. While the results in [39] were new even in the uniformly random case, to the best

²Let us recall that the quenched CLT means that for \mathbb{P} -a.a. ω the sequence of random variables $n^{-1/2}(S_n^\omega - \mathbb{E}[S_n^\omega])$ converges in distribution to a centered normal random variable with variance $\sigma^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}(S_n^\omega)$.

³See the next section for references.

⁴See, for instance, the recent results [20, 33, 22, 21, 34, 25] and references therein.

⁵See [2] and references therein for results for iid maps when u_ω does not depend on ω and [45] for results for iid maps which admit a random tower extensions with sufficiently fast decaying tails.

of our knowledge, there are no examples in literature showing how to apply this method beyond the uniformly random case (where we can take $A = \Omega$). Some of the proofs in this paper will be based on applying the inducting strategy in [39] in the non-uniformly random case (see Section 1.4 for a more detailed discussion).

As will be explained in detail in Section 1.4, our conditions for the CLT and the functional law of iterated logarithm (LIL) will involve some mixing (weak-dependence) related conditions⁶ on sequences (f_n) of random variables of the form $f_n(\omega) = f(a_{1,\theta^n\omega}, \dots, a_{d,\theta^n\omega})$, where f has an explicit form, together with the integrability assumption $\|u_\omega - \mu_\omega(u_\omega)\|_\alpha \in L^p(\Omega, \mathcal{F}, \mathbb{P})$, $p > 2$, where $\|\cdot\|_\alpha$ is the standard Hölder norm corresponding to some exponent α . For instance, when T_ω is a piecewise monotone map on the unit interval with full images on each monotonicity interval, whose Jacobian has sufficiently regular⁷ variation on each monotonicity interval⁸ and its minimal amount of expansion on the monotonicity intervals is denoted by $\gamma_\omega > 1$, then our general conditions yield that the CLT holds true when $\|u_\omega - \mu_\omega(u_\omega)\|_\alpha \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ and the sequence of random variables $(\gamma_{\theta^n\omega})_{n=0}^\infty$ on $(\Omega, \mathcal{F}, \mathbb{P})$ satisfies some weak type of upper mixing condition. Our condition for the other limit theorems are similar (some also require certain integrability conditions, see Section 1.4). We stress that in certain circumstances our mixing assumptions and the other conditions for the limit theorems are essentially independent. For instance, in the above examples the CLT will hold when the random variables γ_ω satisfies conditions similar to the ones imposed on a_ω in the end of Section 1.1.1.

1.2. On the types of Gibbs measures and the smooth case. The equivariant measures μ_ω considered in this manuscript correspond to a piecewise Hölder continuous random potential ϕ_ω and have the property that they are absolutely continuous with respect to a random conformal measure ν_ω , so that $(\mathcal{L}_\omega)^* \nu_\omega = \lambda_\omega \nu_\omega$, where $\mathcal{L}_\omega g(x) = \sum_{y: T_\omega y = x} e^{\phi_\omega(y)} g(y)$ is the transfer operator of T_ω and $\lambda_\omega > 0$ (namely μ_ω is the random Gibbs measure corresponding to ϕ_ω , see [42, 44]). We will call the case “smooth” if the domain of T_ω is a smooth manifold and $e^{-\phi_\omega}$ is the Jacobian of T_ω with respect to the volume measure on the domain (i.e. $\phi_\omega = -\ln J_{T_\omega}$). In this case we have $\lambda_\omega = 1$ and ν_ω is the volume measure m_ω , and so μ_ω is absolutely continuous with respect to m_ω .

Like in [44], for partially expanding maps we impose a certain restriction on the oscillation of the underlying potential ϕ_ω , and, in addition, we will impose a certain restriction on the Hölder constant of ϕ_ω along inverse branches of T_ω , which will be crucial for obtaining the effective RPF rates that will be discussed in the next section. Like in [44], because of the restriction on the oscillation, the results are less applicable in the smooth case, but includes applications for random measures μ_ω of maximal entropy (when $\phi_\omega = 0$) and in the, so-called, high temperature regime when $\phi_\omega = \frac{1}{\beta} \psi_\omega$ for a sufficiently large β and a given random potential ϕ_ω satisfying some regularity conditions.

For properly expanding maps it is unnecessary to directly impose restrictions on the oscillation of the potential ϕ_ω , and we will only impose restrictions on the Hölder constant of the compositions of ϕ_ω with the inverse branches of T_ω . In the smooth case discussed above, this conditions immediately allows applications to piecewise affine maps⁹ (where μ_ω is the Lebesgue measure), since then ϕ_ω is constant on each inverse branch. We will also show that the type of control needed over the local variation in this case is satisfied for fiberwise piecewise C^2 -perturbations of such piecewise affine maps (see Section 2.4), and so we provide several examples in the smooth case, as well.

1.3. On our approach: effective RPF rates. The first part of our approach is based on effective rates in the random version of the (normalized) Ruelle-Perron-Frobenius theorem (RPF), a notion which for the sake of convenience is presented here as a definition.

1. Definition (Effective random rates). Let ϕ_ω be a random potential whose supremum norm is bounded by some random variable b_ω and its “variation” (e.g. local Hölder constant) is bounded

⁶These conditions will always hold true under appropriate restrictions on some upper mixing coefficients related to $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$.

⁷See more details in the paragraph below.

⁸e.g. fiberwise sufficiently small C^2 -perturbations of piecewise linear maps, see Section 2.4.

⁹We will also require that on each injectivity domain has a full image.

by a random variable v_ω . Let μ_ω be the random equivariant (i.e. $(T_\omega)_*\mu_\omega = \mu_{\theta\omega}$) Gibbs measure corresponding to the potential. We say that the transfer operators¹⁰ L_ω of T_ω (with respect to μ_ω) have random effective rates when acting on a space of functions with ‘‘bounded variation’’ if there are random variables $0 < \rho(\omega) < 1$ and $B_\omega \geq 0$ which depend analytically only on the random parameters $a_{1,\omega}, \dots, a_{d,\omega}$ and on b_ω and v_ω so that \mathbb{P} -a.s. for every function g on the domain of T_ω with bounded variation (i.e. $\|g\|_{var} < \infty$) we have

$$(1.1) \quad \left\| L_\omega^n g - \int g d\mu_\omega \right\|_{var} \leq B_{\theta^n \omega} \rho_{\omega,n} \|g\|_{var}$$

where $L_\omega^n = L_{\theta^{n-1}\omega} \circ \dots \circ L_{\theta\omega} \circ L_\omega$ and $\rho_{\omega,n} = \prod_{j=0}^{n-1} \rho(\theta^j \omega)$.

In this paper $\|g\|_{var}$ will always be the Hölder norm corresponding to some exponent α .

2. Example. In the example in Section 1.1.1, the measures $\mu_\omega = \text{Lebesgue}$ are equivariant, the corresponding potential is constant on the monotonicity interval of T_ω and the operator L_ω is defined by

$$L_\omega g(x) = a_\omega g(a_\omega x) + (1 - a_\omega)g(a_\omega + (1 - a_\omega)x).$$

For this example, if $\|\cdot\|_{var}$ denotes the Hölder norm corresponding to some exponent $\alpha \in (0, 1]$ we obtain (1.1) with

$$B_\omega = 24e^{4a_\omega^{-\alpha}}(1 + a_\omega^{-\alpha})^2$$

and

$$\rho(\omega) = \frac{e^{\frac{1}{2}a_{\theta\omega}^{-\alpha}}(1 + a_{\theta\omega}^\alpha) - (1 - a_{\theta\omega}^\alpha)}{e^{\frac{1}{2}a_{\theta\omega}^{-\alpha}}(1 + a_{\theta\omega}^\alpha) + (1 - a_{\theta\omega}^\alpha)}.$$

Note that as $a_\omega \rightarrow 1$ the amount of contraction $\rho(\omega)$ converges to 1, which is expected since when $a_\omega = 1$ we have $T_\omega x = x$ and the maps T_ω are no longer expanding. Observe also that in this example B_ω is actually bounded. This will be the case also for more general classes of random piecewise linear maps (and some of their perturbations), see Section 2.2.3.

We refer to Theorem 47 for a more precise formulation of the effective RPF rates (1.1) obtained in this paper. We also refer to Theorem 49 for effective rates for appropriate complex perturbations of the operators L_ω , which will be crucial for obtaining some of our results (see Section 1.4.1). As noted before, for partially expanding maps in the sense of [44] we obtain effective (real and complex) rates for potentials ϕ_ω with a sufficiently regular oscillation (which has applications for measures of maximal entropy and in the high temperature regime). As we have already mentioned this condition limits the applications to the smooth case, but for properly expanding maps (in the sense of [42, 33]) we will only require that for each inverse branch $y_{i,\omega}$ of T_ω the Hölder constant of $\phi_\omega \circ y_{i,\omega}$ does not exceed $(\gamma_{\theta\omega}^\alpha - 1)$ where γ_ω is the minimal amount of local expansion of T_ω . This condition means that $\phi_\omega \circ y_{i,\omega}$ is close to being a constant when the map $T_{\theta\omega}$ has a small amount of expansion. As mentioned in the end of Section 1.1, the latter condition about the Hölder constants is satisfied for appropriate types of perturbations of piecewise linear or affine maps (see Section 2.4).

The proof of Theorem 47 is based on showing that the non-normalized transfer operator \mathcal{L}_ω of T_ω contracts (w.r.t. the real Hilbert metric) an appropriate family of random cones \mathcal{C}_ω which are defined by means of the parameters $a_{i,\omega}$ (for instance, for properly expanding maps \mathcal{C}_ω is defined only by means of γ_ω , where γ_ω is the minimal amount of local expansion, which in the circumstances of Section 1.1.1 satisfies $\gamma_\omega = a_\omega^{-1}$). This is the main difference here in comparison to many other applications of the contraction properties of real Hilbert metric for random operators (see [38, 40, 42, 33, 44] and references therein), where the cones are usually defined by means of a random variable which can be expressed as a series of known random variables (but with unclear integrability or other regularity properties). As mentioned above, in the setup of [42, 33] the price to pay for being able to use more explicit cones is an additional limitation on the variation of the potential ϕ_ω along inverse branches,

¹⁰ L_ω is the dual of the Koopman operator $g \rightarrow g \circ T_\omega$ with respect to the probability measure μ_ω .

while in the setup of [44] we will also require that the amount of expansion dominates the amount of contraction fiberwise and not only on the average.

We would like to think about $\rho(\omega)$ in (1.1) as the amount of contraction we have on the fiber ω . We refer the readers to Remark 33 for a discussion about situations where B_ω is actually bounded. For instance, for the aforementioned example of perturbations of piecewise affine maps (described in Section 2.4), B_ω is bounded if the Hölder constant of the logarithm of the Jacobian of T_ω (on each inverse branch) is bounded¹¹, while when this Hölder constant is not bounded, we have $B_\omega \leq C\gamma_\omega^\alpha e^{\gamma_\omega^\alpha}$ where γ_ω is the minimal amount of contraction. For more general expanding maps¹² we have $B_\omega \leq C\gamma_\omega^\alpha e^{4\sup|\phi_\omega|+4\gamma_\omega^\alpha} (\deg(T_\omega) \deg(T_{\theta^{-1}\omega}))^2$, where $\deg(T_\omega)$ is the maximal number of preimages that a point x can have under T_ω . Note that under certain types of mixing assumptions on $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ the precise form of $\rho(\omega)$ does not make much difference, and only the fact that it is a function of the parameters $a_i(\omega)$ plays a significant role (still, we refer to Sections 2.3.1 and 2.3.2 for the precise form), since we will work under assumptions guaranteeing that the sequences $(a_{i,\theta^n\omega})_{n=0}^\infty$ are sufficiently fast mixing.

1.3.1. A comparison with existing less explicit random RPF rates. Let us compare (1.1) with a few other random RPF rates in literature. For the maps considered in [42, 40] (see also references therein) and [44] we have (1.1) with a constant ρ but with a random variable B_ω which is defined by means of first hitting times to certain sets and a certain random variable Q_ω which can be expressed as a series of known random variables (see, for instance, [42, Lemma 3.18]). In fact, the proof of these results relies¹³ on obtaining rates of the form (1.1) with random $\rho(\omega)$ which depends on Q_ω (see, for instance, [42, Proposition 3.17]). Note that even though Q_ω has a closed “formula” it is unclear which type of regularity conditions (e.g. integrability) it satisfies. A similar phenomena happens also in the random RPF rates obtained in [4, 5, 6, 14] (note that the third includes results for piecewise monotone interval maps without a random covering assumption), namely one can take $\rho(\omega) = \rho$ to be a constant but with the price of making B_ω less explicit. In any case, since it is not clear which regularity properties B_ω has in the above setups, it is less likely that these rates will be effective for proving limit theorems under explicit conditions, and not conditions involving some restrictions on the random variable B_ω (which are hard to verify).

Another less direct approach is based on an appropriate version of Oseledets theorem for the cocycle of transfer operators $\{L_\omega : \omega \in \Omega\}$, and under certain logarithmic integrability conditions (see [23, Proposition 26] and references therein), it yields that

$$(1.2) \quad \left\| L_\omega^n g - \int g d\mu_\omega \right\|_{var} \leq K(\omega) e^{-n\lambda} \|g\|_{var}$$

for some $\lambda > 0$ and a tempered¹⁴ random variable $K(\omega)$. Notice that once (1.2) is established with some λ then the minimal choice for $K(\omega)$ is

$$K(\omega) = \sup_n \|L_\omega^n - \mu_\omega\|_{var} e^{n\lambda}.$$

Remark also that

$$\lambda \leq \lambda_0(\omega) := -\limsup_{n \rightarrow \infty} \frac{1}{n} \ln \|L_\omega^n - \mu_\omega\|_{var}, \quad \mathbb{P}\text{-a.s.}$$

and so, in a sense, $\lambda = \lambda^{(0)} := \text{ess-inf } \lambda_0(\omega)$ is the smallest possible choice for λ . We note that when $\ln U$ is integrable then (1.1) yields that $\lambda^{(0)} \leq \bar{\lambda} = -\int \ln \rho(\omega) d\mathbb{P}(\omega)$ which is a limitation on the

¹¹We refer to Remarks 9 and 12 for a discussion about this matter and its relation to artificiality limiting the minimal amount of contraction by forcing it to be bounded above. This can always be done, but then stronger conditions on the potential are needed, which essentially reduce to the boundedness of the variation of the potential along inverse branches, which in the smooth case is the negative logarithm of the Jacobian.

¹²More precisely, for the maps described in Remark 4

¹³When (1.1) holds true with any kind of random variables B and ρ , we can replace $\rho(\omega)$ by a constant smaller than 1 by considering the number of visits to a set of the form $A_\varepsilon = \{\rho(\omega) < 1 - \varepsilon\}$ for $\varepsilon \in (0, 1)$ small enough. However, this will make the “new” B_ω less explicit and with unclear regularity properties.

¹⁴Namely, almost surely we have $\lim_{n \rightarrow \infty} \frac{1}{n} \ln K(\theta^n \omega) = 0$.

contraction rate in the exponential convergence towards μ_ω . Even though we have the above explicit form for $K(\omega)$, it is unclear which type of regularity (beyond being tempered) the random variable $K(\omega)$ possesses¹⁵ or if it has a finite upper bound which depends (in a reasonable way) only the parameters $a_i(\omega)$ describing the maps T_ω . Under the conditions of [23, Proposition 26] in [23, 24] limit theorems were obtained in the smooth case for expanding on the average maps T_ω and random potentials u_ω satisfying (roughly speaking) that $K(\omega)\|u_\omega\|_{var} \leq C$ for¹⁶ some constant C . In comparison with the smooth case considered in [23, 24], we restrict ourselves to maps which have some fiberwise expansion (maybe not on the entire space) and not only expansion on the average. Moreover, we will have an additional assumption on the Jacobian (which will be satisfied for certain perturbations of piecewise affine maps) and certain type of upper mixing conditions on the system $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ as well. On the other hand, as noted above, in general $K(\omega)$ does not seem to be “computable”, and we also consider more general families of equivariant measures μ_ω corresponding to potentials with sufficiently regular variation (e.g. the maximal entropy and the high-temperature regime cases discussed above).

Let us also mention related results for (partially hyperbolic) iid maps $\{T_{\theta^j\omega} : j \geq 0\}$ which admit a random (Young) tower extension (see [7, 29, 8, 3]). In this setup estimates of the form

$$(1.3) \quad \left\| L_\omega^n g - \int g d\mu_\omega \right\|_{L^1(\mu_\omega)} \leq K(\omega) a_n \|g\|_{var}$$

were obtained for some sequences $a_n \rightarrow 0$ (the decay rate of a_n is determined by the decay rates of the tails of the random tower) and a random variable $K(\omega)$ which satisfies certain regularity conditions like $K(\omega) \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ for some $p > 1$. While also here $K(\omega)$ does not seem to depend only on some parameters describing the original maps (or something similar), integrability conditions on $K(\omega)$ together with polynomial decay of a_n are sufficient to get appropriate control over the non-uniform decay of correlations, which is enough to prove limit theorems like an almost sure invariance principle (see [45]). However, this is obtained only for iid maps which admit a sufficiently regular random tower extension (and iid functions $\{u_{\theta^j\omega} : 0 \leq k < \infty\}$). Moreover, even for iid maps several other limits theorems like the ones described in Section 1.4.1 seem to require more than (1.3).

1.4. A more detailed discussion on the proofs and conditions of the limit theorems. A major difficulty in proving limit theorems in the non-uniformly random case (beyond the iid case) is that the iterates of the annealed transfer operator (see [2]) do not describe the statistical behavior of the random Birkhoff sums, and due to strong dependence between T_ω, u_ω and $T_{\theta\omega}, u_{\theta\omega}$ it seems less likely that a random tower extension with sufficiently fast decaying tails exists (see again [7, 29, 8, 3] and [45]). Instead, our results will rely on the effective rates (1.1) described in the previous sections, as described in the following paragraphs.

We present two proofs of the central limit theorem (CLT) and the functional law of iterated logarithm (LIL). The first one (i.e. the proof of Theorem 27) is based on inducing, and more precisely we use the inducing strategy in [39, Theorem 2.3]. To the best of our knowledge, this is the first time that a result based on inducing in the Ω direction is applied effectively for expanding maps like the ones considered in this paper (namely, that the required control over the system between two visiting times to the set on the base Ω is achieved). The idea in our proof is that, using (1.1), the conditions of [39, Theorem 2.3] reduce to certain almost sure growth conditions which involve the random variables $\rho(\omega), U_\omega$ and $c_\omega = \|u_\omega - \mu_\omega(u_\omega)\|_{var} = \|u_\omega - \mu_\omega(u_\omega)\|_\alpha$, which in turn can be verified under certain types of mild upper weak-dependence (mixing) assumptions on the sequences¹⁷ $(\rho(\theta^n\omega))_{n=0}^\infty$ and $(B_{\theta^n\omega})_{n=0}^\infty$ and the integrability condition $c_\omega \in L^p(\Omega, \mathcal{F}, \mathbb{P}), p > 2$. We note that when $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ is the shift system generated by a sufficiently fast mixing sequence (e.g. a geometrically ergodic Markov chain or some other exponentially fast mixing sequence) these mixing conditions will always hold true when we can

¹⁵e.g., whether it is in $L^p(\Omega, \mathcal{F}, \mathbb{P})$ for some p or if $(K(\theta^n\omega))$ satisfies some weak-dependence conditions.

¹⁶Let us also note that in [24, Appendix A] it is demonstrated that, in general, scaling conditions of this form are necessary for the validity of certain limit theorems.

¹⁷Recall that $\rho(\omega)$ and B_ω are functions of $a_{i,\omega}$ and so it is enough to impose upper weak-dependence conditions on $(a_{i,\theta^n\omega})_{n=0}^\infty$ for $i = 1, 2, \dots, d$.

approximate $\rho(\omega)$ and B_ω sufficiently fast by functions of $(X_j)_{|j|\leq r}$ as $r \rightarrow \infty$ (in particular, when $\rho(\omega)$ and B_ω depend only on finitely many of the X_j 's). We stress that integrability conditions on B_ω are not required and all that is needed is some type of upper mixing conditions and integrability assumptions on c_ω . In order to illustrate the above scheme, for the example in Section 1.1.1 it will be enough to induce on a set of the form $A = \{\omega : \max(a_\omega, a_{\theta\omega}) \leq 1 - \delta\}$ for a sufficiently small δ such that $\mathbb{P}(A) > 0$ (note that some sufficient mixing conditions were already discussed in Section 1.1.1).

Our second proof of the CLT and LIL (namely, the proof of Theorem 32) is not based on inducing, and instead it exploits (1.1) directly and also requires that $B_\omega \in L^p$ (as noted above, B_ω is even bounded for a wide class of maps, see Remark 33). While in general integrability assumptions on B_ω are true additional requirements, the second type of sufficient conditions for the CLT has two advantages over the first set. First, it requires much weaker restrictions on certain upper mixing coefficients related to the system $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$. Second, it allows weaker approximation rates of $(\rho(\theta^n \omega))_{n=0}^\infty$ and $(B_{\theta^n \omega})_{n=0}^\infty$ than the ones required in the first set of conditions (in the case when the latter sequences can only be approximated by sequences satisfying some type of upper weak-dependence conditions).

We also obtain an almost sure invariance principle (ASIP), see Theorem 34, which concerns strong approximation of the random Birkhoff sums by sums of independent Gaussian random variables (and is a stronger form of the CLT). Under some upper weak-dependence assumptions on $(\rho(\theta^n \omega))_{n=0}^\infty$ and $(U(\theta^n \omega))_{n=0}^\infty$ and some integrability conditions we obtain an ASIP with rates $o(n^{1/4+7p/2+\varepsilon})$, where p is the largest number so that the random variable $Y(\omega)$ described in the last paragraph of Section 1.1 belongs to L^p . For instance (see Remark 36), under certain regularity assumptions on the potential ϕ_ω our integrability conditions are $\|u_\omega - \mu_\omega(u_\omega)\|_\alpha \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ and $N(\omega) \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ where $N(\omega) = \sup\{v_\alpha(g \circ T_\omega) : v_\alpha(g) \leq 1\}$. In the smooth case these conditions hold true for the aforementioned C^2 -perturbations of the piecewise affine maps (where here $N(\omega)$ essentially coincides with the maximal amount of expansion of the map). In more general circumstances we also require that $B_\omega \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ (in the situations discussed before it is bounded).

For non-uniformly random iid maps which admit a random tower extension an ASIP was obtained¹⁸ in [45], while for non uniformly random expanding maps driven by a general ergodic system $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ it was obtained in [26] by inducing on an appropriate set A . The conditions in [26] reduce to certain assumptions on the behavior of the random Birkhoff sums $S_n^\omega u$ when n is smaller than the first visiting time $n_A(\omega)$ of the orbit of ω to A . The proof of the ASIP in this paper is not based on inducing, and instead we apply (1.1) directly, but we still think it could be interesting to check how (1.1) can be combined with an inducing strategy in order to yield some ASIP rates. Finally, we would also like to refer to [24], where an ASIP was obtained under the scaling conditions described in the penultimate paragraph of Section 1.3.1.

1.4.1. *Results which also require random complex effective rates, and the deterministic case.*

In Theorems 39 and 40 we also derive a moderate deviations principle (MDP) which deals with the asymptotic behavior of probabilities of the form $\mu_\omega\{(S_n^\omega u - \mu_\omega(S_n^\omega u))/a_n \in \Gamma\}$ where (a_n) is a certain type of normalizing sequence and $\Gamma \subset \mathbb{R}$ is an arbitrary Borel set (we refer to these results as moderate deviations¹⁹ because of the quadratic rate function involved in the formulation). These results are obtained under an additional condition on the potential u_ω which, roughly speaking, means that either the Hölder norm $\|u_\omega\|_\alpha$ is small when T_ω has some inverse branch with a small amount of contraction, or that it is small when the ratio between the amount of expansion and contraction is close to 1. Such a condition is close in spirit to the scaling conditions in [23, 24] discussed in the penultimate paragraph of Section 1.3.1, but the scaling is done according to the amount of expansion of T_ω .

Under the same additional requirement on the random functions u_ω , we will also obtain self-normalized CLT rates and a moderate version of the local CLT (see Theorems 43 and 45, respectively).

¹⁸The ASIP for uniformly random maps was treated in several papers in different setups, see [20] and the references in [45, 26].

¹⁹As opposed to large deviations.

Our CLT rates are of order $n^{-(1/2-6/p)}$ when appropriate random variables (like the ones discussed in previous paragraphs) belong to $L^p(\Omega, \mathcal{F}, \mathbb{P})$. When these random variables are bounded (i.e. in the uniformly random case) we have $p = \infty$ and this result recovers the Berry-Esseen theorem [33, Theorem 7.1.1] (where the optimal $n^{-1/2}$ rates were obtained, see also [25, 34]). In [33, Theorem 7.15], in the uniformly random case, a local CLT was derived for the type of expanding maps considered in this paper (see also [21, 22, 25, 34]), but the moderate type of local CLT considered here is in a difference scale, it holds true without any additional aperiodicity conditions as in [33, 21, 22, 25, 34] and it is new even in the uniformly random case. On the other hand, it provides local CLT estimates on a weaker scale.

The proofs of the MDP, the CLT rates and the moderate local CLT require effective rates for appropriate complex perturbations of the transfer operators L_ω , which is established Theorem 49. In fact, for partially expanding maps (as in Section 2.2), Theorem 49 is new even in the uniformly random case (in that case B_ω and $\rho(\omega)$ are constants in the appropriate complex version). The proof of Theorem 49 uses Rugh's theory [43] of the contraction properties of complex Hilbert metrics associated with complex cones (see also [27, 28]). For uniformly random properly expanding maps T_ω this method was applied successfully for random complex transfer operators for the first time in [33, Ch. 4-6], and here we show how to apply it when the amount of contraction at the "jump" from ω to $\theta\omega$ depends on ω (roughly speaking, the amount of contraction is $\rho(\omega)$ appearing on (1.1)), as well as for partially expanding maps (for such maps our results are new even in the uniformly random case).

Finally, let us note that for the partially expanding maps considered in this paper, the application of [43] is new even in the deterministic case (i.e. in the setup of [16]). This results in explicit bounds on the spectral gap of appropriate complex perturbations on the dual operator of the Koopman operator corresponding to the deterministic map T and the underlying Gibbs measure. Using such estimates²⁰ we can obtain, for instance, explicit constants in the Berry-Esseen theorem. That is, the methods used in this paper also make it possible to extend [28, Theorem 1.1] from the properly expanding case to the partially expanding case, and we expect other similar quantitative results to follow.

2. RANDOM EXPANDING MAPS

As mentioned in Section 1, similarly to [44] we will consider partially random expanding maps and random Gibbs measures corresponding to random potentials with sufficiently small random oscillation and a small Hölder constant along inverse branches. However, when the maps are properly expanding (i.e. all local inverse branches are strongly contracting) we only need the condition about the Hölder constants, which will allow applications in the smooth case. For that reason we begin the presentation in the setup of [33, Ch. 6] (which is similar to [42]) and only after that we will present the setup of partially expanding maps.

2.1. Properly expanding maps with a local pairing property.

2.1.1. *Random spaces and maps.* Our setup consists of a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ together with an invertible ergodic \mathbb{P} -preserving transformation $\theta : \Omega \rightarrow \Omega$, of a compact metric space (\mathcal{X}, ρ) normalized in size so that $\text{diam}\mathcal{X} \leq 1$ together with the Borel σ -algebra \mathcal{B} , and of a set $\mathcal{E} \subset \Omega \times \mathcal{X}$ measurable with respect to the product σ -algebra $\mathcal{F} \times \mathcal{B}$ such that the fibers $\mathcal{E}_\omega = \{x \in \mathcal{X} : (\omega, x) \in \mathcal{E}\}$, $\omega \in \Omega$ are compact. The latter yields (see [15] Chapter III) that the mapping $\omega \rightarrow \mathcal{E}_\omega$ is measurable with respect to the Borel σ -algebra induced by the Hausdorff topology on the space $\mathcal{K}(\mathcal{X})$ of compact subspaces of \mathcal{X} and the distance function $\rho(x, \mathcal{E}_\omega)$ is measurable in ω for each $x \in \mathcal{X}$. Furthermore, the projection map $\pi_\Omega(\omega, x) = \omega$ on \mathcal{E} is measurable and it maps any $\mathcal{F} \times \mathcal{B}$ -measurable set to a \mathcal{F} -measurable set (see "measurable projection" Theorem III.23 in [15]).

3. Remark. Compactness of either \mathcal{X} or \mathcal{E}_ω will only be needed to insure the measurability of $\omega \rightarrow \mathcal{E}_\omega$ in the above sense. Thus, when $\mathcal{E}_\omega = \mathcal{X}$ for every ω then our results will remain valid for bounded metric spaces \mathcal{X} which are not necessarily compact.

²⁰The idea is that, contrary to the classical perturbative approach based on an appropriate implicit function theorem, we can control the size of perturbation as well as obtain explicit bounds on the corresponding complex RPF triplets.

Next, let

$$\{T_\omega : \mathcal{E}_\omega \rightarrow \mathcal{E}_{\theta\omega}, \omega \in \Omega\}$$

be a collection of maps between the metric spaces \mathcal{E}_ω and $\mathcal{E}_{\theta\omega}$ so that the map $(\omega, x) \rightarrow T_\omega x$ on \mathcal{E} is measurable with respect to the restriction of $\mathcal{F} \times \mathcal{B}$ to \mathcal{E} . For every $\omega \in \Omega$ and $n \in \mathbb{N}$ consider the n -th step iterates T_ω^n given by

$$(2.1) \quad T_\omega^n = T_{\theta^{n-1}\omega} \circ \cdots \circ T_{\theta\omega} \circ T_\omega : \mathcal{E}_\omega \rightarrow \mathcal{E}_{\theta^n\omega}.$$

Our first additional requirement from the maps T_ω is that there is a random variable $\gamma_\omega > 1$ so that for every $x, x' \in \mathcal{E}_{\theta\omega}$ we can write

$$(2.2) \quad T_\omega^{-1}\{x\} = \{y_i = y_{i,\omega}(x) : i < k\} \quad \text{and} \quad T_\omega^{-1}\{x'\} = \{y'_i = y_{i,\omega}(x') : i < k\}$$

and

$$(2.3) \quad \rho(y_i, y'_i) \leq (\gamma_\omega)^{-1} \rho(x, x')$$

for all $1 \leq i < k$ (where either $k \in \mathbb{N}$ or $k = \infty$).

4. Remark. We can also consider a somehow different setup where (2.2) and (2.3) hold true only for two points x, x' such that $\rho(x, x') \leq \xi$ for some fixed constant $\xi < 1$. In this case we would have to assume that $\deg(T_\omega) < \infty$ (so k above is finite) and that $\deg(T_\omega)$ is measurable²¹. Here $\deg(T_\omega) = \max\{|T_\omega^{-1}\{x\}| : x \in \mathcal{E}_{\theta\omega}\}$, where $|A|$ denotes the cardinality of a finite set A . Moreover, we will need to require the following two additional covering assumptions:

(i) there exists $n_0 \in \mathbb{N}$ and an ξ -cover of \mathcal{E}_ω by points $x_i = x_{i,\omega}$ such that for all i we have $T_\omega^{n_0}(B_\omega(x_i, \xi)) = \mathcal{E}_{\theta\omega}$.

(ii) for all $x, x' \in \mathcal{E}_{\theta\omega}$, for every $y \in T_\omega^{-1}\{x\}$ there exists $y' \in T_\omega^{-1}\{x'\}$ so that $\rho(y, y') \leq \xi$.

The main example we have in mind for such types of maps are certain classes of random sub-shifts of finite type (see, for instance [40] but also [42, Section 2.1]). In this case we take ξ small enough so that $\rho(x, x') \leq \xi$ means that the 0-th coordinate of x and x' coincide. Note that for the maps presented before Remark 4 we can just take $\xi = 1$, and in this case there is no need for the additional requirements (i) and (ii). In order to avoid a more complicated presentation of our main results we decided to focus on the case $\xi = 1$, which already includes most of the (non-symbolic) examples we have in mind (e.g. the one in Section 1.1.1).

Next, for every $\omega \in \Omega$ and all $g : \mathcal{E}_\omega \rightarrow \mathbb{C}$ set

$$v(g) = v_{\alpha,\xi,\omega}(g) = \inf\{R : |g(x) - g_\omega(x')| \leq R\rho^\alpha(x, x') \text{ if } \rho(x, x') < \xi\}$$

$$\text{and } \|g\| = \|g\|_{\alpha,\xi} = \|g\|_\infty + v_{\alpha,\xi}(g)$$

where $\|\cdot\|_\infty$ is the supremum norm and $\rho^\alpha(x, x') = (\rho(x, x'))^\alpha$ (and α is the same as in (2.4)).

5. Remark. If $g : \mathcal{E} \rightarrow \mathbb{C}$ is measurable and $g_\omega : \mathcal{E}_\omega \rightarrow \mathbb{C}$ is given by $g_\omega(x) = g(\omega, x)$ then the function $\omega \rightarrow \|g_\omega\|$ is measurable by [33, Lemma 5.1.3].

Next, consider the Banach spaces $(\mathcal{H}_\omega, \|\cdot\|) = (\mathcal{H}_\omega^{\alpha,\xi}, \|\cdot\|_{\alpha,\xi})$ of all functions $h : \mathcal{E}_\omega \rightarrow \mathbb{R}$ such that $\|h\|_{\alpha,\xi} < \infty$ and denote by $\mathcal{H}_{\omega,\mathbb{C}} = \mathcal{H}_{\omega,\mathbb{C}}^{\alpha,\xi}$ the space of all complex-valued functions with $\|h\|_{\alpha,\xi} < \infty$.

2.1.2. The random potential. Let $\phi : \mathcal{E} \rightarrow \mathbb{R}$ be a measurable function so that $\|\phi(\omega, \cdot)\|_\infty < \infty$. Fix some $\alpha \in (0, 1]$, and suppose that for \mathbb{P} -a.e. ω and let H_ω be a random variable such that for all x and x' in $\mathcal{E}_{\theta\omega}$ for all i we have

$$(2.4) \quad |\phi_\omega(y_{i,\omega}(x)) - \phi_\omega(y_{i,\omega}(x'))| \leq H_\omega \rho^\alpha(x, x')$$

where $\phi_\omega(x) = \phi(\omega, x)$. In this paper we assume that

$$(2.5) \quad H_\omega \leq \gamma_{\theta\omega}^\alpha - 1.$$

²¹we can assume that $\deg(T_\omega) \leq d_\omega$ for some random variable d_ω instead.

This condition means that the Hölder constant of each composition $\phi_\omega \circ y_{i,\omega}$ is small when the “next map” $T_{\theta\omega}$ has a small amount of contraction on some piece of the space.

6. Remark. Condition (2.4) holds true when $v(\phi_\omega) \leq \gamma_\omega^\alpha (\gamma_{\theta\omega}^\alpha - 1)$. However, this condition is more general since it only imposes restrictions on the Hölder constant along the inverse branches of T_ω , and, as will be demonstrated in Section 2.4, it allows applications in the smooth case. The idea is that for piecewise affine maps (2.4) holds true with $H_\omega = 0$, and so (2.4) and (2.5) will hold true for appropriate perturbations of such maps (see Section 2.4). Condition (2.5) is also in force when ϕ_ω has the form $\frac{\psi_\omega}{\gamma_\omega^\alpha (\gamma_{\theta\omega}^\alpha - 1)}$ where ψ_ω has Hölder constant (corresponding to the exponent α) smaller than 1 (see also Remark 5).

7. Remark. In principle, we can define

$$(2.6) \quad H_\omega = \sup_i v(\phi_\omega \circ y_{i,\omega})$$

and assume that $H_\omega \leq \gamma_{\theta\omega}^\alpha - 1$. However, the function $\omega \rightarrow H_\omega$ might not be measurable because we did not assume that $\omega \rightarrow \deg(T_\omega)$ is measurable.

Next, we need following summability condition:

8. Assumption. There is a random variable $D_\omega < \infty$ such that

$$\sup_{x \in \mathcal{E}_{\theta\omega}} \sum_{y \in T_\omega^{-1}\{x\}} e^{\phi_\omega(y)} \leq D_\omega.$$

Note that when $\deg(T_\omega)$ is finite and measurable then this assumption trivially holds true $D_\omega = \deg(T_\omega) e^{\|\phi_\omega\|_\infty}$ ($\|\phi_\omega\|_\infty = \sup |\phi_\omega|$ is measurable by [33, Lemma 5.1.3]).

9. Remark. The non-uniform expansion comes from the possibility that γ_ω will be arbitrary close to 1 (when γ_ω is large then T_ω is strongly expanding). Notice that conditions (2.2) and (2.3) remain valid if we replace γ_ω with $\gamma_{\omega,M} = \min(M, \gamma_\omega)$ for some constant $M > 1$. While this limits the amount of expansion, some (but not all) of the conditions of the limit theorems that will be proven in this paper require intergability assumptions which are weaker when γ_ω is bounded. On the other hand, forcing γ_ω to be bounded by replacing it with $\gamma_{\omega,M}$ essentially means that instead of (2.5) we require that $H_\omega \leq (\gamma_{\theta\omega,M}^\alpha - 1)$ for some $M > 0$, and so there is a trade-off between the aforementioned integrability conditions and the latter stronger version of (2.5).

2.1.3. An example and a comparison with [42]. One example of maps which satisfy our conditions are piecewise injective maps. In this case let $\mathcal{E}_\omega = \mathcal{X} = [0, 1]^d$ for some $d \in \mathbb{N}$, and take a random partition of \mathcal{X} into rectangles of the form $[a_1, b_1) \times [a_2, b_2) \times \cdots \times [a_d, b_d)$. Now, on each rectangle we can take a distance expanding map which maps it onto $[0, 1]^d$. Note that since \mathcal{E}_ω does not depend on ω there is no need in compactness to insure its measurability.

The condition (2.5) is a restriction on the potential ϕ_ω . While we can always choose a potential which satisfies this condition, it is interesting to see when this setup applies to the smooth case when $\phi_\omega = -\ln(J_{T_\omega})$ and μ_ω is the unique absolutely continuous invariant measure w.r.t. the volume measure, and we refer to Section 2.4 for examples in the smooth case (fiberwise piecewise C^2 perturbations of certain piecewise linear or affine maps).

2.2. Random maps with dominating expansion.

2.2.1. Random spaces and maps. Let $(\Omega, \mathcal{F}, \mathbb{P}, \theta), (\mathcal{X}, \rho), \{\mathcal{E}_\omega\}$ and $\{T_\omega : \mathcal{E}_\omega \rightarrow \mathcal{E}_{\theta\omega}\}$ satisfy the same properties described in the first paragraph of Section 2.1. In this section, our additional assumptions on the maps T_ω are as follows: we suppose that there exist random variables $l_\omega \geq 1$, $\sigma_\omega > 1$, $q_\omega \in \mathbb{N}$ and $d_\omega \in \mathbb{N}$ so that $q_\omega < d_\omega$ and for every $x \in \mathcal{E}_{\theta\omega}$ we can write

$$(2.7) \quad T_\omega^{-1}\{x\} = \{y_{1,\omega}(x), \dots, y_{d_\omega,\omega}(x)\}$$

where for every $x, x' \in \mathcal{E}_{\theta\omega}$ and for $i = 1, 2, \dots, q_\omega$ we have

$$(2.8) \quad \rho(y_{i,\omega}(x), y_{i,\omega}(x')) \leq l_\omega \rho(x, x')$$

while for $i = q_\omega + 1, \dots, d_\omega$,

$$(2.9) \quad \rho(x_i, x'_i) \leq \sigma_\omega^{-1} \rho(x, x').$$

The above conditions are satisfied in the setup of [44] (see Section 2.2.3 for a discussion). We assume here that

$$(2.10) \quad a_\omega := \frac{q_\omega l_\omega^\alpha + (d_\omega - q_\omega) \sigma_\omega^{-\alpha}}{d_\omega} < 1$$

which is a quantitative estimate on the amount of allowed contraction, given the amount of expansion T_ω has.

Next, denote by \mathcal{H}_ω the space of functions on \mathcal{E}_ω equipped with the norm

$$\|g\| = \|g\|_\infty + v(g)$$

where $\|g\|_\infty = \sup |g|$ and $v(g) = v_\alpha(g)$ is the smallest number so that $|g(x) - g(y)| \leq v(g)(\rho(x, y))^\alpha$ for all x and y in \mathcal{E}_ω (namely $\mathcal{H}_\omega = \mathcal{H}_\omega^{\alpha, 1}$ in the notations of the previous section). In the case when $\alpha = 1$ and each \mathcal{E}_ω is a Riemannian manifold we will also consider the norms $\|g\| = \|g\|_{C^1} = \sup |g| + \sup \|Dg\|$ on the space of C^1 -functions, namely $v(g)$ above is replaced by the supremum norm of the differential of g (so in this case $v(g)$ could either be the Lipschitz constant or $\sup \|Dg\|$).

2.2.2. The random potential. Next, let $\phi : \mathcal{E} \rightarrow \mathbb{E}$ be a measurable function and let $\phi_\omega : \mathcal{E}_\omega \rightarrow \mathbb{R}$ be given by $\phi_\omega(x) = \phi(\omega, x)$. Set

$$\varepsilon_\omega = \text{osc}(\phi_\omega) = \sup \phi_\omega - \inf \phi_\omega$$

be the oscillation of ϕ_ω and for some fixed $\alpha \in [0, 1)$ let

$$(2.11) \quad H_\omega = \max\{v(\phi_\omega \circ y_{i, \omega}) : 1 \leq i \leq d_\omega\}$$

be the maximal Hölder constant along inverse branches. We assume here that both ε_ω and H_ω are finite. Note that if ϕ_ω was Hölder continuous on the entire space \mathcal{E}_ω then $H_\omega \leq l_\omega v(\phi_\omega)$. Our additional requirements from the function ϕ_ω is that

$$(2.12) \quad s_\omega := e^{\varepsilon_\omega} a_\omega < 1 \quad \text{and} \quad e^{\varepsilon_\omega} H_\omega \leq \frac{s_\omega^{-1} - 1}{1 + s_\omega^{-1}}$$

10. Remark. The assumption about H_ω is a version of the combination of conditions (2.4) and (2.5). Let δ_ω be so that $(1 + \delta_\omega) a_\omega < 1$ and suppose that $e^{\varepsilon_\omega} (1 + \delta_\omega) a_\omega < 1$. Then the condition about H_ω is satisfied when $H_\omega \leq \frac{\delta_{\theta_\omega} a_\omega^2}{1 + a_\omega}$. Note that we can always assume that a_ω is bounded below by some positive constant (by replacing a_ω with $\tilde{a}_\omega = \max(a_\omega, 1 - \varepsilon)$ if needed). This will make no difference in our proofs, and in that case the second condition reads $H_\omega \leq C \delta_{\theta_\omega}$ for some C which can be arbitrarily close to 1.

2.2.3. A comparison with [44] and (additional) examples.

On the assumptions. Our assumptions (2.7), (2.8) and (2.9) on the maps correspond to [44, Assumption (H1)] (see also the proof of [44, Proposition 5.4]). Our condition $s_\omega < 1$ is a stronger version of [44, (2.2)] in [44, Assumption (H3)], which requires that $\int \ln s_\omega d\mathbb{P}(\omega) < 0$ instead. In addition to this difference we also have the additional assumption about H_ω , which is an additional fiberwise restriction on the local Hölder constant of the potential on inverse branches. This condition always holds true when ϕ_ω is constant on each inverse branch. On the other hand, in [44] there are several other assumptions on the maps T_ω like [44, Assumptions (H4) and (H5)] or [44, Assumptions (H4) and (H5')] whose purpose is to prove uniqueness of the RPF triplets described in [44, Theorem A] (see also Theorem 47). While it is natural to work under assumptions that guarantee uniqueness of RPF triplets (and equilibrium states), our result will not require such assumptions and all the limit theorems will hold for a certain type of random equilibrium state (Gibbs measure), which coincides with the unique one under the additional assumptions in [44].

Special choices of random potential. Let us discuss two special types of potentials considered in [44, Theorem D]. First, let us consider the case when $\phi_\omega \equiv 0$. This case corresponds to equivariant measure μ_ω of maximal entropy (see [44, Theorem D]), and in our case (2.12) holds true for that choice as long as $a_\omega < 1$. Again, the main difference in this case in comparison with [44] is that the weaker assumption $\int \ln a_\omega d\mathbb{P}(\omega) < 0$ was assumed instead. Another special choice for ϕ_ω is the case when $\phi_\omega = \psi_\omega/T$ for some other random potential and a sufficiently large constant T (this is usually referred to as the high-temperature regime). In the high-temperature regime our results for general potentials are mostly effective in the uniformly random case²², but we note that in the setup of this section most of the results will be new even in then.

Some additional examples of maps. In [44, Section 3] several examples were given, and in our setup we can consider the same examples replacing the assumptions about the integral of $\ln a_\omega$ by almost sure assumptions on a_ω . For instance, let us consider a random finite partition of $[0, 1)$ into intervals $I_{\omega,i} = [a_{\omega,i}, b_{\omega,i})$, $i \leq d_\omega$. On each i let us take a monotone Hölder continuous map $T_{\omega,i} : I_{\omega,i} \rightarrow [0, 1)$ which IS onto $[0, 1)$. Let us assume that the absolute value of the derivatives of $T_{\omega,1}, \dots, T_{\omega,p_\omega}$ is not less than $\sigma_\omega > 1$, while the derivatives of the other $q_\omega = d_\omega - p_\omega$ maps $T_{\omega,i}$ does not exceed l_ω^{-1} for some $l_\omega \geq 1$. Then all the conditions described before are valid if $a_\omega < 1$. A particular case are the, so called, random Manneville–Pomeau maps. Let $\beta(\omega) \in (0, 1)$ be a random variable and let us take $I_{\omega,1} = [0, \frac{1}{2})$ and $I_{\omega,2} = [\frac{1}{2}, 1)$. On the first interval, let $T_{\omega,1}(x) = x(1 + (2x)^{\beta(\omega)})$ while on the second we set $T_{\omega,2}(x) = 2x - 1$. Then $q_\omega = p_\omega = 1$, $\sigma_\omega = 2$ and $l_\omega = 1$. In this case

$$s_\omega = e^{\varepsilon_\omega} \frac{1 + 2^{-\alpha}}{2}, \quad H_\omega = \max(v_\alpha(\phi_\omega \circ T_{\omega,1}^{-1}), v_\alpha(\phi_\omega \circ T_{\omega,2}^{-1})).$$

Similar multidimensional examples can be given, for instance $\mathcal{I}_{\omega,i}$ can be a partition of $[0, 1)^d = [0, 1) \times \dots \times [0, 1)$ into rectangles with disjoint interiors of the form $I_{\omega,i} = [a_{1,\omega}^{(i)}, b_{1,\omega}^{(i)}) \times \dots \times [a_{d,\omega}^{(i)}, b_{d,\omega}^{(i)})$ and on each rectangle we can take an injective map whose image is $[0, 1)^d$, and assume that some of the maps are distance expanding, while others might contract distance on some regions of the rectangle (e.g. we can start with affine maps and perturb). We would also like to refer to [44, Section 3.4] for other multidimensional maps, which are included in our setup when the condition

$$\sup_k \left(\left(1 - \frac{\ell_k}{k} \right) + \frac{\ell_k}{k} L_k \right) < 1$$

mentioned after [44, (3.2)] is satisfied (and $\sigma_k > 1$ for all k). Note that there are additional requirements in [44, Section 3.4] but as mentioned above their purpose is to insure the uniqueness of the RPF triplets, which is not a requirement in this paper.

2.3. Frequently used random variables. In this section we will introduce several random variables and the random cones that will be involved in the formulation of the effective Perron-Frobenius rates (Theorems 47 and 49), as well as in the conditions of the limit theorems.

11. Remark. From now on variables x which involves θ_ω directly will be written in the form $x(\omega)$, while we will use the notation x_ω for a variable x whose definition does not directly involve θ_ω .

2.3.1. Properly expanding maps. In the circumstances of Section 2.1, let \mathcal{C}_ω be the real cone defined by

$$\mathcal{C}_\omega = \{g \in \mathcal{H}_\omega : g \geq 0, g(x) \leq e^{\gamma_\omega \rho^\alpha(x, x')} g(x'), \forall x, x' \in \mathcal{E}_\omega\}.$$

The following random variables are used in the formulation of Theorems 27, 32, 34 (CLT and ASIP) and Theorems 47 and 49 (Perron-Frobenius rates).

²²In the non-uniformly case we can consider ψ_ω which satisfies (2.12) with ε_ω/T instead of T and take $\phi_\omega = \psi_\omega/T$.

Random variable	Role/comments
$B_\omega = 24e^{4\gamma_\omega^\alpha}(1 + \gamma_\omega^\alpha)^2$	appears on the RHS of (1.1)
$q(\omega) = \frac{H_\omega + 1}{\gamma_{\theta_\omega}^\alpha}$	$q(\omega) \in (0, 1)$ because of (2.5)
$D(\omega) = \gamma_{\theta_\omega}^\alpha + 2 \ln \left(\frac{1+q(\omega)}{1-q(\omega)} \right)$	bounds projective diameter, see Corollary 70
$\rho(\omega) = \tanh(D(\omega)/4)$	contraction rate as in the RHS of (1.1); $\rho(\omega) \in (0, 1)$
$B_{\omega,1} = e^{\gamma_\omega^\alpha}$	a lower bound on the random equivariant density h_ω , see Theorem 47
$K_\omega = (1 + \gamma_\omega^\alpha)e^{\gamma_\omega^\alpha}$	bounds aperture of \mathcal{C}_ω , see Theorem 81; see also Theorem 47
$M_\omega = 8(1 - e^{-\gamma_\omega^\alpha})^{-2}$	bounds aperture of the dual of \mathcal{C}_ω , see Theorem 81; $M_\omega \leq 16$

Note that the “the projective diameter” refers to the diameter of the image $\mathcal{L}_\omega \mathcal{C}_\omega$ inside $\mathcal{C}_{\theta_\omega}$ with respect to the (projective) Hilbert metric.

12. Remark. As a continuation of Remark 9, when $H_\omega \leq \min(M, \gamma_{\theta_\omega})^\alpha - 1$ for some $M > 1$ (i.e. $H_\omega \leq \gamma_{\theta_\omega}^\alpha - 1$ and it is bounded) then we can replace γ_ω with $\gamma_{\omega,M} = \min(M, \gamma_\omega)$ (namely assume that γ_ω is bounded above). In this case we have

$$D(\omega) \leq M + 2 \ln \left(\frac{1 + q_M(\omega)}{1 - q_M(\omega)} \right), \quad B_\omega \leq C(M)$$

for some constant $C(M)$, where $q_M(\omega)$ is defined like $q(\omega)$ but with $\gamma_{\omega,M}$ instead of γ_ω .

Next, let $u_\omega \in \mathcal{H}_\omega$ be a random function so that

$$(2.13) \quad \tilde{H}_\omega := \gamma_\omega^{-\alpha} v_\alpha(u_\omega) + H_\omega \leq \gamma_{\theta_\omega}^\alpha - 1$$

and an equivariant family of probability measures μ_ω (i.e. $(T_\omega)_* \mu_\omega = \mu_{\theta_\omega}$). Note that (2.13) is a stronger version of (2.5). Let

$$\tilde{u}_\omega = u_\omega - \mu_\omega(u_\omega).$$

In the circumstances of Section 2.1, the following random variables are used in the formulation of Theorems 39 and 40 (moderate deviations principles), Theorem 43 (CLT rates) and Theorem 45 (moderate local CLT).

Random variable	Role/comments
$\tilde{\rho}(\omega) = \tanh(7D(\omega)/4)$	complex contraction rate as in Theorem 49; $\tilde{\rho}(\omega) \in (0, 1)$
$c_0(\omega) = 3\ \tilde{u}_\omega\ _\infty + \frac{H_\omega}{\gamma_{\theta_\omega}^\alpha - (1 + H_\omega)}$	$c_0(\omega) > 0$ by (2.13)
$E(\omega) = c_0(\omega) (1 + \cosh(7D(\omega)/2))$	determines when transfer operators preserve complex cones
$\bar{D}_\omega = 16e^{\ \tilde{u}_\omega\ _\infty} (1 + v(u_\omega))(1 + \tilde{H}_\omega) D_\omega$	D_ω comes from Assumption 8

2.3.2. Partially expanding maps. In the circumstances of Section 2.2, consider the real cone

$$\mathcal{C}_\omega = \mathcal{C}_{\omega, \kappa_\omega} = \{g \in \mathcal{H}_\omega : g > 0 \text{ and } v(g) \leq s_\omega^{-1} \inf g\}.$$

The following random variables are used in the formulation of Theorems 27, 32, 34 (CLT and ASIP) and Theorems 47 and 49 (Perron-Frobenius rates).

Random variable	Role/comments
$B_\omega = 12(1 + 2/s_\omega)^4$	appears on the RHS of (1.1)
$\zeta_\omega = s_{\theta_\omega} (1 + (1 + s_\omega^{-1})e^{\varepsilon_\omega} H_\omega)$	$\zeta_\omega < 1$ by (2.12)
$D(\omega) = 2 \ln \left(\frac{1+\zeta_\omega}{1-\zeta_\omega} \right) + 2 \ln (1 + \zeta_\omega s_\omega^{-1})$.	bounds projective diameter, see Corollary 70
$\rho(\omega) = \tanh(D(\omega)/4)$	contraction rate as in the RHS of (1.1); $\rho(\omega) \in (0, 1)$
$B_{\omega,1} = 1 + s_\omega^{-1}$	a lower bound on the random equivariant density h_ω , see Theorem 47
$K_\omega = 1 + 2s_\omega^{-1}$	bounds aperture of \mathcal{C}_ω , see Theorem 81; see also Theorem 47
$M_\omega = 6s_\omega^{-1}$	bounds aperture of the dual of \mathcal{C}_ω , see Theorem 81

Finally, let $u_\omega : \mathcal{E}_\omega \rightarrow \mathbb{R}$ be a random function and μ_ω be an equivariant family of probability measures μ_ω . Set $\tilde{u}_\omega = u_\omega - \mu_\omega(u_\omega)$. In the circumstances of Section 2.1, the following random variables are used in the formulation of Theorems 39 and 40 (moderate deviations principles), Theorem 43 (CLT rates) and Theorem 45 (local moderate CLT).

Random variable	Role/comments
$\tilde{\rho}(\omega) = \tanh(7D(\omega)/4)$	complex contraction rate as in Theorem 49; $\tilde{\rho}(\omega) \in (0, 1)$
$c_0(\omega) = \frac{32s_{\theta_\omega}(1+2s_\omega^{-1})e^{\ u_\omega\ _\infty+2\ \phi_\omega\ _\infty}\ \tilde{u}_\omega\ (1+H_\omega)}{1-\zeta_\omega}$	$c_0(\omega) \in (0, \infty)$
$E(\omega) = c_0(\omega) (1 + \cosh(7D(\omega)/2))$	determines when transfer operators preserve complex cones
$\bar{D}_\omega = 16e^{\ \tilde{u}_\omega\ _\infty} (1 + v(u_\omega))(1 + \tilde{H}_\omega)D_\omega$	$D_\omega = \deg T_\omega e^{\ \phi_\omega\ _\infty} = d_\omega e^{\ \phi_\omega\ _\infty}$

2.4. More general (than Section 1.1.1) examples in the smooth case: fiberwise piecewise perturbations of piecewise linear expanding maps. As discussed before, the maps described in Sections 2.1 and 2.2 were essentially considered in [42, 33] and [44], respectively, with the exception that in [42, 33] the potential ϕ_ω did not satisfy (2.4) and (2.5), and in [44] the weaker condition $\int \ln s_\omega d\mathbb{P}(\omega) < 0$ was considered (instead of $s_\omega < 1$), and the potential ϕ_ω did not satisfy the second estimate in (2.12), as well. In comparison with [42], the inequality (2.5) is our main additional assumption on the potential ϕ_ω in the setup of Section 2.1. While we can always work with a random Gibbs measure μ_ω corresponding to a potential ϕ_ω satisfying (2.4) and (2.5), it is interesting to see for which maps these conditions hold true in the smooth case when e^{ϕ_ω} is the Jacobian of T_ω with respect to the volume measure on \mathcal{E}_ω . In this section we will show that (2.4) and (2.5) are valid in the smooth case for certain type of C^2 fiberwise perturbations of piecewise linear maps (and similarly we can consider perturbations of piecewise affine maps, but for the sake of simplicity we will describe only the one dimensional case).

2.4.1. *The piecewise linear case.* The examples in this section are generalizations of the illustrating examples in Section 1.1.1. Let $\mathcal{I}_\omega = \{I_{\omega,i} = [a_i(\omega), b_i(\omega)]\}$ be a (nontrivial) partition of the unit interval $[0, 1)$ into intervals, and on each interval let $\ell_{\omega,i}$ be a linear map that maps $I_{\omega,i}$ to $[0, 1)$ (there are two options, either the decreasing one or the increasing one). Then the slope of $\ell_{\omega,i}$ is $\pm |I_{\omega,i}|^{-1}$, where $|I_{\omega,i}|$ is the length of $I_{\omega,i}$. Let us assume that $I_{\omega,1}$ is the largest interval and set

$$\gamma_\omega(\ell) = |I_{\omega,1}|^{-1} > 1.$$

Next, for each i let $I_{\omega,i_\omega(y)}$ be the unique interval $I_{\omega,i}$ so that $y \in I_{\omega,i}$. Then the map ℓ_ω defined by $\ell_\omega(y) = \ell_{\omega,i_\omega(y)}(y)$ satisfies all the conditions in Section 2.1 in the case $\xi = 1$ with $\gamma_\omega = \gamma_\omega(\ell)$. Moreover, if we consider the smooth case and take e^{ϕ_ω} to be the Jacobian of ℓ_ω then, since the map is piecewise linear we have that H_ω in (2.4) vanishes, and so (2.5) trivially holds true, where we can take the Hölder exponent $\alpha = 1$. Moreover, we have that μ_ω is the Lebesgue measure and $\mathcal{L}_\omega \mathbf{1} = \mathbf{1}$ (and hence $D_\omega = 1$ in this case, and so $\bar{D}_\omega = e^{\|u_\omega\|_\infty} (1 + v_\alpha(u_\omega))$ depends only on u_ω). Recall also that B_ω is bounded (see Remark 12) and note that $q(\omega) = \frac{1}{\gamma_\omega(\ell)\gamma_{\theta\omega}(\ell)}$ in this case.

2.4.2. *Fiberwise piecewise C^2 -perturbations.* Let us explain for which type of piecewise C^2 -perturbations of ℓ_ω the conditions of Section 2.1 with $\xi = 1$ remain true. On each interval $I_{\omega,i}$, without changing the value of $\ell_{\omega,i}$ at the end points, let us take a C^2 perturbation $T_{\omega,i}$ of $\ell_{\omega,i}$ so that

$$(2.14) \quad \|T_{\omega,i} - \ell_{\omega,i}\|_{C^2} \leq \varepsilon_\omega = \frac{1}{2} \min \left(\frac{1}{4} (\gamma_{\theta\omega}(\ell) - 1) (\gamma_\omega(\ell))^2, (\gamma_{\theta\omega}(\ell) - 1), (\gamma_\omega(\ell) - 1) \right).$$

Let us define $T_\omega(y) = T_{\omega,i_\omega(y)}(y)$ (namely by gluing the maps $T_{\omega,i}$). Consider again the smooth case and take e^{ϕ_ω} to be the Jacobian of T_ω . Then μ_ω is equivalent to the Lebesgue measure (since ν_ω in Theorem 47 is the Lebesgue measure).

13. **Lemma.** *Under (2.14) the maps satisfy the conditions in Section 2.1 with $\xi = 1$ and $\gamma_\omega = \gamma_\omega(\ell) - \varepsilon_\omega \geq \frac{\gamma_\omega(\ell)+1}{2}$. Moreover, the potential $\phi_\omega = -\ln J_{T_\omega}$ satisfies (2.4) with $\alpha = 1$ and H_ω so that (2.5) holds true (with $\alpha = 1$).*

Proof. First, it is clear that we can take $\gamma_\omega = \gamma_\omega(\ell) - \varepsilon_\omega \geq \frac{\gamma_\omega(\ell)+1}{2}$. In order to show that condition (2.5) is in force it is enough to show that the derivative of each composition $\phi_\omega \circ y_{\omega,i}$ is bounded by $\gamma_{\theta\omega} - 1$. To establish that, note that $y_{\omega,i}(x) = T_{\omega,i}^{-1}(x)$ and so

$$\sup_x |(\phi_\omega \circ y_{\omega,i})'(x)| = \sup_x (|\phi'_\omega(y_{\omega,i}(x))| \cdot |y'_{\omega,i}(x)|) \leq \sup_y \sup_i \left| \frac{T''_{\omega,i}(y)}{(T'_{\omega,i}(y))^2} \right| \leq \frac{4\varepsilon_\omega}{(\gamma_\omega(\ell))^2}$$

where in the last inequality we have used that $|T''_{\omega,i}(y)| = |T''_{\omega,i}(y) - \ell''_{\omega,i}(y)| \leq \varepsilon_\omega$ and that

$$|T'_{\omega,i}(y)| \geq |\ell'_{\omega,i}(y)| - \varepsilon_\omega \geq |I_{\omega,i}|^{-1} - \varepsilon_\omega \geq \gamma_\omega(\ell) - \varepsilon_\omega \geq \frac{1}{2} \gamma_\omega(\ell).$$

Finally, using that $\varepsilon_\omega \leq \frac{1}{8} (\gamma_{\theta\omega}(\ell) - 1) (\gamma_\omega(\ell))^2$ and that $\varepsilon_\omega \leq \frac{1}{2} (\gamma_\omega(\ell) - 1)$ we see that

$$\sup_x |(\phi_\omega \circ y_{\omega,i})'(x)| \leq \frac{4\varepsilon_\omega}{(\gamma_\omega(\ell))^2} \leq \frac{1}{2} (\gamma_{\theta\omega}(\ell) - 1) \leq \gamma_{\theta\omega}(\ell) - \varepsilon_\omega - 1 \leq \gamma_{\theta\omega} - 1.$$

□

14. **Remark.** As mentioned in Remark 12, when H_ω is bounded then B_ω from (1.1) will be bounded in our applications. Notice that once (2.5) established H_ω will be bounded if γ_ω is. In our case this just means that $\gamma_\omega(\ell)$ is bounded, namely that the number of intervals in the partition \mathcal{I}_ω is bounded.

15. **Remark.** In certain instances $|I_{\omega,1}|$ is bounded away from 1 (e.g. $T_\omega x = (m_\omega x) \bmod 1$, $m_\omega \in \mathbb{N}$). In this case we can take allow larger ε_ω so that the resulting perturbation will not be uniformly expanding (as γ_ω could be arbitrary close to 1).

3. PRELIMINARIES AND MAIN RESULTS

3.1. The random probability space: on the choice of measures μ_ω . For both classes of maps considered in Sections 2.1 and 2.2 let μ_ω be the Gibbs measures corresponding to the potential ϕ_ω . The detailed exposition of these measures is postponed to Section 3.8 (where our results concerning effective rates are described), and for the meanwhile we refer to [42] and [44] for the construction and the main properties of these measures (see also Theorem 47). For instance they are equilibrium states and they have an exponential decay of correlations for Hölder continuous functions. Let us note that the smooth case discussed in Section 1 corresponds to the choice of $\phi_\omega = -\ln(J_{T_\omega})$ (see Section 2.4), while the choice of $\phi_\omega = 0$ corresponds to random measures of maximal entropy (more generally the case $\phi_\omega = \psi_\omega/T$ for a sufficiently large T and a sufficiently regular potential ψ_ω corresponds to the high temperature regime, see [44, Theorem D]).

3.2. Upper mixing coefficients. Let $X = \{X_j : j \in \mathbb{Z}\}$ be a stationary sequence of random variables (taking values on some measurable space) which generates the system $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$, so that θ is the left shift on the paths of X_j , namely $\theta((X_j)) = (X_{j+1})$.

Recall next that the k -th upper α , ϕ and ψ mixing coefficients of the sequence $\{X_j\}$ are the smallest numbers $\alpha_U(k)$, $\phi_U(k)$ and $\psi_U(k)$ so that for every n and a set A measurable²³ with respect to $\sigma\{X_j : j \leq n\}$ and a set B measurable with respect to $\sigma\{X_m : m \geq n+k\}$ we have

$$\mathbb{P}(A \cap B) \leq \mathbb{P}(A)\mathbb{P}(B)(1 + \psi_U(k)),$$

$$\mathbb{P}(A \cap B) \leq \mathbb{P}(A)\mathbb{P}(B) + \phi_U(k)\mathbb{P}(A)$$

and

$$\mathbb{P}(A \cap B) \leq \mathbb{P}(A)\mathbb{P}(B) + \alpha_U(k).$$

Clearly

$$\alpha_U(k) \leq \phi_U(k) \leq \psi_U(k).$$

Notice next that $\alpha_U(k)$, $\phi_U(k)$ and $\psi_U(k)$ are decreasing, and so

$$(3.1) \quad \limsup_{k \rightarrow \infty} \eta(k) = \lim_{k \rightarrow \infty} \eta(k) = \inf_k \eta(k)$$

where η is either α_U , ϕ_U or ψ_U .

16. Remark. Note that due to stationarity we can always consider only $n = 0$ in the definitions of the upper mixing coefficients. We prefer to present the upper mixing coefficients in the above more general form in order to avoid repeating that both forms are equivalents in the course of the proofs.

17. Remark (Two sided mixing coefficients). Recall that the (two sided) mixing coefficients $\alpha(k)$, $\phi(k)$, $\psi(k)$ are defined similarly through the inequalities

$$|\mathbb{P}(A \cap B) - \mathbb{P}(A)\mathbb{P}(B)| \leq \mathbb{P}(A)\mathbb{P}(B)\psi(k),$$

$$|\mathbb{P}(A \cap B) - \mathbb{P}(A)\mathbb{P}(B)| \leq \mathbb{P}(A)\phi(k),$$

and

$$|\mathbb{P}(A \cap B) - \mathbb{P}(A)\mathbb{P}(B)| \leq \alpha(k).$$

Clearly $\psi_U(k) \leq \psi(k)$, $\phi_U(k) \leq \phi(k)$ and $\alpha_U(k) \leq \alpha(k)$. The sequences $\alpha(k)$, $\phi(k)$ and $\psi(k)$ are classical quantities measuring the long range weak-dependence of the sequence $\{X_n\}$ (see [13, 19]).

18. Example. [(properly) mixing examples]

²³Here $\sigma\{X_j : j \in \mathcal{I}\}$ is the σ -algebra generated by $\{X_j : j \in \mathcal{I}\}$, where $\mathcal{I} \subset \mathbb{Z}$.

- (i) $\phi(n)$ decays exponentially fast as $n \rightarrow \infty$ when X_j is a geometrically ergodic Markov chain, namely if R is the transition operator²⁴ of the chain then

$$\|R^n - \mu\|_\infty \leq C\delta^n$$

for some $C > 0$ and $\delta \in (0, 1)$ (this is a consequence of [13, Theorem 3.3]).

- (ii) $\phi(n)$ also decays exponentially fast for uniformly contacting Markov chains in the sense of Dobrushin (see [35, Lemma 3.3]), namely if for some $n_0 \in \mathbb{N}$,

$$\sup_{x, y, \Gamma} |\mathbb{P}(X_{n_0} \in \Gamma | X_0 = x) - \mathbb{P}(X_{n_0} \in \Gamma | X_0 = y)| < 1$$

where Γ ranges over all measurable subsets of the underlying state space and (x, y) ranges over pairs of states.

- (iii) $\psi(n)$ decays exponentially fast as $n \rightarrow \infty$ when X_j is a Markov chain satisfying the two sided Doeblin condition: there exists an $\ell > 0$ such that for any measurable subset Γ on the state space of X_j and a state x we have

$$C_1\eta(\Gamma) \leq \mathbb{P}(X_\ell \in \Gamma | X_0 = x) \leq C_2\eta(\Gamma)$$

for some constants $C_i > 0$ and a probability measure η (see [11]).

- (iv) $\psi(n)$ decays exponentially fast when X_j is the j -th coordinate of a topologically mixing subshift of finite type, see [12]. Therefore, it decays exponentially fast also when X_n is measurable with respect to $T^{-n}\mathcal{M}$, where T is an Anosov map and \mathcal{M} is a Markov partition with a sufficiently small diameter (see also [12]).
- (v) $\psi(n)$ decays exponentially fast also when X_j is the j -coordinate of the symbolic representation of a Gibbs Markov map, see [1] (like in (iv) this has an interpretation involving the Gibbs Markov map itself).
- (vi) $\alpha(n) = O(n^{-(d-1)})$ when $X_j = T^j X_0$ and T is a Young tower whose tails decay like $O(n^{-d})$ and X_0 is measurable with respect to the partition generating the tower (see the proof of [31, Lemma 4] or [34, Proposition 4.14]).
- (vii) $\alpha(n) = O(n^{-(A-1)})$ if (X_j) is a real valued Gaussian sequence such that $\text{Cov}(X_n, X_0) = O(n^{-A})$ for some $A > 1$ (see [19, Section 2.1, Corollary 2] and also [13, Section 7]).

We refer to [19] for additional examples.

While our results are new also if we work only with the usual mixing coefficients α, ϕ, ψ presented above, the proofs only require assumptions on the upper mixing coefficients. In the following remark we will discuss a situation in which the upper mixing coefficients come in handy.

19. Remark. The only place where ergodicity of θ (i.e. of $\{X_j : j \in \mathbb{Z}\}$) is used is to insure that there is a number $\sigma \geq 0$ such that (like in Theorems 27 and 32),

$$\sigma^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}(S_n^\omega u), \mathbb{P} \text{ a.s.}$$

However, without ergodicity, by applying an appropriate ergodic decomposition theorem we will get that the limit²⁵

$$\sigma^2(\omega) = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}(S_n^\omega u)$$

exists, but now it is no longer a constant, and instead it is measurable with respect to the σ -algebra of invariant sets. Moreover, we will get the coboundary characterization (as in Theorems 27 and 32) for the positivity of this limit, but on each ergodic component. Hence our results hold true if we start with a component for which the asymptotic variance $\sigma^2(\omega)$ is positive (when the asymptotic variance vanishes then the CLT is degenerate, namely $S_n^\omega u / \sqrt{n}$ converges to 0 in L^2 , so there is nothing to prove).

²⁴Namley, R maps a bounded function g to a function Rg given by $Rg(x) = \mathbb{E}[g(X_1) | X_0 = x]$.

²⁵i.e. the asymptotic variance.

We conclude that in the non-ergodic case we can just assume that each ergodic component is mixing. For instance, the above modification of our results holds true when X_j is a stationary finite state Markov chain whose transition matrix is composed of blocks which have only positive entries (perhaps after several steps). Indeed, each ergodicity class gives raise to an exponentially fast ψ mixing sequence (see [13]), and thus $\psi_U(n)$ decays exponentially fast (using that for sets from different classes we have $\mathbb{P}(A \cap B) = 0$).

3.3. Quenched limit theorems for random Birkhoff sums. Let $u_\omega : \mathcal{E}_\omega \rightarrow \mathbb{R}$ be a random function (i.e. $u(\omega, x) = u_\omega(x)$ is measurable) so that $u_\omega \in \mathcal{H}_\omega$ (i.e. u_ω is α -Hölder continuous). Let us consider the corresponding random Birkhoff sums

$$S_n^\omega u = \sum_{j=0}^{n-1} u_{\theta^j \omega} \circ T_\omega^j.$$

In this paper, under appropriate assumptions on u_ω , when ω is fixed (chosen from a set of probability one), we will prove limit theorems for the sequence of functions $S_n^\omega u(\cdot)$ considered as random variables on the probability space $(\mathcal{E}_\omega, \mathcal{B}_\omega, \mu_\omega)$, where \mathcal{B}_ω is the Borel σ -algebra on \mathcal{E}_ω .

3.4. The CLT and LIL. First, let us note that, in order to avoid repetitions, in all the result formulated in this section the random variables defined in Sections 2.3.1 and 2.3.2 will be in constant use, sometime without referring to these sections.

Next, in order to formulate our first set of sufficient conditions for the CLT we consider the following assumption.

20. Assumption. There is a measurable set $A \subset \Omega$ with positive probability so that for all $\omega \in A$ we have $\rho(\omega) \leq 1 - \varepsilon$ and $B_\omega \leq M$ for some $\varepsilon, M > 0$ and that for all $r \in \mathbb{N}$ there is a set A_r which is measurable with respect to $\sigma\{X_j, |j| \leq r\}$ so that $\beta_r = \mathbb{P}(A \setminus A_r) \rightarrow 0$ and $\lim_{r \rightarrow \infty} \mathbb{P}(A_r) = \mathbb{P}(A)$.

21. Remark. When $\rho(\omega)$ and B_ω depend only on $X_j, |j| \leq d$ for some d then we can just take A to be a set of the form $A = A_{\varepsilon, M} = \{\omega : \rho(\omega) \leq 1 - \varepsilon, B_\omega \leq M\}$ for a sufficiently small ε and a sufficiently large M (or any measurable subset of such a set with positive probability). In this case $\beta_r = 0$ for all $r > d$. It is important to note that in these circumstances our conditions for the CLT (see Theorems 27 and 32) will only involve some decay rates for $\alpha_U, \phi_{R,U}$ or ψ_U and integrability assumptions on the norm $\|u_\omega - \mu_\omega(u_\omega)\|$, which are independent of all the random variables describing the maps T_ω (the general principle is that if $\omega \rightarrow T_\omega$ depends only on finitely many variables then so are the describing parameters, and then we have no additional requirements from these parameters).

22. Example. Let us consider the example in Section 1.1.1. Then using the formulas for $\rho(\omega)$ and B_ω given in Example 2 we see that $\rho(\omega)$ depends only on $a_{\theta\omega}$ and B_ω depends only on a_ω . Thus, if a_ω depends only on finitely many coordinates then both $\rho(\omega)$ and B_ω depend on finitely many coordinates and, as discussed in Remark 21, Assumption 20 is in force.

To provide examples where Assumption 20 holds true with non-vanishing β_r we will use the following simple result.

23. Lemma. *Suppose that the following approximation condition holds true: there are²⁶ random variables $\rho_r = \rho_r(\omega)$ and $B_r = B_{\omega, r}$ measurable with respect to $\sigma\{X_j, |j| \leq r\}$ so that*

$$(3.2) \quad \max(\|\rho(\omega) - \rho_r(\omega)\|_{L^1}, \|B_\omega - B_{\omega, r}\|_{L^1}) \leq \delta_r \rightarrow 0.$$

Then Assumption 20 holds with $A = A_{\varepsilon, M} = \{\omega : \rho(\omega) \leq 1 - \varepsilon, B_\omega \leq M\}$, $A_r = \{\omega : \rho_r(\omega) \leq 1 - \varepsilon + \sqrt{\delta_r}, B_{\omega, r} \leq M + \sqrt{\delta_r}\}$ and $\beta_r \leq 2\sqrt{\delta_r}$ (where ε is small enough and M is large enough to insure that $\mathbb{P}(A) > 0$).

²⁶Note that condition (3.2) can also be written as

$$\max(\|\rho - \mathbb{E}[\rho|X_{-r}, \dots, X_r]\|_{L^1}, \|B - \mathbb{E}[B|X_{-r}, \dots, X_r]\|_{L^1}) \leq \delta_r$$

where $\rho = \rho(\omega)$ and $B = B_\omega$. This condition is fulfilled when $\rho(\dots, X_{-1}, X_0, X_1, \dots)$ and $B_{\dots, X_{-1}, X_0, X_1, \dots}$ weakly depend (in an L^1 -sense) on the coordinates $X_j, |j| \geq r$. Limit theorems under conditions similar to (3.2) (with some decay rate for δ_r) have been studied extensively in weak dependence theory, see [9, 37] (where iid X_j are considered).

Proof. Using the Markov inequality we see that

$$\mathbb{P}(A_r) \leq \mathbb{P}(|\rho - \rho_r| \geq \sqrt{\delta_r}) + \mathbb{P}(|B - B_r| \geq \sqrt{\delta_r}) + \mathbb{P}(A_{\varepsilon+2\sqrt{\delta_r}, M+2\sqrt{\delta_r}}) \leq 2\sqrt{\delta_r} + \mathbb{P}(A_{\varepsilon+2\sqrt{\delta_r}, M+2\sqrt{\delta_r}})$$

and

$$\mathbb{P}(A \setminus A_r) \leq \mathbb{P}(|\rho - \rho_r| \geq \sqrt{\delta_r}) + \mathbb{P}(|B - B_r| \geq \sqrt{\delta_r}) \leq 2\sqrt{\delta_r}.$$

Hence $\lim_r \mathbb{P}(A_r) = \mathbb{P}(A)$ and $\beta_r = \mathbb{P}(A \setminus A_r) \leq 2\sqrt{\delta_r}$. \square

24. Remark. When $\omega \rightarrow B_\omega$ is in $L^1(\Omega, \mathcal{F}, \mathbb{P})$ then there is always a sequence δ_r satisfying (3.2), but our main results involve certain decay rates for β_r .

25. Example. Let us return to the example in Section 1.1.1, but in addition we suppose that the coordinates ω_j of ω take values on some bounded metric space (Y, d_Y) (not necessarily compact). Let us define a metric on Ω by

$$d_\Omega(\omega, \omega') = \sum_j 2^{-|j|} d_Y(\omega_j, \omega'_j).$$

Then the left shift θ is Lipschitz continuous.

Let us assume that a_ω is a Hölder continuous function of ω with some exponent κ . As discussed in Example 1.1.1, we have

$$B_\omega = 24e^{4a_\omega^{-\alpha}} (1 + a_\omega^{-\alpha})^2$$

and

$$\rho(\omega) = \frac{e^{\frac{1}{2}a_{\theta\omega}^{-\alpha}} (1 + a_{\theta\omega}^\alpha) - (1 - a_{\theta\omega}^\alpha)}{e^{\frac{1}{2}a_{\theta\omega}^{-\alpha}} (1 + a_{\theta\omega}^\alpha) + (1 - a_{\theta\omega}^\alpha)}.$$

Since $a_\omega \in [\frac{1}{2}, 1)$ we see that B_ω is a Lipschitz continuous function of a_ω and $\rho(\omega)$ is a Lipschitz continuous function of $a_{\theta\omega}$. Thus both B_ω and $\rho(\omega)$ are Hölder continuous functions of ω with the same exponent κ .

We claim that condition (3.2) holds true with $\beta_r = O(2^{-\kappa r/2})$ (in fact we will get (stronger) estimates in L^∞). Note that such exponential rate of decay will be more than enough for the decay rates in Theorem 27 (the CLT) to hold. To prove the claim, observe that for every Hölder continuous function $R : \Omega \rightarrow \mathbb{R}$ with exponent $\kappa \in (0, 1]$ and all points ω, ω' such that $\omega_j = \omega'_j$ if $|j| \leq d$ we have

$$|R(\omega) - R(\omega')| \leq L_R D^\kappa 2^{-\kappa(d-1)}$$

where L_R is the Hölder constant of R and $D = \text{Diam}(Y)$. Therefore,

$$\sup_\omega |R(\omega) - R_r(\omega)| = O(2^{-\kappa r})$$

where $R_r(\omega) = \inf\{R(\omega') : \omega'_j = \omega_j \text{ if } |j| \leq r\}$ (apply this with $R(\omega) = B_\omega$ and $R(\omega) = \rho(\omega)$). Note that $R_r(\omega)$ depends only on the coordinates ω_j with $|j| \leq r$. Finally, Assumption 20 holds because of Lemma 23, and, moreover, we have $\beta_r = O(2^{-\kappa r/2})$,

26. Example. In the context of Section 1.1.1, let us consider the case when a_ω has the form

$$a_\omega = \sum_{j=0}^{\infty} f_j(\omega_{-j}, \dots, \omega_0, \dots, \omega_j)$$

for some measurable functions f_j on \mathcal{Y}^{2j+1} , where \mathcal{Y} is the space such that $\Omega = \mathcal{Y}^{\mathbb{Z}}$. Then

$$\left\| a_\omega - \sum_{j \leq r} f_j(\omega_{-j}, \dots, \omega_j) \right\|_{L^1(\mathbb{P})} \leq \sum_{j \geq r} \|f_j\|_{L^1(\mathbb{P})} := A_1(r).$$

Similarly,

$$\left\| a_{\theta\omega} - \sum_{j \leq r} f_j(\omega_{-j+1}, \dots, \omega_{j+1}) \right\|_{L^1(\mathbb{P})} \leq \sum_{j \geq r} \|f_j\|_{L^1(\mathbb{P})} = A_1(r).$$

Now, since B_ω is a Lipschitz continuous function of a_ω and $\rho(\omega)$ is a Lipschitz continuous function of $a_{\theta\omega}$ we conclude that condition (3.2) holds true with $\beta_r = O(A_1(r))$ which converges to 0 if $\|f_j\|_{L^1}$ is summable (to obtain decay rates for β_r we only need to assume some decay rates for the norms $\|f_j\|_{L^1}$ as $j \rightarrow \infty$).

To give a concrete example when $a_\omega \in [\frac{1}{2}, 1)$ we can, for instance, consider the case when

$$f_j(\omega_{-j}, \dots, \omega_j) = a_j + b_j \mathbb{I}(\omega_j \in \mathcal{A})$$

for some measurable set \mathcal{A} such that $0 < \mathbb{P}(\omega_j \in \mathcal{A}) < 1$ and positive sequences (a_j) and (b_j) such that

$$\frac{1}{2} \leq \sum_j b_j \quad \text{and} \quad \sum_j (a_j + b_j) = 1.$$

Then $a_\omega \in (\frac{1}{2}, 1)$ (\mathbb{P} -a.s.) and a_ω can take arbitrarily close to 1 values if $\mathbb{P}(\omega_k \in \mathcal{A}; k \leq d) > 0$ for every $d > 0$ (e.g. in the iid case or for the Markov chains in Example 18 (iv) when $\ell = 1$).

We can also assume that $\omega_j \in [a, b]$, $0 < a < b$ and then take $f_j(\omega_{-j}, \dots, \omega_j) = \alpha_j(\omega_0, \dots, \omega_j)\omega_j$ for some $\alpha_j(\cdot)$ so that $\frac{1}{2a} \leq \sum_j \alpha_j < \frac{1}{b}$, assuming that $b < 2a$. For instance, if also $b < \frac{4a}{3}$ then α_j can have the form $\alpha_j(\cdot) = v_j(\omega_0)\gamma_j$ for a positive series such that $\frac{3}{4a} \leq \sum_j \gamma_j \leq \frac{1}{b}$ and a random variable $v(\omega_0)$ such that $\frac{2}{3} < v(\omega_0) < 1$ and $\|v\|_{L^\infty} = 1$.

27. Theorem (CLT). *Let Assumption 20 be in force. Assume that the random variable $\omega \rightarrow \|u_\omega - \mu_\omega(u_\omega)\|$ is in $L^p(\Omega, \mathcal{F}, \mathbb{P})$ for some $p > 2$ so that $\sum_j (\ln j \beta_{Cj/\ln j})^{1-2/p} < \infty$ for all $C > 0$. In addition, assume that one of the following conditions is in force:*

$$(M1) \sum_j (\alpha_U(Cj/\ln j))^{1-2/p} < \infty \text{ for all } C > 0;$$

$$(M2) \limsup_{k \rightarrow \infty} \phi_U(k) < \mathbb{P}(A) \quad (\text{i.e. } \phi_U(k) < \mathbb{P}(A) \text{ for some } k);$$

$$(M3) \limsup_{k \rightarrow \infty} \psi_U(k) < \frac{1}{1-\mathbb{P}(A)} - 1 \quad (\text{i.e. } \psi_U(k) < \frac{1}{1-\mathbb{P}(A)} - 1 \text{ for some } k).$$

Then:

(i) There is a number $\sigma \geq 0$ so that for \mathbb{P} -a.e ω we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}_{\mu_\omega}(S_n^\omega u) = \sigma^2.$$

Moreover, $\sigma > 0$ if and only if the function $U(\omega, x) = \sum_{j=0}^{n_A(\omega)-1} (u_{\theta^j \omega} \circ T_\omega^j x - \mu_{\theta^j \omega}(u_{\theta^j \omega}))$ has the form $U(\omega, x) = q(\omega, x) - q(\theta^{n_A(\omega)}, T_\omega^{n_A(\omega)} x)$ for \mathbb{P}_A almost every ω and all x , where n_A is the first return time to A , $\mathbb{P}_A(\cdot) = \mathbb{P}(\cdot|A)/\mathbb{P}(A)$ is the conditional measure on A and q is a measurable function so that $\int_A \int_{\mathcal{E}_\omega} |q(\omega, x)|^2 d\mu_\omega(x) d\mathbb{P}(\omega) < \infty$.

(ii) The sequence $S_n^\omega u$ obeys the CLT: for every real t we have

$$\lim_{n \rightarrow \infty} \mu_\omega \{x : n^{-1/2} (S_n^\omega u(x) - \mu_\omega(S_n^\omega u)) \leq t\} = \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^t e^{-\frac{s^2}{2\sigma^2}} ds$$

where if $\sigma = 0$ the above right hand side is interpreted as the distribution function of the constant random variable 0.

(iii) Set $\tau(\omega, x) = (\theta\omega, T_\omega x)$, $\mu = \int_\Omega \mu_\omega d\mathbb{P}(\omega)$ and $\tilde{u}(\omega, x) = u_\omega(x) - \mu_\omega(u_\omega)$. If $\sigma > 0$ then the following functional version of the law of iterated logarithm (LIL) holds true. Let $\zeta(t) = (2t \log \log t)^{1/2}$ and

$$\eta_n(t) = (\zeta(\sigma^2 n))^{-1} \sum_{j=0}^{k-1} (\tilde{u} \circ \tau^j + (nt - k)\tilde{u} \circ \tau^k)$$

for $t \in [\frac{k}{n}, \frac{k+1}{n}]$, $k = 0, 1, \dots, n-1$. Then μ -a.s. the sequence of functions $\{\eta_n(\cdot), n \geq 3/\sigma^2\}$ is relatively compact in $C[0, 1]$ (the space of continuous functions on $[0, 1]$ with the supremum norm), and the set of limit points as $n \rightarrow \infty$ coincides with the set K of absolutely continuous functions $x \in C[0, 1]$ so that $\int_0^1 (\dot{x}(t))^2 dt \leq 1$.

28. **Remark.** In the circumstances of Example 22, β_r vanishes for all r large enough, and so the summability assumption on β_r holds. In the circumstances of Example 25, β_r decays exponentially fast as $r \rightarrow \infty$ and so the latter condition still holds true. In the circumstances of Example 26 the summability condition $\sum_j (\ln j \beta_{Cj/\ln j})^{1-2/p} < \infty$ holds if

$$\sum_{|j| \geq r} \|f_j\|_{L^1} = O(r^{-q})$$

for some q such that $q(1 - 2/p) > 1$.

The proof of Theorem 27 starts in Section 4.1 and it is completed in Section 4.1.2. The proof of Theorem 27 is based on inducing, and more precisely we apply [39, Theorem 2.3]. The role the assumptions on the upper mixing coefficient play is that, together with the effective random RPF rates (1.1), they allow us to verify the abstract conditions of [39, Theorem 2.3] with the set $Q = A$ (where Q is in the notations of [39, Theorem 2.3]).

29. **Remark.** When using condition (M1) we only need that $\sum_j (\ln j \beta_{j/(3C \ln j)})^{1-2/p} < \infty$ and $\sum_j (\alpha_U(Cj/\ln j))^{1-2/p} < \infty$ for some C so that $C|\ln(1 - \mathbb{P}(A)/2)|(1 - 2/p) > 1$.

When using (M2) we only need that $\sum_j (\ln j \beta_{j/(3C \ln j)})^{1-2/p} < \infty$ for some C so that $C|\ln \delta|(1 - 2/p) > 1$, where $\delta = 1 - \mathbb{P}(A) + \limsup_{r \rightarrow \infty} \phi_U(r) < 1$.

When using (M3) we only need that $\sum_j (\ln j \beta_{j/(3C \ln j)})^{1-2/p} < \infty$ for some C so that $C|\ln \delta|(1 - 2/p) > 1$, where $\delta = (1 + \limsup_{r \rightarrow \infty} \psi_U(r))(1 - \mathbb{P}(A)) < 1$.

Next, let us provide alternative conditions for the CLT which involve a stronger type of approximation and moment assumptions on B_ω , but do not require any approximation rates.

30. **Assumption.** There is a sequence $\beta_r \rightarrow 0$ as $r \rightarrow \infty$ so that for every r there is a random variable $\rho_r(\omega)$ which is measurable with respect to $\sigma\{X_j; |j| \leq r\}$ and

$$\|\rho - \rho_r\|_{L^\infty} \leq \beta_r.$$

Namely,

$$\lim_{r \rightarrow \infty} \|\rho - \mathbb{E}[\rho | X_{-r}, \dots, X_r]\|_{L^\infty} = 0.$$

31. **Example.** (i) Assumption 30 holds when $\rho(\dots, X_{-1}, X_0, X_1, \dots)$ depends on finitely many of the X_j 's (in this case we can take $\rho_r = \rho$ for r large enough). We refer to Example 22 for an explicit example (namely the one in Section 1.1.1 with a_ω like in Example 22).

(ii) Assumption 30 also holds true in the context of Example 25 since the estimates obtained there were actually in L^∞ .

(iii) In the context of Example 26 we get that

$$\left\| a_{\theta_\omega} - \sum_{0 \leq j \leq r} f_j(\omega_{-j+1}, \dots, \omega_{j+1}) \right\|_{L^\infty(\mathbb{P})} \leq \sum_{j \geq r} \|f_j\|_{L^\infty(\mathbb{P})} := A_\infty(r).$$

Using that $\rho(\omega)$ is a Lipschitz continuous function of a_{θ_ω} we conclude that

$$\|\rho - \mathbb{E}[\rho | X_{-r}, \dots, X_r]\|_{L^\infty} \rightarrow 0 \text{ as } r \rightarrow \infty$$

if $\sum_j \|f_j\|_{L^\infty} < \infty$ (and so Assumption 30 holds).

32. **Theorem.** *Let Assumption 30 be in force. Moreover, assume that*

$$\limsup_{s \rightarrow \infty} \psi_U(s) < \infty \text{ (i.e. } \psi_U(s) < \infty \text{ for some } s)$$

and that $\|u_\omega - \mu_\omega(u_\omega)\|, B_\omega \in L^{3+\delta}(\Omega, \mathcal{F}, \mathbb{P})$ for some $\delta > 0$.

(i) *There is a number $\sigma \geq 0$ so that for \mathbb{P} -a.a. ω we have*

$$\sigma^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}_{\mu_\omega}(S_n^\omega u).$$

Moreover, $\sigma = 0$ if and only if $\tilde{u}(\omega, x) = q(\omega, x) - q(\theta\omega, T_\omega x)$ for some measurable function $q(\omega, x)$ so that $\int q^2(\omega, x) d\mu_\omega(x) d\mathbb{P}(\omega) < \infty$.

(ii) The CLT as stated in Theorem 27 (ii) is valid.

(iii) The functional LIL as stated in Theorem 27 (iii) is valid.

Note that the results in Theorem 32 are slightly better than Theorem 27 since we obtain a simpler coboundary characterization for the positivity of σ .

The proof of Theorem 27 appears in Section 4.2, and, like the proof of Theorem 27, it is also based on applying [39, Theorem 2.3]. However, even though the conditions of [39, Theorem 2.3] are related to an inducing strategy, we will apply it with the set $Q = \Omega$, namely we will “induce” on Ω , so the proof will not really be based on inducing. In this case the conditions of [39, Theorem 2.3] concern the asymptotic behavior of the system $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ itself (that is, of $\theta^n \omega$ as $n \rightarrow \infty$) and not the induced system. Even though this is a stronger requirement, we will verify these conditions using the assumptions on the upper mixing coefficients together with the effective random RPF rates (Theorem 47).

33. Remark. Besides the additional integrability assumptions in 32, the main difference between Theorems 27 and 32 is that in the former we essentially require certain L^1 -approximation rates (decay rates for β_r), while in the latter we do not require such rates, but instead we work with the stronger L^∞ -approximation coefficients, and only with the upper ψ -mixing coefficients. On the other hand, the restrictions on $\limsup_{k \rightarrow \infty} \psi_U(k)$ in Theorem 32 are much weaker than the ones in Theorem 27. Concerning the additional integrability assumption, as explained in Remark 12 (see also Remark 14), under the additional condition on H_ω described there B_ω is bounded, and so in this case the additional requirement that $B_\omega \in L^{3+\delta}(\Omega, \mathcal{F}, \mathbb{P})$ is always satisfied, and the only true integrability condition in Theorem 32 is $\|u_\omega - \mu_\omega(u_\omega)\| \in L^{3+\delta}(\Omega, \mathcal{F}, \mathbb{P})$ (like in Theorem 27). Finally, recall that B_ω is bounded for the piecewise affine maps and their perturbations consider in Section 2.4. Thus in this case $\|u_\omega - \mu_\omega(u_\omega)\| \in L^{3+\delta}(\Omega, \mathcal{F}, \mathbb{P})$ is the only integrability condition needed.

3.5. The ASIP. In this section we further assume that there is a random variable $N(\omega)$ so that

$$(3.3) \quad v(g \circ T_\omega) \leq N(\omega)v(g)$$

for all functions g with $v(g) < \infty$. Note that when the maps T_ω are piecewise differentiable with bounded derivatives then we can always take $N(\omega) = \sup \|DT_\omega\|$. Our next result is about almost sure approximation of $S_n^\omega u$ by sums of independent Gaussians.

34. Theorem (ASIP). *Let Assumption 30 hold true and suppose that $\sigma > 0$. Suppose²⁷ also that²⁸*

$$(3.4) \quad \limsup_{s \rightarrow \infty} \psi_U(s) < \frac{1}{\mathbb{E}_\mathbb{P}[\rho]} \quad (\text{i.e. } \psi_U(s) < \frac{1}{\mathbb{E}_\mathbb{P}[\rho]} \text{ for some } s).$$

Further assume that $\omega \rightarrow B_\omega$ belongs to L^{2p} and $\omega \rightarrow N(\omega)$ and $\omega \rightarrow \|u_\omega - \mu_\omega(u_\omega)\|$ belong to $L^p(\Omega, \mathcal{F}, \mathbb{P})$ for some $p > 8$. Let $\tilde{u}_\omega(x) = u_\omega(x) - \mu_\omega(u_\omega)$. Then there is a coupling of $u_{\theta^j \omega} \circ T_\omega^j$ (considered as a sequence of random variables on the probability space $(\mathcal{E}_\omega, \mu_\omega)$) with a sequence of independent centered Gaussian random variables Z_j so that for every $\varepsilon > 0$,

$$\max_{1 \leq k \leq n} \left| S_k^\omega \tilde{u} - \sum_{j=1}^k Z_j \right| = O(n^{1/4 + \frac{\varepsilon}{2p} + \varepsilon}), \quad \text{a.s.}$$

and

$$\left\| \sum_{j=1}^n Z_j \right\|_2^2 = \text{Var}_{\mu_\omega}(S_n^\omega u) + O(n^{1/2 + 3/p + \varepsilon}).$$

The proof of Theorem 34 appears in Section 4.3.

²⁷Since $\mathbb{E}_\mathbb{P}[\rho] < 1$ this condition holds true when the limit superior does not exceed 1.

²⁸This condition always holds when (X_j) is ψ -mixing, and we refer to Example 18.

35. Example. For the example described in Section 1.1.1 we have that B_ω is bounded and (3.3) holds with $N(\omega) = (1 - a_\omega)^{-1}$ (i.e. the maximal amount of expansion). We conclude that under (3.4), the ASIP rates $O(n^{1/4 + \frac{9}{2p} + \varepsilon})$ hold if $\omega \rightarrow \|u_\omega\|$ is in L^p and $a_\omega = 1 + 1/R_\omega$ for some $R_\omega \in L^p$. Finally, let us note that this result is already meaningful when X_j are independent. In this case $\psi_U(n) = 0$ and we can just consider any function $\omega \rightarrow a_\omega$, $a_\omega \in (\frac{1}{2}, 1)$ of the Bernoulli shift $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ such that $(1 - a_\omega)^{-1} \in L^p$ (still, we can consider such functions of any shift system generated by an arbitrary ψ -mixing sequence).

36. Remark. More generally, as discussed in Remark 12 (see also Remark 14), when, in addition to (2.5) we have $\|H_\omega\|_{L^\infty} < \infty$ then B_ω is bounded (in particular we). Thus, for such maps the only true integrability conditions in Theorem 34 are $N(\omega), \|\tilde{u}_\omega\| \in L^p$. Finally, recall that B_ω is bounded for the piecewise affine maps and their perturbations considered in Section 2.4. Now, for such maps $N(\omega)$ is the supremum norm of the (piecewise) gradient and so Theorem 34 holds true when the supremum norm and $\omega \rightarrow \|\tilde{u}_\omega\|$ are in L^p .

3.6. Large deviations principles with a quadratic rate function. Consider the following additional condition.

37. Assumption. (i) The random variable E_ω defined in Section 2.3.1 (or Section 2.3.2) is bounded.
(ii) In the setup of Section 2.1, (2.4) is satisfied with some H_ω so that

$$Z_\omega := \gamma_\omega^{-\alpha} v(u_\omega) + H_\omega \leq \gamma_{\theta\omega}^\alpha - 1.$$

38. Remark. In the setup of Section 2.1, the condition that E_ω is bounded essentially means that $\|u_\omega\|_\infty$ and Z_ω are small when $\gamma_{\theta\omega}$ is close to 1. To demonstrate that let us assume that

$$Z_\omega \leq r_\omega(\gamma_{\theta\omega}^\alpha - 1)$$

for some $r_\omega < 1 - \varepsilon$, $\varepsilon \in (0, 1)$. Then by replacing Z_ω with the above upper bound and then using some elementary estimates we see that

$$E_\omega \leq C(\|u_\omega - \mu_\omega(u_\omega)\|_\infty + r_\omega) e^{2\gamma_\omega^\alpha} \gamma_{\theta\omega}^\alpha (\gamma_{\theta\omega}^\alpha - 1)^{-1}.$$

We thus see that E_ω is bounded if $\|u_\omega\|_\infty + r_\omega$ is small enough (fiberwise). For instance, when γ_ω is bounded above we get the sufficient condition

$$\|u_\omega - \mu_\omega(u_\omega)\|_\infty + r_\omega \leq C(\gamma_{\theta\omega}^\alpha - 1)$$

which means that $\|u_\omega - \mu_\omega(u_\omega)\|_\infty + r_\omega$ is small when $T_{\theta\omega}$ has a local inverse branch with a close to 1 amount of contraction $\gamma_{\theta\omega}^{-1}$.

In the setup of Section 2.2, the random variable E_ω is bounded when $\|\phi_\omega\|_\infty, H_\omega$ and $\|u_\omega\|$ are (fiberwise) small enough when ζ_ω is close to 1.

39. Theorem. *Under Assumption 37 we have the following. Assume that $\sigma > 0$ and that for some $p > 4$ we have that $\bar{D}_\omega \in L^{2p}(\Omega, \mathcal{F}, \mathbb{P})$ and $K_\omega, M_\omega, \|\tilde{u}_\omega\|_\infty \in L^p(\Omega, \mathcal{F}, \mathbb{P})$, where $\tilde{u}_\omega = u_\omega - \mu_\omega(u_\omega)$. Moreover, suppose that for some measurable set $A \subset \Omega$ with positive probability we have:*

(i) *A satisfies the approximation properties described in Assumption 20;*

(ii) *the random variable $\max(M_\omega, K_\omega, B_\omega)$ is bounded on A;*

(iii) *A satisfies the assumptions of Theorem 27 with $\frac{p-1}{p} = 1 - 1/p$ instead of $1 - 2/p$ (let us denote the corresponding conditions by $(M1')$, $(M2')$ and $(M3')$, respectively).*

Then the following moderate deviations principle holds true for \mathbb{P} -a.a. ω : for every balanced²⁹ sequence (a_n) so that $\frac{a_n}{\sqrt{n}} \rightarrow \infty$, and $a_n = o(n^{1-6/p})$ and all Borel measurable sets $\Gamma \subset \mathbb{R}$ we have

$$(3.5) \quad - \inf_{x \in \Gamma^o} \frac{1}{2} x^2 \sigma^{-2} \leq \liminf_{n \rightarrow \infty} \frac{1}{a_n^2/n} \ln \mathbb{P}(S_n^\omega \tilde{u}/a_n \in \Gamma) \leq \limsup_{n \rightarrow \infty} \frac{1}{a_n^2/n} \ln \mathbb{P}(S_n^\omega \tilde{u}/a_n \in \Gamma) \leq - \inf_{x \in \bar{\Gamma}} \frac{1}{2} x^2 \sigma^{-2}$$

²⁹We say that a sequence (a_n) of positive numbers is balanced if $\frac{a_n}{a_{c_n}} \rightarrow 1$ for every sequence (c_n) so that $c_n \rightarrow 1$.

where Γ° is the interior of Γ and $\bar{\Gamma}$ is its closure.

Note that as an example of a sequence a_n we can take $a_n = n^q(\ln n)^\theta$ for $\theta \geq 0$ and $\frac{1}{2} < q < \min(1, 2 - p/6)$. As in Example 23 the approximation conditions and (M1')-(M3') hold true when B_ω, M_ω and K_ω can be approximated sufficiently fast by functions of $X_j, |j| \leq r$ and that the upper mixing coefficients of the sequence (X_j) satisfy (M1')-(M3').

The following result provides alternative conditions for the MPD.

40. Theorem. *Under Assumption 37 we have the following. Let the same integrability conditions in Theorem 39 hold with some $p > 8$ and suppose again that $\sigma > 0$. Then the MDP (3.5) holds true with any sequence (a_n) so that $a_n n^{-\max(6/p, 1/2)} \rightarrow \infty$ and $a_n = o(n^{1-8/p})$.*

The proof of Theorems 39 and 40 appear in Section 4.4.

41. Remark. Recall that M_ω is bounded in the setup of Section 2.1, and so the condition $M_\omega \in L^p$ is not really a restriction in that setup. Moreover, as explained in Remark 12 (see also Remark 14) when H_ω is also bounded then the random variables K_ω and B_ω are bounded. In this case also the condition $K_\omega, B_\omega \in L^p$ is not really a restriction, and the only real integrability condition is $\bar{D}_\omega \in L^{2p}$.

42. Remark. The main difference between Theorems 39 and 40 is that Theorem 39 essentially requires some mixing assumptions on the sequences of random variables $(B_{\theta^j \omega}), (M_{\theta^j \omega})$ and $(K_{\theta^j \omega})$, while Theorem 40 does not require mixing assumptions. On the other hand, the integrability conditions in Theorem 39 are weaker than the ones in Theorem 40 (i.e. $p > 4$ versus $p > 8$). Since the first integrability conditions are not much better than the second, Theorem 40 is somehow better than Theorem 39, and the reason that Theorem 39 is included is that its proof is based on a certain inducing strategy, and we find it interesting to present exact conditions which make the method of proof by inducing effective for proving an MDP for random Birkhoff sums.

3.7. Berry Esseen type estimates and moderate local limit theorem. Using the arguments in the proof of Theorems 39 and 40 we can also prove the following results.

43. Theorem (A Berry-Esseen theorem). *Let $\sigma_{\omega, n} = \sqrt{\text{Var}_{\mu_\omega}(S_n^\omega u)}$.*

(i) *Under the assumptions of Theorem 39, when $p > 12$ then \mathbb{P} -a.s. we have*

$$\sup_{t \in \mathbb{R}} |\mu_\omega(S_n^\omega \tilde{u} \leq t \sigma_{\omega, n}) - \Phi(t)| = O(n^{-(1/2-6/p)})$$

where when $p = \infty$ we use the convention $6/p = 0$.

(ii) *Under the assumptions of Theorem 40, when $p > 16$ then \mathbb{P} -a.s. we have*

$$\sup_{t \in \mathbb{R}} |\mu_\omega(S_n^\omega \tilde{u} \leq t \sigma_{\omega, n}) - \Phi(t)| = O(n^{-(1/2-8/p)})$$

where for $p = \infty$ we have $8/p := 0$.

The proof of Theorem 43 appears in Section ??.

44. Remark. In the setup of Section 2.1, the uniformly random case (i.e. $p = \infty$) was covered in [33, Theorem 7.1.1], see also [25, 34] for optimal rates for different types of random maps. However, in the setup of Section 2.2, Theorem 43 is new even in the uniformly random case (so we get the optimal CLT rates $O(n^{-1/2})$ in that case). In the deterministic case when the maps T_ω and the functions u_ω do not depend on ω the arguments in the proof of Theorem 43 provide explicit constants in the Berry-Esseen theorem (similarly to [28, Theorem 1.1]).

45. Theorem (A moderate local central limit theorem). *Under the assumptions of Theorem 40, \mathbb{P} -a.s. we have the following. Let (a_n) be a sequence so that $a_n n^{-2/p} \rightarrow \infty$ and $a_n n^{-1/2} \rightarrow 0$ (where p comes from Theorem 40). Then for every continuous function $g : \mathbb{R} \rightarrow \mathbb{R}$ with a compact support or an indicator function of a bounded interval we have*

$$(3.6) \quad \sup_{v \in \mathbb{R}} \left| \sqrt{2\pi\kappa_{\omega, n}\mu_\omega}(g(S_n^\omega \tilde{u}/a_n - v)) - \left(\int g(y) dy \right) e^{\frac{-v^2}{2\kappa_{\omega, n}}} \right| = o(1)$$

where $\kappa_{\omega,n} = \sigma_{\omega,n}/a_n$. In particular, for every bounded interval I we have

$$\sup_{v \in \mathbb{R}} \left| \sqrt{2\pi\kappa_{\omega,n}} \mu_{\omega}(S_n^{\omega} \tilde{u} \in a_n(v+I)) - |I| e^{\frac{-v^2}{2\kappa_{\omega,n}}} \right| = o(1).$$

The proof of Theorem 45 appears in Section 4.6.

46. Remark. The classical local central limit theorem (LCLT) corresponds to the case when $a_n = 1$, which is excluded in Theorem 45 even when $p = \infty$, where the first requirement on a_n becomes $a_n \rightarrow \infty$. The case $p = \infty$ corresponds to the uniformly random case, and we refer to [33, Ch. 6] for sufficient conditions³⁰ for the validity of (3.6) with $a_n = 1$ in the uniformly random version of the setup of Section 2.1. Relying on Theorem 49 below, the classical LCLT can be obtained in the uniformly random version of the maps considered in Section 2.2 when $l_{\omega} \leq 1$, since in that case we have $\|\mathcal{L}_{\omega}^{it,n}\| \leq C(1+|t|)$ for some $C \geq 1$ and all $t \in \mathbb{R}$, where $\mathcal{L}_{\omega}^{it,n}$ is defined before Theorem 49. Finally, note that Theorem 45 is new even in the uniformly random case, which is important especially when the known sufficient conditions for the classical LCLT fail due to a some type of periodicity exhibited by the random Birkhoff sums.

3.8. Key technical tools: real and complex random RPF theorems with effective rates.

For all the maps T_{ω} considered in Sections 2.1 and 2.2, and every complex number z we consider the random transfer operator $\mathcal{L}_{\omega}^{(z)}$ which maps functions on \mathcal{E}_{ω} to functions on $\mathcal{E}_{\theta\omega}$ according to the formula

$$(3.7) \quad \mathcal{L}_{\omega}^{(z)} g(x) = \sum_{y \in T_{\omega}^{-1}\{x\}} e^{\phi_{\omega}(y) + z\tilde{u}_{\omega}(y)} g(y) = \sum_i e^{\phi_{\omega}(y_{i,\omega}(x)) + z\tilde{u}_{\omega}(y_{i,\omega}(x))} g(y_{i,\omega}(x)).$$

Here $\tilde{u}_{\omega} = u_{\omega} - \mu_{\omega}(u_{\omega})$. We also set $\mathcal{L}_{\omega}^{(0)} = \mathcal{L}_{\omega}$. For each ω, n and z write

$$\mathcal{L}_{\omega}^{z,n} = \mathcal{L}_{\theta^{n-1}\omega}^{(z)} \circ \dots \circ \mathcal{L}_{\theta\omega}^{(z)} \circ \mathcal{L}_{\omega}^{(z)}.$$

It is clear that $\mathcal{L}_{\omega}^{(z)} \mathcal{H}_{\omega} \subset \mathcal{H}_{\theta\omega}$. We will denote by $(\mathcal{L}_{\omega}^{(z)})^*$ the appropriate dual operator. When $z = 0$ we denote $\mathcal{L}_{\omega}^n = \mathcal{L}_{\omega}^{0,n}$.

In [42] it was shown that in the setup of Section 2.1 there is a unique triplet $(\lambda_{\omega}, h_{\omega}, \nu_{\omega})$ consisting of a random variable $\lambda_{\omega} > 0$, a random positive function $h_{\omega} \in \mathcal{H}_{\omega}$ and a probability measure on \mathcal{E}_{ω} so that \mathbb{P} -a.s. we have $\nu_{\omega}(h_{\omega}) = 1$,

$$\mathcal{L}_{\omega} h_{\omega} = \lambda_{\omega} h_{\theta\omega} \quad \text{and} \quad (\mathcal{L}_{\omega})^* \nu_{\theta\omega} = \lambda_{\omega} \nu_{\omega}.$$

Moreover, with $\lambda_{\omega,n} = \prod_{j=0}^{n-1} \lambda_{\theta^j\omega}$, there is a constant $\delta \in (0, 1)$ and a random variable $C(\cdot)$ so that for every $g \in \mathcal{H}_{\omega}$ we have

$$\|(\lambda_{\omega,n})^{-1} \mathcal{L}_{\omega}^n g - \nu_{\omega}(g) h_{\theta^n\omega}\| \leq C(\theta^n\omega) \delta^n \|g\|.$$

The above result is often referred to as a random Ruelle-Perron-Frobenius (RPF) theorem. The random variable $C(\omega)$ can be expressed by means of a first hitting time to a certain set which can be defined by means of some ergodic average and a random variable which can be expressed as a series of known random variables. A similar result follows from [44] in the setup of Section 2.2 (the uniqueness is obtained under additional assumptions but the construction of the RPF triplets proceeds without the additional requirements).

One of the main tools in the proof of all the limit theorems in this paper is the following result, which is an effective version of the above RPF theorem.

47. Theorem (An effective RPF theorem). *The RPF triplets above $(\lambda_{\omega}, h_{\omega}, \nu_{\omega})$ satisfy the following (\mathbb{P} -a.s.):*

- (i) $h_{\omega} \in \mathcal{C}_{\omega}$ and $\|h_{\omega}\| \leq K_{\omega}$;
- (ii) $1 \leq \sup h_{\omega} \leq B_{\omega,1} \inf h_{\omega} \leq B_{\omega,1}$;

³⁰Which involve a period point of θ and some notion of an aperiodicity of the Birkhoff sums.

(iii) for every n and $g \in \mathcal{H}_\omega$,

$$(3.8) \quad \left\| \frac{\mathcal{L}_\omega^n g}{\lambda_{\omega,n}} - \nu_\omega(g) h_{\theta^n \omega} \right\| \leq 4K_{\theta^n \omega} \rho_{\omega,n}$$

where $\lambda_{\omega,n} = \prod_{j=0}^{n-1} \lambda_{\theta^j \omega}$ and $\rho_{\omega,n} = \prod_{j=0}^{n-1} \rho(\theta^j \omega)$;

(iv) let the probability measures μ_ω on \mathcal{E}_ω be given by $\mu_\omega = h_\omega \nu_\omega$. Then \mathbb{P} -a.s. we have $(T_\omega)_* \mu_\omega = \mu_{\theta \omega}$. Moreover, let L_ω be the operator given by $L_\omega g = \frac{\mathcal{L}_\omega g}{\lambda_\omega h_{\theta \omega}}$. Then for every Hölder continuous function g on \mathcal{E}_ω and all $n \geq 1$ we have

$$(3.9) \quad \|L_\omega^n g - \mu_\omega(g)\| \leq B_{\theta^n \omega} \rho_{\omega,n} \|g\|.$$

(v) [exponential decay of correlations] for every natural n and for all $g \in \mathcal{H}_\omega$ and $f \in \mathcal{H}_{\theta^n \omega}$ we have

$$(3.10) \quad |\mu_\omega(g \cdot (f \circ T_\omega^n)) - \mu_\omega(g) \mu_\omega(f \circ T_\omega^n)| \leq B_{\theta^n \omega} \rho_{\omega,n} \|g\| \|f\|_{L^1(\mu_{\theta^n \omega})}.$$

The proof of Theorem 47 appears in Section 5.

48. Remark. Notice that $\lambda_\omega = \nu_{\theta \omega}(\mathcal{L}_\omega \mathbf{1})$, where $\mathbf{1}$ is the function which takes the constant value 1. Hence,

$$(3.11) \quad e^{-\|\phi_\omega\|_\infty} \leq \inf \mathcal{L}_\omega \mathbf{1} \leq \lambda_\omega \leq \sup \mathcal{L}_\omega \mathbf{1}.$$

In the finite degree case $|\mathcal{L}_\omega \mathbf{1}| \leq \deg(T_\omega) e^{\|\phi_\omega\|_\infty}$, while if the degree is not bounded then Assumption 8 insures that $\mathcal{L}_\omega \mathbf{1}$ is a bounded function (recall that for piecewise affine maps we always have $\mathcal{L}_\omega \mathbf{1} = \mathbf{1}$).

In the proof of Theorems 39, 40, 43 and 45 we will also need the following complex version of Theorem 47.

49. Theorem. When the random variable E_ω is bounded we have the following. There is a positive number $r_0 > 0$ so that for any complex number z such that $|z| \leq r_0$ there exist measurable families $\lambda_\omega(z)$, $h_\omega^{(z)}$ and $\nu_\omega^{(z)}$ which are analytic in z , consisting of a nonzero complex number $\lambda_\omega(z)$, a complex function $h_\omega^{(z)} \in \mathcal{H}_\omega$ and a complex continuous linear functional $\nu_\omega^{(z)} \in \mathcal{H}_\omega^*$ such that:

(i) We have

$$(3.12) \quad \mathcal{L}_\omega^{(z)} h_\omega^{(z)} = \lambda_\omega(z) h_{\theta \omega}^{(z)}, \quad (\mathcal{L}_\omega^{(z)})^* \nu_\omega^{(z)} = \lambda_\omega(z) \nu_\omega^{(z)} \quad \text{and} \quad \nu_\omega^{(z)}(h_\omega^{(z)}) = \nu_\omega^{(z)}(\mathbf{1}) = 1.$$

Moreover, $h_\omega^{(0)} = h_\omega$, $\lambda_\omega(0) = \lambda_\omega$ and $\nu_\omega^{(0)} = \nu_\omega$.

(ii) We have $\|\nu_\omega^{(z)}\| \leq M_\omega$ and $h_\omega^{(z)} = \frac{\hat{h}_\omega^{(z)}}{\alpha_\omega(z)}$, where $\alpha_\omega(z) := \nu_\omega^{(z)}(\hat{h}_\omega^{(z)}) \neq 0$ for some analytic in z family of functions $\hat{h}_\omega^{(z)}$ so that $\|\hat{h}_\omega^{(z)}\| \leq 2\sqrt{2}K_\omega$ (note that $\alpha_\omega(0) = 1$ and $\|\alpha_\omega(z)\| \leq 2\sqrt{2}M_\omega K_\omega$).

(iii) Let $n_0(\omega)$ be the first time that $|\alpha_\omega(z)| \geq 2\sqrt{2}M_\omega K_\omega \rho_{\theta^{-n} \omega, n}$. Then for every $n \geq n_0(\omega)$ and all $g \in \mathcal{H}_{\theta^{-n} \omega}$ we have

$$(3.13) \quad \left\| \frac{\mathcal{L}_{\theta^{-n} \omega}^{z,n} g}{\lambda_{\theta^{-n} \omega, n}(z)} - \nu_{\theta^{-n} \omega}^{(z)}(g) h_\omega^{(z)} \right\| \leq 8M_{\theta^{-n} \omega} K_\omega \left(|\alpha_\omega(z)|^{-1} + \frac{M_\omega K_\omega}{|\alpha_\omega(z)|^2} \right) \|g\| \tilde{\rho}_{\theta^{-n} \omega, n} := \mathcal{R}(\omega, n, z)$$

where $\lambda_{\omega, n}(z) = \lambda_\omega(z) \cdot \lambda_{\theta \omega}(z) \cdots \lambda_{\theta^{n-1} \omega}(z)$ (recall that M_ω is bounded in the setup of Section 2.1).

(iv) Let the operators $L_\omega^{(z)}$ be given by

$$L_\omega^{(z)} g = L_\omega(e^{zu_\omega} g) = \frac{\mathcal{L}_\omega^{(z)}(g h_\omega)}{\lambda_\omega h_{\theta \omega}}$$

and set $\bar{\lambda}_\omega(z) = \frac{\lambda_\omega(z)}{\lambda_\omega}$, $\bar{h}_\omega(z) = \frac{h_\omega^{(z)}}{h_\omega}$ and $\bar{\nu}_\omega^{(z)} = h_\omega \cdot \nu_\omega^{(z)}$. Then for all $n \geq n_0(\omega)$,

$$(3.14) \quad \left\| \frac{L_{\theta^{-n} \omega}^{z,n} g}{\bar{\lambda}_{\theta^{-n} \omega, n}(z)} - \bar{\nu}_{\theta^{-n} \omega}^{(z)}(g) \bar{h}_\omega^{(z)} \right\| \leq 6U_\omega \mathcal{R}(\omega, n, z).$$

The proof of Theorem 49 appears in Section 6.

50. **Remark.** Since $z \rightarrow |\alpha_\omega(z)|$ is continuous and positive, we have $\beta_\omega = \sup_{|z| \leq r_0} |\alpha_\omega(z)|^{-1} < \infty$ and so, in principle, we can use that to get an upper bound which does not depend on z . However, β_ω does not have an explicit form. Instead, in the proof of the large deviations theorems (Theorems 39 and 40) we will use that $|\alpha_\omega(z) - 1| \leq C|z|K_\omega M_\omega$ and that, when $K_\omega M_\omega \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ we have $K_{\theta^{n_\omega} \omega} M_{\theta^{n_\omega} \omega} = o(n^{2/p})$, which will produce effective bounds when $|z| = O(n^{-2/p})$.

51. **Remark.** Theorem 49 was proven in [33, Ch.5] in the uniformly random version of the setup in Section 2.1. However, in the setup of Section 2.2 Theorem 49 is new even in the uniformly random case (i.e. when $s_\omega \leq s < 1$ for some constant s and all the other random variables are bounded). In fact, it is new even in the deterministic case when $T_\omega = T, \phi_\omega = \phi$ and $u_\omega = u$ do not depend on ω . As mentioned in Section 1, Theorem 49 makes it possible to extend results like [28, Theorem 1.1] to the partially expanding case. Note that in the uniformly random case $\sup_{|z| \leq r_0} |\alpha_\omega(z)| \leq C$ for some constant $C > 0$, and so, since $\alpha_\omega(0) = 1$, by using the mean value theorem and the Cauchy integral formula and decreasing r_0 if needed, we have that $\frac{1}{2} \leq |\alpha_\omega(z)| \leq \frac{3}{2}$ (and so we can replace the term $|\alpha_\omega(z)|$ with a constant).

4. PROOFS OF THE LIMIT THEOREMS BASED ON THE RANDOM RPF THEOREMS

4.1. **The CLT and LIL: Proof of Theorem 27 by inducing.** The proof of Theorem 27 is based on an application of [39, Theorem 2.3] with the set $Q = A$, where A comes from Assumption 20. Let $c(\omega) = \|u_\omega - \mu_\omega(u_\omega)\|_{L^2(\mu_\omega)}$ and let $n_A(\omega)$ be the first hitting time to the set A . Set $\tilde{u}_\omega = u_\omega - \mu_\omega(u_\omega)$ and

$$(4.1) \quad \Psi_\omega = S_{n_A(\omega)}^\omega \tilde{u}_\omega = \sum_{j=0}^{n_A(\omega)-1} \tilde{u}_{\theta^j \omega} \circ T_\omega^j$$

and let $\Theta : A \rightarrow A$ be given by $\Theta(\omega) = \theta^{n_A(\omega)}(\omega)$. Let us also consider the maps $\mathcal{T}_\omega = T_\omega^{n_A(\omega)}$ and the corresponding transfer operators $\tilde{L}_\omega = L_\omega^{n_A(\omega)}$.

Then the conditions of [39, Theorem 2.3] are met if

$$(4.2) \quad \left\| \sum_{j=0}^{n_A(\omega)-1} c(\theta^j \omega) \right\|_{L^2(\mathbb{P})} < \infty.$$

$$(4.3) \quad \left\| \mathbb{I}(\omega \in A) \sum_{n=0}^{\infty} |\mathbb{E}_{\mu_\omega}[\Psi_\omega \cdot \Psi_{\Theta^n \omega} \circ \mathcal{T}_\omega^n]| \right\|_{L^1(\mathbb{P})} < \infty$$

and

$$(4.4) \quad \left\| \mathbb{I}(\omega \in A) \sum_{n=0}^{\infty} \mathbb{E}_{\mu_\omega}(|\tilde{L}_{\Theta^{-n} \omega}^n \Psi_{\Theta^{-n} \omega}|) \right\|_{L^2(\mathbb{P})} < \infty.$$

4.1.1. *Reduction to tails estimates of the first hitting times.*

52. **Lemma.** *All three conditions (4.2), (4.3) and (4.4) are valid if $c(\omega) \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ and*

$$(4.5) \quad \sum_{j=0}^{\infty} (\mathbb{P}(n_A > j))^{1-2/p} < \infty$$

for some $p > 2$.

Proof. Let us begin with showing that condition (4.2) is in force. Write

$$\sum_{j=0}^{n_A(\omega)-1} c(\theta^j \omega) = \sum_{j=0}^{\infty} c(\theta^j \omega) \mathbb{I}(n_A > j).$$

Then, by the Hölder inequality we have

$$\left\| \sum_{j=0}^{n_A(\omega)-1} c(\theta^j \omega) \right\|_{L^2(\mathbb{P})} \leq \sum_{j=0}^{\infty} \|(c \circ \theta^j) \mathbb{I}(n_A > j)\|_{L^2(\mathbb{P})} \leq \|c\|_{L^p(\mathbb{P})}^2 \sum_{j=0}^{\infty} (\mathbb{P}(n_A > j))^{1-2/p} < \infty.$$

Next, let us show that condition (4.3) is satisfied. First, let $k_n(\omega)$ be so that $\mathcal{T}_\omega^n = T_\omega^{k_n(\omega)}$. Then by using the definition of Ψ_ω and that $\{\mu_\omega\}$ is an equivariant family we see that

$$\mathbb{E}_{\mu_\omega} [\Psi_\omega \cdot \Psi_{\Theta^n \omega} \circ \mathcal{T}_\omega^n] = \sum_{j=0}^{n_A(\omega)-1} \mathbb{E}_{\mu_{\theta^j \omega}} [\tilde{u}_{\theta^j \omega} \cdot \Psi_{\Theta^n \omega} \circ T_{\theta^j \omega}^{k_n(\omega)-j}].$$

Now, by using (3.10), the properties of the set A and that $\Theta^n \omega \in A$ we see that

$$\left| \mathbb{E}_{\mu_{\theta^j \omega}} [\tilde{u}_{\theta^j \omega} \cdot \Psi_{\Theta^n \omega} \circ T_{\theta^j \omega}^{k_n(\omega)-j}] \right| \leq M(1-\varepsilon)^n \|\Psi_{\Theta^n \omega}\|_{L^1(\mu_{\Theta^n \omega})} \|\tilde{u}_{\theta^j \omega}\|$$

where we have used that there are n visits to A between $\theta^j \omega$ and $\Theta^n \omega$ for $j < n_A(\omega)$. Thus,

$$\|\mathbb{E}_{\mu_\omega} [\Psi_\omega \cdot \Psi_{\Theta^n \omega} \circ \mathcal{T}_\omega^n]\| \leq M \|\Psi_{\Theta^n \omega}\|_{L^1(\mu_{\Theta^n \omega})} (1-\varepsilon)^n \sum_{j=0}^{n_A(\omega)-1} \|\tilde{u}_{\theta^j \omega}\|.$$

Next, with $\omega_n = \Theta^n \omega$ we have

$$\|\Psi_{\Theta^n \omega}\|_{L^1(\mu_{\Theta^n \omega})} \leq \sum_{j=0}^{n_A(\omega_n)-1} \|\tilde{u}_{\theta^j \omega_n}\|_{L^1(\mu_{\theta^j \omega_n})} \leq \sum_{j=0}^{n_A(\omega_n)-1} \|\tilde{u}_{\theta^j \omega}\|.$$

Let

$$(4.6) \quad I(\omega) = \sum_{j=0}^{n_A(\omega)-1} \|\tilde{u}_{\theta^j \omega}\| = \sum_{j=0}^{n_A(\omega)-1} c(\theta^j \omega).$$

Then we conclude from the above estimates that

$$\left\| \mathbb{I}(\omega \in A) \sum_{n=0}^{\infty} \|\mathbb{E}_{\mu_\omega} [\Psi_\omega \cdot \Psi_{\Theta^n \omega} \circ \mathcal{T}_\omega^n]\| \right\|_{L^1(\mathbb{P})} \leq M \mathbb{E}[\mathbb{I}(\omega \in A) I(\omega) I(\Theta^n \omega)] \sum_{n=0}^{\infty} (1-\varepsilon)^n.$$

To complete the proof of (4.3) we notice that in the proof of (4.2) we showed that $I(\omega) \in L^2(\Omega, \mathcal{F}, \mathbb{P})$, which together with Cauchy-Schwarz inequality yields that

$$\mathbb{E}[\mathbb{I}(\omega \in A) I(\omega) I(\Theta^n \omega)] = \mathbb{E}[(I(\omega))(\mathbb{I}(\omega \in A) I(\Theta^n \omega))] \leq (P(A))^{-1} \mathbb{E}[I^2(\omega)]$$

where we have used that Θ preserves $\mathbb{P}_A = \mathbb{P}(\cdot|A)$.

Finally, let us verify condition (4.4). First, we have

$$\left\| \mathbb{I}(\omega \in A) \sum_{n=0}^{\infty} \mathbb{E}_{\mu_\omega} (|\tilde{L}_{\Theta^{-n} \omega}^n \Psi_{\Theta^{-n} \omega}|) \right\|_{L^2(\mathbb{P})} \leq \sum_{n=0}^{\infty} \left\| \mathbb{I}(\omega \in A) \mathbb{E}_{\mu_\omega} (|\tilde{L}_{\Theta^{-n} \omega}^n \Psi_{\Theta^{-n} \omega}|) \right\|_{L^2(\mathbb{P})}.$$

Second, since θ preserves \mathbb{P} and $\{\mu_\omega\}$ is an equivariant family for each n we have

$$\begin{aligned} \left\| \mathbb{I}(\omega \in A) \mathbb{E}_{\mu_\omega} (|\tilde{L}_{\Theta^{-n} \omega}^n \Psi_{\Theta^{-n} \omega}|) \right\|_{L^2(\mathbb{P})} &= \left\| \mathbb{I}(\Theta^n \omega \in A) \mathbb{E}_{\mu_{\Theta^n \omega}} (|\tilde{L}_\omega^n \Psi_\omega|) \right\|_{L^2(\mathbb{P})} \\ &\leq \sum_{j=0}^{n_A(\omega)-1} \left\| \mathbb{E}_{\mu_{\Theta^n \omega}} (|L_{\theta^j \omega}^{u_n(\omega)-j} \tilde{u}_{\theta^j \omega}|) \right\|_{L^2(\mathbb{P})} \end{aligned}$$

where $\Theta^n \omega = \theta^{u_n(\omega)} \omega$, and in the last inequality we have used (4.1). Now, since $\theta^{u_n} \omega \in A$ and there are exactly n returns to A between “times” j and u_n (since $j < n_A(\omega)$) we get from (3.9) that

$$\mathbb{E}_{\mu_{\Theta^n \omega}} (|L_{\theta^j \omega}^{u_n(\omega)-j} \tilde{u}_{\theta^j \omega}|) \leq M(1-\varepsilon)^n \|\tilde{u}_{\theta^j \omega}\|.$$

Thus,

$$\left\| \mathbb{I}(\omega \in A) \mathbb{E}_{\mu_\omega}(|\tilde{L}_{\Theta^{-n}\omega}^n \Psi_{\Theta^{-n}\omega}|) \right\|_{L^2(\mathbb{P})} \leq M(1-\varepsilon)^n I(\omega),$$

where $I(\omega)$ was defined in (4.6). Combining the above estimates we conclude that

$$\left\| \mathbb{I}(\omega \in A) \sum_{n=0}^{\infty} \mathbb{E}_{\mu_\omega}(|\tilde{L}_{\Theta^{-n}\omega}^n \Psi_{\Theta^{-n}\omega}|) \right\|_{L^2(\mathbb{P})} \leq \|I(\cdot)\|_{L^2(\mathbb{P})} M \sum_{n=0}^{\infty} (1-\varepsilon)^n < \infty$$

and the proof of (4.4) is completed. \square

4.1.2. *Tails estimates using upper mixing coefficient: proof of Theorem 27.* In this section we will show that condition (4.5) in Lemma 52 is valid under the assumptions of Theorem 27. This together with Lemma 52 and [39, Theorem 2.3] will complete the proof of Theorem 27.

Before we begin with obtaining upper bounds on the tail probabilities $\mathbb{P}(n_A > j)$, let us note that

$$(4.7) \quad \mathbb{P}(n_A > j) = \mathbb{P}\left(\bigcap_{k=1}^j \theta^{-k}(\Omega \setminus A)\right).$$

4.1.3. *Proof of Theorem 27 under Assumption (M1).* We first need the following result.

53. Lemma. *Let I_1, I_2, \dots, I_m , $m \geq 2$ be finite subsets of \mathbb{N} so that I_i is to the left of I_{i+1} and the gap between them is at least L for some $L > 0$. Let A_1, A_2, \dots, A_m be sets of the same probability $p = \mathbb{P}(A_i)$ so that A_i is measurable with respect to $\sigma\{X_j : j \in I_i\}$. Then*

$$\mathbb{P}\left(\bigcap_{i=1}^m A_i\right) \leq p^m + \alpha_P(L) \sum_{j=0}^{m-2} p^j \leq p^m + \alpha_P(L) \frac{1}{1-p}.$$

Proof. We will prove the lemma by induction on m . For $m = 2$ by the definition of $\alpha_U(\cdot)$ we have

$$\mathbb{P}(A_1 \cap A_2) \leq \mathbb{P}(A_1)\mathbb{P}(A_2) + \alpha_P(L)$$

which coincides with the desired upper bound for $m = 2$. Next, suppose that the lemma is true for some $m \in \mathbb{N}$ and let I_1, \dots, I_{m+1} be sets with minimal gap greater or equal to some L , and measurable sets A_1, \dots, A_{m+1} with the same probability p so that A_i is measurable with respect to $\sigma\{X_j : j \in I_i\}$. Then by the definition of $\alpha_U(\cdot)$ we have

$$\begin{aligned} \mathbb{P}\left(\bigcap_{i=1}^{m+1} A_i\right) &\leq \mathbb{P}\left(\bigcap_{i=1}^m A_i\right) \mathbb{P}(A_{m+1}) + \alpha_P(L) \\ &\leq \left(p^m + \alpha_P(L) \sum_{j=0}^{m-2} p^j\right) p + \alpha_P(L) = p^{m+1} + \alpha_P(L) \sum_{j=0}^{m-1} p^j = p^{m+1} + \alpha_P(L) \sum_{j=0}^{(m+1)-2} p^j \end{aligned}$$

where in the last inequality we have used the induction hypothesis with the sets A_1, \dots, A_m and that $\mathbb{P}(A_{m+1}) = p$. \square

54. **Corollary.** *Under Assumption 20, condition (4.5) holds true under the assumption that*

$$(4.8) \quad \sum_j (\ln j \beta_{Cj/(3 \ln j)})^{1-2/p} < \infty \quad \text{and} \quad \sum_j (\alpha_U(j/(3C \ln j)))^{1-2/p} < \infty$$

for some constant C so that $C |\ln(1 - \mathbb{P}(A)/2)| (1 - 2/p) > 1$.

Proof. First, for all integers $s \geq 1$ we have

$$\mathbb{P}(n_A > j) = \mathbb{P}\left(\bigcap_{k=1}^j \theta^{-k}(\Omega \setminus A)\right) \leq \mathbb{P}\left(\bigcap_{k=1}^{\lfloor j/s \rfloor} \theta^{-ks}(\Omega \setminus A)\right).$$

Now, let us take s of the form $s = 3r$ for $r \in \mathbb{N}$. Then

$$\mathbb{P} \left(\bigcap_{k=1}^{\lfloor j/s \rfloor} \theta^{-ks}(\Omega \setminus A) \right) \leq \mathbb{P} \left(\bigcap_{k=1}^{\lfloor j/s \rfloor} \theta^{-ks}(\Omega \setminus A_r) \right) + \lfloor j/s \rfloor \beta_r$$

where A_r and β_r come from Assumption 20. Thus,

$$(4.9) \quad \mathbb{P}(n_A > j) = \mathbb{P} \left(\bigcap_{k=1}^j \theta^{-k}(\Omega \setminus A) \right) \leq \mathbb{P} \left(\bigcap_{k=1}^{\lfloor j/s \rfloor} \theta^{-ks}(\Omega \setminus A_r) \right) + \lfloor j/s \rfloor \beta_r.$$

Next, by Lemma 53 we have

$$\mathbb{P} \left(\bigcap_{k=1}^{\lfloor j/s \rfloor} \theta^{-ks}(\Omega \setminus A_r) \right) \leq (1 - \mathbb{P}(A_r))^{\lfloor j/s \rfloor} + \frac{\alpha_U(r)}{1 - \mathbb{P}(A_r)}.$$

Next, let us take s of the form $s = s_j = C^{-1} \lfloor j / \ln j \rfloor$ for some $C > 0$. Using that $\lim_{r \rightarrow \infty} \mathbb{P}(A_r) = \mathbb{P}(A) > 0$ we get that for all j large enough we have $\frac{1}{2}(1 - \mathbb{P}(A)) \leq 1 - \mathbb{P}(A_r) \leq 1 - \frac{1}{2}\mathbb{P}(A)$. We thus see that for j large enough we have

$$\mathbb{P}(n_A > j) \leq \left(1 - \frac{1}{2}\mathbb{P}(A)\right)^{C \ln j} + \frac{2}{1 - \mathbb{P}(A)} \alpha_U(C \lfloor j / \ln j \rfloor) + \lfloor j/s \rfloor \beta_r.$$

Now let us take C so that $C \ln(1 - \frac{1}{2}\mathbb{P}(A)) > 1 - 2/p$. Then the series $\sum_j (1 - \frac{1}{2}\mathbb{P}(A))^{C(1-2/p) \ln j}$ converges and now the convergence of the series in (4.5) follows from (4.8). \square

Proof of Theorem 27 under Assumption (M1). By Corollary 54 condition (4.5) in Lemma 52 is valid. The proof of Theorem 27 in this case follows now by combining Lemma 52 and [39, Theorem 2.3]. \square

4.1.4. Proof Theorem 27 under Assumption (M2).

55. Lemma. *Let I_1, I_2, \dots, I_m , $m \geq 2$ be finite subsets of \mathbb{N} so that I_i is to the left of I_{i+1} and the gap between them is at least L for some $L > 0$. Let A_1, A_2, \dots, A_m be sets of the same probability $p = \mathbb{P}(A_i)$ so that A_i is measurable with respect to $\sigma\{X_j : j \in I_i\}$. Then*

$$\mathbb{P} \left(\bigcap_{i=1}^m A_i \right) \leq (p + \phi_U(L))^{m-1}.$$

Proof. We will prove the lemma by induction on m . For $m = 2$ the lemma follows from the definition of ϕ_U . Now, suppose that the lemma is true for some m . Let I_1, \dots, I_{m+1} be sets with minimal gap greater or equal to some L , and measurable sets A_1, \dots, A_{m+1} with the same probability p so that A_i is measurable with respect to $\sigma\{X_j : j \in A_i\}$. Then by the definition of ϕ_U we have

$$\mathbb{P} \left(\bigcap_{i=1}^m A_i \cap A_{m+1} \right) \leq \mathbb{P} \left(\bigcap_{i=1}^m A_i \right) (\mathbb{P}(A_{m+1}) + \phi_U(L))$$

and not the proof of the induction step is completed by using the induction hypothesis with the sets A_1, \dots, A_m . \square

56. Corollary. *Suppose that*

$$\limsup_{r \rightarrow \infty} \phi_U(r) < \mathbb{P}(A)$$

and that $\sum_j (\ln j \beta_{j/(3C \ln j)})^{1-2/p} < \infty$ for some C so that $C \ln \delta > 1 - p/2$, where $\delta = 1 - \mathbb{P}(A) + \limsup_{r \rightarrow \infty} \phi_U(r)$. Then the series on the left hand side of (4.5) converges.

Proof. As in the beginning of the proof of Corollary 54, for every $s \in \mathbb{N}$ of the form $s = 3r$ we have

$$\mathbb{P}(n_A > j) = \mathbb{P}\left(\bigcap_{k=1}^j \theta^{-k}(\Omega \setminus A)\right) \leq \mathbb{P}\left(\bigcap_{k=1}^{\lfloor j/s \rfloor} \theta^{-ks}(\Omega \setminus A_r)\right) + \lfloor j/s \rfloor \beta_r.$$

Now, by Lemma 55 we have

$$\mathbb{P}\left(\bigcap_{k=1}^{\lfloor j/s \rfloor} \theta^{-ks}(\Omega \setminus A_r)\right) \leq (1 - \mathbb{P}(A_r) + \phi_U(r))^{\lfloor j/s \rfloor}.$$

Next, since $\lim_{r \rightarrow \infty} \mathbb{P}(A_r) = \mathbb{P}(A)$ and $\limsup_{r \rightarrow \infty} \phi_U(r) < \mathbb{P}(A)$ we see that

$$\limsup_{r \rightarrow \infty} (1 - \mathbb{P}(A_r) + \phi_U(r)) = \delta < 1.$$

Thus, if we take s of the form $s = s_j = C^{-1} \lfloor j / \ln j \rfloor$, then for j large enough we have

$$\mathbb{P}(n_A > j) \leq \delta^{\lfloor j/s \rfloor} + \lfloor j/s \rfloor \beta_{\lfloor s/3 \rfloor}.$$

If we take C so that $C |\ln \delta| (1 - p/2) > 1$ we get that the series $\sum_j \delta^{j(1-2/p)/s_j}$ converges. Now the proof of the lemma is complete since the series $\sum_j (\lfloor j/s_j \rfloor \beta_{\lfloor s_j/3 \rfloor})^{1-2/p}$ converges by the assumptions of the lemma. \square

Proof of Theorem 27 under Assumption (M2). By Corollary 56 condition (4.5) in Lemma 52 is valid. Now the proof of Theorem 27 under (M2) follows by combining Lemma 52 and [39, Theorem 2.3]. \square

4.1.5. *Proof Theorem 27 under Assumption (M3).* We first need the following result.

57. Lemma. *Let I_1, I_2, \dots, I_m , $m \geq 2$ be finite subsets of \mathbb{N} so that I_i is to the left of I_{i+1} and the gap between them is at least L for some $L > 0$. Let A_1, A_2, \dots, A_m be sets so that A_i is measurable with respect to $\sigma\{X_j : j \in I_i\}$. Then*

$$\mathbb{P}\left(\bigcap_{i=1}^m A_i\right) \leq (1 + \psi_U(L))^{m-1} \prod_{j=1}^m \mathbb{P}(A_i).$$

Hence, if $\mathbb{P}(A_i) = p$ for all i and some p then

$$\mathbb{P}\left(\bigcap_{i=1}^m A_i\right) \leq p (p(1 + \psi_U(L)))^{m-1}.$$

Proof. The lemma follows directly by induction and the definition of ψ_U . \square

58. Corollary. *Suppose that $\limsup_{k \rightarrow \infty} \psi_U(k) < \frac{1}{1 - \mathbb{P}(A)} - 1$ and that $\sum_j (\ln j \beta_{j/(3C \ln j)})^{1-2/p} < \infty$ for some C so that $C |\ln \delta| (1 - 2/p) > 1$, where $\delta = (1 + \limsup_{r \rightarrow \infty} \psi_U(r)) (1 - \mathbb{P}(A)) < 1$. Then the series on the left hand side of (4.5) converges.*

Proof. As in the beginning of the proof of Corollary 54, for all $s = 3r$ we have

$$\mathbb{P}(n_A > j) = \mathbb{P}\left(\bigcap_{k=1}^j \theta^{-k}(\Omega \setminus A)\right) \leq \mathbb{P}\left(\bigcap_{k=1}^{\lfloor j/s \rfloor} \theta^{-ks}(\Omega \setminus A_r)\right) + \lfloor j/s \rfloor \beta_r.$$

Next, by applying Lemma 57, we have

$$\mathbb{P}\left(\bigcap_{k=1}^{\lfloor j/s \rfloor} \theta^{-ks}(\Omega \setminus A_r)\right) \leq (1 + \psi_U(r))^{\lfloor j/s \rfloor - 1} (1 - \mathbb{P}(A) + \beta_r)^{\lfloor j/s \rfloor} := q_{s,j}$$

where we have used that $|\mathbb{P}(A) - \mathbb{P}(A_r)| \leq \beta_r$. Now, let us take $s = s_j = C^{-1} \lfloor j / \ln j \rfloor$ for some $C > 0$. Then when j is large enough we see that

$$(1 + \psi_U(r)) (1 - \mathbb{P}(A) + \beta_r) \leq \delta + \varepsilon < 1$$

for some ε small enough, where $\delta = (1 + \limsup_{r \rightarrow \infty} \psi_U(r))(1 - \mathbb{P}(A)) < 1$. Thus, if also $C|\ln \delta|(1 - 2/p) > 1$, by taking a sufficiently small ε we get that both series $\sum_j (q_{s_j, j})^{1-2/p}$ and $\sum_j ([j/s_j] \beta_{[s_j/3]})^{1-2/p}$ converge, and the proof of the corollary is complete. \square

Proof of Theorem 27 under Assumption (M3). By the previous corollary condition (4.5) in Lemma 52 is valid. The proof of Theorem 27 in this case follows now by combining Lemma 52 and [39, Theorem 2.3]. \square

59. Remark. The proofs of Corollaries 54, 56 and 58 show that if A is measurable with respect to $\sigma\{X_j, |j| \leq d\}$ for some d then $\mathbb{P}(n_A > j)$ decays exponentially fast in j under the other assumptions of the corollaries (since we can take $\beta_r = 0$ if $r > d$).

4.2. A second approach to the CLT and LIL: a direct proof of Theorem 32. The idea in the proof of Theorem 32 is to verify the conditions of [39, Theorem 2.3] when $Q = \Omega$, namely when there is no actual inducing involved. This requires us to verify the following three conditions:

$$(4.10) \quad \|c(\omega)\|_{L^2(\mathbb{P})} < \infty, \quad c(\omega) = \|\tilde{u}_\omega\|$$

$$(4.11) \quad \left\| \sum_{n=0}^{\infty} \mathbb{E}_{\mu_\omega} [\tilde{u}_\omega \cdot \tilde{u}_{\theta^n \omega} \circ T_\omega^n] \right\|_{L^1(\mathbb{P})} < \infty$$

and

$$(4.12) \quad \left\| \sum_{n=0}^{\infty} \mathbb{E}_{\mu_\omega} (|L_{\theta^{-n}\omega}^n \tilde{u}_{\theta^{-n}\omega}|) \right\|_{L^2(\mathbb{P})} < \infty.$$

Next, recall our assumption about the existence of a sequence β_r so that $\beta_r \rightarrow 0$ and for every r there is a random variable $\rho_r(\omega)$ which is measurable with respect to $\sigma\{X_j : |j| \leq r\}$ so that

$$(4.13) \quad \|\rho - \rho_r\|_{L^\infty} \leq \beta_r.$$

The first condition (4.10) is a part of the assumptions of Theorem 32. In order to verify conditions (4.11) and (4.12) we first need the following result.

60. Lemma. *Let I_1, \dots, I_d be intervals in the positive integers so that I_j is to the left of I_{j+1} and the distance between them is at least L . Let Y_1, \dots, Y_d be nonnegative bounded random variables so that Y_i is measurable with respect to $\sigma\{X_k : k \in I_i\}$. Then*

$$\mathbb{E} \left[\prod_{i=1}^d Y_i \right] \leq (1 + \psi_U(L))^{d-1} \prod_{i=1}^d \mathbb{E}[Y_i].$$

Proof. Once we prove the lemma for $d = 2$ the general case will follow by induction. Let us assume that $d = 2$. Next, we have

$$Y_i = \lim_{n \rightarrow \infty} Y_i(n) = \lim_{n \rightarrow \infty} \sum_k \mathbb{I}((k-1)2^{-n} < Y_i \leq k2^{-n}) k 2^{-n}$$

and so with $\alpha_i(k, n) = \{(k-1)2^{-n} < Y_i \leq k2^{-n}\}$, by the monotone convergence theorem we have

$$\begin{aligned} \mathbb{E}[Y_1 Y_2] &= \lim_{n \rightarrow \infty} \mathbb{E}[Y_1(n) Y_2(n)] = \lim_{n \rightarrow \infty} \sum_{k_1, k_2} (2^{-n} k_1)(2^{-n} k_2) \mathbb{P}(\alpha_1(k_1, n) \cap \alpha_2(k_2, n)) \\ &\leq \lim_{n \rightarrow \infty} \sum_{k_1, k_2} (2^{-n} k_1)(2^{-n} k_2) (1 + \psi_U(L)) \mathbb{P}(\alpha_1(k_1, n)) \mathbb{P}(\alpha_2(k_2, n)) \\ &= (1 + \psi_U(L)) \lim_{n \rightarrow \infty} \mathbb{E}[Y_1(n)] \mathbb{E}[Y_2(n)] = (1 + \psi_U(L)) \mathbb{E}[Y_1] \mathbb{E}[Y_2] \end{aligned}$$

where in the above inequality we have used the definition of the upper mixing coefficients $\psi_U(\cdot)$. \square

Next, we need the following

61. **Lemma.** *Suppose that*

$$\lim_{s \rightarrow \infty} \Psi_U(s) < \infty$$

and that with some $\delta > 0$ we have $\|u_\omega\|, B_\omega \in L^{3+\delta}(\Omega, \mathcal{F}, \mathbb{P})$. Then conditions (4.11) and (4.12) are in force.

Proof of Theorem 32. The proof of Theorem 32 is completed now by combining Lemma 61 with [39, Theorem 2.3] in the case $Q = \Omega$. \square

Proof of Lemma 61. Since $0 < \rho(\cdot) < 1$ we have $\lim_{q \rightarrow \infty} \rho^q = 0$ and so by the monotone convergence theorem

$$\lim_{q \rightarrow \infty} \mathbb{E}_{\mathbb{P}}[\rho^q] = 0.$$

Thus, since the limit superior of ψ_U is finite, if q is large enough then we have

$$\limsup_{r \rightarrow \infty} \Psi_U(r) < \frac{1}{\mathbb{E}_{\mathbb{P}}[\rho^q]} - 1.$$

Let us take q large enough so that its conjugate exponent p satisfies $3p \leq 3 + \delta$, where δ comes from the assumptions of the lemma (and Theorem 32).

Next, to show that condition (4.11) is in force, let us fix some $n \geq 0$. We first note that by (3.10) we have

$$|\mathbb{E}_{\mu_\omega}[\tilde{u}_\omega \cdot \tilde{u}_{\theta^n \omega} \circ T_\omega^n]| \leq \|\tilde{u}_\omega\| \|\tilde{u}_{\theta^n \omega}\|_{L^1(\mu_{\theta^n \omega})} U(\theta^n \omega) \prod_{j=0}^{n-1} \rho(\theta^j \omega).$$

Next, by applying the generalized Hölder inequality with the exponents $q_1 = q_2 = q_3 = 3p$ and $q_4 = q$ we get that

$$\mathbb{E}_{\mathbb{P}} \left[\|\tilde{u}_\omega\| \|\tilde{u}_{\theta^n \omega}\|_{L^1(\mu_{\theta^n \omega})} U(\theta^n \omega) \prod_{j=0}^{n-1} \rho(\theta^j \omega) \right] \leq \|c(\cdot)\|_{3p}^2 \|U(\cdot)\|_{3p} \left(\mathbb{E}_{\mathbb{P}} \left[\prod_{j=0}^{n-1} \rho^q(\theta^j \omega) \right] \right)^{1/q}$$

where we have used that $\|\tilde{u}_{\theta^n \omega}\|_{L^1(\mu_{\theta^n \omega})} \leq c(\omega)$. Now, for all s of the form $s = 3r$ have

$$\begin{aligned} \mathbb{E}_{\mathbb{P}} \left[\prod_{j=0}^{n-1} \rho^q(\theta^j \omega) \right] &\leq \mathbb{E}_{\mathbb{P}} \left[\prod_{j=1}^{[(n-1)/s]} \rho^q(\theta^{js} \omega) \right] \leq \mathbb{E}_{\mathbb{P}} \left[\prod_{j=1}^{[(n-1)/s]} (\rho_r^q(\theta^{js} \omega) + C_q \beta_r) \right] \\ &\leq (1 + \psi_U(r))^{[(n-1)/s]-1} \prod_{j=1}^{[(n-1)/s]} (\mathbb{E}_{\mathbb{P}}[\rho_r^q] + C_q \beta_r) \end{aligned}$$

where in the last inequality we have used Lemma 60, and C_q is a constant that depends only on q . Since $\lim_{r \rightarrow \infty} \mathbb{E}_{\mathbb{P}}[\rho_r^q] = \mathbb{E}_{\mathbb{P}}[\rho^q]$, $\lim_{r \rightarrow \infty} \beta_r = 0$ and $(1 + \psi_U(r))\mathbb{E}_{\mathbb{P}}[\rho^q] < 1$, by fixing a sufficiently large $s = s_0$ we conclude that

$$\mathbb{E}_{\mathbb{P}}[|\mathbb{E}_{\mu_\omega}[\tilde{u}_\omega \cdot \tilde{u}_{\theta^n \omega} \circ T_\omega^n]|] \leq C(1 - \varepsilon)^n$$

for some constants $\varepsilon \in (0, 1)$ and $C > 0$ (which depend on s_0 and q), and thus Condition (4.11) is in force.

Next, in order to verify condition (4.12), by Theorem 47 we have

$$|L_{\theta^{-n} \omega}^n \tilde{u}_{\theta^{-n} \omega}| \leq B_\omega \|\tilde{u}_{\theta^{-n} \omega}\| \prod_{j=1}^n \rho(\theta^{-j} \omega)$$

and so by the Hölder inequality,

$$\|\mathbb{E}_{\mu_\omega}(|L_{\theta^{-n} \omega}^n \tilde{u}_{\theta^{-n} \omega}|)\|_{L^2(\mathbb{P})} \leq \|B_\omega\|_{L^{2p}(\mathbb{P})} \|\tilde{u}_\omega\|_{L^{2p}(\mathbb{P})} \left\| \prod_{j=1}^n \rho(\theta^{-j} \omega) \right\|_{L^q(\mathbb{P})}.$$

Due to stationarity we have

$$\left\| \prod_{j=1}^n \rho(\theta^{-j}\omega) \right\|_{L^q(\mathbb{P})} = \left\| \prod_{j=0}^{n-1} \rho(\theta^j\omega) \right\|_{L^q(\mathbb{P})} = \left(\mathbb{E}_{\mathbb{P}} \left[\prod_{j=0}^{n-1} \rho^q(\theta^j\omega) \right] \right)^{1/q} = O((1-\varepsilon)^n)$$

where the last estimates was obtained in the course of the proof of (4.11). This completes the proof of (4.12). \square

4.3. An almost sure invariance principle: proof of Theorem 34. Let β_r satisfy (4.13).

4.3.1. *Key auxiliary result.* Before proving Theorem 34 we need the following result.

62. Lemma. *Under the Assumptions of Theorem 34 we have the following.*

(i) Let $R_n(\omega) = \sum_{j=0}^{n-1} \rho(\theta^j\omega) \cdots \rho(\theta^{n-1}\omega)$. Then for every $p \in \mathbb{N}$,

$$(4.14) \quad \mathbb{E}_{\mathbb{P}}[R_n^p] \leq C_p$$

for some constant C_p which does not depend on n . Therefore, for every $\varepsilon > 0$ we have

$$R_n(\omega) = o(n^\varepsilon), \mathbb{P} - a.s.$$

(ii) For every pair of positive integers (n, m) such that $m \leq n$ let

$$R_{m,n}(\omega) = \sum_{k=m}^n \sum_{j=k}^n \rho(\theta^k\omega) \cdot \rho(\theta^{k+1}\omega) \cdots \rho(\theta^j\omega) = \sum_{m \leq k \leq j \leq n} \rho(\theta^k\omega) \cdots \rho(\theta^j\omega).$$

Then for every $\ell \in \mathbb{N}$, $\varepsilon > 0$ and a positive integer p we have

$$(4.15) \quad \mathbb{E}_{\mathbb{P}} \left[\sup_{(n,m): 0 \leq n-m \leq \ell} n^{-(1+\varepsilon)} R_{m,n}^p \right] \leq C_{p,\varepsilon} \ell^{1+p}$$

for some constant $C_{p,\varepsilon} > 0$ which depends only on p and ε . Therefore, \mathbb{P} -a.s. for every $\varepsilon > 0$, uniformly in n and m as $(n-m) \rightarrow \infty$ we have

$$n^{-\varepsilon} R_{m,n}(\omega) = O((n-m)^{1+\varepsilon}), \mathbb{P}\text{-a.s.}$$

Proof. (i) First, the almost sure estimate $R_n(\omega) = o(n^\varepsilon)$ follows from (4.14) and the Borel Cantelli Lemma. Indeed, by taking $p > \frac{1}{\varepsilon}$ and applying the Markov inequality we arrive at

$$\mathbb{P}(R_n \geq n^\varepsilon) = \mathbb{P}(R_n^p \geq n^{\varepsilon p}) \leq C_p n^{-p\varepsilon}.$$

In order to prove (4.14), let us take $s \in \mathbb{N}$ of the form $s = 3r$. Then, since $0 < \rho(\cdot) < 1$,

$$\begin{aligned} \mathbb{E}_{\mathbb{P}}[R_n^p] &= \sum_{0 \leq j_1 \leq j_2 \leq \dots \leq j_p < n} \mathbb{E}_{\mathbb{P}} \left[\prod_{k=1}^p \prod_{u=j_k}^{n-1} \rho(\theta^u\omega) \right] \leq \sum_{0 \leq j_1 \leq j_2 \leq \dots \leq j_p < n} \mathbb{E}_{\mathbb{P}} \left[\prod_{u=j_1}^{n-1} \rho(\theta^u\omega) \right] \\ &\leq \sum_{0 \leq j_1 \leq j_2 \leq \dots \leq j_p < n} \mathbb{E}_{\mathbb{P}} \left[\prod_{v=0}^{[(n-1-j_1)/s]} \rho(\theta^{j_1+sv}\omega) \right] \leq \sum_{0 \leq j_1 \leq j_2 \leq \dots \leq j_p < n} \mathbb{E}_{\mathbb{P}} \left[\prod_{v=0}^{[(n-1-j_1)/s]} (\rho_r(\theta^{j_1+sv}\omega) + \beta_r) \right] \\ &\leq \sum_{0 \leq j_1 \leq j_2 \leq \dots \leq j_p < n} (1 + \psi_U(r))^{[(n-j_1-1)/s]} \prod_{v=0}^{[(n-1-j_1)/s]} \mathbb{E}_{\mathbb{P}} [(\rho_r(\theta^{j_1+sv}\omega) + \beta_r)] \\ &= \sum_{0 \leq j_1 \leq j_2 \leq \dots \leq j_p < n} (1 + \psi_U(r))^{[(n-j_1-1)/s]} a_r^{[(n-1-j_1)/s]+1} \end{aligned}$$

where $a_r = \mathbb{E}[\rho_r] + \beta_r$ and in the last inequality we have used Lemma 60. Taking s large enough so that $a_r(1 + \psi_U(r)) = \delta < 1$ (using (3.4)) and using that $n - 1 - j_1 \geq n - 1 - j_i$ for $i = 1, 2, \dots, d$ we conclude that

$$\mathbb{E}_{\mathbb{P}}[R_n^p] \leq \sum_{0 \leq j_1 \leq j_2 \leq \dots \leq j_p < n} b^{\sum_{i=1}^d (n-1-j_i)} = \left(\sum_{j=0}^{n-1} b^{n-1-j} \right)^p \leq C_p(b) = \frac{1}{(1-b)^p}$$

where $b = b_{p,s} = \delta^{1/sp} \in (0, 1)$.

(ii) First, the almost sure estimate $R_{m,n}(\omega) = O(n^{1/p+\varepsilon}(n-m)^{1+\varepsilon})$ follows from (4.15) and the Borel Cantelli Lemma. Indeed, for all $A > 0$ we have

$$\mathbb{P} \left(\sup_{(n,m): 0 \leq n-m \leq \ell} n^{-(1+\varepsilon)} R_{m,n}^p \geq A^p \right) = O(\ell^{p+1} A^{-p})$$

and so for $A_\ell = \ell^{1+\frac{3}{p}}$ we have

$$\mathbb{P} \left(\sup_{(n,m): 0 \leq n-m \leq \ell} n^{-(1+\varepsilon)/p} R_{m,n} \geq A_\ell \right) \leq C \ell^{-2}.$$

Now, given $\varepsilon > 0$, by taking p large enough we conclude from the Borel Cantelli Lemma that

$$\sup_{(n,m): 0 \leq n-m \leq \ell} n^{-(1+\varepsilon)/p} R_{m,n} = O(\ell^{1+\varepsilon}), \quad \mathbb{P}\text{-a.s.}$$

Thus, \mathbb{P} -a.s. there is a constant C so that for a given n and m with $n - m$ large enough we have $R_{m,n} \leq C(n-m)^{1+\varepsilon} n^{1/p+\varepsilon/p}$. Finally, by taking p large enough we can also insure that $(1+\varepsilon)/p < \varepsilon$.

Next, in order to prove (4.15), we have

$$(4.16) \quad \begin{aligned} \mathbb{E}_{\mathbb{P}} \left[\sup_{(n,m): 0 \leq n-m \leq \ell} n^{-(1+\varepsilon)} R_{m,n}^p \right] &\leq \sum_{(n,m): 0 \leq n-m \leq \ell} n^{-(1+\varepsilon)} \mathbb{E}[R_{m,n}^p] = \sum_{n=1}^{\infty} n^{-(1+\varepsilon)} \sum_{m=n-\ell}^n \mathbb{E}[R_{m,n}^p] \\ &\leq \left(\sum_{n=1}^{\infty} n^{-(1+\varepsilon)} \right) (\ell + 1) \sup_{(n,m): n-\ell \leq m \leq n} \mathbb{E}[R_{m,n}^p] \leq C_\varepsilon \ell \sup_{(n,m): n-\ell \leq m \leq n} \mathbb{E}[R_{m,n}^p]. \end{aligned}$$

Next, let us estimate $\mathbb{E}[R_{m,n}^p]$ for a fixed pair of positive integers (n, m) so that $n - \ell \leq m \leq n$. We first write

$$\mathbb{E}_{\mathbb{P}}[R_{m,n}^p] = \sum_{m \leq k_1, \dots, k_p \leq n} \sum_{k_i \leq j_i \leq n; 1 \leq i \leq p} \mathbb{E}_{\mathbb{P}} \left[\prod_{i=1}^p \prod_{u=k_i}^{j_i} \rho(\theta^u \omega) \right].$$

For a fixed choice of pairs $(k_i, j_i), i = 1, 2, \dots, p$ let $a = a(\{(k_i, j_i) : 1 \leq i \leq p\})$ be an index so that $j_a - k_a$ is the largest among $j_i - k_i$. Since $0 < \rho(\omega) < 1$, by disregarding the products $\prod_{u=k_i}^{j_i} \rho(\theta^u \omega)$ for $i \neq a$ we see that

$$\mathbb{E}_{\mathbb{P}}[R_{m,n}^p] \leq \sum_{m \leq k_1, \dots, k_p \leq n} \sum_{k_i \leq j_i \leq n; 1 \leq i \leq p} \mathbb{E}_{\mathbb{P}} \left[\prod_{u=k_a}^{j_a} \rho(\theta^u \omega) \right].$$

Next, since $0 < \rho(\cdot) < 1$, for all s of the form $s = 3r$, by (4.13) and Lemma 60 we have

$$\begin{aligned} \mathbb{E}_{\mathbb{P}} \left[\prod_{u=k_a}^{j_a} \rho(\theta^u \omega) \right] &\leq \mathbb{E}_{\mathbb{P}} \left[\prod_{u=0}^{[(j_a - k_a)/s]} \rho(\theta^{k_a + su} \omega) \right] \\ &\leq \mathbb{E}_{\mathbb{P}} \left[\prod_{u=0}^{[(j_a - k_a)/s]} (\rho_r(\theta^{k_a + su} \omega) + \beta_r) \right] \leq (1 + \psi_U(r))^{[(j_a - k_a)/s]} (\mathbb{E}_{\mathbb{P}}[\rho_r] + \beta_r)^{[(j_a - k_a)/s] + 1}. \end{aligned}$$

Since $\limsup_{r \rightarrow \infty} (1 + \psi_U(r)) < \frac{1}{\mathbb{E}[\rho]}$ (by (3.4)), by fixing some $s = s_0$ large enough we get that $(1 + \psi_U(r))(\mathbb{E}\rho + \beta_r) = \delta < 1$. Thus, since $j_a - k_a$ is the maximal difference, we have

$$\mathbb{E}_{\mathbb{P}} \left[\prod_{u_a=k_a}^{j_a} \rho(\theta^{u_a} \omega) \right] \leq \delta^{\lfloor (j_a - k_a)/s \rfloor} \leq \varepsilon^{\sum_{i=1}^p (j_i - k_i)}$$

where $\varepsilon = \varepsilon_{p,s} = \delta^{\frac{1}{ps}} < 1$. Hence for all n, m so that $n - \ell \leq m \leq n$ we have

$$\mathbb{E}[R_{m,n}^p] \leq \sum_{m \leq k_1, \dots, k_p \leq n} \sum_{k_i \leq j_i \leq n; 1 \leq i \leq p} \varepsilon^{\sum_{i=1}^p (j_i - k_i)} = \left(\sum_{i=1}^p \sum_{k_i=m}^n \sum_{j_i=k_i}^{j_i} \varepsilon^{j_i - k_i} \right)^p = O((m - n)^p) = O(\ell^p)$$

which together with (4.16) completes the proof of the maximal moment estimates in (ii). \square

4.3.2. A martingale co-boundary representation. Let $\tilde{u}_\omega = u_\omega - \mu_\omega(u_\omega)$. Set

$$G_{\omega,n} = \sum_{j=0}^{n-1} L_{\theta^j \omega}^{n-j}(\tilde{u}_{\theta^j \omega})$$

and

$$M_{\omega,n} = \tilde{u}_{\theta^n \omega} + G_{\omega,n} - G_{\omega,n+1} \circ T_{\theta^n \omega}.$$

Then for every fixed ω we have that $M_{\omega,n} \circ T_\omega^n$ is a reverse martingale difference with respect to the reverse filtration $\mathcal{T}_\omega^n = (T_\omega^n)^{-1}(\mathcal{B}_\omega)$, where \mathcal{B}_ω is the Borel σ -algebra on \mathcal{E}_ω (see [20, Proposition 2]).

63. Lemma. *If $\omega \rightarrow C_\omega$, $\omega \rightarrow U_\omega$, $\omega \rightarrow N(\omega)$ and $\omega \rightarrow \|\tilde{u}_\omega\|$ are in $L^p(\Omega, \mathcal{F}, \mathbb{P})$ for some p then for every $\varepsilon > 0$ for \mathbb{P} -a.e. ω we have*

$$\|M_{\omega,n}^2\| = O(n^{8/p+\varepsilon}).$$

Proof. First by Theorem 47,

$$(4.17) \quad \|G_{\omega,n}\| \leq U(\theta^n \omega) \sum_{j=0}^{n-1} \|\tilde{u}_{\theta^j \omega}\| \rho(\theta^j \omega) \cdots \rho(\theta^{n-1} \omega) \leq U(\theta^n \omega) u_n(\omega) R_n(\omega)$$

where $u_n(\omega) = \sup_{j \leq n} \|\tilde{u}_{\theta^j \omega}\|$ and R_n is defined in Lemma 62 (i). Therefore by the definition (3.3) of $N(\cdot)$ we have

$$\|G_{\omega,n+1} \circ T_{\theta^n \omega}\| \leq \|G_{n+1,\omega}\| N(\theta^n \omega) \leq U(\theta^{n+1} \omega) N(\theta^n \omega) u_{n+1}(\omega) R_{n+1}(\omega).$$

We thus conclude that

$$\|M_{\omega,n}^2\| \leq 3\|M_{\omega,n}\|^2 \leq A (U(\theta^n \omega) + U(\theta^{n+1} \omega))^2 (1 + N(\theta^n \omega))^2 u_{n+1}^2(\omega) (R_n(\omega) + R_{n+1}(\omega))^2$$

where A is an absolute constant. Now the lemma follows by Lemma 62 (i) together with the fact that for any random variable $Q(\omega)$, if $Q(\omega) \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ then $|Q(\theta^n \omega)| = o(n^{1/p})$, \mathbb{P} -almost surely (as a consequence of the mean ergodic theorem). \square

Next, we need the following quadratic variation estimates.

64. Lemma. *If $\omega \rightarrow C_\omega$, $\omega \rightarrow U_\omega$, $\omega \rightarrow N(\omega)$ and $\omega \rightarrow \|\tilde{u}_\omega\|$ are in $L^p(\Omega, \mathcal{F}, \mathbb{P})$ for some p then for every $\varepsilon > 0$, for \mathbb{P} -a.e. ω we have*

$$\sum_{k=0}^{n-1} \mathbb{E}_{\mu_\omega} [M_{\omega,k}^2 \circ T_\omega^k | \mathcal{T}_\omega^{k+1}] = \sum_{k=0}^{n-1} \mathbb{E}_{\mu_\omega} [M_{\omega,k}^2 \circ T_\omega^k] + o(n^{1/2+9/p+\varepsilon} \ln^{3/2+\varepsilon} n), \mu_\omega \text{ a.s.}$$

Proof. Set

$$A_{k,\omega} = \mathbb{E}_{\mu_\omega} [M_{\omega,k}^2 \circ T_\omega^k | \mathcal{T}_\omega^{k+1}], \quad B_{k,\omega} = \mathbb{E}_{\mu_\omega} [A_{k,\omega}] = \mathbb{E}_{\mu_\omega} [M_{\omega,k}^2 \circ T_\omega^k] = \mu_{\theta^k \omega}(M_{\omega,k}^2)$$

and $Y_{k,\omega} = A_{k,\omega} - B_{k,\omega}$. Then, by [26, Lemma 9] in order to prove the lemma it is enough to show that for all $n > m$ and all $\varepsilon > 0$ we have

$$(4.18) \quad \left\| \sum_{k=m}^n Y_{k,\omega} \right\|_{L^2(\mu_\omega)}^2 \leq C(n-m)n^{2/q+\varepsilon}$$

where C is a constant which may depend on ω and ε .

In order to prove (4.18), we first have

$$\left\| \sum_{k=m}^n Y_k \right\|_{L^2(\mu_\omega)}^2 = \text{Var} \left(\sum_{k=m}^n A_k \right) = \left\| \sum_{k=m}^n A_k \right\|_{L^2(\mu_\omega)}^2 - \left(\sum_{k=m}^n B_k \right)^2$$

where we abbreviate $A_{k,\omega} = A_k$ and $B_{k,\omega} = B_k$. Thus,

$$\left\| \sum_{k=m}^n Y_k \right\|_{L^2(\mu_\omega)}^2 \leq \sum_{k=m}^n \mu_\omega(A_k^2) + 2 \left(\sum_{m \leq i < j \leq n} \mu_\omega(A_i A_j) - \sum_{m \leq i < j \leq n} B_i B_j \right).$$

Next, arguing as in [20, Lemma 6] we have

$$A_i = L_{\theta^i \omega}(M_i^2) \circ T_\omega^{i+1}$$

where we abbreviate $M_i = M_{\omega,i}$. Hence, by also using that $(T_\omega^{i+1})_* \mu_\omega = \mu_{\theta^{i+1} \omega}$ and that L_ω is the dual of T_ω (w.r.t. μ_ω) we see that

$$\mu_\omega(A_i A_j) = \int L_{\theta^i \omega}(M_i^2) \cdot (L_{\theta^j \omega}(M_j^2) \circ T_{\theta^{i+1} \omega}^{j-i}) d\mu_{\theta^{i+1} \omega} = \int L_{\theta^i \omega}^{j-i+1}(M_i^2) \cdot L_{\theta^j \omega}(M_j^2) d\mu_{\theta^{j+1} \omega}.$$

Now, by (3.9) we have

$$\left\| L_{\theta^i \omega}^{j-i+1}(M_i^2) - \mu_{\theta^i \omega}(M_i^2) \right\| \leq U(\theta^j \omega) \|M_i^2\| \rho(\theta^i \omega) \cdots \rho(\theta^j \omega).$$

Since

$$\int L_{\theta^j \omega}(M_j^2) d\mu_{\theta^{j+1} \omega} = \mathbb{E}[A_j] = B_j$$

and $B_{\theta^j \omega} = o(j^{2/p})$ (as $B_\omega \in L^{2p}$) we conclude from the above estimates together with Lemma 63 that when $j > i$,

$$|\mu_\omega(A_i A_j) - B_i B_j| \leq B_{\theta^j \omega} \|M_i^2\| \|M_j^2\| \rho(\theta^i \omega) \cdots \rho(\theta^j \omega) = O(n^{18/p+\varepsilon}) \rho(\theta^i \omega) \cdots \rho(\theta^j \omega)$$

for every $\varepsilon > 0$. Thus,

$$\left| \sum_{m \leq i < j \leq n} \mu_\omega(A_i A_j) - \sum_{m \leq i < j \leq n} B_i B_j \right| \leq C n^{18/p+\varepsilon} \sum_{m \leq i < j \leq n} \rho(\theta^i \omega) \cdots \rho(\theta^j \omega) \leq C n^{18/p+\varepsilon} R_{n,m}(\omega).$$

Now the proof of the lemma is completed using Lemma 62 (ii). \square

4.3.3. *Proof of Theorem 34.* First, we have

$$S_n^\omega \tilde{u} = \sum_{j=0}^{n-1} M_{\omega,j} \circ T_\omega^j + G_{\omega,n} \circ T_\omega^n - G_{\omega,0}.$$

Next, by (4.17), Lemma 62 (i) and the assumption that $B_\omega \in L^{2p}(\Omega, \mathcal{F}, \mathbb{P})$ and $\|\tilde{u}_\omega\| \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ we see that for every $\varepsilon > 0$,

$$\|G_{\omega,n}\| = o(n^{3/p+\varepsilon}), \text{ a.s.}$$

and so

$$(4.19) \quad \left\| S_n^\omega \tilde{u} - \sum_{j=0}^{n-1} M_{\omega,j} \circ T_\omega^j \right\|_{L^\infty(\mu_\omega)} = O(n^{3/p+\varepsilon}).$$

In particular, with $\sigma_n^2 = \sigma_{n,\omega}^2 = \mathbb{E}_{\mu_\omega}[(S_n^\omega \tilde{u})^2]$ and ε is small enough we have

$$\sigma_n^2 := \left\| \sum_{j=0}^{n-1} M_{\omega,j} \circ T_\omega^j \right\|_{L^2(\mu_\omega)}^2 = \sum_{j=0}^{n-1} \mathbb{E}_{\mu_{\theta^j \omega}} [M_{\omega,j}^2] = \sigma_n^2 + O(n^{1/2+3/p+\varepsilon}) \asymp \sigma_n^2$$

where we have used that $\sigma_n^2/n \rightarrow \sigma^2 > 0$ and that $p > 3$.

In order to complete the proof we apply [17, Theorem 2.3] (taking into account [17, Remark 2.4]) with the reverse martingale difference $(M_{\omega,n} \circ T_\omega^n)$ and the sequence $a_n = n^{1/2+9/p+\varepsilon} \ln^{3/2+\varepsilon} n$, noticing that $\mathbb{E}_{\mu_{\theta^n \omega}} [M_{\omega,n}^2] = O(n^{8/p+\varepsilon})$ (by Lemma 63), and so when $p > 8$ we have $\mathbb{E}_{\mu_{\theta^n \omega}} [M_{\omega,n}^2] = O(\sigma_n^{2s})$ for some $0 < s < 1$. Taking into account Lemma 64, the first additional condition (i) of [17, Theorem 2.3] holds true. In order to verify the second additional condition (ii) with $v = 2$, for \mathbb{P} -a.a. ω we have

$$\sum_{n \geq 1} a_n^{-2} \mathbb{E}_{\theta^n \omega} [M_{\omega,n}^4] \leq C_\omega \sum_{n \geq 1} a_n^{-2} n^{16/p+\varepsilon}$$

which is a convergent series since $a_n^{-2} n^{16/p+\varepsilon} \leq n^{-1-2/p+2\varepsilon} = O(n^{-1-\delta})$, $\delta > 0$ (assuming that ε is small enough). In the above estimate we used that $\|M_{\omega,n}^4\| \leq 3\|M_{\omega,n}^2\|^2$ together with Lemma 63. We conclude that \mathbb{P} -a.s. there is a coupling of the reverse martingale $(M_{\omega,n} \circ T_\omega^n)$ with a Gaussian independent sequence (Z_n) , so that the ASIP rates in Theorem 34 hold true with $\sum_{j=0}^{n-1} M_{\omega,n} \circ T_\omega^j$ instead of $S_n^\omega u - \mu_\omega(S_n^\omega u) = S_n^\omega \tilde{u}$. Finally, in order to pass from the ASIP for the reverse martingale $(M_{\omega,n} \circ T_\omega^n)$ to the ASIP for the random Birkhoff sums $S_n^\omega u$ we use (4.19) and then the, so-called, Berkes-Philipp lemma (which allows us to further couple $u_{\theta^j \omega} \circ T_\omega^j$ with the Gaussian sequence). \square

4.4. Large deviations principle with quadratic rate function: proof of Theorems 39 and 40. In the circumstances of both Theorems 39 and 40, by the Gartner-Ellis theorem (see [18]) in order to prove the appropriate moderate deviations principle it is enough to show that for all real t we have

$$\lim_{n \rightarrow \infty} \frac{1}{s_n} \ln \mathbb{E}[e^{t a_n S_n^\omega \tilde{u}/n}] = \frac{1}{2} t^2 \sigma^2$$

where the sequence a_n is described in Theorems 39 and 40, $s_n = a_n^2/n$ and $\sigma^2 = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}_{\mu_\omega}(S_n^\omega u)$ (which does not depend on ω and is assumed to be positive). Henceforth we will assume that $\mu_\omega(u_\omega) = 0$, which means that we replace u_ω by $\tilde{u}_\omega = u_\omega - \mu_\omega(u_\omega)$.

4.4.1. Auxiliary estimates. We first need the following result.

65. Lemma. *Let $(\bar{\lambda}_\omega(z), \bar{h}_\omega^{(z)}, \bar{\nu}_\omega^{(z)})$ be the normalized RPF triplets from Theorem 49. There is a constant $r > 0$ so that \mathbb{P} -a.s. for every complex number z with $|z| \leq r$ we have*

$$\|\bar{\nu}_\omega^{(z)}\| \leq M_\omega K_\omega, \quad \|\bar{h}_\omega^{(z)}\| \leq \frac{2\sqrt{2}U_\omega K_\omega}{|\alpha_\omega(z)|}, \quad U_\omega = 6B_{\omega,1}^2 K_\omega$$

and

$$|\bar{\lambda}_\omega(z)| \leq 3e^{\|u_\omega\|_\infty} (1 + 2H_\omega) \|\mathcal{L}_\omega \mathbf{1}\|_\infty \leq \bar{D}_\omega$$

(where K_ω, M_ω and \bar{D}_ω are defined in Sections 2.3.1 and 2.3.2).

Proof. Using the upper and lower bounds in Theorem 47 together with (3.11) we see that

$$\|\bar{\nu}_\omega^{(z)}\| \leq M_\omega \|h_\omega\| \leq M_\omega K_\omega \quad \text{and} \quad \|\bar{h}_\omega^{(z)}\| \leq \frac{U_\omega \cdot (2\sqrt{2}K_\omega)}{|\alpha_\omega(z)|}$$

where we used that $v(1/h) \leq v(h)(\inf h)^{-2}$ for every positive function h (and so $\|1/h_\omega\| \leq U_\omega$). To bound $\bar{\lambda}_\omega(z)$, notice that $\bar{\lambda}_\omega(z) = \nu_{\theta^\omega}^{(z)}(\mathcal{L}_\omega^{(z)} \mathbf{1})$ which yields that

$$|\bar{\lambda}_\omega(z)| \leq M_{\theta^\omega} \|\mathcal{L}_\omega^{(z)} \mathbf{1}\| \leq M_{\theta^\omega} \|e^{zu_\omega}\| \|\mathcal{L}_\omega\|.$$

Firstly, let us bound the norm $\|\mathcal{L}_\omega\|$. Let g be a Hölder continuous function. Then $\|\mathcal{L}_\omega g\|_\infty \leq \|\mathcal{L}_\omega \mathbf{1}\|_\infty \|g\|_\infty$ (this part does not require continuity of g , only boundedness). Secondly, let us estimate

the Hölder constant of $\mathcal{L}_\omega g$. In the setup of Section 2.1 set $c_\omega = \gamma_\omega^{-1}$, while in the setup of Section 2.2 set $c_\omega = l_\omega$. Then for every two points $x, x' \in \mathcal{E}_{\theta_\omega}$ we have

$$\begin{aligned} |\mathcal{L}_\omega g(x) - \mathcal{L}_\omega g(x')| &\leq \sum_i e^{\phi_\omega(y_i)} |g(y_i) - g(y'_i)| + \sum_i |e^{\phi_\omega(y_i)} - e^{\phi_\omega(y'_i)}| |g(y'_i)| \\ &\leq v(g) c_\omega^\alpha \rho^\alpha(x, x') \|\mathcal{L}_\omega \mathbf{1}\|_\infty + 2H_\omega \rho^\alpha(x, x') \|g\|_\infty \|\mathcal{L}_\omega \mathbf{1}\|_\infty \leq (c_\omega^\alpha + 2H_\omega) \|\mathcal{L}_\omega \mathbf{1}\|_\infty \|g\| \rho^\alpha(x, x'). \end{aligned}$$

In the second inequality we have also used that

$$|e^{\phi_\omega(y_i)} - e^{\phi_\omega(y'_i)}| \leq (e^{\phi_\omega(x_i)} + e^{\phi_\omega(y_i)}) H_\omega \rho^\alpha(x, y)$$

which is obtained using the mean value theorem and the definition of H_ω (in either (2.4) or (2.11)). Here $y_i = y_{i,\omega}(x)$ and $y'_i = y_{i,\omega}(x')$ are the inverse images of x and x' under T_ω , respectively. Combining the above estimates we see that

$$(4.20) \quad \|\mathcal{L}_\omega\| \leq (1 + H_\omega + c_\omega^\alpha) \|\mathcal{L}_\omega \mathbf{1}\|_\infty = \tilde{D}_\omega.$$

Finally, using also that $\lambda_\omega \geq e^{-\|\phi_\omega\|_\infty}$ we conclude that when $|z| \leq 1$ then

$$|\bar{\lambda}_\omega(z)| = \frac{|\lambda_\omega(z)|}{\lambda_\omega} \leq M_{\theta_\omega} e^{\|\phi_\omega\|_\infty} \|e^{z u_\omega}\| \tilde{D}_\omega \leq M_{\theta_\omega} e^{\|\phi_\omega\|_\infty} e^{\|u_\omega\|_\infty} (1 + v(u_\omega)) \tilde{D}_\omega \leq \bar{D}_\omega.$$

□

66. Corollary. *There exist constants $r_1, C_1 > 0$ so that \mathbb{P} -a.s. if $|z| \leq r_1$ then*

$$|\bar{\lambda}_\omega(z) - 1| \leq C_1 |z| \bar{D}_\omega$$

and for every $r_2 \leq r_1$ if $|z| \leq r_2/2$ then with $\beta_\omega = \inf_{|z| \leq r_2} |\alpha_\omega(z)|$ we have

$$\|\bar{h}_\omega^{(z)} - \mathbf{1}\| \leq 2\sqrt{2} U_\omega K_\omega |z| \beta_\omega^{-1}$$

where U_ω was defined in Lemma 65.

Proof. Since $z \rightarrow \bar{\lambda}_\omega(z)$ and $z \rightarrow \bar{h}_\omega^{(z)}$ are analytic, the corollary follows from Lemma 65 together with the Cauchy integral formula. □

4.4.2. MDP via inducing: proof of Theorem 39. Let A be the set from the assumptions of Theorem 39. Then there is a constant Q so that for every $\omega \in A$ we have $\max(M_\omega, K_\omega, U_\omega) \leq Q$ (noting that $U_\omega \leq B_\omega$). Let n_A be the first visiting time to A . Then, using the upper bounds on $|\alpha_\omega(z)|$ from Theorem 49 and the Cauchy integral formula, we see that there is a constant $r_0 > 0$ so that if $|z| \leq r_0$ then for every $\omega \in A$ we have

$$|\alpha_\omega(z) - 1| \leq 2\sqrt{2} Q^2 |z| < \frac{1}{2}$$

and so

$$\beta_\omega = \min_{|z| \leq r_0} |\alpha_\omega(z)| \geq \frac{1}{2}.$$

Now, let n be so that $\theta^n \omega \in A$. Then if $|z| \leq r_0$ we have $|\alpha_{\theta^n \omega}(z)| \geq \frac{1}{2}$. On other hand, since K_ω and M_ω are in $L^p(\Omega, \mathcal{F}, \mathbb{P})$ we have $\max(K_{\theta^n \omega}, M_{\theta^n \omega}) = o(n^{1/p})$ (a.s.). Using also that $0 < \rho(\omega) < 1$ we see that $K_{\theta^n \omega} M_{\theta^n \omega} \rho_{\omega, n}$ decays to 0 exponentially fast. In particular, for every n large enough he have

$$\beta_\omega \geq 2\sqrt{2} K_{\theta^n \omega} M_{\theta^n \omega} \rho_{\omega, n}.$$

Hence, by applying (3.14) with $\theta^n \omega$ instead of ω we see that if $\theta^n \omega \in A$ and n is large enough then

$$(4.21) \quad \left\| \frac{L_\omega^{z, n} g}{\bar{\lambda}_{\omega, n}(z)} - \bar{v}_\omega^{(z)}(g) \bar{h}_{\theta^n \omega}^{(z)} \right\| \leq C(Q) M_\omega \|g\| \rho_{\omega, n}$$

for every Hölder continuous function g , where $C(Q)$ is a constant that depends on Q , but not on ω or n .

Next, notice that under the Assumptions of Theorem 39 we have that \bar{D}_ω from Lemma 65 belongs to L^{2p} . Hence $\bar{D}_{\theta^j \omega} = o(|j|^{2/p})$ and by Corollary (66) we have

$$|\bar{\lambda}_{\theta^j \omega}(z) - 1| \leq C |j|^{2/p}.$$

Thus there are uniformly bounded analytic branches (vanishing at the origin) of $\ln \bar{\lambda}_{\theta^j \omega}(z)$ for $j \leq n$ on any domain of the form $|z| = o(n^{-2/p})$. Let us denote these branches by $\Pi_{\theta^j \omega}(z)$.

Now, when $\theta^n \omega \in A$ then by Corollary 66 when $|z|$ is small enough for \mathbb{P} -a.a. ω we have

$$(4.22) \quad \|\bar{h}_{\theta^n \omega}^{(z)} - \mathbf{1}\| \leq \frac{1}{2}.$$

and so

$$\frac{1}{2} \leq |\mu_\omega(\bar{h}_\omega^{(z)})| \leq \frac{3}{2}$$

Therefore we can also develop uniformly bounded branches of $\ln \mu_\omega(\bar{h}_\omega^{(z)})$ around the complex origin which vanishes at the origin.

Next, by using (4.21) we see that for n large enough, if $\theta^n \omega \in A$ and $|z| = O(n^{-2/p})$ then

$$(4.23) \quad \mathbb{E}[e^{z S_n^\omega \tilde{u}}] = \mu_{\theta^n \omega}(L_\omega^{z, n} \mathbf{1}) = \bar{\lambda}_{\omega, n}(z) \left(\mu_{\theta^n \omega}(\bar{h}_{\theta^n \omega}^{(z)}) + O(\rho_{\omega, n} z) \right).$$

Since $\rho_{\omega, n}$ decays exponentially fast to 0 and $|\mu_{\theta^n \omega}(\bar{h}_{\theta^n \omega}^{(z)}) - 1| \leq \frac{1}{2}|z|$ (by (4.22)) by taking the logarithms of both sides and using analyticity (and the Cauchy integral formula) we see that when $\theta^n \omega \in A$ and $|z| = O(n^{-2/p})$ and n is large enough we have

$$\ln \mathbb{E}[e^{z S_n^\omega \tilde{u}}] = \Pi_{\omega, n}(z) + O(|z|) + O(\delta^n)$$

where $\Pi_{\omega, n}(z) = \sum_{j=0}^{n-1} \Pi_{\theta^j \omega}(z)$ and $\delta = \delta_\omega \in (0, 1)$, and we have used that $\ln(1+w) = O(w)$ when $|w|$ is small enough. By taking the derivatives at $z = 0$ and using the Cauchy integral formula on domains of the form $|z| = O(n^{-2/p})$ we see that

$$0 = \mathbb{E}[S_n^\omega \tilde{u}] = \Pi'_{\omega, n}(0) + O(n^{2/p})$$

and

$$\sigma_{\omega, n}^2 = \mathbb{E}[(S_n^\omega \tilde{u})^2] = \Pi''_{\omega, n}(0) + O(n^{4/p}).$$

Moreover, since $|\Pi_{\omega, n}(z)| = O(n)$, by using the Cauchy integral formula to estimate the error term in the second order Taylor expansion of $\Pi_{\omega, n}(z)$ around $z = 0$ we see that when $|z| = O(n^{-2/p})$ then

$$\Pi_{\omega, n}(z) = z \Pi'_{\omega, n}(0) + \frac{1}{2} z^2 \Pi''_{\omega, n}(0) + O(|z|^3) n^{1+6/p}$$

and so

$$\ln \mathbb{E}[e^{z S_n^\omega \tilde{u}}] = O(n^{2/p}) z + \frac{1}{2} z^2 \sigma_{\omega, n}^2 + O(n^{4/p}) z^2 + O(|z|^3) n^{1+6/p} + O(|z|) + O(\delta^n).$$

Let us now fix some $t \in \mathbb{R}$ and take $z = t_n = i t b_n / n$, where b_n satisfies $b_n \gg n^{4/p}$ and $\frac{b_n}{n} = o(n^{-6/p})$. Then, since $p > 4$,

$$\frac{\ln \mathbb{E}[e^{t(b_n/n) S_n^\omega \tilde{u}}]}{b_n^2/n} = o(1) + \frac{1}{2} t^2 (\sigma_{\omega, n}^2 / n), \quad \sigma = \lim_{n \rightarrow \infty} \frac{1}{n} \sigma_{\omega, n}^2.$$

Thus,

$$(4.24) \quad \lim_{n \rightarrow \infty, \theta^n \omega \in A} \frac{1}{b_n^2/n} \ln \mathbb{E}[e^{t_n S_n^\omega \tilde{u}}] = \frac{1}{2} t^2 \sigma^2.$$

The next step will be to use (4.24) to derive a similar result without the restriction $\theta^n \omega \in A$. Let (a_n) be a sequence with the properties described in Theorem 39. Let us take some n so that $\theta^n \omega \notin A$, and let $m = m_n = m_n(\omega)$ be the largest time $m \leq n$ so that $\theta^m \omega \in A$. Then

$$\left| \ln \mathbb{E}[e^{t a_n S_n^\omega \tilde{u}/n}] - \ln \mathbb{E}[e^{t a_n S_{m_n}^\omega \tilde{u}/n}] \right| \leq |t a_n / n| \cdot \left\| \sum_{j=m_n}^{n-1} \tilde{u}_{\theta^j \omega} \circ T_{\theta^{m_n} \omega}^{n-m_n} \right\|_\infty.$$

Now, if we set

$$\tilde{\Psi}_\omega = \sum_{j=0}^{n_A(\omega)-1} \|\tilde{u}_{\theta^j \omega}\|_\infty$$

then

$$\left\| \sum_{j=m_n}^{n-1} \tilde{u}_{\theta^j \omega} \circ T_{\theta^{m_n} \omega}^{n-m_n} \right\|_{\infty} \leq \tilde{\Psi}_{\theta^{m_n}(\omega) \omega}.$$

Observe now that with $c(\omega) = \|\tilde{u}_{\omega}\|_{\infty}$ we have

$$\tilde{\Psi}_{\omega} = \sum_{j=0}^{\infty} c(\theta^j \omega) \mathbb{I}(n_A(\omega) > j)$$

and so by the Hölder inequality, if q denotes the conjugate exponent of p then

$$\|\tilde{\Psi}_{\omega}\|_{L^p(\mathbb{P})} \leq \|c(\cdot)\|_{L^p(\mathbb{P})} \sum_{j=0}^{\infty} (\mathbb{P}(n_A > j))^{1/q}.$$

Arguing as in Section 4.1.2 we see that under each one of the conditions (M1'), (M2') or (M3') we have $\sum_{j=0}^{\infty} (\mathbb{P}(n_A > j))^{1/q} < \infty$. We thus see that $\|\tilde{\Psi}_{\omega}\|_{L^p(\mathbb{P})} < \infty$ and so $\tilde{\Psi}_{\theta^j \omega} = o(j^{1/p})$ almost surely. Thus, since $p > 2$ and $a_n/\sqrt{n} \rightarrow \infty$ we see that

$$\frac{|ta_n/n| \left\| \sum_{j=m_n}^{n-1} \tilde{u}_{\theta^j \omega} \circ T_{\theta^{m_n} \omega}^{n-m_n} \right\|_{\infty}}{s_n} = O(a_n^{-1} n^{1/p}) \rightarrow 0, \quad s_n = a_n^2/n.$$

Finally, since $m_n = n(1 + o(1))$ by the assumptions on the sequence (a_n) in Theorem 39 we have $a_n = a_{m_n}(1 + o(1))$ and $s_n = a_n^2/n = s_{m_n}(1 + o(1))$. Therefore by (4.24) we have

$$\lim_{n \rightarrow \infty} \frac{1}{s_n} \ln \mathbb{E}[e^{ta_n s_n \tilde{u}/n}] = \lim_{n \rightarrow \infty, \theta^n \in A} \frac{1}{s_n} \ln \mathbb{E}[e^{t_n s_n \tilde{u}}] = \frac{1}{2} t^2 \sigma^2$$

and the proof of Theorem 39 is complete. \square

4.4.3. *A direct approach to the MDP: proof of Theorem 40.* Recall that when $|z| \leq r_0$ (for some constant r_0) then $|\alpha_{\omega}(z)| \leq 2\sqrt{2}K_{\omega}M_{\omega}$. Now, using the Cauchy integral formula, when $|z| \leq r_0/2$ we have

$$|\alpha_{\omega}(z) - 1| \leq CK_{\omega}M_{\omega}|z|$$

where $C = C(r_0)$ is some constant. Since K_{ω} and M_{ω} are in $L^p(\Omega, \mathcal{F}, \mathbb{P})$ we have $K_{\theta^j \omega}M_{\theta^j \omega} = o(j^{2/p})$ and so when $|z| = O(n^{-2/p})$, then for every n large enough

$$(4.25) \quad |\alpha_{\theta^n \omega}(z) - 1| \leq \delta_n \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Thus, for such z 's when n is large enough so that $n^{2/p}\rho_{\omega,n} < 1/4$ we can apply (3.14) with $\theta^n \omega$ instead of ω and $|z| = O(n^{-2/p})$ and get that

$$(4.26) \quad \left\| \frac{L_{\omega}^{z,n} g}{\lambda_{\omega,n}(z)} - \bar{v}_{\omega}^{(z)}(g) \bar{h}_{\theta^n \omega}^{(z)} \right\| \leq CM_{\omega} \|g\| \delta_{\omega}^n$$

for some $\delta_{\omega} \in (0, 1)$, where we have used that $K_{\theta^n \omega}, M_{\theta^n \omega}$ and $U_{\theta^n \omega}$ grow at most polynomially fast and $\rho_{\omega,n}$ decays to 0 exponentially fast in n .

Next, by applying the Cauchy integral formula on a domain of the form $\{|z| = O(n^{-2/p})\}$ and using Lemma 65 to bound the derivative of $z \rightarrow \bar{h}_{\theta^n \omega}^{(z)}$ on such domains (taking into account (4.25) and that $U_{\theta^n \omega} K_{\theta^n \omega} = o(n^{2/p})$) we see that when $|z| = O(n^{-2/p})$ then

$$(4.27) \quad \|\bar{h}_{\theta^n \omega}^{(z)} - \mathbf{1}\| = |z|O(n^{4/p}).$$

Thus, we can develop a branch of $\ln \mu_{\theta^n \omega}(h_{\theta^n \omega}^{(z)})$ on a domain of the form $|z| = O(n^{-2/p})$ so that

$$(4.28) \quad \ln \mu_{\theta^n \omega}(h_{\theta^n \omega}^{(z)}) = 1 + |z|O(n^{4/p}).$$

Similarly, by the Assumptions of Theorem 40 we have that \bar{D}_{ω} defined in Lemma 65 belongs to $L^{2p}(\Omega, \mathcal{F}, \mathbb{P})$. Hence $\bar{D}_{\theta^j \omega} = o(|j|^{2/p})$ and by using Corollary (66) we see that

$$|\bar{\lambda}_{\theta^j \omega}(z) - 1| = o(j^{2/p})|z|.$$

Thus there are uniformly bounded branches (vanishing at the origin) of $\bar{\lambda}_{\theta^j \omega}(z)$ for $j \leq n$ on any domain of the form $|z| = O(n^{-2/p})$. Let us denote these branches by $\Pi_{\theta^j \omega}(z)$.

Next, by (4.26) we have

$$\mathbb{E}[e^{zS_n^\omega \tilde{u}}] = \mu_{\theta^n \omega}(L_\omega^n \mathbf{1}) = \bar{\lambda}_{\omega, n}(z) \left(\mu_{\theta^n \omega}(\bar{h}_{\theta^n \omega}^{(z)}) + O(\delta_\omega^n z) \right).$$

Using the above estimates, by taking the logarithm of both sides and using analyticity (and the Cauchy integral formula) we see that when $|z| = O(n^{2/p})$ and n is large enough then

$$\ln \mathbb{E}[e^{zS_n^\omega \tilde{u}}] = \Pi_{\omega, n}(z) + O(|z|n^{4/p}) + O(\tilde{\delta}_\omega^n)$$

where $\Pi_{\omega, n}(z) = \sum_{j=0}^{n-1} \Pi_{\theta^j \omega}(z)$, $\tilde{\delta}_\omega \in (0, 1)$ and we have used that $\ln(1+w) = O(w)$ when $|w|$ is small enough. By taking the derivatives at $z = 0$ and using the Cauchy integral formula we see that

$$(4.29) \quad 0 = \mathbb{E}[S_n^\omega \tilde{u}] = \Pi'_{\omega, n}(0) + O(n^{6/p})$$

and

$$(4.30) \quad \sigma_{\omega, n}^2 = \mathbb{E}[(S_n^\omega \tilde{u})^2] = \Pi''_{\omega, n}(0) + O(n^{8/p}).$$

Moreover, since $|\Pi_{\omega, n}(z)| = O(n)$, by using the Cauchy integral formula to estimate the error term in the second order Taylor expansion of $\Pi_{\omega, n}(z)$ around $z = 0$ we see that when $|z| = O(n^{-2/p})$ then

$$(4.31) \quad \Pi_{\omega, n}(z) = z\Pi'_{\omega, n}(0) + \frac{1}{2}z^2\Pi''_{\omega, n}(0) + |z|^3O(n^{1+8/p})$$

and so

$$\ln \mathbb{E}[e^{zS_n^\omega \tilde{u}}] = O(n^{6/p})z + \frac{1}{2}z^2\sigma_{\omega, n}^2 + O(n^{8/p})z^2 + |z|^3O(n^{1+8/p}) + O(|z|n^{4/p}) + O(\tilde{\delta}_\omega^n).$$

Finally, let us fix some $t \in \mathbb{R}$ and take $z = t_n = ta_n/n$. Then, since $p > 8$, $a_n \gg n^{6/p}$ and $\frac{a_n}{n} = o(n^{-8/p})$ we have

$$\frac{\ln \mathbb{E}[e^{t(a_n/n)S_n^\omega \tilde{u}}]}{a_n^2/n} = o(1) + \frac{1}{2}t^2(\sigma_{\omega, n}^2/n).$$

Thus,

$$\lim_{n \rightarrow \infty} \frac{1}{a_n^2/n} \ln \mathbb{E}[e^{t(a_n/n)S_n^\omega \tilde{u}}] = \frac{1}{2}t^2\sigma^2$$

and the proof of Theorem 40 is complete. \square

4.5. Berry-Esseen type estimates: proof of Theorem 43. In this section we will prove Theorem 43 (ii), and the proof of Theorem 43 (i) is similar (we will provide a few details after completing the proof of the second part).

67. Lemma. *Let $\Pi_{\omega, n}$ be as defined in the proof of Theorem 40.*

(i) *We have*

$$\left| \frac{\Pi''_{\omega, n}(0)}{\sigma_{\omega, n}^2} - 1 \right| = O(n^{8/p-1}) = o(1).$$

(ii) *On any domain of the form $|t/\sigma_{\omega, n}| = O(n^{-2/p})$ we have*

$$|\lambda_{\omega, n}(it/\sigma_{\omega, n})| = |e^{\Pi_{\omega, n}(it/\sigma_{\omega, n})}| \leq e^{-ct^2/2}$$

where $c \in (0, \frac{1}{2})$ is some constant.

Proof. The first part follows from (4.30), and the second part follows from the first and (4.31) together with the fact that $\Pi'_{\omega, n}(0) \in \mathbb{R}$ (since $\Pi_{\omega, n}(t) \in \mathbb{R}$ when t is real) and recalling that $\sigma_{\omega, n}^2$ grows linearly fast in n . \square

Proof of Theorem 43 (ii). Suppose $\mu_\omega(u_\omega) = 0$. Let $d_n = n^{\frac{1}{2}-2/p}$. Then by the Esseen inequality (see [37] or a generalized version [30, §XVI.3]) there is an absolute constant C so that

$$(4.32) \quad \sup_{t \in \mathbb{R}} |\mu_\omega(S_n^\omega u \leq t\sigma_{\omega,n}) - \Phi(t)| \leq \frac{C}{d_n} + \int_{-d_n}^{d_n} \frac{|\mu_\omega(e^{itS_n^\omega u/\sigma_{\omega,n}}) - e^{-t^2/2}|}{|t|} dt.$$

In order to bound the integral on the right hand side, first by (4.31), (4.29), (4.30) and Lemma 67 (i), for every $t \in [-d_n, d_n]$ we have

$$\Pi_{\omega,n}(it/\sigma_{\omega,n}) = -t^2/2 + O(|t|n^{6/p-1/2}) + O(t^2n^{8/p-1}) + O(n^{8/p-1/2}|t|^3)$$

where we have used that $\sigma_{\omega,n}^2$ grows linearly fast in n , which, in particular, insures that $z = it/\sigma_{\omega,n} = O(n^{-2/p})$. Using also Lemma 67 (ii) and the mean value theorem we get that

$$\left| e^{\Pi_{\omega,n}(it/\sigma_{\omega,n})} - e^{-t^2/2} \right| \leq e^{-ct^2/2} \left(O(|t|n^{6/p-1/2}) + O(t^2n^{8/p-1}) + O(n^{8/p-1/2}|t|^3) \right).$$

Using now (4.23), (4.27) and Lemma 67 (ii) we see that

$$(4.33) \quad \left| \mu_\omega(e^{itS_n^\omega u/\sigma_{\omega,n}}) - e^{-t^2/2} \right| \leq C_\omega |t| e^{-ct^2/2} \left(n^{6/p-1/2} + |t|n^{8/p-1} + t^2n^{8/p-1/2} + n^{4/p-1/2} \right)$$

for some constant C_ω which depends on ω but not on t or n . The proof of Theorem 43 (ii) is completed now by combining (4.32) with (4.33). \square

The proof of Theorem 43 (i) proceeds similarly for n 's so that $\theta^n \omega \in A$, and in order to pass to general indexes n we use that $\Psi_{\theta^j \omega} = o(n^{1/p})$ together with [32, Lemma 3.3] (applied with $a = \infty$).

4.6. A moderate local limit theorem: proof of Theorem 45. As in the proof of Theorem 43, let us assume that $\mu_\omega(u_\omega) = 0$. By using a density argument (see [36, Section VI.4]) it is enough to obtain (3.6) for a function $g \in L^1(\mathbb{R})$ whose Fourier transform has a compact support. Note that such a function g satisfies the inversion formula. Let g be a function with these properties and let $L > 0$ be so that $\hat{g}(x) = 0$ if $|x| > L$. Then, by the inversion formula for g , for all $y \in \mathbb{R}$ we have

$$g(y) = \int_{-\infty}^{\infty} \hat{g}(x) e^{iyx} dx = \int_{-L}^L \hat{g}(x) e^{iyx} dx.$$

Taking some $v \in \mathbb{R}$, setting $y = S_n^\omega u/a_n - v$ and then integrating with respect to μ_ω we see that

$$\begin{aligned} \mathbb{E}_{\mu_\omega}[g(S_n^\omega u/a_n - v)] &= \mathbb{E}_{\mu_\omega} \left[\int_{-L}^L \hat{g}(x) e^{ixS_n^\omega u/a_n} e^{-ivx} dx \right] = \int_{-L}^L \hat{g}(x) e^{-ivx} \mu_\omega(e^{itS_n^\omega u/a_n}) dx = \\ &= \frac{a_n}{\sigma_{\omega,n}} \int_{-L\sigma_{\omega,n}/a_n}^{L\sigma_{\omega,n}/a_n} \hat{g}(a_n t/\sigma_{\omega,n}) e^{-iva_n t/\sigma_{\omega,n}} \mathbb{E}[e^{itS_n^\omega u/\sigma_{\omega,n}}] dt \end{aligned}$$

where in the last equality we used the change of variables $x = \frac{a_n}{\sigma_{\omega,n}} t$. Here (a_n) is the sequence specified in Theorem 45. Now, since $a_n n^{-2/p} \rightarrow \infty$, the estimate (4.33) is valid on the domain $\{|t| \leq L\sigma_{\omega,n}/a_n\}$. Therefore, uniformly in $v \in \mathbb{R}$, we have

$$\frac{\sigma_{\omega,n}}{a_n} \mathbb{E}_{\mu_\omega}[g(S_n^\omega u/a_n - v)] - \int_{-L\sigma_{\omega,n}/a_n}^{L\sigma_{\omega,n}/a_n} \hat{g}(a_n t/\sigma_{\omega,n}) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt = o(1).$$

Next, set $\kappa_n = \kappa_{\omega,n} = \sigma_{\omega,n}/a_n$. Then, in order to complete the proof of Theorem 45 we need to show that, uniformly in $v \in \mathbb{R}$, we have

$$\int_{-L\sigma_{\omega,n}/a_n}^{L\sigma_{\omega,n}/a_n} \hat{g}(a_n t/\sigma_{\omega,n}) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt - \frac{1}{\sqrt{2\pi}} e^{-\frac{v^2}{2\kappa_n^2}} = o(1).$$

To prove that let us take an arbitrary small $\varepsilon > 0$ and fix T large enough so that

$$(4.34) \quad \|g\|_{L^1(\mathbb{R})} \int_{|t|>T} e^{-t^2/2} dt < \varepsilon/3.$$

Then, using that $\sup |\hat{g}| \leq \|g\|_{L^1(\mathbb{R})}$ we see that for every n large enough and all $v \in \mathbb{R}$ we have

$$\left| \int_{-L\sigma_{\omega,n}/a_n}^{L\sigma_{\omega,n}/a_n} \hat{g}(a_n t/\sigma_{\omega,n}) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt - \int_{-T}^T \hat{g}(a_n t/\sigma_{\omega,n}) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt \right| < \varepsilon/3$$

where we have used that $\sigma_{\omega,n}/a_n \rightarrow \infty$. Next, since $\lim_{x \rightarrow 0} \hat{g}(x) = \hat{g}(0) = \int g(y) dy$ we see that for every n large enough we have

$$\sup_{v \in \mathbb{R}} \left| \int_{-T}^T \hat{g}(a_n t/\sigma_{\omega,n}) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt - \int_{-T}^T \hat{g}(0) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt \right| < \varepsilon/3.$$

Now, using again (4.34) we see that

$$\sup_{v \in \mathbb{R}} \left| \int_{-T}^T \hat{g}(0) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt - \int_{-\infty}^{\infty} \hat{g}(0) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt \right| < \varepsilon/3.$$

We conclude from the above estimates that, for every n large enough uniformly in v we have

$$\left| \frac{\sigma_{\omega,n}}{a_n} \mathbb{E}_{\mu_\omega} [g(S_n^\omega u/a_n - v)] - \int_{-\infty}^{\infty} \hat{g}(0) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt \right| < \varepsilon.$$

Finally, by the inversion formula

$$\int_{-\infty}^{\infty} \hat{g}(0) e^{-iva_n t/\sigma_{\omega,n}} e^{-t^2/2} dt = \hat{g}(0) \frac{1}{\sqrt{2\pi}} e^{-\frac{v^2}{2\kappa_n^2}}$$

and the proof of Theorem 45 is complete.

5. PROOF OF THE REAL RPF THEOREM (THEOREM 47)

5.1. Effective rates for properly expanding maps: proof of Theorem 47 (i)-(iii) in the setup of Section 2.1. For the sake of completeness, in this section we will also consider the setup described in Remark 4, where for the sake of simplicity we focus on the case $n_0 = 1$ (to consider the case $n_0 > 1$ we essentially need to replace T_ω with $T_\omega^{n_0}$ and ϕ_ω with $S_{n_0}^\omega \phi$). The setup from Section 2.1 will be referred to as “the case $\xi = 1$ ” (as we can pair the inverse images of any two points), while the setup from Remark 4 will be referred to as “the case $\xi < 1$ ”.

5.1.1. The cones. For each $a > 0$ let us consider the real Birkhoff cone

$$\mathcal{C}_{\omega,a} = \{g \in \mathcal{H}_{\omega,\alpha} : g \geq 0, g(x) \leq e^{a\rho(x,x')^\alpha} g(x') \forall x, x' \in \mathcal{E}_\omega \text{ with } \rho(x, x') \leq \xi\}.$$

Set also $\mathcal{C}_\omega = \mathcal{C}_{\omega,\gamma_\omega^\alpha}$.

68. Lemma. *We have*

$$\mathcal{L}_\omega \mathcal{C}_\omega \subset \mathcal{C}_{\theta_\omega, H_\omega + 1} \subset \mathcal{C}_{\theta_\omega}.$$

Proof. First, by (2.5) we have $H_\omega + 1 \leq \gamma_{\theta_\omega}^\alpha$ and so the second inclusion holds true. To prove the first inclusion, let $g \in \mathcal{C}_\omega$ and let $x, x' \in \mathcal{E}_{\theta_\omega}$ be so that $\rho(x, x') \leq \xi$. Then, with $y_i = y_{i,\omega}(x)$ and $y'_i = y_{i,\omega}(x')$ as in (2.2), we have

$$\begin{aligned} \mathcal{L}_\omega g(x) &= \sum_i e^{\phi_\omega(y_i)} g(y_i) \leq \sum_i e^{\phi_\omega(y'_i) + \rho_\omega^\alpha(x, x') H_\omega} e^{\gamma_\omega^\alpha \rho_\omega^\alpha(y_i, y'_i)} g(y'_i) \\ &\leq e^{(H_\omega + \gamma_\omega^\alpha \gamma_\omega^{-\alpha}) \rho_{\theta_\omega}^\alpha(x, x')} \sum_i e^{\phi_\omega(y'_i)} g(y'_i) = e^{(H_\omega + 1) \rho_{\theta_\omega}(x, x')} \mathcal{L}_\omega g(x'). \end{aligned}$$

□

Next,

69. **Lemma.** For all $g \in \mathcal{C}_\omega$ and every $x, x' \in \mathcal{E}_{\theta\omega}$ we have

$$(5.1) \quad \mathcal{L}_\omega g(x) \leq B_{\omega,0} \mathcal{L}_\omega g(x')$$

where when $\xi < \text{diam}(\mathcal{E}_\omega) = 1$

$$B_{\omega,0} = e^{H_\omega \xi^\alpha + \gamma_\omega^\alpha \xi^\alpha} \text{deg}(T_\omega).$$

while when $\xi = 1$ we have

$$B_{\omega,0} = e^{\gamma_{\theta\omega}^\alpha}.$$

Proof. Suppose first that $\xi < 1$. Then

$$\mathcal{L}_\omega g(x) \leq \text{deg}(T_\omega) \max_{y \in T_\omega^{-1}\{x\}} e^{\phi_\omega(y)} g(y) = e^{\phi_\omega(y_0)} g(y_0)$$

for some y_0 . On the other hand, let $y' \in T_\omega^{-1}\{x'\}$ be so that $\rho(y_0, y') \leq \xi$ (existence of such y' follows from our assumptions on the map T_ω). Then, since $g \in \mathcal{C}_\omega$, we have

$$e^{\phi_\omega(y_0)} g(y_0) \leq e^{H_\omega + \gamma_\omega^\alpha \xi^\alpha} e^{\phi_\omega(y')} g(y')$$

where we have also used (2.4). On the other hand,

$$e^{\phi_\omega(y')} g(y') \leq \mathcal{L}_\omega g(x')$$

which together with the previous estimates yields the desired result in the case $\xi < 1$.

When $\xi = 1$ then $\mathcal{L}_\omega g \in \mathcal{C}_{\theta\omega}$ and so (since $\xi = 1$),

$$\mathcal{L}_\omega g(x) \leq e^{\gamma_\omega^\alpha} \mathcal{L}g(x')$$

for all x, x' . □

70. **Corollary.** The the projective diameter of $\mathcal{L}_\omega \mathcal{C}_\omega$ inside $\mathcal{C}_{\theta\omega}$ does not exceed $D(\omega)$ (which was defined in Section 2.3.1).

Proof. The statement follows from Lemma 68 and 69, and it appears in various forms in several places, and we refer to [33, Lemma 5.7.1] or [40]. □

5.1.2. *Reconstruction of ν_ω using dual cones.* Let $\mathcal{C}_\omega = \mathcal{C}_{\omega, \gamma_\omega^\alpha}$ and let \mathcal{C}_ω^* be the dual cone which is given by

$$\mathcal{C}_\omega^* = \{\nu \in \mathcal{H}_\omega^* : \nu(g) \geq 0, \forall g \in \mathcal{C}_\omega\}.$$

Let $\mathcal{L}_\omega^* : \mathcal{H}_{\theta\omega}^* \rightarrow \mathcal{H}_\omega^*$ be the dual operator. Then by [33, Lemma A.2.6] the projective diameter of $\mathcal{L}_\omega^* \mathcal{C}_{\theta\omega}^*$ inside \mathcal{C}_ω^* equals the the projective $\mathcal{L}_\omega \mathcal{C}_\omega$, inside $\mathcal{C}_{\theta\omega}$ (which by Corollary 70 does not exceed $D(\omega)$). Note that [33, Lemma A.2.6] is technically about complex cones, but the arguments needed in the case of real cones are essentially the same³¹.

We need now the following result.

71. **Lemma.** For every $\mu \in \mathcal{C}_\omega^*$ and all $h \in \mathcal{H}$ we have

$$|\mu(h)| \leq 2 \|h\| \mu(\mathbf{1}).$$

Proof. First, let us show that a closed ball of radius 1/2 around $\mathbf{1}$ is contained in \mathcal{C}_ω . Indeed, let $h = 1 + f$ where $\|f\| \leq \frac{1}{2}$. Then h belongs to \mathcal{C}_ω if and only if for all x and x' so that $\rho(x, x') \leq \xi$ we have

$$h(x) \leq e^{\gamma_\omega^\alpha \rho(x, x')^\alpha} h(x'), \quad \rho(x, x')^\alpha = (\rho(x, x'))^\alpha$$

which can also be written as

$$(5.2) \quad f(x) - f(x') \leq (e^{\gamma_\omega^\alpha \rho(x, x')^\alpha} - 1)(1 + f(x')).$$

Now, since $e^t - 1 \geq t$ for all $t \geq 0$ and $1 + f(x') \geq 1 - \|f\|_\infty$ we have

$$(e^{\gamma_\omega^\alpha \rho(x, x')^\alpha} - 1)(1 + f(x')) \geq \gamma_\omega^\alpha \rho(x, x')^\alpha (1 - \|f\|_\infty) \geq \frac{1}{2} \rho(x, x')^\alpha$$

³¹Noticing also that the closure of the cone $\tilde{\mathcal{C}}_\omega = \{\nu \in \mathcal{H}_\omega^* : \nu(g) > 0, \forall g \in \mathcal{C}_\omega \setminus \{0\}\}$ coincides with \mathcal{C}_ω^* (because there is a linear functional which is strictly positive on $\mathcal{C}_\omega \setminus \{0\}$).

where we have used that $\gamma_\omega^\alpha \geq 1$ and $\|f\|_\infty \leq \|f\| \leq \frac{1}{2}$. On the other hand, since $v(f) \leq \|f\| \leq \frac{1}{2}$ we have

$$f(x) - f(x') \leq \rho(x, x')^\alpha v(f) \leq \frac{1}{2} \rho(x, x')^\alpha.$$

Combing the last two estimates we obtain (5.2).

Next, let $\mu \in \mathcal{C}_\omega^*$, and let $h \in \mathcal{H}_\omega$ be so that $\|h\| \leq 1$. Then $1 \pm \frac{1}{2}h \in \mathcal{C}_\omega$ and so

$$\mu(1 \pm \frac{1}{2}h) \geq 0,$$

that is

$$|\mu(h)| \leq 2\mu(\mathbf{1}).$$

□

Next, let $\rho(\omega) = \tanh(D(\omega)/4)$ (as was defined in Section 2.3.1). Let $\mu \in \mathcal{C}_{\theta^n \omega}^*$ and $\nu \in \mathcal{C}_{\theta^m \omega}^*$ for some $m \geq n$. Then by the projective contraction properties of linear maps (see [10] and [41, Theorem 1.1]) the projective distance between $(\mathcal{L}_\omega^n)^* \mu$ and $(\mathcal{L}_\omega^m)^* \nu = (\mathcal{L}_\omega^n)^* (\mathcal{L}_{\theta^{m-n} \omega}^*)^* \nu$ does not exceed $\rho_{\omega, n} = \prod_{j=0}^{n-1} \rho(\theta^j \omega)$. Hence by³² [33, Theorem A.2.3] and [43, Lemma 5.2],

$$\left\| \frac{(\mathcal{L}_\omega^n)^* \mu}{\mu(\mathcal{L}_\omega^n \mathbf{1})} - \frac{(\mathcal{L}_\omega^m)^* \nu}{\nu(\mathcal{L}_\omega^m \mathbf{1})} \right\| \leq \sqrt{2} \rho_{\omega, n}.$$

Notice that $\rho_{\omega, n}$ converges exponentially fast to 0 for \mathbb{P} -a.a. ω (indeed $\rho(\cdot) < 1$ and θ is ergodic). Thus, for any sequence μ_n so that $\mu \in \mathcal{C}_{\theta^n \omega}^*$ the limit

$$\nu_\omega = \lim_{n \rightarrow \infty} \frac{(\mathcal{L}_\omega^n)^* \mu}{\mu(\mathcal{L}_\omega^n \mathbf{1})}$$

exists, belongs to \mathcal{C}_ω^* and it does not depend on the choice of the sequence (hence $\omega \rightarrow \nu_\omega$ is measurable). Moreover, by fixing n and letting $m \rightarrow \infty$ we have

$$(5.3) \quad \left\| \frac{(\mathcal{L}_\omega^n)^* \mu}{\mu(\mathcal{L}_\omega^n \mathbf{1})} - \nu_\omega \right\| \leq \sqrt{2} \rho_{\omega, n}.$$

Note that $\nu_\omega(\mathbf{1}) = 1$. Furthermore, by plugging in $(\mathcal{L}_{\theta \omega}^n)^* \mu$ inside \mathcal{L}_ω^* and using (5.3) with $\theta \omega$ instead of ω we see that there is a number λ_ω so that $\mathcal{L}_\omega^* \nu_{\theta \omega} = \lambda_\omega \nu_\omega$. Plugging in $g = \mathbf{1}$ we also see that $\lambda_\omega = \nu_{\theta \omega}(\mathcal{L}_\omega \mathbf{1})$. Finally, since \mathcal{H}_ω is dense in $C(\mathcal{E}_\omega)$ and ν_ω is positive we get that ν_ω can be extended to a probability measure on \mathcal{E}_ω .

5.1.3. *Reconstruction of h_ω with effective rates.* We first need the following result.

72. **Lemma.** *In the case $\xi < 1$ for every i we have*

$$\nu_\omega(B(x_i, \xi)) \geq \frac{e^{-2\|\phi_\omega\|_\infty}}{\deg(T_\omega)} := b_\omega$$

where the points $x_i = x_{i, \omega}$ are described in Section 2.1.

Proof. First, recall our (covering) assumption that for all i we have $T_\omega(B_\omega(x_i, \xi)) = \mathcal{E}_{\theta \omega}$. Hence, for every $x \in \mathcal{E}_{\theta \omega}$ we have

$$(\mathcal{L}_\omega(\mathbf{1}_{B_\omega(x_i, \xi)}))(x) \geq e^{-\|\phi_\omega\|_\infty}.$$

Next, since $\lambda_\omega = \nu_{\theta \omega}(\mathcal{L}_\omega \mathbf{1})$ we have

$$\lambda_\omega \leq \|\mathcal{L}_\omega \mathbf{1}\|_\infty \leq \deg(T_\omega) e^{\|\phi_\omega\|_\infty}$$

and so

$$\nu_\omega(B(x_i, \xi)) = \nu_{\theta \omega}(\lambda_\omega^{-1} \mathcal{L}_\omega(\mathbf{1}_{B_\omega(x_i, \xi)})) \geq \lambda_\omega^{-1} e^{-\|\phi_\omega\|_\infty} \geq \frac{e^{-2\|\phi_\omega\|_\infty}}{\deg(T_\omega)}.$$

□

³²These results are formulated for complex cones, but a real complex cone is embedded in its canonical complexification (together with the corresponding projective metrics)

73. Lemma. For every $g \in \mathcal{C}_\omega$ we have

$$\|g\| \leq K_\omega \nu_\omega(g)$$

where when $\xi < 1$,

$$K_\omega = e^{2\|\phi_\omega\|_\infty + 2\xi^\alpha \gamma_\omega^\alpha} \deg(T_\omega)(1 + \gamma_\omega^\alpha)$$

while when $\xi = 1$ we have $K_\omega = (1 + \gamma_\omega^\alpha)e^{\gamma_\omega^\alpha}$.

Proof. Fix some $g \in \mathcal{C}_\omega$. Suppose first that $\xi < 1$. Let $x \in \mathcal{E}_\omega$ and let i be so that $\rho(x, x_i) \leq \xi$, $x_i = x_{i,\omega}$. Then since $g \in \mathcal{C}_\omega$ we have

$$\begin{aligned} g(x) &\leq e^{\xi^\alpha \gamma_\omega^\alpha} g(x_i) \leq e^{2\xi^\alpha \gamma_\omega^\alpha} \inf\{g(y) : d(y, x_i) \leq \xi\} \leq \frac{e^{2\xi^\alpha \gamma_\omega^\alpha}}{\nu_\omega(B(x_i, \xi))} \int_{B_\omega(x_i, \xi)} g(z) dz \\ &\leq e^{2\xi^\alpha \gamma_\omega^\alpha} b_\omega^{-1} \nu_\omega(g). \end{aligned}$$

where in the last inequality we have used Lemma 72. By taking the supremum over all possible choices of x we see that

$$\sup g = \|g\|_\infty \leq e^{2\xi^\alpha \gamma_\omega^\alpha} b_\omega^{-1} \nu_\omega(g).$$

When $\xi = 1$ we have

$$\sup g \leq e^{\gamma_\omega^\alpha} \inf g \leq e^{\gamma_\omega^\alpha} \nu_\omega(g).$$

Finally, in both cases, if $g(x) > g(x')$ and $\rho(x, x') \leq \xi$ then

$$g(x) - g(x') = g(x)(1 - g(x)/g(x')) \leq \|g\|_\infty (1 - e^{-\gamma_\omega^\alpha \rho^\alpha(x, x')}) \leq \|g\|_\infty \gamma_\omega^\alpha \rho^\alpha(x, x').$$

Thus,

$$v(g) = v_{\alpha, \xi}(g) \leq \|g\|_\infty \gamma_\omega^\alpha$$

and so

$$\|g\| = v(g) + \|g\|_\infty \leq (1 + \gamma_\omega^\alpha) \|g\|_\infty.$$

Now the lemma follows from the above upper bounds on $\|g\|_\infty$. \square

Next, arguing as in [33, Theorem 5.3.1 (iii)] we can prove the following result.

74. Lemma. For every $f \in \mathcal{H}_\omega$ there are $f_1, f_2 \in \mathcal{C}_\omega$ and negative constants c_1, c_2 so that $f = f_1 - c_1 - (f_2 - c_2)$ and

$$\|f_1\| + \|f_2\| + |c_1| + |c_2| \leq r_\omega \|f\|$$

where $r_\omega = 4\left(1 + \frac{2}{\gamma_\omega^\alpha}\right) \leq 8$.

Next, by applying [33, Theorem A.2.3] and taking into account Corollary 70 and Lemma 73 we see that for every $g \in \mathcal{C}_{\theta^{-n}\omega}$ and $f \in \mathcal{C}_{\theta^{-m}\omega}$ with $m \geq n$ we have

$$\left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^n g}{\nu_{\theta^{-n}\omega}(\mathcal{L}_{\theta^{-n}\omega}^n g)} - \frac{\mathcal{L}_{\theta^{-m}\omega}^n f}{\nu_{\theta^{-m}\omega}(\mathcal{L}_{\theta^{-m}\omega}^n f)} \right\| \leq \frac{1}{2} K_\omega \rho_{\theta^{-n}\omega, n}.$$

Notice that

$$\nu_{\theta^{-n}\omega}(\mathcal{L}_{\theta^{-n}\omega}^n g) = \lambda_{\theta^{-n}\omega, n} \nu_{\theta^{-n}\omega}(g), \quad \nu_{\theta^{-m}\omega}(\mathcal{L}_{\theta^{-m}\omega}^m f) = \lambda_{\theta^{-m}\omega, m} \nu_{\theta^{-m}\omega}(f)$$

where $\lambda_{\omega, n} = \prod_{j=0}^{n-1} \lambda_{\theta^j \omega}$. We thus see that for every sequence (g_n) so that $g_n \in \mathcal{C}_{\theta^{-n}\omega}$ the limit

$$(5.4) \quad h_\omega = \lim_{n \rightarrow \infty} \frac{\mathcal{L}_{\theta^{-n}\omega}^n g_n}{\nu_{\theta^{-n}\omega}(g_n) \lambda_{\theta^{-n}\omega, n}}$$

exists, it does not depend on the sequence (g_n) , it belongs to \mathcal{C}_ω and $\nu_\omega(h_\omega) = 1$ (and so by Lemma 73 we have $\|h_\omega\| \leq K_\omega$). Moreover, by taking $g_n = \mathbf{1}$ for every n and applying \mathcal{L}_ω to both sides of (5.4) we see that $\mathcal{L}_\omega h_\omega = \lambda_\omega h_\omega$. Furthermore, by fixing n , taking $f = f_m \in \mathcal{C}_{\theta^{-m}\omega}$ and letting $m \rightarrow \infty$ we see that for every $g \in \mathcal{C}_{\theta^{-n}\omega}$ we have

$$(5.5) \quad \left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^n g}{\lambda_{\theta^{-n}\omega, n}} - \nu_{\theta^{-n}\omega}(g) h_\omega \right\| \leq \frac{1}{2} K_\omega \nu_{\theta^{-n}\omega}(g) \rho_{\theta^{-n}\omega, n}.$$

In addition, since $h_\omega \in \mathcal{C}_\omega$ by (5.1), for all $x, x' \in \mathcal{E}_{\theta\omega}$ we have

$$h_{\theta\omega}(x) = (\lambda_\omega)^{-1} \mathcal{L}_\omega h_\omega(x) \leq \lambda_\omega^{-1} B_{\omega,0} \mathcal{L}_\omega h_\omega(x') = B_{\omega,0} h_{\theta\omega}(x').$$

Since $\nu_\omega(h_\omega) = 1$ we conclude that $\min h_{\theta\omega} \geq B_{\omega,0}^{-1}$.

Finally, by Lemma 74, for every $g \in \mathcal{H}_{\theta^{-n}\omega}$, there are g_1, g_2 in $\mathcal{C}_{\theta^{-n}\omega}$ and nonnegative constants c_1, c_2 so that $g = g_1 - c_1 - (g_2 - c_2)$ and

$$\|g_1\| + \|g_2\| + |c_1| + |c_2| \leq 8\|g\|.$$

Thus, by applying (5.5) with $g = g_1, g = g_2$ and the constant functions $g = -c_1$ and $g = -c_2$ we see that

$$\left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^n g}{\lambda_{\theta^{-n}\omega, n}} - \nu_{\theta^{-n}\omega}(g) h_\omega \right\| \leq 4K_\omega \|g\| \rho_{\theta^{-n}\omega, n}.$$

5.2. Effective rates for partially expanding maps: proof of Theorem 47 (i)-(iii) in the setup of Section 2.2.

5.2.1. *The cones.* Set $\kappa_\omega = \frac{1}{s_\omega}$ and consider the real cone

$$\mathcal{C}_\omega = \mathcal{C}_{\omega, \kappa_\omega} = \{g \in \mathcal{H}_\omega : g > 0 \text{ and } v(g) \leq \kappa_\omega \inf g\}.$$

75. **Lemma.** *We have*

$$(5.6) \quad \mathcal{L}_\omega \mathcal{C}_\omega \subset \mathcal{C}_{\theta\omega, \zeta_\omega \kappa_{\theta\omega}} \subset \mathcal{C}_{\theta\omega}$$

where³³ $\zeta_\omega = \frac{s_\omega \kappa_\omega + (1 + \kappa_\omega) e^{\varepsilon_\omega} H_\omega}{\kappa_{\theta\omega}} = s_{\theta\omega} (1 + (1 + s_\omega^{-1}) e^{\varepsilon_\omega} H_\omega) < 1$.

Proof of Lemma 68. The proof of (5.6) proceeds similarly to the proof of [16, Theorem 5.1]. Let $g \in \mathcal{C}_\omega = \mathcal{C}_{\omega, \kappa_\omega}$. Fix some ω and two points x, y in $\mathcal{E}_{\theta\omega}$ and denote by (x_i) and (y_i) their inverse images under T_ω , respectively. Then

$$\begin{aligned} \frac{|\mathcal{L}_\omega g(x) - \mathcal{L}_\omega^{(j)} g(y)|}{\inf \mathcal{L}_\omega g} &\leq \frac{|\mathcal{L}_\omega g(x) - \mathcal{L}_\omega g(y)|}{d_\omega e^{\inf \phi_\omega} \inf g} \leq d_\omega^{-1} \sum_{i=1}^{d_\omega} e^{\phi_\omega(x_i) - \inf \phi_\omega} |g(x_i) - g(y_i)| (\inf g)^{-1} \\ &\quad + d_\omega^{-1} \sum_{i=1}^{d_\omega} |(g(y_i) / \inf g) e^{-\inf \phi_\omega} |e^{\phi_\omega(x_i)} - e^{\phi_\omega(y_i)}| := I_1 + I_2. \end{aligned}$$

Next, since $g \in \mathcal{C}_\omega$ for each i we have $|g(x_i) - g(y_i)| \leq v(g) \rho^\alpha(x_i, y_i) \leq \kappa_\omega \inf g \cdot \rho^\alpha(x_i, y_i)$. Moreover, we have $\phi_\omega(x_i) - \inf \phi_\omega \leq \sup \phi_\omega - \inf \phi_\omega = \varepsilon_\omega$. Combining these estimates and taking into account (2.7), (2.8) and (2.9) we get that

$$I_1 \leq \kappa_\omega \rho^\alpha(x, y) e^{\varepsilon_\omega} d_\omega^{-1} (I_\omega^\alpha q_\omega + (d_\omega - q_\omega) \sigma_\omega^{-\alpha}) = \rho^\alpha(x, y) s_\omega \kappa_\omega$$

where we recall that s_ω was defined in (2.12).

In order to bound I_2 , we first observe that $\sup g \leq \inf g + v(g) \leq (1 + \kappa_\omega) \inf g$ and that by the definition 2.11 of the local Hölder constant H_ω and the mean value theorem we see that

$$|e^{\phi_\omega(x_i)} - e^{\phi_\omega(y_i)}| \leq e^{\max(\phi_\omega(x_i), \phi_\omega(y_i))} |\phi_\omega(x_i) - \phi_\omega(y_i)| \leq e^{\inf \phi_\omega + \varepsilon_\omega} H_\omega \rho^\alpha(x, y).$$

Using these estimates we obtain that

$$I_2 \leq \rho^\alpha(x, y) (1 + \kappa_\omega) e^{\varepsilon_\omega} H_\omega.$$

We conclude that

$$v(\mathcal{L}_\omega g) \leq (s_\omega \kappa_\omega + (1 + \kappa_\omega) e^{\varepsilon_\omega} H_\omega) \inf \mathcal{L}_\omega g = \zeta_\omega \kappa_{\theta\omega} \inf \mathcal{L}_\omega g$$

and therefore

$$(5.7) \quad \mathcal{L}_\omega \mathcal{C}_{\omega, \kappa_\omega} \subset \mathcal{C}_{\theta\omega, \zeta_\omega \kappa_{\theta\omega}} \subset \mathcal{C}_{\theta\omega, \kappa_{\theta\omega}} = \mathcal{C}_{\theta\omega}.$$

□

³³The fact that $\zeta_\omega < 1$ follows from the condition on H_ω in (2.12).

Next,

76. Corollary. *The projective diameter of $\mathcal{L}_\omega \mathcal{C}_\omega$ inside $\mathcal{C}_{\theta\omega}$ does not exceed*

$$D(\omega) := 2 \ln \left(\frac{1 + \zeta_\omega}{1 - \zeta_\omega} \right) + 2 \ln (1 + \zeta_\omega \kappa_\omega).$$

Proof. See [16, Section 4] or [44, Section 5] (recalling our assumption that $\text{diam}(\mathcal{E}_\omega) \leq 1$). \square

5.2.2. Reconstruction of ν_ω using dual cones. Let \mathcal{C}_ω^* be the dual cone of \mathcal{C}_ω . Let $\mathcal{L}_\omega^* : \mathcal{H}_{\theta\omega}^* \rightarrow \mathcal{H}_\omega^*$ be the dual operator. Then, as explained in Section 5.1.2, the projective diameter of $\mathcal{L}_\omega^* \mathcal{C}_{\theta\omega}^*$ inside \mathcal{C}_ω^* equals the projective diameter of $\mathcal{L}_\omega \mathcal{C}_\omega$ inside $\mathcal{C}_{\theta\omega}$ (which does not exceed $D(\omega)$).

Next, we need the following result.

77. Lemma. *For every $\mu \in \mathcal{C}_\omega^*$ and all $h \in \mathcal{H}$ we have*

$$|\mu(h)| \leq k_\omega \|h\| \mu(\mathbf{1})$$

where $k_\omega = \frac{1+\kappa_\omega}{\kappa_\omega} = 1 + s_\omega \leq 2$ (and $\kappa_\omega = \frac{1}{s_\omega}$).

Proof. As in the proof of Lemma 71, it is enough to show that a closed ball of radius $\frac{\kappa_\omega}{1+\kappa_\omega}$ around $\mathbf{1}$ is contained in \mathcal{C}_ω . Indeed, let $h = 1 + f$ where $\|f\| < 1$. Then $\inf(1 + f) \geq 1 - \|f\|_\infty \geq 1 - \|f\|$ and $v(1 + f) = v(f) \leq \|f\|$. Hence $h \in \mathcal{C}_\omega$ if

$$\|f\| \leq \kappa_\omega (1 - \|f\|).$$

\square

Finally, let $\rho(\omega) = \tanh(D(\omega)/4)$. Let $\mu \in \mathcal{C}_{\theta^n \omega}^*$ and $\nu \in \mathcal{C}_{\theta^m \omega}^*$ for some $m \geq n$. Then the projective distance between $(\mathcal{L}_\omega^n)^* \mu$ and $(\mathcal{L}_\omega^m)^* \nu = (\mathcal{L}_\omega^n)^* (\mathcal{L}_{\theta^{m-n} \omega}^*)^* \nu$ does not exceed $\rho_{\omega, n} = \prod_{j=0}^{n-1} \rho(\theta^j \omega)$. Thus, as in Section 5.1.2 we conclude that

$$\left\| \frac{(\mathcal{L}_\omega^n)^* \mu}{\mu(\mathcal{L}_\omega^n \mathbf{1})} - \frac{(\mathcal{L}_\omega^m)^* \nu}{\nu(\mathcal{L}_\omega^m \mathbf{1})} \right\| \leq \sqrt{2} k_\omega \rho_{\omega, n}.$$

Thus, for any sequence μ_n so that $\mu \in \mathcal{C}_{\theta^n \omega}^*$ the limit

$$\nu_\omega = \lim_{n \rightarrow \infty} \frac{(\mathcal{L}_\omega^n)^* \mu}{\mu(\mathcal{L}_\omega^n \mathbf{1})}$$

exists, belongs to \mathcal{C}_ω^* and it does not depend on the choice of the sequence. Moreover, by fixing n and letting $m \rightarrow \infty$ we have

$$\left\| \frac{(\mathcal{L}_\omega^n)^* \mu}{\mu(\mathcal{L}_\omega^n \mathbf{1})} - \nu_\omega \right\| \leq \sqrt{2} k_\omega \rho_{\omega, n}.$$

Note that $\nu_\omega(\mathbf{1}) = 1$. Moreover, as in Section 5.1.2 there is a number λ_ω so that $\mathcal{L}_\omega^* \nu_{\theta\omega} = \lambda_\omega \nu_\omega$, where $\lambda_\omega = \nu_{\theta\omega}(\mathcal{L}_\omega \mathbf{1})$. Furthermore, ν_ω is a probability measure.

5.2.3. Reconstruction of h_ω with effective rates. We first need the following result.

78. Lemma. *For every $g \in \mathcal{C}_\omega = \mathcal{C}_{\omega, \kappa_\omega}$ we have*

$$\|g\| \leq K_\omega \inf g \leq K_\omega \nu_\omega(g)$$

where

$$K_\omega = 1 + 2\kappa_\omega.$$

Proof. First, since $g \in \mathcal{C}_\omega$ we have

$$\|g\| = \sup g + v(g) \leq \sup g + \kappa_\omega \inf g.$$

Second, in order to estimate $\sup g$, using that $v(g) \leq \kappa_\omega \inf g$ we see that for every $x \in \mathcal{E}_\omega$ we have $|g(x) - \inf g| \leq v(g) \text{diam}(\mathcal{E}_\omega)^\alpha \leq \kappa_\omega \text{diam}(\mathcal{E}_\omega)^\alpha \inf g$. Taking into account that $\text{diam}(\mathcal{E}_\omega) \leq 1$ we conclude that $\sup g \leq (1 + \kappa_\omega) \inf g$, which completes the proof of the lemma. \square

The next result we need is the following.

79. Lemma. For every $g \in \mathcal{H}_\omega$ there is a constant $c(g) > 0$ and a function $g_1 \in \mathcal{C}_\omega$ so that $g = g_1 - c(g)$ and

$$\|g_1\| + c(g) \leq 3\|g\|.$$

Proof. Let $c(g) = v(g)/\kappa_\omega + \sup|g| \leq \|g\|$. Then $g_1 = g + c(g) \in \mathcal{C}_\omega$ and so $g = g_1 - c_g$ and since $\kappa_\omega \geq 1$ we have

$$\|g_1\| + c(g) = \|g + c(g)\| + c(g) \leq \|g\| + 2c(g) \leq 3\|g\|.$$

□

By repeating the arguments in Section 5.1.3 we see that for every $m, n \in \mathbb{N}$ such that $m \geq n$ and all $g \in \mathcal{C}_{\theta^{-n}\omega}$ and $f \in \mathcal{C}_{\theta^{-m}\omega}$ we have

$$\left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^n g}{\nu_{\theta^{-n}\omega}(\mathcal{L}_{\theta^{-n}\omega}^n g)} - \frac{\mathcal{L}_{\theta^{-m}\omega}^n g}{\nu_{\theta^{-m}\omega}(\mathcal{L}_{\theta^{-m}\omega}^n f)} \right\| \leq \frac{1}{2} K_\omega \rho_{\theta^{-n}\omega, n}.$$

Moreover, for any sequence (g_n) so that $g_n \in \mathcal{C}_{\theta^{-n}\omega}$ the limit

$$h_\omega = \lim_{n \rightarrow \infty} \frac{\mathcal{L}_{\theta^{-n}\omega}^n g_n}{\nu_{\theta^{-n}\omega}(g_n) \lambda_{\theta^{-n}\omega, n}}$$

exists, it does not depend on (g_n) , it belongs to \mathcal{C}_ω and $\nu_\omega(h_\omega) = 1$. Furthermore, $\mathcal{L}_\omega h_\omega = \lambda_\omega h_{\theta\omega}$ and

$$\left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^n g}{\lambda_{\theta^{-n}\omega, n}} - \nu_{\theta^{-n}\omega}(g) h_\omega \right\| \leq \frac{1}{2} K_\omega \nu_{\theta^{-n}\omega}(g) \rho_{\theta^{-n}\omega, n}.$$

In addition, since $h_\omega \in \mathcal{C}_\omega$ we have $\sup h_\omega \leq B_{\omega,1} \inf h_\omega$ where $B_{\omega,1} = 1 + \kappa_\omega$ and we have used that $\text{diam}(\mathcal{E}_\omega) \leq 1$. Since $\nu_\omega(h_\omega) = 1$ we conclude that $\min h_\omega \geq B_{\omega,1}^{-1}$.

Finally, arguing as in Section 5.1.3, by using Lemma 79 instead of Lemma 74 we see that

$$\left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^n g}{\lambda_{\theta^{-n}\omega, n}} - \nu_{\theta^{-n}\omega}(g) h_\omega \right\| \leq 2K_\omega \|g\| \rho_{\theta^{-n}\omega, n}.$$

5.3. Decay of correlations and the normalized transfer operators: proof of Theorem 47 (iv)-(v). Let the operator L_ω be defined by

$$L_\omega g = \frac{\mathcal{L}_\omega(g h_\omega)}{h_{\theta\omega} \lambda_\omega}.$$

Then, using that $\|fg\| \leq 3\|f\|\|g\|$ for every two Hölder continuous functions, we see that

$$\begin{aligned} \|L_\omega^n g - \mu_\omega(g)\| &= \left\| \frac{\mathcal{L}_\omega^n(g h_\omega)}{\lambda_{\omega, n} h_{\theta^n \omega}} - \nu_\omega(g h_\omega) \right\| = \left\| \left(\frac{1}{h_{\theta^n \omega}} \right) (\mathcal{L}_\omega^n(g h_\omega) - \nu_\omega(g h_\omega) h_{\theta^n \omega}) \right\| \\ &\leq 3 \left\| \frac{1}{h_{\theta^n \omega}} \right\| \left\| \frac{\mathcal{L}_\omega^n(g h_\omega)}{\lambda_{\omega, n}} - \nu_\omega(g h_\omega) h_{\theta^n \omega} \right\|. \end{aligned}$$

Now, since $h_\omega \geq B_{\omega,1}^{-1}$ we have $\|1/h_\omega\| \leq v(h_\omega) B_{\omega,1}^2 + B_{\omega,1} \leq 2B_{\omega,1}^2 K_\omega$. Thus, by (3.8) (and recalling the formula of B_ω in Section 2.3.2),

$$\|L_\omega^n g - \mu_\omega(g)\| \leq B_{\theta^n \omega} \rho_{\omega, n}.$$

Now the decay of correlations (3.10) follows from the equality

$$\text{Cov}_{\mu_\omega}(g, f \cdot T_\omega^n) = \int f \cdot (L_\omega^n g - \mu_\omega(g)) d\mu_{\theta^n \omega}.$$

6. RANDOM COMPLEX RPF THEOREMS WITH EFFECTIVE RATES: PROOF OF THEOREM 49

As opposed to the previous sections in this section we will begin with the setup of Section 2.2. The reason is that in the setup of Section 2.1 the appropriate projective estimates needed to prove Theorem 49 are similar to [33, Ch. 4-5] (with some modifications). Henceforth, for the sake of convenience, we will always assume that $\mu_\omega(u_\omega) = 0$, namely we will replace u_ω with $\tilde{u}_\omega = u_\omega - \mu_\omega(u_\omega)$.

6.1. Complex cones contractions for random maps with expansion and contraction. The proof of Theorem 49 relies on the theory of the canonical complexification of real Birkhoff cones. We will give a reminder of the appropriate results concerning this theory in the body of the proof of Theorem 80 below, and the readers are referred to [33, Appendix A] for a summary of the main definitions and results concerning contraction properties of real and complex cones (the properties of the complex cones is essentially a summary of the appropriate results in [43, 27, 28]).

Let $\mathcal{C}_{\omega, \mathbb{C}}$ be the canonical complexification of the cone \mathcal{C}_ω (see [33, Appendix A]), and let $\mathcal{C}_{\omega, \mathbb{C}}^* := \{\nu \in \mathcal{H}_\omega^* : \nu(c) \neq 0 \ \forall \nu \in \mathcal{C}_{\omega, \mathbb{C}} \setminus \{0\}\}$ be its complex dual cone.

80. Theorem. (i) *The cones $\mathcal{C}_{\omega, \mathbb{C}}$ and their duals $\mathcal{C}_{\omega, \mathbb{C}}^*$ have bounded aperture: for all $g \in \mathcal{C}_{\omega, \mathbb{C}}$ and $\nu \in \mathcal{C}_{\omega, \mathbb{C}}^*$ and every point $x_\omega \in \mathcal{E}_\omega$ we have*

$$(6.1) \quad \|g\| \leq Q_\omega |g(x_\omega)| \quad \text{and} \quad \|\nu\| \leq M_\omega |\nu(\mathbf{1})|$$

where $Q_\omega = 2\sqrt{2}(1 + 2\kappa_\omega) = 2\sqrt{2}K_\omega$ and $M_\omega = \frac{1}{6\kappa_\omega}$, $\kappa_\omega = s_\omega^{-1}$

(ii) *The cones $\mathcal{C}_{\omega, \mathbb{C}}$ are linearly convex, namely for every $g \notin \mathcal{C}_{\omega, \mathbb{C}}$ there exists $\mu \in \mathcal{C}_{\omega, \mathbb{C}}^*$ such that $\mu(g) = 0$.*

(iii) *The cones $\mathcal{C}_{\omega, \mathbb{C}}$ are reproducing: for any complex-value function $g \in \mathcal{H}_\omega$ there is are constant $c_1(g), c_2(g) > 0$ and functions $g_1, g_2 \in \mathcal{C}_\omega \subset \mathcal{C}_{\omega, \mathbb{C}}$ so that $g = g_1 - c_1(g) + i(g_2 - c_2(g))$ and*

$$\|g_1\| + c_1(g) + \|g_2\| + c_2(g) \leq 6\|g\|.$$

(iv) *Let*

$$c_0(\omega) = 32\kappa_{\theta_\omega}^{-1}(1 + 2\kappa_\omega)e^{\|u_\omega\|_\infty + 2\|\phi_\omega\|_\infty} \|u_\omega\| (1 + H_\omega)(1 - \zeta_\omega)^{-1}$$

and for all complex z so that $|z| \leq 1$ set

$$\delta_\omega(z) = 2|z|c_0(\omega) (1 + \cosh(D(\omega)/2)).$$

Then, if $\delta_\omega(z) \leq 1 - e^{-D(\omega)}$ we have that

$$\mathcal{L}_\omega^{(z)} \mathcal{C}_{\omega, \mathbb{C}} \subset \mathcal{C}_{\theta_\omega, \mathbb{C}}$$

and the Hilbert diameter of the image with respect to the complex projective metric corresponding to the cone $\mathcal{C}_{\theta_\omega, \mathbb{C}}$ does not exceed $7D(\omega)$.

Proof of Theorem 80. (i) First (see [33, Appendix A] and [43, Section 5]) we have

$$(6.2) \quad \mathcal{C}_{\omega, \mathbb{C}} = \{g \in \mathcal{H}_\omega : \Re(\overline{\mu(g)}\nu(g)) \geq 0 \ \forall \mu, \nu \in \mathcal{C}_{\omega, \mathbb{C}}^*\}.$$

We begin with showing that the complex cones $\mathcal{C}_{\omega, \mathbb{C}}$ and their duals have bounded aperture. First, for every point $a \in \mathcal{E}_\omega$ and $g \in \mathcal{C}_\omega$ we have

$$\|g\| = \sup g + v(g) \leq \inf g + 2v(g) \leq (1 + 2\kappa_\omega) \inf g \leq (1 + 2\kappa_\omega)g(a)$$

where we have used that $g(x) - g(y) \leq (\text{diam}(\mathcal{E}_\omega))^\alpha v(g) \leq v(g)$ for every real-valued function on \mathcal{E}_ω . By applying [43, Lemma 5.2] we conclude that for every $g \in \mathcal{C}_{\omega, \mathbb{C}}$ we have

$$\|g\| \leq 2\sqrt{2}(1 + 2\kappa_\omega)g(a).$$

Next, in order to show that the cone $\mathcal{C}_{\omega, \mathbb{C}}$ has bounded aperture we will apply [33, Lemma A.2.7] which states that

$$\|\nu\| \leq M_\omega \nu(\mathbf{1}), \quad \forall \nu \in \mathcal{C}_{\omega, \mathbb{C}}^*$$

if the complex cone $\mathcal{C}_{\omega, \mathbb{C}}$ contains the ball of radius $1/M_\omega$ around the constant function $\mathbf{1}$. The first step in showing that such a ball exists is the following representation of the real cone:

$$\mathcal{C}_\omega = \{g \in \mathcal{H}_\omega : s(g) \geq 0, \forall s \in \Gamma_\omega\}$$

where $\Gamma_\omega \subset \mathcal{H}_\omega^*$ is the class of linear functionals s which either have the form $s(g) = s_a(g) = g(a)$ for some $a \in \mathcal{E}_\omega$ or have the form

$$s = s_{x,y,t,\kappa_\omega}(g) = \kappa_\omega g(t) - \frac{g(x) - g(y)}{\rho^\alpha(x,y)}$$

for some $x, y, t \in \mathcal{E}_\omega$ so that $x \neq y$. Then (see [33, Appendix A]), since Γ_ω generates the dual cone \mathcal{C}_ω^* , the canonical complexification of \mathcal{C}_ω can be written in the following form:

$$(6.3) \quad \mathcal{C}_{\omega, \mathbb{C}} = \{x \in \mathcal{H}_{\omega, \mathbb{C}} : \Re(\overline{\mu(x)}\nu(x)) \geq 0, \quad \forall \mu, \nu \in \Gamma_\omega\}.$$

Using (6.3), it is enough to show that for all g of the form $g = \mathbf{1} + h$ with $\|h\| \leq \bar{\varepsilon}_\omega := \frac{1}{M_\omega}$, and every $s_1, s_2 \in \Gamma_\omega$ we have

$$\Re(s_1(g) \cdot \overline{s_2(g)}) \geq 0.$$

Notice that $s_i(g) = 1 + s_i(h)$ and so

$$\Re(s_1(g) \cdot \overline{s_2(g)}) \geq 1 - |s_1(h)| - |s_2(h)| - |s_1(h)s_2(h)|.$$

Now, there are four cases. When $s_i(g) = g(a_i)$ for some $a_i \in \mathcal{E}_\omega$ then

$$\Re(s_1(g) \cdot \overline{s_2(g)}) \geq 1 - 2\|h\|_\infty - \|h\|_\infty^2 > 0$$

since $\|h\| \leq \bar{\varepsilon}_\omega < \frac{1}{3}$. Let us suppose next that $s_1(g) = g(a)$ and $s_2(g) = \kappa_\omega g(t) - \frac{g(x)-g(y)}{\rho^\alpha(x,y)}$. Then

$$\Re(s_1(g) \cdot \overline{s_2(g)}) \geq 1 - \kappa_\omega \|h\|_\infty - v(h) - \|h\|_\infty^2 \kappa_\omega - \|h\|_\infty v(h) \geq 1 - 2\|h\|(\kappa_\omega + 1)$$

where we have used that $\|h\| \leq 1$. Notice that the above right hand side is nonnegative if $\|h\| \leq \frac{1}{M_\omega} = \frac{1}{6\kappa_\omega}$ since $\kappa_\omega \geq 1$. A similar inequality holds true when $s_2(g) = g(a)$ and $s_1(g) = \kappa_\omega g(t) - \frac{g(x)-g(y)}{\rho^\alpha(x,y)}$. Let us assume now that $s_i(g) = \kappa_\omega g(t_i) - \frac{g(x_i)-g(y_i)}{\rho^\alpha(x_i,y_i)}$ for appropriate choices of $t_i, x_i, y_i, i = 1, 2$. Then

$$\begin{aligned} \Re(s_1(g) \cdot \overline{s_2(g)}) &\geq 1 - 2(\kappa_\omega \|h\|_\infty + v(h)) - (\kappa_\omega \|h\|_\infty + v(h))^2 \\ &\geq 1 - 2\kappa_\omega \|h\| - (\kappa_\omega \|h\|) - 2\kappa_\omega \|h\| - \|h\| \geq 1 - 6\kappa_\omega \|h\| \end{aligned}$$

where in the last two inequalities we have used that $\kappa_\omega \geq 1$ and $\|h\| \leq \kappa_\omega^{-1} \leq 1$. The above left hand side is nonnegative since $\|h\| \leq \bar{\varepsilon}_\omega = M_\omega^{-1} = \frac{1}{6\kappa_\omega}$.

(ii) By [28, Lemma 4.1], in order to show that the cone $\mathcal{C}_{\omega, \mathbb{C}}$ is linearly convex it is enough to show that there there is a continues linear functional ℓ which is strictly positive on $\mathcal{C}_\omega \setminus \{0\}$. Clearly we can take $\ell(g) = g(a)$ for an arbitrary point a in \mathcal{E}_ω .

(iii) This is a direct consequence of Lemma 79 applied with the real and imaginary parts of g .

(iv) Recall first that by Lemma 68, for all ω ,

$$(6.4) \quad \mathcal{L}_\omega \mathcal{C}_\omega \subset \mathcal{C}_{\theta\omega, \zeta_\omega \kappa_{\theta\omega}}.$$

We will next prove that for every $s \in \Gamma_{\theta\omega}$, $g \in \mathcal{C}_\omega$ (the real cone) and a complex number z so that $|z| \leq 1$ we have

$$(6.5) \quad \left| s(\mathcal{L}_\omega^{(z)} g) - s(\mathcal{L}_\omega g) \right| \leq c_0(\omega) |z| s(\mathcal{L}_\omega g).$$

After this is established we can apply [33, Theorem A.2.4] and obtain item (iv).

Let us first consider the case when $s(f) = f(a)$ for some $a \in \mathcal{E}_{\theta\omega}$. Then

$$\begin{aligned} \left| s(\mathcal{L}_\omega^{(z)} g) - s(\mathcal{L}_\omega g) \right| &= \left| \mathcal{L}_\omega(g(e^{zu_\omega} - 1))(a) \right| \leq \|e^{zu_\omega} - 1\|_\infty \mathcal{L}_\omega g(a) = \|e^{zu_\omega} - 1\|_\infty s(\mathcal{L}_\omega g) \\ &\leq |z| \|u_\omega\|_\infty e^{\|u_\omega\|_\infty} s(\mathcal{L}_\omega g) \leq c_0(\omega) |z| s(\mathcal{L}_\omega g). \end{aligned}$$

Next, let us consider the case when $s = s_{x,y,t,\kappa_{\theta\omega}}$. We first need the following simple result/observation: let A and A' be complex numbers, B and B' be real numbers, and let $\varepsilon_1 > 0$ and $\zeta \in (0, 1)$ be so that

- $B > B'$
- $|A - B| \leq \varepsilon_1 B$
- $|A' - B'| \leq \varepsilon_1 B$
- $|B'/B| \leq \zeta$.

Then

$$\left| \frac{A - A'}{B - B'} - 1 \right| \leq 2\varepsilon_1(1 - \zeta)^{-1}.$$

The proof of this result is elementary, just write

$$\left| \frac{A - A'}{B - B'} - 1 \right| \leq \left| \frac{A - B}{B - B'} \right| + \left| \frac{A' - B'}{B - B'} \right| \leq \frac{2B\varepsilon_1}{B - B'} = \frac{2\varepsilon_1}{1 - B'/B}.$$

Next, fix some nonzero $g \in \mathcal{C}_\omega$ and $(x, y, t) \in \Delta_{\theta\omega}$. Then, in order to obtain (6.5) when $s = s_{x,y,t,\kappa_\omega}$ we need to show that the conditions of the above result hold true with $A = \kappa_{\theta\omega} \mathcal{L}_\omega^{(z)} g(t)$,

$$B = \kappa_{\theta\omega} \mathcal{L}_\omega g(t), \quad A' = \frac{\mathcal{L}_\omega^{(z)} g(x) - \mathcal{L}_\omega^{(z)} g(y)}{\rho^\alpha(x, y)}, \quad B' = \frac{\mathcal{L}_\omega g(x) - \mathcal{L}_\omega g(y)}{\rho^\alpha(x, y)}$$

and $\zeta = \zeta_\omega$ and $\varepsilon_1 = 16\kappa_{\theta\omega}^{-1}(1 + 2\kappa_\omega)(1 + H_\omega)e^{\|u_\omega\|_\infty + 2\|\phi_\omega\|_\infty} \|u_\omega\| |z|$.

We begin by noting that $B > B'$ since the function $\mathcal{L}_\omega g$ is a nonzero member of the cone $\mathcal{C}_{\omega, \mathbb{R}, \zeta_\omega, \kappa_{\theta\omega}}$. In fact, this already implies that

$$|B'/B| \leq \zeta_\omega \inf \mathcal{L}_\omega g / B \leq \zeta_\omega < 1.$$

Next, notice that when $|z| \leq 1$ we have

$$\begin{aligned} |A - B| &= \kappa_{\theta\omega} |\mathcal{L}_\omega^{(z)} g(t) - \mathcal{L}_\omega g(t)| = \kappa_{\theta\omega} |\mathcal{L}_\omega(g(e^{zu_\omega} - 1))(t)| \\ &\leq \kappa_{\theta\omega} \|e^{zu_\omega} - 1\|_\infty \mathcal{L}_\omega g(t) \leq |z| e^{\|u_\omega\|_\infty} \|u_\omega\|_\infty B. \end{aligned}$$

Finally, let us estimate the difference $|A' - B'|$. For each $a, b \in \mathcal{E}_\omega$ we define

$$\Delta_{a,b}(z) = e^{\phi_\omega(a)}(e^{zu_\omega(a)} - 1)g(a) - e^{\phi_\omega(b)}(e^{zu_\omega(b)} - 1)g(b).$$

Denote again by x_i and y_i the preimages of x and y under T_ω , respectively, where $1 \leq i \leq d_\omega$. Then

$$\rho^\alpha(x, y)(A' - B') = \sum_{i=1}^{d_\omega} \Delta_{x_i, y_i}(z).$$

Next, by using the mean value theorem we see that

$$|\Delta_{x_i, y_i}(z)| = |\Delta_{x_i, y_i}(z) - \Delta_{x_i, y_i}(0)| \leq |z| \sup_{|q| \leq |z|} |\Delta'_{x_i, y_i}(q)|.$$

In order to estimate the above derivative, first note that

$$|e^{\phi_\omega(x_i)} - e^{\phi_\omega(y_i)}| \leq (e^{\phi_\omega(x_i)} + e^{\phi_\omega(y_i)}) H_\omega \rho^\alpha(x, y).$$

Therefore, for every complex q so that $|q| \leq 1$ and all $1 \leq i \leq d_\omega$,

$$|\Delta'_{x_i, y_i}(q)| \leq 8(1 + H_\omega)e^{\|u_\omega\|_\infty} \|u_\omega\| (e^{\phi_\omega(x_i)} + e^{\phi_\omega(y_i)}) \|g\| \rho^\alpha(x, y).$$

We conclude that for every $z \in \mathbb{C}$ with $|z| \leq 1$,

$$|A' - B'| \leq 8|z|(1 + H_\omega)e^{\|u_\omega\|_\infty} \|u_\omega\| \|g\| (\mathcal{L}_\omega \mathbf{1}(x) + \mathcal{L}_\omega \mathbf{1}(y)).$$

Now, since $g \in \mathcal{C}_\omega$ we have $\|g\| \leq r_\omega \inf g$, where $r_\omega = (1 + 2\kappa_\omega)$ and

$$\inf g \leq d_\omega^{-1} e^{\|\phi_\omega\|_\infty} \mathcal{L}_\omega g(t) = \kappa_{\theta\omega}^{-1} d_\omega^{-1} e^{\|\phi_\omega\|_\infty} B.$$

Using also that $\mathcal{L}_\omega \mathbf{1} \leq d_\omega e^{\|\phi_\omega\|_\infty}$ we see that

$$|A' - B'| \leq 16\kappa_{\theta\omega}^{-1}(1 + 2\kappa_\omega)(1 + H_\omega)e^{\|u_\omega\|_\infty + 2\|\phi_\omega\|_\infty} \|u_\omega\| |z| B.$$

We thus conclude that we can take

$$\zeta = \zeta_\omega \quad \text{and} \quad \varepsilon_1 = 16\kappa_{\theta\omega}^{-1}(1 + 2\kappa_\omega)(1 + H_\omega)e^{\|u_\omega\|_\infty + 2\|\phi_\omega\|_\infty} \|u_\omega\| |z|$$

in the above general result, which completes the proof of (6.5) in the case $s = s_{x,y,t,\kappa_\omega}$ since $c_\omega = 2\varepsilon_1(1 - \zeta)^{-1}$. \square

6.1.1. *A complex RPF theorem with effective rates.*

Proof of Theorem 49 in the setup of Section 2.2. Set

$$\delta(\omega) = 2c_0(\omega) (1 + \cosh(D(\omega)/2)) = 2E_\omega.$$

Then $\delta_\omega(z) = |z|\delta(\omega)$ and the assumptions in Theorem 49 insure that

$$A := \text{esssup} \left(\delta(\omega)(1 - e^{-D(\omega)})^{-1} \right) < \infty.$$

Hence the condition $\delta_\omega(z) \leq 1 - e^{-D(\omega)}$ holds true when $|z| \leq 1/A$. Relying on Theorem 80, proceeding as in the of the proof of the real RPF theorem (Theorem 47), we see that there is a constant r_0 so that \mathbb{P} -a.s. for every complex number z so that $|z| \leq r_0$ there is a triplet consisting of a nonzero complex random variable $\lambda_\omega(z)$, a random function $\hat{h}_\omega^{(z)} \in \mathcal{C}_{\omega, \mathbb{C}}$ and a random linear functional $\nu_\omega^{(z)} \in \mathcal{C}_{\omega, \mathbb{C}}^*$ so that $\nu_\omega^{(z)}(\mathbf{1}) = \nu_\omega(\hat{h}_\omega^{(z)}) = 1$,

$$(\mathcal{L}_\omega^{(z)})^* \nu_{\theta\omega}^{(z)} = \lambda_\omega(z) \nu_\omega^{(z)}$$

and for every $\mu \in \mathcal{C}_\omega$

$$\left\| \frac{(\mathcal{L}_\omega^{z,n})^* \mu}{\mu(\mathcal{L}_\omega^{z,n} \mathbf{1})} - \nu_\omega^{(z)} \right\| \leq M_\omega \tilde{\rho}_{\omega, n}$$

where $\tilde{\rho}(\omega) = \tanh(7D(\omega)/4)$. Moreover, for every $g \in \mathcal{C}_{\theta^{-n}\omega, \mathbb{C}}$ we have

$$(6.6) \quad \left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^{z,n} g}{\nu_\omega(\mathcal{L}_{\theta^{-n}\omega}^{z,n} g)} - \hat{h}_\omega^{(z)} \right\| \leq \sqrt{2} K_\omega \tilde{\rho}_{\theta^{-n}\omega, n}.$$

Since $\nu_\omega^{(z)}$ and $\hat{h}_\omega^{(z)}$ are uniform limits (in z) of analytic in z measurable functions they are analytic in z and measurable in ω . Similarly, also $\lambda_\omega(z)$ is analytic in z . Since $\nu_\omega^{(z)}(\mathbf{1}) = 1$ we conclude from (6.1) that $\|\nu_\omega^{(z)}\| \leq M_\omega$. Moreover, since $\nu_\omega(\hat{h}_\omega^{(z)}) = 1$ we conclude from (6.1) that $\|\hat{h}_\omega^{(z)}\| \leq 2\sqrt{2}K_\omega$. It is also clear that $\lambda_\omega(0) = \lambda_\omega$, $\nu_\omega^{(0)} = \nu_\omega$ and $\hat{h}_\omega^{(0)} = h_\omega$. To correct the fact that $\nu_\omega^{(z)}(\hat{h}_\omega^{(z)})$ might not equal 1 (notice that it does not vanish since $\nu_\omega^{(z)}$ belongs to the dual cone) let us define

$$h_\omega^{(z)} = \frac{\hat{h}_\omega^{(z)}}{\alpha_\omega(z)}$$

where $\alpha_\omega(z) = \nu_\omega^{(z)}(\hat{h}_\omega^{(z)})$. Notice that $\alpha_\omega(0) = 1$, $\alpha_\omega(z)$ is analytic in z and

$$|\alpha_\omega(z)| \leq \|\nu_\omega^{(z)}\| \|\hat{h}_\omega^{(z)}\| \leq 2\sqrt{2}M_\omega K_\omega.$$

Let us now obtain (3.13), which in particular will yield that $\mathcal{L}_\omega^{(z)}(h_\omega^{(z)}) = \lambda_\omega(z)h_{\theta\omega}^{(z)}$. Let us first prove a version of (3.13) for functions in the cone \mathcal{C}_ω . First, for every complex number z such that $|z| \leq r_0$ and $q \in \mathcal{C}_{\theta^{-n}\omega}$ we have

$$\lambda_{\theta^{-n}\omega, n}(z) \nu_{\theta^{-n}\omega}^{(z)}(q) = \nu_\omega^{(z)}(\mathcal{L}_{\theta^{-n}\omega}^{z,n} q).$$

Next, set

$$b_n(q, z) = b_n(\omega, q, z) = \frac{\nu_\omega(\mathcal{L}_{\theta^{-n}\omega}^{z,n} q)}{\nu_\omega^{(z)}(\mathcal{L}_{\theta^{-n}\omega}^{z,n} q)}.$$

Then,

$$\begin{aligned} & \left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^{z,n} q}{\lambda_{\theta^{-n}\omega, n}(z) \nu_{\theta^{-n}\omega}^{(z)}(q)} - h_\omega^{(z)} \right\| = \left\| b_n(q, z) \cdot \frac{\mathcal{L}_{\theta^{-n}\omega}^{z,n} q}{\nu_\omega(\mathcal{L}_{\theta^{-n}\omega}^{z,n} q)} - h_\omega^{(z)} \right\| \\ & \leq \left\| b_n(q, z) \left(\frac{\mathcal{L}_{\theta^{-n}\omega}^{z,n} q}{\nu_\omega(\mathcal{L}_{\theta^{-n}\omega}^{z,n} q)} - \hat{h}_\omega^{(z)} \right) \right\| + \left\| \left(b_n(q, z) - \frac{1}{\alpha_\omega(z)} \right) \hat{h}_\omega^{(z)} \right\| := I_1 + I_2 \end{aligned}$$

where we have used the above definition of $h_\omega^{(z)}$. Now, by (6.6) we have

$$|(b_n(q, z))^{-1} - \alpha_\omega(z)| = \left| \nu_\omega^{(z)} \left(\frac{\mathcal{L}_{\theta^{-n}\omega}^{z,n} q}{\nu_\omega(\mathcal{L}_{\theta^{-n}\omega}^{z,n} q)} \right) - \nu_\omega^{(z)}(\hat{h}_\omega^{(z)}) \right|$$

$$\leq \|\nu_\omega^{(z)}\| \left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^{z,n} q}{\nu_\omega(\mathcal{L}_{\theta^{-n}\omega}^{z,n} q)} - \hat{h}_\omega^{(z)} \right\| \leq \sqrt{2} M_\omega K_\omega \tilde{\rho}_{\theta^{-n}\omega, n}.$$

Hence, if n satisfies that $\sqrt{2} M_\omega K_\omega \tilde{\rho}_{\theta^{-n}\omega, n} < \frac{1}{2} |\alpha_\omega(z)|$ (which, since $z \rightarrow \alpha_\omega(z)$ is analytic and non-vanishing, is true \mathbb{P} -a.s. for every n large enough uniformly in z) then

$$(6.7) \quad \left| b_n(q, z) - \frac{1}{\alpha_\omega(z)} \right| \leq \frac{2\sqrt{2} M_\omega K_\omega \tilde{\rho}_{\theta^{-n}\omega, n}}{|\alpha_\omega(z)|^2} \leq |\alpha_\omega(z)|^{-1}.$$

Combing this with the upper bound $\|\hat{h}_\omega^{(z)}\| \leq 2\sqrt{2} K_\omega$ we conclude that for such n 's we have

$$I_2 \leq \frac{8M_\omega K_\omega^2 \tilde{\rho}_{\theta^{-n}\omega, n}}{|\alpha_\omega(z)|^2}.$$

Using now (6.6) together with (6.7) and that $|\alpha_\omega(z)| \leq 2\sqrt{2} M_\omega K_\omega$ we see that for n satisfying the above properties we have

$$I_1 \leq |b_n(q, z)| \sqrt{2} K_\omega \rho_{\theta^{-n}\omega, n} \leq 2 |\alpha_\omega(z)|^{-1} \left(\sqrt{2} K_\omega \tilde{\rho}_{\theta^{-n}\omega, n} \right).$$

By combining the above estimates on I_1 and I_2 we see that

$$\left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^{z,n} q}{\lambda_{\theta^{-n}\omega, n}(z) \nu_{\theta^{-n}\omega}^{(z)}(q)} - h_\omega^{(z)} \right\| \leq \left(2\sqrt{2} K_\omega |\alpha_\omega(z)|^{-1} + 8M_\omega K_\omega^2 |\alpha_\omega(z)|^{-2} \right) \tilde{\rho}_{\theta^{-n}\omega, n} := R(\omega, n, z).$$

Finally, using that $\|\nu_{\theta^{-n}\omega}^{(z)}\| \leq M_{\theta^{-n}\omega}$ we conclude that for every $g \in \mathcal{C}_{\theta^{-n}\omega, \mathbb{C}}$ we have

$$\left\| \frac{\mathcal{L}_{\theta^{-n}\omega}^{z,n} q}{\lambda_{\theta^{-n}\omega, n}(z)} - h_\omega^{(z)} \right\| \leq M_{\theta^{-n}\omega} R(\omega, n, z).$$

Now the proof of (3.13) is completed by using the reproducing property stated in Theorem 80 (iii). \square

6.2. Complex cones for properly expanding maps. In this section we will briefly explain how to prove Theorem 49 in the setup of Section 2.1. Since this is completed similarly to the proof in the setup of Section 2.2 (using ideas from [33, Ch.5]) we will formulate the results concerning complex cones without their proofs.

We suppose here that (2.4) holds true and that u_ω satisfies $v(u_\omega) \leq H_\omega$ with some H_ω so that

$$\gamma_\omega^{-\alpha} v(u_\omega) + H_\omega \leq \gamma_{\theta\omega}^\alpha - 1.$$

Then, by replacing ϕ_ω with $\phi_{\omega, t} = \phi_\omega + t u_\omega$ all the results concerning real cones hold true for $\phi_{\omega, t}$ when $t \in [-1, 1]$, with $Z_\omega = \gamma_\omega^{-\alpha} v(u_\omega) + H_\omega$ instead of H_ω and with $\|\phi_\omega\|_\infty + \|u_\omega\|_\infty$ instead of $\|\phi_\omega\|_\infty$.

We will need the following result, whose proof proceeds essentially as in [33, Ch. 5].

81. Theorem. (i) *The cones $\mathcal{C}_{\omega, \mathbb{C}}$ and their duals $\mathcal{C}_{\omega, \mathbb{C}}^*$ have bounded aperture: for all $g \in \mathcal{C}_{\omega, \mathbb{C}}$ and $\nu \in \mathcal{C}_{\omega, \mathbb{C}}^*$ we have*

$$\|g\| \leq 2\sqrt{2} K_\omega |\nu_\omega(g)| \quad \text{and} \quad \|\nu\| \leq M_\omega |\nu(\mathbf{1})|$$

where M_ω and K_ω were defined in Section 2.3.1.

(ii) *The cones $\mathcal{C}_{\omega, \mathbb{C}}$ are linearly convex, namely for every $g \notin \mathcal{C}_{\omega, \mathbb{C}}$ there exists $\mu \in \mathcal{C}_{\omega, \mathbb{C}}^*$ such that $\mu(g) = 0$.*

(iii) *The cones $\mathcal{C}_{\omega, \mathbb{C}}$ are reproducing: for every complex-valued function $g \in \mathcal{H}_\omega$ there are constants $c_1(g), c_2(g) > 0$ and functions $g_1, g_2 \in \mathcal{C}_\omega \subset \mathcal{C}_{\omega, \mathbb{C}}$ so that $g = g_1 - c_1(g) + i(g_2 - c_2(g))$ and*

$$\|g_1\| + c_1(g) + \|g_2\| + c_2(g) \leq 2r_\omega \|g\|$$

where $r_\omega = 8 \left(1 + \frac{2}{\gamma_\omega^\alpha} \right) \leq 24$.

(iv) *Let $c_0(\omega)$ be defined as in Section 2.3.1 and let*

$$\tilde{D}(\omega) = \gamma_{\theta\omega}^\alpha + 2 \ln \left(\frac{1 + \tilde{q}(\omega)}{1 - \tilde{q}(\omega)} \right)$$

where $\tilde{q}(\omega) = \frac{\tilde{H}_\omega + 1}{\gamma_{\theta_\omega}^\alpha}$, Then, if $\delta_\omega(z) \leq 1 - e^{-\tilde{D}(\omega)}$ we have that

$$\mathcal{L}_\omega^{(z)} \mathcal{C}_{\omega, \mathbb{C}} \subset \mathcal{C}_{\theta_\omega, \mathbb{C}}$$

and the Hilbert diameter of the image with respect to the complex projective metric corresponding to the cone $\mathcal{C}_{\theta_\omega, \mathbb{C}}$ does not exceed $7D(\omega)$, with $D(\omega)$ defined in Section 2.3.1.

Relying on Theorem 81 the proof of Theorem 49 in the setup of Section 2.1 proceeds exactly as in Section 6.1.1.

82. Remark. Notice that $\tilde{D}(\omega) \geq 1$ and so $1 - e^{-\tilde{D}(\omega)} \geq 1 - e^{-1}$. Hence, if

$$E_\omega = c_0(\omega) \left(1 + \cosh(\tilde{D}(\omega)/2)\right)$$

is a bounded random variable then there is a constant $r_0 > 0$ so that the condition $\delta_\omega(z) \leq 1 - e^{-\tilde{D}(\omega)}$ holds true when $|z| \leq r_0$. Notice also that $\cosh(\tilde{D}(\omega)/2) \leq e^{\tilde{D}(\omega)/2}$.

6.3. The “normalized” complex operators. In this section we will prove (3.14) relying on (3.13). Let us consider the operators $L_\omega^{(z)}$ given by

$$L_\omega^{(z)} g = \frac{\mathcal{L}_\omega^{(z)}(gh_\omega)}{h_{\theta_\omega} \lambda_\omega}.$$

Then

$$\begin{aligned} \left\| \frac{L_\omega^{z,n} g}{\lambda_{\omega,n}(z)} - \mu_\omega^{(z)}(g) \bar{h}_\omega^{(z)} \right\| &= \left\| \frac{\mathcal{L}_\omega^{z,n}(gh_\omega)}{\lambda_{\omega,n}(z) h_{\theta_\omega}^{n_\omega}} - \nu_\omega^{(z)}(gh_\omega) \frac{h_{\theta_\omega}^{(z)}}{h_{\theta_\omega}^{n_\omega}} \right\| \\ &\leq 3 \left\| \frac{1}{h_{\theta_\omega}^{n_\omega}} \right\| \left\| \frac{\mathcal{L}_\omega^{z,n}(gh_\omega)}{\lambda_{\omega,n}(z)} - \nu_\omega^{(z)}(gh_\omega) h_{\theta_\omega}^{(z)} \right\|. \end{aligned}$$

Now, as in Section 5.3 we have $3\|1/h_\omega\| \leq U_\omega$ and thus (3.14) follows from (3.13).

REFERENCES

- [1] J Aaronson, M Denke, *Local limit theorems for partial sums of stationary sequences generated by Gibbs–Markov maps*, Stochastics and Dynamics, Vol. 01, No. 02, pp. 193–237 (2001).
- [2] R. Aimino, M. Nicol, S. Vaienti, *Annealed and quenched limit theorems for random expanding dynamical systems*, Probab. Theory Relat. Fields 162, 233–274 (2015).
- [3] J. F. Alves, W. Bahsoun and R. Ruziboev, *Almost sure rates of mixing for partially hyperbolic attractors*, Journal of Differential Equations, Volume 311, 2022, Pages 98–157.
- [4] , J. F. Alves, W. Bahsoun, R. Ruziboev and P. Varandas, *Quenched decay of correlations for nonuniformly hyperbolic random maps with an ergodic driving system*, <https://arxiv.org/abs/2205.13424>.
- [5] J. Atnip, J. G. Froyland, C. González-Tokman and S. Vaienti. *Thermodynamic Formalism for Random Weighted Covering Systems*. Commun. Math. Phys. 386, 819–902 (2021).
- [6] J. Atnip, J. G. Froyland, C. González-Tokman and S. Vaienti. *Equilibrium states for non-transitive random open and closed dynamical systems*, ETDS (2022), <https://doi.org/10.1017/etds.2022.68>
- [7] V. Baladi, M. Benedicks and V. Maume-Deschamps. *Almost sure rates of mixing for i.i.d. unimodal maps*. Ann. Sci. Éc. Norm. Supér. (4) 35(1) (2002), 77–126
- [8] W. Bahsoun, C. Bose and M. Ruziboev. *Quenched decay of correlations for slowly mixing systems*, Trans. Amer. Math. Soc. 372(9) (2019), 6547–6587
- [9] P. Billingsley, *Convergence of Probability Measures*, Wiley, New York, 1968.
- [10] G. Birkhoff, *Extension of Jentzsch’s theorem*, Trans. A.M.S. 85 (1957), 219–227.
- [11] J. R. Blum, D.L Hanson, L.H Koopmans, *On the Strong Law of Large Numbers for a Class of Stochastic Processes*, Z. Wahrscheinlichkeitstheorie 1963.
- [12] R. Bowen, *R. Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Lecture Notes in Mathematics, volume 470, Springer Verlag, 1975
- [13] R. Bradley, *Basic properties of strong mixing conditions. A survey and some open questions*, Probability Surveys, Vol. 2 (2005) 107–144.
- [14] J. Buzzi, *Exponential decay of correlations for random Lasota–Yorke maps*, Commun. Math. Phys. 208, pages25–54 (1999).

- [15] C. Castaing and M. Valadier, *Convex analysis and measurable multifunctions*, Lecture Notes Math., vol. 580, Springer, New York, 1977.
- [16] A. Castro, P. Varandas, *Equilibrium states for non-uniformly expanding maps: decay of correlations and strong stability*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 30 (2) (2013), pp. 225–249
- [17] C. Cuny, F. Merlevède *Strong Invariance Principles with Rate for “Reverse” Martingale Differences and Applications*. J Theor Probab 28, 137–183 (2015)
- [18] A. Dembo, O. Zeitouni, *Large deviations techniques and applications*, Springer (2010).
- [19] P. Doukhan, *Mixing, properties and examples*, Springer (1994).
- [20] D. Dragičević, G. Froyland, C. Gonzalez-Tokman and S. Vaienti, *Almost Sure Invariance Principle for random piecewise expanding maps*, Nonlinearity 31 (2018), 2252–2280.
- [21] D. Dragičević, G. Froyland, C. Gonzalez-Tokman and S. Vaienti, *A spectral approach for quenched limit theorems for random expanding dynamical systems*, Comm. Math. Phys. 360 (2018), 1121–1187.
- [22] D. Dragičević, G. Froyland, C. Gonzalez-Tokman and S. Vaienti, *A spectral approach for quenched limit theorems for random hyperbolic dynamical systems*, Trans. Amer. Math. Soc. 373 (2020), 629–664.
- [23] D. Dragičević, J. Sedro, *Quenched limit theorems for expanding on average cocycles*, preprint arXiv:2105.00548.
- [24] D. Dragičević, J. Sedro and Y. Hafouta, *A vector-valued almost sure invariance principle for random expanding on average cocycles*, preprint arXiv:2108.08714.
- [25] D. Dragičević and Y. Hafouta, *Limit theorems for random expanding or Anosov dynamical systems and vector-valued observables*, Ann. Henri Poincaré 21, 3869–3917.
- [26] D. Dragičević and Y. Hafouta, *Almost Sure Invariance Principle for Random Distance Expanding Maps with a Nonuniform Decay of Correlations*, In: Pollicott, M., Vaienti, S. (eds) Thermodynamic Formalism. Lecture Notes in Mathematics, vol 2290. Springer, Cham.
- [27] L. Dubois, *Projective metrics and contraction principles for complex cones*, J. London Math. Soc. 79 (2009), 719–737.
- [28] L. Dubois, *An explicit Berry-Esséen bound for uniformly expanding maps on the interval*, Israel J. Math. 186 (2011), 221–250.
- [29] Z. Du. *On mixing rates for random perturbations*. PhD Thesis, National University of Singapore, 2015.
- [30] W. Feller, *An introduction to probability theory and its applications*, Vol. II., 2d edition, John Wiley & Sons, Inc., New York-London-Sydney, 1971.
- [31] N. Haydn, Y. Psiloyenis *Return times distribution for Markov towers with decay of correlations*, Nonlinearity 27(6), 1323 (2014)
- [32] Y. Hafouta, Yu. Kifer, *Berry–Esseen type estimates for nonconventional sums*, Stoch. Proc. App. Volume 126, Issue 8, August 2016, Pages 2430–2464.
- [33] Y. Hafouta, Yu. Kifer *Nonconventional limit theorems and random dynamics*, World Scientific (2018).
- [34] Y. Hafouta, *Limit theorems for random non-uniformly expanding or hyperbolic maps with exponential tails*, Annales Henri Poincaré 23, 293–332 (2022)
- [35] Y. Hafouta, *Convergence rates in the functional CLT for α mixing triangular arrays*, <https://arxiv.org/pdf/2107.02234.pdf>.
- [36] H. Hennion and L. Hervé, *Limit Theorems for Markov Chains and Stochastic Properties of Dynamical Systems by Quasi-Compactness*, Lecture Notes in Mathematics vol. 1766, Springer, Berlin, 2001.
- [37] I. A. Ibragimov, Yu. V. Linnik *Independent and stationary sequences of random variables*, Wolters-Noordhoff Publishing, Groningen, 1971. 443 pp.
- [38] Yu. Kifer, *Perron-Frobenius theorem, large deviations, and random perturbations in random environments*, Math. Z. 222(4) (1996), 677–698.
- [39] Yu. Kifer, *Limit theorems for random transformations and processes in random environments*, Trans. Amer. Math. Soc. 350 (1998), 1481–1518.
- [40] Yu. Kifer, *Thermodynamic formalism for random transformations revisited*, Stoch. Dyn. 8 (2008), 77–102.
- [41] C. Liverani, *Decay of correlations*, Ann. Math. 142 (1995), 239–301.
- [42] V. Mayer, B. Skorulski and M. Urbański, *Distance expanding random mappings, thermodynamical formalism, Gibbs measures and fractal geometry*, Lecture Notes in Mathematics, vol. 2036 (2011), Springer.
- [43] H.H. Rugh, *Cones and gauges in complex spaces: Spectral gaps and complex Perron-Frobenius theory*, Ann. Math. 171 (2010), 1707–1752.
- [44] M. Stadlbauer, S. Suzuki, P. Varandas *Thermodynamic formalism for random non-uniformly expanding maps*, Commun. Math. Phys. 385, 369–427 (2021)
- [45] Y. Su, *Random Young towers and quenched limit laws*. Ergodic Theory and Dynamical Systems, 1–33. doi:10.1017/etds.2021.164.

Email address: yeor.hafouta@mail.huji.ac.il, yhafouta@umd.edu