

Pole-skipping in holographic theories with bosonic fields

Diandian Wang and Zi-Yue Wang

Department of Physics, University of California, Santa Barbara, CA 93106

We study the phenomenon of pole-skipping in holographic CFTs dual to diffeomorphism invariant theories containing an arbitrary number of bosonic fields in the large N limit. Defining a weight to organize the bulk equations of motion and field components, a set of general pole-skipping conditions are derived. In particular, the frequencies simply follow from general covariance and weight matching. Relating the highest-weight pole-skipping frequency to an exponential growth rate, i.e., the Lyapunov exponent, we show that the chaos bound is generally violated in the presence of finitely many higher spin fields, consistent with existing evidence. In the absence of such pathological fields, we show that the energy density Green's function has its highest-weight pole-skipping happening at a location related to the OTOC for arbitrary higher-derivative gravity, with a Lyapunov exponent saturating the chaos bound and a butterfly velocity matching that extracted from a shockwave calculation. We also suggest a physical explanation for this matching by obtaining the shockwave metric from a regularized limit of the metric perturbation at the skipped pole.

I. INTRODUCTION

The out-of-time-order correlator (OTOC), an important quantity containing characteristics of chaos, can be calculated holographically in a shockwave spacetime [1–5]. For a localized perturbation to a maximally chaotic system at temperature T , the OTOC between a perturbation W at $x = t = 0$ and a probe operator V at a later time t behaves as

$$\langle V(x, t) W V(x, t) W \rangle \sim 1 - e^{\lambda_L(t - t_* - |x|/v_B)}, \quad (1)$$

where t_* is called the scrambling time. This defines the Lyapunov exponent, λ_L , and the butterfly velocity, v_B . For classical bulk gravitational theories, λ_L saturates the chaos bound $\lambda_L \leq 2\pi T$ [6], so they are said to be maximally chaotic. The butterfly velocity, however, depends on the theory [2, 7–11].

More recently, it was discovered that the quantities λ_L and v_B may already show up in features of a two-point function, the energy density retarded Green's function, through a phenomenon called *pole-skipping* [12–14]. It was first found numerically for pure Einstein gravity [12] and later studied analytically for Einstein gravity with matter [14]. See also [15–18] for holographic and [19–22] for boundary studies.

The retarded Green's function is the relation between a source and its response. Holographically, the Green's function of an operator dual to a bulk dynamic field X (suppressing indices) is given by [23, 24]

$$G_R(\omega, k) = \left(\lim_{r \rightarrow \infty} \frac{\Pi(r; \omega, k)|_{X_R}}{X_R(r; \omega, k)} \right) \Big|_{X_0=0}, \quad (2)$$

where X_R is the bulk solution satisfying Dirichlet boundary condition $X_R \rightarrow X_0$ at infinity and ingoing wave boundary condition at the horizon, and Π is its conjugate variable in a radial foliation. In terms of an asymptotic expansion, it is proportional to the ratio between the coefficient of the normalizable falloff and that of the non-normalizable falloff. A quasinormal mode, by definition, does not have a non-normalizable divergence, so

the poles of the Green's function are identified with the quasinormal spectrum.

Generically, X_R is uniquely determined from X_0 , and G_R is therefore well-defined. However, a would-be pole can sometimes get multiplied by a zero, resulting in an ill-defined limit. This happens at a special frequency and momentum,

$$\omega = i\lambda_L, \quad k = \frac{i\lambda_L}{v_B}, \quad (3)$$

where λ_L and v_B are the Lyapunov exponent and the butterfly velocity extracted from a holographic OTOC calculation (1) in Einstein gravity minimally coupled to a large class of matter fields [12, 14].

To explain this universality, [14] discovered a feature of Einstein's equation at the horizon. Expanding metric perturbations around a stationary planar black hole in terms of Fourier modes, a particular component of Einstein's equation evaluated at the horizon was found to be trivial at (3) so that there exists one fewer constraint. This implies an extra degree of freedom of the ingoing modes and consequently an ambiguity in the bulk solution X_R and in turn the Green's function G_R . We will re-assess this universality in the presence of matter. In particular, we show that, even in Einstein gravity, matter fields can enter the expression for the butterfly velocity, but if one restricts the weight (or spin) and regularity of the matter, the butterfly velocity becomes a function of the metric only and thus in this sense more universal.

Later it was discovered that pole-skipping happens more generally at other locations and for other types of Green's functions [16, 17, 25–29]. See also [30, 31] for higher-derivative corrections and [32] for a zero-temperature example. However, unlike the one at (3), the other skipped poles are unrelated to chaos.

We put these in the same framework by considering general diffeomorphism invariant bulk theories with matter fields that are not necessarily minimally coupled. For simplicity, we only consider bosonic fields and leave a discussion of fermionic fields to the last section. By defining a weight, we can separate the equations of motion into

different groups and evaluate them in a given order. This allows us to find the frequencies of the skipped poles and the corresponding momenta in general. This is done in Section II. Furthermore, we point out pathology caused by the presence of higher-weight pole-skipping frequencies and use it to justify the removal of a bounded tower of higher spin fields from consideration in the remaining sections.

In Section III, it is shown that, for general higher-derivative gravitational theories, the butterfly velocity can be obtained from the highest-weight equation of motion, and, for matter fields that are regular enough at the horizon, it agrees with the butterfly velocity obtained via a shockwave calculation. This generalizes the matching for Gauss-Bonnet gravity and Einstein gravity with a string theory correction at $O(\alpha'^3)$ [33]. We also try to establish a more physical connection between pole-skipping and chaos in the same section. By regularizing the metric perturbation at the chaotic skipped pole with a Gaussian distribution in the frequency Fourier space, we obtain a metric that is regular at the horizon. Extending it to a Kruskal-Szekeres coordinate patch and taking the regulator away, we show that this metric perturbation localizes to the past horizon in a distributional sense, just like the shockwave metric. We end with a summary and a discussion of potential future directions in Section IV.

II. GENERAL POLE-SKIPPING CONDITIONS

The metric for a general stationary planar black hole can be written in ingoing Eddington-Finkelstein coordinates as

$$ds^2 = -f(r)dv^2 + 2dvdr + h(r)dx^i dx^i, \quad (4)$$

where $f(r_0) = 0$ at the horizon $r = r_0$ and $i = 1, \dots, d$. The non-vanishing Christoffel components are given by

$$\begin{aligned} \Gamma_{vv}^v &= \frac{1}{2}f', & \Gamma_{ij}^v &= -\frac{1}{2}h'\delta_{ij}, & \Gamma_{vr}^r &= -\frac{1}{2}f', \\ \Gamma_{rj}^i &= \frac{h'}{2h}\delta_j^i, & \Gamma_{vv}^r &= \frac{1}{2}ff', & \Gamma_{ij}^r &= -\frac{1}{2}fh'\delta_{ij}. \end{aligned} \quad (5)$$

For simplicity, we assume that background matter fields are stationary, isotropic and homogeneous in x^i , and regular at both past and future horizons, like the metric.

Now, in this coordinate system, we introduce a *weight* for any tensor component as the number of lower v -indices minus that of lower r -indices, where an upper v is considered a lower r and vice versa. As a consequence of regularity at the bifurcate surface and boost symmetry, any background tensor component (ones constructed from the stationary background metric and matter fields) with positive weight needs to vanish at the horizon [34]. Note that we do not require perturbations to be regular at the bifurcate surface and therefore they do not need to have this property. To describe ingoing quasinormal

modes at the horizon, for any dynamic field X , we expand its perturbation around the stationary background in the Fourier space as

$$\delta X(r, v, x) = \delta X(r) e^{-i\omega v + ikx}. \quad (6)$$

For Einstein gravity, writing Einstein's equation as $E_{\mu\nu} = T_{\mu\nu}$, a particular component under perturbation, δE_{ν}^r , is proportional to

$$\left(k^2 - i\frac{d}{2}\omega h'\right) \delta g_{vv} + (\omega - i2\pi T) [\omega \delta g_{ii} + 2k_i \delta g_{vi}]. \quad (7)$$

On the horizon, for matter perturbations that are regular enough, the stress tensor component $\delta T_{\nu}^r = 0$ [14], and prefactors in δE_{ν}^r can be tuned to zero by choosing (3). As a consequence, Einstein's equation provides one fewer constraint, which serves as an explanation for the universal behaviour of the energy density Green's function with low-spin matter fields coupled to Einstein gravity [14].

Now consider an arbitrary diffeomorphism invariant theory defined with a local action $S = S_g + S_M$ where the gravitational part S_g is given by

$$S_g = \int d^{d+2}x \sqrt{-g} \mathcal{L}(g, R, \nabla, \Phi), \quad (8)$$

and S_M is part of the action with only minimally coupled matter fields, artificially separated from the rest for later convenience. Here \mathcal{L} can be an arbitrary function of the metric, g , and an arbitrary number of bosonic matter fields collectively denoted as Φ . More specifically, \mathcal{L} can be written as a sum of contractions between an arbitrary number of the metric, curvature tensors, matter fields, and an arbitrary number of covariant derivatives of them.

The metric equation of motion is defined as

$$E_{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\delta S_g}{\delta g^{\mu\nu}} = -\frac{2}{\sqrt{-g}} \frac{\delta S_M}{\delta g^{\mu\nu}} = T_{\mu\nu}. \quad (9)$$

The remaining equations of motion are given by $\delta S / \delta \Phi = 0$, indices suppressed. Now, to obtain (2), the idea is to perturb the dynamical fields and apply the equations of motion everywhere. However, it turns out sufficient to consider the near horizon expansion of all perturbations in order to study pole-skipping. For readability, we introduce the following compact notation: we use $\delta \mathcal{E} = 0$ to denote collectively all the perturbed equations of motions and their radial derivatives (∇_r) evaluated on the horizon. These are essentially the coefficients of a near-horizon Taylor expansion. We further define $\delta \mathcal{E}_p^\alpha = 0$ as the α -th independent equation with weight p evaluated on the horizon. We will also suppress the index α when understood, i.e., $\delta \mathcal{E}_p$ should be read as a vector with $|\delta \mathcal{E}_p|$ components. Similarly, we collect perturbations of all dynamics fields (including both the metric and matter) and their radial derivatives with weight q into $\delta \mathcal{X}_q$ (all evaluated on the horizon). For example, $\delta \mathcal{X}_2 = (\delta g_{vv}, \nabla_r \delta B_{vvv}, \dots)$ and $\delta \mathcal{X}_0 = (\delta g_{ij}, \nabla_r \delta g_{vi}, \nabla_r \nabla_r \delta g_{vv}, \delta A_i, \nabla_r \delta A_v, \dots)$.

With these definitions, we can now write

$$\delta\mathcal{E}_p = \sum_q \mathcal{M}_{p,q}(\omega, k) \delta\mathcal{X}_q, \quad (10)$$

where each $\mathcal{M}_{p,q}(\omega, k)$ is a matrix of size $|\delta\mathcal{E}_p| \times |\delta\mathcal{X}_q|$. To arrive at this form, first commute all ∇_r 's through ∇_i 's and ∇_v 's to the rightmost location before substituting the Fourier expansion and evaluating the ∇_i 's and ∇_v 's. By definition, the radial derivatives are then absorbed into $\delta\mathcal{X}_q$. For concreteness, we also commute all ∇_i to the left of ∇_v .

We now prove a useful property that, for $p > q$,

$$\mathcal{M}_{p,q}(\omega, k) \propto [\omega - (p-1)\omega_0] \dots [\omega - q\omega_0], \quad (11)$$

where $\omega_0 = i2\pi T = if'(r_0)/2$.

To begin with, notice that

$$\mathcal{M}_{p,q} = \sum_{k,l} F_{p,q,k,l}(g, R, \nabla, \Phi)(\nabla_i)^k (\nabla_v)^l \quad (12)$$

before substituting the Fourier expansion, where $F_{p,q,k,l}$ is some c -number tensor component constructed out of g, R, ∇ and Φ , such as $R_{virj} A^\mu \nabla_\mu \phi$, evaluated on the horizon of the background configuration. Next, notice that the only way to raise weight is with ∇_v because any background tensor with positive weight vanishes on the horizon. Therefore, to raise the weight of $\delta\mathcal{X}_q$ to that of $\delta\mathcal{E}_p$, one needs at least $(p-q)$ derivatives in v . From (5), it is straightforward to show that, on the horizon,

$$\nabla_v T \propto \left(\partial_v - \frac{n}{2} f'(r_0) \right) T \quad (13)$$

for a general tensor component T with weight n . As an example, for a tensor $T_{\mu\nu\rho}$, the weight zero component T_{vri} behaves as

$$\begin{aligned} \nabla_v T_{vri} &= \partial_v T_{vri} - \Gamma_{vv}^\alpha T_{\alpha ri} - \Gamma_{vr}^\alpha T_{v\alpha i} - \Gamma_{vi}^\alpha T_{vr\alpha} \\ &= \partial_v T_{vri} - \Gamma_{vv}^v T_{vri} - \Gamma_{vr}^r T_{vri} \\ &= [\partial_v - f'(r_0)/2 + f'(r_0)/2] T_{vri} \\ &= \partial_v T_{vri}, \end{aligned} \quad (14)$$

where the second line follows from the vanishing of most Christoffel components on the horizon. Therefore, (11) shows the minimum number of such factors it has to contain for the weight to match.

We now discuss the general conditions for pole-skipping. We take as an assumption that pole-skipping happens whenever an equation of motion becomes trivial [35]. Suppose the highest weight of $\delta\mathcal{X}$ is q_0 , then the highest weight of $\delta\mathcal{E}$ is also q_0 (since the action, being a scalar, has weight zero). Consequently, for any positive integer s , once we set

$$\omega = (q_0 - s)\omega_0, \quad (15)$$

all $\mathcal{M}_{p,q}(\omega, k)$ with $p > q \geq q_0 - s$ are then set to zero (assuming they are not all automatically zero). Now consider the square matrix

$$M_s(k) \equiv \begin{pmatrix} \mathcal{M}_{q_0, q_0} & \dots & \mathcal{M}_{q_0, q_0-s+1} \\ \dots & \dots & \dots \\ \mathcal{M}_{q_0-s+1, q_0} & \dots & \mathcal{M}_{q_0-s+1, q_0-s+1} \end{pmatrix} \quad (16)$$

where (15) has been substituted. The full set of equations of motion $\delta\mathcal{E}_p, \forall p$, does not determine $\delta\mathcal{X}_q, \forall q$, when

$$\det M_s(k) = 0. \quad (17)$$

The equations (15) and (17) are therefore the generalized pole-skipping conditions (for any given $s \geq 1$), assuming the second one has solutions. If the theory has a highest spin field with bounded spin l , then $q_0 = l$ and the pole-skipping frequencies are $(l-s)\omega_0$, consistent with observations made in [19, 29]. The second condition is a polynomial equation for k , and the roots are then the pole-skipping momenta, which could be more than one. The order of the polynomial increases with the size of the matrix, and therefore there will be generically more pole-skipping points at larger s (lower ω).

The first pole skipping happens at $s = 1$ at frequency $\omega = (q_0 - 1)\omega_0 = i(q_0 - 1)2\pi T$. Suppose there exists an equation of motion with e.g. three lower v -indices. In that case, there will be a skipped pole at $2\omega_0 = i4\pi T$, and the field perturbation (6) will grow like $\exp(4\pi Tt)$, corresponding to a Lyapunov exponent that is twice maximally chaotic. On this ground, we expect (finitely many) higher spin fields to violate the chaos bound, consistent with the findings of [19, 36]. This also adds to the existing pathology of such fields caused by causality violation [37]. Notice, however, that equations of motion for fields with no dynamics automatically have $\mathcal{M}_{p,q}(\omega, k) = 0$ for $p > q$ due to the nonappearance of ∇_v , so they do not become trivial from non-trivial; therefore, they do not violate the chaos bound, in agreement with [36] where pure AdS₃ higher spin gravity was exempt from their argument for bound violation.

If $q_0 = 2$, which is the case for an arbitrary metric theory coupled to matter fields of spin no larger than two, then the bound is satisfied and in fact saturated. We will discuss this further in the next section.

For $q_0 < 2$, such as a scalar or vector field without gravitational backreaction, there is no growing mode and therefore no relation to chaos, but an infinite number of skipped poles still exist and constrain the structure of Green's functions [17].

III. CONNECTION TO SHOCKWAVE

We now discuss the case $q_0 = 2$, although the argument here should generalize to higher weights upon a more careful study of higher spin shockwaves. This includes arbitrary higher-derivative gravitational theories coupled to scalar, vector or form fields. These have a skipped pole at $\omega = i2\pi T$, exhibiting maximal chaos. We now show that the corresponding butterfly velocity matches that obtained from the OTOC.

In this case, the only dynamic field with weight 2 is $\delta\mathcal{X}_2 = \delta g_{vv}$, and the corresponding equation of motion is $\delta\mathcal{E}_2 = \delta E_{vv} - \delta T_{vv} = 0$. The perturbation to the stress tensor component δT_{vv} does not necessarily vanish, but

δT_v^r ($= \delta T_{vv} - T_{rv} \delta g_{vv}$) does vanish for matter fields regular on the horizon [14]. We will make this restriction in order to compare results with OTOC: the metric shockwave also has vanishing δT_v^r . To relax this, one would need to first generalize the shockwave solution to allow for more singular matter configurations, which is outside the scope of this paper. Therefore, the pole-skipping conditions with $s = 1$ are given by

$$\omega = \omega_0, \quad \det M_1 = \frac{\delta E_v^r}{\delta g_{vv}} = 0. \quad (18)$$

This gives a polynomial equation for k with only even powers (by symmetry). In cases where the polynomial is of quartic order or higher, one can take the view that all corrections to Einstein gravity should be treated perturbatively so only the roots continuously connected to Einstein gravity are physical. But as we will see, the matching is evident without a perturbative treatment.

From above, we have

$$\delta E_v^r = \sum_{k,l} F_{2,2,k,l}(g, R, \nabla, \Phi)(\nabla_i)^k (\nabla_v)^l \delta g_{vv}, \quad (19)$$

where $F_{p,q,k,l}$ was defined in (12). Evaluating this on the metric (4) and defining new coefficients $H_{k,l}$, it takes the non-covariant form

$$\delta E_v^r = \sum_{k,l} H_{k,l}(f, h, \partial_r, \Phi)(\partial_i)^k (\partial_v)^l \delta g_{vv}. \quad (20)$$

The non-trivial statement that no ∂_r acts on δg_{vv} and none of the other components such as δg_{vi} can appear follow directly from the weight argument [38]. As an example, consider the Einstein gravity equation of motion (7) studied in [14]. Since δg_{ij} has weight zero, it has to pick up a factor of ω to get to weight one and then a factor of $(\omega - \omega_0)$ to get to weight two, similarly for δg_{vi} which only needs to raise its weight by one. Another simplification in Einstein gravity is due to the fact of it being two-derivative. It is not possible for (7) to contain a term like for example $\partial_r \delta g_{vi}$: this quantity has weight zero and therefore needs two v -derivatives to go to two, but it already has one derivative itself.

To compare this with the shockwave calculation, we move to Kruskal-Szekeres coordinates, defined via

$$U = -e^{-f'(r_0)(v-2r_*)/2}, \quad V = e^{f'(r_0)v/2}, \quad (21)$$

where $dr_*/dr = 1/f(r)$. Then $UV = -e^{f'(r_0)r_*}$, and the metric is given by

$$ds^2 = 2A(UV)dUdV + B(UV)dx^i dx^i, \quad (22)$$

where

$$A(UV) = \frac{2}{f'(r_0)^2} \frac{f(r)}{UV}, \quad B(UV) = h(r). \quad (23)$$

In general higher derivative gravity and for a shockwave along $V = 0$, the only non-trivial component of δE_v^μ

perturbed by a local source is δE_V^U [11]. For a general perturbation, δg_{vv} , translating to ingoing Eddington-Finkelstein coordinates, this component is given by

$$\delta E_V^U = \frac{U}{V} \left(\frac{2}{f(r)} \delta E_v^r + \delta E_r^r - \delta E_v^v - \frac{f(r)}{2} \delta E_r^v \right). \quad (24)$$

Compared to the first term, others are suppressed with extra factors of $f(r)$, so they vanish when evaluated on the horizon. Similarly, $\delta T_V^U \propto \delta T_r^r$, but recall that this vanishes for regular matter configurations. Therefore,

$$\begin{aligned} 0 &= \delta E_V^U = \frac{2UV}{f(r)} \frac{1}{V^2} \sum_{k,l} H_{k,l}(\partial_i)^k (\partial_v)^l \delta g_{vv} \\ &= \frac{2UV}{f(r)} \frac{1}{V^2} \sum_{k,l} H_{k,l}(\partial_i)^k \left(\frac{2}{f'(r_0)} V \partial_V \right)^l \delta g_{vv} \\ &= \frac{2UV}{f(r)} \sum_{k,l} H_{k,l}(\partial_i)^k \left(\frac{2}{f'(r_0)} (V \partial_V + 2) \right)^l \frac{\delta g_{vv}}{V^2} \\ &= \frac{2UV}{f(r)} \sum_{k,l} \tilde{H}_{k,l}(\partial_i)^k (V \partial_V)^l \frac{\delta g_{vv}}{V^2} \\ &= \frac{1}{A} \sum_{k,l} \tilde{H}_{k,l}(\partial_i)^k (V \partial_V)^l \delta g_{VV}, \end{aligned} \quad (25)$$

where we used the transformation $\partial_v = \frac{2}{f'(r_0)} V \partial_V$ in going to the second line, and a trick

$$\frac{1}{V^2} V \partial_V = (V \partial_V + 2) \frac{1}{V^2} \quad (26)$$

in going to the third line. The fourth line follows from a reorganization of the sum with new coefficients $\tilde{H}_{k,l}$ and the last line follows from

$$\delta g_{VV}(V, x) = \frac{4\delta g_{vv}(v, x)}{f'(r_0)^2 V^2}. \quad (27)$$

The special thing about $\omega = \omega_0$ is that,

$$\delta g_{vv} \sim e^{-i\omega_0 v} = e^{-\frac{i2\omega_0}{f'(r_0)} \log V} = V, \quad (28)$$

and therefore

$$\delta g_{VV}(V, x) \sim \frac{1}{V} e^{-ikx}. \quad (29)$$

Compare this with a linearized shockwave perturbation

$$\delta g_{VV} \sim \delta(V) e^{-\mu x}, \quad (30)$$

where $\mu = 2\pi T/v_B$ upon using $\delta E_V^U = 0$ (outside of a localized source in the transverse directions). Noticing that $\delta(V)$ has the same distributional behavior as $1/V$ under $V \partial_V$ [12], e.g., $V \delta'(V) = -\delta(V)$ and $V d(1/V)/dV = -1/V$, it follows that $k = i2\pi T/v_B$ upon using (25) for the perturbation (29), thereby extending (3) to general higher-derivative gravity and hence some of the results of [33].

Given the similarity between $1/V$ and $\delta(V)$ and the role this similarity plays in establishing the equivalence of these two calculations of the butterfly velocity, it is natural to wonder whether there is a more direct connection between them. An immediate obstacle is the divergence of the function $1/V$ at the past horizon $V = 0$. We mitigate this problem with an unnormalized regularization of the Fourier space delta function along the real frequency line:

$$\int d\xi \delta(\xi) \rightarrow \int_{-\infty}^{\infty} d\xi e^{-\xi^2/a}, \quad (31)$$

giving rise to a mode

$$\delta g_{vv} = \sqrt{\pi a} e^{-\frac{a}{4}(\lambda_L v)^2} e^{\lambda_L v}. \quad (32)$$

To compare with the shockwave metric (30), we convert this to Kruskal-Szekeres coordinates. Using (27),

$$\delta g_{VV} = \begin{cases} 0, & V < 0 \\ \frac{\sqrt{\pi a}}{\lambda_L^2} \frac{1}{V} e^{-a(\log V)^2/4}, & V \geq 0 \end{cases} \quad (33)$$

where we have used the fact that the perturbation vanishes exactly behind the past horizon. This function is finite and integrates to a constant for finite a , and it vanishes everywhere off the horizon as $a \rightarrow 0$. It therefore behaves as a regularized delta function. Taking the regulator away, this becomes a shockwave localized at $V = 0$.

We should note that the need for the unnormalized regulator arises from the need to remove the divergence. Alternatively, one can use a normalized delta function regulator and remove the divergence at the end using a subtraction not unlike the minimal subtraction in dimensional regularization.

IV. DISCUSSION

We have defined a quantity called weight to organize bulk equations of motion and exploited its convenience to show that pole-skipping happens in holographic CFTs dual to quite general diffeomorphism invariant bulk theories. As a result, the pole-skipping frequencies show up at $(q_0 - s)\omega_0$ for all $s \in \mathbb{Z}^+$, where $\omega_0 = i2\pi T$, and q_0 is defined as the weight of the highest-weight object. In particular, a theory that has a bounded highest spin larger than two in general gives rise to $q_0 > 2$, which leads to very fast scrambling that violates the chaos bound. It is therefore reasonable to disallow a finite tower of higher spin fields, in addition to causality reasons [37]. This brings down q_0 to two, and, with this restriction, the metric is the field that can have the highest weight. This is the main reason behind the universality of the special pole-skipping point at $\omega = i\lambda_L$ and $k = i\lambda_L/v_B$, where $\lambda_L = 2\pi T$, and v_B is defined via a OTOC calculation.

In other words, for maximally chaotic holographic theories, instead of needing to compute a four-point function, the retarded Green's function already knows about

the butterfly velocity, and its dependence on the bulk theory is exactly the same as an OTOC would predict. It would be interesting to test this statement for non-holographic maximally chaotic theories [39]. Furthermore, there are now three ways of computing the butterfly velocity: (i) using entanglement wedge, (ii) using shockwave and (iii) using pole-skipping. Evidence for the equivalence between the first two was presented in [7, 11], and here we proved the equivalence between the last two.

The restriction of the discussion to bosons is for simplicity, and the generalization to include fermions should be completely analogous. For minimally coupled spinors on a fixed background, pole-skipping has been shown to happen at $\omega = (q_0 - s)\omega_0$ for a half integer $q_0 = 1/2$ and positive integers s [26]. If one allows both bosonic and fermionic fields with arbitrary couplings between them, one might expect that both q_0 and s can be half integers. It might be of use to analyze this with the weight argument, perhaps beginning by rephrasing the current discussion in a spin connection language.

We should summarize three assumptions that were used: (i) the existence of a finite q_0 ; (ii) the non-triviality of equation (15), i.e., the entries set to zero by this equations are not already all zero; and (iii) equation (17) has solutions. We expect that assumption i can be lifted with more careful analysis, but assumptions ii and iii are essential. Given any theory, one needs to check whether these are satisfied. For example, Vasiliev gravity which is dual to a sector of free $O(N)$ theory on the boundary [40] contains an infinite tower of higher spin fields but exhibits no chaos [36], so it should be illuminating to understand which assumption's failure is responsible.

Another condition implicit in our discussion is the restriction to finite temperatures. Extremal black holes do not have a bifurcate surface, so the property derived from regularity at the bifurcate surface no longer applies. Furthermore, poles in the Green's function get replaced by branch cuts [32, 41]. Accordingly, a generalization of our argument to zero temperature will be non-trivial.

We also showed that the shockwave metric could be obtained from a regularized mode of metric perturbation. This serves as an explanation for the similarities between the two calculations and the equivalence regardless of the theory. One might try different regulators or use different subtraction schemes to find a more regulator-independent relation.

ACKNOWLEDGMENTS

We thank Xi Dong, Gary Horowitz, Mark Mezei and Aron Wall for valuable feedback. DW is supported by NSF grant PHY2107939.

-
- [1] S. H. Shenker and D. Stanford, Black holes and the butterfly effect, *JHEP* **03**, 067, [arXiv:1306.0622 \[hep-th\]](#).
- [2] D. A. Roberts, D. Stanford, and L. Susskind, Localized shocks, *JHEP* **03**, 051, [arXiv:1409.8180 \[hep-th\]](#).
- [3] D. A. Roberts and B. Swingle, Lieb-Robinson Bound and the Butterfly Effect in Quantum Field Theories, *Phys. Rev. Lett.* **117**, 091602 (2016), [arXiv:1603.09298 \[hep-th\]](#).
- [4] S. H. Shenker and D. Stanford, Multiple Shocks, *JHEP* **12**, 046, [arXiv:1312.3296 \[hep-th\]](#).
- [5] S. H. Shenker and D. Stanford, Stringy effects in scrambling, *JHEP* **05**, 132, [arXiv:1412.6087 \[hep-th\]](#).
- [6] J. Maldacena, S. H. Shenker, and D. Stanford, A bound on chaos, *JHEP* **08**, 106, [arXiv:1503.01409 \[hep-th\]](#).
- [7] M. Mezei and D. Stanford, On entanglement spreading in chaotic systems, *JHEP* **05**, 065, [arXiv:1608.05101 \[hep-th\]](#).
- [8] M. Alishahiha, A. Davody, A. Naseh, and S. F. Taghavi, On Butterfly effect in Higher Derivative Gravities, *JHEP* **11**, 032, [arXiv:1610.02890 \[hep-th\]](#).
- [9] M. M. Qaemmaqami, Criticality in third order Lovelock gravity and butterfly effect, *Eur. Phys. J. C* **78**, 47 (2018), [arXiv:1705.05235 \[hep-th\]](#).
- [10] W.-J. Li, P. Liu, and J.-P. Wu, Weyl corrections to diffusion and chaos in holography, *JHEP* **04**, 115, [arXiv:1710.07896 \[hep-th\]](#).
- [11] X. Dong, D. Wang, W. W. Weng, and C.-H. Wu, A Tale of Two Butterflies: An Exact Equivalence in Higher-Derivative Gravity, preprint (2022), [arXiv:2203.06189 \[hep-th\]](#).
- [12] S. Grozdanov, K. Schalm, and V. Scopelitti, Black hole scrambling from hydrodynamics, *Phys. Rev. Lett.* **120**, 231601 (2018), [arXiv:1710.00921 \[hep-th\]](#).
- [13] M. Blake, H. Lee, and H. Liu, A quantum hydrodynamical description for scrambling and many-body chaos, *JHEP* **10**, 127, [arXiv:1801.00010 \[hep-th\]](#).
- [14] M. Blake, R. A. Davison, S. Grozdanov, and H. Liu, Many-body chaos and energy dynamics in holography, *JHEP* **10**, 035, [arXiv:1809.01169 \[hep-th\]](#).
- [15] M. Natsuume and T. Okamura, Holographic chaos, pole-skipping, and regularity, *PTEP* **2020**, 013B07 (2020), [arXiv:1905.12014 \[hep-th\]](#).
- [16] S. Grozdanov, P. K. Kovtun, A. O. Starinets, and P. Tadić, The complex life of hydrodynamic modes, *JHEP* **11**, 097, [arXiv:1904.12862 \[hep-th\]](#).
- [17] M. Blake, R. A. Davison, and D. Vegh, Horizon constraints on holographic Green's functions, *JHEP* **01**, 077, [arXiv:1904.12883 \[hep-th\]](#).
- [18] M. Blake and R. A. Davison, Chaos and pole-skipping in rotating black holes, *JHEP* **01**, 013, [arXiv:2111.11093 \[hep-th\]](#).
- [19] F. M. Haehl and M. Rozali, Effective Field Theory for Chaotic CFTs, *JHEP* **10**, 118, [arXiv:1808.02898 \[hep-th\]](#).
- [20] F. M. Haehl, W. Reeves, and M. Rozali, Reparametrization modes, shadow operators, and quantum chaos in higher-dimensional CFTs, *JHEP* **11**, 102, [arXiv:1909.05847 \[hep-th\]](#).
- [21] D. M. Ramirez, Chaos and pole skipping in CFT_2 , *JHEP* **12**, 006, [arXiv:2009.00500 \[hep-th\]](#).
- [22] C. Choi, M. Mezei, and G. Sárosi, Pole skipping away from maximal chaos, *JHEP* **02**, 207, [arXiv:2010.08558 \[hep-th\]](#).
- [23] D. T. Son and A. O. Starinets, Minkowski space correlators in AdS / CFT correspondence: Recipe and applications, *JHEP* **09**, 042, [arXiv:hep-th/0205051](#).
- [24] N. Iqbal and H. Liu, Real-time response in AdS/CFT with application to spinors, *Fortsch. Phys.* **57**, 367 (2009), [arXiv:0903.2596 \[hep-th\]](#).
- [25] M. Natsuume and T. Okamura, Nonuniqueness of Green's functions at special points, *JHEP* **12**, 139, [arXiv:1905.12015 \[hep-th\]](#).
- [26] N. Ceplak, K. Ramdial, and D. Vegh, Fermionic pole-skipping in holography, *JHEP* **07**, 203, [arXiv:1910.02975 \[hep-th\]](#).
- [27] Y. Ahn, V. Jahnke, H.-S. Jeong, K.-Y. Kim, K.-S. Lee, and M. Nishida, Pole-skipping of scalar and vector fields in hyperbolic space: conformal blocks and holography, *JHEP* **09**, 111, [arXiv:2006.00974 \[hep-th\]](#).
- [28] Y. Ahn, V. Jahnke, H.-S. Jeong, K.-S. Lee, M. Nishida, and K.-Y. Kim, Classifying pole-skipping points, *JHEP* **03**, 175, [arXiv:2010.16166 \[hep-th\]](#).
- [29] K.-Y. Kim, K.-S. Lee, and M. Nishida, Construction of bulk solutions for towers of pole-skipping points, *Phys. Rev. D* **105**, 126011 (2022), [arXiv:2112.11662 \[hep-th\]](#).
- [30] X. Wu, Higher curvature corrections to pole-skipping, *JHEP* **12**, 140, [arXiv:1909.10223 \[hep-th\]](#).
- [31] M. Natsuume and T. Okamura, Pole-skipping with finite-coupling corrections, *Phys. Rev. D* **100**, 126012 (2019), [arXiv:1909.09168 \[hep-th\]](#).
- [32] M. Natsuume and T. Okamura, Pole-skipping and zero temperature, *Phys. Rev. D* **103**, 066017 (2021), [arXiv:2011.10093 \[hep-th\]](#).
- [33] S. Grozdanov, On the connection between hydrodynamics and quantum chaos in holographic theories with stringy corrections, *JHEP* **01**, 048, [arXiv:1811.09641 \[hep-th\]](#).
- [34] A. C. Wall, A Second Law for Higher Curvature Gravity, *Int. J. Mod. Phys. D* **24**, 1544014 (2015), [arXiv:1504.08040 \[gr-qc\]](#).
- [35] There are points at which triviality of equations of motion does not imply dependence on $\delta\omega/\delta k$ [17], but the limit does depend on higher order quantities such as $(\delta k)^2$ [28]. This justifies our assumption here.
- [36] E. Perlmutter, Bounding the Space of Holographic CFTs with Chaos, *JHEP* **10**, 069, [arXiv:1602.08272 \[hep-th\]](#).
- [37] X. O. Camanho, J. D. Edelstein, J. Maldacena, and A. Zhiboedov, Causality Constraints on Corrections to the Graviton Three-Point Coupling, *JHEP* **02**, 020, [arXiv:1407.5597 \[hep-th\]](#).
- [38] We checked the computation explicitly for $f(\text{Riemann})$ gravity using the method of [11] and the results comply.
- [39] For non-maximally chaotic theories, the predictions from pole-skipping could differ from OTOC results [22, 42].
- [40] I. R. Klebanov and A. M. Polyakov, AdS dual of the critical $O(N)$ vector model, *Phys. Lett. B* **550**, 213 (2002), [arXiv:hep-th/0210114](#).
- [41] T. Faulkner, H. Liu, J. McGreevy, and D. Vegh, Emergent quantum criticality, Fermi surfaces, and AdS(2), *Phys. Rev. D* **83**, 125002 (2011), [arXiv:0907.2694 \[hep-th\]](#).
- [42] M. Mezei and G. Sárosi, Chaos in the butterfly cone, *JHEP* **01**, 186, [arXiv:1908.03574 \[hep-th\]](#).