ON CYLINDRICAL SYMMETRIC PROJECTIVELY FLAT FINSLER METRICS

NEWTON SOLÓRZANO AND VÍCTOR LEÓN

ABSTRACT. We study the cylindrical symmetric Finsler metrics. We obtain the system of differential equations of such metrics which are projectively flat. We give a family of solutions of this system. Examples are included.

1. INTRODUCTION AND MAIN RESULTS

A Finsler metric F(x, y) on an open subset $U \subset \mathbb{R}^n$ is called *projectively flat* if all geodesics are straight lines in U. A Finsler metric F on a manifold M is called *locally projectively flat* if at any point, there is a locally coordinate system (x^i) where F is projectively flat. Hilbert's Fourth Problem, which studies locally projectively flat metrics, is an important problem in Finsler geometry and projectively flat Finsler metrics on a convex domain in \mathbb{R}^n are regular solution to Hilbert's Fourth Problem.

G. Hamel [4] obtained conditions which characterize projectively flat Finsler metric F = F(x, y) on an open subset $U \subset \mathbb{R}^n$. That is

$$F_{x^k y^l} y^k - F_{x^l} = 0.$$

A Randers metric $F = \alpha + \beta$ is locally projectively flat if, and only if, α is locally projectively flat and β is closed [1]. In [6], the authors founded the PDE which characterizes spherically symmetric Finsler metric and projectively flat and given an example using the Gauss error function. In [16], the author studied (α, β) -metrics, and found a necessary and sufficient condition for such metrics to be projectively flat in dimension $n \geq 3$. In [9], the authors studied locally projectively flat warped Finsler metric of constant flag curvature and obtained new examples. In [13], the authors studied the general (α, β) -metrics, and they classify those locally projectively flat when α is projectively flat. New examples were founded.

The following known Finsler metrics:

(1) the known Shen's fish tank metric on $M = I \times \mathbb{B}^2 \subset \mathbb{R}^3$ are of the form:

$$F = \frac{\sqrt{(-x^2y^1 + x^1y^2)^2 + ((y^0)^2 + (y^1)^2 + (y^2)^2)(1 - (x^1)^2 - (x^2)^2)}}{1 - (x^1)^2 - (x^2)^2} - \frac{x^2y^1 - x^1y^2}{1 - (x^1)^2 - (x^2)^2},$$

where $x = (x^0, x^1, x^2) \in \mathbb{R} \times \mathbb{B}^2$ and $y = (y^0, y^1, y^2) \in T_x M$, (2) the particular Randers metric constructed by Z. Shen in [15]:

$$F = \frac{\sqrt{(1 - |x|^4)|y|^2 + (y^0|x|^2 - 2x^0\langle x, y\rangle)^2}}{1 - |x|^4} + \frac{y^0|x|^2 - 2x^0\langle x, y\rangle}{1 - |x|^4},$$

where $x = (x^0, x^1, ..., x^n) \in M = I \times \mathbb{R}^n, y = (y^0, y^1, ..., y^n) \in T_x M$,

Date: August 3, 2022.

²⁰²⁰ Mathematics Subject Classification. 53B40, 53C60.

Key words and phrases. Finsler metric, Cylindrically symmetric, Projectively flat, Warped metric.

(3) the spherically symmetric Finsler metric [5, 14]:

$$F = |y|\phi\left(|x|, \frac{\langle x, y\rangle}{|y|}\right),$$

where $x \in M = \mathbb{R}^{n+1}, y \in T_x M$,

(4) the warped metrics [2, 7, 8, 10] defined on $I \times \mathbb{R}^n$ of the form

$$F = |\overline{y}|\phi\left(x^{0}, \frac{y^{0}}{|\overline{y}|}\right),$$
$$F = |\overline{y}|\phi\left(\frac{y^{0}}{|\overline{y}|}, |\overline{x}|\right),$$

where $x = (x^0, \overline{x}) = (x^0, x^1, ..., x^n) \in M = I \times \mathbb{R}^n, y = (y^0, \overline{y}) = (y^0, y^1, ..., y^n) \in T_x M$,

are important and very studied Finsler metrics. We observed that those above metrics satisfy

(1.1)
$$F((x^0, O\overline{x}), (y^0, O\overline{y})) = F((x^0, \overline{x}), (y^0, \overline{y})),$$

for every $O \in O(n)$. A Finsler metric F is called *cylindrically symmetric* if F satisfies (1.1) for all $O \in O(n)$.

In [17] (see Theorem 4 in section 2) the author showed that every cylindrically symmetric Finsler metric can be written as

$$F(x,y) = |\overline{y}|\phi\left(x^{0}, \frac{y^{0}}{|\overline{y}|}, |\overline{x}|, \frac{\langle \overline{x}, \overline{y} \rangle}{|\overline{y}|}\right),$$

where $|\cdot|$ and $\langle \cdot, \cdot \rangle$ are, respectively, the standard Euclidean norm and inner product on \mathbb{R}^n .

In this paper, we give necessary and sufficient conditions for $F = |\overline{y}|\phi$ to be a Finsler metric. The proof is inspired by the proof given in [12] for (α, β) -metrics.

Theorem 1. Let $F = |\overline{y}| \phi(x^0, z, r, s)$ be a Finsler metric defined on $M = I \times \mathbb{B}^n(\rho)$, where $z = \frac{y^0}{|\overline{y}|}$, $r = |\overline{x}|$, $s = \frac{\langle \overline{x}, \overline{y} \rangle}{|\overline{y}|}$ and TM with coordinates

(1.2)
$$x = (x^0, \overline{x}), \ \overline{x} = (x^1, \dots, x^n), \ y = (y^0, \overline{y}) \ and \ \overline{y} = (y^1, \dots, y^n).$$

Then, F is a Finsler metric if, and only if, the positive function ϕ satisfies $\Lambda > 0$ for n = 2 with additional inequality, $\Omega > 0$ for $n \ge 3$.

Next, we obtain a system of differential equations which characterize cylindrical symmetric projectively flat Finsler metrics:

Theorem 2. Let $F = |\overline{y}| \phi(x^0, z, r, s)$ be a Finsler metric defined on $M = I \times \mathbb{B}^n(\rho)$, where $z = \frac{y^0}{|\overline{y}|}$, $r = |\overline{x}|$, $s = \frac{\langle \overline{x}, \overline{y} \rangle}{|\overline{y}|}$ and TM with coordinates (1.2). Then F is projectively flat if and only if ϕ satisfy

(1.3)
$$\Omega_{x^0} - \phi_{sz} = 0,$$

(1.4)
$$\Omega_r - r\phi_{ss} = 0,$$

where

$$\Omega = \phi - s\phi_s - z\phi_z$$

Studying this PDE, we obtain a family of solutions of this system:

Theorem 3. The family of functions given by

(1.5)
$$\phi = g_1(z) + x^0 g_2(z) + sg_3(z) + zg_4(x^0) + sg_5(r) + \int_0^s \left(\int_0^\eta g_6(r^2 - \xi^2) d\xi \right) d\eta + \int_0^r \xi g_6(\xi^2) d\xi,$$

 $\mathbf{2}$

where $g_i, i = 1, ..., 6$, are arbitrary real differentiable functions with $g_2(z) - zg'_2(z) - g'_3(z) = 0$, are solutions of the system (1.3)-(1.4). Moreover, any Finsler metric on $M = I \times \mathbb{B}^n(\rho)$ defined by

$$F(x,y) = |\overline{y}|\phi\left(x^{0}, \frac{y^{0}}{|\overline{y}|}, |\overline{x}|, \frac{\langle \overline{x}, \overline{y} \rangle}{|\overline{y}|}\right),$$

where the positive function ϕ is given by (1.5) with $g_i, i = 1, ..., 6$ satisfy

$$\Lambda = \left(g_1(z) - zg_1'(z) + (x^0 - sz)g_3'(z) + \frac{1}{2}\int_0^{r^2 - s^2} g_6(\xi) \,d\xi + (r^2 - s^2)g_6(r^2 - s^2)\right) \times (1.6)$$

$$(g_1''(z) + (x^0 - sz)g_2''(z)) - (r^2 - s^2)[g_3'(z)]^2 > 0, \text{ when } n \ge 2,$$

with the additional inequality

(1.7)
$$\Omega = g_1(z) - zg_1'(z) + (x^0 - sz)g_3'(z) + \frac{1}{2}\int_0^{r^2 - s^2} g_6(\xi) \,d\xi > 0, \qquad \text{when } n \ge 3,$$

are locally projectively flat Finsler Metric.

Finally, using results in other works [11, 5, 17] we obtain new examples (see Section 4), which are a generalization of the warped metrics mentioned above.

2. Preliminaries

In this section, we give some notations, definitions and lemmas that will be used in the proof of our main results. Let M be a manifold and let $TM = \bigcup_{x \in M} T_x M$ be the tangent bundle of M, where $T_x M$ is the tangent space at $x \in M$. We set $TM_o := TM \setminus \{0\}$ where $\{0\}$ stands for $\{(x, 0) | x \in M, 0 \in T_x M\}$. A *Finsler metric* on M is a function $F : TM \to [0, \infty)$ with the following properties:

(a) F is C^{∞} on TM_o ;

(b) At each point $x \in M$, the restriction $F_x := F|_{T_xM}$ is a Minkowski norm on T_xM .

Let $\mathbb{B}^n(\rho) \subset \mathbb{R}^n$ the *n* dimensional open ball of radius ρ and centered at the origin $(n \ge 2)$. Set $M = I \times \mathbb{B}^n(\rho)$, with coordinates on TM

$$x = (x^0, \overline{x}), \ \overline{x} = (x^1, \dots, x^n), \ y = (y^0, \overline{y}) \ \text{and} \ \overline{y} = (y^1, \dots, y^n).$$

Consider the Finsler metric F defined on M such that

(2.1)
$$F((x^0, O\overline{x}), (y^0, O\overline{y})) = F((x^0, \overline{x}), (y^0, \overline{y}))$$

for every orthogonal $n \times n$ matrix O.

Inspired by [14], (see also [6]) in [17] was proved the following:

Theorem 4. [17] A Finsler metric F, defined on $M = \mathbb{R} \times \mathbb{B}^n(\rho)$, satisfies (2.1) if, and only if, there exists a positive differentiable function $\phi : \mathbb{R}^4 \to \mathbb{R}$ such that

$$F(x,y) = |\overline{y}|\phi\left(x^0, \frac{y^0}{|\overline{y}|}, |\overline{x}|, \frac{\langle \overline{x}, \overline{y} \rangle}{|\overline{y}|}\right),$$

where $|\cdot|$ and $\langle \cdot, \cdot \rangle$ are, respectively, the standard Euclidean norm and inner product on \mathbb{R}^n .

For $|\overline{y}| \neq 0$, we introduce the notation

$$z := \frac{y^0}{|\overline{y}|}, \ r := |\overline{x}| \ \text{and} \ s := \frac{\langle \overline{x}, \overline{y} \rangle}{|\overline{y}|},$$

where |.| and \langle , \rangle are, respectively, the standard Euclidean norm and inner product on \mathbb{R}^n . Throughout our work, the following convention for indices is adopted:

$$0 \leq A, B, \ldots \leq n$$
 and $1 \leq i, j, \ldots \leq n$.

Defining Ω and Λ by

(2.2)
$$\Omega := \phi - s\phi_s - z\phi_z$$

(2.3) $\Lambda := \Omega \phi_{zz} + (r^2 - s^2)(\phi_{ss}\phi_{zz} - \phi_{sz}^2),$

the matrix
$$(g_{AB}) = \frac{1}{2} [F^2]_{y^A y^B} = \left(\begin{array}{c|c} g_{00} & g_{0j} \\ \hline g_{i0} & g_{ij} \end{array} \right)$$
, is given by
 $g_{00} = \phi_z^2 + \phi \phi_{zz}, \ g_{i0} = g_{0i} = (\phi \Omega)_z u^i + (\phi_s \phi_z + \phi \phi_{sz}) x^i,$

 $\begin{array}{c} \text{and } g_{ij} = \phi \Omega \delta_{ij} + X_{ij}, \\ \text{where } X_{ij} = (u^i, x^i) \left(\begin{array}{c} -[s(\phi \Omega)_s + z(\phi \Omega)_z] & (\phi \Omega)_s \\ (\phi \Omega)_s & \phi_s^2 + \phi \phi_{ss} \end{array} \right) \left(\begin{array}{c} u^j \\ x^j \end{array} \right), \text{ with } u^i = \frac{y^i}{|\overline{y}|}. \\ \text{Note that,} \end{array}$

 $\det(g_{AB}) = (\phi_z^2 + \phi \phi_{zz})^{1-n} \phi^n \det\left((\phi_z^2 + \phi \phi_{zz})\Omega \delta_{ij} + \delta_0 u^i u^j + \delta_1 (x^i + u^i)(x^j + u^j) + \delta_2 x^i x^j\right),$ where,

$$\begin{split} \delta_0 &= -\left[(\phi\Omega)_z \left((\phi\Omega)_z - \phi_s \phi_z - \phi \phi_{sz} \right) + (\phi_z^2 + \phi \phi_{zz}) \left((s+1)(\phi\Omega)_s + z(\phi\Omega)_z \right) \right] \\ \delta_1 &= (\phi\Omega)_s (\phi_z^2 + \phi \phi_{zz}) - (\phi\Omega)_z (\phi_s \phi_z + \phi \phi_{sz}), \\ \delta_2 &= (\phi \phi_{zz} + \phi_z^2) \left(\phi \phi_s + \phi_s^2 - (\phi\Omega)_s \right) - (\phi_s \phi_z + \phi \phi_{sz}) \left(\phi_s \phi_z + \phi_{sz}^2 - (\phi\Omega)_z \right). \end{split}$$

Then, the determinant of g_{AB} is given by

(2.4)
$$\det(g_{AB}) = \phi^{n+2} \Omega^{n-2} \Lambda.$$

As mentioned in [7], the called warped metrics $F = \sqrt{F_1^2 + f^2 F_2^2}$, where $(M, F_1), (N, F_2)$ are Finsler manifolds and f is a smooth function on M, are not necessarily smooth on the vectors of the form $(v_1, 0)$ and $(0, v_2) \in TM \times TN$. However, the first two Shen's examples in the introduction part motivate us to formalize a Finsler metric which satisfies (2.1) although warped type metrics can be obtained by similar technician.

3. Proof of Theorems 1, 2 and 3

Proof of Theorem 1. Suppose that $\Omega, \Lambda > 0$. From $\Lambda > 0$ we conclude that $\phi_{zz} > 0$. In fact, suppose that there exist x_0^0, z_0, r_0, s_0 such that $\phi_{zz}(x_0^0, z_0, r_0, s_0) = 0$, then $\Lambda|_{x_0^0, z_0, r_0, s_0} = -(r^2 - s^2)\phi_{sz}^2 \leq 0$ and, if $\phi_{zz} < 0$, hence $\Lambda|_{r=s=0} = \Omega\phi_{zz}|_{r=s=0} < 0$. Additionally, from $\Lambda > 0$ we obtain $\Omega + (r^2 - s^2)\phi_{ss} > 0$.

Now, we consider the family of functions ϕ_t defined by,

$$\phi_t := (1-t)\sqrt{1+z^2} + t\phi.$$

Let $F_t := |\overline{y}| \phi_t$ and $g_{AB}^t := \frac{1}{2} \left[F_t^2 \right]_{y^A y^B}$. Note that, for any $0 \le t \le 1$, we have

$$\begin{aligned} \Omega_t &:= \phi_t - s(\phi_t)_s - z(\phi_t)_z = \frac{1-t}{\sqrt{1+z^2}} + t\Omega > 0, \\ \Lambda_t &:= \Omega_t(\phi_t)_{zz} + (r^2 - s^2) \left((\phi_t)_{ss}(\phi_t)_{zz} - (\phi_t)_{sz}^2 \right) \\ &= \frac{(1-t)^2}{(1+z^2)^2} + \frac{t(1-t)}{\sqrt{1+z^2}} \phi_{zz} + \frac{t(1-t)}{(1+z^2)^{3/2}} \left(\Omega + (r^2 - s^2) \phi_{ss} \right) + t^2 \Lambda > 0 \end{aligned}$$

Thus $det(g_{AB}^t) > 0$ for all $0 \le t \le 1$. Since (g_{AB}^0) is positive definite, we conclude that (g_{AB}^t) is positive definite for any $t \in [0, 1]$. Therefore, F_t is a Finsler metric for any $t \in [0, 1]$.

Conversely, assume that $F = |\overline{y}|\phi(x^0, z, r, s)$ is a Finsler metric on M. Then $\phi > 0$. By (2.4), $\det(g_{AB}) > 0$ is equivalent to

(3.1)
$$\Omega^{n-2}\Lambda > 0,$$

which implies $\Omega \neq 0$ when $n \geq 3$. Since $\phi|_{s=z=0} > 0$, we have $\Omega|_{s=z=0} > 0$, then $\Omega > 0$. And by (3.1) we have $\Lambda > 0$.

The inverse of (g_{AB}) is given by,

$$\left(g^{AB}\right) = \frac{1}{\phi^4 \Lambda} \left(\begin{array}{c|c} y^{00} & y^{0j} \\ \hline y^{i0} & \frac{\phi^3 \Lambda}{\Omega} \delta_{ij} + Y_{ij} \end{array} \right)$$

where

$$y^{00} = \phi \Omega((\phi - z\phi_z)^2 + z^2\phi\phi_{zz}) + (r^2 - s^2)\phi \left(\phi^2\phi_{ss} + 2z\phi(\phi_s\phi_{sz} - \phi_z\phi_{ss}) + z^2\delta_3\right)$$

$$y^{0i} = \phi \left[-(\Omega + s\phi_s)(\phi\Omega)_z + (r^2 - s^2)(\phi(\phi_s\phi_{sz} - \phi_z\phi_{ss}) + z\delta_3)\right]u^i + \phi^2[\phi_s\Omega_z - \phi_{sz}\Omega]x^i$$

ith

with

$$\delta_{3} = \phi(\phi_{ss}\phi_{zz} - \phi_{sz}^{2}) + \phi_{s}(\phi_{s}\phi_{zz} - \phi_{z}\phi_{sz}) + \phi_{z}(\phi_{ss}\phi_{z} - \phi_{s}\phi_{sz})$$
$$= -\det \begin{pmatrix} -\phi & \phi_{s} & \phi_{z} \\ \phi_{s} & \phi_{ss} & \phi_{sz} \\ \phi_{z} & \phi_{sz} & \phi_{zz} \end{pmatrix},$$

and

$$Y_{ij} = \frac{1}{\phi\Omega} (u^i, x^i) \begin{pmatrix} y_{11} & y_{12} \\ y_{12} & y_{22} \end{pmatrix} \begin{pmatrix} u^j \\ x^j \end{pmatrix}$$

where

$$y_{11} = \phi^2 \left[(\phi \Omega)_z^2 + \phi \phi_{zz} (z(\phi \Omega)_z + s(\phi \Omega)_s) - (r^2 - s^2) (\phi^2 (\phi_{ss} \phi_{zz} - \phi_{sz}^2) - \Omega \delta_2) \right]$$

$$y_{12} = y_{21} = \phi^3 \left[\phi_{sz} (\phi \Omega)_z - \phi_{zz} (\phi \Omega)_s \right]$$

$$y_{22} = -\phi^4 (\phi_{ss} \phi_{zz} - \phi_{sz}^2).$$

The next proposition gives us one the most important quantities in Finsler Geometry: The geodesic coefficients

$$(3.2) G^A = Py^A + Q^A,$$

where

(3.3)
$$P := \frac{F_{x^C} y^C}{2F} \text{ and } Q^A := \frac{F}{2} g^{AB} \left\{ F_{x^C y^B} y^C - F_{x^B} \right\},$$

for a cylindrically simmetric Finsler metric F.

Proposition 1. $F = |\overline{y}| \phi(x^0, z, r, s)$ be a Finsler metric defined on $M = I \times \mathbb{B}^n(\rho)$, where $z = \frac{y^0}{|\overline{y}|}$, $r = |\overline{x}|$, $s = \frac{\langle \overline{x}, \overline{y} \rangle}{|\overline{y}|}$ and TM with coordinates (1.2). Then the geodesic spray coefficients G^A are given by

$$G^{0} = u^{2} \left\{ z(W + sU) + \frac{\Omega}{2\Lambda} (\varphi_{z} - 2\phi_{x^{0}}) - (r^{2} - s^{2})V \right\},\$$

$$G^{i} = uWy^{i} + u^{2}Ux^{i},$$

where Ω, Λ are in (2.2), (2.3) respectively, $u = |\overline{y}|$, and

$$\begin{split} \varphi &:= z\phi_{x^0} + \frac{s}{r}\phi_r + \phi_s, \\ W &:= \frac{1}{\phi} \left\{ \frac{\varphi}{2} - s\phi U - \frac{\phi_z \Omega}{2\Lambda} (\varphi_z - 2\phi_{x^0}) - (r^2 - s^2) \left[\phi_s U - \phi_z V\right] \right\}, \\ U &:= \frac{1}{2\Lambda} \left\{ \left(\varphi_s - \frac{2}{r}\phi_r\right) \phi_{zz} - (\varphi_z - 2\phi_{x^0}) \phi_{sz} \right\}, \end{split}$$

(3.4)
$$V := \frac{1}{2\Lambda} \left\{ \left(\varphi_s - \frac{2}{r} \phi_r \right) \phi_{sz} - \left(\varphi_z - 2\phi_{x^0} \right) \phi_{ss} \right\}.$$

Proof. The derivatives of $F = |\overline{y}|\phi$, with respect to x^0, x^i, y^0 and y^i are given by

$$\begin{split} F_{x^{0}} &= u\phi_{x^{0}}, \\ F_{x^{i}} &= u\left(\phi_{s}u^{i} + \phi_{r}\frac{x^{i}}{r}\right), \\ F_{y^{0}} &= \phi_{z}, \\ F_{x^{0}y^{0}} &= \phi_{x^{0}z}, \\ F_{x^{i}y^{0}} &= \phi_{sz}u^{i} + \phi_{rz}\frac{x^{i}}{r}, \\ F_{y^{i}} &= \Omega u^{i} + \phi_{s}x^{i}, \\ F_{x^{0}y^{i}} &= \Omega_{x^{0}}u^{i} + \phi_{x^{0}s}x^{i}, \\ F_{x^{i}y^{j}} &= \phi_{s}\delta_{ij} + \Omega_{s}u^{i}u^{j} + \phi_{ss}\frac{y^{i}}{u}x^{j} + \Omega_{r}\frac{x^{i}}{r}u^{j} + \frac{1}{r}\phi_{rs}x^{i}x^{j}. \end{split}$$

Then,

$$F_{x^C y^0} y^C - F_{x^0} = u \left(\varphi_z - 2\phi_{x^0}\right),$$

$$F_{x^C y^j} y^C - F_{x^j} = u \left(\left[-s(\varphi_s - \frac{2}{r}\phi_r) - z(\varphi_z - 2\phi_{x^0}) \right] u^j + \left[\varphi_s - \frac{2}{r}\phi_r \right] x^j \right).$$

Thus, from (3.3), we have

$$(3.5) \qquad P = \frac{u^2}{2F}\varphi,$$

$$(3.6) \qquad Q^0 = u^2 \left\{ \frac{1}{2\phi\Lambda} (\phi - z\phi_z)(\varphi_z - 2\phi_{x^0})\Omega - (r^2 - s^2) \left(z\frac{\phi_s}{\phi}U + \frac{\phi - z\phi_z}{\phi}V \right) \right\},$$

(3.7)
$$Q^{i} = -u^{2} \left\{ sU + (r^{2} - s^{2})(\frac{\phi_{s}}{\phi}U - \frac{\phi_{z}}{\phi}V) + \frac{\phi_{z}\Omega}{2\phi\Lambda}(\varphi_{z} - 2\phi_{x^{0}}) \right\} \frac{y^{i}}{u} + u^{2}Ux^{i}.$$

Replacing (3.5), (3.6) and (3.7) into (3.2), we have the result.

A Finsler metric F is projectively flat if, and only if, F satisfies

$$F_{x^k y^l} y^k - F_{x^l} = 0.$$

From (3.6) and (3.7) we obtain a characterization of $F = |\overline{y}|\phi$ which are projectively flat.

Proof of Theorem 2. From (3.6) and (3.7), the Finsler metric $F = |\overline{y}|\phi(x^0, z, r, s)$ is projectively flat if, and only if,

(3.8)
$$\begin{cases} U = 0, \\ (r^2 - s^2)\frac{\phi_z}{\phi}V = \frac{\phi_z}{2\phi\Lambda}\Omega(\varphi_z - 2\phi_{x^0}), \\ (r^2 - s^2)\left(\frac{\phi - z\phi_z}{\phi}\right)V = \frac{\Omega}{2\phi\Lambda}(\phi - z\phi_z)(\varphi_z - 2\phi_{x^0}). \end{cases}$$

Using the fact $\phi - z\phi_z \neq 0$, the system (3.8) is equivalent to

(3.9)
$$\begin{cases} (\varphi_s - \frac{2}{r}\phi_r)\phi_{zz} &= (\varphi_z - 2\phi_{x^0})\phi_{sz}, \\ (\varphi_z - 2\phi_{x^0})\Omega &= 2(r^2 - s^2)\Lambda V. \end{cases}$$

From the first equation of (3.9) and (3.4) we have

$$V = \frac{1}{2\phi_{zz}\Lambda}(\varphi_z - 2\phi_{x^0})(\phi_{sz}^2 - \phi_{ss}\phi_{zz}).$$

Substituting this in the second equation of (3.9), and using the fact $\Lambda \neq 0, \forall y$ we have that the system (3.9) implies

$$\varphi_z - 2\phi_{x^0} = 0,$$

substituting this into the system (3.9), we obtain that the system (3.9) is equivalent to

(3.10)
$$r(\phi_{x^0} - z\phi_{x^0z} - \phi_{sz}) - s\phi_{rz} = 0,$$

(3.11)
$$s\phi_{rs} + r(\phi_{ss} + z\phi_{x^0s}) - \phi_r = 0.$$

Derivating (3.10) and (3.11) in relation to s and z respectively, and subtracting them, we have

(3.12)
$$\phi_{rz} - r\phi_{x^0s} = 0,$$

using (3.12) in equations (3.10) and (3.11) we have the result. The reciprocal is immediately.

Proof of Theorem 3. To construct this family, note that ϕ given by

(3.13)
$$\phi(x^0, z, r, s) = \varepsilon_1(x^0, r) + \varepsilon_2(x^0, z) + \varepsilon_3(r, s) + \varepsilon_4(s, z),$$

where $\varepsilon_1, \varepsilon_2, \varepsilon_3$ and ε_4 are real differentiable functions, satisfy (3.12).

Observe that equations (1.3) and (1.4), in this case, become

(3.14)
$$\Omega_{x^0} - \phi_{sz} = [\varepsilon_1]_{x^0} + [\varepsilon_2]_{x^0} - z[\varepsilon_2]_{x^0z} - [\varepsilon_4]_{sz} = 0,$$

(3.15)
$$\Omega_r - r\phi_{ss} = [\varepsilon_1]_r + [\varepsilon_3]_r - s[\varepsilon_3]_{rs} - r[\varepsilon_3 + \varepsilon_4]_{ss} = 0$$

Taking the derivative of (3.14) with respect to r, we have

$$[\varepsilon_1]_{x^0r} = 0,$$

therefore exists real differential functions $f_1, f_2 : \mathbb{R} \to \mathbb{R}$ such that

(3.16)
$$\varepsilon_1 = f_1(x^0) + f_2(r).$$

Now, taking the derivative of (3.14) with respect to x^0 we get

$$[\varepsilon_2]_{x^0x^0} - z[\varepsilon_2]_{x^0x^0z} = -f_1''(x^0),$$

therefore, there exists real differentiable functions $f_3, f_4, f_5 : \mathbb{R} \to \mathbb{R}$ such that

(3.17)
$$\varepsilon_2 = -f_1(x^0) + zf_3(x^0) + x^0 f_4(z) + f_5(z).$$

Again, taking the derivative of (3.14) with respect to s, we have

$$[\varepsilon_4]_{zss} = 0,$$

therefore there exists real differential functions $f_6, f_7, f_8 : \mathbb{R} \to \mathbb{R}$ such that

(3.18)
$$\varepsilon_4 = f_6(s) + f_7(z) + sf_8(z).$$

Also, taking the derivative of (3.15) with respect to s, we have

$$s\Big[[\varepsilon_3]_{ss}\Big]_r + r\Big[[\varepsilon_3]_{ss}\Big]_s = -rf_6^{\prime\prime\prime}(s),$$

now using characteristic equations in this non-homogeneous quasi linear PDE, we obtain that there exists a real differentiable function $f_9 : \mathbb{R} \to \mathbb{R}$ such that

$$[\varepsilon_3]_{ss} = f_9(r^2 - s^2) - f_6''(s),$$

consequently, together (3.15), there exists real differential function $f_{10} : \mathbb{R} \to \mathbb{R}$ and constant $k \in \mathbb{R}$ such that

(3.19)
$$\varepsilon_3 = k - f_2(r) - f_6(s) + s f_{10}(r) + \int_0^s \left(\int_0^\eta f_9(r^2 - \xi^2) d\xi \right) d\eta + \int_0^r \rho f_9(\rho^2) d\rho.$$

Also substituting (3.16), (3.17), (3.18) and (3.19) in (3.13) and using (3.14), we have

 $f_4(z) - zf'_4(z) - f'_8(z) = 0.$

Finally, the identity

(3.20)
$$\int_0^s \int_0^\eta g_6(r^2 - \xi^2) d\xi d\eta + \int_0^r \rho g_6(\rho^2) d\rho = \frac{1}{2} \int_0^{r^2 - s^2} g_6(\xi) d\xi + s \int_0^s g_6(r^2 - \xi^2) d\xi,$$
and the Theorem 1 give us the conditions (1.6)-(1.7).

Remark 1. As seen in the proof above, we give another useful expression for the general solution of

$$s\phi_{bs} + b\phi_{ss} - \phi_b = 0$$

founded in Proposition 5.1 in [6]:

(3.21)
$$\phi(b,s) = sh(b) - s \int_{s_0}^s t^{-2}g(b^2 - t^2) dt$$

In fact, it is enough to take the derivative of the equation with respect to s, to getting

$$s \Big[\phi_{ss} \Big]_b + b \Big[\phi_{ss} \Big]_s = 0.$$

Using characteristic equations in the homogeneous quasi linear PDE, we obtain that there exists a real differentiable function $f : \mathbb{R} \to \mathbb{R}$ such that

$$\phi_{ss} = f(b^2 - s^2)$$

therefore, there exists real differentiable functions $g, h : \mathbb{R} \to \mathbb{R}$ and constant $k \in \mathbb{R}$ such that

(3.22)
$$\phi = k + sg(b) + \int_0^s \left(\int_0^\eta f(b^2 - \xi^2) d\xi \right) d\eta + \int_0^b \rho f(\rho^2) d\rho$$

by the identity (3.20) we have that conditions $\phi - s\phi_s > 0$ and $\phi - s\phi_s + (b^2 - s^2)\phi_{ss} > 0$ become

(3.23)
$$k + \frac{1}{2} \int_0^{b^2 - s^2} f(u) du > 0, \ \forall \ b \ge |s|,$$

and

(3.24)
$$k + \frac{1}{2} \int_0^{b^2 - s^2} f(\xi) d\xi + (b^2 - s^2) f(b^2 - s^2) > 0, \ \forall \ b \ge |s|,$$

respectively.

The relation between g and f from (3.21) and (3.22) is given by:

$$f = 2g'$$
.

Remark 2. Observe that, for every constant k > 0 and any function f such that $f(\xi) \ge 0, \forall \xi \ge 0$, the conditions (3.23)-(3.24) are satisfied.

Remark 3. The identity (3.20) give us two forms to write the function ϕ in (1.5).

We can use functions founded in [11], [5] and [17] for construct warped-type Finsler metrics, i.e.

Corollary 1. Let $\phi = \phi(x^0, z, r, s)$ be a positive function defined by

(3.25)
$$\phi = k + g_1(z) + zg_4(x^0) + sg_5(r) + \frac{1}{2} \int_0^{r^2 - s^2} g_6(\xi) \, d\xi + s \int_0^s g_6(r^2 - \xi^2) \, d\xi,$$

where $k \in \mathbb{R}$ is constant and $g_1, g_4, g_5, g_6 : \mathbb{R} \to \mathbb{R}$, are differentiable functions, such that (a) $g_1(z) + zg_4(x^0) > 0$, $g_1(z) - zg'_1(z) > 0$ and $g''_1(z) > 0$, $\forall z \in \mathbb{R}$;

(b)
$$k + \frac{1}{2} \int_{0}^{r^{2} - s^{2}} g_{6}(\xi) d\xi + (r^{2} - s^{2}) g_{6}(r^{2} - s^{2}) \ge 0, \ \forall \ r \ge |s|, \ when \ n \ge 2,$$

with additional inequality $k + \frac{1}{2} \int_{0}^{r^{2} - s^{2}} g_{6}(u) du \ge 0, \ \forall \ r \ge |s|, \ when \ n \ge 3,$

then, the warped type Finsler metric on $M = I \times \mathbb{B}^n$ given by

$$F = |\overline{y}| \phi\left(\frac{y^0}{|\overline{y}|}, |\overline{x}|, \frac{\langle \overline{x}, \overline{y} \rangle}{|\overline{y}|}\right),$$

is locally projectively flat.

Proof. In Theorem 3 we considered $g_2(z) = g_3(z) = 0$ (the result is similar if we consider $g_3(z) = 0$ uniquely). By (a), $\phi_{zz} = g''_1(z) > 0$, $\forall z \in \mathbb{R}$. Consequently, from (a), (b), (1.6) and (1.7),

$$\Omega = k + g_1(z) - zg'_1(z) + \frac{1}{2} \int_0^{r^2 - s^2} g_6(\xi) d\xi > 0, \ \forall z \in \mathbb{R}, \ r \ge |s|;$$

$$\Lambda = \Omega \phi_{zz} + (r^2 - s^2) g_6(r^2 - s^2) \phi_{zz} > 0, \ \forall z \in \mathbb{R}, \ r \ge |s|.$$

Observe that if k > 0 and $g_6(\xi) \ge 0, \forall \xi \ge 0$ then, the condition (b), in Corollary 1, is satisfied.

4. Examples

Example 1. In Corollary 1, by choosing $g_1(z) = \sqrt{z^2 + \varepsilon} + \gamma z$ ($\varepsilon > 0, |\gamma| < 1$) (see Example 2 in [17]), $g_6(\xi) = \frac{2-\mu(1+(1+\mu)\xi)}{(1+\mu\xi)^{5/2}}$ (see Corollary 6.2 in [11]) and $g_4(z) = 0$, then, the following warped type Finsler metric

$$F(x,y) = \sqrt{(y^0)^2 + \varepsilon |\overline{y}|^2} + \gamma y^0 + sg_5(|\overline{x}|) + \frac{\left[1 + (1+\mu)|\overline{x}|^2\right] \left[|\overline{y}|^2 |\overline{x}|^2 - \langle \overline{x}, \overline{y} \rangle^2\right] + \langle \overline{x}, \overline{y} \rangle^2}{(1+\mu|\overline{x}|^2)\sqrt{|\overline{y}|^2 + \mu(|\overline{x}|^2|\overline{y}|^2 - \langle \overline{x}, \overline{y} \rangle)}},$$

is locally projectively flat. Here, $g_5(|\overline{x}|)$ is such that $\phi > 0$, for example $g_5(r) = \frac{2\sqrt{1+(1+\mu)r^2}}{(1+\mu r^2)^2}$.

Example 2. In Corollary 1, by choosing $g_1(z) = \sqrt{z^2 + \varepsilon}$ ($\varepsilon > 0$), $g_6(\xi) = 2\xi^m$, m = 0, 1, ... (see similar form in Theorem 1.1 in [5]) and $g_4(z) = 0$, we have that

(4.1)
$$\phi = k + \sqrt{z^2 + \varepsilon} + sg_5(r) + \frac{(r^2 - s^2)^{m+1}}{m+1} + \frac{2s}{1+2m}I_m(r,s),$$

where $I_m = x(r^2 - s^2)^m + 2mr^2 I_{m-1}$, with $I_0(r,s) = s$. Then, the following warped type Finsler metric on $\mathbb{R} \times \mathbb{B}^n$,

$$F(x,y) := |\overline{y}| \phi\left(\frac{y^0}{|\overline{y}|}, |\overline{x}|, \frac{\langle \overline{x}, \overline{y} \rangle}{|\overline{y}|}\right)$$

is locally projectively flat.

Acknowledgements. The first author would like to thank Prof. Keti Tenenblat for helpful conversations during her postdoctoral project at the Universidade de Brasilia (Brazil). Her suggestions were invaluable.

NEWTON SOLÓRZANO AND VÍCTOR LEÓN

References

- Bácsó, S., Matsumoto M.: On Finsler spaces of Douglas type. A generalization of the notion of Berwald space, Publ. Math. Debrecen, 51, 385-406 (1997).
- Chen, B., Shen, Z., Zhao, L.: Constructions of einstein finsler metrics by warped product, International Journal of Mathematics, 29(11): 1850081 (2018) https://doi.org/10.1142/S0129167X18500817
- [3] Douglas, J.: The general geometry of paths, Ann. Math. 29, 143-168 (1927-28).
- [4] Hamel, G.: Über die Geometrieen in denen die Geraden die K
 ürzesten sind, Math. Ann. 57, 231-264 (1903).
- [5] Huang, L., and Mo, X.: On spherically symmetric Finsler metrics of scalar curvature, J. Geom. Phys. 62 (2012), 2279-2287.
- Huang, L., Mo, X.: Projectively flat Finsler metrics with orthogonal invariance, Ann. Polon. Math. 107, 259-270 (2013). https://doi.org/10.4064/ap107-3-3
- [7] Kozma, L., Peter, R., Varga, C.: Warped product of Finsler manifolds. Ann. univ. Sci. Budapest 44, 157-170 (2001).
- [8] Liu, H., Mo, X.: Finsler Warped Product Metrics of Douglas Type. Canadian Mathematical Bulletin, 62(1), 119-130 (2019). https://doi.org/10.4153/CMB-2017-077-0
- [9] Liu, H., Mo, X. On Projectively Flat Finsler Warped Product Metrics of Constant Flag Curvature. J. Geom. Anal. 31, 11471–11492 (2021). https://doi.org/10.1007/s12220-021-00690-5
- [10] Marcal, P., Shen, Z.: Ricci flat Finsler metrics by warped product. Preprint, arXiv:2012.05699.
- [11] Mo, X., Solórzano, N. M., Tenenblat, K.: On spherically symmetric Finsler metrics with vanishing Douglas curvature. Differential Geom. Appl. 31, 746–758 (2013). https://doi.org/10.1016/j.difgeo.2013.09.002
- [12] S. S. Chern and Z. Shen: Riemannian-Finsler geometry. World Scientific Publisher, Singapore, 2005.
- [13] Li, B., Shen, Z.: On a class of locally projectively flat Finsler metrics. International Journal of Mathematics, 27(06), 1650052 (2016).
- [14] Zhou, L.: Spherically symmetric Finsler metrics in \mathbb{R}^n . Publ. Math. Debrecen 80(1-2), 67-77 (2012).
- [15] Shen, Z.: Landsberg curvatur, S-curvature and Riemann curvature in: A Sampler of Finsler Geometry, MSRI series, 50 (Cambridge University Press, Cambridge, 2004).
- [16] Shen, Z: On Projectively Flat (α, β) -metrics. Bulletin Canadien de Mathématiques, **52**(1), 132–144, (2009). doi:10.4153/cmb-2009-016-2
- [17] Solórzano, N.M: On Douglas warped product metrics. Preprint, arXiv:2110.00875v3.

N. SOLÓRZANO. ILACVN - CICN, UNIVERSIDADE FEDERAL DA INTEGRAÇÃO LATINO-AMERICANA, PAR-QUE TECNOLÓGICO DE ITAIPU, FOZ DO IGUAÇU-PR, 85867-970 - BRAZIL *Email address*: nmayer1590gmail.com

V. LEÓN. ILACVN - CICN, UNIVERSIDADE FEDERAL DA INTEGRAÇÃO LATINO-AMERICANA, PARQUE TECNOLÓGICO DE ITAIPU, FOZ DO IGUAÇU-PR, 85867-970 - BRAZIL

Email address: victor.leon@unila.edu.br

10