

History states of one-dimensional Quantum Walks

F. Lomoc¹, A.P. Boette¹, N. Canosa¹, R. Rossignoli^{1,2}

¹ *Instituto de Física La Plata, CONICET, and Departamento de Física, Universidad Nacional de La Plata, C.C. 67, La Plata (1900), Argentina*

² *Comisión de Investigaciones Científicas (CIC), La Plata (1900), Argentina*

We analyze the application of the history state formalism to quantum walks. The formalism allows one to describe the whole walk through a pure quantum history state, which can be derived from a timeless eigenvalue equation. It naturally leads to the notion of system-time entanglement of the walk, which can be considered as a measure of the number of orthogonal states visited in the walk. We then focus on one-dimensional discrete quantum walks, where it is shown that such entanglement is independent of the initial spin orientation for real Hadamard-type quantum coins and real initial states (in the standard basis) with definite site-parity. Moreover, in the case of an initially localized particle it can be identified with the entanglement of the unitary global operator that generates the whole history state, which is related to its entangling power and can be analytically evaluated. Besides, it is shown that the evolution of the spin subsystem can also be described through a spin history state with an extended clock. A connection between its average entanglement (over all initial states) and that of the operator generating this state is also derived. A quantum circuit for generating the quantum walk history state is as well provided.

I. INTRODUCTION

Quantum Walks (QW) were first introduced in [1] as a quantum counterpart of classical random walks. While related ideas can be traced back to the works of Feynman on the discretized Dirac equation [2], interest on QW has grown enormously in the last decades due their relevance in the field of quantum computation and information [3, 4]. They are useful for the development of quantum algorithms [5], with QW based methods [6–8] achieving similar speedup to that of the renowned Grover search algorithm [9]. It has also been proven that both continuous and discrete time QW are universal for quantum computation [10, 11]. They have been employed in network analysis [12] and quantum simulation [13–16], as well as for modeling some biological processes [17, 18]. Besides, QW can be simulated by means of different experimental platforms, such as cold atoms in optical lattices [19–21], trapped ions [22, 23] and photonic setups [24, 25]. Entanglement in QW is also a topic of interest, so far mainly focused on that between coin (spin) and position degrees of freedom [26–36].

The aim of this work is to apply the history state formalism [37–44] to QW. In this formalism, originally proposed by Page and Wootters [37], time is incorporated through a reference quantum clock and the system evolution emerges from an entangled system-clock history state which fulfills a timeless Wheeler-DeWitt-like equation [45]. The approach has attracted much interest in recent years in different areas, including nonrelativistic quantum mechanics (QM) [40–44, 46–49] as well as quantum gravity and relativistic QM [38, 39, 50–53].

In this work we will focus on one-dimensional discrete QW [3, 4, 54–56], in which a spin 1/2 quantum particle undergoes a unitary evolution in a discrete homogeneous lattice in discrete time steps, according to a translation rule controlled at each step by the value of its spin component and the action of a quantum gate (“coin”) on the

spin. We will analyze the walk from the new perspective provided by the history state formalism, here applied in its discrete version [42–44]. The ensuing history state is a pure state of the composite system comprising both the position and spin degrees of freedom on one side (system S), and a quantum clock system T on the other. It contains the information of the whole QW and satisfies a timeless eigenvalue equation, such that the state of S at step n can be obtained by conditioning the history state with the corresponding state of the clock T . It can be generated from an initial product state through a quantum circuit.

The corresponding system-time entanglement is a measure of the degree of evolution of S , i.e., of the number of orthogonal states visited in the QW, and will be shown to be fully determined by the overlaps between the evolved states. Notice that this entanglement is not the spin-position entanglement usually considered [26–28]. We then show that for real Hadamard-type coins such entanglement becomes independent of the initial spin state for a wide class of initial position states, including the standard case of an initially localized particle. In the latter, it also shown that this entanglement is the operator entanglement of the quantum gate generating the history state from the initial product state. Such entanglement defines in fact the entangling power of this operator, which determines the average over all initial spin states of the history state entanglement. In the case of the quadratic entropy it can be evaluated analytically, and an asymptotic expression for large N can be derived in terms of the coin parameter, showing explicitly the deviation from the maximum entropy (i.e., maximally entangled) limit. The associated entanglement spectrum is also analyzed.

Besides, it is also possible to define a spin history state by considering a different partition of the whole system, with the spin on one side and a composite clock (the original clock plus the position degree of freedom) on the other. In the case of an initially localized particle the

ensuing spin-rest entanglement is shown to be related to that of the unitary operator generating the spin history state. An upper bound to the average spin-rest entanglement is thus obtained, arising from the reduced Schmidt rank of the previous operator.

We first briefly review in Section II the main features of discrete one dimensional QW and their exact evolution. The history state formalism for the QW is introduced in section III, where the main results, together with analytic expressions for overlaps and system-time entanglement entropy are provided. The connection with operator entanglement and the spin history state are discussed in section IV. Illustrative numerical results are shown in both III and IV. Finally conclusions are drawn in V.

II. QUANTUM WALKS IN ONE DIMENSION

A. Generalities

Standard one-dimensional quantum walks are processes in which a quantum ‘‘particle’’ (quantum system) with spin $\frac{1}{2}$ and hence internal Hilbert space $\mathcal{H}_s = \mathbb{C}^2$, moves along a one-dimensional lattice spanned by position eigenstates $|x\rangle$, $x \in \mathbb{Z}$, which generate the position Hilbert space \mathcal{H}_p . The full Hilbert space of the system is then $\mathcal{H}_S = \mathcal{H}_p \otimes \mathcal{H}_s$. At each time step two operations are performed such that their composition gives the unitary evolution from one time to the next. The first one acts on the spin component (quantum coin) and leaves the position unchanged. This is usually taken as a kind of generalized Hadamard transform [56], represented in the standard basis (eigenstates $\{|\uparrow\rangle, |\downarrow\rangle\}$ of σ_z) as

$$C = \begin{pmatrix} \sqrt{\rho} & \sqrt{1-\rho} e^{i\gamma} \\ \sqrt{1-\rho} e^{i\phi} & -\sqrt{\rho} e^{i(\gamma+\phi)} \end{pmatrix}, \quad (1)$$

where γ, ϕ are arbitrary angles and $0 \leq \rho \leq 1$, such that $C^\dagger = C$. We will here focus on traceless ($\gamma = -\phi$) real coins, such that it results in a Hadamard-type coin

$$C = \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix} = \sigma_z \cos \theta + \sigma_x \sin \theta \equiv \sigma_\theta, \quad (2)$$

where $\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$, $\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ are the Pauli matrices and $\cos^2 \theta = \rho$. The usual Hadamard coin is recovered for $\theta = \pi/4$. Actually any traceless coin [$C = \sigma_z \cos \theta + (\sigma_x \cos \phi + \sigma_y \sin \phi) \sin \theta$] can be rewritten in the form (2) by redefining the x and y axes.

The second operation is a conditional one-step displacement to the right (left) if spin is up (down) along z , generated by the translation operator T (T^\dagger). The total operator generating the step is then given by

$$U = \frac{1}{2} [T \otimes (\mathbb{1} + \sigma_z) \sigma_\theta + T^\dagger \otimes (\mathbb{1} - \sigma_z) \sigma_\theta] \quad (3)$$

$$= \sum_x [|x+1\rangle \langle x| \otimes |\uparrow\rangle \langle \uparrow| \sigma_\theta + |x-1\rangle \langle x| \otimes |\downarrow\rangle \langle \downarrow| \sigma_\theta], \quad (4)$$

and verifies the unitary condition $U^\dagger U = \mathbb{1}_p \otimes \mathbb{1}_s$.

B. Exact evolution and overlaps

Assuming an initial state which is localized in space, i.e., with support in some finite interval, it is convenient, for obtaining a closed exact analytic expression of the evolved states, to consider a finite position basis $\{|x\rangle, x = 0, \dots, M-1\}$, with $\langle x|x'\rangle = \delta_{xx'}$, which contains the initial state as well as the evolved states up to, say, N steps ($M > 2N$). Then we define the associated discrete Fourier transformed (DFT) basis

$$|k\rangle = \frac{1}{\sqrt{M}} \sum_{x=0}^{M-1} e^{i2\pi x k/M} |x\rangle, \quad k = 0, \dots, M-1, \quad (5)$$

satisfying $\langle k|k'\rangle = \delta_{kk'}$ and $|-k\rangle = |M-k\rangle$. These states are the eigenstates of the cyclic translation operator defined by $T|x\rangle = |x+1\rangle$, with $T|M-1\rangle = |0\rangle$, such that

$$T|k\rangle = e^{-i2\pi k/M} |k\rangle. \quad (6)$$

The operator (4) can then be rewritten as

$$U = \exp\left[-i\frac{2\pi}{M} K \otimes \sigma_z\right] (\mathbb{1} \otimes \sigma_\theta) \quad (7a)$$

$$= \begin{pmatrix} e^{-i2\pi K/M} & 0 \\ 0 & e^{i2\pi K/M} \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix}, \quad (7b)$$

where K is the discrete ‘‘momentum’’ operator satisfying $K|k\rangle = k|k\rangle$, such that $T = \exp[-i2\pi K/M]$.

For each value k of K , (7b) represents, in the standard spin basis, a unitary operator U_k in spin space fulfilling $\text{Det}[U_k] = -1$, with eigenvalues

$$\lambda_k^\pm = \pm e^{\mp i\omega_k} \quad (8a)$$

$$= \pm \sqrt{1 - \cos^2 \theta \sin^2 \phi_k} - i \cos \theta \sin \phi_k, \quad (8b)$$

where $\phi_k = 2\pi k/M$, and eigenvectors

$$|s_k^\pm\rangle = \alpha_k^\pm |\uparrow\rangle + \beta_k^\pm |\downarrow\rangle, \quad \frac{\beta_k^\pm}{\alpha_k^\pm} = \frac{e^{i\phi_k} \lambda_k^\pm - \cos \theta}{\sin \theta}, \quad (9)$$

satisfying $\langle s_k^\nu | s_{k'}^{\nu'} \rangle = \delta_{kk'} \delta^{\nu\nu'}$. Thus,

$$U = \sum_{k=0}^{M-1} |k\rangle \langle k| \otimes U_k, \quad (10a)$$

$$U_k = e^{-i\omega_k} |s_k^+\rangle \langle s_k^+| - e^{i\omega_k} |s_k^-\rangle \langle s_k^-|. \quad (10b)$$

Whereas in the position representation (4) it is the spin which appears as controlling the position displacement, in the momentum representation (10a) based on the eigenbasis of T , it is the momentum K which controls the spin evolution at each step.

Using (10) we can now determine the evolution of a general initial product state

$$|\Psi_0\rangle = |\psi_0\rangle \otimes |\chi_0\rangle, \quad (11a)$$

$$|\psi_0\rangle = \sum_x \psi_0(x) |x\rangle = \sum_k c_k |k\rangle, \quad (11b)$$

$$|\chi_0\rangle = \alpha |\uparrow\rangle + \beta |\downarrow\rangle, \quad (11c)$$

where $\psi_0(x)$ is the initial position state, with

$$c_k = \langle k | \psi_0 \rangle = \frac{1}{\sqrt{M}} \sum_x e^{-i2\pi x k / M} \psi_0(x) \quad (12)$$

the DFT of $\psi_0(x)$ (sums over k, x are from 0 to $M - 1$) and $|\chi_0\rangle$ the initial spin state. The state after n steps is

$$|\Psi_n\rangle = U^n |\Psi_0\rangle = \sum_k c_k |k\rangle \otimes U_k^n |\chi_0\rangle, \quad (13)$$

where the evolved spin state for momentum k is

$$U_k^n |\chi_0\rangle = e^{-in\omega_k} \langle s_k^+ | \chi_0 \rangle |s_k^+\rangle + e^{in\omega_k} \langle s_k^- | \chi_0 \rangle |s_k^-\rangle. \quad (14)$$

A quantity of most importance in this work is the overlap between the evolved states,

$$\langle \Psi_{n'} | \Psi_n \rangle = \langle \Psi_0 | U^{n-n'} | \Psi_0 \rangle = \langle \Psi_0 | \Psi_{n-n'} \rangle \quad (15a)$$

$$= \sum_k |c_k|^2 \langle \chi_0 | U_k^{n-n'} | \chi_0 \rangle \quad (15b)$$

$$= \sum_k |c_k|^2 \sum_{\nu=\pm 1} \nu^{n-n'} e^{i\nu(n-n)\omega_k} |\langle s_k^\nu | \chi_0 \rangle|^2, \quad (15c)$$

which depends just on $n - n'$.

For future use we notice that for $-k \equiv M - k$, $\lambda_{-k}^\pm = \lambda_k^{\pm*}$ ($\omega_{-k} = -\omega_k$) and $|s_{-k}^\pm\rangle = |s_k^\pm\rangle^*$ in the standard basis ($\alpha_{-k}^\pm = \alpha_k^{\pm*}$, $\beta_{-k}^\pm = \beta_k^{\pm*}$ in (9)). Moreover,

$$U_{-k} = -\sigma_y U_k \sigma_y, \quad (16)$$

entailing $U_{-k}^n = (-1)^n \sigma_y U_k^n \sigma_y$. It is also evident from (7b) that for M even,

$$U_{k+M/2} = -U_k. \quad (17)$$

We also notice that in the special case $\theta = \pi/2$, $\lambda_k^\pm = \pm 1$ ($\omega_k = 0$) and $U_k^2 = \mathbb{1}_s \forall k$, i.e. $U^2 = \mathbb{1}$: the system always returns to its initial configuration after two steps, as $C = \sigma_x$ (Eq. (2)) flips the coin at each step.

III. QUANTUM WALK HISTORY STATES

A. Definition and main properties

Let us now consider a quantum walk with $N - 1$ steps, starting from an initial state $|\Psi_0\rangle$ at time $t_0 = 0$ and ending in a state $|\Psi_{N-1}\rangle$ at time $t_{N-1} = (N - 1)\tau$, where τ is a certain time scale. We then consider a quantum clock system T with an orthogonal set of states $|n\rangle$, $n = 0, \dots, N - 1$, representing the scaled time $n = t_n/\tau$ at which n^{th} step takes place. They are eigenstates of a clock operator T satisfying $T|n\rangle = n|n\rangle$, and could represent, for instance, the scaled position of the clock's needle.

We now define the quantum walk history state as

$$|\Psi\rangle = \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} |\Psi_n\rangle \otimes |n\rangle, \quad (18)$$

where $|\Psi_n\rangle$ is the system state (13) at step n . The state (18) contains the whole information of the walk. For example, the time average of an observable O_S of the particle over the complete walk can be expressed as

$$\langle O_S \rangle := \frac{1}{N} \sum_{n=0}^{N-1} \langle \Psi_n | O_S | \Psi_n \rangle \quad (19a)$$

$$= \langle \Psi | O_S \otimes \mathbb{1}_T | \Psi \rangle, \quad (19b)$$

whereas matrix elements between system states at any two times can be written as

$$\langle \Psi_{n'} | O_S | \Psi_n \rangle = N \langle \Psi | (O_S \otimes |n'\rangle \langle n|) | \Psi \rangle. \quad (20)$$

Using Eqs. (13) and (18), the quantum walk history state can be generated from an initial product state $|\psi_0\rangle |\chi_0\rangle |n = 0\rangle$ through the schematic circuit of Fig. 1, where FT implements the transformation $|x\rangle \rightarrow \frac{1}{\sqrt{M}} \sum_k e^{-i2\pi x k / M} |k\rangle$. For a particle initially localized at $x = 0$, $c_k = 1/\sqrt{M} \forall k$ and one may replace the FT gate by a simpler Hadamard gate $H^{\otimes m'}$, with $2^{m'} = M$.




FIG. 1. Schematic circuit representing the generation of the history state (18) through Eq. (13): The Quantum Fourier Transform (FT) is applied to the initial position state $|\psi_0\rangle$, while $H^{\otimes m}$ denotes the Hadamard operator over m qubits, with $2^m = N$, such that $H^{\otimes m}|0\rangle = \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} |n\rangle$. Finally the control- U gate U_k^n acts over the initial spin state $|\chi_0\rangle$.

History states actually *determine* the system evolution when they satisfy a proper *timeless* eigenvalue equation [44]. Defining the system-clock unitary operator

$$\mathcal{U} = \sum_{n=1}^N U_{n,n-1} \otimes |n\rangle \langle n-1|, \quad (21)$$

where $U_{n,n-1} = U$ for $1 \leq n \leq N - 1$ and (identifying $|N\rangle$ with $|0\rangle$) $U_{N,N-1} = (U^{N-1})^\dagger$, the state (18) is first seen to be an exact eigenstate of \mathcal{U} with eigenvalue 1:

$$\mathcal{U}|\Psi\rangle = |\Psi\rangle. \quad (22)$$

Conversely, if $|\Psi\rangle$ is *any* state satisfying Eq. (22), i.e. a state which remains fully invariant under \mathcal{U} , then

$$\langle n | \Psi \rangle = \langle n | \mathcal{U} | \Psi \rangle = U_{n,n-1} \langle n-1 | \Psi \rangle, \quad (23)$$

implying that system states $|\Psi_n\rangle := \sqrt{N} \langle n | \Psi \rangle$ will fulfill the discrete unitary evolution $|\Psi_n\rangle = U_{n,n-1} |\Psi_{n-1}\rangle$, i.e.,

Eq. (13) for constant $U_{n,n-1} = U$ ($1 \leq n \leq N-1$). Normalization $1 = \langle \Psi | \Psi \rangle = \sum_n \langle \Psi | n \rangle \langle n | \Psi \rangle$ then entails $\langle \Psi_n | \Psi_n \rangle = 1$. The whole system evolution up to step $N-1$ is thus determined by Eq. (22) and the initial state $|\Psi_0\rangle$. The same holds for general unitaries $U_{n,n-1}$ provided $U_{N,N-1} \dots U_{1,0} = \mathbb{1}_S$.

Writing $\mathcal{U} = \exp[-i\mathcal{J}]$, Eq. (22) implies $\mathcal{J}|\Psi\rangle = 0$ (or $2k\pi$, k integer), which is a discrete Wheeler-DeWitt-type equation [45]. In fact, if $U_{n,n-1} = U \forall n = 1, \dots, N$ (such that $U^N = \mathbb{1}$) then $\mathcal{J} = H \otimes \mathbb{1}_T + \mathbb{1}_S \otimes P$, where $e^{-iH} = U$ determines the step evolution in S and $e^{-iP} = \sum_{n=1}^N |n\rangle \langle n-1|$ generates the unit translation in the time basis ($|n-1\rangle \rightarrow |n\rangle$) [44].

B. System-time entanglement

The entanglement $E(S, T)$ of the history state (18) can be regarded as a measure of the “*degree of evolution*” of system S , i.e., of the number of orthogonal states visited in the walk: If $|\Psi_0\rangle$ is an eigenstate of U , i.e. a *stationary* state satisfying $U|\Psi_0\rangle = e^{i\gamma}|\Psi_0\rangle$, the history state becomes *separable*: $|\Psi\rangle = |\Psi_0\rangle (\frac{1}{\sqrt{N}} \sum_n e^{in\gamma} |n\rangle)$, and $E(S, T) = 0$ [43]. This case arises here in finite cyclic realizations when $|\psi_0\rangle$ in (14) remains invariant under translations, i.e., when it is a state $|k\rangle$, and $|\chi_0\rangle$ coincides with one of the eigenstates $|s_k^\pm\rangle$ of U_k , such that $U^n |\Psi_0\rangle = e^{\mp in\omega_k} |\Psi_0\rangle$ for $|\Psi_0\rangle = |k\rangle \otimes |s_k^\pm\rangle$.

The opposite situation is an evolution where at each step the system evolves into a new orthogonal state such that $\langle \Psi_n | \Psi_{n'} \rangle = \delta_{nn'}$, in which case $E(S, T)$ is *maximum*. This evolution arises here in the trivial limit $\theta \rightarrow 0$ for an initially localized state $\psi_0(x) = \delta_{xx_0}$ with definite spin along the z axis ($\sigma_z |\chi_0\rangle = \pm |\chi_0\rangle$), such that the particle advances always in the same direction.

In the general case $E(S, T)$ will be determined by the overlaps (15). We can expand (18) as ($\nu = \pm 1$)

$$|\Psi\rangle = \frac{1}{\sqrt{N}} \sum_{k,\nu,n} \langle k, s_k^\nu | \Psi_n \rangle |k, s_k^\nu\rangle \otimes |n\rangle \\ = \sum_{m=1}^{n_s} \sqrt{\lambda_m} |m_S\rangle \otimes |m_T\rangle, \quad (24)$$

where $|k, s_k^\pm\rangle = |k\rangle \otimes |s_k^\pm\rangle$ and (24) is its *Schmidt representation*, obtained from the singular value decomposition (SVD) $M = VDW^\dagger$ of the matrix of elements $M_{k\nu,n} = \langle k, s_k^\nu | \Psi_n \rangle / \sqrt{N}$. Here $D_{mm'} = \sqrt{\lambda_m} \delta_{mm'}$ with λ_m the eigenvalues of the positive semidefinite matrices MM^\dagger or equivalently $M^\dagger M$ (fulfilling $\text{Tr } M^\dagger M = \sum_m \lambda_m = 1$) while V, W are unitary matrices diagonalizing MM^\dagger and $M^\dagger M$ respectively, satisfying $VD = MW$.

In (24) $|m_S\rangle = \sum_{k,\nu} V_{k\nu,m} |k, s_k^\nu\rangle$, $|m_T\rangle = \sum_n W_{nm}^* |n\rangle$ are orthogonal system and clock states respectively ($\langle m_S(T) | m'_S(T) \rangle = \delta_{mm'}$) while n_s is the *Schmidt rank*, i.e. the number of nonzero eigenvalues λ_m , which is just the rank of the matrix M . $|\Psi\rangle$ is then entangled iff $n_s \geq 2$ and separable (product state) if $n_s = 1$.

Eq. (24) shows that the states $|m_S(T)\rangle$ are the eigenvectors of the reduced system and clock states

$$\rho_{S(T)} = \text{Tr}_{T(S)} |\Psi\rangle \langle \Psi| = \sum_{m=1}^{n_s} p_m |m_S(T)\rangle \langle m_S(T)|, \quad (25)$$

which determine the average along the walk of any local observable ($\langle \Psi | O_S \otimes \mathbb{1} | \Psi \rangle = \text{Tr}_S \rho_S O_S$), and have the same nonzero eigenvalues λ_m . Their entropies are then identical and define the *entanglement entropy* of the history state $|\Psi\rangle$ (system-time entanglement [43])

$$E(S, T) = S(\rho_S) = S(\rho_T) = \sum_m f(\lambda_m), \quad (26)$$

where the last expression holds for a general trace-form entropy $S(\rho) = \text{Tr } f(\rho)$, where f is a concave nonnegative function satisfying $f(0) = f(1) = 0$. Eq. (26) vanishes iff the history state is separable.

The reduced states (25) can be here also written as

$$\rho_S = \frac{1}{N} \sum_{n=0}^{N-1} |\Psi_n\rangle \langle \Psi_n|, \quad (27a)$$

$$\rho_T = \frac{1}{N} \sum_{n,n'} \langle \Psi_{n'} | \Psi_n \rangle |n\rangle \langle n'|. \quad (27b)$$

Eq. (27b) shows explicitly that ρ_T , and hence its eigenvalues λ_m and the entanglement (26) of the history state, are *fully determined by the overlaps* (15).

The standard choice for S is the von Neumann entropy

$$S(\rho) = -\text{Tr } \rho \log_2 \rho, \quad (28)$$

which will satisfy

$$0 \leq E(S, T) \leq \log_2 N, \quad (29)$$

such that $2^{E(S,T)}$ is essentially a measure of the number of distinct orthogonal states visited in the evolution. Another convenient choice is the quadratic entropy (also denoted as linear entropy or $q = 2$ Tsallis entropy [57]), which is simply determined by the purity $\text{Tr } \rho^2$:

$$S_2(\rho) = 1 - \text{Tr } \rho^2, \quad (30)$$

and corresponds to $f(\rho) = \rho(\mathbb{1} - \rho)$. It does not require the explicit determination of eigenvalues and can then be here directly evaluated in terms of the overlaps (15):

$$E_2(S, T) = 1 - \frac{1}{N^2} \sum_{n,n'} |\langle \Psi_{n'} | \Psi_n \rangle|^2 \quad (31a)$$

$$= 1 - \frac{1}{N} \left[1 + 2 \sum_{n=1}^{N-1} \left(1 - \frac{n}{N} \right) |\langle \Psi_0 | \Psi_n \rangle|^2 \right]. \quad (31b)$$

It obviously satisfies

$$0 \leq E_2(S, T) \leq 1 - \frac{1}{N}, \quad (32)$$

such that $\frac{1}{1-E_2(S,T)}$ is here the effective number of orthogonal states visited. A directly related quantity is the $q = 2$ Renyi entropy $S_2^R(\rho) = -\log_2(1 - S_2) = -\log_2 \text{Tr} \rho^2$ [58], which satisfies the same bound (29).

The upper limit in (29)–(32) is reached for $\theta \rightarrow 0$ in (2) and an initially localized particle with definite σ_z . On the other hand, for $\theta = \pi/2$ the evolution becomes periodic with period 2 (as $U^2 = \mathbb{1}$) and hence the system-time entanglement entropy will stay trivially bounded $\forall N$:

$$E(S, T) \leq 1, \quad E_2(S, T) \leq 1/2 \quad (\theta = \pi/2), \quad (33)$$

with the upper limit reached for an orthogonal intermediate state. Thus, by varying θ in the interval $[0, \pi/2]$ we can reach, for an initially localized particle, all possible rates of system-time entanglement increase with N , from the maximum rate for $\theta = 0$ to the null increase for $\theta = \pi/2$, entailing in general a decrease of $E(S, T)$ with increasing θ in this interval.

C. Independence of system-time entanglement from initial spin state

1. Real initial states with definite site parity

We now examine the entanglement (26) for some general types of initial states $|\Psi_0\rangle$. We first notice that if $|\Psi_0\rangle$ has a *definite position parity*, such that its support are just even (or odd) sites x ,

$$(e^{i\pi X} \otimes \mathbb{1})|\Psi_0\rangle = \pm|\Psi_0\rangle, \quad (34)$$

the overlap $\langle\Psi_n|\Psi_{n'}\rangle$ will vanish for $n - n'$ odd since at each step the particle will move to sites of opposite parity:

$$(e^{i\pi X} \otimes \mathbb{1})|\Psi_n\rangle = \pm(-1)^n|\Psi_n\rangle. \quad (35)$$

Eq. (34) is trivially fulfilled for an initially localized particle $\psi_0(x) = \delta_{x,x_0}$. In momentum space, Eq. (34) implies $c_{k+M/2} = \pm c_k$ (M even) and (35) follows from Eqs. (17) and (13).

Then, for *real* $\psi_0(x)$ and α, β in (11), such that $c_{-k} = c_k^*$ and $\langle s_{-k}^\pm|\chi_0\rangle = \langle s_k^\pm|\chi_0\rangle^*$, we obtain, using $\sum_{\nu=\pm} |\langle s_k^\nu|\chi_0\rangle|^2 = 1$ and Eq. (15c),

$$\begin{aligned} \langle\Psi_0|\Psi_n\rangle &= \sum_k |c_k|^2 \cos(n\omega_k) (|\langle s_k^+|\chi_0\rangle|^2 + (-1)^n |\langle s_k^-|\chi_0\rangle|^2) \\ &= \begin{cases} \sum_k |c_k|^2 \cos(n\omega_k), & n \text{ even} \\ 0, & n \text{ odd} \end{cases}. \end{aligned} \quad (36)$$

Thus, the overlap becomes *independent of the initial (real) spin state* $|\chi_0\rangle \forall n$. This implies a system-time entanglement entropy *independent* of $|\chi_0\rangle$.

Previous result can be seen more clearly using Eq. (16) and (15b): since $\sigma_y|\chi_0\rangle$ is orthogonal to $|\chi_0\rangle$ for $|\chi_0\rangle$ real (α, β real in (11c), equivalent to $\langle\sigma\rangle$ in the x, z plane) and

$\langle\chi_0|U_{-k}^n|\chi_0\rangle = (-1)^n \langle\chi_0|\sigma_y U_k^n \sigma_y|\chi_0\rangle$, for $|c_{-k}| = |c_k|$ and n even we obtain

$$\begin{aligned} \langle\Psi_0|\Psi_n\rangle &= \frac{1}{2} \sum_k |c_k|^2 (\langle\chi_0|U_k^n|\chi_0\rangle + \langle\chi_0|\sigma_y U_k^n \sigma_y|\chi_0\rangle) \\ &= \frac{1}{2} \sum_k |c_k|^2 \text{Tr} U_k^n = \frac{1}{2} \langle\psi_0|\text{Tr}_S U^n|\psi_0\rangle, \end{aligned} \quad (37)$$

which shows that the even overlap depends just on the *trace* of U_k^n , i.e. on the partial trace of U^n over spin, and is hence independent of the initial spin state $|\chi_0\rangle$. Eq. (37) leads again to (36) for n even and *also* n odd, as $\text{Tr} U_k^n = e^{-in\omega_k} + (-1)^n e^{in\omega_k} = (-1)^n \text{Tr} U_{-k}^n$ and hence the sum in (37) vanishes for n odd when $|c_{-k}| = |c_k|$.

With previous expressions, the sum over N in the quadratic system-time entanglement (31b) can be evaluated analytically:

$$E_2(S, T) = 1 - \sum_{\substack{k, k' \\ \nu=\pm 1}} \frac{|c_k c_{k'}|^2 \sin^2 \frac{N(\omega_k + \nu\omega_{k'})}{2}}{N^2 \sin^2(\omega_k + \nu\omega_{k'})}, \quad N \text{ even} \quad (38a)$$

$$= 1 - \sum_{\substack{k, k' \\ \nu, \nu'=\pm 1}} \frac{|c_k c_{k'}|^2 \sin^2 \frac{(N+\nu')(\omega_k + \nu\omega_{k'})}{2}}{2N^2 \sin^2(\omega_k + \nu\omega_{k'})}, \quad N \text{ odd.} \quad (38b)$$

Here $\frac{\sin^2 mu}{m^2 \sin^2 u}$ is understood as its limit 1 if $\sin u = 0$ (m integer), being a polynomial of degree $m - 1$ in $\cos u$ ($\frac{\sin(n+1)u}{\sin u} = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n-k}{k} (2 \cos u)^{n-2k}$ is a Chebyshev polynomial of the second kind [59]).

As a check, for $\theta = \pi/2$, $\omega_k = 0 \forall k$ and hence, for any initial $\psi_0(x)$ with definite parity, (38a)–(38b) lead to

$$E_2(S, T) = \begin{cases} \frac{1}{2}, & N \text{ even} \\ \frac{1}{2}(1 - \frac{1}{N^2}), & N \text{ odd} \end{cases} \quad (\theta = \pi/2). \quad (39)$$

This means that the system just moves between two orthonormal states (periodic evolution) as previously stated, entailing a non-increasing system-time entanglement entropy. The spectrum of $\rho_S(T)$ is simply $(\frac{1}{2}, \frac{1}{2})$ for N even and $(\frac{N+1}{2N}, \frac{N-1}{2N})$ for N odd.

2. Initially localized particle

For $\psi_0(x) = \delta_{x,x_0}$, $c_k = \frac{1}{\sqrt{M}} e^{-i2\pi x_0 k/M}$ and $|c_k|^2 = \frac{1}{M} \forall k$. Overlaps and system-time entanglement are then determined just by the full trace of U^n and hence the coin angle θ , as implied by (37):

$$\langle\Psi_0|\Psi_n\rangle = \frac{\text{Tr}[U^n]}{2M} = \begin{cases} \frac{1}{M} \sum_k \cos(n\omega_k), & n \text{ even} \\ 0, & n \text{ odd} \end{cases}. \quad (40)$$

We can now evaluate (40) *exactly* $\forall \theta$ and N , using (8). Since the result is independent of M (for $M > 2N$ if $x_0 = M/2$) it can be obtained either summing over k or letting $M \rightarrow \infty$ and replacing the sum by an integral




FIG. 2. System-time entanglement according to the von Neumann entropy (top) and the quadratic entropy (bottom), as a function of the total number N of steps of the quantum walk for different values $\theta = 0, \pi/20, \dots, \pi/2$, of the coin parameter, for an initially localized particle and any real initial spin state.

over $\phi = 2\pi k/M$ (with $d\phi = 2\pi/M$). We obtain

$$\langle \Psi_0 | \Psi_{2n} \rangle = P_n(\cos^2 \theta), \quad (41a)$$

$$P_n(u) = \sum_{j=0}^n (-1)^j \binom{n}{j} \binom{n+j-1}{j} u^j \quad (41b)$$

$$= 1 - n^2 u + \frac{1}{4} n^2 (n^2 - 1) u^2 + \dots \quad (41c)$$

where $P_n(u) = F_{2,1}(-n, n, 1, u) = P_n^{0,-1}(1-2u)$ is a polynomial of degree n in u , with $F_{2,1}$ the hypergeometric function and $P_n^{\alpha,\beta}$ the Jacobi polynomial [59]. It satisfies $P_n(0) = 1$ and $P_n(1) = 0 \forall n \geq 1$.

For large n and $\theta \in (0, \pi/2)$, we can also obtain from (41) and the asymptotics of Jacobi polynomials, the *exact asymptotic expression*

$$\langle \Psi_0 | \Psi_{2n} \rangle \approx (-1)^n \sqrt{\frac{\tan \theta}{n\pi}} \cos(2n\theta + \pi/4) + O(n^{-3/2}), \quad (42)$$

which shows that the overlap fades away as $n^{-1/2}$ for large n and its modulus essentially increases as $\tan^{1/2} \theta$ for increasing $\theta \in (0, \pi/2)$.

Through Eqs. (40)–(41) the quadratic system-time en-




FIG. 3. System-time entanglement according to the von Neumann entropy (top) and quadratic entropy (bottom), as a function of the coin angle θ for different number of steps $N = 2, 10, 20, \dots, 100$, for the same initial states of Fig. 2.

tanglement entropy (31) can be evaluated exactly as:

$$E_2^N(S, T) = 1 - \frac{1}{(2MN)^2} \sum_{n, n'} |\text{Tr}[U^{n-n'}]|^2 \quad (43a)$$

$$= 1 - \frac{1}{N} \left[1 + 2 \sum_{n=1}^{\lfloor \frac{N-1}{2} \rfloor} \left(1 - \frac{2n}{N} \right) P_n^2(\cos^2 \theta) \right]. \quad (43b)$$

As a check, it is verified that for $\theta = 0$, we obtain from (43b) maximum entropy $\forall N$, as $P_n(1) = 0 \forall n > 1$:

$$E_2(S, T) = 1 - \frac{1}{N}, \quad (\theta = 0), \quad (44)$$

in agreement with previous considerations (any entropy is obviously also maximum in this case). And for $\theta = \pi/2$, we recover from (43b) Eq. (39) for both N even or odd, as $P_n(0) = 1 \forall n \geq 1$. Exact results as a function of N and θ are depicted in Figs. 2–3.

For large N , we may use Eq. (42) for $n \neq 0$ and approximate the sum over n in (43b) by an integral over $u = 2n/N$, which leads to the asymptotic expression

$$E_2^N(S, T) \approx 1 - \frac{1}{N} \left[1 + \frac{2}{\pi} \left(\ln \frac{N}{2} + \text{Si}(4\theta) - c \right) \tan \theta \right], \quad (45)$$

where $\text{Si}(x) = \int_0^x \frac{\sin t}{t} dt$ and $c = 1 + \frac{\pi}{2}$ (neglecting terms $O(N^{-2})$). The deviation from maximum entropy $S_2^{\max} = 1 - \frac{1}{N}$ is then $O(\frac{\ln N}{N})$ and proportional to $\tan \theta$, in agreement with previous considerations. This result can also

be obtained from (38) using $\frac{\sin^2(Nu/2)}{N \sin^2 u/2} \xrightarrow{N \rightarrow \infty} 2\pi\delta(u)$ for $|u| < \pi$ and integrating over $\phi = \frac{2\pi k}{m}$. An exact summation using (42) is given in the appendix.

3. Entanglement spectrum

For an initially localized particle, we may also examine the entanglement spectrum, i.e., the common eigenvalues of the reduced densities (27) which determine the entropies of Figs. 2–3, by diagonalizing the overlap matrix (Eqs. (40)–(41))

$$\langle \Psi_{n'} | \Psi_n \rangle = \begin{cases} P_{\frac{|n-n'|}{2}}(\cos^2 \theta) & , n - n' \text{ even} \\ 0 & , n - n' \text{ odd} \end{cases} \quad (46)$$

for $0 \leq n, n' \leq N - 1$. This leads to two similar blocks (n, n' even or odd respectively), identical for N even.




FIG. 4. Entanglement spectrum (eigenvalues of the reduced densities ρ_S or ρ_T) obtained from the overlap matrix (46) for $N = 20$ as a function of the coin parameter θ .

For sufficiently large N the eigenvalues of each block come essentially in almost degenerate pairs for θ not close to $\pi/2$, with the number of non-negligible eigenvalues decreasing with increasing θ , as depicted in Fig. 4. The largest eigenvalue lies close to the Gershgorin upper bound, i.e., using Eq. (42),

$$\lambda_{\max}(\theta) \approx \frac{1}{N} \left[1 + \sqrt{\frac{\tan \theta}{\pi}} \sum_{n=1}^{\frac{N}{2}-1} \frac{|\cos(2n\theta + \pi/4)|}{\sqrt{n}} \right] \quad (47a)$$

$$\approx \frac{1}{N} \left[1 + \sqrt{\frac{\tan \theta}{2\pi}} H_{\frac{1}{2}}\left(\frac{N}{2}\right) \right] \quad (47b)$$

$$\approx \frac{1}{N} + \sqrt{\frac{\tan \theta}{N\pi}} \left[1 + \frac{\zeta\left(\frac{1}{2}\right)}{\sqrt{2N}} \right], \quad (47c)$$

where $H_{1/2}(m) = \sum_{n=1}^m \frac{1}{\sqrt{n}}$ ($\approx 2\sqrt{m} + \zeta(\frac{1}{2})$ for large m) is the generalized harmonic number ($\zeta(1/2) \approx -1.46$ is the Riemann zeta function at $1/2$). Thus, there is always a deviation $O(N^{-1/2})$ from the maximally mixed distribution, proportional to $\tan^{1/2} \theta$.

IV. RELATION WITH OPERATOR ENTANGLEMENT

A. Entanglement of unitary operators

We will show here that the system-time entanglement entropy for the initially localized particle, which is independent of the initial (real) spin state, is the entanglement entropy of the global unitary operator which generates the quantum walk.

First, let us consider a complete set of local orthogonal operators O_i^S (O_j^T) of the system (clock) satisfying

$$\text{Tr}_S [O_i^{S\dagger} O_{i'}^S] = d_S \delta_{ii'}, \quad \text{Tr}_T [O_j^{T\dagger} O_{j'}^T] = d_T \delta_{jj'}, \quad (48)$$

where $d_{S(T)}$ is the Hilbert space dimension of S (T). Any operator \mathcal{U} on the whole system can be expanded as

$$\mathcal{U} = \sum_{i,j} M_{ij} O_i^S \otimes O_j^T, \quad (49)$$

where $M_{ij} = \frac{1}{d_S d_T} \text{Tr} [(O_i^{\dagger} \otimes O_j^{\dagger}) \mathcal{U}]$. Then $\frac{1}{d_S d_T} \text{Tr} [\mathcal{U}^{\dagger} \mathcal{U}] = \sum_{i,j} |M_{ij}|^2 = \text{Tr}[M^{\dagger} M]$. If \mathcal{U} is unitary, $\text{Tr}[M^{\dagger} M] = 1$ and the $|M_{ij}|^2$ become standard probabilities.

Hence, in the same way as done for the history state, through the SVD $M = V D W^{\dagger}$, with $D_{mm'} = \lambda_m \delta_{mm'}$ and V, W unitary, we can also obtain the Schmidt representation of the operator \mathcal{U} ,

$$\mathcal{U} = \sum_{m=1}^{\tilde{n}_s} \sqrt{\tilde{\lambda}_m} \tilde{O}_m^S \otimes \tilde{O}_m^T, \quad (50)$$

where λ_m are the eigenvalues of $M^{\dagger} M$ or $M M^{\dagger}$ and $\tilde{O}_m^S = \sum_i V_{im} O_i^S$, $\tilde{O}_m^T = \sum_j W_{jm}^* O_j^T$ are new local orthogonal operators satisfying (48), with \tilde{n}_s the rank of \tilde{M} . If \mathcal{U} is unitary the eigenvalues λ_m are again standard probabilities ($\lambda_m \geq 0$, $\sum_m \lambda_m = 1$). Thus, for a general trace-form entropy, the entanglement entropy of the unitary operator \mathcal{U} can be defined as

$$E(\mathcal{U}) = \sum_m f(\lambda_m), \quad (51)$$

vanishing iff \mathcal{U} is a product of local unitaries ($\tilde{n}_s = 1$).

The analogy with state entanglement is straightforward if the Choi isomorphism for representing operators is employed. Any operator \mathcal{U} on $\mathcal{H}_{ST} = \mathcal{H}_S \otimes \mathcal{H}_T$ can be associated to a pure state $|\mathcal{U}\rangle$ in $\mathcal{H}_{ST} \otimes \mathcal{H}_{ST}$ given by

$$|\mathcal{U}\rangle = (\mathcal{U} \otimes \mathbb{1}_{ST}) |\mathbb{1}\rangle = \frac{1}{\sqrt{d_S d_T}} \sum_{\mu} (\mathcal{U} |\mu\rangle) \otimes |\mu\rangle \quad (52)$$

where $|\mathbb{1}\rangle = \frac{1}{\sqrt{d_S d_T}} \sum_{\mu} |\mu\rangle \otimes |\mu\rangle$ is a maximally entangled state in $\mathcal{H}_{ST} \otimes \mathcal{H}_{ST}$. In this way, an exact map for inner products is obtained:

$$\langle \mathcal{W} | \mathcal{U} \rangle = \frac{1}{d_S d_T} \text{Tr} [\mathcal{W}^{\dagger} \mathcal{U}], \quad (53)$$

with $\langle \mathcal{U} | \mathcal{U} \rangle = 1$ for a unitary operator \mathcal{U} . Thus, Eqs. (49)–(50) can be recast, noting that $|\mathbb{1}\rangle = |\mathbb{1}_S\rangle \otimes |\mathbb{1}_T\rangle$, as

$$|\mathcal{U}\rangle = \sum_{i,j} M_{ij} |O_i^S\rangle \otimes |O_j^T\rangle = \sum_m \sqrt{\lambda_m} |\tilde{O}_m^S\rangle \otimes |\tilde{O}_m^T\rangle \quad (54)$$

with $|O_i^S\rangle = (O_i^S \otimes \mathbb{1}_S) |\mathbb{1}_S\rangle = \frac{1}{d_S} \sum_\mu (O_i^S |\mu_S\rangle) \otimes |\mu_S\rangle$ and similarly for $|O_j^T\rangle$, such that $M_{ij} = (\langle O_i^S | \otimes \langle O_j^T |) |\mathcal{U}\rangle$. We can now rewrite (51) as

$$E(\mathcal{U}) = S(\rho_S^{\mathcal{U}}) = S(\rho_T^{\mathcal{U}}), \quad (55)$$

where $\rho_{S(T)}^{\mathcal{U}} = \text{Tr}_{T(S)} |\mathcal{U}\rangle \langle \mathcal{U}| = \sum_m \lambda_m |\tilde{O}_m^{S(T)}\rangle \langle \tilde{O}_m^{S(T)}|$ are the local reduced densities derived from $|\mathcal{U}\rangle$. In particular, the quadratic entropy becomes

$$E_2(\mathcal{U}) = 1 - \text{Tr}(\rho_{S(T)}^{\mathcal{U}})^2 = 1 - \text{Tr}[(MM^\dagger)^2]. \quad (56)$$

B. Operator entanglement and quantum walk

If a system initially in a state $|\Psi_0\rangle$ undergoes a discrete unitary evolution through times $n = 0, \dots, N-1$ and states $|\Psi_n\rangle = U_n |\Psi_0\rangle$, its history state can be generated from an initial system-clock product state as (see Fig. 1)

$$|\Psi\rangle = \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} U_n |\Psi_0\rangle \otimes |n\rangle = \mathcal{W}(|\Psi_0\rangle \otimes |\tilde{0}\rangle), \quad (57)$$

where $|\tilde{0}\rangle = H^{\otimes m} |0\rangle = \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} |n\rangle$ and

$$\mathcal{W} = \sum_{n=0}^{N-1} U_n \otimes |n\rangle \langle n| \quad (58)$$

is a control- U_n unitary operator on the whole system. Its state representation (52) is

$$|\mathcal{W}\rangle = \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} |U_n\rangle \otimes |T_n\rangle \quad (59)$$

where $|T_n\rangle = (\sqrt{N} |n\rangle \langle n| \otimes \mathbb{1}) |\mathbb{1}\rangle = |n\rangle \otimes |n\rangle$ and $|U_n\rangle = \frac{1}{\sqrt{d_S}} \sum_\mu (U_n |\mu_S\rangle) \otimes |\mu_S\rangle$. Therefore, the unitary operator \mathcal{W} generating the history state from a product state can itself be represented as an operator history state (59).

The reduced operator state of the clock is here

$$\rho_T^{\mathcal{W}} = \text{Tr}_S |\mathcal{W}\rangle \langle \mathcal{W}| = \frac{1}{N} \sum_{n,n'} \langle U_n | U_{n'} \rangle |T_n\rangle \langle T_{n'}|, \quad (60)$$

in full analogy with (27b), showing again that the entanglement of the generating operator (58)–(59),

$$E(\mathcal{W}) = S(\rho_S^{\mathcal{W}}) = S(\rho_T^{\mathcal{W}}), \quad (61)$$

is fully determined by the overlaps $\langle U_{n'} | U_n \rangle$. In the present random walk, $U_n = U^n$ and

$$\langle U_{n'} | U_n \rangle = \frac{1}{2M} \text{Tr}[U^{n-n'}], \quad (62)$$

is exactly the overlap (40)–(41)–(46) between the evolved system states for an initially localized particle with real initial spin state.

Thus, the operator entanglement (61) is *exactly that of the previous system-clock history state for the initially localized particle*, for any choice of entropy. In particular, the quadratic operator entanglement

$$E_2(\mathcal{W}) = 1 - \text{Tr}(\rho_T^{\mathcal{W}})^2 = 1 - \frac{1}{N^2} \sum_{n,n'} |\langle U_{n'} | U_n \rangle|^2, \quad (63)$$

is just the quadratic S – T entropy (43). And the entanglement spectrum of \mathcal{W} coincides with that of that of the history state (Fig. 4).

The quadratic entropy (63) has an additional meaning: It determines the *entangling power* of the unitary operator \mathcal{W} , i.e. the average quadratic entanglement $E_2(S, T)$ it generates when applied to initial product system-clock states:

$$\begin{aligned} \langle E_2(S, T) \rangle &= \int_{\mathcal{H}} (1 - \text{Tr} \rho_S^2) d\Psi_0 \\ &= \frac{d_S}{d_S + 1} E_2(\mathcal{W}), \end{aligned} \quad (64)$$

with the average taken over the whole set of initial system states $|\Psi_0\rangle$ with the Haar measure $d\Psi_0$ [44]. Thus, for $d_S \gg 1$ the quadratic S – T entanglement entropy (43), equal to (63), is very close to the average value (64).

C. Spin history states in the quantum walk

If we now consider as system only the spin degree of freedom, we first notice from Eq. (13) that for a particle initially localized at $x = 0$, the state after n steps

$$|\Psi_n\rangle = \frac{1}{\sqrt{M}} \sum_k |k\rangle \otimes U_k^n |\chi_0\rangle, \quad (65)$$

is exactly a *spin history state*, with respect to the momentum states $|k\rangle$. The “evolution” operators are here the k -projected unitaries U_k^n acting on the spin, having a nontrivial k -dependence. Thus, the spin-position entanglement at step N is that of a spin history state. The same holds for any initial localization x_0 ($|k\rangle \rightarrow e^{-i2\pi k x_0/M} |k\rangle$).

Therefore, its average $\langle E_2(s, p) \rangle$ over all initial spin states is determined by the entanglement of the operator U^n generating such spin history [from the initial product state $|\Psi_0\rangle = (\frac{1}{\sqrt{M}} \sum_k |k\rangle) |\chi_0\rangle$], itself an operator history state, as is apparent from Eq. (10a):

$$U^n = \sum_{k=0}^{M-1} |k\rangle \langle k| \otimes U_k^n. \quad (66)$$

The associated operator state is $\frac{1}{\sqrt{M}} \sum_k |P_k\rangle |U_k^n\rangle$ with $|P_k\rangle = |k\rangle \otimes |k\rangle$ and $|U_k^n\rangle = \frac{1}{\sqrt{2}} \sum_{\nu=\pm} U_k^n |\nu\rangle |\nu\rangle$ ($|\nu\rangle =$

$\pm\rangle$ are orthogonal spin states). Its entanglement is then determined by the overlaps (see Eqs. (10) and (70) below)

$$\langle U_{k'}^n | U_k^n \rangle = \frac{1}{2} \text{Tr} [U_{k'}^{-n} U_k^n]. \quad (67)$$

While the unitaries U_k^n belong to a four-dimensional space spanned by the identity and the three Pauli operators, limiting the Schmidt rank of (66) to $n_s \leq 4 \forall M$, they actually live within a *three-dimensional* subspace spanned by the identity $\mathbb{1}_s$, the coin $\sigma_\theta \equiv C$ and the orthogonal Pauli operator σ_y : From (7)–(8) we see that

$$U_k = \cos \phi_k \sigma_\theta - i \sin \phi_k \exp[i\theta \sigma_y], \quad (68)$$

since $\sigma_z \sigma_\theta = \cos \theta \mathbb{1}_s + i \sin \theta \sigma_y$. And from (10b) and (68) it follows that

$$|s_k^\pm\rangle \langle s_k^\pm| = \frac{1}{2} (\mathbb{1} \pm \sigma_k), \quad (69a)$$

$$\sigma_k = \frac{\cos \phi_k}{\cos \omega_k} \sigma_\theta + \frac{\sin \phi_k \sin \theta}{\cos \omega_k} \sigma_y, \quad (69b)$$

with $\sigma_k^2 = \mathbb{1}$. Hence all powers

$$U_k^n = e^{-in\omega_k} |s_k^+\rangle \langle s_k^+| + (-1)^n e^{in\omega_k} |s_k^-\rangle \langle s_k^-|$$

$$= \cos(n\omega_k) \mathbb{1} - i \sin(n\omega_k) \sigma_k \quad (n \text{ even}), \quad (70a)$$

$$= -i \sin(n\omega_k) \mathbb{1} + \cos(n\omega_k) \sigma_k \quad (n \text{ odd}), \quad (70b)$$

are spanned just by $\{\mathbb{1}_s, \sigma_\theta, \sigma_y\} \forall k, n$, with $\text{Tr} \sigma_\theta \sigma_y = 0$. This entails a Schmidt rank $n_s \leq 3$ of the n^{th} power (66), implying at most 3 non-zero eigenvalues λ_m in its associated entanglement spectrum, as seen in Fig. 5, and a von Neumann entanglement entropy $E(U^n) \leq \log_2 3$.

For U itself ($n = 1$) just $\{\sigma_\theta, e^{i\theta \sigma_y}\}$ are needed as seen from (68), implying $n_s = 2$, as also evident from the original expression (3). Its entanglement spectrum is just $(1/2, 1/2, 0)$. However, for $n \geq 2$ the rank is $n_s = 3$ if $\theta \in (0, \pi/2)$ (and $M > 3$). Exceptions occur for $\theta = 0$, in which case $\omega_k = \phi_k$ and $\sigma_k = \sigma_\theta = \sigma_z \forall k$, leading to $n_s = 2$ and an entanglement spectrum $(1/2, 1/2, 0) \forall n \geq 1$, as verified in Fig. 5, and also for $\theta = \pi/2$, where $\omega_k = 0 \forall k$ and $\sigma_\theta = \sigma_x$, leading to $n_s = 1 \forall n$ even (as $U_k^n = \mathbb{1} \forall k$) and $n_s = 2$ for n odd (as $U_k^n = \sigma_k$), with spectrum $(1/2, 1/2, 0)$, as also seen in Fig. 5.

Therefore, the entangling power of U^n remains bounded by this rank $n_s \leq 3$: Rescaling the quadratic entropy as $S_2(\rho) = 2(1 - \text{Tr} \rho^2)$, such that $S_2(\rho) = 1$ for a maximally mixed single spin state, it implies $E_2(U^n) \leq 4/3$. Hence, applying the relation (64) to the spin history, the average over all initial spin states of the spin-position entanglement after n steps satisfies

$$\langle E_2(s, p) \rangle = (2/3) E_2(U^n) \leq 8/9. \quad (71)$$

This bound is obviously lower than the maximum $\langle E_2(s, p) \rangle = 1$ reached for a full rank maximally entangled operator U_n ($E_2(U_n) = 3/2$ for the rescaled S_2), for which $|\Psi_n\rangle$ in (65) would be *maximally entangled* ($E_2(s, p) = 1$) for *any* spin state $|\chi_0\rangle$, in agreement with the general results of [44]. The variation of $\langle E_2(s, p) \rangle \propto E_2(U)$ with θ is depicted in Fig. 6.




FIG. 5. Entanglement spectrum of the unitary operator (66) for an odd (top) and even (bottom) N as a function of the coin parameter θ .




FIG. 6. The average over all initial spin states of the spin-position quadratic entanglement entropy $\langle E_2(s, p) \rangle = (2/3) E_2(U^n)$ (Eq. (71)), as a function of the coin parameter θ , for the even and odd cases of Fig. 5. The dotted line indicates the upper bound $8/9$.

Similarly, the full history state (18) for the initially localized particle can also be regarded, when viewed from the spin, as a spin history state with respect to a *composite clock*, comprising both the original clock and the position degrees of freedom:

$$|\Psi\rangle = \frac{1}{\sqrt{NM}} \sum_{k,n} |kn\rangle \otimes U_k^n |\chi_0\rangle, \quad (72)$$

where $|kn\rangle = |k\rangle \otimes |n\rangle$ and for clarity we have altered the order in the tensor product. In this case the unitary




FIG. 7. Average over all initial spin states of the spin–rest quadratic entanglement entropy, $E_2(s, pT) = (2/3)E_2(\mathcal{W}_s)$ (Eq. (75)), as a function of the coin parameter θ , for $N = 20$. The dotted line indicates the upper bound $8/9$. Inset: Entanglement spectrum of the unitary operator \mathcal{W}_s , Eq. (73) generating the full spin history, as a function of the coin parameter θ .

operator generating the full spin history is

$$\mathcal{W}_s = \sum_{k,n} |kn\rangle\langle kn| \otimes U_k^n. \quad (73)$$

The average (over all initial spin states) $\langle E_2(s, pT) \rangle$ of the spin–rest entanglement in the state (72) is then determined by the entanglement of the operator (73), in turn determined by the full set of overlaps

$$\langle U_{k'}^{n'} | U_k^n \rangle = \frac{1}{2} \text{Tr} [U_{k'}^{-n'} U_k^n], \quad (74)$$

for $0 \leq k, k' \leq M$, $0 \leq n, n' \leq N - 1$.

Nonetheless, since all U_k^n are spanned just by $\{\mathbb{1}_s, \sigma_\theta, \sigma_y\}$ (Eq. (68)), the Schmidt rank of (73) is again $n_s = 3$ (or $n_s \leq 3$ in general), and the previous bound, i.e. $E_2(\mathcal{W}_s) \leq 4/3$ still holds. This entails again

$$\langle E_2(s, pT) \rangle = (2/3)E_2(\mathcal{W}_s) \leq 8/9. \quad (75)$$

This average is depicted as a function of θ in Fig. 7, together with the entanglement spectrum of \mathcal{W}_s .

V. CONCLUSIONS

We have analyzed the history state formalism in the context of discrete QW. The history state captures the whole evolution of the system, enabling for instance the evaluation of time averages as single quantum expectation values. It satisfies a timeless eigenvalue equation and can be generated through a quantum circuit from an initial system-clock product state. The formalism leads to the concept of system-time entanglement, which is here a measure of the number of orthogonal system states visited in the QW and is determined by the overlaps between the evolved states. Stationary system states are

then associated with *separable* history states while systems evolving into a new orthogonal state at each step correspond to maximally entangled history states.

We have focused on one dimensional QW with real Hadamard-type coins, where we demonstrated that such entanglement is strictly independent of the initial spin orientation for real initial states with definite parity. We also analyzed its connection with operator entanglement, showing that in the standard case of an initially localized particle it is that of the unitary operator generating the whole QW. Exact analytic results for overlaps and quadratic entropies in this case, including asymptotic expressions for large N , were as well derived, which show a monotonously decreasing entropy with increasing coin angle $\theta \in [0, \pi/2]$ (with $S_2(0) - S_2(\theta) \propto \tan \theta \frac{\ln N}{N}$ for the quadratic entropy). The associated history state entanglement spectrum shows, accordingly, a decreasing rank for increasing θ , with a deviation $O(\frac{\tan \theta}{N})^{\frac{1}{2}}$ from N^{-1} in the largest eigenvalue.

Finally, we have examined the QW from the spin perspective, showing that for an initially localized particle it is also described by a history state with a composite clock and a k -dependent unitary. The average over all initial spin states of the ensuing spin-rest entanglement can then be related to that of the global unitary generating this history. For the present Hadamard-type coin it has a limited rank with just three non-zero eigenvalues in its entanglement spectrum, leading to a bounded average spin-rest entanglement.

In summary, the history state formalism and the ensuing system-time entanglement provide a new perspective for analyzing QW. The extension of present results to more general QW and to open systems is currently under investigation.

ACKNOWLEDGMENTS

Authors acknowledge support from CONICET (F.L., N.C. and A.B.) and CIC (R.R.) of Argentina. Work supported by CONICET PIP Grant No. 11220200101877CO.

Appendix A: Quadratic entanglement entropy summation

The sum for the quadratic entanglement entropy (43b) can be done exactly for large n using the approximation (42) and neglecting terms $O(n^{-\frac{3}{2}})$. This yields

$$E_2(S, T) \approx 1 - [1 + \frac{\tan \theta}{\pi} F(N, \theta)]/N, \quad (A1)$$

$$\begin{aligned} F(N, \theta) &= 2 \sum_{n=1}^{\eta} (\frac{1}{n} - \frac{2}{N}) \cos^2(2n\theta + \pi/4) \quad (A2) \\ &= H(\eta) + \text{Im}[e^{4i\theta\eta} L(e^{4i\theta}, \eta)] + 2\theta - \frac{\pi}{2} \\ &\quad - \frac{2}{N} [\eta - \frac{\sin(2\theta(\eta+1)) \sin(2\theta\eta)}{\sin(2\theta)}], \quad (A3) \end{aligned}$$

where $\eta = \lfloor \frac{N-1}{2} \rfloor$, $H(\eta) = \sum_{n=1}^{\eta} 1/n$ is the harmonic number and $L(z, \eta) := \sum_{k=1}^{\infty} z^k / (k+\eta)$ is directly related

to the Lerch zeta function [59].

-
- [1] Y. Aharonov, L. Davidovich, and N. Zagury, “Quantum random walks,” *Phys. Rev. A* **48**, 1687 (1993).
- [2] R.P. Feynman, A.R. Hibbs, and D.F. Styer, *Quantum mechanics and path integrals* (Courier Corp., 2010).
- [3] J. Kempe, “Quantum random walks: An introductory overview,” *Contemp. Phys.* **44**, 307 (2003).
- [4] S. E. Venegas-Andraca, “Quantum walks: a comprehensive review,” *Quantum Inf. Process.* **11**, 1015 (2012).
- [5] A. Ambainis, “Quantum walks and their algorithmic applications,” *Int. J. Quantum Inform.* **1**, 507 (2003).
- [6] N. Shenvi, J. Kempe, and K. Birgitta Whaley, “Quantum random-walk search algorithm,” *Phys. Rev. A* **67**, 052307 (2003).
- [7] A. M. Childs and J. Goldstone, “Spatial search by quantum walk,” *Phys. Rev. A* **70**, 022314 (2004).
- [8] S. D. Berry and J. B. Wang, “Quantum-walk-based search and centrality,” *Phys. Rev. A* **82**, 042333 (2010).
- [9] L.K. Grover, “Quantum mechanics helps in searching for a needle in a haystack,” *Phys. Rev. Lett.* **79**, 325 (1997).
- [10] A.M. Childs, “Universal computation by quantum walk,” *Phys. Rev. Lett.* **102**, 180501 (2009).
- [11] N. B. Lovett *et al.*, “Universal quantum computation using the discrete-time quantum walk,” *Phys. Rev. A* **81**, 042330 (2010).
- [12] E. Sánchez-Burillo, J. Duch, J. Gómez-Gardeñes, and D. Zueco, “Quantum navigation and ranking in complex networks,” *Sci. Rep.* **2**, 605 (2012).
- [13] S. Lloyd, “Universal quantum simulators,” *Science* **273**, 1073 (1996).
- [14] D. W. Berry and A. M. Childs, “Black-box hamiltonian simulation and unitary implementation,” *Quantum Inf. Comput.* **12**, 29 (2012).
- [15] A. Schreiber *et al.*, “A 2d quantum walk simulation of two-particle dynamics,” *Science* **336**, 55 (2012).
- [16] X. Qiang *et al.*, “Efficient quantum walk on a quantum processor,” *Nat Commun* **7**, 11511 (2016).
- [17] G. Engel *et al.*, “Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems,” *Nature* **446**, 782 (2007).
- [18] P. Rebentrost *et al.*, “Environment-assisted quantum transport,” *New J. Phys.* **11**, 033003 (2009).
- [19] M. Karski *et al.*, “Quantum walk in position space with single optically trapped atoms,” *Science* **325**, 174 (2009).
- [20] P. Preiss *et al.*, “Strongly correlated quantum walks in optical lattices,” *Science* **347**, 1229 (2015).
- [21] S. Muga *et al.*, “Topological bound states of a quantum walk with cold atoms,” *Phys. Rev. A* **94**, 023631 (2016).
- [22] H. Schmitz *et al.*, “Quantum walk of a trapped ion in phase space,” *Phys. Rev. Lett.* **103**, 090504 (2009).
- [23] F. Zähringer *et al.*, “Realization of a quantum walk with one and two trapped ions,” *Phys. Rev. Lett.* **104**, 100503 (2010).
- [24] M. Gräfe *et al.*, “Integrated photonic quantum walks,” *J. Opt.* **18**, 103002 (2016).
- [25] F. Flamini, N. Spagnolo, and F. Sciarrino, “Photonic quantum information processing: a review,” *Rep. Prog. Phys.* **82**, 160001 (2019).
- [26] O. Carneiro *et al.*, “Entanglement in coined quantum walks on regular graphs,” *New. J. Phys.* **7**, 156 (2005).
- [27] S. E. Venegas-Andraca, J.L. Ball, K. Burnett, and S. Bose, “Quantum walks with entangled coins,” *New. J. Phys.* **7**, 221 (2005).
- [28] G. Abal, R. Siri, A. A. Romanelli, and R. Donangelo, “Quantum walk on the line: Entanglement and nonlocal initial conditions,” *Phys. Rev. A* **73**, 042302 (2006), Erratum: *ibid*, *Phys. Rev. A* **73**, 069905 (2006).
- [29] Y. Omar, N. Paunković, L. Sheridan, and S. Bose, “Quantum walk on the line with two entangled particles,” *Phys. Rev. A* **74**, 042304 (2006).
- [30] O. Maloyer and V. Kendon, “Decoherence versus entanglement in coined quantum walks,” *New. J. Phys.* **9**, 87 (2007).
- [31] A. Annabestani, M. R. Abolhasani, and G. Abal, “Asymptotic entanglement in 2d quantum walks,” *J. Phys. A* **43**, 075301 (2010).
- [32] A. Romanelli, “Distribution of chirality in the quantum walk: Markov process and entanglement,” *Phys. Rev. A* **81**, 062349 (2010).
- [33] Y. Ide, N. Konno, and T. Machida, “Entanglement for discrete-time quantum walks on the line,” *Quant. Inf. Comp.* **11**, 761 (2011).
- [34] B. Allés, S. Gündüç, and Y. Gündüç, “Maximal entanglement from quantum random walks,” *Quant. Inf. Process.* **11**, 211227 (2012).
- [35] R. Vieira, E.P.M. Amorim, and G. Rigolin, “Entangling power of disordered quantum walks,” *Phys. Rev. A* **89**, 042307 (2014), *Phys. Rev. Lett.* **111** 180503 (2013).
- [36] A. C. Orthey and E.P.M. Amorim, “Asymptotic entanglement in quantum walks from delocalized initial states,” *Quantum Inf. Process.* **16**, 224 (2017).
- [37] D.N. Page and W.K. Wootters, “Evolution without evolution: Dynamics described by stationary observables,” *Phys. Rev. D* **27**, 2885 (1983), W.K. Wootters, “Time replaced by quantum correlations”, *Int. J. Th. Phys.* **23**, 701 (1984).
- [38] C. Rovelli, “Quantum mechanics without time: a model,” *Phys. Rev. D* **42**, 2638 (1990).
- [39] C. J. Isham, “Quantum logic and the histories approach to quantum theory,” *J. Math. Phys.* **35**, 2157 (1994).
- [40] V. Giovannetti, S. Lloyd, and Maccone L., “Quantum time,” *Phys. Rev. D* **92**, 045033 (2015).
- [41] E. Moreva, G. Brida, M. Gramegna, V. Giovannetti, L. Maccone, and M. Genovese, “Time from quantum entanglement: An experimental illustration,” *Phys. Rev. A* **89**, 052122 (2014).
- [42] J.R. McClean and A. Aspuru-Guzik, “Clock quantum monte carlo technique: An imaginary-time method for real-time quantum dynamics,” *Phys. Rev. A* **91**, 012311 (2015).
- [43] A. Boette, R. Rossignoli, N. Gigena, and M. Cerezo, “System-time entanglement in a discrete-time model,” *Phys. Rev. A* **93**, 062127 (2016).

- [44] A. Boette and R. Rossignoli, “History states of systems and operators,” *Phys. Rev. A* **98**, 032108 (2018).
- [45] B. DeWitt, “Quantum theory of gravity. i. the canonical theory,” *Physical Review* **160**, 1113 (1967).
- [46] E.O. Dias and F. Parisio, “Space-time-symmetric extension of nonrelativistic quantum mechanics,” *Phys. Rev. A* **95**, 032133 (2017).
- [47] A. Nikolova *et al.*, “Relational time in anyonic systems,” *Phys. Rev. A* **97**, 030101(R) (2018).
- [48] P.J. Coles, V. Katariya, S. Lloyd, I. Marvian, and M.M. Wilde, “Entropic energy-time uncertainty relation,” *Phys. Rev. Lett.* **122**, 100401 (2019).
- [49] D. Pabón *et al.*, “Parallel-in-time optical simulation of history states,” *Phys. Rev. A* **99**, 062333 (2019).
- [50] R. Gambini, R. A. Porto, J. Pullin, and S. Tortorolo, “Conditional probabilities with dirac observables and the problem of time in quantum gravity,” *Phys. Rev. D* **79**, 041501(R) (2009).
- [51] Z. Zhao *et al.*, “Geometry of quantum correlations in space-time,” *Phys. Rev. A* **98**, 052312 (2018).
- [52] N. L. Diaz and R. Rossignoli, “History state formalism for Dirac’s theory,” *Phys. Rev. D* **99**, 045008 (2019).
- [53] N. L. Diaz, J. M. Matera, and R. Rossignoli, “History state formalism for scalar particles,” *Phys. Rev. D* **100**, 125020 (2019), “Spacetime quantum actions”, *Phys. Rev. D* **103**, 065011 (2021).
- [54] A. Ambainis, E. Bach, A. Nayak, A. Vishwanath, and J. Watrous, “One-dimensional quantum walks,” in *Proceedings of the Thirty-third Annual ACM Symposium on Theory of Computing*, 37 (2001).
- [55] N. Konno, “Quantum random walks in one dimension,” *Quantum Inf. Proc.* **1**, 345 (2002).
- [56] B. Tregenna, W. Flanagan, R. Maile, and V. Kendon, “Controlling discrete quantum walks: coins and initial states,” *New J. Phys.* **5**, 83 (2003).
- [57] C. Tsallis, *Introduction to Non-Extensive Statistical Mechanics* (Springer-Verlag, N.Y., 2009).
- [58] C. Beck and F. Schlögl, *Thermodynamic of Chaotic Systems* (Cambridge Univ. Press, UK, 1993).
- [59] M. Abramowitz and I.A. Stegun, eds., *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, Applied Mathematics Series*, Vol. 55 (Dover Publications, 1965) F.W.J. Olver, D.M. Lozier, R.F. Boisvert, C.W. Clark eds, *NIST Handbook of Mathematical Functions*, Cambridge Univ. Press (2010).