

# Emerging SYK physics and Wigner-Dyson distribution in polaron system

Chen-Huan Wu \*

College of Physics and Electronic Engineering, Northwest Normal University, Lanzhou 730070, China

October 5, 2022

We study the of SYK interactions and Wigner-Dyson distribution in the polaron system through the theoretical study and numerical simulation. The polaron as a long-lived quasiparticle which can be found in the incompressible state has slow momenta and current relaxation in Fermi liquid phase. We reveal the relation between UV cutoff of polaronic momentum  $\Lambda_q$  and its SYK behavior. The SYK behavior of a polaron system, as well as the relation between scattering momentum and the related statistical behaviors has rarely been investigated before. We found that the inversed momentum cutoff  $\Lambda_q^{-1}$ , which plays the role of an essential degree-of-freedom (DOF) other than the fermions, relates to the distribution and statistical variance of polaronic coupling term. By projecting to a 2d square lattice, we consider this problem in position space where the DOF of polaron scattering momenta is replaced by another flavor (denoted as  $\eta_\Delta$  with flavor number of order of  $O(M)$ ) which is determined by the site potential difference  $\Delta$  as well as the site index, and we also applying the self-attention method to searching for the more efficient route to exploiting the many-body behaviors. The algorithm designed by us allow the automatical optimization and prediction for the resulting spectrum to arbitrary accuracy. Although here  $\eta_\Delta$  (or  $\Lambda_q$  in terms of the momentum space representation) should be a Gaussian variable with zero mean by its own (noninteracting), but it becomes a Chi-square variable when it couples with the fermionic DOF ( $O(N)$ ). The resulting system follows the statistic of between Gaussian symplectic ensemble (GSE) and Gaussian unitary ensemble (GUE) as long as  $O(M) \sim O(N^2/4)$ . While in terms of the momentum space representation, different magnitude of  $\Lambda_q^{-1}$  support to different phases, including non-Fermi liquid phase and (disordered) Fermi liquid phase, which correspond to ill-defined and well-defined polarons, respectively, and the suppression to the Gaussian distribution in non-fermi liquid phase by the pairing condensation-induced local coupling of is also being mentioned.

## 1 Introduction

We consider a two-dimensional  $N(N - 1)$  square lattice consist of the fermion index  $i$  and  $j$  ( $i, j = 1, \dots, N$ ). As we know that to realizing the polaron physics, it requires localization on the momentum space of interacting fermions which leads to vanishingly small scattering momenta  $q \rightarrow 0$ . Then the polarons which play the role of quasiparticles here can be considered in quasi-one-dimensional system in such a 2d lattice by virtue of the chiral character, and we only consider the polarons contributed by the right-direction effects (a combination effect of

---

\*chenhuanwu1@gmail.com

fermion indices difference and on-site potential difference) on each quasi-one-dimensional chain. By considering the vector potential, the random motion of fermions in opposite directions will cancel each other and leads to zero expectation of polaron feature in each quasi-one-dimensional chains. However, by adding another degree-of-freedom (DOF) which is the randomly distributed on-site potential, the nonzero expectation of polaron feature can be generated. All the on-site potential differences in horizontal direction in this 2d system can be regarded as a Gaussian variable which satisfy  $\overline{\Delta_{\alpha\beta}} = 0$ , and its variance is close to (but lower than) the typical value  $\overline{\Delta_{\alpha\beta}^2} = N^2/4$ , where we denote the potential difference on each horizontal lines as  $\Delta_{\alpha\beta}$  ( $\alpha, \beta = 1, \dots, N$ ). Although  $\Delta_{\alpha\beta}$  is a Gaussian variable by its own, it changes to a Chi-square variable (product of two Gaussian ones) with nonzero mean when it couple with the fermionic DOF. We consider that, for each part of fermion indices ( $i, j$ ), the probabilities for them to occupy each part of  $(\alpha, \beta)$  are dominated by the rule that the probabilities for each distinct value of  $\eta_{\Delta} = |\Delta_{\alpha\beta}|\alpha$  must be the same, which means only the quantity  $|\Delta_{\alpha\beta}|\alpha$  (on-site potential difference times the site-index on the left, due to the chiral character of quasi-1d system and vector potential) can be regarded as an independent disorder or DOF relative to fermion DOF, i.e., each part of  $(i, j)$ , instead of the  $\Delta_{\alpha\beta}$ . And the key condition for the realization of polaron behavior the nonzero net potential difference  $\Delta_{\alpha,\beta}$ , as well as the Wigner-Dyson distribution which should be beyond the GUE region (with character level spacing ratio  $\langle r \rangle_{GUE} = 0.59 \sim 0.60$ ), which implies the thermalization and Gaussian randomness on the on-site potential difference DOF. The Hamiltonian describing the effect of polaron reads

$$H = \sum_{ij;\alpha,\beta} = w_{ij}[\Delta_{\alpha\beta}]c_i^{\dagger}c_j, \quad (1)$$

where the potential difference  $\Delta_{\alpha\beta}$  is the weighted Gaussian variable, and it couples with the fermionic DOF represented by the fermionic indices ( $i, j$ ). The weighting function  $w_{ij}$  depends on the fermion DOF by the requirement that the probability  $p_{\eta_{\Delta}}$  must be a constant during the selection by the fermion indices. We summarize our main result [here](#). As long as the system is under this condition, we found that the variance satisfies  $\sigma_q^2 = \overline{\Delta_{\alpha\beta}^2} \sim N^2/4$  and the level spacing ratio  $\langle r \rangle$  will be of the region between GUE and GSE. The value of this variance is also the size of  $M$  (i.e., flavor number of distinct values of  $|\Delta_{\alpha\beta}|\alpha$ ). We also provide another version of the this model in terms of the momentum space representation in Appendix.

## 2 Gaussian-based self-attention

Projection to this 2d lattice model available us to using the transformer method base on (multi-head) self-attention model to solve this problem. Firstly, we consider the case in the absence of Gaussian weighting term, i.e., considering the unbiased potential difference DOF  $\Delta_{\alpha,\beta}$  ( $\alpha, \beta = 1, \dots, N$ ), the self-attention layer reads

$$\mathbf{H} = \text{Attn}(\mathbf{Q}, \mathbf{K}, \mathbf{V}) = \text{Softmax}\left[\frac{\mathbf{Q}\mathbf{K}^T}{\sqrt{\sigma_q^2}}\right]\mathbf{V}, \quad (2)$$

where  $\mathbf{Q} = \mathbf{X}\mathbf{W}^{\mathbf{Q}}$ ,  $\mathbf{K} = \mathbf{X}\mathbf{W}^{\mathbf{K}}$ ,  $\mathbf{V} = \mathbf{X}\mathbf{W}^{\mathbf{V}}$ , are the query, key, and value matrices, which are all  $N \times (N - 1)$  type.  $\mathbf{X}$  is the input  $(\alpha, \beta)$  sequence. It is direct to know that when the potential difference DOF is unweighted (or selected) by the fermionic DOF, it behaves as a Gaussian variable, and the probabilities for  $\Delta_{ij} = 1, \dots, (N - 1)(i \neq j)$  are unequal, as shown in the simulator-obtained spectrum in Fig.1(a,b), lower value of  $\Delta_{ij}$  has bigger weight. In this case the net potential difference is zero and the variance is  $\overline{\Delta_{ij}^2} \sim N^2/8$ , and the level spacing distribution follows the GUE with  $\langle r \rangle = 0.59 \sim 0.60$ .

## 2.1 Gaussian-based self-attention

To see the result when the  $M$ -DOF couples with the fermionic DOF, we firstly simulate the real case according to the calculated exact probabilities for each part of  $(\alpha, \beta)$  according to the selection rule of fermions,  $\partial_{\alpha, \beta} \eta_{\Delta} = 0$ , i.e., each value of  $\eta_{\Delta}$  has the same weight in the final  $\Delta_{\alpha, \beta}$ -spectrum after the selection of fermionic  $(i, j)$  parts. The calculated probabilities for each  $(\alpha, \beta)$ -state (i.e., the weight of each part of  $(\alpha, \beta)$ ) arranged in order of value of  $\eta_{\Delta}$  are shown in Fig.2(c). As we can see, now the probability distribution of potential difference changes and results in nonzero net value of potential difference. The resulting  $\Delta_{\alpha, \beta}$ -spectrum is shown in Fig.3(e,f), and the corresponding level spacing ratio follows a distribution between GUE and GSE.

However, it is low efficiency to calculating one-by-one the probability of each part of  $(\alpha, \beta)$  especially for large value of  $N$ . Next we introduce a Gaussian-weight-based method. Since there will be  $\sim N^2/4$  distinct values of  $\eta_{\Delta}$ , which consist the group  $\{\eta_{\Delta}\}$ . By extracting the mean value and variance of  $\eta_{\Delta}$  from the presence  $\Delta_{\alpha\beta}$ -spectrum, where we denote as  $\mathbf{M} = [\mathbf{QK}^T]_{\eta_{\Delta}}$ , and comparing to those quantities of the group  $\{\eta_{\Delta}\}$  (where we denote as  $m_{\eta}$  and  $\sigma_{\eta}^2$ , respectively), the actual probabilities (weights) for each  $(\alpha, \beta)$ -state can be predicted to high accuracy.

Macroscopically, there will always be more repeating  $\eta_{\Delta}$  in the low-value region as can be seen from the  $\eta_{\Delta}$ -spectrum, thus the mean value and variance of unbiased  $\eta_{\Delta}$ -spectrum will be always lower than the selected one ( $\{\eta_{\Delta}\}$ ). Thus we, in first step, define the mean value  $m_0$  as the maximal value of group  $\eta_{\Delta}$ , i.e.,  $m_0 = N(N-1)$ , and making the inner product

$$\mathbf{M} \cdot \mathbf{G}, \quad (3)$$

where  $[\mathbf{QK}^T]_{\eta_{\Delta}}$  is the  $N(N-1)$  extracted  $\eta_{\Delta}$  arranging according to its value. Next,

$$G_{i \in [1, N(N-1) - |\delta_i|]} = \exp\left[\frac{-\left(\{M\}_i - m_{\text{Max}[i] + \lambda_i}\right)^2}{2\sigma_M^2}\right], \quad (4)$$

$$G_{i \in [(N(N-1) - |\delta_i| + 1, N(N-1))]} = 1,$$

where  $|\delta_i| = 0$  in the beginning and the mean value  $m_{\text{Max}[i] + \lambda_i}$  is the  $\text{Max}[i]$ -th element of the ascending-ordered  $\{M\}_i$  group. We denote the index as  $i = 1, \dots, N(N-1)$  representing the values of  $\eta_{\Delta}$  in ascending order.  $\sigma_M^2 = \frac{1}{M'} \sum_i^{M'} (M_i - m_0)^2$  is variance of elements in the group  $\{M\}_i$  ( $i = 1, \dots, M'$ ) with  $M' \leq M$ .  $\delta_i \in [0, N(N-1))$  is learnable parameter (its variance step is defined according to the feed-back of mean value and variance of the weighted spectrum after the process of last time) which equals to zero in the beginning, and its function here is to restricting the weighted-range to lower- $i$  regions, where has more number of repeating value of  $\eta_{\Delta}$ . While other elements of  $\mathbf{G}$  are setted as 1. Here the mean value  $m_i$  is defined according to the  $(N(N-1) - \delta_i)$ -th value of  $\mathbf{M}$ , and  $\lambda_i$  is a small quantity which is also a learnable parameter. Its sign  $\text{sgn}[\lambda_i]$  is negative when the ratio between mean value (variance) obtained from last iteration (we denote as the  $l$ -th learning) and that of the irreducible group  $\{\eta_{\Delta}\}$  satisfies  $m_l/m_{\eta} \ll \sigma_l^2/\sigma_{\eta}^2$ , and its sign should be positive in the opposite case that  $m_l/m_{\eta} \gg \sigma_l^2/\sigma_{\eta}^2$ . After each time of iteration (learning), the renewed spectrum will be obtained from the matrix  $\mathbf{M}$  weighted by the rescaled  $\prod_{j=1}^l G_j$  (by softmax process).

The iteration process of learning will stop when the  $m_l$  and  $\sigma_l^2$  are close enough to the standard values  $m_{\eta}$  and  $\sigma_{\eta}^2$ , respectively. There are two ways to verify the final results, i.e., the obtained probabilities for each  $(\alpha, \beta)$ -state. First one is comparing the  $\eta_{\Delta}$ -spectrum and the level spacing distribution with the exactly calculated one, the second one is to measuring the distance between the predicted probabilities with the exactly calculated ones (denote as  $p_i^{\eta}$ ). Instead of using the loss function  $L = -\sum_i p_i^{\eta} \ln p_i$ , we choose to calculate the mean

value and variance of the group consist of the elements  $p_i^{p_i^\eta}$ , where both the predict and exact probabilities are arranged in ascending order of  $\eta_\Delta$ . While for the exact case, where all the predicted probabilities are the same with the real one, the mean value and variance of the group  $\{(p_i^\eta)^{p_i^\eta}\}$  are solely determined by the sample size  $N$ , where the mean value ( $\sim 2(e^{1/M} - 1)$ ) exponentially closes to zero and the standard deviation ( $\sim 1 - 2(e^{1/M} - 1)$ ) logarithmically closes to one with the increase of  $N$ .

### 3 Simulations

According to our simulation, the variance of  $\eta_\Delta$  spectrum  $\sigma_\eta^2$  is lower but close to  $\frac{N^4}{16}$  while that of the  $\Delta_{\alpha\beta}$ -spectrum is  $\sigma_\Delta^2 = \frac{N^2}{4}$  which is quite close to the flavor number of  $\eta_\Delta$ -DOF, i.e.,  $M \sim \frac{N^2}{4}$ . Since the  $\eta_\Delta$ -DOF is a independent DOF relative to every  $(i, j)$ -part, this model is a three-DOF (or three-point all-to-all many-body system) system. As shown in Appendix.B, this can be illustrated in the fermion system by adding another scattering momentum-dependent DOF into the four-point fermion-fermion interacting system, and changes the variance of antisymmetry tensor from the four-point one ( $\sim \frac{1}{4!} = \frac{g_4^2}{4N^3}$ ) to the three-point one ( $\sim \frac{1}{3!} = \frac{g_3^2}{3N^2}$  where  $g_n = \sqrt{\frac{n^{n-1}}{(n-1)!}}$  is a constant parameter with dimensions of energy). For example, in Eq.(137) the four Gaussian-independent fermionic indices with  $i < j < k < l$  reduce to two (coarse-grained) Gaussian variables  $(i, k)$  and  $j, l$  (with  $i < j$  and  $k < l$ ) which correspond to the fermions before and after scattering, respectively, when it couples to the  $\eta_\Delta$ -DOF. The ratio between variance of  $\Delta_{\alpha\beta}$ -spectrum and  $\eta_\Delta$  spectrum is close to  $\frac{\sigma_\Delta^2}{\sigma_\eta^2} \sim \frac{4}{N^2}$ . This is because for each fermion part, the probability for  $M$  flavors of  $\eta_\Delta$ -DOF is  $p_\eta = \frac{1}{M} \sim \frac{4}{N^2}$ , while the unequivocal distribution of the probabilities (see Fig.3(a)) for each  $\Delta_{\alpha\beta}$  (whose mean  $\bar{p}_\Delta = \frac{1}{2N(N-1)}$ ) results in larger uncertainty, and this uncertainty results in stronger thermalization effect (compares to the original one) and smaller variance for the  $\Delta_{\alpha\beta}$ -spectrum which is  $\leq \frac{N^2}{4}$ ). In this case the potential difference is a Chi-square variable and has nonzero expectation value. This part of nonzero potential difference corresponds to the UV cutoff in position space as well as the long-wavelength limit for the scattering momenta, which is a signature for the polarons following the Wigner-Dyson distribution. And the resulting distribution is of a intermediate region between the GUE and GSE. While in original  $\Delta_{\alpha\beta}$ -spectrum as shown in Fig.1(a,b), the potential difference is a Gaussian variable with zero mean (and thus zero expectation value for the polaronic effect) and the many-body behavior follows the GUE distribution. Also, this case belongs to the four-point-type fermion interaction, with variance  $\sigma_{4-p}^2 \sim \frac{4N^2}{4!} = \frac{N^2}{6}$ , and its level spacing follows GUE distribution ( $\langle r \rangle \sim 0.59$ ). While the two-point-type fermion interaction (with much stronger thermalization), as shown in Fig.1(c,d), it has a variance which is twice of the four-point one,  $\sigma_{2-p}^2 = \frac{N^2}{2!} \frac{2}{3} = \frac{N^2}{3}$ , and its level spacing follows GSE distribution ( $\langle r \rangle \sim 0.67$ ). Then it is reasonable for the intermediate case we studied here, whose variance  $\sigma_\Delta^2 \sim \frac{N^2}{4}$  is between the two-point one and four-point one, and follows the ensemble between GUE and GSE with  $\langle r \rangle = 0.63 \sim 0.65$ . More critically, the variance  $\sigma_\Delta^2$  is in fact  $N$ -dependent even for  $N \rightarrow \infty$ . If we treating it as an inner product by four-point variance and two-point variance weighted by different probabilities, an experimental expression could be

$$\sigma_\Delta^2 = (1 - \frac{1}{\ln N^2}[\sigma_{2-p}^2, \sigma_{4-p}^2] \cdot \text{Softmax}[1, 2]) \approx (1 - \frac{1}{\ln N^2}(0.2689\sigma_{2-p}^2 + 0.731\sigma_{4-p}^2)). \quad (5)$$

For example, for  $N = 100, 150, 200$ , the corresponding variances  $\{\sigma_{2-p}^2, \sigma_\Delta^2, \sigma_{4-p}^2\}$  are  $\{3333.3, 1954, 1666.5\}$ ,  $\{7500, 4355, 3750\}$ , and  $\{13421, 7674.1, 6666, 6\}$ , respectively. There is a tendency which is persistent for arbitrary large- $N$ : the variance of such intermediate state depends on both the other

two variances,  $\sigma_{2-p}^2$  and  $\sigma_{4-p}^2$  whose values can be solely determined in large- $N$  limit, and with  $N$  increasing, the  $\sigma_{\Delta}^2$  will more and more close to the four-point one  $\sigma_{4-p}^2$  (compared to  $\sigma_{2-p}^2$ ) with a decreasing but never vanished velocity. This indeed reflects a spontaneous localization, and the suppressed thermalization with the increasing  $N$ .

## 4 Mathematical principle behind the learning-algorithm designing

### 4.1 Algorithm

During the steps of automatic iterations, the criterion is to matching the mean value and variance of the predicted spectrum with the reduced one (irreducible group, which has much small sample size). We using the rule that for every weighted group that containing only two elements, its variance will never be changed when we modify there corresponding weights, even if the summation of there corresponding weights not equals to one, by the mean value will changes in this process.

For example, for a group  $k_1, k_2$  weighted by  $w_1, w_2$ , the variance reads

$$\sigma^2 = \frac{\sum_i w_i (k_i - m_k)^2}{\sum_i w_i - \frac{\sum_i w_i^2}{\sum_i w_i}}, \quad (6)$$

where  $m_k = \sum_i k_i w_i / \sum_i w_i$  is the weighted mean value. This expression is equivalent to replacing  $w_2$  by  $(1 - w_1)$  and omit the  $\sum_i w_i$  in the denominator, i.e.,  $\sigma^2 = \frac{\sum_i w_i (k_i - m_k)^2}{1 - \sum_i w_i^2}$ . Thus next we consider only the case  $\sum_i w_i = 1$ . Then we introduce another one with modified weight distribution  $\{w'_1, w'_2\} := \{w_1 + w_d, w_2 - w_d\}$ . These two groups have the same variance

$$\sigma^2 = \frac{1}{2}(k_1 - k_2)^2 = \frac{(k_1 - k_2)^2 \prod_i w_i}{1 - \sum_i w_i^2} = \frac{(k_1 - k_2)^2 (\prod_i w_i - w_d^2 + w_d(1 - 2w_1))}{1 - \sum_i w_i'^2}. \quad (7)$$

This can be explained using the method we discussed in Ref.[1], i.e., expressing this common variance  $\sigma^2$  in terms of a limiting result of the infinitely scaled variable  $\chi$  (the detailed form of this variable does not have to know), which has the form  $\sigma^2 := \lim_{\chi \rightarrow \infty} f = \frac{f}{1 - \partial_{\chi} f}$ . In above expression the summation of squared weights of these two groups can be treated as there corresponding derivative terms with respect to  $\chi$ ,

$$\sum_i w_i^2 = 1 - 2w_1(1 - w_1), \quad \sum_i w_i'^2 = 2(-w_d^2 + (1 - w_1)w_1 + w_d(1 - 2w_1)). \quad (8)$$

For first group, by defining  $\prod_i w_i = w_1(1 - w_1) := f_{\chi}$ , we have

$$\begin{aligned} \frac{1}{2} &= \lim_{\chi \rightarrow \infty} f_{\chi} = \frac{f_{\chi}}{1 - \partial_{\chi} f_{\chi}}, \\ 1 - 2f_{\chi} &= \partial_{\chi} f_{\chi}, \\ \frac{1}{2}(k_1 - k_2)^2 &= \lim_{\chi \rightarrow \infty} (k_1 - k_2)^2 f_{\chi}, \\ 1 - 2f_{\chi} &= \partial_{\chi} [(k_1 - k_2)^2 f_{\chi}]. \end{aligned} \quad (9)$$

There are two possible cases which correspond to different representations. The first one is the representation in unit of  $(k_1 - k_2)^2$  which is independent with  $\chi$ . This case will leads to  $w_i = \frac{1}{2} \pm \frac{i}{2}$ . The second one is the representation in unit of  $\lim_{\chi \rightarrow \infty} f_{\chi} = \frac{1}{2}$ . In this representation, the acceptable results,  $w_i = \frac{1}{2}$  can be solved. Note that for this case, the derivative term  $\partial_{\chi} f_{\chi}$  in the second line of above expression indeed corresponds to  $\partial_{\chi} w_1 =$

$(1 - w_1)$  and  $\partial_\chi w_2 = \partial_\chi(1 - w_1) = w_1$ , which mean the variable  $\chi$  is scalable by 1 (which is no more a constant here), and  $\partial_\chi 1 = 1$ . Here  $1 = e^{i(2\pi - 0^+)}$ , where  $0^+$  denotes an infinitely small shiftment in phase and generate the imaginary part into the previous representation, and the cutoff of this infinitely small imaginary part (which is the UV cutoff in momentum space if we projecting this imaginary part into the position space) determines directly the largest quantity that could be independent with  $\chi$ , i.e., the unit quantity in this representation ( $1/2$ ). Thus more rigorously we have  $\partial_\chi f_\chi = 1^2 - 2f_\chi$  with  $1^2 \neq 1$  and  $\partial_\chi 1 \neq \partial_\chi 1^2$ . Here this cutoff is necessary to fix the tolerance of whole system, i.e.,  $\partial_\chi \frac{1}{2} = \left| \frac{1}{2} - \frac{1}{2 + \frac{1-1^2}{f_\chi}} \right| = 0$ . Substituting the second line of above expressions into the expanded expression in the last line, we have

$$\partial_\chi(k_1 - k_2)^2 = \frac{1 - 2f_\chi}{f_\chi}[1 - (k_1 - k_2)^2] = \frac{\partial_k f_\chi}{f_\chi}[1 - (k_1 - k_2)^2]. \quad (10)$$

In the mean time, due to the invariant property of the infinite scale,  $\partial_\chi \lim_{\chi \rightarrow \infty} f_\chi = 0$ , the first line of Eq.(9) results in the relation

$$\frac{1 - 2f_\chi}{2f_\chi} = -f_\chi \partial_\chi \frac{1}{2f_\chi}. \quad (11)$$

Substituting this into Eq.(10), we can obtain

$$\frac{1 - (k_1 - k_2)^2}{\partial_\chi(k_1 - k_2)^2} = \frac{1}{2} \frac{-1}{f_\chi \partial_\chi \frac{1}{2f_\chi}} = \frac{f_\chi}{\partial_\chi f_\chi}. \quad (12)$$

Thus we can make sure that there must be another  $\prod_i w_i$ -related variable  $\chi'$  satisfying

$$\begin{aligned} \lim_{\chi' \rightarrow \infty} f_{\chi'} &= \frac{\partial f_\chi}{\partial(k_1 - k_2)^2}, \\ \partial_{\chi'} f_{\chi'} &= (k_1 - k_2)^2. \end{aligned} \quad (13)$$

Using the property  $\frac{f_\chi}{\partial_\chi f_\chi} \equiv \lim_{\chi \rightarrow \infty} f_\chi \frac{f_\chi}{\lim_{\chi \rightarrow \infty} f_\chi - f_\chi}$ , we have

$$\begin{aligned} f_\chi &= \frac{-\partial_\chi f_\chi}{1 - \partial_\chi f_\chi} \frac{1}{\partial_\chi \frac{1}{2f_\chi}}, \\ &= \frac{\partial f_\chi}{\partial(k_1 - k_2)^2} [1 - (k_1 - k_2)^2]. \end{aligned} \quad (14)$$

Combined with Eq.(10), we obtain the following form of the limiting scale

$$\lim_{\chi \rightarrow \infty} f_\chi = \frac{1}{2} = f_\chi - f_\chi^2 \partial_\chi \frac{1}{2f_\chi}. \quad (15)$$

Besides, using Eq.(12) and the derivative of the ratio between two limits  $\frac{\lim_{\chi \rightarrow \infty} [(k_1 - k_2)^2 f_\chi]}{\lim_{\chi \rightarrow \infty} f_\chi} = (k_1 - k_2)^2$ , we obtain

$$\begin{aligned} \partial_\chi \ln \frac{1}{\lim_{\chi \rightarrow \infty} f_\chi} &= \partial_\chi \ln 2 \\ &= \frac{\partial_\chi f_\chi}{f_\chi} \frac{1 - (k_1 - k_2)^2}{(k_1 - k_2)^2} \\ &= \frac{\partial_\chi (k_1 - k_2)^2}{(k_1 - k_2)^2}. \end{aligned} \quad (16)$$

Thus we have,  $\frac{\partial_\chi 2}{2} = \frac{\partial_\chi (k_1 - k_2)^2}{(k_1 - k_2)^2}$ , thus in the limit of  $\partial_\chi 2 \rightarrow 2$ ,  $(k_1 - k_2)^2 \rightarrow e^{i(2\pi - 0^+)} = e^\chi$  and  $\frac{\partial_\chi (k_1 - k_2)^2}{(k_1 - k_2)^2 - 1} \rightarrow \frac{\partial_\chi (k_1 - k_2)^2}{(k_1 - k_2)^2}$ . To looking for real solution, in this limit we have  $(k_1 - k_2)^2 = e^\chi \sim 1$ , i.e.,  $\lim_{\chi \rightarrow \infty} |k_1 - k_2| \gg |k_1 - k_2|$ . This fact reveals the reason why the variance of such binary group will always keep invariant when we modify the weight distribution or change the value of elements, as long as the distance between two elements is invariant. This is consistent with the fact that  $|k_1 - k_2| \propto \delta\chi$  as they all describe the distance in the position space, while the corresponding long wavelength limit in momentum space after fourier transformation can be realized by the factor  $[1 - (k_1 - k_2)^2]^{-1}$ . Also, the unit quantity expressed in terms of the limiting relation as shown in eq.(15) allows the easier derivation along the distances in the position space. For example, if we consider beyond the binary groups by replacing  $f_\chi = w_1(1 - w_i)$  as the variance for the binary group in its position, then the  $[\partial_\chi \frac{1}{2f_\chi}]^{-1}$ -term in Eq.(15) can indeed be regarded as the variance of another group (with more elements) in position next to the former one. To illustrating this, we set the binary group in the first position as  $(k_1, k_N)$  whose variance denoted as  $v_1$ , and the one in the next position as  $(k_1, k_1 + 1, k_N - 1, k_N)$  whose variance denoted as  $v_2$ . The other groups with more elements are arranged in the same manner. Then the slope in the position space is equivalent to the variance ratio  $\frac{v_1 - v_2}{v_2 - \frac{v_2^2}{v_1}} = \frac{v_1}{v_2}$ , where  $(v_2 - \frac{v_2^2}{v_1})$  is the smallest step length here appearing in the same form with Eq.(15), i.e., the IR cutoff in the position space. Such unit length as well as the slope are not invariant between arbitrarily two positions with different numbers of element, and this length reduces to zero efficiently for higher-order position derivative. For example, by setting  $k_N = 13$ , the first to fourth-order derivative are  $-6.4918$ ,  $0.84579$ ,  $0.040513$ ,  $0.0093128$ , respectively, and the highest-order one (the twelfth-order derivative) is  $0.000030169$ .

It can be checked that, in this representation, where both the elements and weights are dependent with  $\chi$ , the real solution of weights  $w_i = \frac{1}{2} = \partial_\chi f_\chi = (k_1 - k_2)^2 [1 - \frac{1}{2 \lim_{\chi \rightarrow \infty} (k_1 - k_2)^2}] (f_\chi = \frac{1}{4})$  can be obtained, where  $\partial_\chi \frac{1}{2f_\chi} = -4$ . In the mean time the infinite  $\chi$  corresponds to the simultaneous scalings of  $w_i \rightarrow \frac{1}{2}$  and  $w_d^2 \rightarrow 0$  for the second group, which results in the same result of variance. Thus for such kind of binary groups, we can only modify the mean value and keep the variance be invariant. To modify the variance, the simplest way is by introducing the third element, which can be viewed as a collection of all the other elements in the reducible group  $\{\alpha, \beta\}$  other than the two elements in both sides of the position that the algorithm kernel working on. Once a position is initially determined, the elements in two sides of this position will be treated as a part of binary (summation of weights is one), then the mean value can be increased by giving more weight to the right-side one, which corresponds to larger value of  $\eta_\Delta$ . Then other elements enter this group play the role of third element to available the adjustment for the variance. In Eq.(4), the learnable mean value  $m_{i+\lambda_i}$  at position  $i = i'$  can be adjusted with the range of  $[\{M\}_{i'}, \{M\}_{N(N-1)}]$ . As the subscript of this learnable mean parameter decreasing from the largest one  $N(N-1)$  (corresponds to the position at the end of spectrum), the resulting mean value and variance of the updated  $\eta_\Delta$ -spectrum will both be increased at first and then keep decreasing until the subscript decreases to  $i = i'$ . During this process, there will be a certain position will the mean value is enhanced while the variance is reduced (compares to the former spectrum before the update of parameters).

In this way, the learnable-parameter-dependent step lengths control the positions where the algorithm modifies the weight distribution step-by-step, and the ratio between mean value and variance can be modified until the spectrum exhibits features match enough to the one corresponding to the irreducible group.

## 4.2 Criterion

Except the above-introduced method to estimate the degree of coincidence between the predicted mean value and variance with the exact ones, there is another criterion using the  $\beta_{en}$ -factor ( $\beta_{en} = 1, 2, 4$  for GOE, GUE, and GSE ensembles). As the joint probability density function for Gaussian ensembles[39] satisfies

$$P_{joint} = C_{\beta,N} \prod_{i<j} |k_i - k_j|^{\beta_{en}} e^{-\frac{1}{2} \sum_i^N (k_i - m_k)}, \quad (17)$$

where the normalization parameter reads

$$C_{\beta,N} = \frac{1}{(2\pi)^{N/2}} \prod_j^N \frac{\Gamma[1 + \frac{\beta_{en}}{2}]}{\Gamma[1 + j \frac{\beta_{en}}{2}]}. \quad (18)$$

We note in advance that, as can be seen from Eq.(6) in the above subsection, we are using the bias-corrected sample variance throughout this paper. Then according to the above-introduced algorithm, we start with an ascending-ordered original  $\eta_\Delta$ -group (unreducible)  $\{M\}_{i=1, \dots, N(N-1)} := \{k_1, \dots, k_{N(N-1)}\}$  following GUE distribution (whose mean value and variance are denoted as  $m_o$  and  $\sigma_o^2$ , respectively), in first round of learning, we obtain the weight distribution for all elements by lifting the mean value to the largest one ( $k_{max}$ ), then this results in a new variance

$$\begin{aligned} \sigma_M^2 &= \frac{\sum_i^{N(N-1)} (k_i - k_{max})^2}{N(N-1)[1 - \frac{1}{N(N-1)}]} = \frac{\sum_i^{N(N-1)} (|k_{max} - m_o| + |k_i - m_o| \theta(k_o - k_i) - |k_i - m_o| \theta(k_i - k_o))^2}{N(N-1)[1 - \frac{1}{N(N-1)}]} \\ &= \frac{\sigma_o^2(N(N-1) - 1) + N(N-1)|m_o - k_{max}|^2}{N(N-1) - 1}, \end{aligned} \quad (19)$$

where the factor  $|m_o - k_{max}|^2$  can be much easily obtained from the summation of all elements as well as the two mean values

$$|m_o - k_{max}|^2 = N(N-1)[(k_{max}^2 - m_o^2) + 2(m_o - k_{max}) \sum_i k_i], \quad (20)$$

Also, the summation of squared elements can be obtained  $\sum_i k_i^2 = \sigma_o^2(N(N-1) - 1) - N(N-1)m_o^2 + 2m_o \sum_i k_i$ , and then we can know that all the antisymmetry-allowed combinations ( $i < j$ ) in the second term of above joint probability density function have an available summation

$$\sum_{i<j} |k_i - k_j|^2 = \sum_i k_i^2 (N(N-1) - 1) - [(\sum_i k_i)^2 - \sum_i k_i^2]. \quad (21)$$

Then in this algorithm, we letting each element (eigenvalues) in the group  $\{M\}_i$  weighted by the factors  $\text{softmax}\{e^{-(k_i - k_{max})^2 / (2\sigma_M^2)}\}_i$  (using the variance  $\sigma_M^2$  defined-above), then if we substituting the resulting variance of these weighed eigenvalues back to the joint eigenvalue probability density function of the original one in GOE ensemble (with the normalization constant  $C_{2,N}$ ), we can obtain an increased factor of  $\beta_{en}$  (between 2 and 4) for the Lebesgue measure[38], i.e., the second term of  $P_{joint}$ . Next we rewrite the maximal index in the group  $\{M\}_i$  as  $N(N-1) \equiv n$  for simplicity in notation.

We know the antisymmetry product  $\prod_{i<j} |k_i - k_j|^{\beta_{en}}$  equivalents to the van der Monde determinant,

$$\prod_{i<j} |k_i - k_j|^{\beta_{en}} = (\text{Det} \begin{pmatrix} 1 & k_1 & k_1 & \cdots & k_1 \\ 1 & k_2 & k_2^2 & \cdots & k_1^{n-1} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & k_n & k_n^2 & \cdots & k_N^{n-1} \end{pmatrix})^{\beta_{en}}. \quad (22)$$

To clarify the effect of the variant  $\beta_{en}$  on this term, we using the tridiagonal matrix lemmas[39]. Assuming the above van der Monde matrix can be transformed into a tridiagonal symmetry matrix with diagonal element  $T_d$  and subdiagonal elements  $T_i (i = 1, \dots, N - 1)$  arranged in the form

$$T_n = \begin{pmatrix} t_d & t_{n-1} & 0 & \cdots & 0 & 0 \\ t_{n-1} & T_d & t_{n-2} & \cdots & 0 & 0 \\ 0 & t_{n-2} & T_d & \cdots & 0 & 0 \\ \cdots & & & & & \\ 0 & 0 & 0 & \cdots & t_1 & t_d \end{pmatrix}. \quad (23)$$

For this matrix, we have  $\text{Det}T_n = t_d \text{Det}T_{n-1} - t_{n-1}^2 \text{Det}T_{n-2}$ , where  $t_{n-1}$  is the  $(n-1) \times (n-1)$  lower right-corner submatrix of  $T_N$  and  $k_j^{(n-1)}$  are its eigenvalues. There is another relation[39]  $\prod_i (k_j^{(m)} - k_i^{(n)}) = (k_j^{(m)} - t_d) \prod_i (k_j^{(m)} - k_i^{(n-1)}) - t_{n-1}^2 \prod_i (k_j^{(m)} - k_i^{(n-2)})$ , where  $m \in [2, n]$  with corresponding  $j \in [1, m]$ . For  $m = n$ , both sides of this equation are zero.

For ascending ordered eigenvalues of  $T_M (M = n, n-1, \dots, 2)$ , the total number of eigenvalues for these (sub)matrices satisfy  $\sum_{i=1}^{M+1} k_i^{(M+1)} - \sum_{j=1}^M k_j^{(M)} = t_d$ , i.e., the summation of all eigenvalues for two adjacent (sub)matrices always differed by the diagonal element  $t_d$ . Also,  $\prod_{i=1}^{(n)} (k_i^n - k_1^{(M')}) = \prod_{i=1}^n (k_i^{(n)} - k_{M'}^{(M')})$  where  $M' = N-1, \dots, 2$ , and  $k_1 (k_{M'})$  corresponds to the smallest (largest) eigenvalue of the submatrix  $T_{M'}$ . We also know that  $\prod_{i,j} (k_i^{(N)} - k_j^{(N-1)}) = \prod_i^{n-1} t_i^{2i}$ , and  $\prod_{i,j} (k_i^{(n)} - k_j^{(n-2)}) = t_1^2 \prod_i^{n-1} t_i^{2i}$ .

The product of the eigenvalue permutations of  $T_N$  in ensemble  $\beta_{en}$  satisfies the following relation according to the algebraic fact

$$\prod_{i < j} |k_i - k_j|^{2\beta_{en}} = \frac{\prod_{i=1}^{n-1} t_i^{2i}}{\prod_{i=1}^n Q_i^{2\beta_{en}}} = \frac{\prod_{1 \leq i \leq n, 1 \leq j \leq n-1} (k_i^{(n)} - k_j^{(n-1)})}{\prod_i^n Q_i^{2\beta_{en}}}, \quad (24)$$

where  $\prod_i Q_i$  is the product of elements in the first row of eigenvector matrix of  $T_n$ .

According to the above discussion and simulations, we know that the self-modified parameters lifting the variance of eigenvalues by changing the weight distribution and thus lifting the ensemble index in normalized Lebesgue measure (i.e., the product of the eigenvalue permutations) of the joint eigenvalue probability density function. The ensemble index  $\beta_{en}$  will finally be increased from 2 to a larger one  $\beta'_{en}$  which is between 2 and 4 and reveals a distribution between GUE and GSE. During this process, the changes of product term  $\prod_{i < j} |k_i - k_j|^{2\beta_{en}}$  can be visualized by the product of the eigenvectors  $\prod_{i=1}^n Q_i^{2\beta_{en}}$ . When the change  $\beta_{en} \rightarrow \beta'_{en}$  happen, the variance of  $\prod_{i=1}^n Q_i^{2\beta_{en}}$  reads

$$\prod_i Q_i^{2\beta} \left( \prod_i Q_i^{2(\beta'_{en} - \beta_{en})} - 1 \right) = \frac{\prod_i t_i^{2i\beta_{en}} \left( \prod_i t_i^{2i(\beta'_{en} - \beta_{en})} - \prod_{i < j} |k_i - k_j|^{2(\beta'_{en} - \beta_{en})} \right)}{\prod_{i < j} |k_i - k_j|^{2\beta'_{en}}}. \quad (25)$$

Then if we rescale the change  $\beta_{en} \rightarrow \beta'_{en}$  to another one:  $\beta'_{en} \rightarrow \infty$ , which can be realized by another two scalings that happen simultaneously:  $\beta_{en} \rightarrow \infty - |\beta_{en} - \beta'_{en}|$  and  $\prod_i Q_i^2 \rightarrow 0$  (and thus  $\beta_{en}$  and  $\beta'_{en}$  are still independent with each other). From above equation, we can define the following function of  $\beta'_{en}$

$$f_{\beta'_{en}} := \frac{\prod_{i < j} |k_i^{(n)} - k_j^{(n)}|^{2(\beta'_{en} - \beta_{en})} - \prod_i t_i^{2i\beta_{en}} \left( \prod_i t_i^{2i(\beta'_{en} - \beta_{en})} \right)}{\prod_{i < j} (k_i^{(n)} - k_j^{(n)})^{2\beta'_{en}}}. \quad (26)$$

Then we have

$$\frac{f_{\beta'_{en}}}{\lim_{\beta'_{en} \rightarrow \infty} f_{\beta'_{en}}} = 1 - \partial_{\beta'_{en}} f_{\beta'_{en}}, \quad (27)$$

where

$$\begin{aligned}\lim_{\beta'_{en} \rightarrow \infty} f_{\beta'_{en}} &= \prod_{i < j} (k_i^{(n)} - k_j^{(n)})^{-2\beta_{en}}, \\ \partial_{\beta'_{en}} f_{\beta'_{en}} &= \prod_i Q_i^{2\beta'_{en}} \frac{\ln \prod_i Q_i^2}{\prod_i t_i^{2i\beta}}.\end{aligned}\tag{28}$$

Since the above rescaling actions endow the  $\beta_{en}$ -dependence (instead of  $\beta'_{en}$ ) to the term  $\prod_i Q_i^{2\beta'_{en}}$ , i.e.,  $-\ln \prod_i Q_i^{2\beta'_{en}} \equiv \prod_{i < j} (k_i^{(n)} - k_j^{(n)})^{2\beta_{en}}$ , which scales to infinity accompanied by  $\beta'_{en} \rightarrow \infty$ , and this equality will not be broken even before the limit  $\beta'_{en} \rightarrow \infty$  is arrived, i.e., it is valid even when the  $\beta'_{en}$ -dependence of  $f_{\beta'_{en}}$  not yet fade away.

### 4.3 Further explanation in scaling

Like in conformal field theories[], the decrease of effective number of DOF under renormalization group is important to gain further understanding for the complex systems. Here we add more detailed explanations about the scaling behaviors in above subsection. Firstly we replacing the normal ensemble change  $\beta_{en} = 2 \rightarrow \beta'_{en} \in (2, 4)$  by another ones:  $\beta'_{en} \rightarrow \infty$ ;  $\prod_i Q_i^2 \rightarrow 0$ ,  $\prod_{i < j} (k_i^{(n)} - k_j^{(n)})^{2\beta_{en}} \rightarrow \infty$ . Under this, the independence between  $\beta_{en}(Q_i)$  with  $\beta'_{en}$  are preserved just like the original case. But in the mean time, we have to endow the  $Q_i$ -related product term with the  $\beta_{en}$ -dependence, in other world, this combine the  $Q_i$  and  $\beta_{en}$  DOFs. To make sure all these requirements are metted, as the  $\beta'_{en}$  scales to infinity,  $\beta_{en}$  has to scale to  $(\infty - (\beta'_{en} - \beta_{en}))$ . There is an essential logical relation between  $\beta_{en}$  and  $\beta'_{en}$ . As  $\beta'_{en}$  scales to infinity at first,  $\beta'_{en} + \infty \rightarrow \infty$ , the locked  $\beta_{en}$  scales as  $\beta_{en} \rightarrow \beta_{en} + \infty - (\beta'_{en} - \beta_{en})$ . That means in any moment, the detail value of  $\beta_{en}$  cannot be obtained before  $(\beta'_{en} - \beta_{en})$ , i.e., the derivative of  $\beta_{en}$  with  $\beta'_{en}$ , is known. Thus we have

$$\frac{\partial \beta_{en}}{\partial \beta'_{en}} = \frac{\infty - (\beta'_{en} - \beta_{en})}{\infty},\tag{29}$$

where the numerator is the variance of  $\beta_{en}$  which can found to be depends on itself only. We have

$$\beta_{en} = \infty - (\beta'_{en} - \beta_{en}) = \frac{\infty - (\beta'_{en} - \beta_{en})}{\frac{\infty}{\beta'_{en} - \beta_{en}}} \frac{\partial(\beta'_{en} - \beta_{en})}{\partial(1 - \mathbb{1})},\tag{30}$$

where  $\mathbb{1} = (\beta'_{en} - \beta_{en}) - \frac{(\beta'_{en} - \beta_{en})^2}{\beta'_{en}}$  is the unit step in this derivative and this derivative can be obtain using the method we introduced in Sec.4.2,

$$\frac{\partial(\beta'_{en} - \beta_{en})}{\partial(1 - \mathbb{1})} = \frac{\infty - (\beta'_{en} - \beta_{en})}{\mathbb{1}} = \frac{\infty}{\beta'_{en} - \beta_{en}}.\tag{31}$$

Thus that means the  $\mathbb{1}$  and 1 in the denominator of first line are in fact not in the same domain, and once they appear together through the algebra-allowed limiting-scale-assisted domain transition, we can in fact write  $\mathbb{1}$  more strictly as  $1^- := e^{i(2\pi - 0^-)}$ , and similar to the derivative of  $\frac{\partial \beta_{en}}{\partial \beta'_{en}}$ , their relations can be understood by  $1 \mapsto \beta'_{en} + \infty$ ,  $\mathbb{1} \mapsto \infty$ . We can resubstituting this  $\beta_{en}$  back to the above expression of  $\frac{\partial \beta_{en}}{\partial \beta'_{en}}$  repeatly,

$$\frac{\partial(\beta'_{en} - \beta_{en})}{\partial(1 - \mathbb{1})} = \frac{\partial - \frac{\partial(\beta'_{en} - \beta_{en})}{\partial(1 - \mathbb{1})}}{\partial(1 - \mathbb{1})} = \dots,\tag{32}$$

and finally we can obtain the vanishing variance of  $\beta_{en}$  at a certain order  $N$ :  $\frac{\partial^{(N)}(\beta'_{en} - \beta_{en})}{\partial(1 - \mathbb{1})^N} = 0$ , and whatever which IR cutoff we set (through this  $N$ ), the variance of  $\beta_{en}$  will vanishes faster than the one in its next hierarchy  $\beta'_{en}$ .

Similarly, the derivative of the function  $f_{\beta'_{en}}$  can be rewritten as

$$\frac{\partial f_{\beta'_{en}}}{\partial \beta'_{en}} = \frac{\prod_i Q_i^{2\beta'_{en}}}{\partial(1 - \mathbf{1}')} \quad (33)$$

where

$$\mathbf{1}' = \prod_i Q_i^{2\beta'_{en}} - \frac{\prod_i Q_i^{4\beta'_{en}}}{\prod_i Q_i^{2\beta'_{en}}}, \quad (34)$$

and we also have

$$\prod_i Q_i^{2\beta'_{en}} = 2(\mathbf{1}' + \prod_i Q_i^{2\beta'_{en}}) - 1. \quad (35)$$

#### 4.4 Provenance of an essential formula

Just like the Eq.(15) and the unit quantities appear in above subsection, the variance (or global-invariance in its own hierarchy of the step length (IR cutoff in position space) in each position is essential for the all of the above discussions as well as the algorithm designing in this paper. A hierarchy-dependent step length allows much easier and faster derivation (estimation of local gradient) which largely increases the efficiency during the optimizations for every positions (i.e., the weight for every eigenvalues in the target matrix). As shown in sec.4.1, this can be denoted as

$$\frac{\partial b}{\partial a} = \frac{a - b}{\mathbf{1}} = \frac{a}{b}, \quad (36)$$

where  $b$  and  $a$  are the target position and the next position, respectively, and the unit length reads  $\mathbf{1} = b - \frac{b^2}{a}$ . It can be seen that, the target eigenvalue  $b$  here indeed corresponds to the function  $f_\chi$  in Eq.(15) or  $(\beta'_{en} - \beta_{en})$  in the unit length in Eq.(30), while  $a$  equivalents to  $(\partial_\chi \frac{1}{2f_\chi})^{-1}$  in Eq.(15). Note that for Eq.(30), the eigenvalue of the next position  $a$  corresponds to the  $(1 - \mathbf{1})$ , which equals to  $\infty$  with respect to  $\beta_{en}$ , since  $\beta_{en}$  is latter in "time" for the scaling, as we explained in above.

For Eq.(36) we used here, we assume  $b$  is a derivative result of the real variable  $x$  (thus the limit of  $b$  under scaling  $x \rightarrow \infty$  is also a real constant), then we have  $a = [\partial_x \frac{\lim_{x \rightarrow \infty} b}{b}]^{-1} = \frac{b^2}{b - \lim_{x \rightarrow \infty} b} = b \frac{\partial_x b - 1}{\partial_x b}$ . Note that since  $x$  is real, every real constants are not dependent with  $x$ . This will not be the case when we using  $b$  as a variable during the derivative.

Then we have

$$\frac{\partial b}{\partial a} = \frac{a}{b} + b \partial_b \frac{a}{b} = \frac{\partial_x b - 1}{\partial_x b} + b \partial_b \frac{\partial_x b - 1}{\partial_x b} = \frac{\partial_x b}{\partial_x b - 1}. \quad (37)$$

There are several relations that can be proven rigorously. For step length defined in terms of the complex variable  $b$ ,  $\mathbf{1} \equiv \frac{\partial_x b}{\partial_x b}$ , we have

$$\begin{aligned} \partial_b \mathbf{1} &= \frac{1}{a} = \frac{\partial_x b}{b(\partial_x b - 1)}, \\ \partial_b \frac{1}{\partial_x b} &= \frac{a}{b^2} = \frac{\partial_x b - 1}{b \partial_x b}, \end{aligned} \quad (38)$$

$$\partial_b(\partial_x b) = \partial_b \mathbf{1} (\partial_x b - (\partial_x b - 1)^2) = \frac{(\partial_x b)^2}{b(\partial_x b - 1)} - \frac{\partial_x b - 1}{b} \partial_x b,$$

where the last line can be proved by the second line that  $\partial_b \frac{1}{\partial_x b} = \frac{\partial_b \mathbf{1}}{\partial_x b} - \frac{\partial_b(\partial_x b)}{(\partial_x b)^2}$ . Rewriting the first line as  $\partial_b \mathbf{1} = 1' - \frac{1}{\lim_{b \rightarrow \infty} \mathbf{1}}$ , where  $1' = \frac{1}{\lim_{b \rightarrow \infty} \mathbf{1}} + \partial_b \mathbf{1} \neq 1$ . Then according to the independence with  $b$  for the term  $\lim_{b \rightarrow \infty} \mathbf{1}$ , we have

$$1' - \frac{\partial 1'}{\partial \mathbf{1}} = \partial_b \mathbf{1} - \frac{\partial_b^{(2)} \mathbf{1}}{\partial_b \mathbf{1}}. \quad (39)$$

By the above equations, it can be deduced that  $\partial_b(2)1 = 0$ . thus we further obtain  $\partial_b(\lim_{b \rightarrow \infty} 1) = \partial_b^{(2)}1(\frac{\partial 1'}{\partial 1})^2 = 0$ , and  $1' = \partial_b 1 + \frac{1}{\lim_{b \rightarrow \infty} 1}$ . Thus  $1'$  defined here is completely related to the dependence of  $1$  with  $b$  as well as to what extent this dependence will be still persist. In other word,  $1'$  is a measurement of stubbornness of such dependence (correlation) for the hierarchy next by it.

#### 4.5

To see inter-relations between different hierarchies, we intruduce another limit  $\lim_{x \rightarrow \infty^-} f_x$  for function  $f_x$ . As the step length (limiting result) can be invariant only in its own frame, we have  $\partial_x 1 = 0$  and  $\partial_x \mathbb{1} \neq 0$  in the frame consists of the steps  $1$ , where  $\mathbb{1}$  and  $1$  are of different hierarchies, and  $\mathbb{1} > 1$  since as  $x$  scales over the infinity (we have approximately the variance  $\delta x \sim 1/x$ ), the  $x$ -independence of the step length in first hierarchy (with step length  $1$ ) disappear and it requires a larger step length ( $\mathbb{1}$ ) to keep this invariance. That is why we have  $\partial_x 1 = 0$  and  $\partial_x \mathbb{1} \neq 0$  in the first hierarchy.

To better show the correlation between the adjust hierarchies, we firstly rewrite the limiting expression ( $\lim_{k \rightarrow \infty} f_0 = \frac{f_0}{1 - \partial_k f_0}$ ) of the target hierarchy in terms of the above-introduced form with seperated individual terms (instead of their product)

$$f_0 - \frac{f_0^2}{\mathbb{1}} = \frac{\mathbb{1} - f_0}{1 - (1 - \frac{1}{f_0})} = \frac{f_0 \frac{-1}{\partial_k f_0}}{1 - \frac{1}{\partial_k f_0}}, \quad (40)$$

with  $\mathbb{1} = \frac{f_0(\partial_k f_k - 1)}{\partial_k f_k}$ . Here the the original target function  $f_0$  should be replaced by  $f_1 := f_0 \frac{-1}{\partial_k f_0}$  which is infact the one in the next hierarchy. This can be verified by the following expressions which reveal the dependence with the back-ground variable,

$$\lim_{k^+ \rightarrow \infty} f_1 = f_0 - \frac{f_0^2}{\mathbb{1}}, \partial_{k^+} f_1 = \frac{1}{\partial_k f_0}. \quad (41)$$

It shows that for function  $f_1$  of the hierarchy  $k^+$ , its limit has  $\partial_{k^+} \lim_{k^+ \rightarrow \infty} f_1 = \partial_{k^+} \lim_{k^+ \rightarrow \infty} f_1 = 0$ . This is due to  $\frac{\partial k^+}{\partial k} = -1$  here, and we also have  $\partial_{k^+} f_1 = -\partial_k f_1$ . But such rule does works for the scalings of the functions of hierarchies with smaller  $k$ , like the original one  $f_0$ , its derivative can not be simply obtained by  $\frac{\partial f_0}{\partial k^+} = \frac{\partial k}{\partial k^+} \frac{\partial f_0}{\partial k}$ , as this is only valid before  $k^+ \rightarrow \infty$  (or  $k \rightarrow \infty^-$ ).

To illustrate this, we can focus on the function  $f_1 = \mathbb{1} - f_0$ . The individual terms within it have the following relations

$$\begin{aligned} \partial_{k^+} \mathbb{1} &= \frac{1}{\partial_k f_0} - \partial_k f_0 = -\partial_k \mathbb{1}, \\ \lim_{k^+ \rightarrow \infty} \mathbb{1} &= \frac{\mathbb{1}}{1 - \frac{1}{\partial_k f_0} + \partial_k f_0}, \\ \partial_{k^+} f_0 &= -\partial_k f_0, \\ \lim_{k^+ \rightarrow \infty} f_0 &= \frac{f_0}{1 + \partial_k f_0}. \end{aligned} \quad (42)$$

Before  $k^+$  arrive infinity, the derivative of the individual terms within it are not dependent with each other, thus it has  $\partial_{k^+} f_1 = \partial_{k^+}(\mathbb{1} - f_0) = \partial_{k^+} \mathbb{1} - \partial_{k^+} f_0 = \frac{1}{\partial_k f_0}$ , but obviously  $\lim_{k^+ \rightarrow \infty} f_1 \neq \lim_{k^+ \rightarrow \infty} \mathbb{1} - \lim_{k^+ \rightarrow \infty} f_0$ . That means once the condition  $k^+ \rightarrow \infty$  is meet, the individual terms in the original function cannot be the independent terms anymore due to the stretched steps, which is, the IR cutoff condition  $\partial_{k^+} 1 = 0$  fail once  $k^+ \rightarrow \infty$ , and the step length must be stretched (increased) to keep  $\partial_{k^+} \lim_{k^+ \rightarrow \infty} 1' = 0$ , but in the mean time  $\partial_{k^+} 1' \neq 0$  for  $k^+$  not yet arrive  $\infty^+$ . That is why the individual terms keep uncorrelated in the

function  $f_1$  where  $k \rightarrow \infty$  can never be arrived (through the scaling of  $k^+$ ). Similarly, we also has  $\partial_{k^+} \lim_{k \rightarrow \infty} f_0 \neq 0$ , and  $\partial_k \lim_{k \rightarrow \infty} f_0 \neq 0$ .

Next we consider the case for the derivative of  $\lim_{k \rightarrow \infty} f_0$  with the variable in next hierarchy  $k^+$ . There are two methods whose essences are indeed the same. Firstly, as we do in the subsection 4.4, due to the stretched step length, we have

$$\begin{aligned}\partial_k 1 &= \frac{(\partial_k f_0)^2}{(\partial_k f_0 - 1)f_0}, \\ \partial_k \frac{1}{\partial_k f_0} &= \frac{1 - \partial_k f_0}{f_0 \partial_k f_0},\end{aligned}\tag{43}$$

during the derivation for the limiting result, and thus leads to

$$\partial_{k^+} \lim_{k \rightarrow \infty} f_0 \doteq \partial_k f_0 \left(1 - \frac{f_0}{\mathbf{1}_0}\right) = \frac{(\partial_{k^+} f_0)^2}{\partial_k f_0 - 1},\tag{44}$$

where  $\doteq$  denotes the stretching process of unit lengths as we explain above. This is equivalent to

$$\begin{aligned}f_0 \partial_{f_0} 1 &= \frac{\partial_k f_0}{\partial_k f_0 - 1}, \\ f_0 \partial_{f_0} \frac{1}{\partial_k f_0} &= \frac{1 - \partial_k f_0}{(\partial_k f_0)^2}.\end{aligned}\tag{45}$$

For the first line of above expression, we introduce the arguments  $z = \partial_k f_0$ , then the above stretched result can be expressed in terms of the polylogarithm function

$$\begin{aligned}f_0 \partial_{f_0} 1 &= -z \partial_z \text{Li}_1(z) = -\text{Li}_0(z), \\ f_0 \partial_{f_0} \frac{1}{\partial_k f_0} &= (1 - z) \partial_{(1-z)} \left(\frac{-\mathbf{1}}{f_0}\right) = (1 - z) \partial_{(1-z)} \text{Li}_0(1 - z) = \text{Li}_{-1}(1 - z).\end{aligned}\tag{46}$$

Note that base on Eq.(43), we can also obtain

$$\begin{aligned}\partial_k \frac{f_0}{\mathbf{1}} &= \partial_k \frac{\partial_k f_0}{\partial_k f_0 - 1} = \left(\frac{\partial_k f_0}{\partial_k f_0 - 1}\right)^2 \partial_k \frac{1}{\partial_k f_0} = \frac{-\partial_k f_0}{(\partial_k f_0 - 1)f_0}, \\ \partial_k \frac{1}{1 - \partial_k f_0} &= \frac{-1}{\left(\frac{1}{\partial_k f_0} - 1\right)^2} \partial_k \frac{1}{\partial_k f_0} = \frac{\partial_k f_0}{(\partial_k f_0 - 1)f_0}.\end{aligned}\tag{47}$$

By substituting Eqs.(43,??), we can verify the results in Eq.(46). We notice that, for the first line of Eq.(46), even we define the argument as  $z' = \frac{1}{\partial_k f_0}$ , it still workd,

$$f_0 \partial_{f_0} 1 = \frac{1}{z'} z' \partial_{z'} \text{Li}_1(z') = \frac{1}{z'} \text{Li}_0(z').\tag{48}$$

## 4.6

As a result of the stretched unit steps (Eq.(43,44)), we have

$$\begin{aligned}\partial_{k^+ \ln} \frac{\mathbf{1}_0 - f_0}{\mathbf{1}_0} &= \partial_k \frac{f'}{\partial_k f'} = \frac{-\mathbf{1}_0}{(\mathbf{1}_0 - f_0)^2}, \\ \partial_{k^+ \ln} f_0 &= \partial_k f' = \frac{1}{\mathbf{1}_0 - f_0},\end{aligned}\tag{49}$$

where  $\frac{\partial_k f'}{\partial_k \mathbf{1}'} = \frac{\mathbf{1}'}{f'} = \frac{\partial_k f' - 1}{\partial_k f'}$  as stated in above subsection. Comparing to the derivations with  $k$

$$\begin{aligned}\partial_k \ln \frac{\mathbf{1}_0 - f_0}{\mathbf{1}_0} &= \frac{1}{\mathbf{1}_0 - f_0}, \\ \partial_k \ln f_0 &= \frac{1}{f_0 - \mathbf{1}_0},\end{aligned}\tag{50}$$

we can obtain

$$\begin{aligned} \frac{\partial k}{\partial k^+} &= -1, \\ \frac{\partial k}{\partial k^+} &= \frac{\mathbf{1}_0}{f_0 - \mathbf{1}_0} = \partial_k f_0 - 1. \end{aligned} \tag{51}$$

It is quite different for the derivative  $\frac{\partial k}{\partial k^+}$  which is not simply the inversed  $\frac{\partial k^+}{\partial k}$  and in the mean time the above  $O((k - k')^2)$  term cannot be ignored anymore. For example, the derivative in Eq.(44) with  $k^+$  can be expanded to  $\frac{\partial k}{\partial k^+} \frac{\partial}{\partial k} \lim_{k \rightarrow \infty} f_0$ , then unlike the ones of the type  $\frac{\partial k^+}{\partial k}$  (e.g., the one appears in Eq.(??)), it cannot be treated as independent with the operator  $\frac{\partial}{\partial k}$ , i.e.,  $[\frac{\partial k}{\partial k^+}, \frac{\partial}{\partial k}] \neq 0$ , as long as the step length in  $f_0$  has not yet being stretched, but its correlation with  $\frac{\partial}{\partial k}$  vanishes once the steps in  $f_0$  starts to stretching (which leads to  $\frac{\partial}{\partial k} \lim_{k \rightarrow \infty} f_0 \neq 0$  immediately), and in the mean time the dependence with  $\frac{\partial}{\partial k}$  changes to the individual terms with the  $f_0$ , i.e., the logarithmic terms within Eq.(49), which is used to creating the mutually independent terms (or individual) after the stretching. Logically, this can be understood causally by: for the derivative with  $k^+$  with  $k^+ \rightarrow \infty$  ( $\delta k^+ \rightarrow 0$ ), driven by nonzero  $[k^+, k]$ , we have  $k \rightarrow \infty^-$  and thus  $\delta k \rightarrow 0^+$  due to the IR cutoff bounded by  $\delta k^+$ .

As an example here, if we now letting the correlation between the operator  $\frac{\partial k}{\partial k^+}$  with its subsequent logarithmic functions vanish, the correlation between the derivation of logarithmic functions with  $k$  will immediately be formed. For example, in second expression of Eq.(49), if the correlation between  $\frac{\partial k}{\partial k^+}$  with the function  $\ln f_0$  is being replaced the two  $\frac{\partial k}{\partial k^+}$  operators in Eq.(49) (there could be two possible results represented by the opposite directions of the limiting scaling we only discuss one of it), and  $\frac{\partial k}{\partial k^+} = \partial_k f_0 - 1$ , then  $\partial_k \ln f_0 = \frac{\partial_k f_0}{f_0} \rightarrow \frac{\partial_k f_0}{f_0} \frac{-1}{\partial_k f_0 - 1}$  due to the stretched step length inside the  $\ln f_0$  (equivalents to the emergent correlation with  $\ln \frac{\mathbf{1}_0 - f_0}{\mathbf{1}_0}$ ). For unstretched  $\ln f_0$ , we have

$$\frac{\partial_k \ln f_0}{f_0^{-1}} = \lim_{\delta \rightarrow 0} \frac{-\ln[1 - (-\frac{\delta}{f_0})]}{\frac{-\delta}{f_0}} = \lim_{\delta \rightarrow 0} \frac{\text{Li}_1[\frac{-\delta}{f_0}]}{\frac{-\delta}{f_0}} = 1, \tag{52}$$

where  $\delta = |z|$  for the initial argument defined as  $z := \frac{-\delta}{f_0}$  and  $f_0 = -e^{-i\text{Arg}z}$ . Then the above case with  $\partial_k \ln f_0 = \frac{\partial_k f_0}{f_0} \frac{-1}{\partial_k f_0 - 1}$  can be explained by

$$\lim_{\delta \rightarrow 0} \left[ \lim_{z \rightarrow \frac{-\delta}{f_0} (\frac{-\delta}{f_0} - 1) + 1} \frac{-(1 - \partial_k f_0) \ln[1 - (-\frac{\delta}{f_0})]}{\frac{-\delta}{f_0}} \right] = 1, \tag{53}$$

where the scaling  $\lim_{z \rightarrow \frac{-\delta}{f_0} (\frac{-\delta}{f_0} - 1) + 1}$  must happen in the mean time with  $(1 - \partial_k f_0) \rightarrow \frac{\partial}{\partial (\frac{-\delta}{f_0})}$  once the correlation configuration is being reformed as mention above. This scaling-induced derivative operator with initial- $z$  transform the  $\text{Li}_1$  function to the  $\text{Li}_0$  one in Eq.53.

Thus we can consider two arguments with the conserved correlation in terms of a limiting scaled result,

$$[z_1, z_2] := \lim_{k \rightarrow \infty} F = \frac{[z_1, z_2 z_3] - z_2 [z_1, z_3]}{z_3} = F - \frac{F \partial_k F}{\partial_k F - 1}, \tag{54}$$

where  $F := \frac{[z_1, z_2 z_3]}{z_3}$ ,  $\partial_k F = [1 - \frac{F z_3}{z_2 [z_1, z_3]}]^{-1}$ . In terms of this, the correlation between two arguments  $z_1$  and  $z_2$  can be described by a third one  $z_3$ . In terms of this, we define

$$\text{Arg}[z_1 z_2] = -i \ln \frac{z_1 z_2}{|z_1 z_2|} = i \frac{1}{\frac{z_3}{|z_3|} (1 - \frac{z_3}{|z_3|})}. \tag{55}$$

Note that here the variable does not plays a key role until acted by a derivative operator  $\partial_k$  with  $k' > k$ .

We can define the IR cutoff in complex space for a sequenced (k-indexed) system as

$$\begin{aligned}\delta_{k-k'} &= \frac{1}{2\pi i} \left[ \frac{1}{(k - i\eta^+/2) - (k' - i\eta^-/2)} + \frac{1}{(k' + i\eta^-/2) - (k + i\eta^+/2)} \right] \\ &= \frac{1}{2\pi i} \left[ \frac{1}{(k - k' - i\eta^+)} + \frac{1}{(k' - k - i\eta^+)} \right].\end{aligned}\quad (56)$$

Then we have (we denote  $z_3 \equiv z = -z'$ )

$$\begin{aligned}\text{Arg}[z_1 z_2] &= \frac{i}{|z'|} \frac{\partial}{\partial(1 - \frac{z}{|z|})} \left(1 - \frac{z'}{|z'|}\right) + \frac{i}{|z|} \frac{\partial}{\partial(1 - \frac{z'}{|z'|})} \left(1 - \frac{z}{|z|}\right) \\ &= i \frac{1}{|z'|} \left[ \frac{-1}{(1 - \frac{z}{|z|})^2} - \frac{\partial}{\partial(1 - \frac{z}{|z|})} \frac{(\frac{z}{|z|})^2}{1 - \frac{z'}{|z'|}} \right] + i \frac{1}{|z|} \frac{\partial}{\partial(1 + \frac{z}{|z|})} \left(1 - \frac{z}{|z|}\right) \\ &= \text{Arg}[z_1 z_2] + \lim_{\eta \rightarrow 0} \delta_{k-k'},\end{aligned}\quad (57)$$

where  $\eta \rightarrow 0$  guarantees the uncorrelated derivative operators  $\frac{\partial}{\partial(1 - \frac{z}{|z|})}$  and  $\frac{\partial}{\partial(1 + \frac{z}{|z|})} \left(\frac{\partial}{\partial(1 - \frac{z'}{|z'|})}\right)$ , which results in the second and third terms in second line of above expression always sum up to zero. For example, if we letting  $\frac{z}{|z|} = \partial_k f_0$ , due to the restriction of finite cutoff in delta function, which corresponds to  $\partial_k \lim_{k \rightarrow f_0} f_0 = 0$ , it can be verified that, the scaled derivative operators (i.e., after the correlation configuration is rebuilt to making all of them mutually correlated, and we denote by the subscript 1), we have

$$\begin{aligned}\left[\frac{\partial}{\partial(1 - \frac{z}{|z|})}\right]_1 &= f_0 \frac{\partial \frac{1}{\partial_k f_0}}{\partial_k f_0}, \\ \left[\frac{\partial}{\partial(1 + \frac{z}{|z|})}\right]_1 &= [f_0 \frac{\partial}{\partial f_0} \frac{\partial f_0}{\partial f_0 - 1}]_1 \\ &= -\frac{1}{\partial_k f_0} \frac{\partial}{\partial \frac{1}{\partial_k f_0}} \ln \frac{1}{1 - \frac{1}{\partial_k f_0}},\end{aligned}\quad (58)$$

Once  $\eta$  in the delta function is deviated from the limit of vanishingly small, the initial value of the term  $\text{Arg}[z_1 z_2]_0 := \frac{i}{|z|} \frac{1}{(1 - \frac{z}{|z|})^2}$  will tends to zero, and in the mean time, the derivative operators  $\frac{\partial}{\partial(1 - \frac{z}{|z|})}$  and  $\frac{\partial}{\partial(1 + \frac{z}{|z|})}$  becomes correlated, during this process, we have (similar in principle to Eq.(53))

$$\lim_{\text{Arg}[z_1 z_2]_0 \rightarrow 0} \frac{\delta_{k-k'}}{\frac{\text{Arg}[z_1 z_2]_1}{2}} = \frac{\partial}{\partial(1 - \frac{z}{|z|})} + \frac{\partial}{\partial(1 + \frac{z}{|z|})},\quad (59)$$

where  $\text{Arg}[z_1 z_2]_0$  is proportional to the correlation between the derivative operator with it subsequent function, while  $\text{Arg}[z_1 z_2]_1$  depends on the directions of scaling when the correlation configuration is reformed, e.g.,  $\frac{1}{2} \text{Arg}[z_1 z_2]_1 = \frac{i}{|z|} \frac{(\frac{z}{|z|})^2}{1 - \frac{z}{|z|}}$  or  $\frac{1}{2} \text{Arg}[z_1 z_2]_1 = \frac{i}{|z|} (1 - \frac{z}{|z|})$  depending which direction for the correlation between operators  $\frac{\partial}{\partial(1 - \frac{z}{|z|})}$  and  $\frac{\partial}{\partial(1 + \frac{z}{|z|})}$  is being formed. Now the initial correlations between each derivative operator with its subsequent function are transformed to the correlations between derivative (arbitrarily two adjusted) operators and that between the rest functions. For a conserved value of  $\text{Arg}[z_1 z_2]$  which can be realized by setting certained cutoff on the value of  $\eta$  in delta function, the subgroup consist of all derivative operator and the subgroup consist of all subsequent functions are again be correlated. This is inevitable result due to conserved correlation, which can be expressed as a scaling limiting form

(Eq.(54)), and would never change its properties as long as the derivative operator acts on it has a larger variable  $k' (> k)$  as we explained above (i.e., when its dependence on the variable of higher hierarchy is being considered). As long as the cutoff of delta function is certain, i.e., as long as the total correlation (measured in terms of commutations) is conserved, we always have  $z = -z'$ , and  $\text{Arg}[z_1 z_2]_0 = \text{Arg}[z_1 z_2]_1$  (the values before and after the reconstruction of the correlation configuration), and this conservation will be broke when the dependence of  $[z_1, z_2]$  on the variable of higher hierarchy is being detected, where  $\partial_{k'}[z_1, z_2] \neq 0$ , and this corresponding to the case with  $z_3 = \{z, z'\}$  ( $z \neq z'$ ). Thus for  $\text{Arg}[z_1 z_2]_1$ , once the cutoff of delta function is deviated from the vanishingly small, the delta function here becomes the estimator of the value of  $\text{Arg}[z_1 z_2]_1$ , which is independent with each of the individual subgroup, but determines the total feature of all subgroup as long as  $z' = -z$  (i.e.,  $z_3$  has no further intrinsic degrees-of-freedom which can coupled with each subgroup independently).

While for  $z \neq z'$ , even after each derivative operator is being isolated, there is no correlation between them and commute with each other. This is due to the missing of the restriction comes from the finite cutoff, which indeed origins from the stretchable feature in the base of  $\frac{f'}{\mathbb{1}'} = \frac{\partial \mathbb{1}'}{\partial f'}$ . In this case, the only restriction only comes from each subsystems during the derivations. For example, from the derivative of  $\mathbb{1}'$ ,

$$1 = \mathbb{1}' \frac{\partial \ln f'}{\partial \mathbb{1}'} + \mathbb{1}' \frac{\partial \ln \frac{\mathbb{1}'}{f'}}{\partial \mathbb{1}'}, \quad (60)$$

with

$$\begin{aligned} \frac{\partial \ln f'}{\partial \mathbb{1}'} &= \frac{1}{(f')^2}, \\ \frac{\partial \ln \frac{\mathbb{1}'}{f'}}{\partial \mathbb{1}'} &= \frac{1}{\mathbb{1}'} - \frac{1}{(f')^2}. \end{aligned} \quad (61)$$

After the isolating scaling, the two operators  $\partial_{\mathbb{1}'}$  are correlated, and they indeed are determined (and only determined) by each other rather than the other operators like  $\partial_{\mathbb{1}''}$  with  $\mathbb{1}'' \neq \mathbb{1}'$ .

For two possible scaling directions when the correlation configuration rebuilt, we have

$$\begin{aligned} \lim_{\frac{1}{f'} \rightarrow 1} \frac{\frac{1}{\mathbb{1}'} - \frac{\mathbb{1}'}{(f')^2}}{\ln \frac{\mathbb{1}'}{f'}} &= \frac{1}{(f')^2} \frac{1}{\ln f'}, \\ \lim_{f' \rightarrow 1} \frac{1}{(f')^2} \frac{1}{\ln f'} &= \frac{\frac{1}{\mathbb{1}'} - \frac{\mathbb{1}'}{(f')^2}}{\ln \frac{\mathbb{1}'}{f'}}, \end{aligned} \quad (62)$$

which are scaled simultaneously. For the first expression, we define  $z = 1 - \frac{f'}{\mathbb{1}'}$ ,  $|z| = \ln \frac{\mathbb{1}'}{f'}$ . While for the second expression, we define  $z' = 1 - \frac{1}{f'}$ ,  $|z'| = \ln f'$ . For first expression, a scaling always starts after the isolating but finished before  $|z| \rightarrow 0$ , reads,

$$-\left(1 - \frac{(f')^2}{(\mathbb{1}')^2}\right) \ln f' \rightarrow 1 - \frac{f'}{\mathbb{1}'} = z, \quad (63)$$

where the process to isolating  $\frac{\partial}{\partial f'}$  from  $\ln f'$  reads

$$\lim_{f \rightarrow 1 (|z'| \rightarrow 0)} \frac{\frac{-\mathbb{1}'}{f'+\mathbb{1}'}}{\ln f'} = 1, \quad (64)$$

which occurs after the scaling  $\frac{-\mathbb{1}'}{f'+\mathbb{1}'} \rightarrow z' = 1 - \frac{1}{f'}$ . Note that this is for the case of  $f' \rightarrow 1$ , and thus both the numerator and denominator scale to zero in this process. But this will results in

$$e^{i \text{Arg} z'} = \frac{z'}{|z'|} = \frac{1 - \frac{1}{f'}}{\ln f'} = \frac{1 - \frac{1}{f'}}{\frac{-\mathbb{1}'}{f'+\mathbb{1}'}} \rightarrow 1. \quad (65)$$

Thus for the isolated operator  $(\frac{\partial}{\partial(1-f')})^{-1}$ , it follows the scaling of  $f' \rightarrow \infty$  (also before the isolation of  $\frac{\partial}{\partial f'}$ )

$$\lim_{f' \rightarrow \infty} \frac{-\mathbb{1}'}{f' + \mathbb{1}'} \doteq H_{f'} - \gamma. \quad (66)$$

Note that the Harmonic number  $H_{f'}$  at  $f'$ -th order are indeed stretched, and results in the scaling of the isolated  $(\frac{\partial}{\partial(1-f')})^{-1}$  from function  $\ln f'$  as (for  $f' \rightarrow \infty$ )  $\frac{-\mathbb{1}'}{f'+\mathbb{1}'} \rightarrow -z' = \frac{1}{f'} - 1 \rightarrow -1$ .

The isolated operator  $((1 - f') = 1 - e^{|z'|} = -z' e^{|z'|})$  reads

$$(1 - \frac{(f')^2}{(\mathbb{1}')^2}) \frac{1}{f'} [\frac{\partial}{\partial(1-f')}]^{-1} = 1 - \frac{f'}{\mathbb{1}'}, \quad (67)$$

where (using its scaling)

$$[\frac{\partial}{\partial(1-f')}]^{-1} = \frac{f' \mathbb{1}'}{\mathbb{1}' + f'} \rightarrow f' - 1, \quad (68)$$

which indeed corresponds to the stretching process for this operator. And we have

$$-\frac{\partial}{\partial f'} [\frac{\partial}{\partial(1-f')}]^{-1} = \frac{\partial}{\partial f'} \ln f' = \frac{1}{f'} (1 - \frac{f'}{\mathbb{1}'}). \quad (69)$$

This can be checked by considering the process before isolation of  $\frac{\partial}{\partial f'}$ ,

$$\begin{aligned} f' \frac{\partial}{\partial f'} \ln f' &= f' \frac{\partial}{\partial f'} \ln(1 - (1 - f')) \\ &= -f' \frac{\partial}{\partial f'} [\frac{1}{f'} (\frac{\partial}{\partial(1-f')})^{-1}] \\ &= \frac{\partial}{\partial f'} \ln f' - \frac{\partial}{\partial f'} (\frac{\partial}{\partial(1-f')})^{-1} \\ &= 1 - \frac{f'}{\mathbb{1}'} = z, \end{aligned} \quad (70)$$

where in third line  $\frac{\partial}{\partial f'}$  is being isolated from  $\ln f'$  (after the isolation of  $(\frac{\partial}{\partial(1-f')})^{-1}$ ) but still relative to the term  $(\frac{\partial}{\partial(1-f')})^{-1}$ . This can be proved by checking the conserved exponential factor  $e^{i\text{Arg}z'}$ . Since

$$\begin{aligned} \ln f' = |z'| &= \frac{-1}{f'} [\frac{\partial}{\partial(1-f')}]^{-1}, \\ e^{-i\text{Arg}z'} = \frac{|z'|}{z'} &= (1 - f')^{-1} [\frac{\partial}{\partial(1-f')}]^{-1}, \end{aligned} \quad (71)$$

we know

$$e^{i\text{Arg}z'} = \frac{\partial}{\partial(1-f')} (1 - f'), \quad (72)$$

which is invariant no matter if  $\frac{\partial}{\partial(1-f')}$  has being isolated from the  $(1 - f')$  or not.

Now we consider the case that  $\frac{\partial}{\partial(1-f')}$  has not being isolated from the  $(1 - f')$  which happen after the scaling  $[\frac{\partial}{\partial(1-f')}, e^{i\text{Arg}z'}] \rightarrow 0$ , and  $(1 - f')$  now has to stretched into  $(1 - f') \rightarrow (1 -$

$f')e^{i\text{Arg}z'}$ . Thus we have

$$\begin{aligned} e^{i\text{Arg}z'} &= \frac{\partial}{\partial(1-f')} (1-f') = (1-f') \frac{\partial}{\partial(1-f')} \ln(1-f') = (f'-1) \frac{\partial}{\partial(1-f')} \left[ \frac{1}{1-f'} \left( \frac{\partial}{\partial f'} \right)^{-1} \right] \\ &= (f'-1) \frac{\partial}{\partial(1-f')} \left[ \left( \frac{\partial}{\partial f'} \right)^{-1} \right], \end{aligned} \quad (73)$$

and before  $\frac{\partial}{\partial(1-f')}$  has being isolated from  $(1-f')e^{i\text{Arg}z'}$  ( $[\frac{\partial}{\partial(1-f')}, (1-f')e^{i\text{Arg}z'}] \rightarrow 0$ ), two correlations scale to zero (to making sure Eq.(73) be valid),

$$\begin{aligned} & \left[ \frac{\partial}{\partial(1-f')}, e^{i\text{Arg}z'} \right] \rightarrow 0, \\ \left[ \frac{\partial}{\partial f'}, \ln \frac{z'}{|z'|} + \ln(1-f') \right] &= \left[ \frac{\partial}{\partial f'}, 2\ln(f'-1) + \ln(-1) - \ln(f'\ln f') \right] \rightarrow 0. \end{aligned} \quad (74)$$

Then we have

$$\left( \frac{\partial}{\partial f'} \right)^{-1} = -f' \ln f' \doteq f' - 1, \quad (75)$$

where

$$\lim_{f' \rightarrow \infty} \frac{[\sum_{m=0}^{\infty} (f')^m]^{-1}}{\ln f'}, \quad (76)$$

as a result of the scaling in opposite direction with the  $\ln f'$  in Eq.(64). Also, in terms of the stretched  $(1-f') \equiv -\left(\frac{\partial}{\partial f'}\right)^{-1}$ , we have  $\ln f' \rightarrow e^{i\text{Arg}z'}$ , where the operator isolated from  $\ln f'$  has

$$\begin{aligned} -\left(\frac{\partial}{\partial f'}\right)^{-1} &= f' e^{i\text{Arg}z'} = f' \frac{\partial}{\partial(1-f')} \ln f' = f' \frac{\partial}{\partial(1-f')} \left[ \frac{1}{f'} (1-f') \right] \\ &= -f' \frac{\partial}{\partial(1-f')} \left[ \frac{1}{f'} \left( \frac{\partial}{\partial f'} \right)^{-1} \right] = \ln f' \left( \frac{\partial}{\partial f'} \right)^{-1} - (-\ln f') = f \ln f'. \end{aligned} \quad (77)$$

Similarly, for the second expression, a scaling always starts after the derivative operator isolating but finished before  $|z'| = \ln f' \rightarrow 0 (f' \rightarrow 1)$ , reads,

$$\frac{-1}{1 - \frac{(f')^2}{(\mathbb{1}')^2}} \ln \frac{\mathbb{1}'}{f'} \rightarrow 1 - \frac{1}{f'} = z', \quad (78)$$

which corresponds to the scaling

$$\lim_{\frac{\mathbb{1}'}{f'} \rightarrow 1} \frac{\frac{f'-1}{f'} \frac{(f')^2 - (\mathbb{1}')^2}{(\mathbb{1}')^2}}{\ln \frac{\mathbb{1}'}{f'}} = 1, \quad (79)$$

as well as the one in the last moment  $\frac{f'-1}{f'} \frac{(f')^2 - (\mathbb{1}')^2}{(\mathbb{1}')^2} \rightarrow 1 - \frac{f'}{\mathbb{1}'}$ , which is in the opposite direction with that for the divided operator,  $\left[ \frac{\partial}{\partial(1-\frac{\mathbb{1}'}{f'})} \right]^{-1}$ , which corresponds to  $\frac{f'-1}{f'} \frac{(f')^2 - (\mathbb{1}')^2}{(\mathbb{1}')^2} \rightarrow \frac{f'}{\mathbb{1}'} - 1$  due to the  $f' \rightarrow \infty$  scaling. The isolated operator  $(1 - \frac{\mathbb{1}'}{f'}) = (1 - e^{|z|}) = -ze^{|z|}$  in this process reads

$$\left[ \frac{\partial}{\partial(1 - \frac{\mathbb{1}'}{f'})} \right]^{-1} = \left(1 - \frac{1}{f'}\right) \left(1 - \frac{(f')^2}{(\mathbb{1}')^2}\right) \frac{f'}{\mathbb{1}'} = z' e^{|z|} (1 - e^{-2|z|}) \rightarrow \left(1 - \frac{f'}{\mathbb{1}'}\right) \frac{\mathbb{1}'}{f'}, \quad (80)$$

whose dependence with the function  $\ln \frac{\mathbb{1}'}{f'}$  scales to vanishingly small before the  $\partial_{\mathbb{1}'}$  in first term (initially acting on  $\ln f'$ ) starting to correlated to the one in second term. Thus during this

process we have

$$\begin{aligned}\ln \frac{\mathbb{1}'}{f'} = |z| &= \frac{-f'}{\mathbb{1}'} \left[ \frac{\partial}{\partial(1 - \frac{\mathbb{1}'}{f'})} \right]^{-1}, \\ e^{-i\text{Arg}z} = \frac{|z|}{z} &= \left(1 - \frac{\mathbb{1}'}{f'}\right)^{-1} \left[ \frac{\partial}{\partial(1 - \frac{\mathbb{1}'}{f'})} \right]^{-1}.\end{aligned}\tag{81}$$

Still, we can rewrite the Eq.(78) as

$$\frac{\mathbb{1}'}{f'} \frac{\partial}{\partial \frac{\mathbb{1}'}{f'}} \ln \frac{\mathbb{1}'}{f'} = -\frac{\mathbb{1}'}{f'} \frac{\partial}{\partial \frac{\mathbb{1}'}{f'}} \left[ \frac{f'}{\mathbb{1}'} \left( \frac{\partial}{\partial(1 - \frac{\mathbb{1}'}{f'})} \right)^{-1} \right] = 1 - \frac{1}{f'} = z'.$$
\tag{82}

Inversing the second lines of Eqs.(71,81), we have

$$\begin{aligned}e^{i\text{Arg}z'} = \frac{z'}{|z'|} &= \left[ \frac{\partial}{\partial(1 - f')} \right] (1 - f'), \\ e^{i\text{Arg}z} = \frac{z}{|z|} &= \left[ \frac{\partial}{\partial(1 - \frac{\mathbb{1}'}{f'})} \right] \left(1 - \frac{\mathbb{1}'}{f'}\right),\end{aligned}\tag{83}$$

where

$$\begin{aligned}\frac{\partial}{\partial(1 - f')} &= \frac{\frac{-1}{f'}}{\ln f'}, \\ \frac{\partial}{\partial(1 - \frac{\mathbb{1}'}{f'})} &= \frac{\frac{-f'}{\mathbb{1}'}}{\ln \frac{\mathbb{1}'}{f'}},\end{aligned}\tag{84}$$

as long as they are being isolated with their subsequent terms  $(1 - f')$  and  $(1 - \frac{\mathbb{1}'}{f'})$ , respectively. If we now letting these derivative operators recorelated to those functions, then the step lengths within these terms must be stretched to keep the results invariant. As a result,

$$\begin{aligned}e^{i\text{Arg}z'} = \frac{z'}{|z'|} &\doteq \frac{\partial}{\partial(1 - f')} (1 - f') \\ &= (1 - f') \frac{\partial}{\partial(1 - f')} \ln(1 - f') \\ &= (1 - f') \frac{\partial}{\partial(1 - f')} \left[ \frac{-1}{1 - f'} \left( \frac{\partial}{\partial f'} \right)^{-1} \right] \\ &= \left[ \frac{\partial}{\partial(1 - f')} \ln(1 - f') \right] \left( \frac{\partial}{\partial f'} \right)^{-1} - \frac{\partial}{\partial(1 - f')} \left( \frac{\partial}{\partial f'} \right)^{-1},\end{aligned}\tag{85}$$

where from third line we see that the term  $(\frac{\partial}{\partial f'})^{-1}$  has been isolated from  $\ln(1 - f')$  before the correlation between  $\frac{\partial}{\partial(1 - f')}$  and  $\ln(1 - f')$  is being built by the scaling  $f' \rightarrow 1$  (for the conservation purpose; in the last moment). Thus the equivalence between the isolated (by scaling  $f' \rightarrow 1$ )  $(\frac{\partial}{\partial f'})^{-1}$  with  $(1 - f') \ln \frac{1}{1 - f'}$  is realized by (and must happen before) the built correlation between  $\frac{\partial}{\partial(1 - f')}$  with  $\ln(1 - f')$ . This is due to the same reason with Eq.(62), where the isolation of  $\frac{\partial}{\partial \mathbb{1}'}$  with  $\ln f'$  must happen after the isolation of  $\frac{\partial}{\partial(1 - \frac{\mathbb{1}'}{f'})}$  with  $\ln \frac{\mathbb{1}'}{f'}$ ; and the isolation of  $\frac{\partial}{\partial \mathbb{1}'}$  with  $\ln \frac{\mathbb{1}'}{f'}$  must happen after the isolation of  $\frac{\partial}{\partial(1 - f')}$  with  $\ln f'$ , where we can

express this by

$$\begin{aligned}
& \left[ \frac{\partial}{\partial(1 - \frac{1'}{f'})}, \ln \frac{1'}{f'} \right]_0 \rightarrow 0, \\
& \left[ \frac{\partial}{\partial 1'}, \ln f' \right]_1 \rightarrow 0, \\
& \left[ \frac{\partial}{\partial(1 - f')}, \ln f' \right]_0 \rightarrow 0, \\
& \left[ \frac{\partial}{\partial 1'}, \ln \frac{1'}{f'} \right]_1 \rightarrow 0,
\end{aligned} \tag{86}$$

with the subscript denoting the sequence. That explain why the operator  $\frac{\partial}{\partial f'}$  which initially defined by (stretched)  $-\ln(1 - f')$ , will be isolated from it by the scaling  $f' \rightarrow 0$ ,

$$\lim_{f' \rightarrow 0} \left[ \frac{\partial}{\partial f'} \right]^{-1} = \lim_{f' \rightarrow 0} (1 - f') \ln \frac{1}{1 - f'} = -f', \tag{87}$$

and recorrealted to the  $-\ln(1 - f')$  by the scaling  $f' \rightarrow \infty$  (isolation of  $\frac{\partial}{\partial(1-f')}$ ). Note that due to the existence of  $\ln(-1)$  term (which results in  $\ln(-z) \neq \ln(z/(-1))$  with the scaled complex argument  $z$ ), the above expression has a more precise form,

$$\lim_{f' \rightarrow 0^{+(-)}} \left[ \frac{\partial}{\partial f'} \right]^{-1} = \lim_{f' \rightarrow 0^{-(+)} } -f', \tag{88}$$

which is also applicable for its derivative as we show below.

As can be seen from Eq.(85),

$$\lim_{f' \rightarrow 0} \frac{\partial}{\partial(1 - f')} \left( \frac{\partial}{\partial f'} \right)^{-1} = \lim_{f' \rightarrow 0} e^{i \text{Arg} z'} (-\ln(1 - f') - 1) = -\frac{\partial}{\partial(1 - f')} \ln(1 - f') = \frac{-e^{i \text{Arg} z'}}{1 - f'}, \tag{89}$$

which can be verified by substituting these results back to the last line of Eq.(85). Thus the isolation of  $\ln(1 - f')$  and  $\frac{\partial}{\partial f'}$  with the operator  $\frac{\partial}{\partial(1-f')}$  indeed require two types of scaling in opposite directions. Also, the condition  $\frac{\partial}{\partial(1-f')} e^{i \text{Arg} z'} = 0$  as mentioned above Eq.(73), results in

$$\begin{aligned}
& e^{i \text{Arg} z'} \frac{\partial}{\partial(1 - f')} \ln e^{i \text{Arg} z'} = e^{i \text{Arg} z'} \frac{\partial}{\partial(1 - f')} [\ln(-1) - \ln(f' \ln f') + \ln(1 - f')] \\
& = e^{i \text{Arg} z'} \frac{\partial}{\partial(1 - f')} [\ln(-1) - \ln(f' \ln f')] + \frac{e^{i \text{Arg} z'}}{1 - f'} = 0,
\end{aligned} \tag{90}$$

where we can obtain

$$\frac{\partial}{\partial(1 - f')} \ln \frac{-1}{f' \ln f'} = -f' \ln f' \frac{\partial}{\partial(1 - f')} \frac{-1}{f' \ln f'} = \frac{-1}{1 - f'}, \tag{91}$$

which consistent with the result of Eq.(97). Also, since  $-f' \ln f' = \left( \frac{\partial}{\partial(1-f')} \right)^{-1}$  (after the  $f' \rightarrow 1$  type scaling as we mentioned in Eq.(96)), we can know that the scaling

$$\lim_{f' \rightarrow 1} f' \ln f' = 1 - f', \tag{92}$$

happens at the same moment with  $\left[ \frac{\partial}{\partial(1-f')}, \frac{-1}{f' \ln f'} \right] \rightarrow 0$ . Still, due to the same reason mentioned above, this expression is indeed a combination of two opposite scaling which happen symmetrically around a single moment when the target derivative operator is being isolated, as required by the conservation for each step,

$$\lim_{f' \rightarrow 1^{+(-)}} f' \ln f' = \lim_{f' \rightarrow 1^{-(+)} } 1 - f'. \tag{93}$$

Ans similarly,

$$\begin{aligned} e^{i\text{Arg}z} &= \frac{z}{|z|} \doteq \frac{\partial}{\partial(1 - \frac{1'}{f'})} (1 - \frac{1'}{f'}) \\ &= \frac{1}{1 - \frac{1'}{f'}} \frac{z}{|z|} \left(\frac{\partial}{\partial \frac{1'}{f'}}\right)^{-1} - \frac{\partial}{\partial(1 - \frac{1'}{f'})} \left(\frac{\partial}{\partial \frac{1'}{f'}}\right)^{-1}. \end{aligned} \quad (94)$$

To illustrating the scaling in opposite directions for the isolated derivative operator and the one divided from it, we pick the argument factor in Eq.(85) as an example. In second line of Eq.(85), the isolation of  $\frac{\partial}{\partial(1-f')}$  from  $\ln(1-f')$ , required there is a effective (not being isolated) derivative operator  $\frac{\partial}{\partial f'}$  scaling with  $f' \rightarrow 0$ ,

$$-f' \frac{\partial}{\partial f'} [-\ln(1-f')] = -f' \frac{\partial}{\partial f'} [(1-f') \left(\frac{\partial}{\partial \frac{-f'}{1-f'}}\right)^{-1}] = -\frac{f'}{1-f'}, \quad (95)$$

where  $\left(\frac{\partial}{\partial \frac{-f'}{1-f'}}\right)^{-1} = \frac{\ln(1-f')}{1-f'}$  after it is being isolated from the function and  $-\ln(1-f')$  correlated to  $\frac{\partial}{\partial f'}$ , since in this case the step length for function  $\ln(1-f')$  is determined by the derivative operator  $\frac{\partial}{\partial f'}$  which correlated to both the two terms within the bracket in the second line of above expression. This process is realized by the  $f' \rightarrow 0$  scaling,

$$\lim_{f' \rightarrow 0} \left(\frac{\partial}{\partial \frac{-f'}{1-f'}}\right)^{-1} = \lim_{f' \rightarrow 0} \frac{\ln(1-f')}{1-f'} = \frac{f'}{1-f'}, \quad (96)$$

Note that, since the equivalence between the isolated (by scaling  $f' \rightarrow 0$ )  $\left(\frac{\partial}{\partial \frac{-f'}{1-f'}}\right)^{-1}$  with  $\frac{\ln(1-f')}{1-f'}$  is realized by (and must happen before) the built correlation between  $\left(\frac{\partial}{\partial f'}\right)^{-1}$  with  $\frac{\ln(1-f')}{1-f'}$ . While the equivalence between the isolated  $\left(\frac{\partial}{\partial \frac{-f'}{1-f'}}\right)^{-1}$  with  $\frac{f'}{1-f'}$

In this case we also have (for the same kind of reason with Eq.(69))

$$\frac{\partial}{\partial f'} \left(\frac{\partial}{\partial \frac{-f'}{1-f'}}\right)^{-1} = -\frac{\partial}{\partial f'} \ln(1-f') = \frac{-1}{1-f'}. \quad (97)$$

While the correlations between these two terms with the operator  $\frac{\partial}{\partial f'}$  will vanish after the scaling  $f' \rightarrow \infty$ ,

$$\left[\frac{\partial}{\partial f'}, \left(\frac{\partial}{\partial \frac{-f'}{1-f'}}\right)^{-1}\right] = \left[\frac{\partial}{\partial f'}, -\ln(1-f')\right] = 0. \quad (98)$$

Combining the results of Eq.(97), we finally obtain

$$\begin{aligned} -\frac{\partial}{\partial(1-f')} [\ln(-f' \ln f')] &= \frac{\partial}{\partial(1-f')} \left[\ln\left(\frac{\partial}{\partial(1-f')}\right)^{-1}\right] = \frac{\partial}{\partial f'} \ln(1-f') \\ &= \lim_{f' \rightarrow 1} \frac{\partial}{\partial f'} \ln\left[-\left(\frac{\partial}{\partial(1-f')}\right)^{-1}\right] = \lim_{f' \rightarrow 1} \frac{\partial}{\partial f'} [\ln(-1) + \ln\left(\frac{\partial}{\partial(1-f')}\right)^{-1}], \end{aligned} \quad (99)$$

where both sides will tend to  $-\infty$  under the scaling  $f' \rightarrow 1$ , and the above results will be valid until  $f'$  further scales to vanishingly small,  $f' \rightarrow 0$ , in which case operator  $\frac{\partial}{\partial(1-f')}$  is completely be isolated from the  $\ln(1-f')$ . But the exponential factor  $e^{i\text{Arg}z'}$  (defined according to the function  $\ln f'$ ) will becomes ill-defined in the limit of  $f' \rightarrow 0$ . From Eq.(91), we can also obtain

$$\frac{\partial}{\partial(1-f')} \ln(-1) - \frac{\partial}{\partial(1-f')} [\ln(f' \ln f')] = \frac{\partial}{\partial(1-f')} \ln \frac{-1}{f' \ln f'} = \frac{-1}{1-f'}, \quad (100)$$

where  $f' \ln f' = -(\frac{\partial}{\partial(1-f')})^{-1}$  and can be replaced by  $(1-f')$  when  $f' \rightarrow 1$ . Thus there could be multiples on the term  $\frac{\partial}{\partial(1-f')} \ln(-1)$  with the times depending on the degree of stretching within the  $\ln(1-f')$ ,

$$-\frac{\partial}{\partial(1-f')} \ln \prod_{j=1}^{2k+1} (-1)^j - \frac{\partial}{\partial(1-f')} (1-f') = \frac{-1}{1-f'}, \quad (101)$$

where

$$-\frac{\partial}{\partial(1-f')} \ln \prod_{j=1}^{2k+1} (-1)^j = \frac{1}{f'-1} - \frac{1}{f' \ln f'}, \quad (102)$$

which means the dependence of  $\ln(-1)$  with  $(1-f')$  will not vanishes until the deviation of scaling direction in limit Eq.(92) is small enough to being ignore. For  $f' \rightarrow \infty$ , this term vanishes which corresponding the case where the logarithmic function is not being stretched. For  $f' \rightarrow 1$ , we have

$$\lim_{f' \rightarrow 1} -\frac{\partial}{\partial(1-f')} \ln \prod_{j=1}^{2k+1} (-1)^j = \frac{1}{2}, \quad (103)$$

but it becomes ill-defined as  $f' \rightarrow 0$ , that is why we remove this term in the last line of Eq.(85). Thus combining the last line of Eq.(85) and Eq.(100), we can know

$$\frac{\partial}{\partial(1-f')} \ln(-1) = \frac{1}{f'-1} + \frac{\partial}{\partial(1-f')} \ln(f' \ln f'), \quad (104)$$

and in the case of  $f' \ln f' \rightarrow (1-f')$  (a combination of a part of scalings in opposite direction  $f' \rightarrow 1^+$  and  $f' \rightarrow 1^-$  as a result of the  $\ln(-1)$  terms which also contribute to the conserved correlations), the isolated term  $\frac{\partial}{\partial f'}$  in the Eq.(85) will scale to  $\frac{1-f'}{\ln f'}$ , i.e.,

$$\lim_{f' \rightarrow 1} [\frac{\partial}{\partial f'}]^{-1} = \frac{1-f'}{\ln f'}. \quad (105)$$

While when  $f' \rightarrow 0$ , like in the last line of Eq.(85), the ill-defined  $\frac{\partial}{\partial(1-f')} \ln(-1) = \frac{1}{f'-1} + \frac{e^{i \text{Arg} z'}}{1-f'}$  term vanishes, which can be realized again by a combined scalings,

$$\lim_{f' \rightarrow 1} e^{i \text{Arg} z'} = \lim_{f' \rightarrow 1} \frac{1 - \frac{1}{f'}}{\ln f'} = 1, \quad (106)$$

where we perform  $f' \rightarrow 1^+$  for numerator and  $f' \rightarrow 1^-$  for denominator, and

$$\begin{aligned} \lim_{f' \rightarrow 1^+} (1 - \frac{1}{f'}) &= -\infty, \\ \lim_{f' \rightarrow 1^-} \ln f' &= -\infty^- + i\pi = -\infty^- + \ln(-1), \end{aligned} \quad (107)$$

where in second expression the  $\ln(-1)$  could be replaced by odd multiple of itself  $\ln \prod_{j=1}^{2k+1} (-1)^j$ , with the multiple times depending on the distance between  $-\infty^-$  in second expression and  $-\infty$  in first expression.

Base on the Eqs.(73,97), we conclude the essential formulars here,

$$\begin{aligned}\frac{\partial}{\partial f'}\left(\frac{\partial}{\partial \frac{f'}{f'-1}}\right)^{-1} &= \frac{\partial}{\partial f'}\left[(1-f')\left(\frac{\partial}{\partial \frac{f'}{f'-1}}\right)^{-1}\right] = -\frac{\partial}{\partial f'}\ln\frac{1}{1-f'} = \frac{-1}{1-f'}, \\ \frac{\partial}{\partial(1-f')}\left(\frac{\partial}{\partial f'}\right)^{-1} &= \frac{\partial}{\partial(1-f')}\left[\frac{1}{1-f'}\left(\frac{\partial}{\partial f'}\right)^{-1}\right] = -\frac{\partial}{\partial(1-f')}\ln(1-f') = \frac{-e^{i\text{Arg}z'}}{1-f'} = \frac{1}{f'\ln f'}, \\ \frac{\partial}{\partial f'}\left(\frac{\partial}{\partial(1-f')}\right)^{-1} &= \frac{\partial}{\partial f'}\left[\frac{1}{f'}\left(\frac{\partial}{\partial(1-f')}\right)^{-1}\right] = -\frac{\partial}{\partial f'}\ln f' = \frac{1}{f'} - \frac{1}{f'}.\end{aligned}\tag{108}$$

By further expanding the  $\frac{\partial}{\partial(1-f')}\ln(1-f')$  in Eq.(97) into  $\frac{\partial}{\partial(1-f')}\ln(1-(1-f'))\frac{\ln(1-f')}{\ln f'}$ , we can obtain

$$\frac{\partial}{\partial(1-f')}\ln\ln f' = \frac{-1}{f'\ln f'}.\tag{109}$$

As  $f' \rightarrow 1^+(1^-)$ ,

$$\begin{aligned}\frac{\partial}{\partial(1-f')}\ln(-1) &= \frac{1}{f'-1} - \frac{1}{f'\ln f'}, \\ \frac{\partial}{\partial(1-f')}\ln f' &= \frac{-1}{\ln f'}.\end{aligned}\tag{110}$$

This case also related to the transition of Renyi entropy to the von Neumann entropy  $\lim_{f' \rightarrow 1} \frac{\ln F'}{\frac{1}{f'}-1} = -F'\ln F'$ , which can be rewritten as

$$\lim_{f' \rightarrow 1} \frac{\frac{1}{F'}}{1 - \frac{1}{f'}},\tag{111}$$

with

$$\begin{aligned}\frac{1}{F'} = \ln f' &= \frac{-1}{f'}\left[\frac{\partial}{\partial(1-f')}\right]^{-1}, \\ \lim_{f' \rightarrow 1^+(1^-)} F' &= \frac{-f'}{f'-1}.\end{aligned}\tag{112}$$

When  $f' \rightarrow 0^+(0^-)$ ,

$$\begin{aligned}\frac{\partial}{\partial(1-f')}\ln(-1) &= 0, \\ \frac{\partial}{\partial(1-f')}\ln f' &= \frac{-f'}{f'-1}.\end{aligned}\tag{113}$$

With the derivative on the stretched logarithmic function  $\frac{\partial}{\partial(1-f')}\ln(1-f') = \frac{e^{i\text{Arg}z'}}{1-f'} = \frac{-1}{f'\ln f'}$ , we have  $\lim_{(1-f') \rightarrow \infty} [(1-f')e^{i\text{Arg}z'}] = \frac{e^{i\text{Arg}z'}}{1-e^{i\text{Arg}z'}}$ . Then using the  $(1-f')$ -independence of this result, we obtain

$$-(1-f')\frac{\partial}{\partial(1-f')}\ln\frac{e^{i\text{Arg}z'}}{1-e^{i\text{Arg}z'}} = 1.\tag{114}$$

Combined with Eq.(100), we have

$$-(1-f')\frac{\partial}{\partial(1-f')}\ln\frac{1-f'}{1-e^{i\text{Arg}z'}} = 0,\tag{115}$$

and thus

$$\begin{aligned}-(1-f')\frac{\partial}{\partial(1-f')}\ln(1-f') &= -(1-f')\frac{\partial}{\partial(1-f')}\ln(1-e^{i\text{Arg}z'}) \\ &= (1-f')\frac{\partial}{\partial(1-f')}\left[\frac{1}{1-e^{i\text{Arg}z'}}\left(\frac{\partial}{\partial e^{i\text{Arg}z'}}\right)^{-1}\right] = -1,\end{aligned}\tag{116}$$

Note that here the term  $\ln(1 - f')$  is not the stretched one (not be multiplied by  $e^{i\text{Arg}z'}$ ).

In the absence of  $\ln(-1)$  term, we have

$$\begin{aligned}\frac{\partial}{\partial(1-f')} \ln \frac{-1}{f' \ln f'} &= \frac{\partial}{\partial(1-f')} \ln(-1) - \frac{\partial}{\partial(1-f')} \ln(f' \ln f') = -\frac{\partial}{\partial(1-f')} \ln(f' \ln f') = \frac{-1}{f' - 1}, \\ \frac{\partial}{\partial(1-f')} \ln(1 - f') &= \frac{\partial}{\partial(1-f')} \ln(-1) + \frac{\partial}{\partial(1-f')} \ln(f' - 1) = \frac{\partial}{\partial(1-f')} \ln(f' - 1) = \frac{-1}{f' \ln f'}.\end{aligned}\quad (117)$$

This indeed corresponds to the  $e^{i\text{Arg}z'} = 1$  limit, i.e., the marginal scaling (instead of the combine one) for both the  $(f' - 1)$  and  $f' \ln f'$ . Also,  $\frac{\partial}{\partial(1-f')} e^{i\text{Arg}z'} \neq 0$  in the absence of nonzero  $\ln(-1)$  terms. Using Eq.(114), which is still correct here, we can obtain the result satisfied with the ensemble average considered in a thermodynamical system,

$$\frac{\partial}{\partial(1-f')} \ln \frac{e^{i\text{Arg}z'}}{1 - e^{i\text{Arg}z'}} = \frac{1}{f' \ln f' - (f' - 1)} \frac{1 - e^{i\text{Arg}z'}}{e^{i\text{Arg}z'}} = \frac{1}{e^{\ln f'} - 1}, \quad (118)$$

which can be rewritten as

$$\frac{\partial}{\partial(1-f')} \ln \frac{e^{i\text{Arg}z'}}{1 - e^{i\text{Arg}z'}} = \frac{-\sum_{\gamma=0}^{\infty} e^{-i\text{Arg}z'\gamma} \frac{i\text{Arg}z'\gamma}{\ln f'}}{-\sum_{\gamma=0}^{\infty} e^{-i\text{Arg}z'\gamma}}, \quad (119)$$

with

$$i\text{Arg}z' = \ln \frac{z'}{|z'|} = \ln \frac{f' - 1}{f' \ln f'} = \frac{f'}{f' - 1} \frac{(f' - 1) - f' \ln f'}{(f' \ln f')^2}, \quad (120)$$

where two  $(1 - f')$ -independent terms can be extracted from the last line of Eq.(118),

$$\begin{aligned}\alpha &= \frac{1 - f'}{1 - e^{i\text{Arg}z'}}, \\ \beta &= \frac{f'}{(1 - f')^2} (1 - e^{-i\text{Arg}z'}),\end{aligned}\quad (121)$$

where  $\alpha$  is the derivative of the logarithm of the summation over microscopic states while  $\beta$  is similar to the inverse temperature in a system satisfies the Bose-Einstein distribution. While the result of Eq.(118) indeed estimating if the step length has being stretched or not, in terms of the  $(-1)$  in scaled form  $\exp[\ln(-1) - \ln(1 - f')]$ . Also, it is related to the entropy and randomness of the targets with large amount in this system.

Note that Eq.(109) is valid in both the cases where  $\ln(-1)$  plays role or not, i.e., it can coexist and independent with the  $\ln(-1)$  term. From Eq.(109), we can obtain the following expression,

$$\lim_{(1-f') \rightarrow \infty} \ln \ln f' = \ln \ln f' \frac{f' \ln f'}{f' \ln f' + 1}, \quad (122)$$

whose  $(1 - f')$ -independence leads to

$$\frac{\partial}{\partial(1-f')} \frac{f' \ln f'}{f' \ln f' + 1} = \frac{1}{f' \ln f' + 1} \frac{1}{\ln \ln f'}. \quad (123)$$

Then we have

$$f' \ln f' \frac{\partial}{\partial(1-f')} \ln \frac{f' \ln f'}{f' \ln f' + 1} = f' \ln f' \frac{\partial}{\partial(1-f')} \left( \frac{1}{1 - (\frac{-1}{f' \ln f'})} \left[ \frac{\partial}{\partial(\frac{-1}{f' \ln f'})} \right]^{-1} \right) = \frac{1}{\ln \ln f'}. \quad (124)$$

As discussed before, as long as here  $\frac{\partial}{\partial(1-f')}$  has not being isolated, the above expression always equivalent to

$$f' \ln f' \frac{\partial}{\partial(1-f')} \left[ \frac{\partial}{\partial(\frac{-1}{f' \ln f'})} \right]^{-1} = \frac{1}{\ln \ln f'}, \quad (125)$$

which is guaranteed by the following expression,

$$-f' \frac{\partial}{\partial(1-f')} \left[ \frac{\partial}{\partial(\frac{-1}{f' \ln f'})} \right]^{-1} = -f' \frac{\partial}{\partial(1-f')} \text{Li}_1\left(\frac{-1}{f' \ln f'}\right) = \frac{-1}{(\ln f') \ln(\ln f')}, \quad (126)$$

which is the inversed entropy for the system distribution (or degree of stretching) described by  $\ln f'$ . Then by substituting this into Eq.(124), we obtain

$$\left[ \frac{\partial}{\partial(\frac{-1}{f' \ln f'})} \right]^{-1} = \frac{1}{f' \ln f'}. \quad (127)$$

As in this case the  $\ln(-1)$  could be nonzero, and  $e^{i\text{Arg}z'} \neq 1$  (whose derivative with respect to  $(1-f')$  is zero). Thus we can regard  $\left[ \frac{\partial}{\partial(\frac{-1}{f' \ln f'})} \right]^{-1}$  as a Dirac delta function  $\delta(1-f')$ , and  $f' \ln f'$  as the text function. Then from Eq.(124), we have

$$\begin{aligned} & - \int d(1-f') f' \ln f' \frac{\partial}{\partial(1-f')} \delta(1-f') = \int d(1-f') \delta(1-f') \frac{\partial}{\partial(1-f')} (f' \ln f') \\ & = \int d(1-f') \delta(1-f') \frac{f' \ln f'}{1-f'} + \int d(1-f') \delta(1-f') (1-f') \frac{\partial}{\partial(1-f')} \frac{f' \ln f'}{1-f'} = \frac{-1}{\ln \ln f'}. \end{aligned} \quad (128)$$

As this expression is valid in both the  $\ln(-1) = 0$  and  $\ln(-1) \neq 0$  cases. For the first case, with  $e^{i\text{Arg}z'}$  can be trivially treated as 1, the operator  $\left[ \frac{\partial}{\partial(\frac{-1}{f' \ln f'})} \right]^{-1}$  follows exactly the property of Dirac delta function, and we have

$$- \lim_{\ln(-1) \rightarrow 0} \int d(1-f') f' \ln f' \frac{\partial}{\partial(1-f')} \delta(1-f') = \lim_{\ln(-1) \rightarrow 0} \int d(1-f') \frac{\partial}{\partial(1-f')} \ln(f' \ln f') = \frac{1}{1-f'}, \quad (129)$$

in which case

$$(1-f') \frac{\partial}{\partial(1-f')} \delta(1-f') = -\delta(1-f') = \frac{-1}{f' \ln} f'. \quad (130)$$

For the second case with  $e^{i\text{Arg}z'} \neq 1$ , we have

$$\begin{aligned} \frac{-1}{\ln \ln f'} &= \int d(1-f') \left[ \frac{1}{1-f'} + \frac{\partial}{\partial(1-f')} \ln(-1) \right], \\ \int d(1-f') \delta(1-f') \frac{f' \ln f'}{1-f'} &= \int d(1-f') \frac{e^{i\text{Arg}z'}}{1-f'}. \end{aligned} \quad (131)$$

#### 4.7 Summary for the algorithm part

One of the most salient feature of this algorithm is the different form of the limiting result due to the stretched step length, that is, if we define  $f_\infty = \lim_{k \rightarrow \infty} f_k$ ,  $F_k = f_\infty f_k$ , it has

$$F_\infty := \lim_{k \rightarrow \infty} F_k = \frac{f_\infty f_k}{1 - f_\infty \partial_k f_k}, \quad (132)$$

instead of the traditional result with fix IR cutoff throughout all the target samples of the system  $\frac{f_\infty f_k}{1 - \partial_k f_k}$ . Our criterion that the correlations between arbitrary two quantities is only

determined by the variable-dependence. It is easy to verified that the Eq.(132) shows the same  $k$ -independence with  $\lim_{k \rightarrow \infty} f_k$ , and as we discuss above, as long as  $k' < k$ , we always have

$$\begin{aligned}\partial_{k'} f_\infty &= \partial_k f_\infty = 0, \\ \partial_{k'} F_\infty &= \partial_k F_\infty = 0,\end{aligned}\tag{133}$$

but for  $k' > k$ ,

$$\begin{aligned}\partial_{k'} f_\infty &= \frac{(\partial_k f_k)^2}{\partial_k f_k - 1}, \\ \lim_{k' \rightarrow \infty} F_k &= \frac{F_k}{1 - (\partial_{k'} f_k) f_\infty - \frac{f_k (\partial_k f_k)^2}{\partial_k f_k - 1}} = \frac{F_k}{1 - \partial_k f_k f_\infty [-(1 + \partial_k f_k)]}.\end{aligned}\tag{134}$$

We can write the derivative of Eq.(132) as

$$\partial_k \left[ \ln \frac{f_\infty f_k e^{F_\infty}}{1 - f_\infty \partial_k f_k} \right] = 0,\tag{135}$$

which can be connected to Eq.(115) by setting  $f_\infty f_k e^{i \text{Arg} z'} = (1 - f')$ ,  $\frac{-1}{f' \ln f'} = \frac{\partial_k f_k}{f_k} e^{-i \text{Arg} z'}$ . As all these factors in Eq.(115) are divided from the exponential factor  $e^{i \text{Arg} z'}$ , there is not the stretched step length. In this case, combined with Eq.(100), we can see as long as the step length is not be stretched (in which case the  $\ln(-1)$  terms plays no role),

$$\begin{aligned}\lim_{f' \rightarrow 1^-} \ln \frac{-1}{f' \ln f'} &= -\ln(1 - f'), \\ \lim_{f' \rightarrow 1^-} \ln \frac{e^{i \text{Arg} z'}}{1 - e^{i \text{Arg} z'}} &= \frac{1}{1 - e^{i \text{Arg} z'}} \left( \frac{\partial}{\partial e^{i \text{Arg} z'}} \right)^{-1},\end{aligned}\tag{136}$$

where the limiting scales becomes a marginal one, instead of the combined one ( $f' \rightarrow 1^+/1^-$ ) for the case where  $\ln(-1)$  plays a role, as we introduced above.

## 5 Conclusion

We reveal the relation between UV cutoff of polaronic momentum  $\Lambda_q$  and its SYK behavior. The SYK behavior of a polaron system, as well as the relation between scattering momentum and the related statistical behaviors has rarely been investigated before. By projecting to a 2d square lattice, The algorithm designed by us allow the automatically optimization and prediction for the resulting spectrum. In this 2d lattice model with position space representation, the emergent nonzero expectation value of site potential difference  $\Delta_{\alpha\beta}$  shows the existence of polaron. For eigenvalues arranged in ascending order in the position space, the learning program devotes to predict the weight distribution of each eigenvalue in the final spectrum, which is realized by modifying the relative distance between each pair of the two elements (groups) in both sides of the selected positions. Indeed this is equivalent to modifying the step length for each position, i.e., the IR cutoff in the position space, but this cutoff is no longer a invariant constant here. One of the obvious advance of this is the much easier estimation of the derivative of potential difference in each position, which is directly related to the variance as well as mean value of the final spectrum. Also, the large distance in the lattice position space (like the scaled variable  $\chi$ ) corresponds to the long wavelength limit in the momentum space.

In momentum space, as the polaron emerges at the pole of scattering amplitude, the polaronic momentum  $q$ , which is also the relative momentum during scattering between impurity and majority particles (or holes), reads  $q = ia^{-1}$ . The quantity  $qa$  is also an important characteristic scale in predicting its many-body behaviors. As we show in the Appendix, the polaron

system in momentum space with four-fermion interactions, the modified level distribution (from GUE to another one between GUE and GSE) can be described and explained in terms of the additional momentum  $q$ -dependent Gaussian wave functions  $\Phi_{ik}$  and  $\Phi_{jl}$ .

## 6 Appendix: SYK-type polaron behavior in terms of momentum space representation

Similar to Eq.(1), we can illustrating this model in momentum space as, (for fermions with scaling dimension  $1/4$ ),

$$H_{SYK} = \sum_{i<j;k<l} \Phi_{-q}^{ij} \Phi_q^{kl} c_i^\dagger c_k^\dagger c_l c_j = \sum_{i<j;k<l} J_{ik;jl} \Phi_{-q} \Phi_q c_i^\dagger c_k^\dagger c_l c_j, \quad (137)$$

where  $i, j, k, l = 1 \cdots N$ .  $J_{ijkl}$  is an antisymmetry tensor which follows a Gaussian distribution with zero mean. To adding the DOF related to scattering momenta, we introduce the following  $s$ -wave operators which can expressed in terms of the eigenfunctions of SYK Hamiltonian

$$\begin{aligned} \Phi_{-q}^{ij} &= \Phi_{-q} c_i^\dagger c_j, \\ \Phi_q^{kl} &= \Phi_q c_k^\dagger c_l. \end{aligned} \quad (138)$$

Here the Gaussian variables  $(i, k)$  and  $(j, l)$  with, respectively, the scattering momenta  $-q$  and  $q$ , correspond to the  $i$  and  $j$  in Eq.(1), and the polaronic momentum wavefunction  $\phi_{-q} \phi_q$  related to the potential difference-DOF in Eq.(1). The coupling satisfies

$$\delta_{ik} \delta_{jl} J_{i<j;k<l} = - \sum_q^{\Lambda_q} \Phi_{ij}^{-q} g_q \Phi_{kl}^q. \quad (139)$$

We restrict that, projecting to momentum space, the flavor number of the third DOF, which is the  $O(M) \sim \frac{N^2}{4}$  in Eq.(1). As the variance  $\sigma_\Delta^2 \sim \frac{N^2}{4} = \frac{g_{(3)}^2}{3N^2} \frac{3N^2}{2} = \frac{g_{(3)}^2}{2}$ , the inverse momentum cutoff reads  $\Lambda_q^{-1} = \frac{2J}{N^2}$ , where  $g_{(3)}^2 = \frac{N^2}{2}$  is the constant parameter as stated in Sec.3, and  $J^2$  is the corresponding coupling in momentum space but also in dimensions of energy. This result is in consistent with the final variance in Eq.(148).

As we stated in above, the calculations related to the polaron dynamics usually requires momentum cutoff  $\Lambda_q$ . The polaron coupling reads (with  $E_b$  the binding energy and  $W$  the bandwidth)

$$g_q^{-1} = - \sum_{kp}^{\Lambda} \frac{1}{E_b + \varepsilon_p + \varepsilon_k + W}, \quad (140)$$

which vanishes in  $\Lambda \rightarrow \infty$  limit. Similarly, in two space dimension, the polaron corresponds to the pole  $q = ia^{-1}$  where  $a$  is the scattering length (or scattering amplitude), which proportional to the polaronic coupling strength, and the strongest polaronic coupling realized at  $q \sim a^{-1}$  while the weakest coupling realized at  $qa \ll 1$ . This is a special property of polaron formation and is important during the following analysis. Now we know that  $g_q$  is inversely proportional to the value of polaronic exchanging momentum  $q$ , then by further remove the  $q$ -dependence of polaronic coupling

$$g^{-1} = - \sum_q^{\Lambda_q} \sum_{kp}^{\Lambda} \frac{1}{E_b + \varepsilon_{p-q} + \varepsilon_{k+q} + W}, \quad (141)$$

the integral in Eq.(141) is vanishingly small when  $\Lambda_q \rightarrow \infty$ . Note that in the following we may still use  $g_q$  to denote the polaronic coupling to distinct it from the SYK one.

In opposite limit, when  $\Lambda_q \rightarrow 0$ , both the couplings  $J_{ijkl}$  and  $g_q$  become very strong (thus enters the SYK regime). Similar to the disorder effect from temperature (which is lower than the coherence scale but higher than other low energy cutoff) to Fermi liquid, the fermion frequency can be treated as a disorder to non-Fermi liquid SYK physics, (the pure SYK regime can be realized in  $\omega, \omega_c \rightarrow 0$  limit and extended to zero temperature) thus we can write the essential range of parameter to realize SYK physics,

$$\Lambda_q^{-1} \sim \frac{g_q}{\omega} v_F \gg N, \quad (142)$$

this is one of the most important results of this paper which relates the polaron physics to the SYK physics, and in the mean time, it is surely important to keep  $N \gg U \gg \omega \gg \omega_c \gg U/N$ . Here the  $\omega \sim v_F q$  plays the role of disorder in frequency space. Note that here vanishingly small cutoff in momentum space  $\Lambda_q$  corresponds to vanishing spatial disorder  $\Lambda_r$  which is the lattice spacing in two-dimensional lattice in real space[14]. That is, in the presence of short range interaction, by reducing the distance between two lattice sites (and thus enlarging the size of hole), the size number as well as the coupling is increased. Thus in this case the polaron term becomes asymptotically Gaussian distributed due to the virtue of the central limit theorem.

In the  $\Lambda_q = 0$  limit (SYK), the Hamiltonian can be rewritten in the exact form of SYK $_{q=2}$  mode.

$$\begin{aligned} H_p &= g_q \sum_{ik} \sum_{\sigma, \sigma'} \Phi_i c_i^\dagger c_i \Phi_k c_{k\sigma'}^\dagger c_k \\ &\equiv g_q \sum_{ik} \sum_{\sigma, \sigma'} b_i b_{k\sigma'}, \end{aligned} \quad (143)$$

where  $b_i = \Phi_i c_i^\dagger c_i$  and  $[b_i, H_p] = 0$ . After disorder average in Gaussian unitary ensemble (GUE), for Gaussian variable  $\Phi_{ik} \Phi_{jl}$ , we have

$$\overline{\Phi_{ik} \Phi_{jl}} = 0, \quad (144)$$

and the replication process reads

$$g_q \Phi_{il} \Phi_{jk} \mathcal{O}_{il,jk} \rightarrow -\frac{J^2 \delta_{ij} \delta_{jk} 2\mathcal{O}_{il,jk}^\dagger \mathcal{O}_{il,jk}}{16N^2} = -\frac{J^2 \delta_{ij} \delta_{jk} \mathcal{O}_{il,jk}^\dagger \mathcal{O}_{il,jk}}{8N^2}. \quad (145)$$

Note that before replication, the number of observable  $\mathcal{O}$  should equal to the number of Gaussian variables, which is one in the above formula. While in the finite but small  $\Lambda_q$  case

$$H_p = g_q \sum_{il,jk} \sum_{\sigma, \sigma'} \Phi_{il}^{-q} \Phi_{jk}^q c_i^\dagger c_l c_j^\dagger c_k, \quad (146)$$

where we have

$$\begin{aligned} g_q \Phi_{il}^{-q} \Phi_{jk}^q \mathcal{O}_{ik}^\dagger \mathcal{O}_{lj} 2\mathcal{O}_q^\dagger \mathcal{O}_q &\rightarrow \frac{J_{il,jk}^2 \delta_{il} \delta_{jk} \Lambda_q^2 4\mathcal{O}_{ik} \mathcal{O}_{ik}^\dagger \mathcal{O}_{lj}^\dagger \mathcal{O}_{lj} (2\mathcal{O}_q^\dagger \mathcal{O}_q)^2}{16N^2} \\ &= \frac{J_{il,jk}^2 \delta_{il} \delta_{jk} \Lambda_q^2 \mathcal{O}_{ik} \mathcal{O}_{ik}^\dagger \mathcal{O}_{lj}^\dagger \mathcal{O}_{lj} (\mathcal{O}_q^\dagger \mathcal{O}_q)^2}{N^2}. \end{aligned} \quad (147)$$

Note that each operator  $\mathcal{O}$  must contain  $q$  completely independent (uncorrelated) indices, and before replication, the indices of each operator  $\mathcal{O}$  must not be completely the same. For example, in the finite (although small)  $\Lambda_q$  case,  $i$  ( $l$ ) is completely independent of  $k$  ( $j$ ), but index  $ik$  is not completely uncorrelated with  $lj$  because mapping to their momentum space they have  $i - l = k - j$  due to the fixed polaronic momentum  $q$ , in other words, the mechanism that transforms  $i$  to  $l$  is the same with that to transform  $k$  to  $j$ , thus  $ik$  can be continuously mapped to  $lj$ .

In case of finite (but small)  $\Lambda_q$  with approximately uncorrelated random Gaussian variables  $\Phi_{ik}$  and  $\Phi_{jl}$  ( $\overline{\Phi_{ik}} = \overline{\Phi_{jl}} = 0$ ,  $g_q \overline{\Phi_{ik}^2} = g_q \overline{\Phi_{jl}^2} = \frac{J}{2N} 2\Lambda_q = \frac{J}{N} \Lambda_q$ ), we can perform the disorder averages over fermion indices and the  $q$  in the same time, which leads to the following mean value and variance of Chi-square random variable  $g_q \Phi_{il}^{-q} \Phi_{jk}^q$

$$\begin{aligned} \overline{g_q \Phi_{il}^{-q} \Phi_{jk}^q} &= g_q \overline{\Phi_{il}^{-q} \Phi_{jk}^q} = 0, \\ \overline{(g_q \Phi_{il}^{-q} \Phi_{jk}^q)^2} &= g_q^2 \overline{(\Phi_{il}^{-q})^2 (\Phi_{jk}^q)^2} \\ &= \frac{J^2 4\Lambda_q^2}{4N^2} \\ &= \frac{J^2 \Lambda_q^2}{N^2}. \end{aligned} \quad (148)$$

The second line is valid because  $(\Phi_{il}^{-q} \Phi_{jk}^q)^2 = (\Phi_{il}^{-q})^2 (\Phi_{jk}^q)^2$  when  $\Phi_{il}^{-q}$  is independent of  $\Phi_{jk}^q$ , and we assume the variance of  $\Phi^{-q} \Phi^q$  is zero throughout this paper. Note that this is only valid in the case that the disorder average over  $q$  fermion indices are done separately, i.e., the degree of freedom of  $q$  will not affect the correlation between  $\Phi_{il}$  and  $\Phi_{jk}$ , and vice versa. Besides,  $q$  must be integrated in the same dimension of  $N$ , i.e., one dimension (which is the case we focus on in this paper), and thus  $\Lambda_q$  is in the same scale with  $N$ . If the  $q$ -integral is carried out in  $d$ -space dimension, then the above result becomes

$$\overline{(g_q \Phi_{il}^{-q} \Phi_{jk}^q)^2} = \frac{J^2 \Lambda_q^{2d}}{N^2}. \quad (149)$$

because the sample number of  $q$  is related to spacial dimension  $d$ .

Next we take the spin degree of freedom into account. To understand the effects of perturbation brought by finite small  $q$  (where the polaronic coupling is still approximately viewed as a constant), we use the SU(2) basis to deal with the degree of freedom of spin (i.e., of impurity and majority particles),  $\Phi_{\sigma=\pm} = \frac{1}{\sqrt{2}}(\Phi_1 \pm i\Phi_2)$ . Then we have

$$\begin{aligned} &g_q \sum_{\sigma, \sigma'=\pm} \sum_{il;jk} \sum_q^{\Lambda_q} \Phi_{il}^{-q} \Phi_{jk}^q \\ &= g_q \sum_q^{\Lambda_q} \sum_{il;jk} \left( \frac{1}{2} (\Phi_{il1}^{-q} + i\Phi_{il2}^{-q}) (\Phi_{jk1}^q - i\Phi_{jk2}^q) - \frac{1}{2} (\Phi_{il1}^{-q} - i\Phi_{il2}^{-q}) (\Phi_{jk1}^q + i\Phi_{jk2}^q) \right) \\ &= g_q \sum_q^{\Lambda_q} \sum_{il;jk} (i\Phi_{il2}^{-q} \Phi_{jk1}^q - i\Phi_{il1}^{-q} \Phi_{jk2}^q). \end{aligned} \quad (150)$$

And we obtain the variance

$$\begin{aligned} g_q \Phi_{il1}^{-q} \Phi_{jk1}^q 2\mathcal{O}_{il}^\dagger \mathcal{O}_{il} &= g_q \Phi_{il2}^{-q} \Phi_{jk2}^q 2\mathcal{O}_{il}^\dagger \mathcal{O}_{il} = \frac{J}{4N} \delta_{il,jk} 2\Lambda_q 2\mathcal{O}_{il}^\dagger \mathcal{O}_{il}, \\ \overline{g_q \Phi_{il1}^{-q} \Phi_{jk1}^q} &= \overline{g_q \Phi_{il2}^{-q} \Phi_{jk2}^q} = \frac{J\Lambda_q}{N} \delta_{il,jk}, \\ \overline{\Phi_{il1}^{-q} \Phi_{jk2}^q} &= 0, \end{aligned} \quad (151)$$

where  $J > 0$ . Thus  $\Phi_{il1}^{-q}$  is orthogonal with  $\Phi_{jk2}^q$  (as long as the  $\Phi_{ik}$  is approximately treated as independent of  $\Phi_{jl}$  in the small  $\Lambda_q$  limit (e.g., the SYK limit)). Combined with Eq.(148), we know the variance  $\text{Var}(g_q \Phi_{il1}^{-q} \Phi_{il1}^q) = \text{Var}(g_q \Phi_{jk2}^{-q} \Phi_{jk2}^q) = 0$ , which is different to the result of next section.

In the small (but finite)  $\Lambda_q$  limit, according to semicircle law, the spectral function of single fermion reads

$$\rho(\varepsilon) = \frac{1}{\pi J} \sqrt{1 - \frac{\varepsilon^2}{4J^2}} = \frac{1}{2\pi J^2} \sqrt{4J^2 - \varepsilon^2}, \quad (152)$$

with

$$J^2 = \overline{(g_q \Phi_{1il}^{-q} \Phi_{2jk}^q)^2} \frac{N^2}{\Lambda_q^2}. \quad (153)$$

The mean value of eigenvalues is thus

$$\left| \int_{\varepsilon < 0} d\varepsilon \varepsilon \rho(\varepsilon) \right| \approx \frac{4J}{3\pi}. \quad (154)$$

Then we obtain that the matrices  $g_q \Phi_{1il}^{-q} \Phi_{2jk}^q$  and  $\Phi_{1il}^{-q}$  and  $\Phi_{2jk}^q$  are  $\frac{N^2}{\Lambda_q} \times \frac{N^2}{\Lambda_q}$  (a  $\infty \times \infty$ ) matrix, and now these matrices are automatically diagonalized. In such a configuration constructed by us,  $g_q \Phi_{1il}^{-q} \Phi_{2jk}^q$  can not be simply viewed as a product of matrices  $g_q \Phi_{il}^{-q}$  and  $g_q \Phi_{jk}^q$ , since  $g_q \Phi_{il}^{-q} \Phi_{jk}^q$  is a  $N^2 \times N^2$  matrix while  $g_q \Phi_{il}^{-q} \Phi_{jk}^q$  is a  $\Lambda_q^{-1} \times \Lambda_q^{-1}$  diagonal matrix ( $q \neq 0$  here). Instead, it requires mapping  $il(jk) \rightarrow \frac{il(jk)}{\sqrt{\Lambda_q}}$  and  $q \rightarrow q \frac{1}{N^2}$  (to realizes  $\delta\Lambda'_q = \Lambda_q'^2 = \Lambda_q^2 \frac{1}{N^2}$ ). This is the SYK phase with gapless SYK mode, and it requires  $\Lambda_q^{-1} \gg N$ .

## 6.1 Removable the correlation between $\Phi_{ik}$ and $\Phi_{jl}$ by summing over $q$

The SYK phase can be gapped out due to the broken symmetry by finite eigenvalue of  $(g_q \Phi_{1ik} \Phi_{2jl})^2$  (or  $(g_q \Phi_{il} \Phi_{jk})^2$ ). To understand this, it is more convenient to use another configuration, where we carry out the summation over  $q$  first in Eq.(34), instead of carrying out the disorder averages over  $ijkl$  and  $q$  in the same time. Then the disorder average over fermion indices  $i, j, k, l$  simply results in

$$\overline{(g_q \Phi_{1ik} \Phi_{2jl})^2} = \frac{J_{il;jk}^2}{4N^2}, \quad (155)$$

which can be calculated as

$$\begin{aligned} \overline{(g_q \Phi_{1ik} \Phi_{2jl})^2} &= \overline{\sum_q^{\Lambda_q^{-1}} g_q \Phi_{1il}^{-q} \Phi_{1jk}^q)^2} \\ &= \overline{(g_q \Phi_{1il}^{-\delta\Lambda_q} \Phi_{1jk}^{\delta\Lambda_q} + g_q \Phi_{1il}^{-2\delta\Lambda_q} \Phi_{1jk}^{2\delta\Lambda_q} + \dots)^2} \\ &= \overline{(g_q \Phi_{1il}^{-\delta\Lambda_q} \Phi_{1jk}^{\delta\Lambda_q})^2 + (g_q \Phi_{1il}^{-2\delta\Lambda_q} \Phi_{1jk}^{2\delta\Lambda_q})^2 + \dots} \\ &= \frac{J^2}{4N^2} \sum_q (\Phi^{-q} \Phi^q)^2 \\ &= \frac{J^2 4\Lambda_q}{4N^2} \\ &= \frac{J^2 \Lambda_q}{N^2}, \end{aligned} \quad (156)$$

i.e.,  $J_{il;jk}^2 = 4\Lambda_q J_{il;jk}^2$ . The third line is due to the fact about variance of Gaussian variables:  $\text{Var}(A+B) = \text{Var} A + \text{Var} B$  where  $A$  and  $B$  are independent with each other. The fourth line

is because

$$\begin{aligned}
\text{Var}(\Phi^q) &= \overline{(\Phi^q)^2} - (\overline{\Phi^q})^2 = 2\Lambda_q - 0 = 2\Lambda_q, \\
\text{Var}(\Phi^{-q}\Phi^q) &= \overline{(\Phi^{-q}\Phi^q)^2} - (\overline{\Phi^{-q}\Phi^q})^2 = 4\Lambda_q^2 - (2\Lambda_q)^2 = 0, \\
\sum_q \Phi^{-q}\Phi^q &= \overline{\Phi^{-q}\Phi^q}\Lambda_q^{-1} = 2, \\
\sum_q (\Phi^{-q}\Phi^q)^2 &= \overline{(\Phi^{-q}\Phi^q)^2}\Lambda_q^{-1} = 4\Lambda_q,
\end{aligned} \tag{157}$$

where  $\Phi^{-q}\Phi^q$  is obviously not a Gaussian variable (just like the  $\Phi_{ik}\Phi_{jl}$  except in the SYK limit), and  $\overline{\Phi^{-q}\Phi^q} = \overline{(\Phi^q)^2} \neq \overline{\Phi^{-q}\Phi^q} \neq 0$  because it is impossible to make  $\Phi^{-q}$  and  $\Phi^q$  orthogonal to each other due to the connection between them  $q = -(-q)$ . That is to say, although  $\Phi^{-q}$  and  $\Phi^q$  are Gaussian variables with zero mean  $\overline{\Phi^{-q}} = \overline{\Phi^q} = 0$ , their product  $\Phi^{-q}\Phi^q$  is not a Gaussian variable and do not have zero mean. This is different to the variance of Chi-square variable  $\Phi_{1ik}\Phi_{2jl}$  which is finite due to the zero mean  $\overline{\Phi_{1ik}\Phi_{2jl}} = \overline{\Phi_{1ik}}\overline{\Phi_{2jl}} = 0$ , by treating them to be approximately mutually orthogonal (i.e.,  $\Phi_{ik}$  approximately independent with  $\Phi_{jl}$ ). Here we note that following relations in new configuration

$$\begin{aligned}
\sqrt{|g_q|}\Phi_{1ik} &= \sqrt{g_q}\Phi_{2jl} = 0, \\
\text{Var}(\sqrt{|g_q|}\Phi_{1ik}) &= \overline{g_q\Phi_{1ik}^2} = \frac{J}{2N}2 = \frac{J}{N}, \\
\text{Var}(\sqrt{|g_q|}\Phi_{2jl}) &= \overline{g_q\Phi_{2jl}^2} = \frac{J}{2N}2 = \frac{J}{N}, \\
\text{Var}(g_q\Phi_{1ik}\Phi_{1jk}) &= \overline{(g_q\Phi_{1ik}\Phi_{1jk})^2} - \overline{g_q\Phi_{1ik}\Phi_{1jk}}^2 \\
&= \frac{J^2}{4N^2}4\Lambda_q\delta_{il,jk} - \left(\frac{J}{2N}2\right)^2\delta_{il,jk} \\
&= \frac{J^2}{N^2}(\Lambda_q - 1)\delta_{il,jk}, \\
\overline{g_q\Phi_{1ik}\Phi_{2jl}} &= 0.
\end{aligned} \tag{158}$$

Under this configuration, by approximately treating  $\Phi_{1ik}$  and  $\Phi_{2jl}$  to be mutually orthogonal, they can be viewed as two vectors, and each of them contains  $N^2$  components, then  $(g_q\Phi_{1ik}\Phi_{2jl})$  is a  $N^2 \times N^2$  matrix with complex eigenvalues. But note that, away from the  $\Lambda_q^{-1} \rightarrow \infty$  limit, this construction fail because exactly speaking,  $(g_q\Phi_{1ik}\Phi_{2jl})$  (after summation over  $q$ ) is a  $N \times N \times N = N^3$  matrix (unlike the  $N \times N$  SYK $_{q=2}$  or the  $N \times N \times N \times N = N^4$  SYK $_{q=4}$ ) due to the polaron property, i.e., over the fermion indices  $i, j, k, l$ , one of them is always identified by the other three, so there are at most three independent indices (degrees of freedom).

A precondition to treat Gaussian variables  $i\Phi_{ik}$  and  $\Phi_{jl}$  mutually orthogonal (independent), is that it must away from the  $\Lambda_q^{-1} \rightarrow \infty$  limit, since too small sample number will makes the matrix  $g_q\Phi_{-q}\Phi_q$  leaves away from the Gaussian distribution according to central limit theorem, and thus the disorder average over  $q$  can not be successively carried out, that is why we instead make the summation over  $q$ . Then, the relation  $\Lambda_q^{-1} < N^2/2$  ( $\Lambda_q > 2/N^2$ ) indicates the large number of  $N$ , which preserves the Gaussian distribution of  $\Phi_{ik}$  and  $\Phi_{jl}$  and also makes the disorder average over fermion indices to matrix  $g_q\Phi_{il}\Phi_{jk}$  more reliable, despite the indices  $il$  and  $jk$  are not completely independent but correlated by some certain mechanism before the summation over  $q$ .

Then we turning to the matrix

$$g_q\Phi_{il}^{-q}\Phi_{jk}^q = ig_q\Phi_{1il}^{-q}\Phi_{2jk}^q - ig_q\Phi_{2il}^{-q}\Phi_{1jk}^q, \tag{159}$$

which is also a  $N^2 \times N^2$  matrix now and is Hermitian (whose eigenvectors and eigenvalues are much more easy to be solved) with all diagonal elements be zero. In this scheme, to make sure  $g_q \Phi_{il} \Phi_{jk}$  is a  $N^2 \times N^2$  matrix, the disorder average over  $i, j, k, l$  must be done after the summation over  $q$ . Then to diagonalizing the  $N^2 \times N^2$  matrix  $g_q \Phi_{il}^{-q} \Phi_{jk}^q$ , it requires  $\Lambda_q^{-1} < N^2/2$  to make sure all the vectors  $\Phi_{1ik}$  and  $\Phi_{2jl}$  are orthogonal with each other within the matrix  $g_q \Phi_{il}^{-q} \Phi_{jk}^q$ . This is because there at most exists  $N^2$  vectors can orthogonal with each other in  $N^2$ -dimensional space (formed by  $N^2$ -component vectors). In other word, the propose of this is to make sure vectors  $\Phi_{il}^{-\delta\Lambda_q}, \Phi_{il}^{-2\delta\Lambda_q}, \dots, \Phi_{ik}^{-\Lambda_q}, \Phi_{jk}^{\delta\Lambda_q}, \Phi_{jk}^{2\delta\Lambda_q}, \dots, \Phi_{jl}^{\Lambda_q}$  are orthogonal to each other.

Then there are  $N^2 - 2\Lambda_q^{-1}$  eigenvectors with eigenvalues equal zero (correspond to the ground state), i.e.,

$$\text{Det}[g_q \Phi_{il}^{-q} \Phi_{jk}^q] = 0, \quad (160)$$

and  $2\Lambda_q^{-1}$  eigenvectors  $\Phi_\sigma = \frac{\Phi_1^{-q} \pm i\Phi_2^q}{\sqrt{2}}$  with eigenvalue  $\pm \frac{J}{2\Lambda_q^{-1/2}N}$ . This can be verified by the rule that for Hermitian matrix the eigenvectors corresponding to different eigenvalues are orthogonal to each other,

$$\left(\frac{\Phi_1^{-q} + i\Phi_2^q}{\sqrt{2}}\right)^H \cdot \frac{\Phi_1^{-q} - i\Phi_2^q}{\sqrt{2}} = \frac{1}{2}(\Phi_1^{-q} \cdot \Phi_1^{-q} - i\Phi_1^{-q} \cdot \Phi_2^q - i\Phi_1^{-q} \cdot \Phi_2^q - \Phi_2^q \cdot \Phi_2^q) = 0, \quad (161)$$

where  $\Phi_1^{-q} \cdot \Phi_2^q = \Phi_1^{-q} \cdot \Phi_2^q = 0$  since they are orthogonal to each other, and superscript  $H$  denotes the transpose conjugation (Hermitian conjugate). The result of Eq.(35) is used here.

In the special case of  $\Lambda_q^{-1} = N^2/2$ , we have, in matrix  $g_q \Phi_{il} \Phi_{jk}$ ,  $N^2$  eigenvectors  $\frac{\Phi_1^{-q} \pm i\Phi_2^q}{\sqrt{2}}$  with eigenvalue  $\pm \frac{J\Lambda_q}{2\sqrt{2}}$ . Then processing the disorder average over  $ijkl$  to  $g_q \Phi_\sigma^{-q} \Phi_\sigma^q$ , we have the variance

$$\text{Var}(g_q \Phi_{il}^{-q} \Phi_{jk}^q) = \overline{(g_q \Phi_{il}^{-q} \Phi_{jk}^q)^2} = \overline{(ig_q \Phi_{1il}^{-q} \Phi_{2jk}^q - ig_q \Phi_{2il}^{-q} \Phi_{1jk}^q)^2} = \frac{1}{4} \frac{J^2 \Lambda_q}{N^2} = \frac{J^2}{2N^4}, \quad (162)$$

since  $\overline{g_q \Phi_{il}^{-q} \Phi_{jk}^q} = 0$ . The factor  $1/4$  origin from the spin degrees of freedom, and can be verified by the square of eigenvalue  $\lambda^2$ . Note that here the overline denotes only the disorder average over  $ijkl$  index. This result is in consistent with the property of Wigner matrix in GUE

$$\overline{\lambda^2} = \overline{(g_q \Phi_{il}^{-q} \Phi_{jk}^q)^2} = \frac{J^2 \Lambda_q}{4N^2} \sim O(N^{-2}), \quad (163)$$

in contrast with that in Gaussian orthogonal ensemble (GOE) which reads  $O(N^{-2})(1 + \delta_{il,jk})$ . Here  $\lambda$  denote the eigenvalues. The GUE with  $\sum \lambda^2 = \frac{J^2 \Lambda_q}{4}$  thus corresponds to the SYK non-Fermi liquid case, with continuous distributed peaks in the SYK fermion spectral function, i.e., the level statistics agree with the GUE distribution, and the set of eigenvalues follow an ascending order. In GUE, we also have the relation

$$\langle \mathcal{O}_{il} \mathcal{O}_{jk}^\dagger \rangle = \langle c_i^\dagger c_l c_j^\dagger c_k \rangle = \langle c_i^\dagger c_j^\dagger c_k c_l \rangle = \langle c_i^\dagger c_l \rangle \langle c_j^\dagger c_k \rangle - \langle c_i^\dagger c_k \rangle \langle c_j^\dagger c_l \rangle, \quad (164)$$

at zero temperature. While the GOE correponds to the case of nonzero pairing order parameter (in which case pair condensation happen at temperature lower than the critical one), and thus admit the anomalous terms. In GOE we have

$$\overline{\lambda^2} = \overline{(g_q \Phi_{il}^{-q} \Phi_{jk}^q)^2} = \frac{J^2 \Lambda_q}{4N^2} + \frac{O(\delta_{il,jk})}{N^2}, \quad (165)$$

$$\sum \lambda^2 = \frac{J^2 \Lambda_q}{4} + O(\delta_{il,jk}).$$

Thus the GOE has a level repulsion slightly larger than that of GUE in the small level spacing limit during the level statistic. Then if we turn to the many-body localized phase where the

thermalization (chaotic) is being suppressed by the stronger disorder, the level statistic follows the Poisson distribution. Since  $g\Phi_{il}\Phi_{jk}$  is not a positive-define matrix, the largest eigenvalues splitting happen which corresponds to the discrete spectrum with the level statistics agree with Poisson distribution, i.e., it has the largest eigenvalues  $\lambda_{max} = \pm \frac{J\Lambda_q^{1/2}}{2\sqrt{2}}$  and  $\frac{N^2-2}{2}$  eigenvalues  $\frac{1}{N^2-2}$  and  $\frac{N^2-2}{2}$  eigenvalues  $\frac{-1}{N^2-2}$ . Such a distribution of eigenvalues implies the existence of off-diagonal long range order. When the pair condensation happen, the above relation becomes  $\langle c_i^\dagger c_j^\dagger \rangle \langle c_k c_l \rangle$ , with the pairing order parameter

$$\Delta_0 = \langle c_k c_l \rangle = \frac{\sum_{kl} \sum_{\sigma\sigma'} c_k c_l}{2N^{1/2}} \quad (166)$$

where the factor  $(2N^{1/2})^{-1}$  origin from the result of disorder average

$$\begin{aligned} \overline{\Phi_{kl}^2} &= \frac{J}{2N} 2 = \frac{J}{N}, \\ \overline{\Phi_{kl,\sigma\sigma'}^2} &= \frac{J}{4N}. \end{aligned} \quad (167)$$

The  $N^2 \times N^2$  positive-define matrix  $\langle c_i^\dagger c_j^\dagger \rangle \langle c_k c_l \rangle$  has summation of eigenvalues corresponds to the total number of pairs  $\sum \lambda \leq N^2$  and thus  $\bar{\lambda} \leq 1$ , i.e.,  $\lambda_{max} = N^2 = \text{Sp} c_i^\dagger c_j^\dagger \langle c_k c_l \rangle \equiv \text{Sp} b_{ij}^\dagger b_{kl}$ . We also found that, once the boson-fermion interacting term is taken into account, the maximum eigenvalue reduced to: For  $ij \neq kl$ ,  $\lambda_{max} = \text{Sp} b_{ij}^\dagger c^\dagger c b_{kl} \leq \text{Sp}(b_{ij}^\dagger c^\dagger c b_{kl} + c b_{kl} b_{ij}^\dagger c^\dagger) = 0$ ; For  $ij = kl$  ( $\Lambda_q = 0$ ),  $\lambda_{max} = \text{Sp} b_{ij}^\dagger c^\dagger c b_{kl} \leq \text{Sp}(b_{ij}^\dagger c^\dagger c b_{kl} + c b_{kl} b_{ij}^\dagger c^\dagger) = \text{Sp} c c^\dagger = \text{Sp}(1 - c^\dagger c) = 1 - \text{Sp} c^\dagger c \leq 1$ . The superconductivity emerge when  $\Delta_0$  condenses, and in large -N limit, the renormalized Green's function reads

$$G(i\omega)' = \frac{G(i\omega)}{1 + J^2 |\Delta_0^2| G^2(i\omega)}. \quad (168)$$

For a further study about this renormalization effect, see Ref.[27, 24].

In this case, the coupling  $J$  within spectral function reads

$$J^2 = 2N^4 \overline{(g_q \Phi_{il}^{-q} \Phi_{jk}^q)^2}, \quad (169)$$

In the  $\Lambda_q^{-1} \ll N^2/2$  limit, we can easily know that  $J$  is vanishingly small, and the polaronic dynamic then dominates over the SYK dynamic, and the system exhibits Fermi liquid feature. While for  $0 \ll \Lambda_q^{-1} < N^2/2$ , the system exhibits disordered Fermi liquid feature with sharp Landau quasiparticles, and for positive define matrix  $g_q \Phi_{il} \Phi_{jk}$ , since every zero eigenvalue corresponds to a ground state, there are  $N^2 - 2\Lambda_q^{-1}$  ground states, and thus the system exhibits degeneracy  $2^{N^2 - 2\Lambda_q^{-1}}$ . While in the case of  $\mu \gg g_q$ , the bilinear term as a disorder will gap out the system and lift the degeneracy in ground state, although in some certain systems[22] the near nesting of Fermi surface sheets can prevent the increase of degeneracy by disorder. Here the bilinear term is absent but the finite value of variance  $\text{Var}(g_q \Phi_{il} \Phi_{jk})$  with  $\Lambda_q^{-1} < N^2/2$  plays its role and drives the SYK non-Fermi liquid state toward the disordered Fermi liquid ground state.

Finally, we conclude that in  $\Lambda_q^{-1} \ll N^2/2$  case, although  $g_q \Phi_{il} \Phi_{jk}$  is a Hermitian matrix with randomly independent elements and large  $N$ , and each of its matrix elements follows the same distribution (distribution of Chi-square variables), the eigenvalue distribution does not follows the semicircle law. This is because, for  $\Lambda_q^{-1} < N^2/2$ ,  $N^2$ -component vectors  $\Phi_1^q, \Phi_2^q$  are mutually orthogonal, i.e.,  $\Phi_1^q \cdot \Phi_2^{q'} = \Phi_2^q \cdot \Phi_1^{q'} = \Phi_1^q \cdot \Phi_1^{q'} = \Phi_2^q \cdot \Phi_2^{q'} = \delta_{q,q'}$ , which leads to large degeneracy in ground state. In this case, the spectral function does not follows the semicircle law, but exhibits three broadened peaks locate on the energies  $\varepsilon = 0, \pm \frac{J\Lambda_q^{1/2}}{2N}$ , with heights correspond to the numbers of the corresponding eigenvectors.

During the above basis transformation between the original polaron momentum basis and the SYK fermion indices basis, in the  $\Lambda_q^{-1} \rightarrow \infty$  limit, only the statistical relations between  $\phi_i$  and  $\phi_l$  or  $\phi_j$  and  $\phi_k$  depends on the value of  $\Lambda_q^{-1}$ , but this dependence on  $\Lambda_q^{-1}$  is also being replaced by number  $N$  after the transformation. While the disorder average over  $q$  is carried out separately with that of  $N$ , which is allowed only under the condition  $\Lambda_q^{-1} \gg N^2$ , and in this case, the SYK physics cannot be realized if we do the summation over  $q$  first which requires  $\Lambda_q^{-1} < N^2/2$  to realize SYK physics.

## References

- [1] Wu C H. Statistic behaviors of gauge-invariance-dominated 1D chiral current random model[J]. arXiv preprint arXiv:2207.07821, 2022.
- [2] Alexandrov A S. Transition from Fermi liquid to charged Bose liquid: A possible explanation of the isotope shift in high-T c oxides[J]. Physical Review B, 1992, 46(22): 14932.
- [3] Gu Y, Qi X L, Stanford D. Local criticality, diffusion and chaos in generalized Sachdev-Ye-Kitaev models[J]. Journal of High Energy Physics, 2017, 2017(5): 1-37.
- [4] Kohstall C, Zaccanti M, Jag M, et al. Metastability and coherence of repulsive polarons in a strongly interacting Fermi mixture[J]. Nature, 2012, 485(7400): 615-618.
- [5] Chowdhury D, Werman Y, Berg E, et al. Translationally invariant non-fermi-liquid metals with critical fermi surfaces: Solvable models[J]. Physical Review X, 2018, 8(3): 031024.
- [6] Culcer D. Linear response theory of interacting topological insulators[J]. Physical Review B, 2011, 84(23): 235411.
- [7] Sears V F, Svensson E C, Martel P, et al. Neutron-Scattering Determination of the Momentum Distribution and the Condensate Fraction in Liquid He 4[J]. Physical Review Letters, 1982, 49(4): 279.
- [8] Schick M. Two-dimensional system of hard-core bosons[J]. Physical Review A, 1971, 3(3): 1067.
- [9] Field B, Levinsen J, Parish M M. Fate of the Bose polaron at finite temperature[J]. Physical Review A, 2020, 101(1): 013623.
- [10] Christensen R S, Levinsen J, Bruun G M. Quasiparticle properties of a mobile impurity in a Bose-Einstein condensate[J]. Physical review letters, 2015, 115(16): 160401.
- [11] Allen S, Tremblay A M S. Nonperturbative approach to the attractive Hubbard model[J]. Physical Review B, 2001, 64(7): 075115.
- [12] Sedrakyan T A, Efetov K B. Supersymmetry method for interacting chaotic and disordered systems: The Sachdev-Ye-Kitaev model[J]. Physical Review B, 2020, 102(7): 075146.
- [13] Gu Y, Kitaev A, Sachdev S, et al. Notes on the complex Sachdev-Ye-Kitaev model[J]. Journal of High Energy Physics, 2020, 2020(2): 1-74.
- [14] Lantagne-Hurtubise é, Li C, Franz M. Family of Sachdev-Ye-Kitaev models motivated by experimental considerations[J]. Physical Review B, 2018, 97(23): 235124.
- [15] Pikulin D I, Franz M. Black hole on a chip: proposal for a physical realization of the Sachdev-Ye-Kitaev model in a solid-state system[J]. Physical Review X, 2017, 7(3): 031006.
- [16] Sachdev S. Bekenstein-Hawking entropy and strange metals[J]. Physical Review X, 2015, 5(4): 041025.
- [17] Butera D, Sarma S D. Coupled electron-impurity and electron-phonon systems as trivial non-Fermi liquids[J]. Physical Review B, 2019, 100(23): 235149.
- [18] Mahajan R, Barkeshli M, Hartnoll S A. Non-Fermi liquids and the Wiedemann-Franz law[J]. Physical Review B, 2013, 88(12): 125107.

- [19] Cha P, Patel A A, Gull E, et al. Slope invariant T-linear resistivity from local self-energy[J]. *Physical Review Research*, 2020, 2(3): 033434.
- [20] Xu C. Gapless bosonic excitation without symmetry breaking: An algebraic spin liquid with soft gravitons[J]. *Physical Review B*, 2006, 74(22): 224433.
- [21] Chen-Huan WU. SYK behaviors generated by on-site Hubbard-type four-fermion on-site random interactions and the related phase transitions. arXiv: 2105.12844.
- [22] Graser S, Maier T A, Hirschfeld P J, et al. Near-degeneracy of several pairing channels in multiorbital models for the Fe pnictides[J]. *New Journal of Physics*, 2009, 11(2): 025016.
- [23] Patel A A, Lawler M J, Kim E A. Coherent superconductivity with a large gap ratio from incoherent metals[J]. *Physical review letters*, 2018, 121(18): 187001.
- [24] Pikulin D I, Franz M. Black hole on a chip: proposal for a physical realization of the Sachdev-Ye-Kitaev model in a solid-state system[J]. *Physical Review X*, 2017, 7(3): 031006.
- [25] Wang Y. Solvable strong-coupling quantum-dot model with a non-Fermi-liquid pairing transition[J]. *Physical review letters*, 2020, 124(1): 017002.
- [26] Chen A, Ilan R, De Juan F, et al. Quantum holography in a graphene flake with an irregular boundary[J]. *Physical review letters*, 2018, 121(3): 036403.
- [27] Patel A A, Sachdev S. Critical strange metal from fluctuating gauge fields in a solvable random model[J]. *Physical Review B*, 2018, 98(12): 125134.
- [28] You Y Z, Ludwig A W W, Xu C. Sachdev-Ye-Kitaev model and thermalization on the boundary of many-body localized fermionic symmetry-protected topological states[J]. *Physical Review B*, 2017, 95(11): 115150.
- [29] Hu X, Pajerowski D M, Zhang D, et al. Freezing of a Disorder Induced Spin Liquid with Strong Quantum Fluctuations[J]. *Physical Review Letters*, 2021, 127(1): 017201.
- [30] Merz F, Chalker J T. Negative scaling dimensions and conformal invariance at the Nishimori point in the J random-bond Ising model[J]. *Physical Review B*, 2002, 66(5): 054413.
- [31] Levy R, Clark B K. Entanglement Entropy Transitions with Random Tensor Networks[J]. arXiv preprint arXiv:2108.02225, 2021.
- [32] Bi Z, Jian C M, You Y Z, et al. Instability of the non-Fermi-liquid state of the Sachdev-Ye-Kitaev model[J]. *Physical Review B*, 2017, 95(20): 205105.
- [33] Skinner B, Ruhman J, Nahum A. Measurement-induced phase transitions in the dynamics of entanglement[J]. *Physical Review X*, 2019, 9(3): 031009.
- [34] Ikegaya S, Kobayashi S, Asano Y. Symmetry conditions of a nodal superconductor for generating robust flat-band Andreev bound states at its dirty surface[J]. *Physical Review B*, 2018, 97(17): 174501.
- [35] Konig E J, Ostrovsky P M, Protopopov I V, et al. Metal-insulator transition in two-dimensional random fermion systems of chiral symmetry classes[J]. *Physical Review B*, 2012, 85(19): 195130.
- [36] Wei C, Sedrakyan T A. Optical lattice platform for the Sachdev-Ye-Kitaev model[J]. *Physical Review A*, 2021, 103(1): 013323.
- [37] Iyoda E, Katsura H, Sagawa T. Effective dimension, level statistics, and integrability of Sachdev-Ye-Kitaev-like models[J]. *Physical Review D*, 2018, 98(8): 086020.
- [38] Tracy C A, Widom H. The distributions of random matrix theory and their applications[J]. *New trends in mathematical physics*, 2009: 753-765.
- [39] Dumitriu I, Edelman A. Matrix models for beta ensembles[J]. *Journal of Mathematical Physics*, 2002, 43(11): 5830-5847.




Figure 1:  $\Delta_{\alpha\beta}$ -spectrum for four-point interaction (a,b) which follows GUE distribution and two-point interaction (c,d) which follows GSE distribution. In this article we only show the results simulated by setting  $N = 100$ .




Figure 2:  $\eta_{\Delta}$ -spectrum for the unweighted  $(\alpha, \beta)$ -states (a) and the weighted one (b) which corresponds to irreducible group  $\{\eta_{\Delta}\}$ . For  $N = 100$ , there will be nearly  $O(M) \sim \frac{N^2}{4}$  flavors corresponding to distinct values of  $\eta_{\Delta}$ . (c) and (d) show the exact probabilities  $p_i^j$  and  $(p_i^j)p_i^j$  for each  $(\alpha, \beta)$ -state.




Figure 3: (a) Probability distribution of potential difference  $\Delta_{\alpha\beta}$ . (b) The softmax-renormalized results (the  $QT^T$  matrix) which is available to inner product with the value matrix. (c,d) show the exactly calculated  $\Delta_{\alpha\beta}$ -spectrum and the level spacing distribution, while (e,f) show the predicted one using the learnable parameters in self-attention method.




Figure 4: The first, second, and third rows show the three times of learning process using the self-attention method. The learnable parameters are automatically modified after each iteration. The first, second, and third columns are the predicted probability distributions, resulting  $\eta_{\Delta}$ -spectrums, and the  $p_i^{D_i}$ -spectrum which plays the role of loss function here to better identify the accuracy after each step.