

Universal anomalous fluctuations in charged single-file systems

Žiga Krajnik,¹ Johannes Schmidt,^{2,3} Vincent Pasquier,⁴ Tomaž Prosen,¹ and Enej Ilievski¹

¹*Faculty for Mathematics and Physics, University of Ljubljana, Jadranska ulica 19, 1000 Ljubljana, Slovenia*

²*Technische Universität Berlin, Institute for Theoretical Physics, Hardenbergstr. 36, D-10623 Berlin, Germany*

³*Bonacci GmbH, Robert-Koch-Str. 8, 50937 Cologne, Germany*

⁴*Institut de Physique Théorique, Université Paris Saclay, CEA, CNRS UMR 3681, 91191 Gif-sur-Yvette, France*

Conventional classification of dynamical phenomena is based on universal hydrodynamic relaxation characterized by algebraic dynamical exponents and asymptotic scaling of the dynamical structure factor. This work uncovers a novel type of dynamical universality reflected in statistical properties of macroscopic fluctuating observables such as the transmitted charge. By considering a general class of one-dimensional single-file systems (meaning that particle crossings are prohibited) of interacting hardcore charged particles, we demonstrate that stringent dynamical constraints give rise to universal anomalous statistics of cumulative charge currents manifested both on the timescale characteristic of typical fluctuations and also in the rate function describing rare events. By computing the full counting statistics of net transferred charge between two extended subsystems, we establish a number of unorthodox dynamical properties in an analytic fashion. Most prominently, typical fluctuations in equilibrium are governed by a universal distribution that markedly deviates from the expected Gaussian statistics, whereas large fluctuations are described by an exotic large-deviation rate function featuring an exceptional triple critical point. Far from equilibrium, competition between dynamical phases leads to dynamical phase transitions of first and second order. Despite dynamical criticality, we find the large-deviation rate function of the joint particle-charge transfer obeys the fluctuation relation. Curiously, the univariate charge-current rate function experiences a spontaneous breaking of fluctuation symmetry upon varying the particle and charge densities in a nonequilibrium initial state. The rich phenomenology of the outlined dynamical universality is exemplified on an exactly solvable classical cellular automaton of charged hardcore particles. We determine the dynamical phase diagram in the framework of Lee–Yang’s theory of phase transitions and exhibit a hyper-dimensional diagram of distinct dynamical regimes.

CONTENTS

I. Introduction	1	A. Exact full counting statistics	28
II. Setting and background	4	1. Localization	28
A. Representative models	6	2. Scaled cumulants	29
B. Full counting statistics	7	3. Typical fluctuations	30
C. Multivariate fluctuation relation	9	B. Dressing the particle rate function	30
D. Central Limit Theorem	10	C. The Lee–Yang theory	33
III. Summary of the results	10	1. Contour representation	33
A. Large fluctuations: the dressing formalism	15	2. Phase boundaries and the Stokes phenomenon	33
1. Dressing the particle rate function	15	3. Equilibrium and exceptional triple point	34
2. Dressing the moment generating function	17	VI. Conclusion	37
B. Universal anomalous fluctuations in equilibrium	17	A. Dressing the moment generating function	39
C. Dynamical phases	19	B. Fluctuations at equal densities	40
IV. Hardcore cellular automaton	20	References	41
A. Particle fluctuations	20		
B. Joint particle-charge fluctuations	20		
C. Anomalous charge fluctuations	21		
D. Fluctuation symmetry	23		
1. Multivariate fluctuation relation	23		
2. Spontaneous breaking of the univariate fluctuation relation	23		
E. Phase diagrams	25		
V. Technical background	28		

I. INTRODUCTION

The core question in the area of statistical physics is to explain how irreversible phenomenological laws of nature emerge out of microscopic, often highly complex, time-reversible microscopic laws of motion. An overarching goal is to reconcile all our empirical observations within a common quantitative mathematical framework. In this regard, fluctuations theorems offer many

invaluable insights into the foundational aspects of statistical mechanics. One of the general and well-known principles in nonequilibrium physics is the fluctuation-dissipation theorem (FDT), relating fluctuations in thermal equilibrium to how a system responds to small external perturbations. Another profound, albeit somewhat less known, symmetry principle concerns relative probabilities of time-averaged entropy production far away from equilibrium where the condition of detailed balance is no longer satisfied. Specifically, it asserts that the ratio between the probabilities of observing a positive production rate σ_S and a negative rate of the same magnitude takes a *universal* form, exponential in the rate σ_S [1]. To phrase it simply, observing a flow of entropy opposite to what the second law of thermodynamics dictates becomes exponentially unlikely. This fundamental law, known as the transient fluctuation theorem (or the finite-time Evans–Searles symmetry [2, 3]), is universally obeyed in dynamical systems with time-reversal invariant (TRI) evolution laws.

In the context of (quasi)stationary current-carrying steady states, there is an emergent symmetry principle that concerns the structure of temporal fluctuations of macroscopic charge transfer at late times, originally empirically discovered in shear fluids in Ref. [4]. Afterwards, Gallavotti and Cohen [5] provided a rigorous derivation for a certain class of strongly chaotic dynamical systems known as Anosov systems (on compact manifolds). Building on the results of [6], the Gallavotti–Cohen relation (GCR) has soon afterwards been established in Ref. [7] as a general property of finite-state irreducible and aperiodic (ergodic and mixing) Markov chains. The GCR is a symmetry that emerges in nonequilibrium states at late times and, unlike the transient fluctuation theorem, is generally not valid at finite times. As an introduction to the subject, we can recommend Ref. [8], while more comprehensive and technical expositions can be found in Refs. [9–13].

Anomalous transport. The study of nonequilibrium phenomena in many-body systems is currently an active area of theoretical research. The interest has been greatly amplified in recent years thanks to many major experimental advancements in the fields of quantum optics and cold atoms [14–18]. While in generic ergodic many-body systems, the focus has traditionally been on developing efficient numerical tools for computing frequency-resolved conductivities associated with local conserved charges such as energy, particle number, magnetization etc, in recent years, the focus of attention has shifted to non-ergodic phenomena in one-dimensional quantum and classical many-body systems. Particularly, integrable models exhibit anomalous relaxation to generalized Gibbs ensembles [19, 20] and, owing to long-lived ballistic quasiparticle excitations [21–24], feature *anomalous* transport properties such as finite Drude weights [25] or superdiffusive transport of nonabelian charges [26–30] reviewed in [31]. Presence of weak integrability-breaking interactions [32–34] is not always sufficient to restore nor-

mal transport properties [35–37], see Ref. [38] for a review. There are other types of mechanisms that lead to ergodicity breaking, such as e.g. kinetic constraint [39–42] or strong disorder, leading either to a slowdown or a completely inhibited transport.

Our current understanding of transport phenomena is largely centered around the framework of linear response theory. In our view, the reason is mainly two-fold. On the one hand, numerically extracting information beyond two-point dynamical correlation functions such as e.g. the full counting statistics (FCS) is a daunting challenge. On the other hand, large-deviation rate functions are, besides being extremely hard to probe experimentally, believed to be non-universal objects. In this paper, our goal is to demonstrate that

*the notion of dynamical universality reaches
beyond asymptotic scaling of dynamical structure factors,*

quantified by algebraic dynamical exponents and scaling functions.

To corroborate our claim, we introduce a general class of non-ergodic one-dimensional systems of interacting charged particles that display universal properties at the fluctuating level, irrespectively of the algebraic dynamical exponent. On the one hand, we establish that the distribution of the cumulative charge current computed in equilibrium without charge bias is distinctly non-Gaussian (including for charge dynamics with a diffusive dynamical exponent) and rigorously derive its *universal* form that only depends on the scaling variable. Other universal properties are only manifested far from equilibrium. This requires computing the large-deviation rate function (encoding probabilities of exponentially rare events) and analyzing its analytic structure, where we encounter several dynamical regimes. The outlined anomalous statistical properties are unrelated to integrability but rather stem from strongly broken ergodicity due to exponential foliation (or ‘fragmentation’) of the classical many-body phase space.

Large deviation principle. Fluctuating space-time trajectories can be, broadly speaking, divided into two types. When measuring the total charge transferred through the origin, typical deviations from the mean value are of the order $\mathcal{O}(\sqrt{t^{1/z}})$, corresponding to standard deviation of the dynamical process (with dynamical exponent z which governs the relaxation timescale of small perturbations in equilibrium). By contrast, the probability of observing atypically large fluctuations decays to zero (exponentially) at large times. In quantitative terms, large deviations of the order $\mathcal{O}(t^\alpha)$ (for $\alpha > 1/2z$) are expected to obey the so-called large deviation principle (LDP) [43–45], asserting that probabilities of such atypical events satisfy an asymptotic law of the form $\mathbb{P}(j) \asymp \exp[-t^\alpha I(j)]$, where $j = t^{-\alpha} J(t)$ is the rescaled current and $I(j)$ is the large-deviation rate function (LDF) [43].

The Gallavotti–Cohen relation manifests itself as a particular type of inversion symmetry for the probabilities of such rare events. In equilibrium, there is no directionality and hence we simply have that $\mathbb{P}(j) = \mathbb{P}(-j)$. Quite strikingly, the ratio of probabilities $\mathbb{P}(j)/\mathbb{P}(-j)$ tends to a universal function for asymptotically large times even away from equilibrium (i.e. in the absence of local detailed balance) under fairly general conditions provided that the microscopic evolution law exhibits time-reversal symmetry. In this view, the Gallavotti–Cohen fluctuation symmetry may be regarded as a natural extension of the second law of thermodynamics away from equilibrium. In equilibrium, one recovers the Green–Kubo fluctuation-dissipation relation and the Onsager reciprocal relations [7, 13].

Beyond Macroscopic Fluctuation Theory. Accessing dynamical properties beyond the linear response regime by direct simulations is very demanding or even prohibitively difficult in practice. To bypass this obstacle, one commonly resorts to the formalism of Macroscopic Fluctuation Theory (MFT) [46, 47]. Over the years MFT has established itself as a highly versatile theoretical tool for predicting statistical properties of fluctuating macroscopic quantities in generic *diffusive* (mesoscopic or extended) particle-conserving systems. One may regard it as a phenomenological extension of Fick’s law promoted to the level of fluctuating space-time trajectories.

Within the scope of MFT, statistical properties of large fluctuations associated with the transferred charge are generated by variational optimization of a *universal* functional that is fully determined by the linear-response data. The saddle-point equations can be cast as a pair of coupled Hamiltonian PDEs. The MFT field equations do not in general admit an analytic solution and currently there is only a few exactly solvable instances, such as noninteracting random walkers, the simple asymmetric exclusion process [48] and the KMP model [49–51]. By assuming time-stationarity of the optimal trajectories (equivalent to demanding that the LDF obeys the so-called additivity principle [52–56]), the MFT field equations simplify quite dramatically.

There are nonetheless several limitations of the MFT approach. To begin with, the MFT effective action has been originally proposed to capture fluctuations in ergodic and *stochastic* many-body systems governed by *Markovian* evolution laws with a *single* local conserved quantity. Apart from very recent attempts as e.g. Ref. [57], systems with multiple coupled conservation laws have so far remained virtually unexplored. In addition, applicability of MFT technique hinges on a set of underlying assumptions that may sometimes be violated. In fact, a violation of the GCR has already been demonstrated in a number of instances: in non-ergodic Markov chains or Markov chains with absorbing states [58], in Markov processes on semi-infinite lattices [59] or with unbounded local state spaces, and in zero-range processes [60, 61] featuring the condensation transition. The fluctuation symmetry was also found to break down

in memory-dependent (i.e. non-Markovian) systems [62], alongside other examples [63].

Fluctuation theorem in deterministic systems. Using that fluctuation relations rely fundamentally only on time-reversibility of the microscopic evolution laws (and not on noise, for example), one expects them to remain valid in *deterministic* dynamical many-body systems as well. In comparison to stochastic dynamical systems, statistical properties of fluctuating macroscopic quantities in *deterministic* dynamical many-body systems are much less explored. One of the major stumbling blocks is that computing the FCS in generic, ergodic interacting deterministic systems poses a formidable challenge.

The FCS has nevertheless been computed analytically in a wide array of deterministic systems governed by time-reversible microscopic evolution laws, predominantly in the domain of *noninteracting* exactly solvable models including, among others, the transverse Ising chain [64], coupled harmonic oscillators [65], the free Dirac theory [66], two-dimensional conformal field theories [67, 68], the XX quantum chain [69] and the free fermionic chain [70], boundary-driven free fermions [71] and interacting quantum spin chains weakly coupled to Lindbladian baths [72]. These models permit an explicit validation of the GCR. More recently, pivotal progress has been made in certain simple *interacting* exactly solvable classical cellular automata [73–75]. Moreover, a general approach for computing the exact FCS in ballistic systems has been developed in a series of papers [76–78], formulated in the language of generalized hydrodynamics (GHD) [21, 22]. The GCR has been explicitly verified to hold in the classical interacting hard-rod gas and in the quantum Lieb–Liniger model. The GCR has also been recently derived within the so-called ballistic MFT approach in Ref. [79]. The main upshot here is that GCR is seemingly immune to the lack of ergodicity, in contrast with other fundamental principles of statistical physics (such as absence of thermalization, for example).

A general and rigorous proof of the fluctuation theorem in deterministic TRI systems is, to the best of authors’ knowledge, still lacking. Besides the pioneering work [5], it is worth highlighting the scattering theory approach to the fluctuation relation developed in Ref. [80], which establishes the GCR (associated with transmission of energy and charge) in infinitely extended Hamiltonian dynamical systems with time-reversal invariant dynamics. Even though Ref. [80] does not invoke ergodicity or make other assumptions on the underlying dynamics, there is nevertheless a loophole in the proof. The derivation in Ref. [80] rests on an assumption that the generating function for the FCS of the transmitted charge is an everywhere differentiable function of the counting field. It is well known that this technical condition is not fulfilled in systems that can sustain first-order dynamical phase transitions [81–85]. In fact, dynamical phase transitions (DPTs) are known to ruin the additivity property, and thus the GCR could be in jeopardy. While dynamical phase transitions can indeed arise from the MFT func-

tional [86] (commonly associated with time-dependent optimal fluctuating trajectories leading to emergent self-organizing coherent patterns), a detailed analysis shows that critical points are always produced in *pairs* [87–89], thereby leaving the GCR intact [87, 90, 91]. By contrast, in this work we shall discuss dynamical phase transitions of fundamentally different kind that cannot be captured with the the standard framework of the MFT.

Spontaneous breaking of fluctuation symmetry. This work is devoted to a specific class of one-dimensional interacting single-file systems of hardcore particles with internal (charge) degrees of freedom. Owing to the single-file property, particles are purely reflective, i.e. their worldlines never cross. A particularly simple realization of such dynamics is a classical reversible deterministic cellular automaton describing interacting charged particles. In our previous work [74], we have computed the exact FCS of the transferred charge in equilibrium ensembles where, remarkably, we discovered a *non-Gaussian* probability distribution of charge-current fluctuations. There, we attributed this to the absence of sufficient ‘regularity conditions’. In this paper, we further advance this view and gain a more comprehensive understanding behind such unorthodox behavior. More importantly, we go on to demonstrate that consequences of the lack of regularity are even more pronounced away from equilibrium.

It is worth emphasizing that the single-file property not only renders the dynamics (strongly) non-ergodic but additionally, as we explain subsequently, allows for dynamical phase transitions between distinct competing dynamical phases. For this reason, and due to the lack of differentiability of the LDF, the validity of the universal fluctuation relations could be questioned.

By explicitly computing the full counting statistics of joint particle-current fluctuations we uncover two remarkable properties:

- (A) Provided that the univariate particle LDF obeys the GCR, we find that the *joint* LDF invariably obeys the fluctuation symmetry. It is remarkable that the symmetry persists even for non-convex or non-differentiable LDFs resulting from dynamical phase transition of first or second order, respectively (which, to our knowledge, does not follow from any general formal proof).
- (B) The univariate LDF associated to the charge current exhibits an intricate, highly complex structure and, more strikingly, generically undergoes a *spontaneous* breaking of the univariate (Gallavotti–Cohen) fluctuation relation (UFR). The word ‘spontaneous’ signifies that violation occurs at the level of ensemble averages upon varying the control parameters such as e.g. the particle and charge densities (as opposed to explicit breaking of microscopic reversibility).

We wish to emphasize that time-reversal invariance of microscopic dynamics alone is not sufficient to guarantee the Gallavotti–Cohen symmetry of the univariate LDFs.

This has been confirmed in numerous explicit examples of stochastic systems with two coupled currents, for example in molecular motors [92, 93]. In our models, the situation is markedly different. While (by assumption) the univariate GCR is always satisfied for particle dynamics, the univariate GCR of transported charge only holds in certain region of the parameter space bounded by the (dynamical) critical manifolds.

The class of models we consider have not yet, to the best of our knowledge, been investigated previously in the literature. We shall demonstrate that, owing to the distinct lack of ergodicity, the models reveal a *universal*, hitherto undisclosed, non-ergodic behaviour beyond the conventional approaches (such as MFT), including universal non-Gaussian probability distribution of (typical) charge-current fluctuations, coexistence of dynamical phases and novel types of dynamical phase transitions, distinct ‘dynamical regimes’, and emergence of exceptional critical points. We guide the reader through the multifaceted phenomenology of the novel dynamical universality while systematically uncovering the mathematical underpinnings behind the symmetry-breaking mechanism.

Outline. The paper is structured as follows. In Section II we first introduce a novel class of classical one-dimensional systems comprising of hard-core interacting particles carrying internal degrees of freedom that are subjected to a non-crossing (i.e. single-file) constraint. To set the stage we proceed by outlining the general setting which includes a quick introduction into the basic concepts of the large-deviation principle and the Gallavotti–Cohen relation. For concreteness, we also briefly discuss two simple representative examples. Section III contains an exposition of our main results. We open the section by a non-technical overview of the most salient universal features and subsequently zero in on various aspects. We first outline the ‘dressing formalism’ and provide a succinct summary of key results inferred from asymptotic analysis, independently at the level of the moment generating and large-deviation rate functions. We close the section by discussing several remarkable interrelated features, including universal non-Gaussian fluctuations in equilibrium and several competing coexisting dynamical phases. Section IV is devoted to exhibiting our formalism on a concrete model. We choose an exactly solvable classical cellular automaton, which has a remarkably simple (analytic) structure. In Section V we spell out the derivations and supply additional technical details. In Section VI we conclude by summarizing the key results and briefly discuss why our results can be likely further generalized by relaxing the underlying assumptions.

II. SETTING AND BACKGROUND

We begin by outlining the general setting and introducing the basic concepts of fluctuation theory. Once the stage has been set, we shall proceed with a concise

summary of the main results and discuss the key physical features.

We specialize this study exclusively to classical dynamical systems of interacting distinguishable particles in one spatial dimension that conserve the number of particles. Let accordingly $x_\ell = x_\ell(t)$ label the trajectory of the ℓ -th particle and $\mathbf{x}(t) = (x_1(t), x_2(t), \dots, x_{Q_p}(t))$ be the position vector of Q_p particles at time t . Space and time can be continuous or discrete (our main working example, introduced below in Sec. II A, is however a fully discrete model). In addition, particles carry *discrete* internal degrees of freedom c_ℓ , e.g. charge or color, which for simplicity we assume to be binary, $c_\ell \in \mathbb{Z}_2$. Charges can be assigned to particles in an unconstrained manner.

The dynamics can be either deterministic (not necessarily Hamiltonian) or stochastic. In deterministic models, dynamics of particles is governed by the time-local bijective phase-space mapping (propagator) \mathcal{U}_t , formally expressed as $\mathbf{x}(t) = \mathcal{U}_t \circ \mathbf{x}(0)$, that constitutes a group, $\mathcal{U}_{t+t'} = \mathcal{U}_t \circ \mathcal{U}_{t'}$, for both continuous ($t \in \mathbb{R}$) or discrete ($t \in \mathbb{Z}$) times; in the discrete time case the dynamics corresponds to iteration of the one-step propagator \mathcal{U}_1 . For definiteness, we shall confine ourselves to dynamical systems with short range interaction.

More importantly, we now impose two additional dynamical constraints:

- (I) **single-file condition:** particles are not allowed to jump across (or pass by) one another, meaning that their trajectories are subject to the ordering condition

$$x_{\ell-1}(t) < x_\ell(t) \quad (1)$$

at all times t and for all $\ell \in \{2, \dots, Q_p\}$.

- (II) **charge inertness:** charge degrees of freedom remain attached to particles at all times and have no dynamics of their own. Therefore, the c_ℓ carried by ℓ -th particle does not depend on time and the total charge is conserved. Equipping particles with charges does not affect the underlying dynamics of particles governed by the propagator \mathcal{U}_t .

In conjunction the properties (I) and (II) imply that *any* configuration of charges $\mathbf{c} = (c_1, c_2, \dots, c_{Q_p})$ in an initial configuration remains preserved throughout the time evolution. In this view, interaction among the charged particles can be regarded as ‘purely reflective’. Inertness of charge can be also perceived as an extreme version of ‘charge separation’ with trivial (charge) dynamics.

Neglecting the internal degrees of freedom, condition (I) is the defining property of the so-called *single-file dynamics* (see e.g. Refs. [94–97] discussing exact FCS), re-

ferring in general to quasi-1D systems of particles clogged in a narrow channel (so that they are unable to overtake each other). Conservation of the initial charge pattern implies that the entire phase space foliates into 2^{Q_p} dynamical subsectors, closely related to the notion of Hilbert space fragmentation studied in the context of certain non-ergodic quantum dynamics [39, 98].

Separation of charge from matter permits us to integrate out the charge degrees of freedom in an exact analytic fashion and will thus be of pivotal importance for computing the moment generating function (MGF) encoding the statistics of charge transfer.

Time-reversal symmetry. We additionally demand the time evolution to be *time-reversible*: in deterministic dynamics, this requires an involutive phase-space mapping Θ such that

$$\Theta \circ \mathcal{U}_t = \mathcal{U}_{-t} \circ \Theta. \quad (2)$$

In stochastic systems, described by a master equation $dP_t/dt = \mathcal{M}P_t$ with Markov operator \mathcal{M} , TRI follows directly from *detailed balance* (see e.g. [99])

$$\mathcal{M}(C, C')P_{\text{eq}}(C') = \mathcal{M}(C', C)P_{\text{eq}}(C), \quad (3)$$

where C describes the complete configuration of the system (phase space point), e.g. $C = (\mathbf{x}, \mathbf{c})$, and $M(C', C)dt$ is the probability of transition $C \rightarrow C'$ during dt and P_{eq} is the stationary probability measure, $(d/dt)P_{\text{eq}} = 0$, representing e.g. the Gibbs equilibrium state, $P_{\text{eq}} \simeq \exp(-\beta E(C))$. By virtue of detailed balance, the probability of finding a ‘forward’ trajectory $C_{\rightarrow}(t)$ is equal to that of the time-reversed trajectory $C_{\leftarrow}(t)$, that is $\mathbb{P}(C_{\rightarrow}(t)) = \mathbb{P}(C_{\leftarrow}(t))$.

Grand-canonical equilibrium. The total particle number and total charge, denoted by $Q_p = \sum_\ell 1$ and $Q_c = \sum_\ell c_\ell$ respectively, are two general conserved quantities of our models (expressible as spatial sums of local densities). There could in principle be additional local conservation laws, which nevertheless have no impact on our findings. By coupling the two conserved charges to chemical potentials β_p and β_c , we consider grand-canonical ensembles corresponding to the stationary measure $P_{\text{gc}} = \mathcal{Z}^{-1} \exp(\beta_p Q_p + \beta_c Q_c)$, normalized by partition sum \mathcal{Z} . Since both conserved quantities $Q_{p,c}$ represent homogeneous sums of local (one-site) densities, the partition sum factorizes into one-body terms $\mathcal{Z} = \mathcal{Z}_1^N$, $\mathcal{Z}_1 = \exp(f_{\text{eq}})$, where f_{eq} is free energy density per unit length,

$$f_{\text{eq}}(\beta_p, \beta_c) = \log [1 + e^{\beta_p} (e^{\beta_c} + e^{-\beta_c})]. \quad (4)$$

For our convenience, we shall parametrize equilibrium states by specifying averages, $\rho \equiv \langle q_p \rangle_{\text{gc}} = \partial_{\beta_p} f_{\text{eq}}$, and $\rho b \equiv \langle q_c \rangle_{\text{gc}} = \partial_{\beta_c} f_{\text{eq}}$, related to chemical potentials via

$$\beta_p = \log \left[\frac{\rho \sqrt{1 - b^2}}{2(1 - \rho)} \right], \quad \beta_c = \frac{1}{2} \log \left[\frac{1 + b}{1 - b} \right]. \quad (5)$$

By inverting the second relation, we have $b = \tanh(\beta_c)$. The covariance matrix χ of (static) charge susceptibilities is a 2×2 matrix with matrix elements $\chi_{ij} = \partial_{\beta_i} \partial_{\beta_j} f_{\text{eq}}(\beta_p, \beta_c)$ (for $i, j \in \{p, c\}$), reading

$$\chi = \begin{pmatrix} \chi_{pp} & \chi_{pc} \\ \chi_{cp} & \chi_{cc} \end{pmatrix} = \begin{pmatrix} \rho(1-\rho) & \rho(1-\rho)b \\ \rho(1-\rho)b & \rho - (\rho b)^2 \end{pmatrix}. \quad (6)$$

Full counting statistics: bipartitioning protocol. There exist two widely popular settings to compute the full counting statistics of cumulative currents far away from equilibrium. In the mesoscopic approach, a system of finite length is attached between two effective baths which drive it away from equilibrium. At late times, the system relaxes into a stationary current-carrying state, and one can monitor the net charge transfer during a finite interval of time. To eliminate the role of stochastic baths, we shall use an alternative technique. In this work, we consider infinitely extended systems prepared initially in a nonstationary initial state consisting of two thermalized semi-infinite partitions joined at the origin that are released to evolve under a TRI equation of motion. Each partition is initialized in a stationary ensemble characterized by a finite density of particles and charge. The density of particles in the left (right) partition is $\rho_-, (\rho_+) \in [0, 1]$, while $\bar{\rho}_\pm \equiv 1 - \rho_\pm \in [0, 1]$ are the corresponding densities of vacant space. Similarly, we introduce the biases $b_\pm \in [-1, 1]$, such that $\rho_\pm(1+b_\pm)/2$ are the densities of positively charged particles in the left or right partition.

In the general case of unequal densities and biases, $\rho_- \neq \rho_+$ and $b_- \neq b_+$, we find a finite net current of particles and charge flowing across the origin, leaving behind an ever-expanding dynamical interface. The time-integrated particle and charge currents will accordingly grow as $\mathcal{O}(t^\alpha)$ at late times, for some (model-dependent) algebraic dynamical exponent $\alpha \leq 1$.

A. Representative models

The entire class of models that respect the defining conditions (I) and (II) (as specified in Sec. I) display *universal* anomalous fluctuations of macroscopic transferred charge. Before summarizing our key findings in Sec. III, we wish to emphasize that all of our main conclusion hold irrespective of the underlying particle dynamics (apart from minor technical assumptions to ensure that fluctuations of particle trajectories behave sufficiently regularly). Before delving into technical aspects, we find it instructive to introduce two simple models that belong to this particular class of single-file systems.

Exactly solvable hardcore charge gas. Arguably the simplest dynamical system complying with all the above requirements are free, ballistically propagating ‘matter’ particles that carry an internal binary (\mathbb{Z}_2) degree of freedom (which one may think of as charge or color). A




FIG. 1. Hardcore charged gas in discrete space-time, representing a classical cellular automaton of interacting charged particles that propagating ballistically between their collisions. Scattering between positively (red) and negatively (blue) charged particles is purely reflective, yielding the single-file property of non-intersecting particles’ worldlines.

discrete space-time version of such a model is the *hardcore gas cellular automaton*, illustrated in Fig. 1, was introduced and studied initially in Ref. [100] (see also the follow-up works [101–103]).

The $\mathbb{Z} \times \mathbb{Z}$ space-time lattice is occupied with one of the three local states: a positive charge (+), a negative charge (−), or a charge-neutral vacant site \emptyset . The time evolution is realized by sequential ‘brickwork’ application of local two-body maps given by the following rules, the ‘interaction vertex’ $(q_c, q'_c) \leftrightarrow (q_c, q'_c)$ for $q_c, q'_c \in \{+, -\}$, and ‘free vertex’ $(\emptyset, q_c) \leftrightarrow (q_c, \emptyset)$ for $q_c \in \{+, -, \emptyset\}$ (see Ref. [100] for more precise definitions). The propagator \mathcal{U}_{2t} , consisting of alternating odd \mathcal{U}^o and even layers \mathcal{U}^e (satisfying $\mathcal{U}^o \circ \mathcal{U}^o = \mathcal{U}^e \circ \mathcal{U}^e = \text{Id}$), $\mathcal{U}_{2t} = (\mathcal{U}^e \circ \mathcal{U}^o)^t$, is invariant under time-reversal, with $\Theta = \mathcal{U}^o$.

The model, which belongs to a family of (super)integrable classical cellular automata [104], exhibits ballistic charge transport with diffusive corrections. The basic properties of charge transport can be inferred from the asymptotic scaling form of the dynamical structure factor (DSF) of charge density, $S_c(x, t) \equiv \langle q_c(x, t) q_c(0, 0) \rangle_{\text{gc}}$ (normalized by $\sum_x S_c(x, t) = \chi_{cc}$), computed in Ref. [101]

$$S_c(x, t) = \rho(1-b^2) \mathcal{G}_{\text{dif}}(x, t) + \mathcal{D}_c \mathcal{G}_{\text{bal}}(x, t), \quad (7)$$

where $\mathcal{G}_{\text{dif}}(x, t) \equiv (2\pi t D_c)^{-1/2} \exp[-x^2/(2D_c t)]$ denotes a central Gaussian peak that broadens diffusively with diffusion constant $D_c = (1-\rho)/\rho$, whereas $\mathcal{G}_{\text{bal}}(x, t) \equiv \frac{1}{2}(\delta(x-t) + \delta(x+t))$ are two side ballistic ‘sound’ peaks whose magnitude corresponds to the charge Drude weight $\mathcal{D}_c = \rho(1-\rho)b^2$.

The charge diffusion constant D_c (determined from the variance of the asymptotic DSF) and the DC charge conductivity (related to D_c via Einstein relation) can be likewise retrieved from a quenched inhomogeneous profile or in the mesoscopic setup with boundary driving (see Refs. [100–102]). The fluctuation-dissipation relation therefore remains intact.

In order to detect signatures of the fragmented phase

space we have to look beyond the DSF $S_c(x, t)$. To this end, we shall compute the full counting statistics of the cumulative charge current. In our previous study [74], we have established that the stationary PDF of the time-integrated charge current in equilibrium at finite density and no charge bias takes a *non-Gaussian* form. This result is somewhat at odds with the general prediction of ballistic fluctuation theory for integrable models developed in Refs. [76, 77].

Stimulated by the peculiarities observed in Ref. [74], in Sec. IV we revisit the problem of anomalous charge-current fluctuations in the hardcore gas and use this opportunity to clarify the theoretical underpinnings behind the observed anomalous behavior in full capacity. By generalizing the computation of the FCS to nonequilibrium states, we map out an unexpectedly rich phenomenology of the charge-current LDF associated with rare fluctuations.




FIG. 2. Stochastic two-species simple symmetric exclusion process in discrete time with a parallel update scheme: at each timestep, a particle is activated with a finite probability (here $p_{\text{act}} = 1$) and jumps in a random unbiased direction to the target neighboring site provided the target site is unoccupied. Potential conflicts (marked by crosses) are resolved democratically.

Two-species symmetric exclusion process with parallel updates. The other representative example is a stochastic model. We consider the simplest two-species variant of the *symmetric simple exclusion process* (SSEP), realized with a parallel update rule (see Fig. 2 for an illustration): by randomly distributing charged particles on a discrete lattice, at every time-step each particle attempts to jump with probability p to one of its adjacent sites (drawn uniformly at random), subjected to the exclusion rule that prevents particles from jumping to non-vacant sites. When two different particles attempt to jump to the same site, the ‘winner’ is chosen randomly with equal probability (ensuring detailed balance) We have explicitly verified (see Fig. 3) that typical fluctuations on a timescale $t^{1/4}$ are normally distributed.

We stress that the proposed model differs crucially from the two-component AHR model [105] where charge-

carrying particles are allowed to swap their positions with a finite probability. Despite the extra selection rule, dynamics of particles remain diffusive and, similarly to the ordinary simple (symmetric) exclusion process [106], fluctuations of the transmitted particles are normally distributed.




FIG. 3. Typical fluctuations of the particle transfer in the two-species symmetric simple exclusion process in discrete time with a parallel update rule, showing the time-dependent PDF $\mathcal{P}_p(\mathcal{J}_p|t)$ of the rescaled cumulative particle current $\mathcal{J}_p(t) = t^{-1/2z_p} J_p(t)$ for different times (in normal and logarithmic scale), with system size $L = 10^6$ and particle density $\rho = 0.5$ (with zero charge bias $b = 0$). Particles undergo diffusive motion with dynamical exponent $z_p = 2$. Black curves correspond to the asymptotic stationary PDF $\mathcal{P}_p^{\text{typ}}(j_p) = \lim_{t \rightarrow \infty} t^{1/4} \mathcal{P}_p(j_p = \mathcal{J}_p|t)$. A least-squares fit to a Gaussian profile gives an estimated standard deviation $\sigma_p \approx 0.441 \pm 0.002$.

B. Full counting statistics

We now make a slight digression to introduce the key objects for computing the full counting statistics of charge transfer. In the following we assume space and time to be continuous. We make this choice solely for compactness of notation, and adapting our construction to lattice models in continuous time or to the fully discrete setting is straightforward.

We consider an extended system with a finite number N of globally conserved charges $Q_i = \int dx q_i(x, t)$ enumerated by label i , with $q_i(x, t)$ denoting the local densities at position x and time t . The associated current densities, denoted by $j_i(x, t)$, are determined from the local continuity relations, $\partial_t q_i(x, t) + \partial_x j_i(x, t) = 0$.

The aim is to quantify fluctuations of the total time-integrated current density flowing through the origin in a time interval t . To this end, we introduce cumulative currents (also known as *Helfand moments* [107])

$$J_i(t) = \int_0^t d\tau j_i(0, \tau). \quad (8)$$

The full set of temporally extensive dynamical variables $\mathbf{J}(t) = (J_1(t), \dots, J_N(t))$ quantifies the net transferred charges through the origin over a time period t . By prescribing an initial state (an ensemble of phase space points i.e. configurations), the task is to derive the joint PDF $\mathcal{P}(\mathbf{J}|t)$.

We focus our attention to two physically distinguished timescales. Helfand moments $J_i(t)$ associated to *typical* trajectories are of the order $\mathcal{O}(t^{1/2z_i})$, corresponding to the standard deviation, i.e. the square root of the second cumulant (variance)

$$c_2^{(i)}(t) = \left\langle J_i^2(t) \right\rangle_{\text{init}}^c \sim t^{1/z_i}. \quad (9)$$

In diffusive systems, the characteristic dynamical exponent equals $z = 2$. By contrast, due to ballistically propagating quasiparticles in integrable systems one generically finds ballistic scaling $z = 1$, see e.g. Refs. [76–78]. There are nevertheless important exceptions to this generic behavior, for example, charge transport in the presence of charge-conjugation symmetry [74, 108].

Coupling $J_i(t)$ to ‘counting fields’ λ_i , we introduce the *multivariate moment generating function* (MMGF),

$$G(\boldsymbol{\lambda}|t) = \left\langle e^{\boldsymbol{\lambda} \cdot \mathbf{J}(t)} \right\rangle_{\text{init}}, \quad (10)$$

where $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$ and the ensemble averaging is done with respect to a prescribed *non-stationary* initial state. For discrete variable systems, $J_i(t) \in \mathbb{Z}$, hence periodicity in the imaginary direction permits us to restrict λ_i to the infinite cylinder, $\lambda_i \in \mathbb{R} \times S^1 \subset \mathbb{C}$ (we shall largely restrict our considerations to real λ_i , and only relax this condition in Sec. VC). The multivariate MGF is given by the multivariate Laplace transform of the time-dependent joint PDF $\mathcal{P}(\mathbf{J}|t)$,

$$G(\boldsymbol{\lambda}|t) = \int d\mathbf{J} e^{\boldsymbol{\lambda} \cdot \mathbf{J}} \mathcal{P}(\mathbf{J}|t), \quad (11)$$

where $d\mathbf{J} \equiv \prod_{i=1}^N \int dJ_i$. All the higher (finite-time) cumulants $c_{n>2}^{(i)}(t)$ of the integrated current density $J_i(t)$ can be computed with aid of the *multivariate cumulant generating function* (MCGF),

$$F(\boldsymbol{\lambda}|t) = \log G(\boldsymbol{\lambda}|t). \quad (12)$$

via

$$c_n^{(i)}(t) = \left. \frac{\partial^n F(\boldsymbol{\lambda}|t)}{\partial \lambda_i^n} \right|_{\boldsymbol{\lambda}=0}. \quad (13)$$

Furthermore, let $\mathcal{P}_i(J_i|t)$ denote the univariate (time-dependent) PDFs obtained by marginalization of the full joint PDF $\mathcal{P}(\mathbf{J}|t)$,

$$\mathcal{P}_i(J_i|t) = \int \prod_{l \neq i} dJ_l \mathcal{P}(\mathbf{J}|t). \quad (14)$$

The associated univariate MGFs $G_i(\lambda_i|t)$ are obtained by setting all $\lambda_{l \neq i}$ to zero, $G_i(\lambda_i|t) \equiv G(\boldsymbol{\lambda}|t)|_{\lambda_{l \neq i} \rightarrow 0}$; they

correspond to the Laplace transform of the PDFs $\mathcal{P}_i(J_i|t)$

$$G_i(\lambda_i|t) = \int dJ_i e^{\lambda_i J_i} \mathcal{P}_i(J_i|t). \quad (15)$$

Typical fluctuations. To infer the *stationary* PDFs $\mathcal{P}(\mathcal{J}_i|t)$ associated with typical fluctuations, we first rescale the cumulative current $J_i(t)$ as $\mathcal{J}_i(t) \equiv t^{-1/2z_i} J_i(t)$, and subsequently take the large-time limit,

$$\mathcal{P}_i(j_i) = \lim_{t \rightarrow \infty} t^{1/2z_i} \mathcal{P}_i(\mathcal{J}_i = j_i|t). \quad (16)$$

Large deviation principle. The theory of large deviations [43–45] deals with probabilities of exponential form. According to the large deviation principle, atypical fluctuations of the cumulative currents $J_i(t)$ decay exponentially with time,

$$\mathbb{P}(J_i(t)) \asymp e^{-t^{\alpha_i} I_i(j)}, \quad (17)$$

where exponents α_i are referred to as ‘speeds’. Large fluctuations are in general reserved for fluctuations of the largest magnitude $\mathcal{O}(t^{\alpha_i})$ (whereas fluctuations larger than typical are sometimes referred to as ‘moderate fluctuations’).

Under certain technical conditions, the LDFs $I_i(j_i)$ can be extracted from the associated MGFs. The asymptotic scaling of the univariate MGFs,

$$G_i(\lambda_i|t) \asymp e^{t^{\alpha_i} F_i(\lambda_i)}, \quad (18)$$

is governed by *scaled* cumulant generating functions (SCGFs)

$$F_i(\lambda_i) = \lim_{t \rightarrow \infty} t^{-\alpha_i} \log G_i(\lambda_i|t). \quad (19)$$

Note that dynamical exponents α_i are (by definition) the largest exponents such that $F_i(\lambda_i)$ exists and are non-trivial. Moreover, $F_i(\lambda_i)$ are convex functions of the counting field $\lambda_i \in \mathbb{R}$. Extended diffusive systems are characterized by $\alpha = 1/2$, see e.g. [97, 109]. By contrast, systems that support long-lived (free or interacting) quasiparticle excitations generically exhibit growth with ballistic exponent $\alpha = 1$.

By taking into account that all the currents are mutually coupled we subsequently put $\alpha_i = \alpha$ for all i . To properly exhibit the symmetry properties of the counting process, it is crucial to treat all the (cumulative) currents on equal footing. The joint LDF $I(\mathbf{j}) \equiv I(j_1, \dots, j_N)$ associated to $\mathcal{P}(\mathbf{J}|t)$ is accordingly given by

$$I(\mathbf{j}) = \lim_{t \rightarrow \infty} t^{-\alpha} \log \mathcal{P}(\mathbf{j} = \mathcal{J}|t). \quad (20)$$

Provided that the multivariate SCGF $F(\boldsymbol{\lambda})$ is everywhere differentiable on its domain \mathbb{R}^N , the Gärtner–Ellis theorem ensures that the LDF $I(\mathbf{j})$ is given by the Legendre–Fenchel transform

$$I(\mathbf{j}) = F^*(\mathbf{j}) = \sup_{\boldsymbol{\lambda} \in \mathbb{R}^N} \{ \boldsymbol{\lambda} \cdot \mathbf{j} - F(\boldsymbol{\lambda}) \}, \quad (21)$$

representing a convex, lower-semicontinuous and non-negative multivariate function obeying $\inf_{\mathbf{j} \in \mathbb{R}^N} I(\mathbf{j}) = 0$. At late times, the expectation values of currents are encoded in the first moment,

$$\bar{\mathbf{j}} = \lim_{t \rightarrow \infty} t^{-\alpha} \langle \mathbf{J}(t) \rangle_{\text{init}} = \left. \frac{dF(\boldsymbol{\lambda})}{d\boldsymbol{\lambda}} \right|_{\boldsymbol{\lambda}=\mathbf{0}}, \quad (22)$$

such that $I(\bar{\mathbf{j}}) = 0$.

Regularity. It is instructive to briefly discuss the formal properties of the univariate SCGFs $F(\lambda)$ (suppressing the subscript label). In physics literature, $F(\lambda)$ is introduced as the generating function of *scaled* cumulants s_n through the series expansion

$$F(\lambda) = \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} s_n. \quad (23)$$

It is important to keep in mind, however, that this is *not* unconditionally true. Instead, only when all cumulants $c_n(t)$ grow asymptotically with a common algebraic exponent, that is $c_n(t) \sim t^\alpha$, we have that $s_n = \lim_{t \rightarrow \infty} t^{-\alpha} c_n(t)$. As emphasized in [74], only provided that the large-time limit $F(\lambda) = \lim_{t \rightarrow \infty} t^{-\alpha} F(\lambda|t)$ can be exchanged with an infinite sum, it is guaranteed that s_n exist and correspond to the series coefficients of the SCGF $F(\lambda)$. Establishing that $F(\lambda)$ is a *real* analytic function around the origin $\lambda = 0$ (with a finite radius of convergence) is, perhaps non-intuitively, not enough to ensure interchangeability of limits. Rather, a stronger sufficient ‘regularity condition’ is required, as first pointed out by Bryc [110]: if $F(\lambda|t)$ is *holomorphic* at all times in some finite fixed neighborhood around the origin in the complex λ -plane, then $F(\lambda)$ represents a *faithful* generating function of scaled cumulants (see also [12]). To our knowledge, Bryc regularity does not follow from a more general principle and thus it remains an open question whether faithfulness of $F(\lambda)$ can be formally established without invoking any model-specific information. Lastly, we note that even an unfaithful SCGF is still physically meaningful. Assuming it is everywhere differentiable on its domain, it provides the LDF via the Legendre transform, $I(j) = F^*(j) \equiv \sup_{\lambda \in \mathbb{R}} \{\lambda j - F(\lambda)\}$.

While the reader might be under the impression that establishing faithfulness of $F(\lambda)$ offers very little besides obfuscating a conceptually simple formalism, we in fact wish to argue exactly the opposite. Breakdown of regularity can have severe consequences and therefore should not be disregarded. For some reason, this subtle aspect has apparently gone completely unnoticed (in physics applications at least) prior to our recent work [74] where, by studying charge-current fluctuations in the hard-core automaton, we have rigorously shown that the MGF in equilibrium (at finite particles density and general bias) fails to satisfy the aforementioned regularity condition, which can be traced back to emergent dynamical criticality (attributed to Lee–Yang zeros colliding with the real axis at the origin of the λ -plane, see Sec. V).

C. Multivariate fluctuation relation

By virtue of *detailed balance* there is, on average, no flow of currents in equilibrium states. Consequently, fluctuations (both typical and large) of magnitudes $\pm j_i$ away from the mean value $\bar{j}_i = 0$ are equiprobable. This is no longer the case away from equilibrium as $\bar{j}_i \neq 0$ implies a preferred direction for particle and current fluxes. Accordingly, observing a large deviation from the mean current in the direction of the flow is exponentially more likely than observing a current of the same magnitude flowing in the opposite direction. Remarkably, the probabilities of the two events can be related by exploiting time-reversibility of the microscopic evolution law, yielding a *universal* ratio of the form for asymptotically large times

$$\frac{\mathbb{P}(\mathbf{j})}{\mathbb{P}(-\mathbf{j})} \asymp e^{t^\alpha \boldsymbol{\sigma}_\varepsilon(\mathbf{j})}, \quad (24)$$

with a linear form $\boldsymbol{\sigma}_\varepsilon(\mathbf{j}) = \boldsymbol{\varepsilon} \cdot \mathbf{j}$ and the vector of ‘thermodynamic forces’ $\boldsymbol{\varepsilon} \equiv (\varepsilon_1, \dots, \varepsilon_N) \in \mathbb{R}^N$, customary called *affinities*. In the rest of the paper, we shall refer to Eq. (24) as the *multivariate fluctuation relation* (MFR), see e.g. Ref. [13]. The average of the exponent in Eq. (24) can be interpreted as the rate of thermodynamic entropy production $\sigma_S \equiv dS(t)/dt \geq 0$,

$$\sigma_S = \boldsymbol{\varepsilon} \cdot \bar{\mathbf{j}} = \lim_{t \rightarrow \infty} t^{-\alpha} D_{\text{KL}}(\mathcal{P}(\mathbf{J}|t) || \mathcal{P}(-\mathbf{J}|t)), \quad (25)$$

coinciding with the rescaled (in units of t^α) *relative entropy* (also known as the Kullback–Leibler divergence) of $\mathcal{P}(\mathbf{J}|t)$ and its time-reversed counterpart $\mathcal{P}(-\mathbf{J}|t)$, $D_{\text{KL}}(\mathcal{P}(\mathbf{J}|t) || \mathcal{P}(-\mathbf{J}|t)) = \int d\mathbf{J} \mathcal{P}(\mathbf{J}|t) \log \left[\frac{\mathcal{P}(\mathbf{J}|t)}{\mathcal{P}(-\mathbf{J}|t)} \right]$. The LDF $I(\mathbf{j})$ therefore satisfies the relation

$$I(-\mathbf{j}) - I(\mathbf{j}) = \boldsymbol{\varepsilon} \cdot \mathbf{j}. \quad (26)$$

Expressed in terms of the multivariate SCGF, the MFR is manifested as an *inversion symmetry* around $\boldsymbol{\lambda} = -\boldsymbol{\varepsilon}/2$,

$$F(\boldsymbol{\lambda}) = F(-\boldsymbol{\lambda} - \boldsymbol{\varepsilon}). \quad (27)$$

This is a multivariate generalization of the celebrated Gallavotti–Cohen fluctuation relation [7, 13].

In stochastic systems, the Gallavotti–Cohen relation is a corollary of the additivity principle [52]. The GCR is however obeyed even in the absence of additivity principle (e.g. in systems supporting dynamical phase transitions), see. Ref. [111]. In higher-dimensional TRI systems there exists more general, so-called isometric, fluctuation relations [8]. In the scope of MFT, the fluctuation symmetry for particle-conserving TRI diffusive systems takes a universal form (see Ref. [112]), reading $\varepsilon(\rho_-, \rho_+) = \int_{\rho_+}^{\rho_-} d\rho [2D(\rho)/\sigma(\rho)] = \int_{\rho_+}^{\rho_-} d\rho f'_{\text{eq}}(\rho) = f'_{\text{eq}}(\rho_-) - f'_{\text{eq}}(\rho_+)$.

Univariate fluctuation relations. Unlike the multivariate SCGF $F(\boldsymbol{\lambda})$, univariate SCGFs $F_i(\lambda_i)$ will not in general display any particular symmetry property in spite of TRI [7]. Such lack of symmetry is simply due to the fact that all currents flip sign under the time-reversal, that is $\mathbf{j}_i \circ \Theta = -\mathbf{j}_i$. One can nonetheless identify situations when even the marginalized SCGFs $F_i(\lambda_i)$ possess a Gallavotti–Cohen symmetry of the form

$$F_i(\lambda_i) = F_i(-\lambda_i - \tilde{\varepsilon}_i), \quad (28)$$

for some ‘effective’ univariate affinity $\tilde{\varepsilon}_i$ (in general differing from the affinity component ε_i). This happens, for example, when (i) the nonequilibrium state is induced by a single thermodynamic force (corresponding to bias $\delta\mu_i \neq 0$), (ii) whenever the i th cumulative current $J_i(t)$ is parametrically slower or faster than other cumulative currents (see e.g. Ref. [113]) or (iii) under the ‘tight-coupling condition’ $J_i \simeq J_\ell$ for all $\ell \neq i$ (see e.g. Ref. [114]).

In Section III, we describe another, different, dynamical mechanism involving two coupled currents that obey both the univariate and joint fluctuation relations. However, while the UFR associated to the particle current is always obeyed, the UFR of the charge current can undergo spontaneous breaking.

D. Central Limit Theorem

The celebrated Central Limit Theorem – another cornerstone result in probability theory – states that empirical means obtained by sampling identical and independent random variables become normally distributed when the number of samples grows large. More remarkably, even in strongly interacting particle systems subjected to highly non-trivial (temporal) correlations (as commonly found in physics applications), one empirically finds that temporal clustering of correlations is typically strong enough to preserve the central limiting behavior. In other words, if the memory of all initially correlated local observables (the current density, for example) decays sufficiently fast, fluctuations of the associated macroscopic dynamical quantity (the time-integrated current density, for example) cannot be distinguished from those of a random process.

Typical values of the cumulative current $J(t)$ at large t are proportional to the standard deviation $\sqrt{c_2(t)} \sim t^{1/2z}$. To infer the associated *stationary* PDF, the time-integrated current density has to be rescaled as $\mathcal{J}(t) = t^{-1/2z}J(t)$, yielding

$$\mathcal{P}^{\text{typ}}(j) \equiv \lim_{t \rightarrow \infty} t^{1/2z} \mathcal{P}(j = \mathcal{J}(t)). \quad (29)$$

Cumulants of $\mathcal{P}^{\text{typ}}(j)$, denoted by κ_n , are (assuming the limits exist, i.e. $\kappa_n < \infty$ for all n) accordingly given by

$$\kappa_n(t) \equiv t^{-n/2z} c_n(t), \quad \kappa_n = \lim_{t \rightarrow \infty} \kappa_n(t). \quad (30)$$

Then, the central limit behavior (CLT property) holds if and only if κ_2 is finite and non-zero ($0 < \kappa_2 < \infty$) and $\kappa_{n>2} = 0$, implying $\mathcal{P}^{\text{typ}}(j)$ is a Gaussian distribution of zero mean and finite variance κ_2 . Fluctuations whose PDFs deviate from Gaussianity can be regarded as *anomalous*.

Bryc regularity provides a *sufficient* condition of the CLT property. More specifically, complex analyticity of $G(\lambda|t)$ within a disc centered around $\lambda = 0$ guarantees existence of scaled cumulants, i.e. $s_n < \infty$ for all $n \in \mathbb{N}$, which in turn implies that $\lim_{t \rightarrow \infty} \kappa_{n>2} = 0$, while $\kappa_2 = s_2$ stays finite. Lack of regularity on the other hand opens the door for anomalous statistics associated to typical events. Beware however, that absence of regularity is not necessary detrimental for the CLT property (see Ref. [74] for an example). As pointed out previously in Refs. [74, 108],

singular scaled cumulants arise if $z > 1/\alpha$,

signifying that $c_2(t) \sim t^{1/z}$ grow asymptotically *slower* than t^α , namely the exponent governing the asymptotic growth of $G(\lambda|t)$. Absence of regularity is manifested, for example, in certain widely studied integrable systems that support subballistic (either diffusive [23, 25, 115] or superdiffusive [27, 29, 31]) charge transport. It is nonetheless not inherently linked to integrability. For instance, the proposed parallel-update SSEP is one of the simplest stochastic models violating Bryc regularity: while particles diffuse through the system ($\alpha = 1/2$), charge is slowed down by the exclusion rule and instead spreads subdiffusively with dynamical exponent $z = 4$.

III. SUMMARY OF THE RESULTS

In this section, we give a non-technical exposition of our main findings. We mostly focus on anomalous *rare* charge-current fluctuations in the considered charged single-file systems, describe their universal properties, the underlying dynamical phases and transition among them.

In a nutshell, the unexpectedly rich behavior comes from an interplay of the two defining properties: the single-file constraint and inertness of charge. More specifically, since particles are unable to overtake each other, we can readily conclude that any increase of $J_p(t)$ in a given time interval t is due to right-moving particles originating from the negative half of the system ($x < 0$) and, conversely, that any decrease comes from particles that have started from the positive half ($x > 0$). Indeed, the same logic applies to charge degrees of freedom (owing to their inertness), where additionally we have to account for the charge-dependent sign. The simple picture that emerges is that each of the two partitions acts as a separate source of fluctuations, each attempting to enforce its own fluctuations on the whole system. Although the basic picture is deceptively simple, there is no obvious way to tell a priori which of the two halves eventually

prevails upon averaging over an ensemble of initial configurations. As we now describe, large charge-current fluctuations have quite a complex structure and intricately depend on the initial condition (i.e. the density of particles and average charge) *and* the magnitude of fluctuations.

Dressing approach. Our main goal is to compute the full time-dependent joint PDF $\mathcal{P}_{c,p}(J_c, J_p|t)$ (here $N = 2$) and subsequently infer from it the joint LDF $I_{c,p}(j_c, j_p)$. Here variables j_i pertain to dynamically rescaled cumulative (i.e. time-integrated) particle and charge currents, $\mathcal{J}_i(t) = t^{-\alpha} J_i(t)$, where α quantifies the timescale of large (exponentially rare) events.

Computing exact finite-time rate function in genuinely *interacting* systems seems a rather hopeless task. Even in ‘exactly solvable’ models, computing the MGF at finite times presents a daunting challenge. Fortunately however, the principal characteristic features of the considered dynamically constrained models can be *qualitatively* described in a fully analytic and rigorous fashion, provided that the underlying statistical properties of particle dynamics are supplied as a phenomenological input (similarly as in the MFT, where one provides the diffusion constant and conductivity).

Using that charges have no effect on the underlying particle dynamics, the counting statistics for the charge degrees of freedom can be resolved in a purely combinatorial fashion. For this reason, we suggestively call this technique the “dressing” approach. Here we only briefly describe the basic idea and leave a detailed analysis to Section V. The main object is the time-independent ‘dressing factor’ $I_{c|p}(j_c, j_p)$, representing the conditional probability for observing j_c for a given value of j_p . By adjoining the particle-current rate function $I_p(j_p)$, we obtain a joint bivariate LDF of the form $I_{c,p}(j_c, j_p) = I_{c|p}(j_c, j_p) + I_p(j_p)$. In Sec. V, we establish that a fluctuation relation of the form $I_{c,p}(-j_c, -j_p) - I_{c,p}(j_c, j_p) = \varepsilon_c j_c + \varepsilon_p j_p$ is satisfied provided that $I_p(j_p)$ obeys the univariate Gallavotti–Cohen relation $I_p(-j_p) - I_p(j_p) = \tilde{\varepsilon}_p j_p$.

In the following, we confine our analysis mostly to the univariate LDF $I_c(j_c)$. The reason is two-fold. Firstly, since $I_c(j_c)$ inherits the most salient qualitative properties of $I(j_c, j_p)$, we find it better suited to exhibit the underlying dynamical criticality. Secondly, $I_c(j_c)$ can undergo spontaneous breaking of fluctuation symmetry. To compute $I_c(j_c)$, the joint LDF $I_{c,p}(j_c, j_p)$ has to be minimized over the range of j_p . The biphasic structure of the dressing factor allows us to perform a ‘chiral decomposition’ into two separate optimizations $I_{c,p}^{(\pm)}(j_c, j_p)$ associated with two branches of the rate function, $I_c^{(\pm)}(j_c) = \sup_{\lambda} \{\lambda j_c - F_{\pm}(\lambda)\}$. In practice, one first carries out ‘inner optimizations’ on $I_{c,p}^{(\pm)}(j_c, j_p)$ yielding $I_c^{(\pm)}(j_c)$, and finally selects the optimal global value (for fixed j_c), namely $I_c(j_c) = \min\{I_c^{(\pm)}(j_c)\}$.

Coexisting dynamical phases and first-order dynamical phase transition. We now describe the main universal characteristics of the charge LDF $I_c(j_c)$. We find it convenient to discuss it in terms of its Legendre-dual $F_c(\lambda_c)$, which we formally view as the dynamical free energy density governing the asymptotic growth of the dynamical partition sum $G_c(\lambda_c|t)$. To lighten our notation, we subsequently drop the subscript label ‘ c ’ from the univariate charge MGF and LDF (while making the identifications $\lambda_c \rightarrow \lambda, j_c \rightarrow j$).

During a finite window of time t , the associated ‘dynamical free energy’ $F(\lambda|t)$ thus receives contributions $F_{\pm}(\lambda)$ from two distinct *dynamical phases*. We can picture them as distinct branches of the dynamical free energy (measured in units of t^α), see Eq. 19. However, only the larger (in magnitude) of the two branches is physically relevant at late times. The other phase is subleading and merely visible as a transient finite-time correction that fades away exponentially with time. The exception to this are equilibrium states, where detailed balance ensures that both branches contribute equally.

Characterizing the nature of charge-current fluctuations boils down to determining which of the two competing branches F_{\pm} dominates for any specified value of the counting field $\lambda \in \mathbb{R}$. Based on this, we can thus anticipate two intervals, denoted by $\mathcal{I}_{\pm} \subset \mathbb{R}$, along which the respective branches F_{\pm} dominate the asymptotic growth of (charge) MGF $G(\lambda|t)$. On purely formal grounds, we can regard these two (meta)stable branches $F_{\pm}(\lambda)$ as distinct *dynamical phases*. We thus deal with a scenario that closely resembles the physics of first-order thermodynamic phase transitions [116].

Suppose that $F_{\pm}(\lambda)$ exchange dominance at $\lambda = \lambda_{\bowtie}$. If $F_{\pm}(\lambda)$ are strictly convex, then λ_{\bowtie} represents a non-differentiable (corner) point in $F(\lambda)$. Non-differentiable points are typically a precursor of a *first-order* (dynamical) phase transition.

How emergence of non-differentiable points affects the large-deviation rate function is less obvious and requires a careful analysis. To begin with, presence of a corner no longer guarantees that the rate function $I(j)$ coincides with the Legendre dual $F^*(j)$ of the SCGF $F(\lambda)$. In general, $F^*(j)$ is only a convex hull that bounds the physical rate function $I(j)$ from below [43]. Conversely, the Legendre–Fenchel transform of $I(j)$ *always*, regardless of convexity, yields the physical $F(\lambda)$. A non-differentiable corner point in $F(\lambda)$ translates to an *affine* (i.e. linear) segment in $F^*(j) = \sup_{\lambda \in \mathbb{R}} \{\lambda j - F(\lambda)\}$, spanning a contiguous range of values in j between the left and right derivatives of $F(\lambda)$ at the corner λ_{\bowtie} . Any non-differentiable point in $F(\lambda)$ therefore erases some information about the rate function, meaning that computing the rate function $I(j)$ necessitates additional information beyond that provided by $F(\lambda)$ alone. In general, $F(\lambda)$ with a corner point implies that the charge-current rate function $I(j)$ cannot be strictly convex everywhere; it is either non-convex or it contains an affine part, in formal analogy to non-concave microcanonical

entropies that imply inequivalent (thermodynamic) ensembles [117, 118].

Based on the general formal analysis of the solutions to the optimization problem (see Section V) we conclude that $I(j)$ never develops an affine parts. Instead, it simply consists of two ‘patches’ of *locally* convex branches, meaning that convexity of $I(j)$ will not be preserved globally (i.e. for the entire admissible range of large integrated currents j). Until the critical value j_{\bowtie} , large charge-current fluctuations are realized by one of the partitions (branches) $I^{(\pm)}(j < j_{\bowtie})$, beyond which the events from the other branch $I^{(\mp)}(j > j_{\bowtie})$ become more probable and take over. We have thus eliminated the possibility of stable phase coexistence. Coexistence of dynamical phases, emerging in certain systems supporting second-order DPTs associated with particle-hole symmetry breaking, is associated with *convex* rate functions possessing affine parts, corresponding to the Legendre–Fenchel transform $F^*(j)$. In our case, the single-file constraint on particle trajectories prohibits phase coexistence. We find that the rate function $I(j)$ is strictly convex everywhere, except at the critical point $j = j_{\bowtie}$ where both branches intersect, $I^{(+)}(j_{\bowtie}) = I^{(-)}(j_{\bowtie})$.




FIG. 4. Continuous symmetry breaking of convexity of a regular bulk branch: upon smoothly varying b_+ , a (strictly) convex bulk branch $F_+(\lambda)$ (light red) with a single minimum undergoes a continuous transition to a symmetry-broken non-convex shape (dark red) with a doubly degenerate minimum at value F_0 . The middle red curve shows the critical shape when the two minima $\lambda_{\pm}^{(\pm)}$ of $F_+(\lambda)$ (either real or purely imaginary) coalesce. The other bulk branch $F_-(\lambda)$ (blue) always remains in a symmetry-broken form and is unaffected by the change of b_+ .

The central question now is whether non-differentiable points in $F(\lambda)$ have any adverse consequence for the UFR. A crucial observation in this respect is that the first-order DPTs emerge due to appearance of a single critical point. This is to be contrasted with other symmetry-breaking scenarios discussed previously in the literature (see Refs. [88, 89]) where critical points are produced in pairs, that is symmetrically with respect to

inversion point of an unbroken phase. In the absence of the second-order critical points, the UFR will still hold locally for all j that are smaller in magnitude than the distance of the non-differentiable point in $I(j)$ from the origin. In contrast, for large charge-current fluctuations in the direction of the flow that in magnitude exceed the critical value j_{\bowtie} , fluctuations in the opposite direction of the same size are realized by a different bulk branch and hence the GCR (26) will no longer be satisfied *globally* for all values of j .

Dynamical phase transition of second order. The class of models we consider supports another type of dynamical phase transitions. There is a subtle symmetry-breaking mechanism that induces a DPT of the *second order*, signaled by the emergence of (strictly) flat segments in $F(\lambda)$. In fact, individual (meta)stable branches $F_{\pm}(\lambda)$ may develop such a flat piece, arising as a consequence of a *continuous* (i.e. second-order) phase transition of Landau–Ginzburg type from a strictly convex, called regular, form to a *symmetry-broken* non-convex shape with a doubly-degenerate minimum. Such a transition from a regular to a symmetry-broken branch is illustrated in Fig. 4.

Under certain (mild) ‘regularity assumptions’ on the particle rate function $I_p(j_p)$ (cf. Sec. V), *at least* one of the branches $F_{\pm}(\lambda)$ has broken symmetry. This property can be again deduced by investigating the formal structure of solutions to the outlined optimization problem. It can be shown that convexity at $j_c = 0$ enforces that at least one of the branches attains the minimum at the boundary of the optimization domain at $j_p = 0$. Lack of convexity precludes a symmetry-broken branch to be physical for the entire range of real counting fields $\lambda \in \mathbb{R}$. We must then distinguish between the following cases: (i) one of the bulk branches F_{\pm} is regular and thus directly corresponds to $F(\lambda)$, or (ii) only the strictly convex parts of $F_{\pm}(\lambda)$ that has experienced a symmetry-breaking transition are physical. In the latter case, the missing range of λ (lying along the unphysical part of the broken branch) must then be identified with the flat branch $F_0 < 0$, being the dominant contribution to the MGF $G(\lambda|t)$ at late times.

We next discuss the distinguished role of the constant branch. Unlike $F_{\pm}(\lambda)$, it appears (for physical values $\lambda \in \mathbb{R}$) only away from equilibrium. The corresponding ‘interval of dominance’ $\mathcal{S}_0 \subset \mathbb{R}$ is a *single* compact interval on the real λ -axis (always excluding the origin) located in between two doubly-degenerate minima of the physical branch $F_{\pm}(\lambda \in \mathbb{R} \setminus \mathcal{S}_0)$. Another important general property (see Sec. V) is that only a regular dominant (i.e. physical) branch can undergo a symmetry-breaking transition, whereas the subdominant branch remains broken throughout. This scenario is visualized in Fig 4. By continuously varying the counting field along the real λ -axis, the physical bulk branch F_{\pm} reaches its minimum $\lambda = \lambda_{\pm}^{(-)}$, ‘jumps’ over to F_0 and, upon reaching another minimum at $\lambda_{\pm}^{(+)} > \lambda_{\pm}^{(-)}$, returns back to

the same branch F_{\pm} . There is however another possible scenario when both branches F_{\pm} have broken symmetry; it may occur that the closest (degenerate) minimum to $\lambda_{\pm}^{(-)}$ belongs to the other branch at $\lambda_{\mp}^{(+)}$, in which case for $\lambda > \lambda_{\mp}^{(+)}$ (assuming absence of first-order phase transitions upon further increasing λ) the dominant branch is F_{\mp} .

We have thus far established the following picture. Either of the bulk branches $F_{\pm}(\lambda)$ may, upon varying the densities or biases in the initial state, experience a second-order transition to a symmetry-broken form with two-fold degenerate minima. The transition occurs when the unique minimum of an unbroken physical branch $F_{\pm}(\lambda)$ decreases to F_0 (see Figure 4), giving rise a constant branch in $F(\lambda)$. The *physical* $F(\lambda)$ thus becomes continuously degenerate along a compact interval (extending between two adjacent minima of $F_{+}(\lambda)$ or $F_{-}(\lambda)$) where F_0 dominates the growth of $G(\lambda|t)$. The boundaries of \mathcal{S}_0 , marking transitions between F_{\pm} and F_0 , are critical points associated with a second-order DPT.

The main distinction with the first-order transitions is that $F(\lambda)$ now remains differentiable everywhere, including at the two second-order critical points. Since $F'_0 = 0$ and $F''(\lambda) > 0$, the second derivatives however experience a discontinuity at the minima. This time (unlike in the case of first-order DPTs) differentiability of $F(\lambda)$ ensures that its Legendre dual $F^*(j) = \sup_{\lambda \in \mathbb{R}} \{\lambda j - F(\lambda)\}$ coincides with the LD rate function, $F^*(j) = I(j)$. Recall that upon performing a Legendre transform of $F(\lambda)$, any flat or affine part with slope j_{aff} is mapped to a *single* point $I(j_{\text{aff}})$. The Legendre counterpart of $F(\lambda)$ with a flat segment will thus feature a *corner* at the origin $j = 0$. This further means that the pair of dynamical critical points associated with the second-order DPT manifests itself as an isolated non-differentiable point in the rate function. Following Ehrenfest's classification scheme, such points would correspond to critical point of first order. In this work, we follow the 'canonical' terminology of criticality and classify phase transitions in terms of differentiability (or lack thereof) of the dynamical free energy (sometimes referred to as ' λ -ensembles', as e.g. in Ref. [89]).


Dynamical regimes

By continuously changing the counting field along the real λ -axis, dynamical phases $F_{+}(\lambda)$, $F_{-}(\lambda)$ and F_0 shown, respectively, as red, blue and gray curves in the left column figures and the corresponding rate functions $I_{+}(j)$, $I_{-}(j)$, shown as red and blue curves in the right column figures, display different interweaving patterns. We shall explicitly distinguish between *four* different scenarios which we hereafter refer to as 'dynamical regimes':


- **regular regime:** one of the convex bulk branches $F_{\pm}(\lambda)$ dominates over the entire physical range of counting fields $\lambda \in \mathbb{R}$. Correspondingly, the LD rate function is Legendre dual to $F(\lambda)$, $I(j) = F^*(j)$, and involves a single physical (strictly convex) branch $I(j) = I^{(\pm)}(j)$.

We can offer another, perhaps more physically suggestive, perspective on the emergence of a flat part. The constant branch F_0 dominates along an interval \mathcal{S}_0 , signifying that the main contributions to $G(\lambda \in \mathcal{S}_0|t)$ at late times are due to phase-space trajectories that differ in a *subextensive* (i.e. for J_i on scales asymptotically smaller than $\sim t^{\alpha}$) amount of transported charge. For any finite large current $j \neq 0$, the rate function $I_c(j)$ is instead differentiable and consequently the relevant rare trajectories concentrating around the maximum of MGF $G(\lambda|t)$ carry integrated currents of the order $\mathcal{O}(j t^{\alpha})$. By contrast, the subextensive rare events associated to the second-order criticality are not associated with bulk-extremum contributions but rather stem from the global maxima at the boundary of the integration domain (see Sec. V). Despite being exponentially unlikely, with a probability decaying with a rate of $I_0(j) = -F_0 > 0$, one would need to look at subleading orders in time to gain further insight into the finer structure of such trajectories.


Lastly, we examine the validity of the univariate fluctuation relation. We need to explicitly distinguish between the following two cases (i) the flat part connect both the degenerate minima $\lambda_{\pm}^{(\pm)}$ on the same bulk branch F_{\pm} or (ii) F_0 interpolates between two degenerate minima of different bulk branches. In case (i), the UFR for the charge LDF $I(j)$ remains intact (provided that F_{\pm} individually obey the symmetry), as evidently both degenerate minima appear symmetrically with respect to the inversion point $\lambda_{\pm}^{(0)} = -\tilde{\varepsilon}_c/2$, irrespective of the extent of the flat branch F_0 . Analogously, the appearance of a flat part preserves the UFR of the LDF in spite of a corner at $j = 0$. The situation is different if the flat part connects two degenerate minima on different bulk branches. Even when F_{\pm} both have inversion points $\lambda = -\tilde{\varepsilon}_c^{(\pm)}/2$, the UFR ceases to hold simply because the two reflection points in general do not coincide, $\tilde{\varepsilon}_c^{(+)} \neq \tilde{\varepsilon}_c^{(-)}$. In this case, the presence of F_0 spoils the inversion symmetry of $F(\lambda)$. What is less obvious is that there are no direct transitions from regime (i) to (ii), or vice-versa, but only via the regime that features a first-order criticality. Finally, we also mention that out of equilibrium with uniform particle density, $\rho_{-} = \rho_{+}$, and arbitrary charge biases b_{\pm} , the UFR is always violated.



- **corner regime:** there is an exchange of dominance between the bulk branches $F_{\pm}(\lambda)$. Their respective regions of dominance \mathcal{S}_{\pm} meet at the critical point $\lambda_{>} \in \mathbb{R}$ where the physical SCGF $F(\lambda)$ develops a non-differentiable (corner) point. The latter is the critical point of a *first-order* dynamical phase transition. The rate function $I(j)$ is no longer the Legendre transform of $F(\lambda)$ but instead a *non-convex* function with a non-differentiable (corner) points at the critical large current $j_{>}$, corresponding to the minimum of the two branches $I(j) = \min_j \{I^{\pm}(j)\}$.



- **tunneling regime:** the flat branch F_0 sets in, arising from a continuous symmetry-breaking transition of a convex regular branch into a non-convex form with a doubly degenerate minimum. The flat part of $F(\lambda)$ connects between two degenerate minima of symmetry-broken branches F_{\pm} . We call this phenomenon ‘tunneling’ (and symbolize it by \asymp). There are two subregimes: the tunneling transition via F_0 connecting two degenerate minima of the same branch F_{\pm} , labeled by $\llbracket \pm \asymp \pm \rrbracket$, and the transition connecting the left minimum of F_{\pm} to the right minimum of another branch F_{\mp} , labeled by $\llbracket \pm \asymp \mp \rrbracket$. Every transition between the minima of F_{\pm} and F_0 is a dynamical phase transitions of second order. In both subregimes, the LD rate function $I(j) = \min_j \{I_{\pm}(j)\}$, each of which has a non-differentiable point (corner) at $j = 0$.



- **mixed regime:** apart from tunneling between two degenerate minima of the same bulk branch F_{\pm} , there is a transfer to another branch via a non-differentiable corner point, or in the opposite order. Intersection of bulk branches F_{\pm} can never coexist with tunneling to another branch.

Recap. Before detailing out various formal aspects of the outlined dynamical criticality, we take an opportunity to succinctly summarize the key findings and state the general conclusions:

- ♣ The single-file property (I) combined with inertness of charge (II) implies fragmentation of the classical phase space, foliating into exponentially many sectors characterized by conserved charge patterns. Dynamical systems of this type display strongly non-ergodic behavior manifested through the competition of dynamical phases. For a given value of the counting field $\lambda \in \mathbb{R}$, the finite-time CGF $F(\lambda|t)$ receives contributions from both branches $F_{\pm}(\lambda)$ attributed to each of the two partitions in the initial non-stationary state. While away from equilibrium only one of them dominates the growth of MGF at late times, this delicately depends on the average value of particle and charge densities characterizing initial non-stationary states. An exchange of dominance between $F_+(\lambda)$ and $F_-(\lambda)$ introduces a non-differentiable (corner) point in the SCGF $F(\lambda)$, signaling a DPT of the first order. In this event, the Gallavotti–Cohen relation breaks for large currents $|j| > |j_{\bowtie}|$, but survives for subcritical values $|j| < |j_{\bowtie}|$. The large-deviation rate function $I(j)$ corresponds to taking the minimum of two convex branches $I_{\pm}(j)$.
- ♣ There exists a region in the parameter space of initial bipartitioned states where a new constant branch F_0 emerges as a part of the physical SCGF $F(\lambda)$. The flat segments arise when the physical branch experiences a symmetry-breaking phase transition into a non-convex form. The flat branch interpolates between two degenerate minima – either of the same symmetry-broken branch $F_{\pm}(\lambda)$ or two adjacent minima of the opposite branches – along a compact interval \mathcal{S}_0 where it dominates over $F_{\pm}(\lambda)$. While transitions from the bulk branches $F_{\pm}(\lambda)$ to the flat branch F_0 (or vice-versa) do not spoil differentiability of $F(\lambda)$, the second derivative $F''(\lambda)$ features a discontinuity at the minima of $F_{\pm}(\lambda)$. The boundaries of \mathcal{S}_0 are accordingly interpreted as critical points of a second order DPT, transcribing into a single non-differentiable (corner) point in the associated LDF $I(j)$ at $j = 0$.
- ◆ Depending on the presence and type of dynamical criticality, there are four qualitatively distinct dynamical regimes, dubbed as **regular** (of type $\llbracket + \rrbracket$ or $\llbracket - \rrbracket$), **tunneling** (of types $\llbracket \pm \asymp \pm \rrbracket$ or $\llbracket \pm \asymp \mp \rrbracket$), **corner** $\llbracket \pm \bowtie \mp \rrbracket$ and finally **mixed** $\llbracket \pm \bowtie \pm \asymp \mp \rrbracket$. These four regimes provide the full partitioning of the parameter space. The UFR of the SCGF $F(\lambda)$ is globally preserved in $\llbracket \pm \rrbracket$ and $\llbracket \pm \asymp \pm \rrbracket$ regimes, locally preserved for subcritical large currents in $\llbracket \pm \bowtie \mp \rrbracket$ regime and fully violated in $\llbracket \pm \asymp \mp \rrbracket$ regime.

A. Large fluctuations: the dressing formalism

We now describe the *dressing procedure* that permits one to compute the charge-current rate function from that of the particle-current. This can be achieved, remarkably, without resorting to any model-specific input. For technical reasons, we shall only assume certain minimal ‘regularity properties’ on the particle rate function. We first outline the procedure at the level of the rate function by expressing the charge-current rate function as the solution to a convex optimization problem (53) and systematically examine its structure. Here we only provide a succinct summary and postpone a detailed analysis to Sec. V.

1. Dressing the particle rate function

By prescribing $\mathcal{P}_p(\mathbf{J}_p|t)$, the joint PDF $\mathcal{P}_{c,p}(\mathbf{J}_p, \mathbf{J}_c|t)$ can be computed with aid of the conditional PDF $\mathcal{P}_{c|p}(\mathbf{J}_c|\mathbf{J}_p)$ according to the main axiom of probability,

$$\mathcal{P}_{c,p}(\mathbf{J}_c, \mathbf{J}_p|t) = \mathcal{P}_{c|p}(\mathbf{J}_c|\mathbf{J}_p)\mathcal{P}_p(\mathbf{J}_p|t). \quad (31)$$

This can be formally viewed as an operator (with the kernel $\mathcal{P}_{c|p}(\mathbf{J}_c|\mathbf{J}_p)$) which we regard as the dressing operator $\mathfrak{D}_{\mathcal{P}}$, namely

$$\mathcal{P}_{c,p}(\mathbf{J}_c, \mathbf{J}_p|t) = \mathfrak{D}_{\mathcal{P}}[\mathcal{P}_p(\mathbf{J}_p|t)](\mathbf{J}_c). \quad (32)$$

The univariate PDF $\mathcal{P}(\mathbf{J}_c|t)$ can be obtained by integrating out the particle current,

$$\mathcal{P}_c(\mathbf{J}_c|t) = \int d\mathbf{J}_p \mathcal{P}_{c,p}(\mathbf{J}_c, \mathbf{J}_p). \quad (33)$$

Since the assignment of internal charge degrees of freedom is, by virtue of the inertness property, uncorrelated with particles’ positions, $\mathcal{P}_{c|p}(\mathbf{J}_c|\mathbf{J}_p)$ is indeed merely a combinatorial factor. We shall suggestively refer to it as the ‘‘dressing factor’’. Below we compute its general form for the class of bipartitioned grand-canonical ensembles.

We begin by introducing a few auxiliary objects. For simplicity, we assume in the following that space is discrete. Fixing a window of time $[0, t]$, we denote by Λ_{\pm} the sublattices occupied by particles at initial time $t = 0$ that have crossed the origin during that time interval, with $n_- = |\Lambda_-|$ particles initially in the left subsystem and similarly $n_+ = |\Lambda_+|$ particles in the right subsystem. Moreover, we denote by $p_{L,\pm} \equiv (1 \pm b_-)/2$ and $p_{R,\pm} \equiv (1 \pm b_+)/2$ the probabilities of finding a charge + or – in the left (L) and right (R) partitions, respectively. Combinatorial counting yields

$$\begin{aligned} \mathcal{P}_{c|p}(\mathbf{J}_c|\mathbf{J}_p) &= \sum_{n_-, n_+ = 0}^{\infty} \delta_{n_- - n_+, \mathbf{J}_p} \sum_{m_-}^{n_-} \sum_{m_+}^{n_+} \binom{n_-}{m_-} \binom{n_+}{m_+} \\ &\times p_{L,+}^{m_-} p_{L,-}^{n_- - m_-} p_{R,-}^{m_+} p_{R,+}^{n_+ - m_+} \delta_{2m_- - n_-, \mathbf{J}_c}, \quad (34) \end{aligned}$$

where the outer double summation over n_- and n_+ goes over all possible combinations of crossings, while the inner double summation counts over all the colorings of charges. Note that we have also incorporated the Kronecker δ -constraint to ensure the difference $n_- - n_+ = J_p$, with $n \equiv n_+ + n_-$, $m \equiv m_+ + m_-$.

By further demanding the single-file property (1), the general form of Eq. (34) simplifies significantly. The central observation is that *at most one* of the sets Λ_\pm is non-empty which, after explicitly resolving the Kronecker constraints, brings us to a far simpler, factorizable expression for the conditional probability,

$$\mathcal{P}_{c|p}(J_c|J_p) = \mathcal{P}_{c|p}^{[0]}(J_c|J_p) \mathcal{B}(J_p, J_c). \quad (35)$$

Largely for convenience, we have separated out the conditional probability in the absence of biases ($b_\pm = 0$),

$$\mathcal{P}_{c|p}^{[0]}(J_c|J_p) \equiv 2^{-|J_p|} \left(\frac{|J_p|}{|J_p| + J_c} \right), \quad (36)$$

and introduced the ‘biasing factor’ $\mathcal{B}(J_p, J_c)$ of the form

$$\mathcal{B}(J_p, J_c) \equiv 2^{|J_p|} \left(\frac{1 - b_\zeta^2}{4} \right)^{|J_p|/2} \left(\frac{1 - b_\zeta}{1 + b_\zeta} \right)^{\zeta J_c/2}, \quad (37)$$

depending on bias parameters b_\pm implicitly through the direction of the integrated particle current J_p via signature $\zeta \equiv -\text{sgn}(J_p)$. Owing to the single-file property, positive (negative) number of transferred particles are associated with the negative (positive) partitions.

Large fluctuations. We are now in a position to infer the exact LDF of the transferred charge from the asymptotic behavior of the dressing kernel $\mathcal{P}_{c|p}(J_c|J_p)$. To this end, we first pick two arbitrary dynamical exponents ζ_i in the range $1/2z_i \leq \zeta_i \leq \alpha_i$ (with $i \in \{p, c\}$) and introduce the corresponding rescaled cumulative currents, $\mathcal{J}_i(t) = t^{-\zeta_i} J_i(t)$. In terms of the rescaled currents, we have the following asymptotic formula (abusing notation for PDFs with scaled arguments)

$$t^{-\zeta_c} \mathcal{P}_{c|p}(j_c|t) \asymp \int dj_p \mathcal{P}_{c|p}(j_c|j_p) \mathcal{P}_{\zeta_p}(j_p|t). \quad (38)$$

We are mainly interested in asymptotic behavior associated with the *largest* timescale $\zeta_p = \alpha_p \equiv \alpha$, pertaining to rare space-time trajectories in which the transferred particle number scales asymptotically as $\sim t^\alpha$. We remind the reader that the timescale α is fixed by the rate of growth of $G_p(\lambda|t) \equiv \langle e^{\lambda J_p(t)} \rangle_{\text{init}}$ at late times. Inertness of charge immediately implies that the net charge current carried by those rare events is of the same order, that is $\alpha_c = \alpha_p$. When we wish to infer the statistics of large charge-current fluctuations we therefore set $\zeta_c = \zeta_p = \alpha$.

In the following, we further make the following assumption on the cumulative particle current $J_p(t)$:

- (a) $J_p(t)$ obeys the LD principle on the *large* timescale α , with the SCGF given by

$$F_p(\lambda_p) = \lim_{t \rightarrow \infty} t^{-\alpha} \log G_p(\lambda_p|t). \quad (39)$$

Let moreover $\mathcal{P}_\alpha(\mathcal{J}_p(t)|t)$ denote the PDF associated to the rescaled time-integrated particle current $\mathcal{J}_p(t) = t^{-\alpha} J_p(t)$. At late times, the probability for observing a value j_p of the rescaled cumulative particle current $\mathcal{J}_p(t)$ is characterized by the rate function

$$I_p(j_p) = - \lim_{t \rightarrow \infty} t^{-\alpha} \log \mathcal{P}_\alpha(\mathcal{J}_p = j_p|t). \quad (40)$$

- (b) in equilibrium, *typical* fluctuations of the cumulative particle current $J_p(t)$, characterized by scaling exponent $\zeta_p = 1/2z_p$ (where $z_p = 1/\alpha$ is the algebraic dynamical exponent associated with the asymptotic temporal growth of the second cumulant of $G_p(\lambda_p|t)$), are *Gaussian* with zero mean and variance of σ_p^2 ,

$$\mathcal{P}_{1/2z_p}(j_p) = \frac{1}{\sqrt{2\pi\sigma_p^2}} \exp \left[-\frac{j_p^2}{2\sigma_p^2} \right]. \quad (41)$$

We proceed by approximating the binomial weights in Eq. (36) using the Stirling formula. To facilitate the computation, it is convenient to introduce a new dynamical variable

$$\xi(t) \equiv t^{\zeta_c - \zeta_p} \frac{j_c}{|j_p|}, \quad (42)$$

in terms of which the exact asymptotic expression for the conditional probability (suppressing subexponential terms) takes the form

$$\mathcal{P}_{c|p}^{[0]}(j_c|j_p) \asymp \exp \left[-t^{\zeta_p} |j_p| \Xi(\xi) \right], \quad (43)$$

with

$$\Xi(\xi) \equiv \frac{1}{2} \sum_{\epsilon \in \{\pm\}} [(1 + \epsilon \xi) \log(1 + \epsilon \xi)] - \log(2). \quad (44)$$

Similarly, the (rescaled) biasing weight, denoted hereafter by $\mathcal{B}(j_p, j_c)$, can be presented in a factorized form,

$$\mathcal{B}(j_p, j_c) \asymp \mathcal{B}_p(j_p) \mathcal{B}_c(j_c) \quad (45)$$

with

$$\mathcal{B}_p(j_p) = \exp \left[t^{\zeta_p} \frac{|j_p|}{2} \log \left(\frac{1 - b_\zeta^2}{4} \right) \right], \quad (46)$$

$$\mathcal{B}_c(j_c) = \exp \left[t^{\zeta_c} \frac{\zeta j_c}{2} \log \left(\frac{1 - b_\zeta}{1 + b_\zeta} \right) \right]. \quad (47)$$

We now observe that for moderate fluctuations associated with timescales $\zeta_c < \zeta_p$, only the lowest non-trivial

order in ξ in Eq. (43) remains relevant at late times, yielding a remarkably simple result

$$\mathcal{P}_{c|p}^{[0]}(j_c|j_p) \asymp \exp\left[-t^{2\zeta_c - \zeta_p} \frac{j_c^2}{|j_p|}\right]. \quad (48)$$

By contrast, in the case of large fluctuations $\zeta_c = \zeta_p$ we have $\xi \in \mathcal{O}(t^0)$, i.e. $\xi \equiv \xi(0) = j_c/|j_p|$ becomes independent of time. The charge-current univariate PDF $\mathcal{P}_c(\mathcal{J}_c|t)$ of the dynamically rescaled cumulative charge current $\mathcal{J}_c(t) = t^{-\alpha} J_c(t)$ is accordingly given by the following asymptotic expression (for compactness suppressing irrelevant subexponential terms in the integrand)

$$e^{-t^\alpha I_c(j_c)} \asymp t^{-\alpha/2} \int dj_p e^{-t^\alpha I_{c,p}(j_c, j_p)}. \quad (49)$$

The above expression can be viewed as marginalization of the *joint* rate function $I_{c,p}(j_c, j_p)$, see e.g. [119]. The latter can be naturally decomposed as

$$I_{c,p}(j_c, j_p) = I_{c|p}(j_c, j_p) + I_p(j_p), \quad (50)$$

where $I_p(j_p)$ is interpreted as the *marginal* rate function, while $I_{c|p}(j_c|j_p)$ is the *conditional rate function*

$$I_{c|p}(j_c|j_p) = \sum_{\epsilon \in \{\pm\}} \frac{|j_p|}{2} (1 + \epsilon \xi) \log \left[\frac{1 + \epsilon \xi}{1 - \epsilon \zeta b_\zeta} \right], \quad (51)$$

with signature $\zeta = -\text{sgn}(j_p)$.

In summary, provided the particle SCGF $F_p(\lambda_p)$ as an input, one can retrieve the joint LDF $I_{c,p}(j_c, j_p)$ and SCGF $F_{c,p}(\lambda_c, \lambda_p)$ via the following sequence of explicit transformations

$$F_p(\lambda_p) \xrightarrow{*} I_p(j_p) \xrightarrow{\mathfrak{D}_p} I_{c,p}(j_c, j_p) \xrightarrow{*} F_{c,p}(\lambda_c, \lambda_p) \quad (52)$$

where the action of \mathfrak{D}_p on a rate function is given by Eq. (50). Moreover, if $F_p(\lambda_p)$ fulfills the assumptions of the Gärtner–Ellis theorem, the corresponding rate function $I_p(j_p)$ is simply given by the Legendre transform of the particles SCGF $F_p(\lambda)$, i.e. $I_p(j_p) = F_p^*(j_p)$.

The univariate charge-current LDF $I_c(j_c)$ can be straightforwardly retrieved by marginalization. By invoking the Laplace principle, in the large-time limit $t \rightarrow \infty$ the integral (49) localizes around the extremum, implying

$$I_c(j_c) = \inf_{j_p} I_{c,p}(j_c, j_p). \quad (53)$$

To finally obtain the SCGF $F_c(\lambda)$ one can make use of the Legendre–Fenchel transform,

$$F_c(\lambda) = I_c^*(\lambda) \equiv \sup_{j_c} \{\lambda j_c - I_c(j_c)\}. \quad (54)$$

2. Dressing the moment generating function

The dressing procedure described in Sec. III A 1 can be alternatively formulated at the level of the moment generating functions. Here we derive a simple correspondence between the finite-time particle-current MGF $G_p(\lambda_p|t)$ and the joint particle-charge MGF $G_{c,p}(\lambda_c, \lambda_p|t)$. In the following computations, we employ the multiplicative counting fields $z_p \equiv e^{\lambda_p}$ and $z_c \equiv e^{\lambda_c}$ and assume that the integrated currents J_p, J_c take only integer values (as is the case for point particle or discrete variable systems).

Computing the joint *finite-time* MGF $G_{c,p}(\lambda_c, \lambda_p|t)$ amounts to acting with the dressing operator \mathfrak{D}_G ,

$$G_{c,p}(z_c, z_p|t) = \mathfrak{D}_G[G_p(z_p|t)], \quad (55)$$

on the particle MFG

$$G_p(z_p|t) = \langle z_p^{J_p(t)} \rangle = \mathfrak{L}[\mathcal{P}(J_p|t)](z_p^{-1}). \quad (56)$$

The dressing operator can be most conveniently expressed as a composition $\mathfrak{D}_G = \mathfrak{L} \circ \mathfrak{D}_p \circ \mathfrak{L}^{-1}$, representing conjugation of $\mathcal{P}(J_c|t)$ by the *bilateral* Laplace transform \mathfrak{L} ,

$$\mathfrak{L}[\mathcal{P}(J_p|t)](z_p^{-1}) \equiv \int dJ_p z_p^{J_p} \mathcal{P}(J_p|t), \quad (57)$$

whose inverse satisfies $\mathcal{P}(J_p|t) = \mathfrak{L}^{-1}[G_p(z_p|t)](J_p)$.

Evaluating the action of the dressing operator (55) requires a few technical steps which are spelled out in Appendix A. There we demonstrate that acting with \mathfrak{D}_G corresponds to applying the following simple *substitution rule*:

$$G_{c,p}(z_c, z_p|t) = G_p(z_p|t) \Big|_{z_p^{\pm n} \rightarrow z_p^{\pm n} [\mu_{\pm}(z_c)]^n}, \quad (58)$$

with the ‘dressed counting fields’

$$\mu_{\pm}(z_c) = \frac{1}{2}(z_c + z_c^{-1}) \mp b_{\pm} \frac{1}{2}(z_c - z_c^{-1}). \quad (59)$$

In summary, we have thus established that

the finite-time joint MGF $G_{c,p}(z_c, z_p|t)$ is given by the Laurent series expansion of the particle MGF $G_p(z_p|t)$ upon multiplying all positive and negative integral powers of (exponential) counting fields $z_p^{\pm n}$ by the corresponding dressed counting fields $[\mu_{\pm}(z_c)]^n$.

B. Universal anomalous fluctuations in equilibrium

In this section, we consider the univariate PDF of the cumulative charge current rescaled to the timescale of typical fluctuations $\zeta_c = 1/2z_c$,

$$\mathcal{P}_c^{\text{tYP}}(j_c) = \lim_{t \rightarrow \infty} t^{-\zeta_c} \mathcal{P}_{\zeta_c=1/2z_c}(\mathcal{J}_c|t). \quad (60)$$

We shall now establish the following remarkable property: in *equilibrium* ensembles with finite particle density and without charge bias ($b = 0$), the PDF $\mathcal{P}_c^{\text{typ}}(j_c)$ takes a universal *non-Gaussian* form in spite of detailed balance. This property has been previously observed and explained in our recent paper [74], where we computed the FCS for an exactly solvable classical automaton of hardcore charged particles. In Sec. IV we revisit the model and compute the FCS of charge transfer with respect to non-stationary bipartitioned initial states.

We wish to stress that the observed anomalous fluctuations found in unbiased equilibrium ensembles are a general feature of dynamical systems that are subjected to the constraints specified in Sec. II. In other words, absence of the so-called CLT property is a corollary of the imposed constraints, namely (I) the single-file property and (II) inertness of charge.

We can infer directly from Eqs. (38) and Eq. (48) that convergence of the rescaled charge-current PDF $t^{-z_c} \mathcal{P}_c(\mathcal{J}_c|t)$ towards a non-trivial *stationary* PDF can be achieved only provided that the particle and charge dynamical exponents obey

$$z_c = 2z_p. \quad (61)$$

We have thus inferred that inert charges are slowed down and spread through the system on a timescale given by the square root of that associated with particle transport.

The PDF $\mathcal{P}_c^{\text{typ}}(j_c)$ takes the *universal* form with the following integral representation

$$\mathcal{P}_c^{\text{typ}}(j_c) = \frac{1}{\pi\sigma_p} \int_{-\infty}^{\infty} \frac{dj_p}{\sqrt{|j_p|}} \exp\left[-\frac{j_p^2}{2\sigma_p^2} - \frac{j_c^2}{2|j_p|}\right]. \quad (62)$$

The corresponding MGF $G_c^{\text{typ}}(\eta)$ is given by the bilateral Laplace transform of $\mathcal{P}_c^{\text{typ}}(j_c)$, namely $G_c^{\text{typ}}(\eta) = \mathcal{L}[\mathcal{P}_c^{\text{typ}}(j_c)](\eta) = \int_{-\infty}^{\infty} dj_c e^{-\eta j_c} \mathcal{P}_c^{\text{typ}}(j_c)$, yielding the following integral representation

$$G_c^{\text{typ}}(\eta) = \int_{-\infty}^{\infty} dy \frac{2|y|}{\sqrt{\pi}\sigma_p} \exp\left[-\frac{y^4}{2\sigma_p^2} + \frac{\eta^2 y^2}{2}\right]. \quad (63)$$

Switching to a new variable $u = y^2$, splitting the integral into two separate integrals over the real semi-axes \mathbb{R}_{\pm} , and using the identity $\int_0^{\infty} du \pi^{-1/2} \exp[vu - (u/2)^2] = \exp(v^2) \text{erfc}(v)$, we arrive at the compact explicit expression

$$G_c^{\text{typ}}(\eta) = E_{1/2}\left(\frac{\eta^2 \sigma_p}{4}\right), \quad (64)$$

where $E_{1/2}(y)$ belong to a one-parameter family of Mittag-Leffler functions $E_a(y)$ (see e.g. [120])

$$E_a(y) = \sum_{n \geq 0} \frac{y^n}{\Gamma(an + 1)}, \quad (65)$$

widespread in applications of fractional calculus [121, 122] (Γ denotes Euler's Gamma function), while $\sigma_p \equiv \sqrt{2}\sigma_p$ sets the characteristic width. In particular, the second cumulant κ_2 of $\mathcal{P}_c^{\text{typ}}(j_c)$ equals $\kappa_2 = \sqrt{2/\pi}\sigma_p$. For example, in the hardcore automaton $\sigma_p(\rho) = \sqrt{\rho(1-\rho)}$.

Since $E_a(y)$ are entire functions, their inverse (bilateral) Laplace transform is essentially the Fourier transform, yielding $\mathcal{L}^{-1}[E_{2a}(y^2)] = \frac{1}{2}M_a(|y|)$. Here $M_a(y)$ denote a one-parameter family of PDFs known by the name of *symmetrized M-Wright function* [123],

$$M_a(y) = \sum_{n \geq 0} \frac{(-y)^n}{n! \Gamma((1-a) - na)}, \quad (66)$$

belonging to a subfamily of special functions called Wright functions. The final result is a closed-form universal expression for the PDF,

$$\mathcal{P}_c^{\text{typ}}(j_c) = \frac{1}{\sqrt{\sigma_p}} M_{1/4}\left(\frac{2|j_c|}{\sqrt{\sigma_p}}\right), \quad (67)$$

shown in Fig. 5. The M-Wright function of the scaling variable $2J_c/\sqrt{\sigma_p}t^{1/2z_c}$ indeed plays the role of the Green's function of the Cauchy problem associated with the time-fractional (in Caputo sense) diffusion equation $\partial^\beta \psi / \partial t^\beta = D_\beta \partial^2 \psi / \partial x^2$ of fractional order $\beta = 2/z_c$. The analogy is not exact, however; in the above PDF, the index of the function is always (i.e. irrespective of exponent z_p) equal to $z_p/(2z_c) = 1/4$. This ratio is presently uniquely fixed by demanding time-stationarity of the appropriately rescaled dynamical PDF $\mathcal{P}_c(\mathcal{J}_c|t)$.




FIG. 5. Time-dependent probability density functions $t^{1/2z_c} \mathcal{P}_c(\mathcal{J}_c|t)$ (colored dashed curves, computed using Eqs. (31), (34) and a Gaussian distribution of J_p) of the rescaled cumulative charge current $\mathcal{J}_c(t) = t^{-1/2z_c} J_c(t)$ associated with typical fluctuations with $z_c = 4$, shown for the two-species simple symmetric exclusion process with parallel update rule at different times (displayed on normal and logarithmic scale). Solid black curves are the theoretically predicted M-Wright distribution corresponding to the stationary PDF $\mathcal{P}_c^{\text{typ}}(j_c) = \lim_{t \rightarrow \infty} t^{1/8} \mathcal{P}(\mathcal{J}_c|t)$.

C. Dynamical phases

In this section, we take a closer look at the univariate SCGF $F_c(\lambda_c) \equiv F(\lambda)$. Given that the function $F(\lambda)$ governs the asymptotic growth of the charge-current MGF $G_c(\lambda|t) \equiv G(\lambda|t)$, we may regard it – *mutatis mutandis* – as a dynamical analogue of (thermodynamic) free energy. Supplying the particle univariate MGF $G_p(\lambda|t)$, the exact $G(\lambda|t)$ is obtained by substituting the counting fields $\exp(\pm n\lambda)$ with their biased counterparts $(\mu_{\pm}(\lambda))^n$, see Eq. (58). The task boils down to inferring the asymptotic behavior of $G(\lambda|t)$, by e.g. applying the methods of localization or steepest descent. To this end, one first replaces the discrete summation and combinatorial weights with appropriate continuous counterparts, yielding a two-dimensional integral over a (possibly non-compact) integration domain \mathcal{D} . Then, we take into account that *for asymptotically large times, the integral concentrates about the extremal points (local maxima) in the integration domain \mathcal{D}* . In general, we have to distinguish between two types of extrema: (i) local maxima in the interior of the domain \mathcal{D} , and (ii) maxima located at the boundary $\partial\mathcal{D}$.

Whenever the MFG $G(\lambda|t)$ receives contributions from multiple extrema, we may asymptotically approximate it by

$$G(\lambda|t) \asymp \sum_{k=1}^K W_k(\lambda) e^{t^\alpha F_k(\lambda)}. \quad (68)$$

Here $k = 1, \dots, K$ enumerates all (possibly degenerate) critical points (including the boundary extrema), while expansion coefficients $W_k(\lambda)$ will be referred to as weighting functions. Functions $F_k(\lambda)$ in the exponents can be suggestively interpreted as coexisting (stable or metastable) *branches* of the dynamical free energy $F(\lambda)$.

By fixing the value of the counting field λ , a single branch $F_k(\lambda)$ eventually dominates at late times (save for degenerate cases which we neglect for the time being). All other subdominant branches are only visible in the transient dynamics (in the form of corrections that are exponentially suppressed as $t \rightarrow \infty$). Nothing however prevent a subdominant branch from taking over the dominant one upon varying the counting field. It is thus conceivable that by virtue of multiple coexisting competing dynamical phases, different branches dominate the late-time growth of $G(\lambda|t)$, depending of the value of the counting field λ . As a matter of fact, such behavior may depend on the parameters of the initial state.

Allowing for the most general scenario, we can envisage a partitioning of the *complex* λ -plane $\cup_{k=1}^K \mathcal{R}_k \cong \mathbb{C}$, where $\mathcal{R}_k \in \mathbb{C}$ denote the corresponding ‘regions of dominance’. If we are only interested in reconstructing the LD rate function $I(j)$, it suffices to compute $F(\lambda)$ along the real λ -axis (with an extra technical assumption that that $F(\lambda)$ is everywhere differentiable). Let accordingly \mathcal{S}_k denote the regions of dominance pertaining to $F_k(\lambda)$. Intervals \mathcal{S}_k are given by intersections of \mathcal{R}_k with the

real λ -axis, $\cup_{k=1}^K \mathcal{S}_k \cong \mathbb{R}$. Certain \mathcal{S}_k can be empty, attributed to those $F_k(\lambda)$ that remain subdominant for all values of $\lambda \in \mathbb{R}$. The structure of the associated LD rate function can be, as it turns out, quite non-trivial; although there are K dynamical phases involved, the total number of distinct sequences of interweaving branches $F_k(\lambda)$ in the direction of the (fictitious) ‘ λ -flow’ (see Fig. 6 for an illustration), can be much larger.

Transitions between dynamical phases occur at the boundaries of intervals \mathcal{S}_k , identified with dynamical critical points. While the SCGF $F(\lambda)$ is a continuous and everywhere convex function of $\lambda \in \mathbb{R}$, it will in general feature non-analytic behavior at the critical points. Judging from the asymptotic form (68) of the multi-branched structure of the MGF, we anticipate DPTs of the *first order* with one of several critical points λ_{\triangleright} . Such point show up as *non-differentiable* (corner) point in the physical $F(\lambda)$. We remind the reader that Legendre duality is no longer guaranteed to hold for any SCGF $F(\lambda)$ that develops a non-differentiable point [43]; it is only when $F(\lambda)$ is differentiable everywhere on its domain that there is a guarantee (in the form of the Gärtner–Ellis theorem) that the Legendre transform of $F(\lambda)$ yields the large-deviation rate function, i.e. that $I(j) = F^*(j)$. Otherwise, $F^*(j)$ is merely the convex hull of $I(j)$.

As we explain next, exploiting additional analytic input stored in the weighting functions $W_k(z)$ provides an elegant route that bypasses the need for the Gärtner–Ellis theorem or a fully-fledged steepest descent analysis. By adopting the asymptotic representation (68) as a starting point, the inverse Laplace transform can be computed for each branch individually as follows [124]: by decomposing the PDF $\mathcal{P}(j|t) = \sum_k \mathcal{P}_k(j|t)$ and introducing potentials $\Phi_k(\lambda; j) \equiv \lambda j - F_k(\lambda)$, each term is asymptotically of the form

$$\mathcal{P}_k(j|t) \asymp \frac{1}{2\pi i} \int_{\mathcal{D}_k} d\lambda W_k(\lambda) e^{t^\alpha \Phi_k(\lambda; j)}, \quad (69)$$

where \mathcal{D}_k denote *deformed* integration (Bromwich) contours passing through saddle points λ_k^* of $\Phi_k(\lambda; j)$, given by the unique solutions to $F_k'(\lambda) = j$. We have assumed that functions $W_k(\lambda)$ are either free of poles (in the complex λ -plane) or that in the process of deforming the original Bromwich integration contours to \mathcal{D}_k we have not crossed any poles. In the opposite case, the asymptotics of $\mathcal{P}_k(j|t)$ would pick up additional contributions stemming from the residues of the integrand in Eq. (69) (see Ref. [124] for more details). At late times, we thus have

$$\mathcal{P}_k(j|t) \asymp e^{t^\alpha \Phi_k(\lambda_k^*; j)}, \quad (70)$$

and the rate function $I(j) = -\lim_{t \rightarrow \infty} t^{-\alpha} \log \mathcal{P}(j|t)$ is simply the infimum over the branches

$$I(j) = \inf_k \{\Phi_k(\lambda_k^*; j)\}. \quad (71)$$

We note that a similar type of large-deviation rate functions describe mixtures of Bernoulli trials [125].

IV. HARDCORE CELLULAR AUTOMATON

Having finally put the formal framework in place, we now turn to practical applications and exemplify how the outlined techniques can be applied to a specific model. For demonstrative purposes, we present the full solution of the classical deterministic time-reversible classical cellular automaton introduced in Sec. II.

We devote the remainder of the paper to a comprehensive analysis of charge fluctuations *away* from equilibrium, thereby expanding on our previous work [74]. We first briefly present, as a warm-up, how to infer the full counting statistics associated with particle number. By applying the dressing method, we then compute the exact statistics of charge and exhibit its most salient features. Finally, we carry out a systematic analysis of various dynamical phases and their interweaving patterns by studying the corresponding phase diagrams in the complex λ -plane within the scope of Lee–Yang theory.

A. Particle fluctuations

We consider an inhomogeneous initial state consisting of two semi-infinite partitions, each initialized in a grand-canonical equilibrium state with respective particles densities ρ_- and ρ_+ . For later convenience, we introduce the ratio of hole to particle densities, $\nu = \bar{\rho}/\rho$. The latter obeys $1/\nu = \rho^{-1} \partial f_{\text{eq}}(\rho)/\partial \rho$, where $f_{\text{eq}} = \log(1 + e^{\beta \nu})$ is the free-energy density parametrized by chemical potential $\beta_p = \log(1/\nu)$. The static susceptibility accordingly reads $\chi_p(\rho) = \partial^2 f_{\text{eq}}(\rho)/\partial \beta_p^2 = \rho(1 - \rho)$.

To compute fluctuations of net particle transfer, we may simply ignore the charge degrees of freedom. In the hardcore automaton, this effectively eliminates interaction among the particles, and hence computing the corresponding FCS becomes a simple exercise; in the absence of relaxation, fluctuations can be read off directly from the initial condition, yielding the univariate SCGF of the form

$$F_p(\lambda_p) = \log [(\rho_+ + e^{\lambda_p} \bar{\rho}_+)(\rho_- + e^{-\lambda_p} \bar{\rho}_-)], \quad (72)$$

associated with a sum of independent Bernoulli processes characterized by state densities ρ_{\pm} . The result is in agreement with the celebrated Levitov–Lesovik formula [126, 127] (specialized for the case of perfect transmission). Since particles propagate freely, the MGF $G_p(\lambda_p|t) = \exp(t F_p(\lambda_p))$ is indeed exact at all times and not just asymptotically. Moreover, $F_p(\lambda_p)$ depends only on a single parameter

$$\varpi \equiv \rho_+ \bar{\rho}_- (e^{-\lambda_p} - 1) + \rho_- \bar{\rho}_+ (e^{\lambda_p} - 1), \quad (73)$$

in terms of which it reads $F_p(\lambda_p) = \log(1 + \varpi(\lambda_p))$. It is instructive to compare this result with the SCGF of the simple symmetric exclusion process, reading [109] $F_{\text{SSEP}}(\lambda_p) = \frac{1}{\pi} \int_{-\infty}^{\infty} du \log[1 + \varpi(\lambda_p) \exp(-u^2)]$.

The first two scaled cumulants as functions of particle densities read

$$s_1^{(p)} = \lim_{t \rightarrow \infty} \frac{c_1^{(p)}(t)}{t} = \bar{j}_p = \rho_- - \rho_+, \quad (74)$$

$$s_2^{(p)} = \lim_{t \rightarrow \infty} \frac{c_2^{(p)}(t)}{t} = \rho_- \bar{\rho}_- + \rho_+ \bar{\rho}_+. \quad (75)$$

Setting $\rho_- - \rho_+ \equiv \delta\rho$ and expanding $F_p(\lambda)$ around equilibrium, $\delta\rho \rightarrow 0$, we can extract ‘diffusivity’ $D_*(\rho) = 1$ and ‘mobility’ $\sigma_*(\rho) = 2\rho(1 - \rho)$, obeying the local Einstein relation $2\chi_p(\rho)D_*(\rho) = \sigma_*(\rho)$. Curiously, we have retrieved the exact same dependence as found in the SSEP (on an infinite line, see Ref. [109]), with the proviso that SSEP is a diffusive system ($\alpha = 1/2$) while free particles are ballistic ($\alpha = 1$). Beware not to confuse $\sigma_*(\rho)$ with conductivity $\sigma(\rho)$; in ballistic systems, the second cumulant instead gives the first absolute moment of the DSF called the Drude self-weight $s_2^{(p)} = \lim_{t \rightarrow \infty} \int_{-t}^t d\tau \langle j_p(0, \tau) j_p(0, 0) \rangle^c$, see Ref. [128].

Fluctuations of the particle transfer evidently satisfy the GCR. In analogy with the SSEP, the fluctuation symmetry is inherited from reflection symmetry of the reduced variable ϖ . Moreover, the univariate affinity $\tilde{\varepsilon}_p$ is simply given by the difference of chemical potentials,

$$\tilde{\varepsilon}_p = \beta_{p,-} - \beta_{p,+} = \log \nu_+ - \log \nu_- = \log \kappa^2, \quad (76)$$

implying the UFR of the form

$$F_p(\lambda) = F_p(-\lambda - \tilde{\varepsilon}_p). \quad (77)$$

Using further that freely propagating particles obey pure transmission, the SCGF $F_p(\lambda)$ satisfies the so-called *extended fluctuation relation* [80]. The latter states that $F_p(\lambda)$ is fully determined already by the first scaled cumulant $s_1^{(p)} = \bar{j}_p$, namely

$$F_p(\lambda) = \int_0^{\lambda} d\gamma \bar{j}_p(\log \nu_- - \gamma, \log \nu_+ + \gamma). \quad (78)$$

Plugging in $\bar{j}_p = (1 + \nu_-)^{-1} - (1 + \nu_+)^{-1}$ correctly reproduces Eq. (72).

B. Joint particle-charge fluctuations

We now consider the FCS of joint particle-charge transfer encoded in $G_{c,p}(\lambda_c, \lambda_p|t)$. The exact finite-time joint MFG $G_{c,p}(\lambda_c, \lambda_p|t)$ can be calculated from first principles (see Sec. V), by following the lines of Ref. [74] adapted to the bipartitioned initial state (see Sec. III). Alternatively, $G_{c,p}(\lambda_c, \lambda_p|t)$ can be obtained from the univariate particle-current MGF $G_p(\lambda_p|t)$ by applying the general replacement rule (58). The associated joint SCGF $F_{c,p}(\lambda_c, \lambda_c) = \lim_{t \rightarrow \infty} t^{-1} \log G_{c,p}(\lambda_c, \lambda_p|t)$ is subsequently computed by applying the Laplace’s method (the derivations are presented in Sec. V).

By introducing the ‘dressed counting fields’

$$\mu_{\pm}(\lambda_c) \equiv \cosh(\lambda_c) \mp b_{\pm} \sinh(\lambda_c), \quad (79)$$

and a pair of auxiliary functions depending on the densities and λ_p ,

$$\nu_{\pm}(\lambda_p) \equiv \frac{\bar{\rho}_{\pm}}{\rho_{\pm}} e^{\pm \lambda_p}, \quad \nu_{\pm} \equiv \nu_{\pm}(0), \quad (80)$$

we find $F_{c,p}^{(\pm)}(\lambda_c, \lambda_p) \equiv \log(f_{c,p}^{(\pm)}(\lambda_c, \lambda_p))$ with

$$f_{c,p}^{(\pm)}(\lambda_c, \lambda_p) = \prod_{\epsilon \in \{\pm\}} \frac{[\mu_{\pm}(\lambda_c)]^{\pm \epsilon} + \nu_{\epsilon}(\lambda_p)}{1 + \nu_{\epsilon}}. \quad (81)$$

The ‘bulk branches’ $F_{c,p}^{(\pm)}(\lambda_c, \lambda_p)$ are attributed to two isolated local maxima residing in the interior of the upper (lower) triangular subdomain \mathcal{D}_- (\mathcal{D}_+) of \mathcal{D}_{\square} , of the integration domain \mathcal{D}_{\square} of $G_{c,p}(\lambda_c, \lambda_p|t)$, see Sec. V. On the other hand, $F^{(0)} \equiv F_0$ represents a constant branch that only depends solely on particle densities,

$$F_0 = 2 \log [\sqrt{\rho_- \rho_+} + \sqrt{\bar{\rho}_- \bar{\rho}_+}]. \quad (82)$$

Such a flat branch occurs when the maximum of $G_{c,p}(\lambda_c, \lambda_p|t)$ it attained exactly on the diagonal of the integration domain \mathcal{D}_{\square} (as represented by the ‘ λ -flow’ of the bulk maxima in Fig. 6). This may be interpreted as bulk maxima escaping ‘out-of-bounds’ upon colliding with the diagonal, playing the role of the domain boundary $\partial\mathcal{D}$. More importantly, the bulk maxima are not unconditionally present in the integration domain \mathcal{D}_{\square} . Instead, they only appear within certain regions of the parameter space specified below. Introducing $\boldsymbol{\lambda} \equiv (\lambda_c, \lambda_p)$ and another auxiliary function

$$\kappa(\lambda_p) \equiv \sqrt{\frac{\nu_+(\lambda_p)}{\nu_-(\lambda_p)}}, \quad \kappa \equiv \kappa(0), \quad (83)$$

we find that in the regions

$$\mathcal{E}_{\pm} = \{\mu_{\pm}(\lambda_c) \geq \kappa^{\pm 1}(\lambda_p) \wedge \mu_{\mp}(\lambda_c) < \kappa^{\mp 1}(\lambda_p)\}, \quad (84)$$

in the two-dimensional $\boldsymbol{\lambda}$ -plane, only a single extremum contribution $F_{\pm}(\lambda)$ appears in the bulk of subdomains \mathcal{D}_{\pm} , whereas both extrema coexist within

$$\mathcal{E}_{+-} = \{\mu_+(\lambda_c) \geq \kappa(\lambda_p) \wedge \mu_-(\lambda_c) \geq \kappa^{-1}(\lambda_p)\}. \quad (85)$$

Competition between the two bulk extrema and the flat branch can be summarized in terms of the following *selection rules*

$$\mathcal{I}_{\pm} = \mathcal{E}_{\pm} \vee \left(\mathcal{E}_{+-} \wedge f_{c,p}^{(\pm)}(\boldsymbol{\lambda}) \geq f_{c,p}^{(\mp)}(\boldsymbol{\lambda}) \right), \quad (86)$$

$$\mathcal{I}_0 = \{\mu_+(\lambda_c) < \kappa(\lambda_p) \wedge \mu_-(\lambda_c) < \kappa^{-1}(\lambda_p)\}, \quad (87)$$

which permit us to write compactly

$$F_{c,p}(\boldsymbol{\lambda}) = \left\{ F_{c,p}^{(k)}(\boldsymbol{\lambda}); \boldsymbol{\lambda} \in \mathcal{I}_k \right\}_{k \in \{+, 0, -\}}. \quad (88)$$

We postpone a more detailed study of $F_{c,p}(\boldsymbol{\lambda})$ to Sec. IV D 1, where we establish the fluctuation relation.




FIG. 6. Visualization of the ‘ λ -flow’ over the integration domain \mathcal{D}_{\square} of MGF $G(\lambda|t)$, parametrized in terms of continuous scaling coordinates $x_1 = l/t$ and $x_2 = r/t$ (with l and r representing the number of transferred ballistic holes from the left and right partitions, respectively, in units of t). Trajectories along the direction of arrows represent the motion of bulk maxima from the interior of the integration domain, associated with the bulk branches, namely $F_+(\lambda)$ (red) and $F_-(\lambda)$ (blue). When the bulk maxima reach the diagonal, representing the boundary \mathcal{D}_{\square} for each of the bulk subdomains \mathcal{D}_{\pm} , the constant (flat) branch F_0 sets in. Solid (dashed) curves designate the physical (subleading, metastable) contributions to the SCGF $F(\lambda)$. Four different dynamical regimes are shown: regular regime $[[+]]$, where a single physical branch $F_+(\lambda \in \mathbb{R})$ dominates for all $\lambda \in \mathbb{R}$; two types of tunneling (sub)regimes, to the same branch $[[+ \times +]]$ and to a different branch $[[+ \times -]]$ via F_0 supported on $[\lambda_{\pm}^{(-)}, \lambda_{\pm}^{(+)})$, and a corner regime $[[+ \times -]]$ with F_+ jumping over to F_- at corner point λ_{∞} .

C. Anomalous charge fluctuations

In the following, we focus our discussion to the univariate charge-current SCGF $F_c(\lambda_c)$. Although $F_c(\lambda_c)$ is merely a specialization of the full joint SCGF $F_{c,p}(\lambda_c, \lambda_p)$ obtained by putting $\lambda_p = 0$ (corresponding to integrating out the cumulative particle current), the univariate function $F_c(\lambda_c)$ reveals (as already summarized earlier in Sec. III) an intricate structure. Unlike $F_{c,p}(\lambda_c, \lambda_p)$, $F_c(\lambda_c)$ does not necessarily involve all three dynamical phases when restricted to $\lambda_c \in \mathbb{R}$.




FIG. 7. Dynamical phase diagram of the joint particle-charge SCGF $F_{c,p}(\lambda_c, \lambda_p)$ in the real λ -plane, exemplified for different dynamical regimes of $F_c(\lambda_c)$: (a) **regular regime** $[[+]]$, (b) **corner regime** $[[+\bowtie-]]$, (c) **tunneling subregime** $[[+\asymp+]]$ and (d) **tunneling subregime** $[[+\asymp-]]$. The bulk dynamical phases $F_{c,p}^{(\pm)}(\lambda_c, \lambda_p)$ are shown in red (+) and blue (-). Their level sets are shown by solid lines of the corresponding color. The constant phase $F^{(0)}$ is shown in white. Coordinate axes $\lambda_p = 0$ ($\lambda_c = 0$) are represented by solid horizontal (vertical) black lines. The inversion point $\lambda^{\text{inv}} = -\varepsilon/2$ of $F_{c,p}(\lambda_c, \lambda_p)$ lies in the geometric center of $F^{(0)}$, while λ_p^{inv} and λ_c^{inv} are inversion points associated to the univariate SCGFs $F_p(\lambda_p)$ and $F_c(\lambda_c)$, respectively. Parameters values in (a), (b), (c), (d) read respectively: $\rho_- = 0.18, 0.34, 0.28, 0.34$, $\rho_+ = 0.4, 0.4, 0.66, 0.43$, $b_- = 0.68, 0.99, -0.9, -0.73$, $b_+ = -0.26, 0.3, 0.97, 0.85$.

To lighten our notations, we shall subsequently drop the subscript by identifying $F_c(\lambda_c) \equiv F(\lambda)$, where

$$F(\lambda) = \{F_k(\lambda); \lambda \in \mathcal{S}_k^c\}_{k \in \{+,0,-\}}, \quad (89)$$

for $\mathcal{S}_k^c \equiv \mathcal{S}_k(\lambda_p = 0)$. Moreover, in the limit $|\lambda| \rightarrow \infty$, the bulk branches $F_{\pm}(\lambda)$ grow linearly as

$$F_{\pm}(\lambda) \asymp |\lambda| + \log \left[\frac{(1 \mp b_{\pm})\nu_{\pm}}{2(1 + \nu_+)(1 + \nu_-)} \right] + \mathcal{O}(e^{-\lambda}). \quad (90)$$

which is a manifestation of strict causality (with unit maximal velocity) of the time evolution.

For any fixed choice of the state parameters ρ_{\pm}, b_{\pm} , the physical value of the SCGF $F(\lambda)$ for a given counting field λ is given by the dominant branch. Which of the branches $F_k(\lambda)$ dominates the growth of the MGF $G(\lambda|t)$ at large times is encoded in the selection rules derived in Sec. IV B.

To obtain an explicit parametrization of the separatrices between different dynamical regimes in the hard-core automaton, we now explicitly work out the selection rules for univariate charge-current SCGF $F(\lambda)$. The bulk branches $F_{\pm}(\lambda)$ attain their extrema at $\lambda_{\pm}^{(\varepsilon)}$, $\varepsilon \in \{+, -\}$,

$$\lambda_{\pm}^{(\varepsilon)} = \log \left[\frac{\kappa^{\pm 1} + \varepsilon \sqrt{\kappa^{\pm 2} + b_{\pm}^2 - 1}}{1 \mp b_{\pm}} \right], \quad (91)$$

and a central extremum at

$$\lambda_{\pm}^{(0)} = \frac{1}{2} \log \left[\frac{1 \pm b_{\pm}}{1 \mp b_{\pm}} \right] = \beta_c(\mp b_{\pm}). \quad (92)$$

Importantly, while $\lambda_{\pm}^{(0)}$ is always real for any $\lambda \in \mathbb{R}$, the other pair of extrema $\lambda_{\pm}^{(\pm)}$ can be either real (with \pm

referring to the left (-) and right (+) real minima) or imaginary depending on parameters κ, b_{\pm} of the initial state.

For compactness of presentation, we now assume (with no loss of generality) that $\rho_+ > \rho_-$. Introducing new auxiliary parameters

$$\gamma_{\pm} \equiv \kappa^2(b_{\pm} \pm 1), \quad (93)$$

the phase boundary for the transition $F_+ \leftrightarrow F_-$ (for $\lambda \in \mathbb{R}$) is inferred by equating $F_+(\lambda) = F_-(\lambda)$, yielding $\gamma_{\pm} = \pm 1 - b_+$ for $\pm \lambda > 0$. Similarly, the phase boundaries between the bulk branches $F_{\pm}(\lambda)$ and flat branch F_0 occurs when $F_{\pm} = F_0$, i.e. at the threshold value $b_{\pm} = \pm \sqrt{1 - \kappa^2}$. The limiting curves for the tunneling transitions $F_{\pm} \leftrightarrow F_{\mp}$ are determined from the conditions

$$b_+ > 0 : \quad F_-(\lambda_-^{(+)}) = F_+(\lambda_+^{(\pm)}), \quad (94)$$

$$b_+ < 0 : \quad F_-(\lambda_-^{(-)}) = F_+(\lambda_+^{(\pm)}), \quad (95)$$

yielding

$$b_+ = \sqrt{1 + \gamma_-}, \quad b_+ = -\sqrt{1 + \gamma_-}, \quad (96)$$

respectively. This allows us to determine the separatrices between different regimes and their support (in the parameter space of initial states):

(I) **regular regime** $[[+]]$

$$|b_+| < \sqrt{1 - \kappa^2} \quad \wedge \quad b_+ \lesssim -\gamma_{\pm} \pm 1, \quad (97)$$

(II) **tunneling regimes**

- subregime $[[\pm \asymp \pm]]$, with tunneling from F_{\pm} via F_0 back to the same bulk branch F_{\pm}

$$|b_+| < \sqrt{1 - \kappa^2} \quad \wedge \quad b_+ \lesssim -\gamma_{\pm} \pm 1. \quad (98)$$

- subregime $[\pm \asymp \mp]$, with tunneling from F_{\pm} via F_0 to a different bulk branch F_{\mp}

$$b_+ \lesseqgtr \pm \sqrt{1 + \gamma_-} \quad \wedge \quad |b_+| > \sqrt{1 + \gamma_-}. \quad (99)$$

(III) **corner regime** $[\pm \bowtie \mp]$

- *unbroken* branch F_+ with a unique minimum

$$|b_+| < \sqrt{1 - \kappa^2} \quad \wedge \quad b_+ \lesseqgtr -\gamma_{\pm} \mp 1. \quad (100)$$

- *regular* branch F_+ with a doubly degenerate minimum

$$|b_+| > \sqrt{1 - \kappa^2} \quad \wedge \quad b_- \lesseqgtr 0 \quad \wedge \quad b_+ \gtrless \pm \sqrt{1 + \gamma_-}. \quad (101)$$

(IV) **mixed regime** $[\mp \bowtie \pm \asymp \pm]$

$$|b_+| < \sqrt{1 - \kappa^2} \quad \wedge \quad b_- \gtrless 0 \quad \wedge \quad (-\gamma_{\mp} \mp 1 \gtrless b_+ \gtrless \pm \sqrt{1 + \gamma_-}). \quad (102)$$

D. Fluctuation symmetry

In this section, we examine the fluctuation symmetry of the joint MGF $G(\lambda_c, \lambda_p|t)$ (see Sec. V for the derivation) and univariate charge-current SCGF $F_c(\lambda_c)$.

1. Multivariate fluctuation relation

By explicit computations, we established that the joint particle-charge SCGF $F(\lambda_c, \lambda_p) = \lim_{t \rightarrow \infty} t^{-1} \log G(\lambda_c, \lambda_p|t)$ obeys the multivariate fluctuation relation of the form

$$F(\lambda_c, \lambda_p) = F(-\lambda_c - \varepsilon_c, -\lambda_p - \varepsilon_p), \quad (103)$$

with affinities $\varepsilon \equiv (\varepsilon_c, \varepsilon_p)$ reading

$$\varepsilon_c = \frac{1}{2} \log \left[\frac{1 + b_-}{1 - b_-} \frac{1 - b_+}{1 + b_+} \right] = \delta\beta_c, \quad (104)$$

$$\varepsilon_p = \tilde{\varepsilon}_p + \frac{1}{2} \log \left[\frac{1 - b_-^2}{1 - b_+^2} \right] = \delta\beta_p. \quad (105)$$

Recalling that $\delta\beta_i = \beta_{i,-} - \beta_{i,+}$ for $i \in \{c, p\}$ correspond to differences of the particle and charge chemical potentials in the two initial partitions, the affinities indeed take a canonical form [77, 80, 129]. In the hard-core automaton (and other charged single-file systems) $\tilde{\varepsilon}_p = \log \kappa^2$. We have therefore established that in the considered models with time-reversal invariant dynamics

the joint particle-charge fluctuation relation (103) is unconditionally satisfied in the entire parameter space despite dynamical phase transitions.

To illustrate the geometric meaning of the MFR, we next examine a diagram of dynamical phases in the two-dimensional λ -plane, depicted in Fig. 7. Notably, all three distinct dynamical phases $F^{(k)}(\lambda)$, with $k \in \{+, 0, -\}$, are always present in the joint SCGF $F_{c,p}(\lambda)$. The constant phase $F^{(0)}(\lambda)$ appears as a single compact ‘island’ somewhere between $F_{c,p}^{(\pm)}(\lambda)$. The one-dimensional boundaries $\mathcal{L}_{k\ell}$ separating the k th and ℓ th can be deduced from the balancing conditions $F^{(k)}(\lambda) = F^{(\ell)}(\lambda)$ (provided that both phases coexist, i.e. obey the selection rules given by Eq. (87)). As shown in Fig. 7, outside of the constant branch $F^{(0)}$, the level sets of $F_{c,p}(\lambda_c, \lambda_p)$ form closed contours. On such contours, any two points symmetric with respect to the inversion point λ^{inv} always belong to distinct bulk phases $F_{c,p}^{(\pm)}(\lambda)$.

The MFR implies the existence of an inversion point $\lambda^{\text{inv}} = -\varepsilon/2$. Curiously, the latter is always situated precisely in the geometric center of the constant phase F_0 (see Fig. 7), exactly halfway between two *triple* critical points – one of them being the inversion point of $F_p(\lambda_p)$ located on the vertical ($\lambda_c = 0$) axis at $\lambda_p^{\text{inv}} = -\log(\kappa)$, whereas the other is situated at

$$\lambda_{\text{triple}} = \left(-\delta\beta_c, \lambda_p^{\text{inv}} + \frac{1}{2} \log \left[\frac{1 - b_+^2}{1 - b_-^2} \right] \right). \quad (106)$$

Upon approaching equilibrium ($\delta\beta_p = \delta\beta_c = 0$), both triple points move towards the origin $\lambda = (0, 0)$ causing $F^{(0)}$ to disappear. It is also worth noticing that the effective affinity $\tilde{\varepsilon}_p$ (located on the $\lambda_c = 0$ axis in the λ -plane) must be a triple point at the junction of three phase boundaries $\mathcal{L}_{k\ell}$ where $F^{(+)}(\lambda) = F^{(-)}(\lambda) = F^{(0)}$. This readily follows from the fact that $F_p(\lambda_p)$ is (by assumption) strictly convex and twice differentiable.

2. Spontaneous breaking of the univariate fluctuation relation

We have already demonstrated that the univariate SCGFs of the particle current $F_p(\lambda)$ exhibits the Gallavotti–Cohen symmetry, cf. Eq. (77). We now investigate the symmetry properties of the univariate charge current SCGF $F_c(\lambda_c) = F_{c,p}(\lambda_c, \lambda_p = 0)$. In the following, we write shortly $F_c(\lambda_c) \equiv F(\lambda)$, and systematically examine each of the dynamical regimes.

In **regular regimes** $[\pm]$, $F(\lambda)$ coincides with one of the bulk branches $F_{\pm}(\lambda)$. Therefore, $F(\lambda)$ is strictly convex and everywhere differentiable and hence satisfies the following UFR

$$F_{\pm}(\lambda) = F_{\pm}(-\lambda - \tilde{\varepsilon}_c^{(\pm)}), \quad \tilde{\varepsilon}_c^{(\pm)} = \mp 2\beta_c(b_{\pm}). \quad (107)$$

Here the effective affinities $\tilde{\varepsilon}_c^{(\pm)} = -2\lambda_c^{\text{inv}}$, with $\lambda_{\pm}^{\text{inv}} = \lambda_{\pm}^{(0)}$ being the unique minima of $F_{\pm}(\lambda)$, are different from the actual charge affinity ε_c . Indeed, $\tilde{\varepsilon}_c^{(\pm)}$ only depend on a single chemical potential (that of the dominant dynamical phase) and not on the canonical ‘gradient’ $\delta\beta_c$ given by Eq. (104) that enters in the MFR.




FIG. 8. Dynamical regimes of the hardcore cellular automaton, represented by a two-dimensional cross section in the b -plane cross section of the parameter space (shown for $\rho_- = 0.45$ and $\rho_+ = 0.5$) with SCGF $F(\lambda)$ exhibiting different interweaving patterns of dynamical phases: **regular regime** (blue), **corner regime** (red), two distinct types **tunneling regimes** via F_0 – to the same branch (yellow) and to another branch (green) – and **mixed regime** (purple), involving transfer via a corner followed by tunneling to the same branch or vice-versa.

In **corner regimes**, denoted by $[\pm \bowtie \mp]$, there is a non-differentiable point (a convex corner) in $F(\lambda)$ at $\lambda_{\bowtie} \in \mathbb{R}$ where the bulk branches $F_{\pm}(\lambda)$ interchange their dominance, signaling a DPT of the first order. In effect, the UFR of $F_c(\lambda_c)$ is invariably violated (seen as the absence of the inversion point on the $\lambda_p = 0$ axis in panel (b) in Fig. 7).

Despite that, the UFR is preserved *locally* within the ‘subcritical region’, i.e. an interval of counting fields λ centered at $\lambda_{\pm}^{(0)} = -\varepsilon_c^{(\pm)}/2$, such that

$$|\lambda - \lambda_{\pm}^{(0)}| < |\lambda_{\bowtie} - \lambda_{\pm}^{(0)}|. \quad (108)$$

Accordingly, the LDF $I(j)$ develops a non-differentiable point (a concave corner) at the critical large current j_{\bowtie} , and hence the UFR is only satisfied locally for subcritical (large) currents $|j| < |j_{\bowtie}|$. Exceeding the critical current j_{\bowtie} , namely for $|j| > |j_{\bowtie}|$, the GCR is no longer satisfied as large fluctuations in the opposite direction belong to a different dynamical phase.

In **tunneling regime**, the appearance of the flat branch F_0 (assuming $F(\lambda)$ is everywhere differentiable), require us to differentiate between two scenarios, here-

after referred to as **tunneling subregimes**: (i) tunneling from F_{\pm} via F_0 back to the same phase (denoted by $[\pm \times \pm]$), or (ii) tunneling to the other bulk branch (denoted by $[\pm \times \mp]$). In each of these subregimes, both bulk branches F_{\pm} reside in a symmetry-broken phase with a two-fold degenerate minima at $\{\lambda_{\pm}^{(\epsilon)}; \epsilon \in \{\pm\}\}$. As we elaborate in turn, validity of the univariate Gallavotti–Cohen relation depends crucially on the dynamical subregime.

Depending on the state parameters, a second-order DPT takes place at two out of these four degenerate minima. We recall that at the second-order critical point the second derivative of the physical (dynamical) free energy $F''_{\pm}(\lambda)$ experiences a jump discontinuity. The pair of critical points always consists of two adjacent minima. In fact, there are only two possible scenarios: (i) the pair of second-order critical points coincides with two minima of the same bulk branch or (ii) there is a tunneling transition from the left minimum $\lambda_{\pm}^{(-)}$ of F_{\pm} to the right minimum $\lambda_{\mp}^{(+)}$ of another branch F_{\mp} . Using that the symmetry-broken bulk branches $F_{\pm}(\lambda)$ each enjoys reflection symmetry with respect to the central maximum $\lambda_{\pm}^{(0)}$, the inversion symmetry $F(\lambda)$ clearly remains intact in case (i) despite the presence of the constant phase F_0 . In effect, the UFR in the form (107) remain valid. This is however no longer true in case (ii) where, upon approaching the boundaries of the flat segment \mathcal{S}_0 from the outside, we find a mismatch in the second derivatives evaluated in both minima,

$$\lim_{\lambda \rightarrow \lambda_{\pm}^{(-)}} F''_{\pm}(\lambda) \neq \lim_{\lambda \rightarrow \lambda_{\mp}^{(+)}} F''_{\mp}(\lambda). \quad (109)$$

This means that neither of the effective charge affinities $\varepsilon_c^{(\pm)}$ (with the exception of equilibrium, when they are both equal) can be the inversion point of $F(\lambda)$. In effect, the UFR (107) is violated for the entire range of counting fields $\lambda \in \mathbb{R}$. Spontaneous breaking of the UFR can be alternatively discussed from the viewpoint of the LDF $I(j) \equiv I_c(j_c)$. In **tunneling regime**, $I(j)$ is not differentiable at $j = 0$ in both subregimes (i) and (ii). In subregime (i), lack of differentiability has no implications on the validity of the UFR since $I(j)$ involves only a single branch $I^{(\pm)}(j)$. In subregime (ii), positive and negative (large) currents are realized in different dynamical phases and consequently the UFR breaks down on the entire support of j . The same reasoning applies to **mixed regime**, where both types of dynamical criticality are simultaneously present.

To conclude the section, we briefly recapitulate the key findings. Unlike the joint particle-charge SCGF, the univariate charge-current SCGF $F(\lambda)$ does not *globally* (i.e. for the entire interval of counting fields $\lambda \in \mathbb{R}$) satisfy a fluctuation relation of the Gallavotti–Cohen type. The relation (28) is found to be obeyed only in a finite fraction of the parameter space, comprising the union of **regular regimes** ($[\pm]$ and $[\mp]$) and **tunneling subregime** $[\pm \times \pm]$.




FIG. 9. The b -plane cross section of various dynamical regimes (shown for $\rho_+ = 0.5$ and several values of ρ_-) in the hardcore automaton: regular $[\pm]$ (blue), corner $[\pm \bowtie -]$ (red), tunnelling regime, with subregimes $[\pm \times \mp]$ (green) and $[\pm \times \mp]$ (yellow) and mixed (purple). Separatrices, marked by white lines, represent phase boundaries associated with dynamical phase transitions of first or second order. The univariate fluctuation relation of the charge-current large-deviation function holds globally in $[\pm]$ and $[\pm \times \pm]$, and is violated in $[\pm \times \mp]$ (globally) and $[\pm \bowtie \mp]$ (locally).

As represented in Fig. 8, these regimes are confined within a diagonal belt region in the b -plane cross section. In other words, a continuous (second-order) convexity-breaking transition experienced by regular bulk branches, inducing dynamical phase transitions between **regular regimes** $[\pm]$ and **tunneling subregimes** $[\pm \times \pm]$, preserves the UFR. By contrast, the UFR is violated globally whenever the system undergoes a dynamical phase transition of first order, namely in **corner regimes** $[\pm \bowtie \mp]$ and **mixed regime**. In these two dynamical regime, the UFR still holds *locally* for subcritical large currents. Finally, the UFR is completely broken in the **tunneling subregime** $[\pm \times \mp]$. We also note that $[\pm \times \mp]$ cannot be reached from $[\pm]$ directly (as seen in Fig. 8) but only via a first-order DPT to another regime. The dependence of dynamical regimes on state parameters is illustrated in Fig. 9.

E. Phase diagrams

Thus far we have been almost exclusively concerned with the late-time behavior of the (univariate) charge-current MGF $G(\lambda|t)$ restricted to real-valued counting fields. Using asymptotic analysis, we have inferred the general structure of the SCGF $F(\lambda)$ and the corresponding large-deviation rate function $I(j)$. As already emphasized in Sec. II, the counting field λ is merely a formal complex control variable with no direct physical meaning. As we now demonstrate, the outlined physical phases and interweaving dynamical regime admit analytic continuation to complex λ . In the conventional theory of thermodynamic phase transitions, such an analytic continuation of couplings or state parameters, such as temperature and chemical potentials, goes under the name of Lee–Yang analysis [130]. To elucidate the interplay of the three dynamical phases, the Lee–Yang approach is tremendously helpful. Here we only provide a broader overview of the main results, postponing a detailed analysis to Sec. V.




FIG. 10. A representative example of a phase diagram (selected from the mixed regime, with $\rho_- = 0.75$, $\rho_+ = 0.77$, $b_- = 0.78$, $b_+ = -0.47$), representing analytic continuation of dynamical phases into the complex v -plane. Shaded regions indicate the regions of dominance \mathcal{R}_\pm belonging to the bulk dynamical phases (F_+ in red, F_- in blue), while white background is the region of dominance \mathcal{R}_0 associated with flat branch F_0 . Algebraic curves corresponding to extended anti-Stokes lines that separate distinct dynamical phases apart from one another: $F_+ \leftrightarrow F_0$ (red), $F_- \leftrightarrow F_0$ (blue) and $F_+ \leftrightarrow F_-$ (gray). Physical parts of the curves, marked by solid lines, coincide with the Lee–Yang contours (with dashed lines marking unphysical parts). The Lee–Yang zeros (shown for $t = 32$) are marked with gray dots.

The Lee–Yang theory offers a powerful theoretical framework for characterizing thermodynamic phase transitions using the language of complex analysis, based on the singularity structure of the thermodynamic free

energy [131–133], analytically continued to the complex temperature or fugacity plane. In finite systems, every zero contained in the polynomial part of the partition sum show up as an isolated logarithmic singularity in the free energy extended to the complex fugacity plane. Although singularities of this type are always away from the physical real axis and thus unphysical, they are nevertheless responsible for the appearance of critical points at large time. Specifically, the number of singularities grows extensively (i.e. linearly) with system size and, in the thermodynamic limit, the singularities tend to form clusters. One commonly finds that the majority of singularities arrange densely along contours or inside two-dimensional subdomains. Such condensates form natural boundaries between distinct thermodynamic phases.




FIG. 11. Phase diagram in the **regular regime** $\llbracket + \rrbracket$ (with state parameters $\rho_- = 0.18$, $\rho_+ = 0.4$, $b_- = 0.28$, $b_+ = -0.14$) in complex v -plane, comprising a single contiguous region of dominance $\mathcal{R}_+^{(v)}$ associated to F_+ (red), and the domain $\mathcal{R}_0^{(v)}$ of the constant phase F_0 (white background). The extended anti-Stokes line $\mathcal{A}_{+,0}^{(v)}$ ($\mathcal{A}_{-,0}^{(v)}$), separating F_+ (F_-) from F_0 , is shown by the red (blue) curve, with solid part corresponding to the Lee–Yang contour $\mathcal{C}_{+,0}^{(v)}$ (and gray dots marking the Lee–Yang zeros of $G(v|t)$ at time $t = 40$). Dashed blue line is the ghost anti-Stokes line $\mathcal{A}_{-,0}^{(v)}$ violating the selection criterion.

Introducing a complex counting field $z \equiv e^\lambda$, we can apply the Lee–Yang approach to the rescaled CGF $F(z|t)$ by maintaining analogy with the thermodynamic free energy. Taking advantage of the fact that in the hardcore automaton the MGF $G(z|t)$ enjoys a particularly simple analytic structure, we managed to obtain a fully general solution. By exploiting the formal analogy with two-dimensional electrostatics [133, 134], we succeeded in explicitly parametrizing the Lee–Yang contours, permitting us to compute the associated density of zeros from the jump discontinuities of the imaginary poten-

tial of the complexified dynamical free energy density $F(z)$ upon transversing the phase boundaries in the z -plane. In the hardcore automaton, we find a unit fraction of Lee–Yang zeros condensing along certain closed contours in the complex z -plane, separating different regions of dominance. This implies that Lee–Yang contours along which the zeros condense naturally play the role of phase boundaries between distinct (complexified) dynamical phases, obstructing analytic continuation of $F(z)$ across the phase boundaries. In the language of Morse theory, they are commonly referred to as the anti-Stokes lines (see e.g. Refs. [135, 136]).




FIG. 12. Phase diagram in **corner regime** $\llbracket + \bowtie - \rrbracket$ (with state parameters $\rho_- = 0.34$, $\rho_+ = 0.74$, $b_- = 0.97$, $b_+ = -0.72$) in complex v -plane, comprising two disjoint regions of dominance $\mathcal{R}_\pm^{(v)}$ associated to F_+ (red) and F_- (blue), next to domain $\mathcal{R}_0^{(v)}$ of the constant phase F_0 (white background). Curves represent the extended anti-Stokes lines $\mathcal{A}_{k,\ell}^{(v)}$, separating F_k from F_ℓ , for $k, \ell \in \{+, 0, -\}$ with $k \neq \ell$, namely $\mathcal{A}_{+,0}^{(v)}$ (red), $\mathcal{A}_{-,0}^{(v)}$ (blue) and $\mathcal{A}_{+,-}^{(v)}$ (gray). Solid parts of the curves belong to the Lee–Yang contours. Gray dots mark the Lee–Yang zeros of $G(v|t)$ at time $t = 46$.

In terms of the multiplicative counting field z , the MGF $G(z|t)$ represents a Laurent polynomial with $2t$ terms. By exploiting the underlying symmetries, there is however a more convenient (bijective) reparametrization in terms of a complex counting variable v , defined via a conformal (Möbius) transformation $v(z) = (z-1)/(z+1)$, which we adopt in the following analysis. The physical part of $F(v)$ (real λ, z) then maps to the compact interval $\varphi \equiv [-1, 1]$, whereas the regions of dominance associated with dynamical phases F_k will be denoted by $\mathcal{R}_k^{(v)} \subset \mathbb{C}$. It is worth noticing that the extended phase diagram in the complex v -plane comprises all three phases, with the sole exception being the regular regimes $\llbracket \pm \rrbracket$. In addition, each of the two bulk phases forms a single compact region of dominance, $F_\pm(v \in \mathcal{R}_\pm^{(v)})$, with the exception of the tunneling regimes $\llbracket \pm \asymp \pm \rrbracket$. The flat phase F_0 may be seen as the background.

In **regular regime**, the physical interval φ is con-

tained entirely within a single region of dominance $\mathcal{R}_{k \in \{\pm\}}^{(v)}$, depending on the selection criteria. Notice that both the dominant and subdominant bulk branches F_{\pm} are always simultaneously present in the v -plane, with each of them becoming physical along $\mathcal{S}_{\pm}^{(v)} = \wp \cap \mathcal{R}_{\pm}^{(v)}$ when there is a non-trivial intersection with the physical line \wp .




FIG. 13. Phase diagram in subregime $[[+ \times +]]$ of **tunneling regime** (with state parameters $\rho_- = 0.23$, $\rho_+ = 0.33$, $b_- = -0.67$, $b_+ = 0.69$), comprising two disjoint regions of dominance $\mathcal{R}_{\pm}^{(v)}$ associated to F_+ (red), and the domain $\mathcal{R}_0^{(v)}$ of the constant phase F_0 (white background). The extended anti-Stokes lines $\mathcal{A}_{+,0}^{(v)}$ ($\mathcal{A}_{-,0}^{(v)}$), separating F_+ (F_-) from F_0 , are shown by red (blue) curve. Solid curves indicates the Lee–Yang contour $\mathcal{C}_{+,0}^{(v)}$. Gray dots mark the Lee–Yang zeros of $G(v|t)$ at time $t = 36$.

In **corner regime**, the physical interval comprises two regions (intervals) of dominance, $\wp = \mathcal{S}_+^{(v)} \cup \mathcal{S}_-^{(v)}$, with $\mathcal{S}_{\pm}^{(v)}$ meeting at the corner point $v_{\bowtie} \in \wp$, being a dynamical critical point of first order where the bulk branches interchange their dominance. The Lee–Yang contour intersects the physical interval \wp with an impact angle of $\pi/2$ (see Figure 12), such that the density of zeros at the intersection is strictly positive, consistent with first-order criticality [132]. Curiously, the Lee–Yang contour emanating from the physical first-order critical point terminates in the complex v -plane at a (unphysical) *triple point* where all three distinct phase boundaries meet at one point.

In **tunneling regime**, the flat branch F_0 acquires a non-empty overlap with the physical interval \wp . Disregarding degenerate scenarios, there are only two possibilities of entering tunneling regimes $[[\pm \asymp \pm]]$ by varying parameters of the initial state, either from regular regimes $[[\pm]]$ or from mixed regimes $[[\mp \bowtie \pm \asymp \pm]]$ or $[[\pm \asymp \pm \bowtie \mp]]$. The other tunneling subregimes $[[\pm \asymp \mp]]$ can be entered from $[[\pm \asymp \pm \bowtie \mp]]$ or $[[\pm \bowtie \mp]]$.

Entering from **regular regime**, the region $\mathcal{R}_0^{(v)}$ approaches close to \wp until it eventually opens up a ‘corri-

dor’ through $\mathcal{R}_{\pm}^{(v)}$, splitting it into two disjoint compact regions (see Fig. 13). In this process, a physical regular bulk branch experiences a symmetry-breaking transition to a non-convex shape. Note that the pairs of degenerate minima associated with the bulk branches F_{\pm} show up as singular nodal points of the algebraic curves that parametrize the extended anti-Stokes lines (by disregarding the selection rules). The second-order transition can hence be also be seen as a conjugate pair of complex nodal points of the extended anti-Stokes lines colliding with \wp and thus becoming physical dynamical critical points.




FIG. 14. Phase diagram in subregime $[[+ \times -]]$ of **tunneling subregime** (with state parameters $\rho_- = 0.34$, $\rho_+ = 0.43$, $b_- = -0.68$, $b_+ = -0.85$), comprising two disjoint regions of dominance $\mathcal{R}_{\pm}^{(v)}$ associated to both bulk dynamical phases F_+ (red) and F_- (blue), and the domain $\mathcal{R}_0^{(v)}$ of the constant phase F_0 (white background). The extended anti-Stokes lines $\mathcal{A}_{\pm,0}^{(v)}$ separating F_+ (F_-) from F_0 are shown by red (blue) algebraic curves. The respective Lee–Yang contour $\mathcal{C}_{+,0}^{(v)}$ and $\mathcal{C}_{-,0}^{(v)}$ are marked by solid curves. Gray dots mark the Lee–Yang zeros of $G(v|t)$ at time $t = 36$.

Entering from the mixed regime corresponds to the first and second order critical points swapping places, see Fig. 10.

Lastly, we note that two tunneling subregimes, depicted in Figures 13 and 14, are disconnected. Naïvely, one would expect that a direct transition between these two regimes takes place when adjacent degenerate minima of the bulk branches F_{\pm} swap places. This is not what happens, however. Suppose that, for definiteness, we start in $[[+ \times +]]$: before $\lambda_-^{(+)}$ can pass through $\lambda_+^{(+)}$, a transition to $[[+ \times + \bowtie -]]$ is inevitable simply due to the fact that branch F_+ is steeper than F_- at large λ .

V. TECHNICAL BACKGROUND

In this section, we provide additional technical information and spell out the derivations. Considering a bipartitioned nonequilibrium initial state, the main object of study is the finite-time joint MGF

$$G(\lambda_c, \lambda_p | t) = \left\langle e^{\lambda_c J_c(t) + \lambda_p J_p(t)} \right\rangle_{\text{init}}. \quad (110)$$

By applying methods of localization, we compute large- t limit of $G(\lambda_c, \lambda_p | t)$ and infer the exact joint SCGF $F(\lambda_c, \lambda_p)$. We subsequently restrict our considerations to the univariate SCGF $F_c(\lambda_c)$ encoding the FCS of the charge transfer, while computing the scaled cumulants and deduce the probability distribution of typical fluctuations. We proceed afterwards by carefully examining the general solutions of the minimization problem arising in the dressing procedure applied to the large-deviation rate function (53) and outline the main formal properties of such a dressing transformation. We also include illustrative examples of the emergent dynamical regimes. We provide a short overview of Lee–Yang theory, which we afterwards apply to the hardcore cellular automaton. The exact finite-time MGF allows us to determine the complex regions of dominance and parametrize the Lee–Yang contours that separate between them. We conclude the section by resolving the degenerate case of grand-canonical equilibrium states that feature an exceptional triple critical point.

A. Exact full counting statistics

The exact formula for the finite-time charge-current MGF $G_c(\lambda_c | t)$ in the hardcore automaton, computed for the grand-canonical equilibrium states, is already known from Ref. [74]. The computation can be easily adapted to bipartitioned initial ensembles. To keep the notation in line with Ref. [74], we present the bivariate MGF $G(\lambda_c, \lambda_p | t)$ as a double sum by summing over contribution of freely propagating *holes* (as opposed to particles).

Let $l \equiv \bar{n}_-$ and $r \equiv \bar{n}_+$ denote the number of *holes* crossing the origin in a time interval t starting from the left and right partitions, respectively, and similarly $n_{\mp} = |\Lambda_{\mp}|$ for particles. Note that a particle crossing the origin requires a hole passing in the opposite direction, and thus we have $|\Lambda_{\pm}| = [|l - r| \pm (l - r)]/2$. By averaging over initial configurations, the exact finite-time joint MGF $G(\lambda_c, \lambda_p | t)$ is given by a double sum of the form

$$G(\lambda_c, \lambda_p | t) = \sum_{\bar{n}_-, \bar{n}_+ = 0}^t \prod_{\epsilon \in \{\pm\}} \binom{t}{\bar{n}_{\epsilon}} \rho_{\epsilon}^t [\nu_{\epsilon}(\lambda_p)]^{\bar{n}_{\epsilon}} \mu_{\epsilon}^{|\Lambda_{\epsilon}|}, \quad (111)$$

where $\mu_{\pm}(\lambda_c)$ is the ‘dressed’ counting fields (given by Eq. (79)). Under charge conjugation, $C : b_{\pm} \mapsto -b_{\pm}$, the joint MGF transforms as

$$C[G(\lambda_c, \lambda_p | t)] = G(-\lambda_c, \lambda_p | t). \quad (112)$$

Similarly, space reflection $R : x \mapsto -x$ interchanges the state parameters, $R : b_{\pm} \mapsto b_{\mp}$ and $R : \rho_{\pm} \mapsto \rho_{\mp}$, and hence the MGF satisfies

$$R[G(\lambda_c, \lambda_p | t)] = G(-\lambda_c, -\lambda_p | t). \quad (113)$$

The univariate MGF $G_c(\lambda_c | t)$ is simply obtained by marginalization, namely $G_c(\lambda_c | t) = G(\lambda_c, 0 | t)$. An alternative way to derive $G_c(\lambda_c | t)$ is to initially compute the PDF associated to the transferred particles, which is easily accomplished by identifying the cumulative particle current $J_p = r - l$, yielding

$$\mathcal{P}_p(J_p | t) = \sum_{\bar{n}_-, \bar{n}_+ = 0}^t \delta_{J_p, \bar{n}_+ - \bar{n}_-} \prod_{\epsilon \in \{\pm\}} \binom{t}{\bar{n}_{\epsilon}} \nu_{\epsilon}^{\bar{n}_{\epsilon}} \rho_{\epsilon}^t. \quad (114)$$

The associated MGF $G_p(\lambda_p | t)$ is given by the Laplace transform, $G_p(\lambda_p | t) = \mathcal{L}[\mathcal{P}_p(J_p | t)](-\lambda_p)$, yielding

$$G_p(\lambda_p | t) = \sum_{\bar{n}_-, \bar{n}_+ = 0}^t \prod_{\epsilon \in \{\pm\}} \binom{t}{\bar{n}_{\epsilon}} [\nu_{\epsilon}(\lambda_p)]^{\bar{n}_{\epsilon}} \rho_{\epsilon}^t. \quad (115)$$

Applying the dressing operator \mathfrak{D}_G (see Eq. (55)), amounts to using the substitution rule Eq. (58) in Eq. (115), which recovers Eq. (111).

By applying the particle-hole transformation to Eq. (114), we obtain the following equivalent double-sum representation of $\mathcal{P}_p(J_p | t)$ involving summation over the transferred particles instead of vacancies (cf. Eq. (34))

$$\mathcal{P}_p(J_p | t) = \sum_{n_-, n_+ = 0}^t \delta_{J_p, n_- - n_+} \prod_{\epsilon \in \{\pm\}} \binom{t}{n_{\epsilon}} \bar{\rho}_{\epsilon}^t \nu_{\epsilon}^{-n_{\epsilon}}. \quad (116)$$

The obtained result is very transparent and intuitive: computing $\mathcal{P}_p(J_p | t)$ for non-interacting ballistically propagating particles entails summing over all the contributions from n_- right movers from the left partition and subtracting n_+ left-movers from the right partition, weighted with appropriate statistical factors.

1. Localization

Asymptotic growth of MGF $G(\lambda | t)$ can be computed using Laplace’s method of localization as we now describe. We begin by first listing the general formulae, which we subsequently apply to our specific case. Let \mathcal{D} be a d -dimensional domain parameterized by coordinates $\mathbf{x} = (x_1, x_2, \dots, x_d)$. We consider two dummy functions $f(\mathbf{x})$ and $g(\mathbf{x})$ and an integral of the type $\mathfrak{G}_g[f] \equiv \int_{\mathcal{D}} d\mathbf{x} g(\mathbf{x}) e^{t f(\mathbf{x})}$. For large t , the integral localizes around the point \mathbf{x}_0 , $\nabla f|_{\mathbf{x}_0} = 0$, corresponding to the global maximum of f in the bulk (i.e. interior) of the integration domain \mathcal{D} . We have the following asymptotic approximation

$$\mathfrak{G}_g[f] \asymp \left(\frac{2\pi}{t} \right)^{d/2} \frac{g(\mathbf{x}_0) e^{t f(\mathbf{x}_0)}}{\sqrt{\det \mathcal{H}[f](\mathbf{x}_0)}}, \quad (117)$$

with $\mathcal{H}[f]$ being the Hessian of function f evaluated at point \mathbf{x}_0 . When \mathcal{D} is a compact domain, it might also happen that maximum is attained at the boundary $\partial\mathcal{D}$. In this case, assuming a non-vanishing derivative in the normal direction to the boundary with unit normal \mathbf{n}_\perp , we use the following formula for $\mathbf{x}_0 \in \partial\mathcal{D}$,

$$\mathfrak{G}_g[f] \asymp \frac{1}{2\pi} \left(\frac{2\pi}{t} \right)^{\frac{d+1}{2}} \frac{g(\mathbf{x}_0)e^{t f(\mathbf{x}_0)}}{|\nabla_{\mathbf{n}_\perp} f(\mathbf{x}_0)| \sqrt{\det \mathcal{H}_\parallel[f](\mathbf{x}_0)}}, \quad (118)$$

with $\mathcal{H}_\parallel[f](\mathbf{x}_0)$ denoting the Hessian submatrix evaluated in the remaining (non-normal) coordinates (i.e. subspace orthogonal to \mathbf{n}_\perp) at \mathbf{x}_0 .

Returning to our working example, we now infer the behavior of $G_{c,p}(\lambda_c, \lambda_p|t)$ at large times. We introduce the rescaled continuum coordinates

$$\frac{l}{t} \rightarrow \mathbf{x}_1, \quad \frac{r}{t} \rightarrow \mathbf{x}_2, \quad (119)$$

and subsequently convert the double sum in Eq. (111) into a two-dimensional ($d = 2$) integral over the square domain $\mathcal{D}_\square \equiv [0, 1]^2$ and use Stirling's approximation $n! \asymp \sqrt{2\pi n} (n/e)^n$ applied to the binomials in Eq. (111), in the regime where all t , l and their difference $t - l$ become large

$$\binom{t}{l} \asymp \sqrt{\frac{t}{2\pi l(t-l)}} \left(\frac{l}{t}\right)^{-l} \left(1 - \frac{l}{t}\right)^{l-t}. \quad (120)$$

In this manner we deduce the following exact asymptotic expression for the joint particle-charge MGF

$$G_{c,p}(\lambda_c, \lambda_p|t) \asymp \frac{t}{2\pi} \int_{\mathcal{D}_\square} d\mathbf{x} \frac{e^{t(f_c(\mathbf{x}_1, \mathbf{x}_2) + f_p(\mathbf{x}_1, \mathbf{x}_2))}}{\prod_{i=1}^2 \sqrt{x_i(1-x_i)}}, \quad (121)$$

with exponents

$$e^{f_c(\mathbf{x}_1, \mathbf{x}_2)} = \frac{\nu_-^x \nu_+^y \prod_{\epsilon \in \{\pm\}} \mu_\epsilon^{(|x_1 - x_2| + \epsilon(x_1 - x_2))/2}}{(\rho_+ \rho_-)^{-1} \prod_{i=1}^2 x_i^{x_i} (1-x_i)^{1-x_i}}, \quad (122)$$

$$e^{f_p(\mathbf{x}_1, \mathbf{x}_2)} = e^{\lambda_p(x_2 - x_1)}. \quad (123)$$

Bulk maxima. When $f(\mathbf{x})$ attains its maximum in the bulk of \mathcal{D}_\square , we find two maxima corresponding to two critical points located at $\mathbf{x}_\pm \equiv (x_{1,\pm}, x_{2,\pm})$, with coordinates

$$x_{1,\pm} = \frac{\nu_-(\lambda_p)}{\nu_-(\lambda_p) + [\mu_\pm(\lambda_c)]^{\mp 1}}, \quad (124)$$

$$x_{2,\pm} = \frac{\nu_+(\lambda_p)}{\nu_+(\lambda_p) + [\mu_\pm(\lambda_c)]^{\pm 1}}. \quad (125)$$

Critical points \mathbf{x}_\pm are however not always simultaneously present. In fact, depending on the ratio of densities $\kappa(\lambda_p) = \sqrt{\nu_+(\lambda_p)/\nu_-(\lambda_p)}$, extremal point \mathbf{x}_+ appears only below the diagonal of domain \mathcal{D}_\square , namely for $x_{1,+} \geq x_{2,+}$, equivalent to inequality $\mu_+(\lambda_c) \geq \kappa(\lambda_p)$; analogously \mathbf{x}_- appears above the diagonal for $x_{1,-} \leq x_{2,-}$ or,

equivalently, $\mu_-(\lambda_c) \geq \kappa^{-1}(\lambda_p)$. The two separatrices are determined by the conditions $\mu_\pm(\lambda_\pm) = [\kappa(\lambda_p)]^{\pm 1}$. For example, for $\lambda_p = 0$ this yields two solutions $\lambda_\pm^{(\epsilon)}$ (with $\epsilon \in \{\pm\}$) given by Eq. (91). At critical points \mathbf{x}_\pm , the exponent in Eq. (121) evaluates to

$$f_{c,p}^{(\pm)}(\lambda_c, \lambda_p) \equiv [f_c(\lambda_c) + f_p(\lambda_p)]|_{\mathbf{x}_\pm}, \quad (126)$$

given by the expressions in Eq. (81). Moreover, the Hessian precisely cancels out the square-root factor in Eq. (121). Writing $\boldsymbol{\lambda} \equiv (\lambda_c, \lambda_p)$, the joint MGF therefore assumes the following simple asymptotic form

$$G_{c,p}(\boldsymbol{\lambda}|t) \asymp \begin{cases} e^{t f_{c,p}^{(\pm)}(\boldsymbol{\lambda})} & \boldsymbol{\lambda} \in \mathcal{E}_\pm \\ \sum_{\epsilon \in \{\pm\}} e^{t f_{c,p}^{(\epsilon)}(\boldsymbol{\lambda})} & \boldsymbol{\lambda} \in \mathcal{E}_{+,-} \end{cases}, \quad (127)$$

where \mathcal{E}_\pm and $\mathcal{E}_{+,-}$ are given by Eq. (84) and Eq. (85), respectively. The above asymptotic formula does not apply for $\lambda_c \in \mathcal{S}_0$

Localization along the diagonal of \mathcal{D}_\square . When both conditions $\mu_+(\lambda_c) < \kappa(\lambda_p)$ and $\mu_-(\lambda_c) < \kappa^{-1}(\lambda_p)$ are simultaneously satisfied, the interior of \mathcal{D}_\square is devoid of any critical points. In this case, function f in the exponent in Eq. (121) attains a maximum on the diagonal at $\mathbf{x}_0 \equiv (x_0, x_0)$ with parallel and normal unit directions $\mathbf{n}_\parallel = \frac{1}{\sqrt{2}}(1, 1)$ and $\mathbf{n}_\perp = \pm \frac{1}{\sqrt{2}}(1, -1)$ (for $x_1 \gtrless x_2$), respectively. Solving for $\nabla_{\mathbf{n}_\parallel} f(\mathbf{x})|_{\mathbf{x}_0} = 0$, we find $x_0 = \sqrt{\nu_- \nu_+} / (1 + \sqrt{\nu_- \nu_+})$, with a non-zero second derivative $\nabla_{\mathbf{n}_\parallel}^2 f(\mathbf{x})|_{\mathbf{x}_0} = -(2 + \sqrt{\nu_- \nu_+} + 1/\sqrt{\nu_- \nu_+})$ and $\nabla_{\mathbf{n}_\perp} f(\mathbf{x})|_{\mathbf{x}_0} = -\sqrt{2} \log(\kappa^{\pm 1}(\lambda_p)/\mu_\pm(\lambda_c))$. For $\boldsymbol{\lambda} \in \mathbb{R} \times \mathbb{R}$, the MGF $G_{c,p}(\boldsymbol{\lambda}|t)$ restricted to intervals $\mathcal{S}_k \in \{+, 0, -\}$ (cf. Eqs. (87)) takes the form

$$G_{c,p}(\boldsymbol{\lambda} \in \mathcal{S}_k|t) \asymp W_k(\boldsymbol{\lambda}) e^{t F_k(\boldsymbol{\lambda})}, \quad (128)$$

with bulk branches $F_{c,p}^{(\pm)}(\boldsymbol{\lambda})$ (cf. Eq. (81)) and flat branch $F^{(0)}$ (cf. Eq. (82)), respectively. The 'weighting functions' read explicitly

$$W_\pm = 1, \quad (129)$$

$$W_0(\boldsymbol{\lambda}) = \frac{\prod_{\epsilon \in \{\pm\}} [\log(\kappa^\epsilon(\lambda_p) \mu_\epsilon^{-1}(\lambda_c))]^{-1}}{[4\pi t (2 + \sqrt{\nu_- \nu_+} + 1/\sqrt{\nu_- \nu_+})]^{1/2}}. \quad (130)$$

2. Scaled cumulants

In this section, we compute the scaled cumulant $s_n \equiv s_n^{(c)}$ for the general case of unequal densities $\rho_- \neq \rho_+$. Invoking Bryc's regularity condition (which follows from the Lee-Yang analysis carried out in Sec. V C below), the dominant branch (81) is always faithful and consequently all cumulants $c_n(t)$ grow linearly with time. In effect, scaled cumulants s_n are precisely the Taylor series coefficients of the charge-current SCGF $F(\lambda)$. The first two scaled cumulants are of particularly simple form

$$s_1 = (\rho_- - \rho_+) b_\pm, \quad (131)$$

$$s_2 = \mp(\rho_- - \rho_+) - b_\pm^2(\rho_+^2 + \rho_-^2 - 2\rho_\mp), \quad (132)$$

where the signature depends on which of the two bulk branches $F_{\pm}(\lambda)$ dominates at $\lambda = 0$ (recall that F_0 can never be dominant at $\lambda = 0$).

The degenerate case of equal densities, $\rho_{\pm} = \rho$, is rather exceptional. To begin with, the first cumulant $c_1(t)$ no longer grows linearly with time. As discussed in Sec. II, the lack of Bryc regularity implies that the SCGF $F(\lambda)$ is no longer a faithful generating function. In Appendix B, we infer how cumulants $c_n(t)$ grow at large times with aid of localization and Faà di Bruno's formula, establishing that $c_n \sim t^{n/2}$. To give a flavor, the first two cumulants read

$$c_1(t) \asymp (b_- - b_+) \frac{\chi_p(\rho)}{\pi} t^{1/2}, \quad (133)$$

$$c_2(t) \asymp \chi_p(\rho) (b_-^2 + b_+^2 - \frac{1}{\pi} (b_- - b_+)^2) t. \quad (134)$$

While in equilibrium, that is for $b_- = b_+$, all odd cumulants vanish identically, even cumulant still behave anomalously.

3. Typical fluctuations

We have already established that $G(\lambda|t)$ grows asymptotically with exponent $\alpha = 1$. This means that typical values of the integrated charge current through the origin in a time interval t are of the order $J_c(t) \sim \sqrt{t}$. By taking full advantage of the exact expression for the finite-time charge-current MGF $G(\lambda|t)$, we are in a position to deduce the associated PDF, denoted subsequently by $\mathcal{P}^{\text{typ}}(j)$ (here we have once again dropped the subscript 'c' by identifying $\lambda_c \rightarrow \lambda$ and $j_c \rightarrow j$). In the following, we treat the generic case with unequal densities $\rho_- \neq \rho_+$. The special case of equal densities is worked out in Appendix B.

Introducing the shifted cumulative charge current $\hat{J}_c(t) \equiv J_c(t) - c_1(t)$ by subtracting the first cumulant $c_1(t)$, we first define the dynamically rescaled time-dependent PDF

$$\mathcal{P}_{1/2}(\hat{J}_c|t) \equiv t^{1/2} \mathcal{P}(\hat{J}_c = t^{-1/2} \hat{J}_c(t)|t), \quad (135)$$

such that

$$e^{-\lambda c_1(t)} G(\lambda|t) = \int d\hat{J}_c \mathcal{P}_{1/2}(\hat{J}_c|t) e^{\lambda t^{1/2} \hat{J}_c}. \quad (136)$$

This yields, in the limit of large times, the stationary PDF

$$\mathcal{P}^{\text{typ}}(j) = \lim_{t \rightarrow \infty} \mathcal{P}_{1/2}(j = \hat{J}_c|t). \quad (137)$$

By introducing a dynamically rescaled counting field η via

$$\eta \equiv t^{1/2} \lambda, \quad (138)$$

and using $\lim_{t \rightarrow \infty} e^{-t^{1/2} \eta c_1(t)} G(t^{-1/2} \eta|t) = e^{s_2 \eta^2}$, we can express $\mathcal{P}^{\text{typ}}(j)$ via the inverse Laplace transform

(acting on a function of variable η)

$$\mathcal{P}^{\text{typ}}(j) = \mathcal{L}^{-1} \left[e^{s_2 \eta^2} \right] (j), \quad (139)$$

yielding a *Gaussian* PDF

$$\mathcal{P}^{\text{typ}}(j) = \frac{1}{\sqrt{2\pi\sigma_{\text{typ}}^2}} \exp \left[-\frac{j^2}{2\sigma_{\text{typ}}^2} \right], \quad \sigma_{\text{typ}}^2 = s_2. \quad (140)$$

Here $s_2 = F''(0)$ is the second scaled cumulant s_2 of the (faithful) SCGF $F(\lambda)$. Importantly, the obtained result implicitly depends on the dominant branch of $F_{\pm}(\lambda)$ at $\lambda = 0$. We have thus recovered the CLT behavior, as indeed guaranteed by Bryc regularity.




FIG. 15. Two branches of the joint particle-charge LDF $I_{c,p}(j_c, j_p)$. For $j_c \neq 0$, the two branches $I_{c,p}^{(\pm)}(j_c, j_p)$ are supported on disjoint domains \mathcal{J}_{\pm} separated by a gap, attaining their minima $j_p^{\pm}(j_c)$ in the interior of \mathcal{J}_{\pm} . For $j_c = 0$, the gap between the two domains closes while preserving convexity of $I_{c,p}(j_c, j_p)$. This time, one of the branches (in Figure \mathcal{J}_{-} , shown in blue) attains the minimum at $j_c = 0$ and since $j_p^{\pm}(j_c)$ is not differentiable at $j_c = 0$ the rate function $I_c(j_c)$ can acquire a corner point at the origin.

B. Dressing the particle rate function

This section includes a formal analysis of the solutions to the optimization problem given below by Eq. (147). Since now the particle and charge current both play a role and appear in the formulae simultaneously, we shall reinstate the subscript indices to avoid ambiguities. As we explain in turn, the computation of $I_c(j_c)$ can be formulated as a nested, *two-stage* optimization procedure. In the following, the LD rate function of the particle transfer $I_p(j_p)$ provides an input to the outlined optimization.

Our starting point will be the following joint rate function

$$I_{c,p}(j_c, j_p) = I_{c|p}(j_c, j_p) + I_p(j_p), \quad (141)$$

allowing us to express $I_c(j_c)$ through marginalization,

$$I_c(j_c) = \inf_{j_p} \{I_{c,p}(j_c, j_p)\}, \quad (142)$$

namely by minimizing the joint rate function over the domain \mathcal{J} of time-integrated rescaled particle current j_p , with j_c kept fixed.

By exploiting the fact that $|j_c| \leq |j_p|$, we split \mathcal{J} into two *disjoint* intervals \mathcal{J}_\pm such that $\mathcal{J} = \mathcal{J}_- \cup \mathcal{J}_+$. Assuming strict causality, such that $j_p \in [j_p^{\min}, j_p^{\max}]$, we define

$$\mathcal{J}_+(j_c) \equiv -[j_p^{\min}, |j_c|], \quad \mathcal{J}_-(j_c) \equiv [|j_c|, j_p^{\max}], \quad (143)$$

and correspondingly split the bivariate function $I_{c,p}(j_c, j_p)$ into two separate branches

$$I_{c,p}^{(\pm)}(j_c, j_p) \equiv I_{c|p}^{(\pm)}(j_c, j_p) + I_p(j_p). \quad (144)$$

The *split* conditional rate functions

$$I_{c|p}^{(\pm)}(j_c, j_p) \equiv I_{c|p}(j_p | j_c) \Big|_{j_p \in \mathcal{J}_\pm}, \quad (145)$$

have the following explicit form

$$I_{c|p}^{(\pm)}(j_c, j_p) = \sum_{\epsilon \in \{\pm\}} \frac{|j_p|}{2} (1 + \epsilon \xi) \log \left[\frac{1 + \epsilon \xi}{1 \mp \epsilon b_\pm} \right]. \quad (146)$$

The first stage of optimization involves minimizing the problems on the separate non-overlapping subdomains \mathcal{J}_\pm ,

$$I_c^{(\pm)}(j_c) \equiv \inf_{j_p \in \mathcal{J}_\pm} \{I_{c,p}^{(\pm)}(j_c, j_p)\}, \quad (147)$$

resulting in two branches of the rate function $I_c^{(\pm)}(j_c)$. In the second stage of optimization, the physical rate function $I_c(j_c)$ is determined by picking the minimum of the two branches $I_c^{(\pm)}$ for every given value of j_c ,

$$I_c(j_c) = \min\{I_c^{(\pm)}(j_c)\}. \quad (148)$$

We shall not attempt to give a fully general classification of solutions. Instead, we proceed with certain mild simplifying assumptions on the input rate function $I_p(j_p)$. We require that $I_p(j_p)$ is a strictly convex and twice differentiable everywhere on its domain, ensuring $I_p''(j_p) \equiv \partial_{j_p}^2 I_p(j_p) \geq 0$. Generic rate functions are expected to fulfill this technical assumptions, which appreciably simplifies the ongoing analysis. If the particle dynamics itself sustains a dynamical phase transition, there might of course be additional physical features that are not captured by our analysis. This is however beyond the scope of the present work. Our objective is to understand how the single-file and inertness properties lead to anomalous fluctuations associated with the charge transfer.

We proceed by investigating the minimization problems given by Eq. (147). Let j_p^\pm denote the minima of

$I_{c,p}^{(\pm)}(j_c, j_p)$. The solutions to Eqs. (147) are then given by $I_{c,p}^{(\pm)}(j_c, j_p^\pm) = I_c^{(\pm)}(j_c)$. It is important to stress that minima j_p^\pm are functions of j_c , i.e. $j_p^\pm = j_p^\pm(j_c)$. Noting that $I_{c,p}^{(\pm)}(j_c, j_p)$ are differentiable in both arguments, it follows that when $j_p^\pm(j_c)$ are differentiable functions of variable j_c , then so are the functions $I_c^{(\pm)}(j_c)$.

The second derivative with respect to the variable j_p of the conditional rate function $I_{c|p}(j_c, j_p)$, that is $I_{c|p}''(j_c, j_p) \equiv \partial_{j_p}^2 I_{c|p}(j_c, j_p)$, reads explicitly (with $\xi = j_c/|j_p|$, see Eq. (42))

$$I_{c|p}''(j_c, j_p) = \frac{1}{|j_p|} \frac{\xi^2}{1 - \xi^2}, \quad (149)$$

and is *non-negative* everywhere except at $j_c = 0$ where it vanishes, $I_{c|p}''(0, j_p) = 0$. Using that $I_p(j_p)$ is convex, this further implies that $I_{c,p}^{(\pm)}(j_c \neq 0, j_p)$ is strictly convex as a function of j_p . It then follows that the functions $I_{c,p}^{(\pm)}(j_c, j_p)$ possess unique minima at j_p^\pm , respectively. For this reason, we exclude for the moment the case $j_c = 0$ and continue with the general case $j_c \neq 0$.

Noting that for $|\xi| \rightarrow 1$ (i.e. when approaching the boundaries of domains \mathcal{J}_\pm from the left/right respectively) the magnitudes of the first derivatives

$$I_{c|p}'^{(\pm)}(j_c, j_p) = \pm \frac{1}{2} \log \left(\frac{1 - b_\pm^2}{1 - \xi^2} \right), \quad (150)$$

are unbounded and of different sign, we conclude (using that $I_{c,p}^{(\pm)}$ is convex and $I_p(j_p)$ bounded) that the minima j_p^\pm reside in the interior of domains $\mathcal{J}_\pm(j_p)$ (see Fig. 15). By invoking the implicit function theorem, it then follows from $I_{c,p}''^{(\pm)}(j_c \neq 0, j_p) > 0$, that the minima $j_p^\pm(j_c)$ are differentiable with respect to j_c , $j_p^\pm = j_p^\pm(j_c)$ for $j_c \neq 0$. Consequently, both branches $I_c^{(\pm)}(j_c \neq 0)$ in Eq. (147) are also differentiable functions.

Differentiability of $I_c(j_c)$ is however no longer guaranteed at the origin $j_c = 0$, representing a distinguished point. We note that at $j_c = 0$ ($\xi = 0$) we have a finite first derivative (see Eq. (150)) and a vanishing second derivative (see Eq. (149)). This means that the first derivatives of $I_{c,p}^{(\pm)}(0, j_p)$ with respect to j_p can, depending on state parameters, be either negative or positive, respectively, and consequently the corresponding minima of $I_{c,p}^{(\pm)}(0, j_p)$ can be attained at the boundaries of the optimization domains \mathcal{J}_\pm , implying that $j_p^\pm = 0$. Notice that even when $j_c = 0$, the minima j_p^\pm can still reside in the interior of $\mathcal{J}_\pm(0)$, in which case it follows immediately that $I_c^{(\pm)}(j_c)$ are differentiable by essentially reiterating the argument for the $j_c \neq 0$ case. Otherwise, when the minimum of $I_{c,p}^{(\pm)}(0, j_p)$ is located precisely on the boundary $\partial \mathcal{J}_\pm$ of the minimization domains \mathcal{J}_\pm (i.e. when $j_p^\pm = 0$), functions $j_p^\pm(j_c)$ cease to be differentiable at the origin, $j_c = 0$, where they develop a (convex) corner.




FIG. 16. Two branches $I_{c,p}^{(\pm)}(j_c, j_p)$ (with + and - shown by red and blue curves, respectively) of the joint particle-charge LDF $I_{c,p}(j_c, j_p)$, depicted for different dynamical regimes of the hardcore automaton: (a) regular regime, (b) tunneling subregime to the same bulk branch, (c) tunneling subregime to the other bulk branch and (d) corner regime. Thin curves correspond to different values of j_c with gray dots marking the location of minima. Thick curves indicate the flow of minima in each of the respective domains \mathcal{J}_{\pm} while continuously varying j_c . The global minima on $\mathcal{J} = \mathcal{J}_+ \cup \mathcal{J}_-$ follow the solid thick lines, while the dashed thick lines trace the subdominant minima. In regular regime (a), the global minimum always, i.e. for all j_c , belongs to one of the branches $I_{c,p}^{(\pm)}$, yielding a differentiable rate function $I_c(j_c)$. Likewise, the global minimum belongs to the same branch in tunneling subregime $[[+ \times +]]$ shown in panel (b), except that at $j_c = 0$ the minimum is found at $j_p = 0$, resulting in a non-differentiable (corner) point in the rate function. Panel (c) shows tunneling to another bulk branch labelled by $[[+ \times -]]$, where at $j_p = 0$ the global minimum shifts from $[[+]]$ over to $[[-]]$, also inducing a corner in the rate function. In the corner regime, shown in panel (d) for $[[+ \times -]]$, the trajectories of the local minima $j_p^{\pm}(j_c)$ do not meet at $j_p = 0$. Instead global minimum jumps from one branch to another at the critical charge current $j_{c\Delta}$, producing a non-differentiable (corner) point in the physical rate function $I(j_c)$ at $j_c = j_{c\Delta}$. Insets show the two branches $I_c^{(\pm)}(j_c)$ of the corresponding large-deviation rate function, with solid curves marking the physical rate function $I_c(j_c)$.

Inserting it into Eqs. (147), we conclude that the branches $I_c^{\pm}(j_c)$ are non-differentiable, also possessing a convex corner at $j_c = 0$.

Having worked out the formal structure of $I_c^{\pm}(j_c)$, we can now return to the outer optimization (148). We first establish that at least one of the branches $I_c^{(\pm)}(j_c)$ is not differentiable at $j_c = 0$, which readily follows from the fact that $\lim_{j_p \rightarrow 0} I_{c|p}^{(+)}(0, j_p) = \lim_{j_p \rightarrow 0} I_{c|p}^{(-)}(0, j_p)$, in turn implying that $I_{c,p}(0, j_p) = I_p(j_p)$ is continuous on its entire domain, i.e. for $j_p \in \mathcal{J}_+(0) \cup \mathcal{J}_-(0) = [-j_p^{\min}, j_p^{\max}]$. Writing shortly $I_{c,p}^{(\pm)}(0, j_p) \equiv \partial_{j_p} I_{c,p}^{(\pm)}(0, j_p)$ and taking into account that

$$\lim_{j_p \rightarrow 0^{\mp}} I_{c,p}^{(\pm)}(0, j_p) = \lim_{j_p \rightarrow 0^{\mp}} I_p'(j_p) \pm \frac{1}{2} \log(1 - b_{\pm}^2), \quad (151)$$

in conjunction with differentiability of $I_p(j_p)$ at the origin

$j_p = 0$, we infer the following inequality

$$\lim_{j_p \rightarrow 0^+} I_{c,p}^{(-)}(0, j_p) \geq \lim_{j_p \rightarrow 0^-} I_{c,p}^{(+)}(0, j_p). \quad (152)$$

Based on this, we conclude that $I_{c,p}(0, j_p)$ is a convex function on the entire domain of j_p , i.e. for $j_p \in [-j_p^{\min}, j_p^{\max}]$, despite non-differentiability of the branches $I_c^{(\pm)}(j_c)$ at $j_c = 0$. Since a convex function has only one minimum, it follows that at least one of the minima of $I_{c,p}^{\pm}(0, j_p)$ occur at the boundary $j_p = 0$, as illustrated in Figure 15.

The outlined nested minimization problem, given by Eq. (147) and Eq. (148), thus gives rise to several qualitatively different types of rate functions $I_c(j_c)$. There are four main cases which are exemplified in Figure 16.

C. The Lee–Yang theory

We now exhibit the dynamical critical phenomena in the framework of the Lee–Yang theory of phase transitions. To perform this analysis, we first express the MGF in terms of the exponential counting field $z \equiv e^\lambda$, yielding a Laurent series

$$G(z|t) = \sum_{n=-t}^t g_n(t) z^n = \mathcal{Z}_0(t) z^{-t} \sum_{j=1}^{2t} (z - z_j(t)). \quad (153)$$

The MGF $G(z|t)$ involves $2t$ dynamical (i.e. time-dependent) zeros $\{z_j(t)\}$ called Lee–Yang zeros. Due to reality of $J(t)$, we have that every zero z_j is paired with its complex-conjugate counterpart \bar{z}_j . In equilibrium, the detailed balance condition forces the zeros to combine into pairs z_j and $1/z_j$, implying a reflection symmetry $G(z|t) = G(z^{-1}|t)$. To facilitate the asymptotic analysis of $F(z|t) = t^{-1} \log G(z|t)$, it proves useful to separate the contributions of zeros, denoted hereafter by $F_{\text{LY}}(z|t)$, from the ‘regular potential’ $V(z|t) = -\log(z) + t^{-1} \log \mathcal{Z}_0(t)$ attributed to the non-polynomial prefactor in Eq. (153), i.e. perform the splitting $F(z|t) = V(z|t) + F_{\text{LY}}(z|t)$. The Lee–Yang zeros enter into the ‘singular part’

$$F_{\text{LY}}(z|t) = \sum_{j=1}^{2t} \log(z - z_j(t)). \quad (154)$$

in the form of logarithmic singularities.

The total number of zeros grows extensively with time t . We are interested in asymptotic growth of $G(z|t)$ at late times. In this limit, one commonly finds that the full fraction of zeros gradually approach each other before eventually forming condensates. One can expect (neglecting the possibility of accumulation points) the zeros to distribute along certain one-dimensional segments in the complex z -plane, or to (densely) concentrate inside certain two-dimensional domains. In the hardcore automaton, the latter scenario can be ruled out based on the asymptotic form of $G(z|t)$ given by Eq. (68). Notice that in the limit of large time any finite number (i.e. of vanishing fraction, subextensive in t) of zeros may be harmlessly disregarded.

1. Contour representation

We now return to our main example – the classical hardcore automaton. The task at hand is to explicitly determine the Lee–Yang contours. It turns out that the full fraction of zeros condense along a single contour or a union of contours which we denote hereafter by \mathcal{C}_z .

Introducing the (line) density $\rho(z)$ of Lee–Yang zeros,

$$\rho(z) = \lim_{t \rightarrow \infty} \frac{1}{t} \sum_{j=1}^{2t} \delta(z - z_j(t)), \quad (155)$$

distributed along a contour \mathcal{C}_z (with normalization $\int_{\mathcal{C}_z} dz \rho(z) = 2$), the limiting SCGF $F_{\text{LY}}(z) = \lim_{t \rightarrow \infty} F_{\text{LY}}(z|t)$ can be cast as a contour integral

$$F_{\text{LY}}(z) = \int_{\mathcal{C}_z} dw \mathcal{G}(z, w) \rho_z(w), \quad (156)$$

with respect to $\rho(w)$, where $\mathcal{G}(z, w) \equiv \log(z - w)$ is the Green’s function. In terms of the arc-length parameter s , the total number of zeros in an infinitesimal line element ds at position s on the contour \mathcal{C}_z is $t \rho_z(s) ds \sim \mathcal{O}(t)$.

Exploiting the formal analogy to the problem of 2D electrostatics, we further split $F_{\text{LY}}(z)$ into two real-valued potentials, $F_{\text{LY}}(z) = \Phi(z) + i\Psi(z)$. Interpreting the density $\rho_z(z)$ as a source, the real component $\Phi(z)$ fulfills the Poisson equation $\frac{1}{2\pi} \nabla^2 \Phi(z) = \rho_z(z)$. Potential $\Phi(z)$ is *continuous* across \mathcal{C}_z . In contrast, $\Psi(z)$ experiences a jump,

$$\delta\Psi(z(s)) \equiv \Psi(z(s) + i0) - \Psi(z(s) - i0), \quad (157)$$

proportional to the local density of zeros,

$$\rho_z(z(s)) = \frac{1}{2\pi} \frac{d}{ds} \delta\psi(z(s)), \quad z(s) \in \mathcal{C}_z. \quad (158)$$

By convention, infinitesimal shifts $\pm i0$ in the definition of $\delta\Psi$ pertain to values below (i.e. to the left) and above (to the right) of \mathcal{C}_z upon tracing the contour in the positive direction of increasing s .

2. Phase boundaries and the Stokes phenomenon

The function $G(z|t)$, now understood as a complex function of complex counting field z , can exhibit different asymptotic behavior depending on the state parameters. As briefly discussed in Sec. II and further expanded upon in Sec. IV, this can be traced back to coexistence of multiple competing extremal points (maxima) in the bulk of the integration domain \mathcal{D}_\square of $G(z|t)$, attributed to individual (meta)stable branches $F_k(z)$. To determine their ‘strength’, one has to compare the respective ‘Morse potentials’

$$\phi_k(z) = \text{Re } F_k(z), \quad (159)$$

obtained by analytic continuation of $F_k(z \in \mathbb{R})$ to the complex z -plane. Individual maxima of $G(z|t)$ undergo a non-trivial motion upon varying the counting variable $z \in \mathbb{C}$, and one can anticipate different competing maxima to dominate in different regions in the z -plane. Such behavior is generally known as the *Stokes phenomenon*. Alternatively, for certain values of z the dominant contribution to MGF $G(z|t)$ might not be due to localization around a bulk maximum but rather a round a maximum at the boundary $\partial\mathcal{D}_\square$.

The exchange of dominance associated with dynamical phases $F_k(z)$ and $F_\ell(z)$ occurs along certain one-dimensional contours, denoted by $\mathcal{C}_{k,\ell}^{(z)}$, separating apart

their respective regions of dominance \mathcal{R}_k and \mathcal{R}_ℓ . To determine $\mathcal{C}_{k,\ell}^{(z)}$, we first identify the ‘extended anti-Stokes lines’

$$\mathcal{A}_{k,\ell}^{(z)} = \{z \in \mathbb{C} \mid \phi_k(z) - \phi_\ell(z) = 0\}. \quad (160)$$

Here ‘extended’ signifies that $\mathcal{A}_{k,\ell}^{(z)}$ are not necessary the true anti-Stokes lines since we do not additionally account for the selection criteria, namely whether the compared k th and ℓ th phases even coexist. The latter can be inferred by analytic continuation of the selection rules, see Eq. (87). The Lee–Yang contour \mathcal{C} is then given by a union of anti-Stokes lines $\mathcal{C}_{k,\ell}^{(z)}$,

$$\mathcal{C}_z = \bigcup_{k,\ell \in \{+,0,-\}} \mathcal{C}_{k,\ell}^{(z)}. \quad (161)$$

We reemphasize that $\mathcal{C}_{k,\ell}^{(z)}$ are only a part of the extended anti-Stokes lines $\mathcal{A}_{k,\ell}^{(z)}$, whereas the remaining ‘ghost’ parts of $\mathcal{A}_{k,\ell}^{(z)}$ (depicted in figures by dashed lines, as e.g. shown in Figure 10) are of no physical relevance. We nonetheless choose to display them for convenience, given that full $\mathcal{A}_{k,\ell}^{(z)}$ in fact represent algebraic curves of finite genus.

3. Equilibrium and exceptional triple point

It turns out that several salient features of competing branches $F_k(z)$ neatly manifest themselves already in equilibrium states. However, equilibrium states are, as we corroborate in this section, rather singular. Here we perform the Lee–Yang analysis by putting $b_\pm \rightarrow b$ and $\rho_\pm \rightarrow \rho$, and writing shortly $\chi \equiv \chi_\rho(\rho) = \rho(1 - \rho)$.

To facilitate the computations, we find it most convenient to perform a change of variables $z \mapsto v(z)$ via a fractional linear transformation,

$$v(z) = \frac{z - 1}{z + 1}, \quad (162)$$

bijectionally transforming z -plane to v -plane in such a way that the real z -axis gets mapped to the open interval $(-1, 1)$ while the origin $z = 0$ stays put, that is $v(0) = 0$. The preimages $z = z(v)$ of the edge points $v = \pm 1$ are $z(-1) = 0$ and $z(1) = \infty$ (corresponding to $\lambda \rightarrow \pm\infty$). By the detailed balance condition, the set of zeros is invariant under reflection $v \mapsto -v$.

In the complex v -plane, the Lee–Yang contour by \mathcal{C}_v is a union of two closed contours $\mathcal{C}_{\pm,0}^{(v)}$, that is $\mathcal{C}_v = \mathcal{C}_{-,0}^{(v)} \cup \mathcal{C}_{+,0}^{(v)}$. It is sufficient to focus on the single physical contour, say $\mathcal{C}_{-,0}^{(v)}$ in the right-half of the v -plane with $\text{Re}(v) > 0$. The other half of \mathcal{C}_v is then simply obtained by reflecting $\mathcal{C}_{-,0}^{(v)}$ across the imaginary axis, yielding $\mathcal{C}_{+,0}^{(v)}$. We recall that in equilibrium the constant branch F_0 can never appear for $v \in \wp$. In fact, the anti-Stokes contour $\mathcal{C}_{+,-}^{(v)}$ is trivial.

Finite bias. We initially consider a general equilibrium case with a non-vanishing bias in the range $0 < b < 1$. The two bulk branches $F_\pm(v) = \log f_\pm(v)$ read explicitly

$$f_\pm(v) = 1 + \frac{4\chi v^2(b \mp v)^2}{(1 - v^2)(1 \mp 2bv + v^2)}. \quad (163)$$

The corresponding extended anti-Stokes lines $\mathcal{A}_{\pm,0}^{(v)[b]}$ are determined by the algebraic equations

$$|f_\pm(v)| = 1. \quad (164)$$

Writing $v \equiv v_1 + iv_2$, the above condition specifies an algebraic plane curve $P^{[b]}(v_1, v_2) = 0$ of degree $\deg P^{[b]} = 8$, with two real double (nodal) points located on the real axis at $v = 0$ and $v = -b$. The latter however lie on the ghost part the Lee–Yang contour, see Figure 17. The condensate of zeros is entirely supported on two mirror-symmetric contours $\mathcal{C}_{v,\pm}^{[b]}$ with $v \geq 0$, representing closed contours in the shape of a droplet that emanate out from the double point at the origin at an angle $\pi/4$.




FIG. 17. Complex phase diagram in v -plane (shown for equilibrium at finite bias, with parameters $\rho_\pm = 0.5$ and $b_\pm = 0.7$), featuring two bulk degenerate dynamical phases, $F_+(v)$ (red) and $F_-(v)$ (blue), and constant phase F_0 (white). The respective phase boundaries $\mathcal{C}_{\pm,0}^{(v)[b]}$ are shown by solid curves, while the ghost part of the extended anti-Stokes lines are marked by dashed curves. Gray dots show the Lee–Yang zeros for $t = 34$.

Having $F_k(v)$ available in a closed analytic form, one should in principle be able to find the expressions for density of zeros $\rho_v^{[b]}(s)$. We have not been able to derive a closed-form expression for the line density, mainly because we lack a useful local parametrization of the physical contours $\mathcal{C}_{v,\pm}^{[b]}$. Nonetheless, the density near the critical point origin at $v_0 = 0$ can be readily obtained by employing a local series expansion $v_2 = v_1 + [2(b^2 - 1)/b]v_1^2 + \mathcal{O}(v_1^3)$, allowing to subsequently compute $\nabla\psi \cdot \hat{t}|_{v=v_1}$ (with unit tangent $\hat{t} = (1, 1)/\sqrt{2}$ at $v_0 = 0$) to

the leading order in the arc-length parameter s , yielding

$$\rho_v^{[b]}(s) = \frac{4\chi(\rho)b^2}{\pi}s + \mathcal{O}(s^2). \quad (165)$$

We have thereby established *linear* dependence near the critical point v_0 at small s , namely $\rho_v^{[b]}(s) \simeq s$, and the impact angle of $\pi/4$, indicating that the phase transition across the anti-Stokes lines $\mathcal{A}_{\pm,0}^{(v)[b]}$ (and hence the physical phase boundary \mathcal{C}_v) is of the *second order*.




FIG. 18. Complex phase diagram in v -plane (shown for equilibrium with particle density $\rho = 0.5$ and no bias, $b = 0$), featuring two bulk degenerate dynamical phases, $F_+(v)$ (red) and $F_-(v)$ (blue), and constant phase F_0 (white). The respective phase boundaries $\mathcal{C}_{\pm,0}^{(v)[b]}$ are shown by solid curve. The ghost parts of the extended anti-Stokes lines, marked by dashed curves, overlap with each other. Gray dots show the Lee–Yang zeros for $t = 34$.

Zero bias. Upon taking the limit of zero bias, $b \rightarrow 0$, the MGF $G(v|t)$ acquires (next to retaining mirror symmetry $v \rightarrow -v$) an extra symmetry under charge conjugation $b \rightarrow -b$. As we now explain, this additional symmetry has important consequences. Firstly, the two phases $F_{\pm}(v)$ now become exactly degenerate, implying that at late times we have the following asymptotic form

$$G(v|t) \asymp e^{tF_+(v)} + e^{tF_-(v)} = 2[f_{\pm}(v)]^t, \quad (166)$$

with

$$f(v) = \frac{(1-4\chi)v^4 - 1}{v^4 - 1}. \quad (167)$$

Note also that the ghost parts of the anti-Stokes lines $\mathcal{A}_{\pm,0}^{(v)[b]}$ glue together into a single contour, (as depicted

in Figure 18) while the nodal points on the ghost contours approach the origin. Exactly for $b = 0$, we end up with a curve in the shape of a four-blade ‘boat propeller’, see Figure 18. This symmetry reflects the fact that $G(v|t)$ is a polynomial of v^2 of degree t , allowing for a reduction by performing another variable substitution $w = v^2$. This means that the algebraic curves $\mathcal{A}_{\pm,0}^{[b]}$ in the v -plane turn into quartic curves in the w -plane. By accordingly parametrizing $w = w_1 + iw_2$, the anti-Stokes line (once again determined by condition $|f_{\pm}^{[0]}| = 1$ with $f_{\pm}^{[0]}(w) = 1 + 4\chi w^2/(1-w^2)$) is equivalent to the algebraic curve

$$P^{[0]}(w_1, w_2) \equiv (1-2\chi)(w_2^2 + w_1^2)^2 + w_2^2 - w_1^2 = 0. \quad (168)$$

In practice, it proves convenient to use polar coordinates, namely writing $w = re^{i\vartheta}$, yielding the following compact parametrization of $P^{[0]}(w) = 0$,

$$r^2(\vartheta) = \frac{\cos(2\vartheta)}{1-2\chi}. \quad (169)$$

The reader can now recognize the famous lemniscate of Bernoulli – a rational quartic algebraic curve of genus zero. It is important to stress that the physical Lee–Yang contour $\mathcal{C}_w^{[0]}$ is not the whole lemniscate, but only the part lying in the right half-plane $\text{Re}(w) \geq 0$, i.e. along the interval of angles $\vartheta \in [-\pi/4, \pi/4]$. Moreover, contour $\mathcal{C}_w^{[0]}$ encloses a compact region in the complex w -plane attributed to the region of dominance of (analytically continued) degenerate bulk phases $F_{\pm}(w)$. As before, the exterior of $\mathcal{C}_w^{[0]}$ is the region of dominance of the flat branch.

This time, in the absence of bias, the line density $\rho_w(s)$, parametrized in terms of arc-length s , can actually be computed explicitly. In the interior of $\mathcal{C}^{[0]}$, the ψ -potential $\psi(r, \vartheta) \equiv \arctan(\Upsilon(r, \vartheta))$ can be parametrized explicitly

$$\Upsilon(r, \vartheta) = \frac{4\chi r^2 \cos(4\vartheta)}{r^4(1-4\chi) - 2r^2(1-2\chi)\cos(2\vartheta) + 1}, \quad (170)$$

From the jump discontinuity at the phase boundary $\mathcal{A}_w^{[0]}$ we deduce the line density as a function of polar angle ϑ ,

$$\rho_w(\vartheta) = \frac{1}{2\pi} \frac{16\chi(1-\chi)}{1-4\chi+8\chi^2+(4\chi-1)\cos(4\vartheta)}, \quad (171)$$

which now (by virtue of symmetry reduction) is normalized as $\int_{-\pi/4}^{\pi/4} d\vartheta \rho_w(\vartheta) = 1$. In the arc-length parametrization, $\rho_w(s) = (ds/d\vartheta)^{-1} \rho_w(\vartheta)$, the line density takes the form

$$\rho_w(s) = \frac{4\sqrt{2}\chi}{\pi} \sqrt{\frac{\frac{\pi}{4} - \theta(s)}{1-2\chi}}, \quad (172)$$

where $\vartheta(s) = -\text{am}((1-2\chi)^{-1/2}s - \text{F}(\frac{\pi}{4}, 2), 2)$ has been obtained by inverting $s(\vartheta) = (1-2\chi)^{-1/2}(\text{F}(\frac{\pi}{2}, 2) -$

$F(\vartheta, 2)$ (with $F(\vartheta, k)$ denoting the elliptic integral of the first kind and $\text{am}(u, k)$ the corresponding inverse function – the Jacobi amplitude). By finally expanding $\rho_w(s)$ in a Taylor series, we once more find the *linear* density of zeros near the origin,

$$\rho_w(s) = \frac{4\chi}{\pi}s + \mathcal{O}(s^5). \quad (173)$$




FIG. 19. Reduced complex phase diagram in w -plane (shown for equilibrium at zero bias with density $\rho = 1/2$) with a single merged dynamical phase (purple) invariant under charge conjugation, with the background constant phase F_0 (white). The extended anti-Stokes line is the lemniscate of Bernoulli, representing a quartic curve of genus zero. The physical phase boundary, shown by solid curve, is the right half. gray dots show the Lee–Yang zeros for $t = 34$.

The main upshot of the above analysis is that *in equilibrium*, dynamical phase transitions across the phase boundaries, $\mathcal{C}_{v,\pm}^{[b]}$ and $\mathcal{C}_{v,\pm}^{[b]}$, are always of the *second* order, irrespective of bias. According to Ehrenfest classification, a transition is characterized as order $n \geq 1$ if the density $\rho(s)$ (in v -plane or, equivalently, z -plane) diminishes as

$$\rho(s) \sim s^{n-1}, \quad (174)$$

in the proximity of a critical point. A first-order transition is associated with non-zero constant density across the critical point v_c at $s = 0$, while a second-order transition is characterized by a linearly decreasing density as small s . By expanding the $F(v)$ phase around v_c on both sides of the contour $\mathcal{C}_{v,\pm}^{[b]}$, namely

$$F_k(v) = \log f_k(v) = \sum_{n=0}^{\infty} \frac{f_n^{(k)}}{n!} (v - v_c), \quad (175)$$

the density of zeros $\rho(s)$ can be (to the leading order in s) read off from the jump discontinuity

$$\delta F_{k,\ell}(v) = \sum_{n \geq 0} \frac{\delta f_n^{(k,\ell)}}{n!} (v - v_c). \quad (176)$$

Using that $f_2^{(0)} = 0$ and $\delta f_2^{(\pm,0)} = f_2^{(\pm)} - f_2^{(0)}$, we correctly retrieve

$$\rho(s) = \frac{\delta f_2^{(\pm,0)}}{\pi} s + \mathcal{O}(s^2). \quad (177)$$

Employing polar coordinates, $v(r, \varphi) = r e^{i\varphi}$, the *impact angle* can be determined by tracing the contours,

$$\varphi_{\text{imp}} = \lim_{r \rightarrow 0} \varphi(r), \quad v \in \mathcal{C}_{\pm,0}^{[b]}, \quad (178)$$

By requiring the $\phi(v)$ is continuous across the phase boundaries, the ensuing condition $\sum_{n=-\infty}^{\infty} r^n \cos(n\varphi) = 0$ reduces, upon taking the limit $r \rightarrow 0$, to $\cos(n\varphi(r)) = 0$, yielding a finite set of n admissible impact angles of the form $\varphi_{\text{imp}}^{(k)} = [(2k+1)/n] \frac{\pi}{2}$ ($k \in \mathbb{N}$). For $n = 2$ specifically, the allowed angles are $\varphi_{\text{imp}} = \pm\pi/4$.

Exceptional triple point. Let us take a closer look at the emergent dynamical critical point v_t located at the origin of the complex v -plane and examine its formal properties. This critical point is distinguished by the fact that it lies on two distinct Lee–Yang contours. We note that in equilibrium at finite density and any bias, the SCGF $F(v)$ in the physical interval $v \in \wp$ features a discontinuity at $v_t = 0$.

For finite bias, $b > 0$, SCGF $F(v)$ has a discontinuous *third* derivative at v_t . In this view, v_t is criticality of *third order*, as already advocated in our previous work [74]. There is no controversy in this statement provided we stick with the conventional definition of a n -th order phase transition based on a discontinuity in the n -th derivative in the (dynamical) free-energy density (with all derivative or lower order being continuous). One should nevertheless be careful with reversing the logic. Contrary to a widely held belief,

an analytic real-valued (thermodynamic or dynamical) free energy $F(v)$ does not preclude emergence of criticality nor do the Lee–Yang zeros approaching and colliding with the physical axis necessarily render the free energy singular.

Indeed, $F(v)$ in equilibrium at zero bias provides an explicit counterexample.

The outlined Lee–Yang analysis helps resolving this subtle aspect. Let us first recall that in equilibrium, $F(v)$ for $v \in \wp$ involves only two bulk branches F_{\pm} that are degenerate in magnitude, while F_0 branch is entirely invisible. By contrast, any neighbourhood within a finite radius encompasses *three* distinct phases, indicating that v_t is indeed a *triple* point. It is not an ordinary triple point though, but rather an *exceptional* one; there are only two distinct phase boundaries (contours) emanating out of v_t .

We wish to emphasize that only upon analytically continuing $F(v)$ into complex v -plane we have enough information at hand for an unambiguous classification of a (dynamical) criticality. Specifically, since there are three phases involved, there are likewise three types of phase transitions: the pair of transitions $F_{\pm} \leftrightarrow F_0$ across the contours $\mathcal{C}_{v,\pm}^{[b]}$ (expelling the phase F_0 away from the physical interval), closing into the critical point v_t at the

origin at an angle of $\pm\pi/4$, and the ‘direct transition’ $F_+ \leftrightarrow F_-$ between the degenerate bulk phases mediated by v_t . In the former case, we find the difference of dynamical free energies,

$$\delta F_{\pm,0}(v) = \sum_{n=0}^{\infty} \frac{\delta f_n^{(\pm,0)}}{n!} v^n, \quad \delta f_{n<2}^{(\pm,0)} = 0, \quad (179)$$

is discontinuous at the second order, whereas in the case of $\delta F_{+,-}(v)$ we instead find, for finite bias $b > 0$, a third-order discontinuity. By contrast, for $b = 0$ we find $\delta F_{+,-} \equiv 0$ by virtue of the charge-conjugation symmetry.

To briefly summarize: in equilibrium states, $v_t = 0$ is (dynamical) critical triple point lying at the intersection of two distinct (extended) anti-Stokes lines (separating F_{\pm} phases encoding large deviations of the positive and negative charge transfer from the flat branch F_0) where two different (albeit degenerate) dynamical phases touch at a point. Upon crossing from F_- to F_+ through a singular (nodal) point one passes across two phase boundaries at the same time, see Fig. 18 and Fig. 19. We are unaware of a thermodynamic counterpart of such singular behavior.

VI. CONCLUSION

Studying classical single-file dynamical systems comprising of interacting charged particles, we uncovered a new type of dynamical universality reflected through anomalous statistical properties of fluctuating time-integrated current (charge transfer) in equilibrium and quasistationary nonequilibrium states. The unconventional behavior can be attributed to strongly broken ergodicity induced by the combination of the (I) *single-file property* and (II) *inertness of charges*. Any initial charge pattern therefore remains conserved throughout the time evolution, implying foliation of the phase space into exponentially many decoupled sectors.

By extending our previous work [74] to the nonequilibrium setting and generalizing it to a broad class of models, we have computed the statistics of particle and charge transfer between two semi-infinite partitions initially prepared in two different grand-canonical equilibrium states with a finite density of particles and charge. We derived the exact form of the finite-time joint moment generating function and examined its behavior at late times.

Our formal analysis of the scaled cumulant generating function and the associated large-deviation rate function revealed multifaceted structure. Our result concern two relevant timescales. In equilibrium states with vanishing bias, we analytically obtain a universal non-Gaussian probability distribution of typical cumulative charge currents and thereby rigorously established violation of the CLT property for the entire class of considered models. Anomalous fluctuations however become even more prominent away from equilibrium, where it turns out that

our systems generically sustain dynamical phase transitions of two sorts – either a first-order transition associated with an exchange of dominance among competing branches of the moment generating function attributed to distinct dynamical phases, or a second-order transition induced by a spontaneous breaking of convexity of a stable branch. The latter produces a finite flat segment in the scaled cumulant generating function, manifesting itself as a non-differentiable corner point (at the origin) in the large-deviation rate function.

The full counting statistics of charge transfer can be neatly obtained by applying the dressing operator to the particle counting statistics in a manner that avoids any model-specific information. Indeed, the entire phenomenology of dynamical phase transitions can be inferred from a formal analysis of the solutions to a problem of convex optimization associated with the joint particle-charge large-deviation rate function.

The most prominent universal (anomalous) feature of our models is a peculiar form of fluctuation relations. While the existing fluctuation theorems rely on either ergodicity or lack of first-order dynamical phase transitions, an explicit computation nevertheless confirms that the multivariate fluctuation relation of the Gallavotti–Cohen type is indeed inherited from that of the particle current and hence unconditionally satisfied, including in the dynamical regimes featuring dynamical criticality. What is more puzzling is the structure of the univariate rate function encoding large fluctuations of the cumulative charge current which in spite of time-reversal invariance generically experiences *spontaneous breaking* of the fluctuation relation mediated by dynamical phase transitions of first order. More specifically, we find that the univariate fluctuation relation of the charge-current rate function is only obeyed in a finite region in the global parameter space. Apart from that, even where it holds (in the so-called ‘regular regime’ and a subregime of the ‘tunneling regime’), the associated affinity depends solely on the state parameters of the dominant dynamical phase. In the complementary irregular regimes (involving a transition between the bulk branches), the univariate fluctuation relation is found to be violated. In the so-called ‘corner regime’, however, the symmetry is still obeyed locally for subcritical values of large currents.

To exemplify our methods, we devoted the second part of the paper to the study of a simple representative example of an exactly solvable classical deterministic and reversible cellular automaton. By deriving the exact FCS in bipartitioned nonequilibrium states, we expounded the salient features of the novel non-equilibrium universality and, by complexifying the counting field, computed the phase diagrams in different dynamical regimes. The equilibrium case, already studied in our previous work [74], is then recovered as a special degenerate case – the flat branch entirely disappears (and only remains visible for complex counting fields), while first-order critical points flow towards the origin of the complex plane. Restoration of detailed balance ensures that the bulk branches

become exactly degenerate. Meanwhile, the Lee–Yang contours – separating the bulk branches from the flat one – meet at the origin, giving rise to an exceptional triple critical point on the physical axis. The latter is in turn responsible for the lack of regularity, yielding divergent scaled cumulants. Curiously, in spite of dynamical criticality the fluctuation-dissipation relation remains preserved.

Outlook. A important aspect that remains to be addressed in future work is whether the outlined dynamical universality pertains strictly to the considered class of systems or whether it can be extended to a larger class of non-ergodic systems subjected to kinetic constraints. In this respect, we note that the property of pattern conservation (leading to exponential foliation of the phase space) alone is weaker than independently requiring both defining constraints (I) and (II). One could, for example, consider one-dimensional models where particle trajectories are correlated with the assignment of internal (e.g. charge) degrees of freedom. Our expectation is that such ‘pattern-conserving’ dynamical systems are most promising candidates to reveal similar anomalous behavior. Various instances of such models have already appeared in the recent literature, mainly in the context of quantum Hamiltonian systems and unitary circuits. Owing to presence of exponentially many dynamical subsectors, the model belongs to a large class of systems exhibiting ‘Hilbert space fragmentation’ [39, 98, 137].

To corroborate our view, we briefly discuss the prototypical example of the ‘folded’ XXZ quantum chain – an exactly solvable model with a local four-site interaction introduced in Refs. [138, 139], arising as an effective model for the large-anisotropy limit of the gapped Heisenberg spin-1/2 chain. A classical cellular automaton analogue of the folded XXZ model has been constructed in Ref. [140]. Apart from conserving the total number of particles (magnons), the number of kinks (separating the boundaries of uniformly magnetized domains made out of ‘frozen’ Bethe strings) becomes an additional (emergent) conserved quantity. The model consequently features an exponentially degenerate eigenspectrum owing to immobile spin-conserving patterns, in Ref. [138] dubbed ‘topological invariants’ (in the classical automaton, these are related to an exponential number of static domain-wall configurations [141]). There exists a dual version of the model with a three-site local update rule (not distinguishing between spin-flip conjugate configurations) obtained by performing the ‘site-bond transformation’ [140], where kinks become immobile particles and (mobile) solitons manifest themselves as bound states of kinks. As shown in Ref. [138], the dual folded model can be reinterpreted in terms charged degrees of freedom living on macrosites, directly exhibiting preservation of the charge pattern. A related exact mapping, yielding the so-called Massarani–Mathieu model, was given in Ref. [141] and employed recently in Ref. [42] to explain the subdiffusive nature of charge transport. According to a phenomenological hydrody-

dynamic picture proposed in Ref. [142], the statistics of the transferred charge, computed in equilibrium at vanishing density of magnetization, appears to match the M-Wright distribution derived earlier in Ref. [74] in the hardcore automaton (now established as one of the key universal properties of charged single-file systems studied in this work). Although this seems to suggest that the folded XXZ automaton (and perhaps many other related models) belong to the same universality class, our ‘dressing approach’ (which crucially rest on the inertness of charge degrees of freedom) is not directly applicable as, among other technical nuances, particles’ worldlines cannot be equipped with charges in an uncorrelated manner. Directly transcribing the full counting statistics of magnetization transfer in the original folded XXZ automaton to the counting statistics of charge transport in its dual counterpart is further obstructed by length-changing effects [42, 138].

Relatedly, we note that the folded XXZ model, together with other pattern-conserving systems, violate Bryc regularity. This follows from the fact that charge transport (in the unbiased sector) obeys the inequality $z_c > 1/\alpha$, where exponent α governs the particle dynamics, see Ref. [42]. This in turn implies divergent scaled cumulants arising due to an emergent dynamical critical point at the origin (assuming that the Lee–Yang zeros of the moment generating function approach the origin of the complex z -plane sufficiently rapidly [74], this would further imply lack of the CLT property). Based on this, we expect pattern-conserving systems to support phase transitions between two or more dynamical phases. Away from equilibrium, the critical point might leave the physical (i.e. real) axis in the z -plane. In the opposite scenario, a system would sustain a first-order dynamical phase transition leading to a spontaneous breaking of the univariate Gallavotti–Cohen relation. The Lee–Yang formalism offers a powerful tool to investigate these aspects.

Even though the considered models are interacting, the imposed conditions allow us to analytically compute the fluctuations of transmitted charge. As such they could serve as useful minimal model for addressing other questions in interacting dynamics. For example, the conditional distribution $\mathcal{P}_{p|c}^{[0]}(J_p|J_c)$ is bi-modal as a function of the integrated particle current in unbiased equilibrium. Interpreting the latter as an order-parameter similar to [143], this demonstrates that the order-parameter may become bi-modal when conditioning on anomalous currents even in equilibrium.

Before closing, we wish to shortly discuss the experimental relevance of our findings. Our hope is that the universal anomalous distribution of typical fluctuations is, thanks to its strongly non-Gaussian character, amenable to direct experimental detection. By analogy with conventional single-file systems, the aim would be to realize a Brownian motion of two different species of particles clogged in a narrow channel or particles that spread diffusively on a one-dimensional substrate. The proposed version of a simple symmetric exclusion process

of non-crossing Brownian trajectories of two colors (or its continuum limit) provides one basic mathematical model of such a process. A direct experimental detection of dynamical phase transitions in the structure of rare events seems less feasible. Nevertheless, one can hope to at least detect some traces of dynamical criticality or emergence of phase boundaries via probing temporal growth of the higher cumulants, following the lines of Refs. [144–149].

Acknowledgments. We thank J. M. Luck for correspondence. The authors benefited from discussions with K. Mallick, B. Doyon, B. Derrida and H. Spohn at the workshop on “Integrable and Chaotic dynamics” in Pokljuka, Slovenia (in July 2022). The work has been supported by ERC (European Research Council) Advanced grant 694544-OMNES (TP), by ARRS (Slovenian research agency) research program P1-0402 (ŽK, TP, EI), and by the SFB910 (project number 163436311) of the DFG (German Research Foundation) (JS). ŽK acknowledges support of the Milan Lenarčič foundation.

Appendix A: Dressing the moment generating function

To evaluate \mathfrak{D}_G , defined in Eq. (55), the main step involves computing the inverse bilateral Laplace transform $\mathcal{P}(J_p|t) = \mathfrak{L}^{-1}[G_p(w^{-1}|t)](J_p)$. This involves integrating along a counter-clockwise circular contour of radius R within the region of convergence of $G_p(w^{-1}|t)$,

$$\frac{1}{2\pi i} \oint_{|w|=R} \frac{dw}{w} w^{J_p} G_p(w^{-1}|t). \quad (\text{A1})$$

Using that PDF $\mathcal{P}(J_p|t)$ has finite support (implying that the region of convergence is $\mathbb{C} \setminus \{0, \infty\}$) for any finite time, we can interchange the order of contour integration and summations over J_p and J_c in \mathfrak{D}_G . Writing $\mathbf{z} \equiv (z_c, z_p)$, we thus arrive at the following integral representation

$$G_{c,p}(\mathbf{z}|t) = \frac{1}{2\pi i} \oint_{|w|=R} \frac{dw}{w} K(\mathbf{z}|w) G_p(w^{-1}|t), \quad (\text{A2})$$

with the time-independent kernel

$$K(\mathbf{z}|w) \equiv \sum_{J_p, J_c} z_c^{J_c} \mathcal{P}_{c|p}(J_c|J_p) z_p^{J_p} w^{J_p}. \quad (\text{A3})$$

The trouble we now face with evaluating the contour integral in Eq. (A2) is that the kernel $K(\mathbf{z}|w)$ has a vanishing region of convergence. This can be overcome by splitting the kernel into three separate pieces as follows. We introduce a pair of auxiliary projectors \mathcal{P}_{\pm} such that, when acting on a Laurent series in variable w , they project out only the polynomial of (strictly) positive powers in $w^{\pm 1}$. Similarly, let \mathcal{P}_0 projects out only the w^0 term, namely $K^{(0)}(\mathbf{z}|w) = 1$. In terms of the projectors we can perform the decomposition

$$K^{(k)}(\mathbf{z}|w) \equiv \mathcal{P}_k[K(\mathbf{z}|w)], \quad (\text{A4})$$

implying

$$G_{c,p}(\mathbf{z}|t) = \sum_{k \in \{+, 0, -\}} G_{c,p}^{(k)}(\mathbf{z}|t). \quad (\text{A5})$$

In particular, we have

$$G_{c,p}^{(0)}(\mathbf{z}|t) = \frac{1}{2\pi i} \oint_{|w|=R} \frac{dw}{w} G_p(w^{-1}|t), \quad (\text{A6})$$

Crucially, positive and negative parts $K^{(\pm)}(\mathbf{z}|w)$, when viewed as functions of w , now enjoy finite regions of convergence around $w = 0$ and $w = \infty$, respectively. To ensure that both regions of convergence have a non-vanishing overlap, we apply the variable transformation $w \mapsto w^{-1}$ to $K^{(+)}(\mathbf{z}|w)$ (as depicted in Fig. 20), obtaining the following representation

$$G_{c,p}^{(\pm)}(\mathbf{z}|t) = \oint_{|w|=R} \frac{dw}{2\pi i w} K^{(\pm)}(\mathbf{z}|w^{\mp 1}) G_p(w^{\pm 1}|t). \quad (\text{A7})$$




FIG. 20. Regions of convergence of integration kernels $K^{(-)}(\mathbf{z}|w)$ (blue) and $K^{(+)}(\mathbf{z}|w^{\pm 1})$ (red) in complex w -plane. Grey counter-clockwise circular contour represents the integration path within the intersection of both regions of convergence. The red and blue dots indicated the locations of poles of $w^{-1}K^{(+)}(\mathbf{z}|w^{-1})$ and $w^{-1}K^{(-)}(\mathbf{z}|w)$, respectively, while $w^{-1}K^{(0)}(\mathbf{z}|w) = w^{-1}$ is the simple pole at the origin (black dot).

In terms of linear combinations $J_{\pm} \equiv (J_p \pm J_c)/2 \in \mathbb{Z}_{\geq 0}$, we then have

$$1 + K^{(\pm)}(\mathbf{z}|w) = \sum_{J_+, J_-} \binom{J_p}{J_+} \prod_{\epsilon \in \{\pm\}} \left[\frac{1 \mp \epsilon b_{\pm} z_p^{\pm 1}}{2w} \frac{z_p^{\pm 1}}{z_c^{-\epsilon}} \right]^{J_{\epsilon}}, \quad (\text{A8})$$

and by evaluating the sum arrive at a compact expression

$$K^{(\pm)}(\mathbf{z}|w) = \frac{1}{1 - z_p^{\pm 1} \mu_{\pm}(z_c)/w} - 1, \quad (\text{A9})$$

with

$$|z_p^{\pm 1} \mu_{\pm}(z_c)| < |w|, \quad (\text{A10})$$

where $\mu_{\pm}(z_c) = \frac{1}{2}(z_c + z_c^{-1}) \mp b_{\pm} \frac{1}{2}(z_c - z_c^{-1})$. Notice that functions $w^{-1}K^{(\pm)}(\mathbf{z}|w)$ in the integrand in

Eq. (A7) involve two simple poles located at $w = 0$ and $w = z_p^{\pm 1} \mu_{\pm}(z_c)$. Using further that $G_p(w^{-1})$ is a Laurent series, the contour integrals can be evaluated simply by collecting the residues. Specifically, a circular integration contour within the common region of convergence,

$$|w| = R > \max\{|z_p \mu_+(z_c)|, |z_p^{-1} \mu_-(z_c)|\}, \quad (\text{A11})$$

encircles the three poles at $w = 0$ and at $w = z_p^{\pm 1} \mu_{\pm}(z_c)$, see Fig. 20. Computing the integrals (A7) therefore amounts to simply replacing the positive (+) and negative (-) powers of $z_p^{\pm n}$ with $z_p^{\pm n} [\mu_{\pm}(z)]^n$. As the outcome, applying the dressing operator \mathcal{D}_G is equivalent to the replacement rule (58).

Appendix B: Fluctuations at equal densities

Asymptotic moments. To infer the asymptotic behavior of $c_n(t)$ at equal density ($\rho_- = \rho_+$), we first compute the asymptotic of moments of the MGF $G(\lambda|t)$,

$$G^{(n)} \equiv \left. \frac{d^n G(\lambda|t)}{d\lambda^n} \right|_{\lambda=0}, \quad (\text{B1})$$

with the double-sum representation

$$G^{(n)}(t) = \rho^{2t} \sum_{l=0}^t \sum_{r=0}^t \binom{t}{l} \binom{t}{r} \nu^{l+r} \gamma b_{\gamma} d_n^{|l-r|}(b), \quad (\text{B2})$$

where $\gamma = \text{sgn}(l-r)$ and $d_n^k(b) = [d^n \mu_b^k(\lambda)/d\lambda^n]_{\lambda=0}$ with

$$\mu_b(\lambda) \equiv \cosh(\lambda) - b \sinh(\lambda). \quad (\text{B3})$$

We separate the computation into several steps. To begin with, we define a sequence of auxiliary double sums

$$S_n = \rho^{2t} \sum_{l=0}^t \sum_{r=0}^t \binom{t}{l} \binom{t}{r} \nu^{l+r} |l-r|^n. \quad (\text{B4})$$

Rescaling the coordinates as $l/t \rightarrow x_1$, $r/t \rightarrow x_2$ and approximating binomials using the Stirling's approximation, the double sum can be cast as an integral over the square domain \mathcal{D}_{\square}

$$S_n \asymp \frac{t^{n+1}}{2\pi} \int_{\mathcal{D}_{\square}} dx_1 dx_2 h_n(x_1, x_2) e^{t f(x_1, x_2)}, \quad (\text{B5})$$

with $h_n(x_1, x_2) = |x_1 - x_2|^n / \sqrt{x_1(1-x_1)x_2(1-x_2)}$ and

$$f(x_1, x_2) = \log \left[\frac{\rho^2 \nu^{x_1+x_2}}{\prod_{i=1}^2 x_i^{x_i} (1-x_i)^{1-x_i}} \right]. \quad (\text{B6})$$

The critical point $\mathbf{x}_{\star} = (x_{1,\star}, x_{2,\star})$, determined from the solution to $\nabla f(\mathbf{x}_{\star}) = 0$, sits at $x_{1,\star} = x_{2,\star} = 1 - \rho$, while $\det H[f](\mathbf{x}_{\star}) = [\rho(1-\rho)]^{-2}$. Here we run in an obstacle, as functions $h_n(x_1, x_2)$ in the integrand evaluate to zero,

i.e. $h_n(\mathbf{x}_{\star}) = 0$. This can be overcome by developing $f(x_1, x_2)$ as a Taylor series up to quadratic order about the critical points

$$f(x_1, x_2) \approx \frac{[x_1 - (1-\rho)]^2 + [x_2 - (1-\rho)]^2}{2\rho(1-\rho)}, \quad (\text{B7})$$

and subsequently performing integration in the rotated frame, with coordinates $u \equiv x_1 + x_2$, $v \equiv x_1 - x_2$, yielding

$$S_n \asymp \frac{t^{n+1}}{\pi} \int du \int_{v>0} dv \mathcal{S}(u, v), \quad (\text{B8})$$

with

$$\mathcal{S} \equiv \frac{v^n e^{-t[(u-u_{\star})^2 + v^2]/4\chi_p}}{\sqrt{(u^2 - v^2)(1 - \frac{1}{2}(u+v))(1 - \frac{1}{2}(u-v))}}. \quad (\text{B9})$$

Since both maxima lie symmetrically about $v = 0$, it sufficient to keep only the $v > 0$ region. Using that the expression under the square root localizes to the value χ_p^{-1} and, anticipating localization, extending the range of integration, we have

$$S_n(t) \asymp \frac{t^{n+1}}{2\pi\chi_p} \int_{-\infty}^{\infty} du e^{-\frac{t(u-u_{\star})^2}{4\chi_p}} \int_0^{\infty} dv v^n e^{-\frac{tv^2}{4\chi_p}}. \quad (\text{B10})$$

After computing the Gaussian integrals we arrive at a compact expression

$$S_n(t) \asymp s_n (\chi_p t)^{n/2}, \quad s_n \equiv \frac{2^{n-1}}{\sqrt{\pi}} \Gamma\left(\frac{n+1}{2}\right). \quad (\text{B11})$$

The asymptotics of cumulants $c_n(t)$ can be deduced from that of the moments $G^{(n)}$. Writing shortly $g^{(p)}(x) \equiv d^p g/dx^p$, and introducing the incomplete Bell polynomials $B_{n,m} = B_{n,m}(\{x_j\}_{j=1}^{n-m+1})$ with the generating series

$$\exp \left[z \sum_{k=1}^{\infty} x_k \frac{y^k}{k!} \right] = 1 + \sum_{n=1}^{\infty} \frac{y^n}{n!} \sum_{m=1}^n z^m B_{n,m}, \quad (\text{B12})$$

the asymptotic behavior of higher moments $G^{(n)}(t)$ can be expressed in terms of $S_m(t)$ with aid of the Faá di Bruno's formula

$$\frac{d^n f(g(x))}{dx^n} = \sum_{m=1}^n f^{(m)}(g(x)) B_{n,m}, \quad (\text{B13})$$

where $B_{n,m} \equiv B_{n,m}(\{g^{(p)}\}_{p=1}^{n-m+1})$. Coefficients d_n^k are polynomials in variable k of degree $n/2$,

$$d_n^k = \sum_{m=1}^n \frac{k}{(k-m)!} B_{n,m}(-b, 1, -b, 1, \dots). \quad (\text{B14})$$

Replacing monomials k^m by S_m in Eq. (B2) we obtain $G^{(n)}(\{S_j\})$. With another application of Eq. (B13), we finally obtain

$$c_n(t) \asymp \sum_{m=1}^n (-1)^{m-1} (m-1)! B_{n,m}(\{G^{(j)}(t)\}). \quad (\text{B15})$$

Generating function for asymptotic cumulants. Introducing dynamically rescaled counting fields,

$$\omega_{\pm}(\lambda) \equiv b_{\pm}(\chi_p t)^{1/2} \lambda, \quad (\text{B16})$$

the asymptotic moments

$$G^{(n)}(t) \asymp c_n (b_-^n + (-b_+)^n) (\chi_p t)^{n/2}, \quad (\text{B17})$$

can be resummed into the following generating series

$$\Omega(\{\omega_{\pm}(\lambda)\}) = \log \left[\frac{E_{1/2}(\omega_-) + E_{1/2}(-\omega_+)}{2} \right], \quad (\text{B18})$$

where $E_{1/2}(\pm x) = e^{x^2} [1 \pm \text{erf}(x)]$. Asymptotics of the cumulants is retrieved as $c_n(t) \asymp (d^n/d\lambda^n) \Omega(\{\eta_{\pm}(\lambda)\})|_{\lambda=0}$ or, using the chain rule as

$$c_n(t) \asymp \left[\sum_{\epsilon \in \{\pm\}} \left(\frac{d\omega_{\epsilon}}{d\lambda} \right)^n \frac{\partial^n}{\partial \omega_{\epsilon}^n} \right] \Omega(\{\omega_{\pm}(\lambda)\}). \quad (\text{B19})$$

Typical fluctuations. Setting the densities ρ_{\pm} to be equal (while leaving b_{\pm} arbitrary), the PDF $\mathcal{P}_{\text{typ}}^{(\rho)}(j)$ of typical charge-current fluctuations can be recovered from Eq. (B18). By dynamically rescaling the generating function $\Omega(\{\omega_{\pm}(\lambda)\})$ with $\lambda \rightarrow t^{-1/2} \eta$, we explicitly compute

$$\mathcal{G}^{(\rho)}(\eta) \equiv \lim_{t \rightarrow \infty} \left[e^{-c_1(t) t^{1/2} \eta} G(\lambda(\eta)|t) \right]. \quad (\text{B20})$$

In terms of $\omega_{\pm} = \sqrt{\chi_p} b_{\pm} \eta$, we find

$$\log \mathcal{G}^{(\rho)}(\eta) = \frac{\omega_+ - \omega_-}{\sqrt{\pi}} \Omega(\omega_{\pm}). \quad (\text{B21})$$

The PDF $\mathcal{P}_{\text{typ}}^{(\rho)}(j_c)$ is given implicitly by the inverse Laplace transform $\mathfrak{L}^{-1}[\mathcal{G}^{(\rho)}(\eta)]$ of the MGF, reading compactly

$$\mathcal{G}^{(\rho)}(\eta) = e^{\pi^{-1/2}(\omega_+ - \omega_-)} \frac{1}{2} \sum_{\epsilon \in \{\pm\}} E_{1/2}(-\epsilon \omega_{\epsilon}). \quad (\text{B22})$$

Notice that the extra multiplicative factor relative to the generating series, cf. Eq. (B18), comes from the first cumulant $c_1(t)$ (i.e. average time-integrated current) which has been subtracted. The results for the biased equilibrium ensembles is recovered in the limit $b_{\pm} \rightarrow b$, yielding a Gaussian MGF $G^{(\rho)}(\eta) = \exp[b^2 \chi_p \eta^2]$ and hence the PDF is a normal distribution with variance $\sigma = 2\chi_p b^2$.

-
- [1] D. J. Evans and D. J. Searles, *Advances in Physics* **51**, 1529 (2002).
- [2] D. J. Evans and D. J. Searles, *Phys. Rev. E* **50**, 1645 (1994).
- [3] E. Sevcik, R. Prabhakar, S. R. Williams, and D. J. Searles, *Annual Review of Physical Chemistry* **59**, 603 (2008).
- [4] D. J. Evans, E. G. D. Cohen, and G. P. Morriss, *Physical Review Letters* **71**, 3616 (1993).
- [5] G. Gallavotti and E. G. D. Cohen, *Physical Review Letters* **74**, 2694 (1995).
- [6] J. Kurchan, *Journal of Physics A: Mathematical and General* **31**, 3719 (1998).
- [7] J. L. Lebowitz and H. Spohn, *Journal of Statistical Physics* **95**, 333 (1999).
- [8] P. I. Hurtado, C. Pérez-Espigares, J. J. del Pozo, and P. L. Garrido, *Proceedings of the National Academy of Sciences* **108**, 7704 (2011).
- [9] C. Maes, *Journal of Statistical Physics* **95**, 367 (1999).
- [10] D. Andrieux and P. Gaspard, *Journal of Statistical Mechanics: Theory and Experiment* **2006**, P01011 (2006).
- [11] D. Andrieux and P. Gaspard, *Journal of Statistical Mechanics: Theory and Experiment* **2008**, P11007 (2008).
- [12] V. Jakšić, Y. Ogata, Y. Pautrat, and C.-A. Pillet, in *Quantum Theory from Small to Large Scales* (Oxford University Press, 2012) pp. 213–410.
- [13] P. Gaspard, *New Journal of Physics* **15**, 115014 (2013).
- [14] S. Hild, T. Fukuhara, P. Schauß, J. Zeiher, M. Knap, E. Demler, I. Bloch, and C. Gross, *Phys. Rev. Lett.* **113**, 147205 (2014).
- [15] Y. Tang, W. Kao, K.-Y. Li, S. Seo, K. Mallayya, M. Rigol, S. Gopalakrishnan, and B. L. Lev, *Phys. Rev. X* **8**, 021030 (2018).
- [16] P. N. Jepsen, J. Amato-Grill, I. Dimitrova, W. W. Ho, E. Demler, and W. Ketterle, *Nature* **588**, 403 (2020).
- [17] S. Scherg, T. Kohler, P. Sala, F. Pollmann, B. H. M. I. Bloch, and M. Aidelsburger, *Nature Communications* **12**, 4490 (2021).
- [18] D. Wei, A. Rubio-Abadal, B. Ye, F. Machado, J. Kemp, K. Srakaew, S. Hollerith, J. Rui, S. Gopalakrishnan, N. Y. Yao, I. Bloch, and J. Zeiher, *Science* **376**, 716 (2022).
- [19] L. Vidmar and M. Rigol, *Journal of Statistical Mechanics: Theory and Experiment* **2016**, 064007 (2016).
- [20] E. Ilievski, J. De Nardis, B. Wouters, J.-S. Caux, F. H. L. Essler, and T. Prosen, *Phys. Rev. Lett.* **115**, 157201 (2015).
- [21] O. A. Castro-Alvaredo, B. Doyon, and T. Yoshimura, *Phys. Rev. X* **6**, 041065 (2016).
- [22] B. Bertini, M. Collura, J. De Nardis, and M. Fagotti, *Phys. Rev. Lett.* **117**, 207201 (2016).
- [23] J. De Nardis, D. Bernard, and B. Doyon, *Physical Review Letters* **121** (2018), 10.1103/physrevlett.121.160603.
- [24] B. Doyon, *SciPost Physics Lecture Notes* (2020), 10.21468/scipostphyslectnotes.18.
- [25] B. Bertini, F. Heidrich-Meisner, C. Karrasch, T. Prosen, R. Steinigeweg, and M. Žnidarič, *Reviews of Modern Physics* **93** (2021), 10.1103/revmodphys.93.025003.
- [26] E. Ilievski, J. De Nardis, M. Medenjak, and T. Prosen, *Physical Review Letters* **121** (2018), 10.1103/physrevlett.121.230602.

- [27] M. Ljubotina, M. Žnidarič, and T. Prosen, *Physical Review Letters* **122** (2019), 10.1103/physrevlett.122.210602.
- [28] J. De Nardis, M. Medenjak, C. Karrasch, and E. Ilievski, *Phys. Rev. Lett.* **123**, 186601 (2019).
- [29] Ž. Krajnik, E. Ilievski, and T. Prosen, *SciPost Phys.* **9**, 38 (2020).
- [30] E. Ilievski, J. D. Nardis, S. Gopalakrishnan, R. Vasseur, and B. Ware, *Physical Review X* **11** (2021), 10.1103/physrevx.11.031023.
- [31] V. B. Bulchandani, S. Gopalakrishnan, and E. Ilievski, *Journal of Statistical Mechanics: Theory and Experiment* **2021**, 084001 (2021).
- [32] A. J. Friedman, S. Gopalakrishnan, and R. Vasseur, *Phys. Rev. B* **101**, 180302 (2020).
- [33] J. Lopez-Piqueres, B. Ware, S. Gopalakrishnan, and R. Vasseur, *Phys. Rev. B* **103**, L060302 (2021).
- [34] J. Lopez-Piqueres, S. Gopalakrishnan, and R. Vasseur, *Journal of Physics A: Mathematical and Theoretical* **55**, 234005 (2022).
- [35] J. De Nardis, S. Gopalakrishnan, R. Vasseur, and B. Ware, *Phys. Rev. Lett.* **127**, 057201 (2021).
- [36] P. W. Claeys, A. Lamacraft, and J. Herzog-Arbeitman, *Phys. Rev. Lett.* **128**, 246603 (2022).
- [37] J. De Nardis, S. Gopalakrishnan, R. Vasseur, and B. Ware, arXiv preprint arXiv:2109.13251 (2021).
- [38] A. Bastianello, A. D. Luca, and R. Vasseur, *Journal of Statistical Mechanics: Theory and Experiment* **2021**, 114003 (2021).
- [39] P. Sala, T. Rakovszky, R. Verresen, M. Knap, and F. Pollmann, *Physical Review X* **10** (2020), 10.1103/physrevx.10.011047.
- [40] A. Gromov, A. Lucas, and R. M. Nandkishore, *Phys. Rev. Research* **2**, 033124 (2020).
- [41] K. T. Grosvenor, C. Hoyos, F. Peña Benitez, and P. Surówka, *Phys. Rev. Research* **3**, 043186 (2021).
- [42] J. Feldmeier, W. Witczak-Krempa, and M. Knap, arXiv preprint arXiv:2205.07901 (2022).
- [43] H. Touchette, *Physics Reports* **478**, 1 (2009).
- [44] R. S. Ellis, *Entropy, Large Deviations, and Statistical Mechanics* (Springer Berlin Heidelberg, 2006).
- [45] M. Esposito, U. Harbola, and S. Mukamel, *Reviews of Modern Physics* **81**, 1665 (2009).
- [46] L. Bertini, A. D. Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, *Physical Review Letters* **87** (2001), 10.1103/physrevlett.87.040601.
- [47] L. Bertini, A. D. Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, *Journal of Statistical Physics* **107**, 635 (2002).
- [48] K. Mallick, H. Moriya, and T. Sasamoto, arXiv preprint arXiv:2202.05213 (2022).
- [49] L. Bertini, D. Gabrielli, and J. L. Lebowitz, *Journal of Statistical Physics* **121**, 843 (2005).
- [50] E. Bettelheim, N. R. Smith, and B. Meerson, *Phys. Rev. Lett.* **128**, 130602 (2022).
- [51] E. Bettelheim, N. R. Smith, and B. Meerson, arXiv preprint arXiv:2204.06278 (2022).
- [52] T. Bodineau and B. Derrida, *Physical Review Letters* **92** (2004), 10.1103/physrevlett.92.180601.
- [53] P. I. Hurtado and P. L. Garrido, *Physical Review E* **81** (2010), 10.1103/physreve.81.041102.
- [54] K. Saito and A. Dhar, *Physical Review Letters* **107** (2011), 10.1103/physrevlett.107.250601.
- [55] M. Žnidarič, *Physical Review E* **89** (2014), 10.1103/physreve.89.042140.
- [56] O. Shpielberg and E. Akkermans, *Physical Review Letters* **116** (2016), 10.1103/physrevlett.116.240603.
- [57] C. Gutiérrez-Ariza and P. I. Hurtado, *Journal of Statistical Mechanics: Theory and Experiment* **2019**, 103203 (2019).
- [58] F. Angeletti, H. Touchette, E. Bertin, and P. Abry, *Journal of Statistical Mechanics: Theory and Experiment* **2014**, P02003 (2014).
- [59] H. G. Duhart, P. Mörters, and J. Zimmer, *Potential Analysis* **48**, 301 (2017).
- [60] R. J. Harris, A. Rákos, and G. M. Schütz, *Journal of Statistical Mechanics: Theory and Experiment* **2005**, P08003 (2005).
- [61] A. Rákos and R. J. Harris, *Journal of Statistical Mechanics: Theory and Experiment* **2008**, P05005 (2008).
- [62] R. J. Harris and H. Touchette, *Journal of Physics A: Mathematical and Theoretical* **42**, 342001 (2009).
- [63] R. van Zon and E. G. D. Cohen, *Physical Review Letters* **91** (2003), 10.1103/physrevlett.91.110601.
- [64] A. D. Luca, J. Viti, D. Bernard, and B. Doyon, *Physical Review B* **88** (2013), 10.1103/physrevb.88.134301.
- [65] K. Saito and A. Dhar, *Physical Review Letters* **99** (2007), 10.1103/physrevlett.99.180601.
- [66] T. Yoshimura, *Journal of Physics A: Mathematical and Theoretical* **51**, 475002 (2018).
- [67] D. Bernard and B. Doyon, *Journal of Physics A: Mathematical and Theoretical* **45**, 362001 (2012).
- [68] D. Bernard and B. Doyon, *Journal of Statistical Mechanics: Theory and Experiment* **2016**, 033104 (2016).
- [69] H. Moriya, R. Nagao, and T. Sasamoto, *Journal of Statistical Mechanics: Theory and Experiment* **2019**, 063105 (2019).
- [70] O. Gamayun, O. Lychkovskiy, and J.-S. Caux, *SciPost Physics* **8** (2020), 10.21468/scipostphys.8.3.036.
- [71] M. Žnidarič, *Physical Review Letters* **112** (2014), 10.1103/physrevlett.112.040602.
- [72] B. Buča and T. Prosen, *Physical Review Letters* **112** (2014), 10.1103/physrevlett.112.067201.
- [73] B. Buča, J. P. Garrahan, T. Prosen, and M. Vanicat, *Phys. Rev. E* **100**, 020103 (2019).
- [74] Ž. Krajnik, J. Schmidt, V. Pasquier, E. Ilievski, and T. Prosen, *Physical Review Letters* **128** (2022), 10.1103/physrevlett.128.160601.
- [75] A. Kuniba, G. Misguich, and V. Pasquier, *Journal of Physics A: Mathematical and Theoretical* (2022), 10.1088/1751-8121/ac6d8c.
- [76] J. Myers, J. Bhaseen, R. J. Harris, and B. Doyon, *SciPost Physics* **8** (2020), 10.21468/scipostphys.8.1.007.
- [77] B. Doyon and J. Myers, *Annales Henri Poincaré* **21**, 255 (2019).
- [78] G. Peretto and B. Doyon, *SciPost Physics* **10** (2021), 10.21468/scipostphys.10.5.116.
- [79] B. Doyon, G. Peretto, T. Sasamoto, and T. Yoshimura, arXiv e-prints, arXiv (2022).
- [80] D. Bernard and B. Doyon, *Journal of Physics A: Mathematical and Theoretical* **46**, 372001 (2013).
- [81] J. P. Garrahan, R. L. Jack, V. Lecomte, E. Pitard, K. van Duijvendijk, and F. van Wijland, *Physical Review Letters* **98** (2007), 10.1103/physrevlett.98.195702.
- [82] J. P. Garrahan, R. L. Jack, V. Lecomte, E. Pitard, K. van Duijvendijk, and F. van Wijland, *Journal of Physics A: Mathematical and Theoretical* **42**, 075007 (2009).

- [83] P. T. Nyawo and H. Touchette, *EPL (Europhysics Letters)* **116**, 50009 (2016).
- [84] A. Lazarescu, *Journal of Physics A: Mathematical and Theoretical* **50**, 254004 (2017).
- [85] R. L. Jack, T. Nemoto, and V. Lecomte, *Journal of Statistical Mechanics: Theory and Experiment* **2020**, 053204 (2020).
- [86] T. Bodineau and B. Derrida, *Physical Review E* **72** (2005), 10.1103/physreve.72.066110.
- [87] C. P. Espigares, P. L. Garrido, and P. I. Hurtado, *Physical Review E* **87** (2013), 10.1103/physreve.87.032115.
- [88] Y. Baek, Y. Kafri, and V. Lecomte, *Physical Review Letters* **118** (2017), 10.1103/physrevlett.118.030604.
- [89] Y. Baek, Y. Kafri, and V. Lecomte, *Journal of Physics A: Mathematical and Theoretical* **51**, 105001 (2018).
- [90] P. I. Hurtado and P. L. Garrido, *Physical Review Letters* **107** (2011), 10.1103/physrevlett.107.180601.
- [91] D. Manzano and P. I. Hurtado, *Physical Review B* **90** (2014), 10.1103/physrevb.90.125138.
- [92] A. W. C. Lau, D. Lacoste, and K. Mallick, *Physical Review Letters* **99** (2007), 10.1103/physrevlett.99.158102.
- [93] D. Lacoste and K. Mallick, *Physical Review E* **80** (2009), 10.1103/physreve.80.021923.
- [94] P. Krapivsky, K. Mallick, and T. Sadhu, *Physical Review Letters* **113** (2014), 10.1103/physrevlett.113.078101.
- [95] P. L. Krapivsky, K. Mallick, and T. Sadhu, *Journal of Statistical Mechanics: Theory and Experiment* **2015**, P09007 (2015).
- [96] P. L. Krapivsky, K. Mallick, and T. Sadhu, *Journal of Statistical Physics* **160**, 885 (2015).
- [97] T. Imamura, K. Mallick, and T. Sasamoto, *Physical Review Letters* **118** (2017), 10.1103/physrevlett.118.160601.
- [98] V. Khemani, M. Hermele, and R. Nandkishore, *Physical Review B* **101** (2020), 10.1103/physrevb.101.174204.
- [99] K. Mallick, *Physica A: Statistical Mechanics and its Applications* **418**, 17 (2015).
- [100] M. Medenjak, K. Klobas, and T. Prosen, *Phys. Rev. Lett.* **119**, 110603 (2017).
- [101] K. Klobas, M. Medenjak, and T. Prosen, *Journal of Statistical Mechanics: Theory and Experiment* **2018**, 123202 (2018).
- [102] M. Medenjak, V. Popkov, T. Prosen, E. Ragoucy, and M. Vanicat, *SciPost Physics* **6** (2019), 10.21468/scipostphys.6.6.074.
- [103] M. Medenjak, arXiv preprint arXiv:2201.00395 (2022).
- [104] T. Gombor and B. Pozsgay, *SciPost Physics* **12** (2022), 10.21468/scipostphys.12.3.102.
- [105] P. F. Arndt, T. Heinzl, and V. Rittenberg, *Journal of Physics A: Mathematical and General* **31**, L45 (1998).
- [106] R. Arratia, **11** (1983), 10.1214/aop/1176993602.
- [107] E. Helfand, *Physical Review* **119**, 1 (1960).
- [108] Krajnik, Žiga and Ilievski, Enej and Prosen, Tomaž, *Phys. Rev. Lett.* **128**, 090604 (2022).
- [109] B. Derrida and A. Gerschenfeld, *Journal of Statistical Physics* **137**, 978 (2009).
- [110] W. Bryc, *Statistics & Probability Letters* **18**, 253 (1993).
- [111] L. Bertini, A. D. Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, *Reviews of Modern Physics* **87**, 593 (2015).
- [112] B. Derrida, *Journal of Statistical Mechanics: Theory and Experiment* **2007**, P07023 (2007).
- [113] G. B. Cuetara, M. Esposito, G. Schaller, and P. Gaspard, *Physical Review B* **88** (2013), 10.1103/physrevb.88.115134.
- [114] P. Gaspard and E. Gerritsma, *Journal of Theoretical Biology* **247**, 672 (2007).
- [115] J. D. Nardis, B. Doyon, M. Medenjak, and M. Panfil, *Journal of Statistical Mechanics: Theory and Experiment* **2022**, 014002 (2022).
- [116] M. Biskup, C. Borgs, J. T. Chayes, L. J. Kleinwaks, and R. Kotecký, *Physical Review Letters* **84**, 4794 (2000).
- [117] M. Costeniuc, R. S. Ellis, and H. Touchette, *Journal of Mathematical Physics* **46**, 063301 (2005).
- [118] M. Costeniuc, R. S. Ellis, H. Touchette, and B. Turkington, *Physical Review E* **73** (2006), 10.1103/physreve.73.026105.
- [119] N. R. Chaganty, *Sankhyā: The Indian Journal of Statistics, Series A (1961-2002)* **59**, 147 (1997).
- [120] H. J. Haubold, A. M. Mathai, and R. K. Saxena, *Journal of Applied Mathematics* **2011**, 1 (2011).
- [121] F. Mainardi, in *Fractals and Fractional Calculus in Continuum Mechanics* (Springer Vienna, 1997) pp. 291–348.
- [122] R. Gorenflo, A. A. Kilbas, F. Mainardi, and S. Rogosin, *Mittag-Leffler Functions, Related Topics and Applications* (Springer Berlin Heidelberg, 2020).
- [123] F. Mainardi and A. Consiglio, *Mathematics* **8**, 884 (2020).
- [124] H. Touchette, R. J. Harris, and J. Tailleur, *Physical Review E* **81** (2010), 10.1103/physreve.81.030101.
- [125] I. H. Dinwoodie and S. L. Zabell, *The Annals of Probability* **20**, 1147 (1992).
- [126] G. B. Lesovik and L. S. Levitov, *Physical Review Letters* **72**, 538 (1994).
- [127] L. S. Levitov, H. Lee, and G. B. Lesovik, *Journal of Mathematical Physics* **37**, 4845 (1996).
- [128] B. Doyon and H. Spohn, *SciPost Phys.* **3**, 039 (2017).
- [129] D. Bernard and B. Doyon, *Annales Henri Poincaré* **16**, 113 (2014).
- [130] C. N. Yang and T. D. Lee, *Physical Review* **87**, 404 (1952).
- [131] M. Suzuki, *Progress of Theoretical Physics* **38**, 1225 (1967).
- [132] W. Janke and R. Kenna, *Nuclear Physics B - Proceedings Supplements* **106-107**, 905 (2002).
- [133] I. Bena, M. Droz, and A. Lipowski, *International Journal of Modern Physics B* **19**, 4269 (2005).
- [134] R. Blythe and M. Evans, *Brazilian Journal of Physics* **33**, 464 (2003).
- [135] C. Itzykson, R. Pearson, and J. Zuber, *Nuclear Physics B* **220**, 415 (1983).
- [136] C. Pisani and E. R. Smith, *Journal of Statistical Physics* **72**, 51 (1993).
- [137] T. Rakovszky, P. Sala, R. Verresen, M. Knap, and F. Pollmann, *Physical Review B* **101** (2020), 10.1103/physrevb.101.125126.
- [138] L. Zadnik and M. Fagotti, *SciPost Physics Core* **4** (2021), 10.21468/scipostphyscore.4.2.010.
- [139] L. Zadnik, K. Bidzhiev, and M. Fagotti, *SciPost Physics* **10** (2021), 10.21468/scipostphys.10.5.099.
- [140] B. Pozsgay, *Journal of Physics A: Mathematical and Theoretical* **54**, 384001 (2021).
- [141] B. Pozsgay, T. Gombor, A. Hutsalyuk, Y. Jiang, L. Pristýák, and E. Vernier, *Physical Review E* **104** (2021), 10.1103/physreve.104.044106.

- [142] S. Gopalakrishnan, A. Morningstar, R. Vasseur, and V. Khemani, arXiv preprint arXiv:2203.09526 (2022).
- [143] J. Schmidt and A. Schadschneider, arXiv preprint arXiv:2205.06062 (2022).
- [144] C. Flindt and J. P. Garrahan, *Phys. Rev. Lett.* **110**, 050601 (2013).
- [145] X. Peng, H. Zhou, B.-B. Wei, J. Cui, J. Du, and R.-B. Liu, *Physical Review Letters* **114** (2015), 10.1103/physrevlett.114.010601.
- [146] K. Brandner, V. F. Maisi, J. P. Pekola, J. P. Garrahan, and C. Flindt, *Physical Review Letters* **118** (2017), 10.1103/physrevlett.118.180601.
- [147] A. Deger, K. Brandner, and C. Flindt, *Physical Review E* **97** (2018), 10.1103/physreve.97.012115.
- [148] A. Deger and C. Flindt, *Physical Review Research* **1** (2019), 10.1103/physrevresearch.1.023004.
- [149] T. Kist, J. L. Lado, and C. Flindt, *Phys. Rev. Research* **3**, 033206 (2021)