

Emergent quantum mechanics of the event-universe, quantization of events via Denrographic Hologram Theory

Oded Shor ^{2,3}, Felix Benninger ^{1,2,3}, and Andrei Khrennikov ^{4,*}

¹ Department of Neurology, Rabin Medical Center, Petach Tikva, Israel

² Felsenstein Medical Research Center, Beilinson Hospital, Petach Tikva, Israel

³ Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

⁴ Faculty of Technology, Department of Mathematics Linnaeus University, Växjö, Sweden

Abstract. Quantum mechanics (QM) is derived on the basis of a *universe composed solely of events*, for example, outcomes of observables. Such an event universe is represented by a dendrogram (a finite tree) and in the limit of infinitely many events by the p-adic tree. The trees are endowed with an ultrametric expressing *hierarchical relationships between events*. All events are coupled through the tree structure. Such a holistic picture of event-processes was formalized within the *Dendrographic Hologram Theory* (DHT). The present paper is devoted to the *emergence of QM* from DHT. We used the generalization of the QM-emergence scheme developed by Smolin. Following this scheme, we did not quantize events but rather the differences between them and through analytic derivation arrived at *Bohmian mechanics*. Previously, we were able to embed the basic elements of general relativity (GR) into DHT, and now after Smolin-like quantization of DHT, we can take a step toward quantization of GR. Finally, we remark that DHT is *nonlocal in the treelike geometry*, but this nonlocality refers to relational nonlocality in the space of events and not Einstein's spatial nonlocality.

I. INTRODUCTION

One of the main problems in the unification of quantum and classical mechanics and general relativity (GR) is the *quantization scheme* that is applied to classical observables (see section B for the extended discussion).

The important lesson of quantum foundational studies is that quantum mechanics (QM) is about events, namely, the outcomes of measurements (*phenomena* in Bohr's terminology, [1,2]). Therefore, QM can be treated as a special formalism for event representation of physical processes (see, for example, several studies [3–10]). This is the basic idea of *the relational*

interpretation of QM (due to Rovelli). Thus, one can try merging quantum theory and general relativity as two event-based theories (5–7,9).

From our viewpoint, one of the obstacles for event reconstruction of QM is the common use of spatial representation and Cartesian coordinates. Even Smolin who tried to consistently develop the event QM started with the spatial coordinates (6).

In recently developed *Dendrographic Hologram Theory* (DHT) as described in several studies (11–13), we proceeded with the reconstruction without using the spatial or even the temporal picture, at least as the theory's starting point (later the spatial and temporal coordinate description can emerge from the purely event-based theory [13]). The 5 cornerstones DHT relies on are as follows:

1. *Leibniz principle* (identity of indiscernibles) which states that If, for every property F, object x has F if and only if object y has F, then x is identical to y.
2. *Relational and event interpretations of physical theories.*
3. *Bohr's contextuality.* According to Bohr the outcomes of measurements are not the objective properties of systems. They quantitatively represent interrelation between a system S and an observer O (the measurement device of O).
4. *Ontic-epistemic structuring of scientific theories.* Ontic description is observer independent description of reality - as it is. Epistemic description is based on knowledge which O can extract within experiments. The ontic theory is not verifiable experimentally. It is unapproachable by the observer; the observer constructs its approximate epistemic representation of the ontic description by collecting data.
5. *P-adic (ultrametric) theoretical physics.* (See, e.g. [14-23]).

In DHT, events (Bohr's phenomena) are portrayed as branches of a dendrogram (finite tree). For an illustration that describes how to construct the dendrographic tree from data please see figure 1. These finite trees, which are the observer's epistemic description of reality, are constructed as follows:

First an observer collects data by performing measurements. Then he applies a *heirarchical clustering algorithm*. By choosing a distance matrix and a certain clustering (linkage function) algorithm, one constructs an agglomerative hierarchical cluster tree. In this tree each event, which is pictured by the unique branch, can be represented by a binary string corresponding to a finite sequence of questions with the answers yes/no (or 1/0). The set of branches of the tree (or

the strings of ones and zeros that fully describe them) is the dendrogram. Each branch runs from tree's root to a leaf – the end point of the branch. The leaves encode the events generated by measurements. For two such event-branches, the longer the initial common part corresponds to a closer relation between these two events. This common path closeness relation determines the *2-adic metric* on a dendrogram. This is very special metric, so called *ultrametric*.¹

The strings of ones and zeros corresponding to the branches of a dendrogram can be identified with the 2-adic (binary) expansions of natural numbers. So, at the epistemic level events can be mathematically encoded by natural numbers. The crucial point is that the distance between these natural numbers is determined by the tree structure, this is 2-adic ultrametric.


Within the limits of an infinite number of events, the ontic description of the event-universe is portrayed as an infinite tree. The simplest class of such trees are p-adic trees, which are homogeneous trees with $p > 1$ edges branched from any vertex. Such trees can be endowed with the algebraic structure and the topology consistent with this structure; the p-adic topology is given by the p-adic ultrametric, namely, the strong triangle inequality holds (14).² P-adic distance between two branches of the tree is determined by their common initial path: a longer common path represents a shorter distance. Infinite *branches represent events*, so the space of events, a finite or infinite tree, is endowed with a p-adic ultrametric. Hence, DHT does not deal with space–time localization of events but with p-adic distance encoding hierarchic relations between events. This common initial path distance determines *the degree of similarity between events*. This field is endowed with strange geometry where, for instance, all triangles are isosceles. This is a consequence of the strong triangle inequality. Moreover, upon defining “open” and “closed” balls as $B(R, a) = \{ x: r_p(a, x) \leq R \}$ and $B-(R, a) = \{ x: r_p(a, x) < R \}$, both balls are at the same time closed and open sets of the metric space, as such each point in a ball can be selected as its center. Geometrically a ball is a batch of infinite branches having the finite common initial segment. Such spaces are disordered, totally disconnected, and having zero

¹ Of course, this distance between events depends on dendrogram's construction based on the distance on the space of measurement's outcomes which was the initially chosen for clustering algorithm. For example, we can start with the Euclidean distance on the data-set. Then it is transferred into the 2-adic distance. One can use more complex clustering algorithms generating trees with p-edges leaving each vertex, where $p > 1$ is a natural number. These are p-adic trees and they are endowed with p-adic ultrametric: closeness based on the common initial part of two branches.

² Thus, the Dendrographic Hologram Theory (DHT) is part of well-established branch of theoretical physics, p-adic (non-Archimedean) physics [14–22], which was widely explored in string theory, cosmology, GR, theory of complex disordered systems (including spin glasses which were studied with p-adic methods by Parisi et al. [23] and others).

topological dimension. However, such geometries arise very naturally from data series with application of clustering algorithms. For an illustration that describes how to construct dendrographic tree from data please see figure 1.

Fig 1. Illustration that describes the construction of a dendrographic tree and its 2-adic representation from a single discrete geodesic with 8 events. A. An observer "O" measures discrete dynamics along a spacetime geodesic .B choosing a distance metric he calculates pairwise distances between the 8 discrete events along a geodesic and constructs a distance matrix . C by applying a hierarchical clustering algorithm on the distance matrix he constructs an Agglomerative hierarchical cluster tree. D in this hierarchical tree each event, which have a unique branch, can be represented by a binary string or p-adic expansion. E A numerical example is given for calculating the sum of 2-adic expansions of each event 2-adic representation. More over 2-adic differences between two events, which are marked as q_{ik} , are shown as a numerical example.



Based on extensive numerical simulations DHT has been successful in simplifying nontrivial concepts, for instance, the identification of the Bohmian explicate-implicate order with the dendrographic epistemic-ontic description of universe (11). Moreover this connection of the explicate-implicate order with its epistemic-ontic showed that each explicate level contains the information of the lower explicate order and vice versa, Thus suggesting a holographic principle contained in each explicate order on the implicate order (11). The illusive Bohmian quantum potential reduces in DHT to a simple structural property of the dendrographic relational structure. In article (12) it was shown (with extensive numerical simulations) that by using dendrogram representation of classical data it is possible to violate the CHSH inequality. We found that the seed of the CHSH-violation is violation of ergodicity -- the hidden hierarchic relations are non-ergodic. In article (12) the degree of “quantumness” was determined by the degree of the violation of the Bell type inequalities. Moreover, the degree of classicality is based on system’s complexity. The latter is the size and topological complexity of its dendrogram representation. where quantum systems are characterized by a low complexity of their dendrogram topology. Another study (13) (again with extensive numerical simulations) represented GR dynamics along geodesics in the relational p-adic framework. Here the dynamics on dendrograms are emergent from the simple action principle. These studies also showed that p-adic coordinates emergent from the action principle correspond the real spacetime coordinates.

DHT is not classical nor quantum theory in the conventional sense. Connection with classical theory (in the GR framework) was established in article (13) and will be discussed in more detail in section III. In the present article, we want to establish connection between DHT and QM and to emerge QM from DHT (see references (5,6,24–29) for various suggestions for emergent QM).

Based on a scheme developed by Smolin (5), we demonstrate that QM can be emerged from the event model given by DHT. Using this scheme for each event i , a set of *views* V_i^k from i to other events $k=1, \dots, N$ is defined, where N is the total number of events. In this situation, the views rather than the spatial coordinates are the basic quantities. The view V_i^k encodes the difference between the events, i and k . This kind of quantization is not quantization of positions and momenta but of the differences between events. In particular, the event-momentum is expressed via views. However, in Smolin’s approach, the Cartesian coordinates still play the crucial role. In DHT, we start directly with the event representation given by a dendrogram. The

views are determined by the hierarchic relational structure of the event-tree and the differences between events are also with respect to this structure. So, our approach is more consistent as it solely explores the event picture without a direct connection to the space-time aspect. On the other hand, although we used different mathematical apparatus, we perfectly matched Smolin's ideology of event-physics.

We will generalize Smolin's quantization scheme to match DHT. This process allows us to use any form of distribution on the space of views and many differences/relational functions acting on the differences of events. In article (5), a very special distribution was in the use and its properties played crucial role in generating the quantum potential in the Bohmian form.

Emergence of QM in the Bohmian mechanics form immediately rises the issue of *nonlocality* and its meaning in our theory. DHT is fundamentally non-local since all events are coupled via the hierarchical relational tree-like structure. This coupling is especially evident in DHT-dynamics (30) in which appearance of a new event generates reconstruction of the whole dendrographic universe via a recombination of the tree branches. However, this non-locality is not the same as Einstein's space-time non-locality rather it is *relational nonlocality*. As is shown in article (13), starting with the dendrogram, one can reconstruct (but not uniquely) space-time representation. In that study, the relational non-locality would be expressed in the form of apparent spatial non-locality.

II. CANONICAL QUANTIZATION SCHEME

This section is aimed at reminding the reader that although QM functions well in numerous applications, it has some foundational problems. The reader who is not surprised by this statement can omit this section.

The quantum mechanical scheme was and still is very productive, and its consistent application led to successful development of quantum theory. One of its main advantages is the coupling to classical phase space mechanics and theory of Hamiltonian equations. This coupling in the form of deformation quantization leads to the *correspondence principle*, which is so important for heuristic justification of the quantization scheme. Deformation quantization is a rigorous mathematical procedure.

The quantum algebra of pseudo-differential operators and functions of operators of position and momentum within the limits of $\hbar \rightarrow 0$ coincide with the classical symbol algebra of these operators, namely, the algebra of functions on the phase space.

At the same time, it is well known that the quantization procedure has some problems that seem to be unsolvable, at least within the internal framework of QM. One such problem is the impossibility of constructing a relativistic theory for QM for which a relativistic theory was built only for quantum field theory (QFT). However, QFT suffers from infinities. Although this problem can be smoothed via numerous renormalization procedures, and these procedures are not so natural based on the heuristic viewpoint.³ We also note that the basic notions of quantum information theory, such as entanglement, are naturally formulated within the QM framework but not within the QFT framework [32].

Now, we point to difficulties in quantization of GR. It seems that GR is not quantizable, at least in a straightforward manner. In fact, the failure in unification of QM (or QFT) with GR is really the big black cloud on the sunny quantum sky.

We also recall the almost forgotten paper of Zeilinger in which he stressed that QM was not derived from some physically natural fundamental principles, such as the principles of special relativity, namely, the constancy of the light velocity or the relativity principle (33). Zeilinger considered the absence of the fundamental principles as one of the main difficulties in foundational justification of QM and QFT (33).

Finally, we turn to deformation quantization and point out that this excellent mathematical theory has a big interpretational problem, namely, the treatment of the Planck constant \hbar as a variable parameter.

The aforementioned problems led to a variety of attempts to emerge quantum theory from heuristically more acceptable theories (24–29).

III. QUANTIZATION OF EVENTS OR VIEWS

We claim that in the event-universe in which each event is unique, the process of quantization of concrete events (and not differences between them, namely, the views) would lead to inconsistency in the GR. The reason is that from a unique-events universe, the only

³ Nowadays the problem of quantum field theory (QFT) infinities is practically ignored, but Dirac was so disturbed by it during his lifetime that he permanently demanded that the theory based on the procedure of canonical quantization is inconsistent and it should be replaced with a totally new theory [31] (see [32] for a discussion of this issue).

available non-coarse-grained distribution is the trivial distribution. A N event universe will generate a probability distribution for which each event probability to occur is 1/N. In article [5], this distribution appeared in equation 22, but totally different and very special distribution then appeared in the formula 24 which imposes constraints on the possible configuration space. The latter leads to the right expression for the quantum potential, while QM cannot emerge with uniform distribution

It is evident that in a unique-events theory such as GR, the Leibnitz principle (11), *the identity of indiscernible*, is maintained. Practically, *this principle argues that if there are two or more identical events, they are in fact the same event*. This concept in our understanding is the main problem in uniting QM and GR. QM allows many events that are the same to construct a probabilistic theory which will have other distributions than the trivial, 1/N, one as mentioned above. Therefore, we should allow many identical events to concurrently have the same view (or “they agree on the view”) of the rest of the universe; in return, this process contradicts the relativity element in GR. A way out of this situation exists in which GR allows identical differences between events. Thus, in GR we can construct the distributions of differences. In return, this step might lead us to an emergent QM compatible with GR.

In article [13], GR was embedded into DHT. In this article, GR’s geometry was represented by its geodesics. Temporal discretization of geodesics generated the data set, which was transformed into a dendrogram. Its branches encode events, which are the points of discretized geodesics. The DHT-dendrogram expressed all relationships between events, including the space–time structure corresponding to the concrete GR-metric. By quantizing DHT resulting in the emergence of QM (via the scheme described in article [5]), we actually quantize GR event data via its DHT representation. Of course, these schemes are just schematic manipulations, which are far from the rigid basis of quantum gravity.

IV. EMERGENCE OF QUANTUM REPRESENTATION FROM THE DH THEORY

We follow the line developed by Smolin (5) with slight modifications and conversion to our DH-theory point of view.

We start with the *view* of the *i*’th event, which is according to Smolin:

$$V_i^k = \frac{x_i - x_k}{|x_i - x_k|^2} (R - |x_i - x_k|) \text{ for } k \neq i$$

and x_k is any of the rest $N - 1$ coordinates of phase space

With $(R - |x_i - x_k|)$ as a cut-off for how far the view can be influenced from the other x_k

We did not apply this cut-off. For our purposes, we used the equation shown below:

$$edge_i = \sum_{j=0}^k a_j \times 2^j$$

is a dendrogram branch 2 adic expansion and its monna map conversion is defined as $event_i =$

$$\sum_{j=0}^k a_j \times 2^{-j-1}, a_j = 1,0,$$

$$q_{ik} = (event_i - event_k) \text{ thus :}$$

$$V_i^k = 1/q_{ik} \text{ in which } k \text{ is any of the remaining } N - 1 \text{ edges } k \neq i \quad (1)$$

We then define the measure of differences called distinctiveness, same as in (5):

$$I_{ij} = \frac{1}{N} \sum_{ik} (V_i^k - V_j^k)^2 \quad (2)$$

Thus, the smaller I_{ij} , the more easily $event_i$ and $event_j$ can be differentiated by their views. We

then defined the variety as shown below, same as in (5):

$$v = \frac{A}{N^2} \sum_{i \neq j} I_{ij} \quad (3)$$

Or as inter ensemble potential energy.

We start with constructions of the “differences pdf”, \tilde{p} . This will be our fundamental distribution. Thus, given a dendrogram that describes, p-adically, the relations between m events we calculate all possible pairwise differences of the p-adic edges representation and represent them as events through monna map as above:

$$q_{ik} = (event_i - event_k) \quad (4)$$

for $p = 2$,

$$edge_i = \sum_{j=0}^k a_j \times 2^j \rightarrow event_i = \sum_{j=0}^k a_j \times 2^{-j-1}, a_j = 1,0, \quad (5)$$

the Monna map maps natural numbers into rational numbers belonging the segment $[0,1]$. A dendrogram is mapped into a subset of $[0,1]$. This map can be extended to the infinite p-adic tree where its branches are represented by infinite series; for $p=2$,

$$edge_i = \sum_{j=0}^{\infty} a_j \times 2^j \rightarrow event_i = \sum_{j=0}^{\infty} a_j \times 2^{-j-1}, a_j = 1,0, \quad (6)$$

The latter is important for considering the limit of the universe with infinitely many events; they are portrayed on an infinite 2-adic tree.

Thus, from all q_{ik} we have a discrete “difference pdf”, \tilde{p} , which is the fraction of each unique value of q_{ik} defined For the set of the unique q_{ik} values Q we have:

$$\tilde{p}_j = \frac{(\text{number of different } q_{ik} \text{ that equal } Q_j)}{\text{total number of } q_{ik}}$$

A. The event differences distribution ρ

By using the above \tilde{p} distribution, we can construct the ρ distribution, which is the distribution of events in the given dendrogram. So, the x axis will be the edges with values of the Monna map and the $y = \rho(x)$ axis will be calculated as follows:

$edge_i$ has $m-1$ values of q_{ik} in which $k \neq i$, thus:

$$\rho(edge_i) = \left(\prod_{k \neq i} \tilde{p}(q_{ik}) \right) / \sum_{j=1}^n \left(\prod_{k \neq j} \tilde{p}(q_{jk}) \right)$$

in which $n = \text{number of edges in dendrogram}$ (7)

Of course, this $\rho(edge)$ is only one possible pdf from a vast space of possible pdf's made possible by the \tilde{p} distribution (it will be interesting to understand what characterizes this ρ distribution from the rest of all possible distributions).

Please note that based on this equation, we can calculate any possible edge (even if it is not in the dendrogram) as long as all of its q_{ik} $k \neq i$ are present in the \tilde{p} distribution.

B. The differences energy

To define the differences energy, which will be equivalent to the usual kinetic energy, we define the equation:

$$p_j = \frac{1}{N} \sum_{k \neq j} edge_j - edge_k \text{ in which } N \text{ is number of edges minus } 1. \quad (8)$$

This formula represents the mean difference between event j and the remaining events. Thus, p_j takes the part as the momentum in Bohmian mechanics.

Now, we introduce the phase S as the new parameter of the model:

$$\text{We set } p_j = \partial S_j \text{ and } \omega_j = e^{iS_j}. \quad (9)$$

We have our equivalent of momentum of an $event_j$; hence, the viewed differences of $event_j$, is the sum of all views of the $event_j$ multiplied by the differences of S_j to the rest S_k .

$$p_{\text{viewed differences of event } j} = -i \left(\frac{1}{N} \right) \sum_{j \neq k} V_k^j \ln \left(\frac{\omega_j}{\omega_k} \right) = -i \left(\frac{1}{N} \right) \sum_{k \neq j} V_k^j (S_j - S_k) \quad (10)$$

We can then calculate the “differences energy” as the real part of:

$$T_{\text{differences energy}} = \frac{1}{N^2} \sum_{k \neq j} \left(\frac{1}{q_{jk}} \right)^2 \left(\ln \ln \frac{\omega_j}{\omega_k} \right)^2 \quad (11)$$

Where $T_{differences\ energy}$ serves as our equivalent of kinetical energy

Thus, our fundamental action would be the same as Smolins

$$S(w, q_{jk}) = Z(\sum_{k \neq j} dge_k(1/q_{jk}) \frac{d}{d(dendrogram)} (\ln \ln \frac{\omega_j}{\omega_k})^2) - H(w, q_{jk}) \quad (12)$$

In which

$$H(w, q_{jk}) = Z_k \sum_{k \neq j} (1/q_{jk})^2 Re(\ln \ln \frac{\omega_j}{\omega_k})^2 - Z_V \sum_k \sum_{i \neq j} (\frac{1}{q_{jk}} - \frac{1}{q_{ik}})^2 + \sum_k U \quad (13)$$

and Z , Z_V and Z_Q are some normalization factors.

C. The origin of the quantum potential

$$\text{We started with } View_{ik} = (edge_i - edge_k)/|edge_i - edge_k|^2 \quad (14)$$

We identify $edge_i$ as z in the continuous limit, and we identify $edge_k$ as $z + x$ in the continuous limit

Thus,

$$\Phi(z, z + x) = View_{z, z+x} \int \Phi(z, z + x) \rho(z, z + x) dx = \text{all views of } i \text{ (} z \text{ is fixed)} \quad (15)$$

$$\text{Similarly, with } View_{jk} = (edge_j - edge_k)/|edge_j - edge_k|^2 \quad (16)$$

We identify $edge_j$ as z in the continuous limit, and we identify $edge_k$ as $z + y$ in the continuous limit

$$\text{Thus, } \Phi(z, z + y) = View_{z, z+y} \int \Phi(z, z + y) \rho(z, z + y) dx = \text{all views of } j \text{ (} z \text{ is fixed)} \quad (17)$$

(z is fixed).

$$\text{Now, the variety is defined as } \sum_i \sum_{j \neq i} \sum_k (View_{ik} - View_{jk})^2 \quad (18)$$

$$\text{Or in the continuous form } \int dz \int dy \int dx (\Phi(z, z + y) - \Phi(z, z + x)) \quad (19)$$

$$\text{However, } \Phi(z, z + y) = View_{z, z+y} = (z - (z + y))/|z - (z + y)|^2 = -y/(y)^2 \quad (20)$$

$$\text{and } \Phi(z, z + x) = View_{z, z+x} = (z - (z + x))/|z - (z + x)|^2 = -x/(x)^2 \quad (21)$$

Thus, we have

$$\begin{aligned} & \int z \rho(z) Z_V \int dy \int dx ((\Phi(z, z + y) - \Phi(z, z + x)) \rho(z, z + y) \rho(z, z + x)) \\ & = \int dz \rho(z) Z_V \int_a^R dx \int_a^R dy (x/(x)^2 - y/(y)^2)^2 \rho(z, z + y) \rho(z, z + x) \end{aligned}$$

$$\text{where } a = \frac{1}{\rho_2 \text{maximal } 2\text{-adi ball of dendrogram}} \text{ and } R = 2^{\text{minimal } 2\text{-adi ball of all } q_{jk}} \quad (22)$$

which is equivalent to $Z_V \sum_{j \neq k} \sum_k \sum_{i \neq j} (\frac{1}{q_{jk}} - \frac{1}{q_{ik}})^2$ where Z_V is a normalization factor.

We show now the emergence of the quantum potential term in our action

where $\rho(z+x)$ and $\rho(z+y)$ are expanded in x and y around z . we take the two integral terms with $(\rho(x))^2$ and $(\partial\rho(x))^2$

$$Z_V(\rho(z))^2 \int_a^R dx \int_a^R dy (\frac{1}{x} - \frac{1}{y})^2 \left[= \frac{1}{R^2} \right] + \\ + Z_V(\partial\rho(z))^2 \int_a^R dx \int_a^R dy (\frac{1}{x} - \frac{1}{y})^2 xy \quad [= (\partial\rho(z))^2 / (\rho(z))^2] \quad (23)$$

$$\text{Thus } \int_a^R dx \int_a^R dy (\frac{1}{x} - \frac{1}{y})^2 \left[= \frac{1}{R^2 Z_V(\rho(z))^2} \right] \text{ and } \int_a^R dx \int_a^R dy (\frac{1}{x} - \frac{1}{y})^2 xy \quad [= \frac{1}{Z_V(\rho(z))^2}] \quad (24)$$

As the arguments and bounds of the definite integrals are the same we calculate the indefinite integrals then substitute $x=y=(R-a)$

Thus, we obtain two equations:

$$y - 2 - 2\ln^2(y) = \frac{1}{R^2 Z_V(\rho(x))^2} \text{ in which } y = R - a \\ y + y^2 \ln(y) - y^2 = \frac{1}{R^2 Z_V(\rho(x))^2} \text{ in which } y = R - a \\ (25)$$

Thus

$$R - a - 2 - 2\ln^2\left(\frac{R}{a}\right) = \frac{1}{R^2 Z_V(\rho(z))^2} \\ R - a + (R - a)^2 \ln\left(\frac{R}{a}\right) - 2(R - a)^2 = \frac{1}{Z_V(\rho(z))^2} \quad (26)$$

We previously defined a and R in which $a = \frac{1}{\rho^{2\text{maximal } 2-a} \text{ ball of dendrogram}}$ and

$R = 2^{\text{minimal } 2-a} \text{ ball of all } q_{jk}$. We also defined $1/N$ as $1/2^{\text{maximal } 2-a} \text{ ball of dendrogram}$,

which tends to move toward zero as we increase event numbers and thus the dendrogram size.

using our ρ distribution that we constructed as described above

$$\text{We are left with equation 27:} \quad R - a - 2 = \frac{1}{R^2 Z_V(\rho(z))^2} \quad (27)$$

and equation 28:

$$R - a - 2R^2 + 2a^2 = \frac{1}{Z_V(\rho(z))^2} \quad (28)$$

By selecting Z_V in the appropriate way, we obtained in both equations terms that result (equation 27 the term -2 and in equation 28 the term $-2R^2$) in $-2R^2Z_V = \frac{1}{(\rho(z))^2}$ while all other terms tended to move toward zero by $1/N$.

Thus, we have two arguments in the integral of equation 22: $\frac{1}{R^2}$ and $(\partial\rho(z))^2/(\rho(z))^2$

This mathematical trick can be implemented on other arguments then $(\frac{1}{x} - \frac{1}{y})^2$. For instance, for: $(x - y)^2$ and even $(xy)^2$

We will show that we can generate a quantum potential term for $(x - y)^2$

$$\int dz \rho(z) Z_V \int_a^R dx \int_a^R dy (x - y)^2 \rho(z + x) \rho(z + y) \quad (29)$$

Using the same steps, we obtain two equations

$$y - \left(\frac{1}{6}\right) y^4 = \frac{1}{R^2 Z_V (\rho(x))^2}$$

$$y - \left(\frac{5}{6}\right) y^6 = \frac{1}{R^2 Z_V (\rho(x))^2} \quad (30)$$

And

$$R - a - \left(\frac{1}{6}\right) (R - a)^4 = \frac{1}{R^2 Z_V (\rho(x))^2}$$

$$R - a - \left(\frac{5}{6}\right) (R - a)^6 = \frac{1}{R^2 Z_V (\rho(x))^2} \quad (31)$$

However, in the first equation we obtain the term R^4 and in the second R^6 and by selecting the right Z_V we get for those two terms $R^6 Z_V = \frac{1}{(\rho(z))^2}$; thus we obtain the correct terms for the quantum potential and a constant in the principle action.

D. The continuous differences energy

From the term:

$\sum_{k \neq j} (1/q_{jk})^2 Re(\ln \frac{\omega_j}{\omega_k})^2$ we reconstructed the probabilistic equivalent as shown below:

we construct again the probabilistic equivalent as:

$$RE \int dz \rho(z) Z_d \int_a^R dx \rho(x) \frac{1}{z - x} \left(\ln \frac{\omega(x)}{\omega(z)}\right)^2 =$$

$$= \int dz \rho(z) \int_a^R dx \rho(x) Z_d \frac{1}{z - x} (S(x) - S(z))^2 \quad (32)$$

And by expanding around z in x we arrive at

$$differences\ energy = \int dz \rho(z) \frac{(\partial S)^2}{2} \quad (33)$$

E. The continuous dendrogram differences energy

Constructing the probabilistic equivalent of

$$Z \sum_{k \neq j} edge_k (1/q_{jk}) \frac{d}{d(dendrogram)} (\ln \frac{\omega_j}{\omega_k})^2 \quad (34)$$

$$\text{we obtain } \frac{dm}{d(dendrogram)} = \sum_{j \neq k} -i(1/q_{jk}) \left(\frac{ds_j}{d(dendrogram)} - \frac{ds_k}{d(dendrogram)} \right) \quad (35)$$

where upon transforming it to the continuum we obtained:

$$N \int d(dendrogram) \int dz Z_0 \rho(z) z \frac{dp(z)}{d(dendrogram)}$$

$$\text{Where } \frac{dm(z)}{d(dendrogram)} = (C/\rho(z)) \partial \rho(z) \frac{ds(z)}{d(dendrogram)} \quad (36)$$

The full continuous action is thus:

$$S = \int dendrogram \int dz \rho(z) \left[\frac{dS(z)}{d(dendrogram)} + (\partial S)^2/2 - \left(\frac{\partial \rho(z)}{\rho(z)} \right)^2 + U \right] \quad (37)$$

Where $U = \rho(z)$.

The Hamilton Jacobi equation of the action (notice we are on the straight line [0 1] now in all our pdfs).

$$-\dot{S} = (\partial_{edge} S)^2 + U + U^Q \quad (38)$$

In which $U^Q = (\Delta^2 \sqrt{\rho})/\sqrt{\rho}$ the quantum potential

And $U = \rho$, the potential

And $\dot{S} = S(\text{present dendrogram}) - S(\text{previous dendrogram})$

All parameters are well-defined.

The probability conservation law is

$$\dot{\rho} = \partial_{edge} (\rho \partial_{edge} S)^2$$

In which $\dot{\rho} = \rho(\text{present dendrogram}) - \rho(\text{previous dendrogram})$ (39)

Equations 38 and 39 are the real and imaginary parts of the Schrodinger equation for $\psi = \sqrt{\rho} e^{iS}$

please notice the arguments in the action must follow least action principle thus they are all following best match (or best distance) distance between two zero dimensional dendrogram structures. This is with agreement with particle shape dynamics theory ideology. In our case the particles are the events encoded by dendrograms p-adically. ρ is the probabilistic representation of the dendrographic structure where events correspond to particles in shape dynamics. S is again a property of the events encoded in the dendrogram. Thus, we closely linked casual set theory with particle shape dynamics under our formulation with an emergent QM. Notice that shape dynamics do not yet have a quantum theory.

We now turn to the link with the gravitational field as stated above

We use as our probabilistic representation (or our quantum representation) the Hamiltonian:

$$H = \rho(z)[(\partial S)^2/2 - (\frac{\partial \rho(z)}{\rho(z)})^2 + U]$$

With the canonical variables ρ and S and where $U = \rho$. Please note that ρ represents the variable *edge*, and S the variable $\int p$ where p is defined as $\frac{1}{N} \sum_{k \neq j} edge_j - edge_k$.

but in our construction of the dendrogram we have used a clustering function, F , so

$$\rho(edge) = F\left(\frac{d^2 x^\sigma}{ds^2} + \Gamma_{\mu\nu}^\sigma \frac{dx^\mu}{ds} \frac{dx^\nu}{ds}\right)$$

where F is a function which clusters spacetime coordinates, x , by using some distance metric and clustering algorithm, and then constructs the agglomerative hierarchical cluster tree from which the dendrogram structure is defined by some algorithm and then gives each single cluster a p-adic representation. Again, x are spacetime coordinates. The procedure applies also to

$$S(edge) = \int \int \int F\left(\frac{d^2 x^\sigma}{ds^2} + \Gamma_{\mu\nu}^\sigma \frac{dx^\mu}{ds} \frac{dx^\nu}{ds}\right) - F\left(\frac{d^2 y^\sigma}{ds^2} + \Gamma_{\mu\nu}^\sigma \frac{dy^\mu}{ds} \frac{dy^\nu}{ds}\right) dx dy d(edge)$$

and where $y=x+z$ and x and y are spacetime coordinates

In which $\Gamma_{\mu\nu}^\sigma$ is the gravitational force.

V. DISCUSSION

DHT [11–13,30] presents the event picture of the universe. This picture does not contain the physical space, rather, it has the treelike geometry expressing hierarchical relationships between events. The relational structure generates a type of nonlocality effect. This effect is especially evident in the dynamical model for DHT [30] in which the appearance of a new event induces recombination of all events on the tree and interrelation between them.

The DHT model is not classical or quantum in the sense of conventional physics. The discrete structure of branching on the event-tree suggests quantization. In article [12], we found some quantum-like effects within DHT, for example, the possibility to violate the Bell type inequalities, which were treated as the tests of the level of quantum properties.

However, coupling of DHT with standard QM was missing up to that point. This coupling was established in the present paper based on Smolin's scheme for emergence of QM from the view-universe. Although ideologically Smolin advertised the purely event pictured universe, in concrete calculations he needed the standard mathematical apparatus with real Cartesian coordinates, at least in article (5). We start without space-time, and just used a hierarchical tree of events. However, found that the Smolin quantization scheme can be generalized for state space of a dendrogram within the limitations of the p-adic type.

One of the aims of quantization of DHT is to take a step toward quantization of GR, which was coupled to DHT in article [13]. As explained in sections II and III, the individual structure of events in GR cannot be expressed in the probabilistic framework of QM (see figure 2) . Therefore, to emerge QM from DHT, we should try not to operate with events distributions. Following Smolin, we operated with distributions of differences between events expressed in terms of views from one event to another.

We can say that the mission of DHT quantization was successfully completed. Of course, this completion is just the first step. In future work, we should represent the basic DHT quantities in standard quantum terms. One of interesting problem is construction of the quantum representation of correlations between views.

Further studies should address the formulation of other properties of QM in our approach such the emergence of spin and the characterization of different field types along with their corresponding QM properties. Moreover, we speculate that the connection to shape dynamics, in terms of dynamics of relations structures would increase our predictive powers of the dynamical processes in the event universe




Fig 2. An abstract illustration that suggests DHT can describe quantum phenomena, general relativity phenomena and quantize gravitation field. Direct quantization from GR to QM is still missing.

References

1. N. Bohr, (1987). *The Philosophical Writings of Niels Bohr*, 3 vols. (Ox Bow Press, Woodbridge, CT).
2. A. Plotnitsky, (2012). *Niels Bohr and Complementarity: An Introduction* (Springer, Berlin and New York).
3. Rovelli, C. (1996), "Relational quantum mechanics", *International Journal of Theoretical Physics*, 35: 1637–1678.
4. Laudisa, Federico, and Carlo Rovelli. "Relational quantum mechanics." (2002).
5. L. Smolin, Quantum mechanics and the principle of maximal variety. *Found Physics*, 46, 736-758 (2016).
6. Smolin, L. (2018). The dynamics of difference. *Foundations of Physics*, 48(2), 121-134.
7. Martin-Dussaud, P., Rovelli, C., & Zalamea, F. (2019). The notion of locality in relational quantum mechanics. *Foundations of Physics*, 49(2), 96-106.
8. Biagio, A. D., & Rovelli, C. (2022). Relational Quantum Mechanics is about Facts, not States: A reply to Pienaar and Brukner. *Foundations of Physics*, 52(3), 1-21.
9. L. Smolin, *Three roads to quantum gravity*. Hachette UK (2008).
10. G. Brida Experimental test of an event-based corpuscular model modification as an alternative to quantum mechanics. *J Phys Soc Japan*, 82, 034004 (2013).
11. Shor, O.; Benninger, F.; Khrennikov, A. Representation of the Universe as a Dendrographic Hologram Endowed with Relational Interpretation. *Entropy* 2021, 23, 584. <https://doi.org/10.3390/e23050584>.

12. Shor, O.; Benninger, F.; Khrennikov, A. Dendrographic Representation of Data: CHSH Violation Vs. Nonergodicity. *Entropy* **2021**, *23*, 971. <https://doi.org/10.3390/e23080971>.
13. Shor, O.; Benninger, F.; Khrennikov, A. Towards Unification of General Relativity and Quantum Theory: Dendrogram Representation of the Event-Universe. *Entropy* **2022**, *24*, 181. <https://doi.org/10.3390/e2402018>.
14. Khrennikov, A. *p-Adic Valued Distributions in Mathematical Physics*; Springer: Berlin/Heidelberg, Germany, 1994.
15. Vladimirov, V.S.; Volovich, I.V.; Zelenov, E.I. *p-Adic Analysis and Mathematical Physics*; World Scientific: Singapore, 1994.
16. Parisi, G.. On p-adic functional integrals. *Modern Physics Letters A* **1988**, *3*(06), 639-643.
17. García-Compeán, H., Edgar Y. López, and W. A. Zúñiga-Galindo. p-Adic open string amplitudes with Chan-Paton factors coupled to a constant B-field. *Nuclear Physics B* **951** (2020): 114904.
18. Dragovich, B. "A p-Adic Matter in a Closed Universe." *Symmetry* **14.1** (2022): 73.
19. Djordjević, G. S., Dragovich, B., Nešić, L. D., & Volovich, I. V. (2002). p-Adic and adelic minisuperspace quantum cosmology. *International Journal of Modern Physics A*, *17*(10), 1413-1433.
20. Chen, L.; Liu, X.; Hung, L. Emergent Einstein Equation in P-Adic Conformal Field Theory Tensor Networks. *Phys. Rev. Lett.* **2021**, *127*, 221602. [[Google Scholar](#)] [[CrossRef](#)]
21. Hung, L.-Y.; Li, W.; Melby-Thompson, C.M. P-Adic CFT is a Holographic Tensor Network. *J. High Energy Phys.* **2019**, *2019*, 170. [[Google Scholar](#)] [[CrossRef](#)]
22. Gubser, S.S.; Heydeman, M.; Jepsen, C.; Marcolli, M.; Parikh, S.; Saberi, I.; Stoica, B.; Trundy, B. Edge Length Dynamics on Graphs with Applications to P-Adic AdS/CFT. *J. High Energy Phys.* **2017**, *2017*, 1–35. [[Google Scholar](#)] [[CrossRef](#)]
23. Parisi, G.; Sourlas, N. P-adic numbers and replica symmetry breaking. *Eur. Phys. J. B Condens. Matter Complex. Syst.* **2000**, *14*, 535–542.
24. t' Hooft, G. Emergent Quantum Mechanics and Emergent Symmetries. AIP Conference Proceedings **957**, 154 (2007);
25. Elze, H.T.: Quantum models as classical cellular automata, arXiv:1701.02252
26. Elze, H.T. (2008). Note on the existence theorem in ‘emergent quantum mechanics and emergent symmetries’. *Journal of Physics A: Mathematical and Theoretical*, *41*(30), 304020.

27. 't Hooft, G.: The cellular automaton interpretation of quantum mechanics. In: *Fundamental Theories in Physics*, vol. 185, Springer, Berlin (2016).
28. Khrennikov, A. *Beyond quantum*. Jenny Stanford Publ., Singapore (2014).
29. Khrennikov, A. Quantum epistemology from subquantum ontology: Quantum mechanics from theory of classical random fields. *Ann. Phys.* **2017**, *377*, 147–163. [[Google Scholar](#)] [[CrossRef](#)]
30. Shor, O., Benninger, F., & Khrennikov, A. (2022). Dendrographic Hologram Theory: Predictability of Relational Dynamics of the Event Universe and the Emergence of Time Arrow. *Symmetry*, *14*(6), 1089.
31. Dirac, P.A.M. *Lectures on Quantum Mechanics and Relativistic Field Theory*. Martino Fine Books, Eastford (2012).
32. Khrennikov, A. (2017). The present situation in quantum theory and its merging with general relativity. *Found. Phys.*, *47*(8), 1077-1099.
33. Zeilinger, A. (1999). A foundational principle for quantum mechanics, *Found. Phys.* *29* (4), 631--643.