EFFECTIVE VERSION OF RATNER'S EQUIDISTRIBUTION THEOREM FOR $SL(3, \mathbb{R})$

LEI YANG

ABSTRACT. In this paper, we will prove an effective version of Ratner's equidistribution theorem for unipotent orbits in $SL(3,\mathbb{R})/SL(3,\mathbb{Z})$ with a natural Diophantine condition.

1. Introduction

1.1. Ratner's theorem and its effective versions. In 1991, Ratner [Rat91b] proved the following fundamental theorem on equidistribution of unipotent orbits in homogeneous spaces:

Theorem 1.1. Let G be a Lie group and Γ be a lattice in G, namely, the homogeneous space $X = G/\Gamma$ admits a G-invariant probability measure. Let $U = \{u(r) : r \in \mathbb{R}\}$ be a one-parameter unipotent subgroup of G. For any $x \in X$, the closure of the U-orbit Ux of x is a closed L-orbit Lx for some Lie subgroup $L \subset G$. Moreover, Ux is equidistributed in Lx with respect to the unique L-invariant probability measure μ_L , namely, for any $f \in C_c^{\infty}(X)$ (where $C_c^{\infty}(X)$ denotes the space of smooth functions on X with compact support),

$$\lim_{T \to \infty} \frac{1}{T} \int_{-T/2}^{T/2} f(u(r)x) dr = \int_{Lx} f d\mu_L.$$

Since Ratner's proof relies on her measure classification theorem (cf. [Rat90a], [Rat90b], [Rat91a]) which proves a conjecture by Dani [Dan81], it does not tell how fast it tends to equidistribution. Therefore, a natural question is whether we can make the equidistribution effective, namely, giving an explicit upper bound on the difference

$$\left| \frac{1}{T} \int_{-T/2}^{T/2} f(u(r)x) dr - \int_{Lx} f d\mu_L \right|$$

for given T>0. Moreover, effective versions of equidistribution of U-orbits have many applications to number theory, see [LM14], [EM22], [LM21], [LMW22], [Ven10], [NV21], [EMV09], [CY19], [BV16] and references therein for details. As a result, proving effective versions of Ratner's theorem has attracted much attention and has been a major challenge in homogeneous dynamics.

In recent years, people have made significant progress in establishing effective equidistribution results. For G being nilpotent, the effective version of

Key words and phrases. homogeneous dynamics, unipotent orbits, equidistribution, effective Ratner's theorem.

Ratner's theorem was established by Green and Tao [GT12]. For $G = SL(2, \mathbb{R})$, since the one-parameter unipotent subgroup is horospherical with respect to the diagonal subgroup, one can apply the thickening argument developed in Margulis's thesis (cf. [Mar04], [KM12]) and spectral gap results for unitary representations of semisimple groups to establish effective equidistribution. See [Sar82], [Bur90], [FF03], [SU15], [Str04] and [Str13] for works in this setting. In fact, equidistribution for this case was proved before Ratner's theorem, see [Fur73], [DS84]. Using similar arguments one can also establish effective equidistribution results for horospherical unipotent orbits in homogeneous spaces. See [KM12], [Edw21] for results in this setting. Using techniques from analytic number theory, Strombergsson [Str15] established the effective equidistribution for $G/\Gamma = \mathrm{SL}(2,\mathbb{R}) \ltimes \mathbb{R}^2/\mathrm{SL}(2,\mathbb{Z}) \ltimes \mathbb{Z}^2$ and U being a oneparameter unipotent (and horospherical) subgroup in the semisimple part. Building on Strombergsson's result, Chow and the author [CY19] proved an effective equidistribution result for a special family of one-parameter unipotent orbits in $G/\Gamma = \mathrm{SL}(3,\mathbb{R})/\mathrm{SL}(3,\mathbb{Z})$. As an application, we proved that Gallagher's theorem in multiplicative Diophantine approximation holds for almost every point on any given planar straight line. The reader is also referred to [BV16] for a similar effective equidistibution result which has applications to number theory. Strombergsson's result was recently generalized by Kim [Kim21] to $SL(n,\mathbb{R}) \ltimes \mathbb{R}^n/SL(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$ with the unipotent subgroup being horospherical in the semisimple part. Recently, Lindenstrauss, Mohammadi and Wang [LMW22] established effective Ratner's theorem for one-parameter unipotent orbits in G/Γ where $G = \mathrm{SL}(2,\mathbb{R}) \times \mathrm{SL}(2,\mathbb{R})$ or $\mathrm{SL}(2,\mathbb{C})$. The reader is referred to [EMV09] and [EMMV20] for effective equidistribution of closed orbits of maximal semisimple subgroups which can be regarded as an effective version of a result by Mozes and Shah [MS95].

Concerning effective density, there are also significant results. Lindenstrauss and Margulis proved [LM14] an effective density result for one-parameter unipotent orbits in $SL(3,\mathbb{R})/SL(3,\mathbb{Z})$ and applied it to prove an effective version of Oppenheim's conjecture. Recently, Lindenstrauss and Mohammadi [LM21] established an optimal effective density result or unipotent orbits in G/Γ for $G = SL(2,\mathbb{R}) \times SL(2,\mathbb{R})$ or $SL(2,\mathbb{C})$.

1.2. **Notation.** Throughout this paper we will fix the following notation.

Given a normed vector space V and r > 0, let $B_V(r)$ denote the set of vectors in V with norm $\leq r$. Given a Lie group J and r > 0, let $B^J(r)$ denote the r-neighborhood of the identity in J. Given a quantity \mathcal{A} , let $O(\mathcal{A})$ denote a quantity whose absolute value is $\leq C\mathcal{A}$ where C > 0 is an absolute constant. By $\mathcal{A} \ll \mathcal{B}$, we mean $\mathcal{A} = O(\mathcal{B})$. For an interval $I \subset \mathbb{R}$, let |I| denote the length of I. For I > 0, let I denote the interval $I \subset I$ denote the interval I denote the interval $I \subset I$ denote the int

1.3. **Main results.** In this paper we will focus on the following case: Let $G = \mathrm{SL}(3,\mathbb{R}), \ \Gamma = \mathrm{SL}(3,\mathbb{Z}), \ X = G/\Gamma.$ Let $U = \{u(r) : r \in \mathbb{R}\}$ be a

one-parameter unipotent subgroup of G defined as follows:

$$u(r) := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & r \\ 0 & 0 & 1 \end{bmatrix}. \tag{1.1}$$

Let μ_G denote the unique G-invariant probability measure on X.

For $\beta > 0$, let us define

$$X_{\beta} := \left\{ x = g\Gamma \in X : ||g\mathbf{v}|| \geqslant \beta \text{ for any } \mathbf{v} \in \mathbb{Z}^3 \text{ or } \bigwedge^2 \mathbb{Z}^3. \right\}$$
 (1.2)

By Mahler's criterion, every compact subset of X is contained in some X_{β} . For $t \in \mathbb{R}$, let us denote

$$a_0(t) := \begin{bmatrix} e^{t/3} & & \\ & e^{-t/6} & \\ & & e^{-t/6} \end{bmatrix} \in G,$$
 (1.3)

and

$$a(t) := \begin{bmatrix} 1 & & & \\ & e^{t/2} & & \\ & & e^{-t/2} \end{bmatrix} \in G.$$
 (1.4)

We will prove the following result concerning effective equidistribution of U-orbit:

Theorem 1.2. There exist $\eta > 0$ and $t_0 > 1$ such that for any $t \ge t_0$ and any $x \in X$, one of the following holds:

(1) For any $f \in C_c^{\infty}(X)$, we have

$$\left| \int_{[1]} f(u(re^t)x) dr - \int f d\mu_G \right| \ll e^{-\eta t} ||f||_S,$$

where $\|\cdot\|_S$ denotes a fixed Sobolev norm;

(2)
$$a(-t)a_0(\ell)x \notin X_{e^{-|\ell|/3}}$$
 for some $0.99t \leqslant |\ell| \leqslant 1.01t$.

Proving Theorem 1.2 is equivalent to proving the following theorem:

Theorem 1.3. Let $\eta, t_0 > 0$ be as above. For any $t \ge t_0$, and any $x \in G/\Gamma$, one of the following holds:

(1) For any $f \in C_c^{\infty}(X)$, we have

$$\left| \int_{[1]} f(a(t)u(r)x) dr - \int f d\mu_G \right| \ll e^{-\eta t} ||f||_S;$$

(2) $a_0(\ell)x \notin X_{e^{-|\ell|/3}}$ for some $0.99t \leqslant |\ell| \leqslant 1.01t$.

In fact, it follows from the following equality:

$$\int_{[1]} f(a(t)u(r)x) dr = \int_{[1]} f(u(re^t)y) dr,$$

where y = a(t)x.

Using Theorem 1.2, we can prove the following two corollaries on effective equidistribution of expanding curves under translates of diagonal subgroups:

Corollary 1.4. Let us denote

$$a_1(t) := a(t)a_0(t) = \begin{bmatrix} e^{t/3} & & \\ & e^{t/3} & \\ & & e^{-2t/3} \end{bmatrix}.$$

Let $\mathbf{w} = (w_1, w_2) \in \mathbb{R}^2$ be a vector with Diophantine exponent $\omega(\mathbf{w}) = \kappa \leq 0.6$, namely, for any positive integer $q \in \mathbb{Z}_+$, we have

$$\max\{\langle qw_1\rangle, \langle qw_2\rangle\} \gg q^{-\kappa},$$

where $\langle \cdot \rangle$ denotes the distance to the nearest integer. For $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$ let us denote

$$n(\mathbf{v}) := \begin{bmatrix} 1 & v_2 \\ 1 & v_1 \\ & 1 \end{bmatrix}.$$

Let us define

$$\varphi:[1]\to\mathbb{R}^2$$

by $\varphi(r) := (w_1r + w_2, r)$. Then there exist $\eta, t_0 > 0$, such that for any $t \ge t_0$ and any $f \in C_c^{\infty}(X)$ we have

$$\left| \int_{[1]} f(a_1(t)n(\varphi(r))\Gamma) dr - \int f d\mu_G \right| \ll e^{-\eta t} ||f||_S.$$

Proof assuming Theorem 1.3. Note that

$$a_1(t)n(\varphi(r)) = z(w_1)a(t)u(r)a_0(t)n^*(-w_1, w_2),$$

where

$$z(w_1) := \begin{bmatrix} 1 & w_1 \\ & 1 \\ & & 1 \end{bmatrix},$$

and

$$n^*(-w_1, w_2) := \begin{bmatrix} 1 & -w_1 & w_2 \\ & 1 & \\ & & 1 \end{bmatrix}.$$

Then for any $f \in C_c^{\infty}(X)$ with zero integral, we have that

$$\int_{[1]} f(a_1(t)n(\varphi(r))\Gamma) dr = \int_{[1]} f_{w_1}(a(t)u(r)a_0(t)n^*(-w_1, w_2)\Gamma) dr,$$

where $f_{w_1}(x) = f(z(w_1)x)$. Note that since μ_G is G-invariant, f_{w_1} also has zero integral.

Let $x = a_0(t)n^*(-w_1, w_2)\Gamma$. Note that the Diophantine condition on **w** ensures that $a_0(\ell)x \in X_{e^{-|\ell|/3}}$ for any $0.99t \leq |\ell| \leq 1.01t$. Applying Theorem 1.3 with $x = a_0(t)n^*(-w_1, w_2)\Gamma$ and f_{w_1} , we have that

$$\left| \int_{[1]} f_{w_1}(a(t)u(r)a_0(t)n^*(-w_1, w_2)\Gamma) dr \right| \ll e^{-\eta t} ||f_{w_1}||_S.$$

Noting that $||f_{w_1}||_S \ll ||f||_S$, we complete the proof.

Remark 1.5. The ineffectively version of Corollary 1.4 was proved in [KdSSY21] for $\omega(\mathbf{w}) \leq 2$ using Ratner's theorem. The authors also proved that the equidistribution does not hold for $\omega(\mathbf{w}) > 2$. By analyzing the Diophantine condition in Theorem 1.3 more carefully, one can easily get a better upper bound on $\omega(\mathbf{w})$ which ensures effective equidistribution.

Corollary 1.6. Let us use the same notation as in Corollary 1.4. Let ψ : $[1] \to \mathbb{R}^2$ be a smooth non-degenerate curve in \mathbb{R}^2 , namely, derivatives of ψ span the whole space \mathbb{R}^2 at every $r \in [1]$. Then there exist $\eta, t_0 > 0$ such that for any $t \geq t_0$ and any $f \in C_c^{\infty}(X)$ we have

$$\left| \int_{[1]} f(a_1(t)n(\psi(r))\Gamma) dr - \int f d\mu_G \right| \ll e^{-\eta t} ||f||_S.$$

Proof. Let us denote $\psi(r) = (r, \psi_2(r))$. Let us divide [0, 1] into small pieces of size $e^{-t/2}$. Let us fix a small piece $\Delta(r_0) = [r_0 - 1/2e^{-t/2}, r_0 + 1/2e^{-t/2}]$. Then

$$\{a_1(t)n(\psi(r))\Gamma : r \in \Delta(r_0)\}\$$

can be approximated by

$$\{z(\psi_2'(r_0))a(t/2)u(r)x : r \in [-1/2, 1/2]\}$$

with exponentially small error, where

$$z(\psi_2'(r_0)) = \begin{bmatrix} 1 & \psi_2'(r_0) \\ & 1 \\ & & 1 \end{bmatrix},$$

as defined in Corollary 1.4, and

$$x = a_0(t/2)a_1(t/2)z(-\psi_2'(r_0))n(\psi(r_0))\Gamma.$$

Then if we can show that

$${a(t/2)u(r)x : r \in [1]}$$

effectively equidistributed, we will get that

$$\{a_1(t)n(\psi(r)): r \in \Delta(r_0)\}\$$

is effectively equidistributed. By Theorem 1.3, if we have that for any $0.49t \le |\ell| \le 0.51t$,

$$a_0(\ell)a_0(t/2)a_1(t/2)z(-\psi_2'(r_0))n(\psi(r_0))\Gamma \in X_{e^{-t/10}},$$

then we are done. Now for a fixed $|\ell| \in [0.49t, 0.51t]$, let us estimate the measure of

$$\mathfrak{m}_{\ell} := \{ r_0 \in [1] : a_0(\ell) a_0(t/2) a_1(t/2) z(-\psi_2'(r_0)) n(\psi(r_0)) \Gamma \not \in X_{e^{-t/10}} \}.$$

By [BKM01, Theorem 1.4], we have that $|\mathfrak{m}_{\ell}| = O(e^{-\eta_2 t})$ for some constant $\eta_2 > 0$ independent of ℓ . Let us remove all \mathfrak{m}_{ℓ} from [1] and get a subset $\mathfrak{m} \subset [1]$. Then we have that

$$|[1] \setminus \mathfrak{m}| = O(te^{-\eta_2 t}),$$

and for any $r_0 \in \mathfrak{m}$,

$$\{a_1(t)n(\psi(r))\Gamma : r \in \Delta(r_0)\}\$$

is effectively equidistributed in X. Combining these two facts we conclude that the whole orbit

$$\{a_1(t)n(\psi(r))\Gamma: r \in [1]\}$$

is effectively equidistributed. This completes the proof.

Remark 1.7. The ineffective version of Corollary 1.6 was proved in [Sha09] using Ratner's theorem.

Acknowledgements. The author thanks Wen Huang, Dmitry Kleinbock, Elon Lindenstrauss, Amir Mohammadi, Nimish Shah, Ralf Spatzier, Zhiren Wang and Barak Weiss for valuable discussions and Victor Beresnevich, Sam Chow, Wen Huang, Elon Lindenstrauss, Jens Marklof and Pengyu Yang for valuable comments on an earlier version of the paper. The author is supported in part by NSFC grant No. 12171338.

2. Preliminaries

In this section we recall some basic facts on $SL(3,\mathbb{R})$ and its Lie algebra which will be used in the proof of our main theorem.

Let $G = SL(3, \mathbb{R})$, and $H \subset G$ be the following subgroup of G:

$$H := \left\{ \begin{bmatrix} 1 \\ h \end{bmatrix} \in G : h \in \mathrm{SL}(2, \mathbb{R}) \right\}. \tag{2.1}$$

Clearly H is isomorphic to $SL(2, \mathbb{R})$.

Let \mathfrak{g} and \mathfrak{h} denote the Lie algebras of G and H, respectively.

Consider the adjoit action of H on \mathfrak{g} , we have the following decomposition:

$$\mathfrak{g} = \mathfrak{h} + \mathfrak{r}_0 + \mathfrak{r}_1 + \mathfrak{r}_2, \tag{2.2}$$

where $\mathfrak{r}_0, \mathfrak{r}_1, \mathfrak{r}_2$ are invariant subspaces with respect to the adjoint action of H, and $\dim \mathfrak{r}_1 = \dim \mathfrak{r}_2 = 2$, $\dim \mathfrak{r}_0 = 1$. In particular, $\mathfrak{r}_0 = \mathbb{R}\mathfrak{a}_0$ where

$$\mathfrak{a}_0 = \begin{bmatrix} 1/3 & 0 & 0 \\ 0 & -1/6 & 0 \\ 0 & 0 & -1/6 \end{bmatrix};$$

 $\mathfrak{r}_1 = \mathbb{R}\mathbf{v}_1 + \mathbb{R}\mathbf{v}_2$ where

$$\mathbf{v}_1 = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

and

$$\mathbf{v}_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix};$$

 $\mathbf{r}_2 = \mathbb{R}\mathbf{w}_1 + \mathbb{R}\mathbf{w}_2$ where

$$\mathbf{w}_1 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

and

$$\mathbf{w}_2 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

The adjoint action of H on \mathfrak{r}_1 and \mathfrak{r}_2 are given as follows: the adjoint action of H on \mathfrak{r}_1 is the same as the standard action of $\mathrm{SL}(2,\mathbb{R})$ on \mathbb{R}^2 if we choose

$$\{\mathbf{v}_1, \mathbf{v}_2\}$$
 as the basis; for h corresponding to $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and $\mathbf{w} = x_1 \mathbf{w}_1 + x_2 \mathbf{w}_2$,

$$Ad(h)\mathbf{w} = (ax_1 - bx_2)\mathbf{w}_1 + (-cx_1 + dx_2)\mathbf{w}_2.$$

Let $\{\mathfrak{a},\mathfrak{u},\mathfrak{u}^*\}$ denote the standard basis of \mathfrak{h} , where \mathfrak{a} corresponds to

$$\begin{bmatrix} 1/2 & 0 \\ 0 & -1/2 \end{bmatrix} \in \mathfrak{sl}_2(\mathbb{R}),$$

u corresponds to

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \in \mathfrak{sl}_2(\mathbb{R}),$$

and \mathfrak{u}^* corresponds to

$$\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \in \mathfrak{sl}_2(\mathbb{R}).$$

Let us denote

$$A := \{a(t) = \exp(t\mathfrak{a}) : t \in \mathbb{R}\} \subset H,$$

$$U := \{u(r) := \exp(r\mathfrak{u}) : r \in \mathbb{R}\} \subset H.$$

and

$$U^* := \{u^*(t) = \exp(r\mathfrak{u}^*) : r \in \mathbb{R}\} \subset H.$$

The adjoint action of $a_0(t)$ on \mathfrak{g} is as follows:

$$\operatorname{Ad}(a_0(t))\mathbf{h} = \mathbf{h}, \text{ for } \mathbf{h} \in \mathfrak{h},$$

$$\operatorname{Ad}(a_0(t))\mathbf{v} = e^{-t/2}\mathbf{v}, \text{ for } \mathbf{v} \in \mathfrak{r}_1,$$

and

$$Ad(a_0(t))\mathbf{w} = e^{t/2}\mathbf{w}, \text{ for } \mathbf{w} \in \mathfrak{r}_2.$$

Let us denote

$$b(t) := a(t)a_0(-t) = \begin{bmatrix} e^{-t/3} & & \\ & e^{2t/3} & \\ & & e^{-t/3} \end{bmatrix} \in G,$$
 (2.3)

and

$$a_1(t) := a(t)a_0(t) = \begin{bmatrix} e^{t/3} & & \\ & e^{t/3} & \\ & & e^{-2t/3} \end{bmatrix} \in G.$$
 (2.4)

Let us denote

$$\mathbf{r}^+ := \mathbb{R}\mathbf{w}_1 + \mathbb{R}\mathbf{v}_1, \tag{2.5}$$

and

$$\mathbf{r}^- := \mathbb{R}\mathbf{w}_2 + \mathbb{R}\mathbf{v}_2,\tag{2.6}$$

Let us denote by $p_{1,2}: \mathfrak{g} \to \mathfrak{r}_1 + \mathfrak{r}_2$ the projection of \mathfrak{g} to $\mathfrak{r}_1 + \mathfrak{r}_2$. For j = 0, 1, 2, let $p_j: \mathfrak{g} \to \mathfrak{r}_j$ denote the projection from \mathfrak{g} to \mathfrak{r}_j . Let $p_+, p_-, p_{\mathbf{w}_j}, p_{\mathbf{v}_j}$ and $p_{\mathfrak{u}^*}$ denote the projection to $\mathbf{r}^+, \mathbf{r}^-, \mathbb{R}\mathbf{w}_j, \mathbb{R}\mathbf{v}_j$ and $\mathbb{R}\mathfrak{u}^*$, respectively.

2.1. **Outline of the proof.** In this subsection we will give the outline of the proof.

The proof is inspired by Ratner's original proof of her measure rigidity theorem [Rat90a] and recent papers by Lindenstrauss, Mohammadi and Wang [LM21, LMW22]. However, compared with previous works, we take a quite different approach in this paper.

We will start with $\mathcal{F}_0 = a(s)u([1])x$ for $s = t - \delta_1 t$ and analyze the dimension in directions transversal to the H-orbit direction.

In §5.3, we will show that \mathcal{F}_0 has certain dimension control in the transversal direction unless condition (2) in Theorem 1.3 holds. Here we are allowed to remove an exponentially small proportion from \mathcal{F}_0 . At this step quantitative non-divergence results proved in §4 are needed. This is the starting point of our proof. The argument in this part is similar to the corresponding parts in [LM21, LMW22].

§5.4 is the crucial part. In this subsection, we will construct a sequence $\{\mathcal{F}_i: i \in \mathbb{N}\}$ of U-orbits starting from \mathcal{F}_0 where $\mathcal{F}_{i+1} = a(s_i)\mathcal{F}_i$ for each $i \geq 0$. We will prove that \mathcal{F}_{i+1} has a better dimension control compared with \mathcal{F}_i (at a larger scale as a cost). To prove this, we introduce a Kakeya-type model to study the divergence of nearby U-orbits and calculate the weighted intersection number of the Kakeya-type model. The outcome of the calculation is the following: Either \mathcal{F}_{i+1} has a better dimension control, or the orbit is concentrated on a small neighborhood of a piece of an orbit of a subgroup isomorphic to $\mathrm{SL}(2,\mathbb{R})\ltimes\mathbb{R}^2$. The latter case can be ruled out by our Diophantine condition (see Proposition 5.3). At each step we are allowed to remove an exponentially small proportion. This part is novel compared with previous works.

The inductive construction given as above will provide a \mathcal{F}_n which has good dimension control along the directions transversal to H-orbit direction. Then we can apply Proposition 3.1 from §3 to conclude the effective equidistribution. Proposition 3.1 can be proved by following a Van der Corput argument due to Venkatesh (see [Ven10, §3] and [LM21, Proposition 4.2] for details).

3. High transversal dimension to effective equidistribution

We will need the following statement to get effective equidistribution from high transversal dimension control:

Proposition 3.1. There exists $\theta > 0$ such that the following holds: For any $\varepsilon > 0$, there exist $\eta, s_0 > 0$ depending on $\varepsilon > 0$ such that for any $s' \geqslant s_0$ with

 $\beta = e^{-\varepsilon^2 s'}$, any Borel probability measure ρ on $[\beta]^2$ with dimension larger than $2 - \theta$ at scale s', that is, for any interval $I \subset [\beta]^2$ of length $\beta e^{-s'}$, we have

$$\rho(I) \leqslant |I|^{2-\theta},$$

any function $f \in C_c^{\infty}(X)$, and any $x \in X$, we have

$$\left| \int_{[1]^2} \int_{[1]} f(a(2s')u(r) \exp(\beta w_1 \mathbf{w}_1 + \beta v_1 \mathbf{v}_1) x) dr d\rho(w_1, v_1) - \int f d\mu_G \right| \ll e^{-\eta s'} ||f||_S.$$

Proof. The statement can be proved by following the proof of [LM21, Proposition 4.2] step by step. \Box

4. Quantitative non-divergence

This section is devoted to proving the following quantitative non-divergence result:

Proposition 4.1. There exists $\alpha > 0$ such that the following holds: For any $\beta, s > 0$, if $x \in X$ satisfies that $a_0(-s)x, a_0(s)x \notin X_{e^{-s/3}}$, then,

$$|\{r \in [1] : a(s)u(r)x \notin X_{\beta}\}| \ll \beta^{\alpha}.$$

Proof. Let us denote $x=g\Gamma$. By the Kleinbock-Margulis quantitative non-divergence theorem (cf. [Kle08, Theorem 2.2], [KM98, Theorem 5.2]), if the statement does not hold then there exists $\mathbf{v} \in \bigwedge^j \mathbb{Z}^3$ (where j=1 or 2), such that

$$\max_{r \in [1]} \|a(s)u(r)g\mathbf{v}\| \leqslant 1.$$

Case 1. $\mathbf{v} \in \mathbb{Z}^3$: Let us denote $\mathbf{v}' = a(s)g\mathbf{v} = (v_1', v_2', v_3')$. Then

$$a(s)u(r)g\mathbf{v} = u(re^s)\mathbf{v}' = (v_1', v_2' + re^s v_3', v_3').$$

Then $||a(s)u(r)g\mathbf{v}|| \leq 1$ implies that $|v_1'| \leq 1$, $|v_2' + re^sv_3'| \leq 1$ for any $r \in [1]$. The latter easily implies that $|v_2'| \leq 1$ and $|v_3'| \leq e^{-s}$. This implies that $||a_1(-s)\mathbf{v}'|| \leq e^{-s/3}$. Noting that $a_1(s) = a(s)a_0(s)$, we complete the proof for Case 1.

Case 2. $\mathbf{v} \in \bigwedge^2 \mathbb{Z}^3$: Let us denote $\mathbf{v}' = a(s)g\mathbf{v} = (v_1', v_2', v_3')$ where we use the coordinates with respect to the basis $\{\mathbf{e}_2 \wedge \mathbf{e}_3, \mathbf{e}_1 \wedge \mathbf{e}_3, \mathbf{e}_1 \wedge \mathbf{e}_2\}$. Then

$$a(s)u(r)g\mathbf{v} = u(re^s)\mathbf{v}' = (v_1', v_2', v_3' + re^s v_2').$$

By repeating the same argument as in Case 1, we have $|v_1'|, |v_3'| \le 1$ and $|v_2'| \le e^{-s}$ which implies that $||b(-s)\mathbf{v}'|| \le e^{-s/3}$. Noting that $b(s) = a(s)a_0(-s)$, we complete the proof for **Case 2**.

Note that for any $\beta > 0$ small enough and any $x \in X_{\beta^{1/4}}$, $B^G(\beta)x$ embeds into X injectively.

We will need the following lemma:

Lemma 4.2. For any $\beta > 0$, any $\ell > 1$ large enough (depending on β), and any $x \in X$ with $a(-\ell)x \in X_{\beta^{1/4}}$, if

$$\exp(r_1 e^{\ell} \mathbf{u} + r_2 e^{\ell/2} \mathbf{v}_1 + r_3 e^{\ell/2} \mathbf{w}_1) x = \exp(\mathbf{v}) x$$

where $r_1, r_2, r_3 \in [\beta]$, $\|(r_1, r_2, r_3)\| \ge \beta/4$, and $\mathbf{v} \in B_{\mathfrak{g}}(\beta)$. Let us write

$$\mathbf{v} = u\mathbf{u} + u^*\mathbf{u}^* + a\mathbf{a} + a_0\mathbf{a}_0 + \sum_{j=1}^2 w_j\mathbf{w}_j + v_j\mathbf{v}_j.$$

if we have $|u^*| < \beta e^{-\ell}$, then $||(v_2, w_2)|| \geqslant \beta e^{-\ell/2}$.

Proof. For a contradiction, let us assume that $|u^*| < \beta e^{-\ell}, |v_2, w_2| < \beta e^{-\ell/2}$. Let us denote $x = g\Gamma$. Then

$$\exp(r_1 e^{\ell} \mathbf{u} + r_2 e^{\ell/2} \mathbf{v}_1 + r_3 e^{\ell/2} \mathbf{w}_1) g = \exp(\mathbf{v}) g \gamma,$$

for some $\gamma \in \Gamma$. For $\ell > 1$ large enough, we have $\gamma \neq e$. Then we have

$$\exp(-\mathbf{v}) \exp(r_1 e^{\ell} \mathbf{u} + r_2 e^{\ell/2} \mathbf{v}_1 + r_3 e^{\ell/2} \mathbf{w}_1) = g \gamma g^{-1}.$$

Then

$$a(-\ell)g\gamma g^{-1}a(\ell) = a(-\ell)\exp(-\mathbf{v})\exp(r_1e^{\ell}\mathbf{u} + r_2e^{\ell}\mathbf{w}_1)a(\ell)$$

= $\exp(-\mathrm{Ad}(a(-\ell))\mathbf{v})\exp(r_1\mathbf{u} + r_2\mathbf{v}_1 + r_3\mathbf{w}_1).$

Since $|u^*| < \beta e^{-\ell}$, $|v_2, w_2| < \beta e^{-\ell/2}$, we have $\|\operatorname{Ad}(a(-\ell))\mathbf{v}\| < \beta$. This implies that

$$a(-\ell)g\gamma g^{-1}a(\ell) \in B^G(\beta),$$

which means that $B^G(\beta)a(-\ell)x$ does not embed into X injectively. This contradicts to that $a(-\ell)x \in X_{\beta^{1/4}}$. This completes the proof.

5. Dimension Control

5.1. **List of constants.** We first list all constants we will use later.

Let $\delta_1, \delta_2 = 10^{-5}$ and $d_0 = 10^{-6}$. Let $\theta > 0$ be the constant from Theorem 3.1. Without loss of generality, we can assume that $\theta < 10^{-10}$. Let $\alpha > 0$ be the constants as in Proposition 4.1 and $\alpha_2 = \alpha/4$.

Let
$$C_1 = 10d_0^{-1} = 10^7$$
, $\varepsilon_1 = 10^{-20}d_0\theta$, $N_1 = 10^5\varepsilon_1^{-1}$.
Let $\varepsilon = e^{-10^5N_1}$, $\beta = e^{-\varepsilon^3t}$, and $\delta_3 = 10^{-5}\theta\varepsilon$.

5.2. **Initial construction of transversal sets.** We first introduce some notation.

For $\ell > 1$ and a subspace $V \subset \mathfrak{r}_1 + \mathfrak{r}_2$ generated by some vectors from the canonical basis of $\mathfrak{r}_1 + \mathfrak{r}_2$, namely, $\{\mathbf{v}_{1,2}, \mathbf{w}_{1,2}\}$, let us define

$$Q(V, \ell) := \{ \exp(\mathbf{w}) : \mathbf{w} \in B_{\mathbf{r}_1 + \mathbf{r}_2}(\beta), \| p_V(\mathbf{w}) \| \le \beta e^{-\ell} \},$$
 (5.1)

where p_V denotes the projection to V with respect to the canonical basis.

Let us denote

$$Q^{H}(\ell) := \{ h = \exp(\mathbf{h}) \in B^{H}(\beta), \|p_{\mathfrak{u}^{*}}(\mathbf{h})\| \leqslant \beta e^{-\ell} \}, \tag{5.2}$$

and

$$R^{H}(\ell) := \{ h = \exp(\mathbf{h}) \in B^{H}(\beta), \|p_{\mathfrak{u}}(\mathbf{h})\| \leqslant \beta e^{-\ell} \}.$$
 (5.3)

For $s_1, s_2 \geqslant 0$, let us denote

$$Q_{s_2}^{s_1} := \{ \exp(\mathbf{w}) : \mathbf{w} \in \mathfrak{r}_1 + \mathfrak{r}_2 : \|p_+(\mathbf{w})\| \leqslant \beta e^{-s_1}, \|p_-(\mathbf{w})\| \leqslant \beta e^{-s_2} \}, \quad (5.4)$$
 and $Q_s := Q_s^s$.

Let $x \in X$ be such that (2) in Theorem 1.3 does not hold. Let us start with the normalized measure on $\mathcal{E} = a(s)u([1])x$ where $s = (1 - \delta_1)t > 0$. This is a U-orbit of length e^s . Let us define $\mathcal{F} \subset \mathcal{E}$ as follows:

Definition 5.1. $\mathcal{F} \subset \mathcal{E}$ is defined by removing from \mathcal{E} points $y = a(s)u(r)x \in \mathcal{E}$ satisfying $a_0(\ell')a(-\ell)y \notin X_{\beta^{1/4}}$ for some $0 \leqslant \ell \leqslant 2\delta_2 t$ and $-2\delta_1 t \leqslant \ell' \leqslant 2\delta_1 t$. Let $\mu_{\mathcal{F}}$ denote the normalized U-orbit measure on \mathcal{F} .

By Proposition 4.1 and our assumption on x, it is easy to show that the removed proportion is $O(s^2\beta^{\alpha_2})$. Therefore,

$$\mu_{\mathcal{E}} = \mu_{\mathcal{F}} + O(s^2 \beta^{\alpha_2}), \tag{5.5}$$

where $\mu_{\mathcal{E}}$ denotes the normalized *U*-invariant measure on \mathcal{E} .

5.3. **Initial dimension bound.** We shall prove the following proposition:

Proposition 5.2. For any $\ell \in [\delta_3 t, \delta_1 t]$, and any $x_0 \in \mathcal{F}$, we have

$$\mu_{\mathcal{F}}(Q_{\ell}B^{A_0H}(\beta)x_0) \leqslant e^{-d_0\ell}.$$

Proof. Let $\eta_1 = 100d_0$. Let us divide a(s)u([1])x into pieces of length $e^{\eta_1\ell}$.

It suffices to show that for each piece $\tilde{\mathcal{E}}$ with $\tilde{\mathcal{F}} := \tilde{\mathcal{E}} \cap \mathcal{F} \neq \emptyset$, and any $x_0 \in \tilde{\mathcal{F}}$,

$$|\tilde{\mathcal{F}} \cap Q_{\ell} B^{A_0 H}(\beta) x_0| \leqslant e^{-d_0 \ell} e^{\eta_1 \ell}.$$

For a contradiction, let us assume that the statement does not hold for some $x_0 \in \tilde{\mathcal{F}}$. Then we can find at least $e^{\eta_2 \ell}$ (where $\eta_2 = \eta_1/4$) many $x \in \tilde{\mathcal{F}} \cap Q_\ell B^{A_0 H}(\beta) x_0, r_x \in [1]$ such that

$$u(r_x e^{2d_0 \ell})x = \exp(\mathbf{v}_x)a_0^x h_x x,$$

where $a_0^x \in B^{A_0}(\beta)$, $h_x \in B^H(\beta)$, and $\mathbf{v}_x \in B_{\mathfrak{r}_1+\mathfrak{r}_2}(\beta e^{-\ell})$. Then x satisfies $x \in X_{\beta^{1/4}}$ and $a(-2d_0\ell)x \in X_{\beta^{1/4}}$. By Lemma 4.2, if we write $h_x = \exp(\mathbf{h}_x)$, we have $\|p_{\mathfrak{u}^*}(\mathbf{h}_x)\| \geqslant \beta e^{-2d_0\ell}$.

Let us write $x = u_x x_0$ where $u_x = u(re^{\eta_1 \ell})$ for some $r \in [1]$. Let us fix a representative $g_0 \in G$ of x_0 , then we have

$$u(r_x e^{d_0 \ell}) u_x q_0 = \exp(\mathbf{v}_x) a_0^x h_x u_x q_0 \gamma_x,$$

for some $\gamma_x \in \Gamma$. The above equality is equivalent to

$$g_0 \gamma_x g_0^{-1} = (a_0^x h_x u_x)^{-1} \exp(-\mathbf{v}_x) u(r_x e^{2d_0 \ell}) u_x.$$
 (5.6)

Without loss of generality, we can assume that for different x and x' as above, $|u_{x'}^{-1}u_x| \ge e^{5d_0\ell}$.

For each x we get $\gamma_x \in \Gamma$. We claim that those γ_x 's are different. In fact, if there exist x, x' such that $\gamma_x = \gamma_{x'}$, then we have

$$(a_0^x h_x u_x)^{-1} \exp(-\mathbf{v}_x) u(r_x e^{2d_0 \ell}) u_x$$

= $(a_0^{x'} h_{x'} u_{x'})^{-1} \exp(-\mathbf{v}_{x'}) u(r_{x'} e^{2d_0 \ell}) u_{x'}.$

This implies that

$$\exp(\mathbf{v}_x')(a_0^x)^{-1}u_x^{-1}h_x^{-1}u_xu(r_xe^{2d_0\ell})$$

$$=\exp(\mathbf{v}_{x'}')(a_0^{x'})^{-1}u_{x'}^{-1}h_{x'}^{-1}u_{x'}u(r_{x'}e^{2d_0\ell}),$$

where $\mathbf{v}'_x = -\mathrm{Ad}((a_0^x h_x u_x)^{-1})\mathbf{v}_x$ and $\mathbf{v}'_{x'}$ denotes the same expression with x replaced by x'. By comparing the $\mathfrak{r}_1 + \mathfrak{r}_2$, A_0 and H components of both sides, we have $\exp(\mathbf{v}'_x) = \exp(\mathbf{v}'_{x'})$, $a_0^x = a_0^{x'}$ and

$$u_x^{-1}h_x^{-1}u_xu(r_xe^{2d_0\ell}) = u_{x'}^{-1}h_{x'}^{-1}u_{x'}u(r_{x'}e^{2d_0\ell}).$$

This implies that

$$u(-L)h_x u(L) = u(re^{2d_0\ell})h_{x'}, (5.7)$$

where $u(L) = u_x u_{x'}^{-1}$ and $r = r_x - r'_{x'}$. Note that $|r| \leq 1$ and $|L| \geq e^{10d_0\ell}$. $|r| \leq 1$ implies that the norm of the right hand side is $\leq e^{2d_0\ell}$. On the other hand, the left hand side is equal to

$$\exp(\operatorname{Ad}(u(-L))\mathbf{h}) = \exp(\operatorname{Ad}(u(-L))(a_0\mathfrak{a} + a_1\mathfrak{u} + a_2\mathfrak{u}^*))$$

whose \mathfrak{u} coordinate is $a_1 + a_0 L + a_2 L^2$. Since $|a_2| \geqslant \beta e^{-2d_0 \ell}$ and $|L| \geqslant e^{5d_0 \ell}$, we have that

$$|a_1 + a_0 L + a_2 L^2| \geqslant \beta e^{8d_0 \ell}$$
.

Therefore, we have that the equality (5.7) is impossible to hold and conclude the claim.

Let us consider the adjoint action of G on $\bigoplus_{i=1}^{8} \wedge^{j} \mathfrak{g}$ and denote

$$v_H = \mathfrak{a} \wedge \mathfrak{u} \wedge \mathfrak{u}^*$$
.

Then the stabilizer of v_H

$$\operatorname{Stab}(v_H) = A_0 H.$$

Then (5.6) implies that

$$\gamma_x g_0^{-1} v_H = \exp(\operatorname{Ad}(g_0^{-1}) \mathbf{v}_x') g_0^{-1} v_H.$$

The norm of $g_0^{-1}v_H$ is $\leqslant \|g_0^{-1}\|^{300}$. Let us estimate the norm of \mathbf{v}_x' . Note that $\|u_x\| \leqslant e^{\eta_1 \ell}$, we have that the norm of $(a_0^x h_x u_x)^{-1}$ is bounded by $2e^{\eta_1 \ell}$. Therefore, $\|\mathbf{v}_x'\| \leqslant e^{-\ell + \eta_1 \ell}$. Let us denote $v_0 = g_0^{-1} v_H / \|g_0^{-1} v_H\|$, then we have

$$\|\gamma_x v_0 - v_0\| \le \|\mathbf{v}_x'\| \le e^{-(1-\eta_1)\ell}.$$

Considering the group generated by γ_x 's, we have the following two cases:

Case 1: $\langle \gamma_x \rangle$ is abelian.

Case 2: $\langle \gamma_x \rangle$ is not abelian.

For Case 1, we claim that there exists γ_x whose unipotent part is not trivial. In fact, if every γ_x is diagonizable, then they belong to a maximal torus in G. Note that $\|\gamma_x\| \leq e^{\eta_1 \ell}$, we have that there are at most ℓ^{100} different

 γ_x 's which contradicts to the fact that there are $e^{\eta_2\ell}$ different γ_x 's. Note that $\gamma_x \in \mathrm{SL}(3,\mathbb{Z})$, if γ_x has nontrivial unipotent part, it must be unipotent. By repeating the same argument as in [EMV09], we can find $g' \in G$ satisfying that $\gamma_x g'^{-1} v_H = g'^{-1} v_H$ and

$$||g' - g_0|| \le ||\gamma_x v_0 - v_0|| ||\gamma_x||^{10} \le e^{-(1-\eta_1)\ell} e^{10\eta_1 \ell} \le e^{-\ell/2}$$

Then $g'\gamma_xg'^{-1} \in A_1H$. Since γ_x is unipotent, we have $g'\gamma_xg'^{-1} \in H$. We claim that the lattice $g'\mathbb{Z}^3$ contains a nonzero vector \mathbf{p} satisfying $\|\mathbf{p}\| \leq e^{2\eta_1\ell}$ and $\mathbf{p} \in \mathbb{R}\mathbf{e}_2 + \mathbb{R}\mathbf{e}_3$. In fact, we can find a basis $\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$ of $g'\mathbb{Z}^3$ such that $\|\mathbf{p}_j\| \leq \|g'\| \leq 2\|g_0\| \leq 2\beta^{-1}$ for j = 1, 2, 3. Since $g'\gamma_xg'^{-1}$ is a unipotent element in H, its fixing vectors

$$V(g'\gamma_x g'^{-1}) := \{ \mathbf{p} \in \mathbb{R}^3 : g'\gamma_x g'^{-1} \mathbf{p} = \mathbf{p} \}$$

has dimension 2. Therefore, $g'\gamma_xg'^{-1}\mathbf{p}_j \neq \mathbf{p}_j$ for some $j \in \{1, 2, 3\}$. On the other hand, we have $g'\gamma_xg'^{-1}\mathbf{p}_j \in g'\mathbb{Z}^3$. Thus,

$$g'\gamma_x g'^{-1}\mathbf{p}_j - \mathbf{p}_j \in g'\mathbb{Z}^3.$$

Moreover, since $g'\gamma_x g'^{-1} \in H$, it is easy to see that $\mathbf{p} := g'\gamma_x g'^{-1}\mathbf{p}_j - \mathbf{p}_j \in \mathbb{R}\mathbf{e}_2 + \mathbb{R}\mathbf{e}_3$. Its norm is bounded by

 $\|\mathbf{p}\| \leqslant \|\mathbf{p}_j\| + \|g'\gamma_x g'^{-1}\|\|\mathbf{p}_j\| \leqslant \|\mathbf{p}_j\| + \|g'\|\|\gamma_x\|\|g'^{-1}\|\|\mathbf{p}_j\| \leqslant 2\beta^{-1} + e^{\eta_1 \ell}\beta^{-10} \leqslant e^{2\eta_1 \ell}.$

Then for any $\eta_3 > 0$,

$$||a_0(\eta_3\ell)\mathbf{p}|| = e^{-\eta_3\ell/6}||\mathbf{p}|| \le e^{-(\eta_3/6 - 2\eta_1)\ell}.$$

Let us choose $\eta_3 = 24\eta_1$. Then it is easy to see that

$$||a_0(\eta_3\ell)g' - a_0(\eta_3\ell)g_0|| \le 1,$$

and $a_0(\eta_3\ell)g'\Gamma \not\in X_{e^{-2\eta_1\ell}}$. This implies that $a_0(\eta_3\ell)g_0\Gamma \not\in X_{\beta^{1/4}}$ which contradicts to that $g_0\Gamma = x_0 \in \mathcal{F}$.

Let us consider Case 2. Let us take $\gamma = \gamma_x$ and $\gamma' = \gamma_{x'}$ not commuting. Using the same argument as in Case 1 (again using the argument in [EMV09]), we can find we can find $g' \in G$ satisfying that $\gamma g'^{-1}v_H = g'^{-1}v_H$, $\gamma' g'^{-1}v_H = g'^{-1}v_H$ and

$$||g' - g_0|| \leqslant e^{-\ell/2}$$
.

This implies that $g'\gamma g'^{-1}, g'\gamma' g'^{-1} \in A_1H$. Then their commutator

$$g'\gamma\gamma'\gamma^{-1}\gamma'^{-1}g'^{-1}\in H.$$

Then we can use the same argument as in Case 1 to deduce that $a_0(\eta_3 \ell)g_0\Gamma \notin X_{\beta^{1/4}}$ leading to the same contradiction.

This completes the proof of Proposition 5.2.

The following statement tells that the orbit can not be concentrated on a small neighborhood of a piece of $SL(2,\mathbb{R}) \ltimes \mathbb{R}^2$ orbit:

Proposition 5.3. For any $x \in \mathcal{F}$, j = 1, 2, and any $\ell \in [2\delta_3 t, \delta_1 t]$

$$\mu_{\mathcal{F}}(Q(\mathfrak{r}_j,\ell)B^{A_0H}(\beta)x) \leqslant e^{-d_0\ell/2}$$

Proof. Let us prove the statement for j = 1. The proof for j = 2 is similar. For a contradiction, let us assume that there exists $x \in \mathcal{F}$ such that

$$\mu_{\mathcal{F}}(Q(\mathfrak{r}_1,\ell)B^{A_0H}(\beta)x) > e^{-d_0\ell/2}.$$

Let us denote $\tilde{x} = a_0(\ell)x$ and $\tilde{\mu} = a_0(\ell)_*\mu_{\mathcal{F}}$. Noting that $a_0(\ell)Q(\mathfrak{r}_1,\ell)B^{A_0H}(\beta)x = Q_{\ell/2}B^{A_0H}(\beta)\tilde{x}$, we have that

$$\tilde{\mu}(Q_{\tilde{\ell}}B^{A_0H}(\beta)\tilde{x}) > e^{-d_0\tilde{\ell}},$$

where $\tilde{\ell} = \ell/2 \in [\delta_3 t, \delta_1 t]$. By repeating the argument in the proof of Proposition 5.2, we can show that $a_0(\eta_3 \tilde{\ell})\tilde{x} = a_0(\ell + \eta_3 \tilde{\ell})x \notin X_{\beta^{1/4}}$, which contradicts to that $x \in \mathcal{F}$. This completes the proof.

5.4. **Dimension Improvement.** This subsection is devoted to dimension improvement under the action of a(t).

Let us first introduce some definitions:

Definition 5.4. Given a probability measure μ on X, we say that μ is (d, ℓ) -good if for any $x_0 \in X$,

$$\mu(Q_{\ell}B^{A_0H}(\beta)x_0) \leqslant e^{-d\ell}.$$

We say that μ is $(d, [\ell_1, \ell_2])$ -good if it is (d, ℓ) -good for any $\ell \in [\ell_1, \ell_2]$.

According to this definition, Proposition 5.2 says that $\mu_{\mathcal{F}}$ is $(d_0, [\delta_3 t, \delta_1 t])$ -good.

Starting from $s_0 = \delta_1 t/2$ and d_0 , let us define $\{(d_i, s_i) : i \in \mathbb{N}\}$ by $s_{i+1} = s_i/2$ and $d_{i+1} = d_i + \varepsilon_1$.

Starting from $\mathcal{F}_0 = \mathcal{F}$, we will construct a sequence $\{\mathcal{F}_i : i \in \mathbb{N}\}$ such that every \mathcal{F}_{i+1} is constructed by removing a small proportion from $a(s_i)\mathcal{F}_i$.

In this subsection, we will prove that $\mu_{\mathcal{F}_{i+1}}$ has better dimension control than $\mu_{\mathcal{F}_i}$. Similarly to Definition 5.1, by removing an exponentially small proportion, we can assume that for any $x \in \text{supp}\mu_{\mathcal{F}_i}$, any $0 \leq \ell \leq 2\delta_2 t$ and $-2\delta_1 t \leq \ell' \leq 2\delta_1 t$, $a_0(\ell')a(-\ell)x \in X_{\beta^{1/4}}$. By repeating the proof of Proposition 5.2, we have that $\mu_{\mathcal{F}_i}$ is $(d_0, [\delta_3 t, \delta_1 t])$ -good.

Before stating the main statement of this subsection, let us introduce some notation.

For notational simplicity, for $s' > 0, \ell \ge 0$, let us denote

$$\Pi_{s',\ell} := a(2s')Q_{\ell}B^{A_0H}(\beta)a(-2s'), \tag{5.8}$$

$$\Pi_{s'} := \Pi_{s',0} = a(2s')B^G(\beta)a(-2s'), \tag{5.9}$$

$$\Omega_{s'} := Q_{3s'}^{s'} B^{A_0}(\beta) Q^H(2s'), \tag{5.10}$$

and

$$\Theta_{s'} := Q_{s'} B^{A_0}(\beta) Q^H(2s'). \tag{5.11}$$

For s' > 0, let us denote

$$\Sigma_{s'} := B^{A_0 A}(\beta) B^{U^*}(\beta e^{-2s'}) B_{\mathbf{r}^-}(\beta e^{-s'}) B_{\mathbf{r}^+}(\beta e^{s'}) B^U(\beta). \tag{5.12}$$

For $x \in X$, let us denote

$$\Pi(x) = \Pi_{s'}x,$$

$$\Sigma(x) = \Sigma_{s'}x.$$

 $\Sigma(x)$ can be regarded as a cross-section of $\Pi(x)$ along the *U*-orbits. For $\bar{x} \in \Sigma(x) \subset \Pi(x)$, let us denote

$$\Pi_{\ell}(\bar{x}) = \Pi_{s',\ell}\bar{x},$$

and

$$\Sigma_{\ell}(\bar{x}) = \Pi_{\ell}(\bar{x}) \cap \Sigma(x).$$

Then $\Sigma_{\ell}(\bar{x})$ can also be regarded as a cross-section of $\Pi_{\ell}(\bar{x})$ along the *U*-orbits. We first prove the following lemma:

Lemma 5.5. Let $\mu = a(s_i)_* \mu_{\mathcal{F}_i}$, $x \in X$ and $s' = s_i/2$. For any $\delta_3 t \leqslant \ell \leqslant \delta_1 t$ and any $\bar{x} \in \Pi(x)$,

$$\mu(\Sigma_{\ell}(\bar{x})) \leqslant e^{-2s'} e^{-d_0 \ell}.$$

Proof. Note that $\Sigma_{\ell}(\bar{x})$ is a cross-section of $\Pi_{\ell}(\bar{x})$ along the *U*-orbits, and μ is Lebesgue along the *U*-orbits. We have

$$\mu(\Sigma_{\ell}(\bar{x})) = e^{-2s'}\mu(\Pi_{\ell}(\bar{x})).$$

Note that

$$\mu(\Pi_{\ell}(\bar{x})) = \mu_{\mathcal{F}_i}(a(-2s')\Pi_{\ell}(\bar{x})),$$

and

$$a(-2s')\Pi_{\ell}(\bar{x}) = Q_{\ell}B^{A_0H}(\beta)x'$$

where $x' = a(-2s')\bar{x}$. By repeating the proof of Proposition 5.2 to \mathcal{F}_i we have

$$\mu_{\mathcal{F}_i}(Q_\ell B^{A_0 H}(\beta) x') \leqslant e^{-d_0 \ell},$$

which concludes the proof.

We then introduce a Kakeya-type model to analyze the dimension change along the U-orbits.

Definition 5.6. Let s' > 0 and μ be a probability measure on X defined by taking the normalized measure on a subset consisting of U-orbits of length $e^{2s'}$. Let us fix a playground of the form $\Pi(x) = \Pi_{s'}x \subset X$ and a cross-section $\Sigma(x) = \Sigma_{s'}x$. Let us cut $\Pi(x)$ into small pieces each of which is of the form $\Omega(y) = \Omega_{s'}y$. For each piece $\Omega(y) \in \Sigma(x)$, let us write

$$y = a_0(y)a(y)u^*(r^*(y))\exp(\mathbf{w}_y^-)\exp(\mathbf{w}_y^+)u(r(y))x,$$

where $a_0(y) \in B^{A_0}(\beta)$, $a(y) \in B^A(\beta)$, $\mathbf{w}_y^- = v_{2,y}\mathbf{v}_2 + w_{2,y}\mathbf{w}_2 \in \mathbf{r}^-$ and $\mathbf{w}_y^+ = v_{1,y}\mathbf{v}_1 + w_{1,y}\mathbf{w}_1 \in \mathbf{r}^+$, $r^*(y) \in [\beta e^{-2s'}]$, and $r(y) \in [\beta]$. By replacing y with another element in $\Omega(y)$, we may ignore $a_0(y)a(y)u^*(r^*(y))$ and write

$$y = \exp(\mathbf{w}_y^-) \exp(\mathbf{w}_y^+) u(r(y)) x.$$

Then we assign $\Omega(y)$ with a curve:

$$\mathcal{L}(y) := \{ (f_y(r), f_{v,y}(r), f_{w,y}(r) : r \in [e^{2s'}] \},$$

where f_y $f_{v,y}$ and $f_{w,y}$ are given as follows: Let us write

 $u(r) \exp(\mathbf{w}_{y}^{-}) \exp(\mathbf{w}_{y}^{+}) = a_{0}(r)a(r)u^{*}(f_{y}^{*}(r)) \exp(\mathbf{f}_{-,y}(r)) \exp(\mathbf{f}_{+,y}(r))u(f_{y}(r)),$

where $a(r) \in B^A(\beta)$, $a_0(r) \in B^{A_0}(\beta)$, $f_y^*(r) \in [\beta e^{-2s'}]$, $\mathbf{f}_{-,y}(r) \in \mathbf{r}^-$ and $\mathbf{f}_{+,y}(r) \in \mathbf{r}^+$. Then $f_{v,y}$ and $f_{w,y}$ are \mathbf{v}_1 and \mathbf{w}_1 , respectively, coordinates of $\mathbf{f}_{+,y}$. In fact, it is straightforward to check that

$$f_{v,y}(r) = \frac{2rv_{2,y}}{2 + rv_{2,y}w_{2,y}} + v_{1,y},$$
(5.13)

$$f_{w,y}(r) = \frac{-2rw_{2,y}}{2 - rv_{2,y}w_{2,y}} + w_{1,y},$$
(5.14)

and

$$f_y(r) = \frac{2r + 2rw_{1,y}v_{2,y}}{2 + rv_{2,y}w_{2,y}} + \frac{w_{1,y}v_{1,y} - f_{v,y}(r)f_{w,y}(r)}{2}$$
(5.15)

Let $\mathcal{T}(y)$ denote the $\beta e^{-s'}$ -neighborhood of $\mathcal{L}(y)$. We then assign $\mathcal{T}(y)$ with a weight

$$\mathcal{W}(y) = \mu(\Omega(y)).$$

For $\mathcal{T}(y_1)$ and $\mathcal{T}(y_2)$, let us define $\Delta(y_1, y_2)$ as follows: By replacing y_2 with another element in $\Omega(y_2)$, we can write

$$y_2 = \exp(\mathbf{w}^-) \exp(\mathbf{w}^+) u(r) y_1,$$

where $\mathbf{w}^- = v_2 \mathbf{v}_2 + w_2 \mathbf{w}_2 \in \mathbf{r}^-$. Then we define

$$\Delta(y_1, y_2) = (v_2, w_2). \tag{5.16}$$

Let us call the collection of tubes $\{\mathcal{T}(y)\}$ with weights $\{\mathcal{W}(y)\}$ the Kakeya-type model for $(\mu, \Pi(x))$.

The following lemma tells the connection between the dimension change along the U-orbits and the Kakeya-type model:

Lemma 5.7. Let us fix μ , s' > 0 and $\Sigma(x) \subset \Pi(x)$ as above. Then for any $\Omega(y_1)$ and $\Omega(y_2)$ from $\Sigma(x)$, the following two statements are equivalent:

- (1). There exist $L_1, L_2 \in [e^{2s'}]$ with $|L_1 L_2| \leq \beta |L_1|$ and $y'_j \in \Omega(y_j)$ for j = 1, 2 such that $u(L_1)y'_1$ and $u(L_2)y'_2$ are contained in the same neighborhood of the form $\Theta(z)$;
- (2). $\mathcal{T}(y_1)$ intersects $\mathcal{T}(y_2)$ at $\tilde{L} = f_{y_1}(\tilde{L}_1) = f_{y_2}(\tilde{L}_2)$ where $|\tilde{L} \tilde{L}_j| \leq \beta |\tilde{L}_j|$ and $|L_j \tilde{L}_j| \leq \beta |L_j|$ for j = 1, 2.

Moreover, if one of the above holds, then for any $|L'_1| \leq e^{-s'} ||\Delta(y_1, y_2)||^{-1}$, there exists L'_2 with $|L'_1 - L'_2| \leq \beta |L'_1|$ such that $u(L_1 + L'_1)y'_1$ and $u(L_2 + L'_2)y'_2$ are contained in the same neighborhood of the form $\Theta(z')$.

Proof. Without loss of generality, let us assume that $y_1 = x$ and denote y_2 by y.

By replacing y with another element in $\Omega(y)$, we can write

$$y = \exp(\mathbf{w}_y^-) \exp(\mathbf{w}_y^+) u(r(y)) x.$$

Suppose statement (1) holds for L_1, L_2 . Then there exist \tilde{L}_1, \tilde{L}_2 such that $|\tilde{L}_j - L_j| \leq \beta |L_j|$ for j = 1, 2 and $u(\tilde{L}_1)x, u(\tilde{L}_2)y$ are contained in the same $\Theta(z')$. Let us consider $u(\tilde{L}_2)y$:

$$u(\tilde{L}_2)y = u(\tilde{L}_2) \exp(\mathbf{w}_y^-) \exp(\mathbf{w}_y^+) u(r(y)) x$$

= $a_0(\tilde{L}_2) a(\tilde{L}_2) u^*(f_y^*(\tilde{L}_2)) \exp(\mathbf{f}_{-,y}(\tilde{L}_2)) \exp(\mathbf{f}_{+,y}(\tilde{L}_2)) u(f_y(\tilde{L}_2)) x.$

Note that

$$\|\mathbf{f}_{-,y}(\tilde{L}_2)\| \leqslant \beta e^{-s'}.$$

We then have

$$\|\mathbf{f}_{+,y}(\tilde{L}_2)\| \leqslant \beta e^{-s'}$$

which implies $\mathcal{T}(y)$ and $\mathcal{T}(x)$ intersect at $\tilde{L} = f_y(\tilde{L}_2)$. This proves that statement (2) holds.

Using similar argument, we can prove that (2) implies (1) and the second part of the lemma. \Box

Let us introduce a coordinate system for different $\Omega(z')$'s in a fixed $\Theta(z)$:

Definition 5.8. Given $\Theta(z)$, we can assign every $\Omega(z') \subset \Theta(z)$ a box of size $\beta e^{-3s'}$ in \mathbb{R}^2 as follows. By replacing z' with another element in $\Omega(z')$, we can write

$$z' = \exp(\mathbf{w}^-)z$$

for some $\mathbf{w}^- = v_2 \mathbf{v}_2 + w_2 \mathbf{w}_2 \in \mathbf{r}^-$. Then we can assign $\Omega(z')$ with the box of size $\beta e^{-3s'}$ centered at (v_2, w_2) and call it the coordinate box of $\Omega(z')$. Let us call (v_2, w_2) the coordinate of $\Omega(z')$.

We also need the following lemma:

Lemma 5.9. Let us fix a cross-section $\Sigma(\bar{x})$. For $\Omega(y_1), \Omega(y_2) \in \Sigma(\bar{x})$, we say that $\mathcal{T}(y_1)$ and $\mathcal{T}(y_2)$ intersect with parameter r if there exists r_2 such that $u(r)y_1$ and $u(r_2)y_2$ are contained in the same $\Theta(z)$. Note that r is not unique. But any two parameters r, r' satisfy $|r - r'| \leq \beta |r|$.

Let $\Theta(z)$ and $\Omega(y)$ be in $\Sigma(\bar{x})$. Suppose there exists $z' \in \Theta(z)$ such that

$$z' = \exp(\mathbf{w}^+)u(r_y)y,$$

where $\mathbf{w}^+ = v_1 \mathbf{v}_1 + w_1 \mathbf{w}_1$. Let (X_0, Y_0) denote the coordinate of $\Omega(z')$. Then the coordinate box of all $\Omega(\bar{z}) \in \Theta(z)$ such that $\mathcal{T}(\bar{z})$ and $\mathcal{T}(y)$ intersect with parameter $|r| \ge \beta^2 e^{2s'}$ is contained in a $\beta^{-2} e^{-3s'}$ neighborhood of a hyperbola

$$(X - X_0)(Y - Y_0) + v_1^{-1}(X - X_0) + w_1^{-1}(Y - Y_0) = 0,$$

or a straight line

$$v_1^{-1}(X - X_0) + w_1^{-1}(Y - Y_0) = 0,$$

 $in \ [\beta]^2 \subset \mathbb{R}^2.$

Proof. Without loss of generality, let us assume $|w_1| \ge |v_1|$. The opposite case can be handled similarly.

Suppose $z'' = \exp(\mathbf{w}^-)z'$ (where $\mathbf{w}^- = v_2\mathbf{v}_2 + w_2\mathbf{w}_2$) satisfies that $\mathcal{T}(z'') \cap \mathcal{T}(y) \neq \emptyset$. Then by (5.13), (5.14), and (5.15), we have that

$$u(r)z'' = a_0(r)a(r)u^*(f^*(r)) \exp(\mathbf{f}_{-}(r)) \exp(\mathbf{f}_{+}(r))u(f(r))y,$$

where $\mathbf{f}_{+}(r) = f_v(r)\mathbf{v}_1 + f_w(r)\mathbf{w}_1$,

$$f_v(r) = \frac{2rv_2}{2 + rv_2w_2} + v_1,$$

and

$$f_w(r) = \frac{-2rw_2}{2 - rv_2w_2} + w_1.$$

Suppose for some $|r| \ge \beta^2 e^{2s'}$, we have $|f_v(r)|, |f_w(r)| \le e^{-s'}$. Let us denote $\Delta = v_2 w_2/2$. Then we have

$$r^{-1} + \Delta = -\frac{v_2}{v_1} (1 + O(e^{-s'} v_1^{-1})), \tag{5.17}$$

and

$$r^{-1} - \Delta = \frac{w_2}{w_1} (1 + O(e^{-s'} w_1^{-1})). \tag{5.18}$$

Note that $|r^{-1}| \ge e^{-2s'} \ge \beta^2 e^{-2s'} \ge |\Delta|$. Since $|r| \ge \beta^2 e^{2s'}$, we get

$$|v_2/v_1|, |w_2/w_1| \le \beta^{-2} e^{-2s'}.$$
 (5.19)

Moreover, (5.18) and (5.17) imply that

$$v_2w_2 + v_1^{-1}v_2 + w_1^{-1}w_2 = O(\beta^{-2}e^{-3s'}v_1^{-1}) + O(\beta^{-2}e^{-3s'}w_1^{-1}).$$

This implies that (v_2, w_2) is contained in $\beta^{-2}e^{-3s'}$ neighborhood of the hyperbola

$$(X - X_0)(Y - Y_0) + v_1^{-1}(X - X_0) + w_1^{-1}(Y - Y_0) = 0.$$

Also note that when $|v_1| \leq \beta^{-4} e^{-s'} |w_1|$, the hyperbola is $\beta^{-2} e^{-3s'}$ close to the straight line

$$v_1^{-1}(X - X_0) + w_1^{-1}(Y - Y_0) = 0.$$

In this case we can consider it as $\beta^{-2}e^{-3s'}$ neighborhood of the straight line. \square

Remark 5.10. In fact, by (5.19), we have that if $||(v_1, w_1)|| \leq e^{s'-\ell}$, then

$$||(v_2, w_2)|| \le \beta^{-2} e^{-s'-\ell}.$$

This implies that the curve can be restricted to $(X_0, Y_0) + [\beta^{-2}e^{-s'-\ell}]^2$.

Let us denote the neighborhood of the curve by $\mathcal{C}(\mathcal{T}(y), z) \subset [\beta e^{-s'}]^2$. By $\Omega(\bar{z}) \in \mathcal{C}(\mathcal{T}(y), z)$, we mean the coordinate box of $\Omega(\bar{z})$ is contained in $\mathcal{C}(\mathcal{T}(y), z)$. It is easy to see that there exists a surface, whose $\beta^{-2}e^{-s'}$ -neighborhood is denoted by $\mathcal{S}(\mathcal{T}(y), z) \in \mathbb{R}^3$ such that

$$\mathcal{T}(\bar{z}) \subset \mathcal{S}(\mathcal{T}(y), z).$$

Let us introduce the following structure-randomness decomposition of a measure μ :

Definition 5.11. Let s' > 0 and μ be a probability measure on X defined by taking the normalized measure on a subset consisting of U-orbits of length $e^{2s'}$. For a fixed playground $\Pi(x) = \Pi_{s'}x$, let us consider its corresponding Kakeya-type model. To fit our case, let us assume that the weight of $\mathcal{T}(y)$ satisfies that $\mathcal{W}(y) \leq e^{-2d_i s'} e^{-2s'}$ for every tube $\mathcal{T}(y)$. Then every neighborhood $\Theta(z) = \Theta_{s'}z \subset \Pi(x)$ corresponds to a collection of tubes passing through $\mathbf{p}_z = (r_z, v_{1,z}, w_{1,z})$. By Lemma 5.9, for every $\Omega(y)$, there is a corresponding curve

$$\mathcal{C}(\mathcal{T}(y), z) \subset [\beta]^2 \subset \mathbb{R}^2$$
.

It is easy to see that the sum of weights of tubes in $\mathcal{C}(\mathcal{T}(y), z)$ is at most

$$\beta^{-2}e^{2s'}e^{-2d_is'}e^{-2s'} = \beta^{-2}e^{-2d_is'}.$$

A curve C(T(y), z) is called highly concentrated if the sum of weights of tubes in C(T(y), z) is at least $e^{-7C_1\varepsilon_1 s'}e^{-2d_i s'}$. Let us take all U-orbits corresponding to tubes contained in highly concentrated curves and define a measure $\mu_{\mathcal{P}}$. Then μ admits the following decomposition:

$$\mu = \nu_{\mathfrak{r}} \mu_{\mathfrak{r}} + \nu_{\mathcal{P}} \mu_{\mathcal{P}}.$$

Let us call μ_{τ} the random component of μ and $\mu_{\mathcal{P}}$ the structured component of μ .

Recall that starting from $\mathcal{F}_0 = \mathcal{F}$, we want to construct a sequence $\{(\mathcal{F}_i, s_i)\}$ such that \mathcal{F}_{i+1} is a subset of $a(s_i)\mathcal{F}_i$ with a better dimension control.

The following proposition is the goal of this subsection:

Proposition 5.12. Suppose that \mathcal{F}_i is (d_i, s_i) -good. Then $\mu = a(s_i)_* \mu_{\mathcal{F}_i}$ admits a decomposition:

$$\mu = \nu_{\mathcal{R}_i} \mu_{\mathcal{R}_i} + \nu_{i+1} \mu_{i+1} + \nu_{\mathcal{P}} \mu_{\mathcal{P}},$$

such that $\nu_{\mathcal{R}_i} = O(\beta)$, $\mu_{\mathcal{P}}$ is the structured component of μ as in Definition 5.11, and μ_{i+1} is $(d_i + \varepsilon_1, s_{i+1})$ -good. Recall that $s_{i+1} = s_i/2$.

Proof. Let us denote $s' = s_{i+1}$. We claim that our hypotheses on \mathcal{F}_i implies that for any $\Omega(y) = \Omega_{s_i} y$, we have $\mu(\Omega(y)) \leq e^{-2s'} e^{-2d_i s'}$. In fact,

$$\mu(\Omega(y)) = \mu_{\mathcal{F}_i}(a(-2s')\Omega(y)).$$

Note that

$$a(-2s')\Omega(y) = Q_{2s'}B^{A_0}(\beta)R^H(2s')y'$$

where y' = a(-2s')y. Noting that $\mu_{\mathcal{F}_i}$ is a normalized measure on *U*-orbits, we have

$$\mu_{\mathcal{F}_i}(Q_{2s'}B^{A_0}(\beta)R^H(2s')y') \leqslant e^{-2s'}\mu_{\mathcal{F}_i}(Q_{2s'}B^{A_0H}(\beta)y').$$

By our hypothesis, we have

$$\mu_{\mathcal{F}_i}(Q_{2s'}B^{A_0H}(\beta)y') \leqslant e^{-2d_i s'},$$

which implies the claim.

Let us write

$$\mu = \nu_{\mathfrak{r}} \mu_{\mathfrak{r}} + \nu_{\mathcal{P}} \mu_{\mathcal{P}},$$

as in Definition 5.11.

For $\nu_{\mathfrak{r}}\mu_{\mathfrak{r}}$, we have two cases:

Case 1. $\nu_{\mathfrak{r}} < \beta$;

Case 2. $\nu_{\rm r} \geqslant \beta$.

For Case 1, we can put

$$\mu_{\mathcal{R}_i} = \mu_{\mathfrak{r}} \text{ and } \mu_{i+1} = 0,$$

and we are done.

Now let us handle **Case 2**, namely, $\nu_{\tau} \geqslant \beta$. In this case, by the definition of μ_{τ} , we have the corresponding Kakeya-type model does not have any highly concentrated curves.

For every neighborhood $\Theta(z) = \Theta_{s'}z$, we call it a small neighborhood if

$$\mu_{\mathfrak{r}}(\Theta(z)) \leqslant e^{-2s'} e^{-(d_i + \varepsilon_1)s'}.$$

Otherwise we call it a big neighborhood.

Let $\bar{\mathcal{R}}_i$ denote the union of all orbits contained in big neighborhoods. Then it suffices to show that $\mu_r(\bar{\mathcal{R}}_i) < \beta$.

For integer $j \ge 1$, let us denote

$$\Xi_i(z') := \Xi_{s',i} z' \subset \Theta(z),$$

where

$$\Xi_{s',j} := Q_{s'+j}^{s'} Q^H(s').$$

Given a big neighborhood $\Theta(z)$, we call $\Theta(z)$ bad if there exists $\Xi_{C_1\varepsilon_1s'}(z') \subset \Theta(z)$ such that

$$\mu_{\mathfrak{r}}(\Xi_{C_1\varepsilon_1s'}(z')) \geqslant (1-\beta)\mu_{\mathfrak{r}}(\Theta(z)).$$

Otherwise we call it a good neighborhood. Let \mathcal{R}_i denote the union of good (and big) neighborhoods.

Let us first prove $\mu_{\mathfrak{r}}(\mathcal{R}_i) < \beta$.

For a contradiction, let us assume $\mu_{\mathfrak{r}}(\mathcal{R}_i) \geqslant \beta$.

For every $y \in \mathcal{R}_i$, if

$$|u([e^{2s'}])y \cap \mathcal{R}_i| \geqslant \beta^2 e^{2s'},$$

we will call it a good point. Otherwise we call it a bad point. Then it is easy to see that the collection of bad points has $\mu_{\rm r}$ -measure $\leq \beta^2$. Therefore, the collection of good points, denoted by \mathcal{R}'_i , has $\mu_{\rm r}$ -measure $\geq \beta$.

Now let us cut $\Pi(x)$ into pieces each of which is a cross-section with respect to the U-direction of the form

$$\Sigma(\bar{x}) := \Sigma_{s'}\bar{x}$$

for some $\bar{x} \in u([e^{2s'}])x$. We call $\Sigma(\bar{x})$ good if

$$\mu_{\mathfrak{r}}(\mathcal{R}'_i \cap \Sigma(\bar{x})) \geqslant \beta^2 \mu_{\mathfrak{r}}(\Sigma(\bar{x})).$$

Otherwise we called it bad.

Now let us fix a good cross-section $\Sigma(\bar{x})$. Let us consider its corresponding Kakeya-type model and calculate the following weighted intersection number:

We take the sum of $W(y_1)W(y_2)$ for all pairs $\mathcal{T}(y_1)$ and $\mathcal{T}(y_2)$ intersecting each other with parameter $|r| \ge \beta^2 e^{2s'}$ and $y_1 \in \mathcal{R}_i$ or $y_2 \in \mathcal{R}_i$. Let us denote it by $\mathcal{S}(\Sigma(\bar{x}))$.

Now for a fixed $\mathcal{T}(y)$ with $y \in \mathcal{R}'_i$, we have that

$$|u([e^{2s'}])y \cap \mathcal{R}'_i| \geqslant \beta^2 e^{2s'}.$$

For every $\tilde{y} \in u([e^{2s'}])y \cap \mathcal{R}'_i$, we have

$$\mu_{\mathfrak{r}}(\Theta(\tilde{y})) \geqslant e^{-2s'} e^{-(d_i + \varepsilon_1)s'}.$$

By Lemma 5.7, every $\Omega(\tilde{z}) \subset \Theta(\tilde{y})$ corresponds to some $\Omega(z)$ from $\Sigma(\bar{x})$ with $\mathcal{T}(z)$ interesecting $\mathcal{T}(y)$.

Let us only consider $\Omega(z)$ such that $\Omega(\tilde{z}) \subset \Theta(\tilde{y}) \setminus \Xi_{C_1 \varepsilon_1 s'}(\tilde{y})$, or equivalently, $\Omega(z)$ such that

$$\|\Delta(y,z)\| \geqslant e^{-(1+C_1\varepsilon_1)s'}. (5.20)$$

Noting that $\mu_{\mathfrak{r}}(\Theta(\tilde{y}) \setminus \Xi_{C_1\varepsilon_1s'}(\tilde{y})) \geqslant \beta\mu_{\mathfrak{r}}(\Theta(\tilde{y}))$ by the definition of \mathcal{R}_i , by running over all $\Omega(\tilde{z}) \subset \Theta(\tilde{y}) \setminus \Xi_{C_1\varepsilon_1s'}(\tilde{y})$, we have that the contribution from $\Theta(\tilde{y})$ to $\mathcal{S}(\Sigma(\bar{x}))$ is at least

$$\beta \mathcal{W}(y) \mu_{\mathfrak{r}}(\Theta(\tilde{y})) \geqslant \beta \mathcal{W}(y) e^{-2s'} e^{-(d_i + \varepsilon_1)s'}.$$

By (5.20) and Lemma 5.7, each such intersection can cover at most $e^{C_1\varepsilon_1s'}$ of the whole $u([e^{2s'}])y$ orbit. Therefore, from the orbit we can find at least $\beta^2 e^{(2-C_1\varepsilon_1)s'} \ge e^{(2-2C_1\varepsilon_1)s'}$ different \tilde{y} 's. Therefore, we have that the contribution from the whole orbit $u([e^{2s'}])y$ to $\mathcal{S}(\Sigma(\bar{x}))$ is at least

$$\beta \mathcal{W}(y)e^{-2s'}e^{-(d_i+\varepsilon_1)s'}e^{(2-2C_1\varepsilon_1)s'} = \mathcal{W}(y)e^{-(d_i+4C_1\varepsilon_1)s'}.$$

By running over all $\Omega(y) \in \mathcal{R}'_i \cap \Sigma(\bar{x})$, we get

$$S(\Sigma(\bar{x})) \geqslant \tilde{\mathfrak{m}}e^{-4C_1\varepsilon_1s'}e^{-d_is'}$$
.

where $\tilde{\mathfrak{m}} = \mu_{\mathfrak{r}}(\Sigma(\bar{x})).$

On the other hand, let us get an upper bound on $S(\Sigma(\bar{x}))$.

For any $\mathcal{T}(y)$ and $\Theta(z)$ with $z \in \mathcal{R}_i$, let us consider all $\Omega(z') \subset \Theta(z)$ such that the corresponding tubes intersect with parameter $|r| \ge \beta^2 e^{2s'}$. Note that $\Omega(z') \subset \mathcal{C}(\mathcal{T}(y), z)$. Since μ_{τ} does not have any highly concentrated curves, we have that the contribution to $\mathcal{S}(\Sigma(\bar{x}))$ from $\mathcal{T}(y)$ and all such $\mathcal{T}(z')$ is

$$\leqslant \mathcal{W}(y)e^{(2-7C_1\varepsilon_1)s'}e^{-2d_is'}e^{-2s'} = \mathcal{W}(y)e^{-7C_1\varepsilon_1s'}e^{-2d_is'}.$$

Since $z \in \mathcal{R}_i$, the μ_{r} -measure of each $\Theta(z)$ is at least $e^{-(d_i+\varepsilon_1)s'}e^{-2s'}$ and the measure of $\Sigma(\bar{x})$ is at most $e^{-2s'}$, we have that there are at most $e^{d_is'}e^{\varepsilon_1s'}$ different $\Theta(z)$ from $\Sigma(\bar{x})$. Thus the total contribution to $\mathcal{S}(\Sigma(\bar{x}))$ with a fixed $\mathcal{T}(y)$ and all possible $\Theta(z)$ is at most

$$\mathcal{W}(u)e^{-7C_1\varepsilon_1 s'}e^{-2d_i s'}e^{d_i s'}e^{\varepsilon_1 s'} = \mathcal{W}(u)e^{-6C_1\varepsilon_1 s'}e^{-d_i s'}.$$

By running over all W(y), we get

$$S(\Sigma(\bar{x})) \leqslant \tilde{\mathfrak{m}}e^{-6C_1\varepsilon_1 s'}e^{-d_i s'}.$$

This implies that

$$\tilde{\mathfrak{m}}e^{-4C_1\varepsilon_1s'}e^{-d_is'} \leqslant \tilde{\mathfrak{m}}e^{-6C_1\varepsilon_1s'}e^{-d_is'}.$$

which leads to a contradiction. This proves that $\mu_{\mathfrak{r}}(\mathcal{R}_i) < \beta$.

We then prove that $\mu_{\mathfrak{r}}(\bar{\mathcal{R}}_i \setminus \mathcal{R}_i) < \beta^2$.

Let us divide $\bar{\mathcal{R}}_i \setminus \mathcal{R}_i$ into

$$ar{\mathcal{R}}_i \setminus \mathcal{R}_i = igcup_{j=C_1arepsilon_1 s}^{3s} \mathcal{R}_{i,j},$$

where $\mathcal{R}_{i,j}$ denotes the union of big $\Theta(z)$'s such that there exists $\Xi_j(z') \subset \Theta(z)$ satisfying

$$\mu_{\mathfrak{r}}(\Xi_j(z')) \geqslant (1-\beta)^j \mu_{\mathfrak{r}}(\Theta(z)),$$

and

$$\mu_{\mathfrak{r}}(\Xi_{j+1}(z'')) < (1-\beta)\mu_{\mathfrak{r}}(\Xi_{j}(z'))$$

for any $\Xi_{j+1}(z'') \subset \Xi_j(z')$.

Then it suffices to show that $\mu_{\mathfrak{r}}(\mathcal{R}_{i,j}) < \beta^3$ for all $j \geqslant 1$.

Let $\mathcal{R}'_{i,j}$ denote the union of $\Xi_j(z)$ with

$$\mu_{\mathfrak{r}}(\Xi_j(z)) \geqslant (1-\beta)^j e^{-2s'} e^{-(d_i+\varepsilon_1)s'}$$

and

$$\mu_{\mathfrak{r}}(\Xi_{j+1}(z')) < (1-\beta)\mu_{\mathfrak{r}}(\Xi_{j}(z))$$

for any $\Xi_{j+1}(z') \subset \Xi_j(z)$. To prove $\mu_{\mathfrak{r}}(\mathcal{R}_{i,j}) < \beta^3$, it suffices to show that $\mu_{\mathfrak{r}}(\mathcal{R}'_{i,j}) < \beta^3$.

For a contradiction, let us assume that $\mu_{\mathfrak{r}}(\mathcal{R}'_{i,j}) \geqslant \beta^3$. Using similar arguments as above, we can get a subset $\mathcal{R}''_{i,j} \subset \mathcal{R}'_{i,j}$ such that $\mu_{\mathfrak{r}}(\mathcal{R}''_{i,j}) \geqslant \beta^4$ and for any $y \in \mathcal{R}''_{i,j}$,

$$|u([e^{2s'}])y \cap \mathcal{R}_{i,j}''| \geqslant \beta^4 e^{2s'}.$$

Let us choose a cross-section $\Sigma(\bar{x})$ such that

$$\mu_{\mathfrak{r}}(\Sigma(\bar{x}) \cap \mathcal{R}''_{i,j}) \geqslant \beta^5 \mu_{\mathfrak{r}}(\Sigma(\bar{x})).$$

Now let us consider the sum $S_j(\Sigma(\bar{x}))$ of weighted intersection number $W(y_1)W(y_2)$ where $\Omega(y_1), \Omega(y_2) \subset \Sigma(\bar{x})$ satisfies $\mathcal{T}(y_1)$ and $\mathcal{T}(y_2)$ intersect with parameter $|r| \geq \beta^2 e^{2s'}$, one of $\Omega(y_1), \Omega(y_2)$ is contained in $\mathcal{R}'_{i,j}$, and

$$\|\Delta(y_1, y_2)\| \in [e^{-s'-j}, 2e^{-s'-j}].$$

Let us first get the lower bound on $S_j(\Sigma(\bar{x}))$. For any fixed $\Omega(y) \subset \mathcal{R}''_{i,j} \cap \Sigma(\bar{x})$, we have that

$$|u([e^{2s'}])y \cap \mathcal{R}_{i,j}''| \geqslant \beta^4 e^{2s'}$$
.

For any $y' \in u([e^{2s'}])y \cap \mathcal{R}''_{i,j}$, we have that

$$\mu_{\mathfrak{r}}(\Xi_j(y')) \geqslant (1-\beta)^j e^{-2s'} e^{-(d_i+\varepsilon_1)s'}$$

and

$$\mu_{\mathfrak{r}}(\Xi_{j+1}(y')) < (1-\beta)\mu_{\mathfrak{r}}(\Xi_{j}(y')).$$

It is easy to see that every $\Omega(y_2') \subset \Xi_j(y') \setminus \Xi_{j+1}(y')$ corresponds to some $\Omega(y_2) \subset \Sigma(\bar{x})$ with $\mathcal{T}(y) \cap \mathcal{T}(y_2) \neq \emptyset$ and

$$\|\Delta(y, y_2)\| \in [e^{-s'-j}, 2e^{-s'-j}].$$

This will contribute at least

$$\mathcal{W}(y)\mu_{\mathfrak{r}}(\Xi_{j}(y')\setminus\Xi_{j+1}(y')) > \mathcal{W}(y)\beta(1-\beta)^{j}e^{-2s'}e^{-(d_{i}+\varepsilon_{1})s'} > \mathcal{W}(y)e^{-2s'}e^{-(d_{i}+2\varepsilon_{1})s'}$$

to $S_j(\Sigma(\bar{x}))$. Since $\|\Delta(y, y_2)\| \geqslant e^{-s'-j}$, each intersection can cover at most e^j of the orbit $u([e^{2s'}])y$. Therefore, there are at least $\beta^4 e^{2s'-j}$ different $\Xi_j(y')$'s. This implies that for this fixed $\Omega(y)$, the contribution to $S_j(\Sigma(\bar{x}))$ is at least

$$\beta^4 e^{2s'-j} \mathcal{W}(y) e^{-2s'} e^{-(d_i+2\varepsilon_1)s'} > \mathcal{W}(y) e^{-(d_i+3\varepsilon_1)s'-j}.$$

This implies that

$$S_j(\Sigma(\bar{x})) \geqslant \mu_{\mathbf{r}}(\mathcal{R}''_{i,j} \cap \Sigma(\bar{x}))e^{-(d_i+3\varepsilon_1)s'-j} \geqslant \beta^5 \tilde{\mathbf{m}} e^{-(d_i+3\varepsilon_1)s'-j},$$

where $\tilde{\mathfrak{m}} = \mu_{\mathfrak{r}}(\Sigma(\bar{x})).$

Now let us get an upper bound on $S_i(\Sigma(\bar{x}))$.

For a fixed $\Omega(y_2) \in \Sigma(\bar{x})$ and $\Xi_j(z) \subset \Sigma(\bar{x}) \cap \mathcal{R}'_{i,j}$, let us consider all $\Omega(y_1) \subset \Xi_j(z)$ such that $\mathcal{T}(y_1)$ and $\mathcal{T}(y_2)$ intersect with parameter $|r| \geq \beta^2 e^{2s'}$, and

$$\|\Delta(y_1, y_2)\| \in [e^{-s'-j}, 2e^{-s'-j}].$$

It is easy to see that if $\Xi_j(z)$ is not contained in $\Sigma_j(y_2)$, then $\Xi_j(z)$ does not contain any $\Omega(y_1)$ satisfying the conditions. For $\Xi_j(z) \subset \Sigma_j(y_2)$, such $\Omega(y_1)$'s are contained in $\mathcal{C}(\mathcal{T}(y_2), z)$ as in Definition 5.11. The sum of those $\mathcal{W}(y_1)$'s is at most

$$e^{2s'-j}e^{-2s'}e^{-2d_is'} = e^{-j}e^{-2d_is'}.$$

By Lemma 5.5, we have

$$\mu_{\mathfrak{r}}(\Sigma_j(y_2)) \leqslant e^{-2s'} e^{-d_0 j}.$$

Noting that $\mu_{\mathfrak{r}}(\Xi_j(z)) \geqslant (1-\beta)^j e^{-2s'} e^{-(d_i+\varepsilon_1)s'}$, there are at most

$$\frac{e^{-2s'}e^{-d_0j}}{(1-\beta)^j e^{-2s'}e^{-(d_i+\varepsilon_1)s'}} = (1-\beta)^{-j}e^{-d_0j}e^{(d_i+\varepsilon_1)s'}$$

different $\Xi_j(z)$ contained in $\Sigma_j(y_2)$. Thus, for the fixed $\Omega(y_2)$, the contribution to $S_j(\Sigma(\bar{x}))$ is at most

$$\mathcal{W}(y_2)(1-\beta)^{-j}e^{-d_0j}e^{(d_i+\varepsilon_1)s'}e^{-j}e^{-2d_is'} \leqslant 2\mathcal{W}(y_2)e^{-d_0j}e^{(d_i+\varepsilon_1)s'}e^{-j}e^{-2d_is'},$$

since $(1-\beta)^{-j} \leq 2$. Therefore, we get

$$S_j(\Sigma(\bar{x})) \leqslant 2\tilde{\mathfrak{m}}e^{-d_0j}e^{(d_i+\varepsilon_1)s'}e^{-j}e^{-2d_is'} = 2\tilde{\mathfrak{m}}e^{-d_0j}e^{-(d_i-\varepsilon_1)s'-j}$$

This upper bound is larger than the lower bound because $d_0j \ge d_0C_1\varepsilon_1s' \ge 10\varepsilon_1s'$. This leads to a contradiction.

This completes the proof.

6. The structured component

This section is devoted to the study of the structured component $\mu_{\mathcal{P}}$ of μ . First note that by repeating the argument in the proof of Proposition 5.12, we can prove the following statement for $\mu_{\mathcal{P}}$:

Proposition 6.1. Suppose $\mu_{\mathcal{F}_i}$ is (d, s_i) -good. Let $\mu = a(s_i)_*\mu_{\mathcal{F}_i}$, $\mu_{\mathfrak{r}}$, and $\mu_{\mathcal{P}}$ be as above. Then after removing an exponentially small proportion, $\mu_{\mathcal{P}}$ is $(d - C_1\varepsilon_1, s_{i+1})$ -good.

Let us decompose $\mu_{\mathcal{F}_i}$ as

$$\mu_{\mathcal{F}_i} = \mu_{\mathcal{F}_i,1} + \mu_{\mathcal{F}_i,2}$$

where $\mu_{\mathcal{F}_{i},1}$ is $(d_{i} + (C_{1} + 1)\varepsilon_{1}, s_{i})$ -good and $\mu_{\mathcal{F}_{i},2}$ satisfies that for any $x \in \text{supp}\mu_{\mathcal{F}_{i},2}$,

$$\mu_{\mathcal{F}_{i,2}}(Q_{s_i}B^{A_0H}(\beta)x) \geqslant e^{-(d_i + (C_1 + 1)\varepsilon_1)s_i}.$$
(6.1)

By Proposition 6.1, we have that $a(s_i)_*\mu_{\mathcal{F}_i,1}$ is $(d_i + \varepsilon_1, s_{i+1})$ -good. Therefore, it suffices to show that $a(s_i)_*\mu_{\mathcal{F}_i,2}$ is $(d_i + \varepsilon_1, s_{i+1})$ -good. Later in this paper, we can assume that $\mu_{\mathcal{F}_i} = \mu_{\mathcal{F}_i,2}$, namely, for any $x \in \text{supp}\mu_{\mathcal{F}_i}$,

$$\mu_{\mathcal{F}_i}(Q_{s_i}B^{A_0H}(\beta)x) \geqslant e^{-(d_i + (C_1 + 1)\varepsilon_1)s_i}.$$
(6.2)

Under this assumption, we have that for any $x \in \text{supp}\mu$,

$$\mu(\Omega(x)) \geqslant e^{-2s'} e^{-2(d_i + (C_1 + 1)\varepsilon_1)s'}.$$
 (6.3)

By Proposition 6.1, there exists $\mathfrak{E} \subset X$ with exponentially small measure such that for $x \in X \setminus \mathfrak{E}$,

$$\mu_{\mathcal{P}}(\Theta(x)) \leqslant e^{-2s'} e^{-(d_i - C_1 \varepsilon_1)s'}. \tag{6.4}$$

Thus, if $d_i < 2 - 7C_1\varepsilon_1$, then we will not have any highly concentrated surfaces. Therefore, later in this paper, we can assume that

$$d_i \geqslant 2 - 7C_1 \varepsilon_1. \tag{6.5}$$

We will prove the following statement:

Proposition 6.2. For $2 - 7C_1\varepsilon_1 \le d_i \le 4 - 10^{11}C_1\varepsilon_1$, we have $\mu_{\mathcal{P}}$ is $(d_i + \varepsilon_1, s_{i+1})$ -good.

We first define the following property:

Definition 6.3. Given $\Omega(y)$ and $\Theta(z)$, a surface $\mathcal{S} = \mathcal{S}(\mathcal{T}(y), z)$ is called integrable if the corresponding curve $\mathcal{C} = \mathcal{C}(\mathcal{T}(y), z)$ is highly concentrated and there are at least $e^{(4-120C_1\varepsilon_1)s'}$ different $\Omega(y')$ such that $\mathcal{T}(y')$ is contained in \mathcal{S} .

We need the following lemma on integrable surfaces.

Lemma 6.4. Suppose a highly concentrated curve C = C(T(y), z) has equation

$$(X - X_0)(Y - Y_0) + v_1^{-1}(X - X_0) + w_1^{-1}(Y - Y_0) = 0,$$

with $||(v_1, w_1)|| \ge e^{s'-7C_1\varepsilon_1 s'}$. Then the corresponding surface $\mathcal{S} = \mathcal{S}(\mathcal{T}(y), z)$ is integrable only if

$$|v_1| \leqslant e^{120C_1\varepsilon_1 s'},$$

or

$$|w_1| \leqslant e^{120C_1\varepsilon_1 s'}.$$

Proof. By Lemma 5.9, for $|v_1| \leq e^{-7C_1\varepsilon_1 s'}$ or $|w_1| \leq e^{-7C_1\varepsilon_1 s'}$, \mathcal{C} is a straight line

$$v_1^{-1}(X - X_0) + w_1^{-1}(Y - Y_0) = 0.$$

It is easy to see that this case C can be contained in an integrable surface.

Now let us assume that $|v_1|, |w_1| \ge e^{-7C_1\varepsilon_1 s'}$.

Let us fix $\Omega(z_0) \subset \mathcal{C}$ and consider all $\Omega(y')$ such that

$$C = C(T(y'), z)$$

and

$$z_0 = \exp(v_{1,y'}\mathbf{v}_1 + w_{1,y'}\mathbf{w}_1)u(r_{y'})y'.$$

Let us denote by $\mathcal{M}(\mathcal{C}, z_0)$ the collection of those $\Omega(y')$'s. Let us fix $\Omega(y_0) \in \mathcal{M}(\mathcal{C}, z_0)$ and write

$$z_0 = \exp(v_1 \mathbf{v}_1 + w_1 \mathbf{w}_1) u(r) y_0.$$

Then \mathcal{C} has equation

$$(X - X_0)(Y - Y_0) + v_1^{-1}(X - X_0) + w_1^{-1}(Y - Y_0) = 0,$$

where (X_0, Y_0) is the coordinate of $\Omega(z_0)$. Without loss of generality, let us assume that $(X_0, Y_0) = \mathbf{0}$. Then the equation reads

$$XY + v_1^{-1}X + w_1^{-1}Y = 0.$$

Then for any $\Omega(y') \in \mathcal{M}(\mathcal{C}, z_0)$ such that

$$z_0 = \exp(v_{1,y'}\mathbf{v}_1 + w_{1,y'}\mathbf{w}_1)u(r_{y'})y',$$

we get the equation of \mathcal{C} also reads

$$XY + v_{1,y'}^{-1}X + w_{1,y'}^{-1}Y = 0.$$

In other words, this curve is contained in $\beta^{-2}e^{-3s'}$ neighborhood of the curve

$$XY + v_1^{-1}X + w_1^{-1}Y = 0.$$

Let us assume that $|w_1| \ge |v_1|$. The opposite case can be handled similarly.

Let us denote

$$k = \frac{w_1}{v_1}$$
, and $k' = \frac{w_{1,y'}}{v_{1,y'}}$.

Then the curves can be written as

$$X = \frac{-Y}{w_1 Y + k}$$
, and $X = \frac{-Y}{w_{1,y'} Y + k'}$.

Then we have for all $Y \in [e^{-s'-7C_1\varepsilon_1 s'}]$,

$$\left| \frac{Y}{w_1 Y + k} - \frac{Y}{w_{1,y'} Y + k'} \right| \le \beta^{-2} e^{-3s'}.$$

This implies that

$$|Y| \cdot |w_1 - w_{1,y'}| \frac{\left|Y + \frac{k' - k}{w_{1,y'} - w_1}\right|}{|w_1 Y + k| \cdot |w_{1,y'} Y + k'|} \le \beta^{-2} e^{-3s'}.$$

Let us denote

$$|k| \in [e^{\ell}, 2e^{\ell}].$$

Then it is easy to show that

$$|k'| \in [e^{\ell}/2, 4e^{\ell}].$$

We have the following two cases:

CASE 1: $|k - k'| < e^{-s'} |w_1 - w_{1,y'}|$;

CASE 2:
$$|k - k'| \ge e^{-s'} |w_1 - w_{1,y'}|$$
.

Let us first consider CASE 1. In this case we have

$$|w_1 - w_{1,y'}|e^{-2s'-14C_1\varepsilon_1 s'-2\ell} \leqslant \beta^{-2}e^{-3s'},$$

namely,

$$|w_1 - w_{1,y'}| \le \beta^{-2} e^{-s'} e^{14C_1 \varepsilon_1 s'} e^{2\ell}$$

Then

$$|k - k'| < e^{-s'} |w_1 - w_{1,y'}| \le \beta^{-2} e^{-2s'} e^{14C_1 \varepsilon_1 s'} e^{2\ell}.$$

Then there are at most $e^{60C_1\varepsilon_1 s'}e^{2\ell}$ different $\Omega(y')$ contained in $\mathcal{M}(\mathcal{C}, z_0)$. If $|v_1| > e^{120C_1\varepsilon_1 s'}$, then

$$e^{\ell} \leqslant |k| < e^{s'-120C_1\varepsilon_1 s'},$$

which implies that there are at most $e^{2s'-120C_1\varepsilon_1s'}$ different $\Omega(y')$ contained in $\mathcal{M}(\mathcal{C}, z_0)$. By running over all $\Omega(z_0) \subset \mathcal{C}$, we have that there are at most

$$e^{4s'-120C_1\varepsilon_1s'}$$

different $\Omega(y')$ contained in $\mathcal{S}(\mathcal{T}(y), z)$. This completes the proof for **CASE** 1.

Now let us consider CASE 2. In this case we have

$$|k - k'|e^{-s' - 7C_1\varepsilon_1 s' - 2\ell} \le \beta^{-2}e^{-3s'},$$

namely,

$$|k - k'| \le \beta^{-2} e^{-2s'} e^{7C_1 \varepsilon_1 s'} e^{2\ell}$$
.

Then

$$|w_1 - w_{1,u'}| \le e^{s'} |k - k'| \le \beta^{-2} e^{-s'} e^{7C_1 \varepsilon_1 s'} e^{2\ell}$$

Then there are at most $e^{60C_1\varepsilon_1 s'}e^{2\ell}$ different $\Omega(y')$ contained in $\mathcal{M}(\mathcal{C}, z_0)$. If $|v_1| > e^{120C_1\varepsilon_1 s'}$, then

$$e^{\ell} \leqslant |k| < e^{s'-120C_1\varepsilon_1 s'},$$

which implies that there are at most $e^{2s'-120C_1\varepsilon_1s'}$ different $\Omega(y')$ contained in $\mathcal{M}(\mathcal{C}, z_0)$. By running over all $\Omega(z_0) \subset \mathcal{C}$, we have that there are at most

$$e^{4s'-120C_1\varepsilon_1s'}$$

different $\Omega(y')$ contained in $\mathcal{S}(\mathcal{T}(y), z)$. This completes the proof for **CASE** 2.

Combining these two cases, we complete the proof.

Remark 6.5. In fact, we show the following: If a highly concentrated curve

$$C: XY + v_1^{-1}X + w_1^{-1}Y = 0$$

satisfies that

$$|k| = |w_1/v_1| \in [e^{\ell}, 2e^{\ell}],$$

then there are at most $e^{2s'+2\ell+60C_1\varepsilon_1s'}$ different $\Omega(y')$ contained in \mathcal{S} (the surface determined by \mathcal{C}).

Let \mathcal{P}_1 denote the collection of integrable surfaces with $|v_1| \leq e^{120C_1\varepsilon_1 s'}$ and \mathcal{P}_2 denote the collection of integrable surfaces with $|w_1| \leq e^{120C_1\varepsilon_1 s'}$.

 \mathcal{P}_1 corresponds to the subgroup

$$H_1 := \left\{ \begin{bmatrix} 1 & \mathbf{w} \\ & h \end{bmatrix} : \mathbf{w} \in \mathbb{R}^2, h \in \mathrm{SL}(2, \mathbb{R}) \right\} \cong \mathrm{SL}(2, \mathbb{R}) \ltimes \mathbb{R}^2,$$

and \mathcal{P}_2 corresponds to the subgroup

$$H_2 := \left\{ \begin{bmatrix} 1 \\ \mathbf{v} & h \end{bmatrix} : \mathbf{v} \in \mathbb{R}^2, h \in \mathrm{SL}(2, \mathbb{R}) \right\} \cong \mathrm{SL}(2, \mathbb{R}) \ltimes \mathbb{R}^2.$$

For $2 - 7C_1\varepsilon_1 \leq d_i \leq 2 + 800C_1\varepsilon_1$, it is possible that $\mu_{\mathcal{P}}$ is concentrated on an integrable surface. The following statement tells this obstruction is ruled out by Proposition 5.3.

Proposition 6.6. For $2 - 7C_1\varepsilon_1 \leqslant d_i \leqslant 2 + 800C_1\varepsilon_1$, μ_P does not admit an integrable surface.

Proof. For a contradiction, let us assume that there exists an integrable surface S from \mathcal{P}_j for μ . Note that $\mu_{\mathcal{F}_i} = a(-2s')_*\mu$ and $a(-2s')\mathcal{S}$ is contained in a neighborhood of the form $Q(\mathfrak{r}_j,\ell)B^{A_0H}(\beta)\tilde{x}$ where $\ell=s'-120C_1\varepsilon_1s'$ and $\tilde{x}=a(-2s')x$. Then it implies that

$$\mu_{\mathcal{F}_i}(Q(\mathfrak{r}_i,\ell)B^{A_0H}(\beta)\tilde{x}) \geqslant e^{-920C_1\varepsilon_1s'} > e^{-d_0\ell/2}$$

which contradicts to Proposition 5.3.

Now we have equipped to prove Proposition 6.2

Proof of Proposition 6.2. For a contradiction, let us assume that there is a subset $\mathcal{R} \subset \text{supp}\mu_{\mathcal{P}}$ with measure $\geqslant \beta$ such that for each $z \in \mathcal{R}$,

$$\mu_{\mathcal{P}}(\Theta(z)) \geqslant e^{-2s'} e^{-(d_i + \varepsilon_1)s'}.$$

By replacing \mathcal{R} with $\mathcal{R} \setminus \mathfrak{E}$, we can assume that (6.4) holds for any $z \in \mathcal{R}$.

By repeating the argument in the proof of Proposition 5.12, we have that to avoid a contradiction, there must be at least

$$e^{2d_is'}e^{d_is'}e^{-14C_1\varepsilon_1s'}$$

highly concentrated curves, counting multiplicity. Therefore, there exists $\mathcal{R}' \subset \mathcal{R}$ with measure $\geqslant \beta^2$, such that for any $\Theta(z) \subset \mathcal{R}'$, counting multiplicity, there are at least $e^{2d_is'}e^{-15C_1\varepsilon_1s'}$ highly concentrated curves contained in $\Theta(z)$.

First note that it is enough to show that those highly concentrated curves can cover at least $e^{d_i s' + 20C_1 \varepsilon_1 s'}$ different $\Omega(z')$. In fact, if this holds, then by (6.3), we have that

$$\mu_{\mathcal{P}}(\Theta(z)) \geqslant e^{-2s'} e^{-(d_i - 16C_1 \varepsilon_1)s'},$$

which contradicts to (6.4).

Now we aim to show that these curves cover at least $e^{d_i s' + 20C_1 \varepsilon_1 s'}$ different $\Omega(z')$.

We first consider the case where most of the curves are contained in integrable surfaces. For $2 - 7C_1\varepsilon_1 \le d_i \le 2 + 800C_1\varepsilon_1$, Proposition 6.6 above tells that there is no integrable surfaces. So, let us assume $d_i > 2 + 800C_1\varepsilon_1$.

Then there are at least $e^{2d_is'-4s'}e^{-15C_1\varepsilon_1s'}$ different highly concentrated curves. Without loss of generality, we can assume that those curves are all contained in integrable surfaces of type \mathcal{P}_1 . Note that in this case every highly concentrated curve is contained in a neighborhood of the form $(X_0,Y_0)+[\beta e^{-2s'+120C_1\varepsilon_1s'}]\times [\beta e^{-s'}]$. Let us denote the neighborhood by $\mathcal{N}(X_0,Y_0)$ and cut $[\beta e^{-s'}]^2$ into union of those neighborhoods:

$$[\beta e^{-s'}]^2 = \bigcup_{i \le e^{\ell}} \mathcal{N}_i,$$

where $\ell = s' - 120C_1\varepsilon_1 s'$.

Now let us estimate the number of $\Omega(z')$'s covered by these highly concentrated curves. Our goal is to show that they cover at least

$$e^{d_i s' + 20C_1 \varepsilon_1 s'}$$

different $\Omega(z')$'s.

Counting multiplicity, these curves cover at least

$$e^{2d_is'-2s'}e^{-15C_1\varepsilon_1s'}$$

 $\Omega(z') \subset \Theta(z)$. If every $\Omega(z')$ is counted for at most $e^{d_i s' - 2s'} e^{-35C_1 \varepsilon_1 s'}$ times, then we are done. Now let us assume that some $\Omega(z_1) \subset \Theta(z)$ is counted for at least $e^{d_i s' - 2s'} e^{-35C_1 \varepsilon_1 s'}$ times. Then the highly concentrated curves containing $\Omega(z_1)$ cover at least $e^{d_i s'} e^{-35C_1 \varepsilon_1 s'}$ different $\Omega(z'')$'s. Moreover, all these highly concentrated curves are contained in some \mathcal{N}_{i_1} . Let us consider all highly concentrated curves contained in \mathcal{N}_{i_1} . We claim that counting multiplicity, there are at most $e^{2d_i s' - 800C_1 \varepsilon_1 s'}$ highly concentrated curves. In fact, fixing $\Omega(z') \subset \mathcal{N}_{i_1}$, we have that for any $\Omega(y)$ such that $\mathcal{C}(\mathcal{T}(y), z) \subset \mathcal{N}_{i_1}$, after replacing y with another element in $\Omega(y)$

$$y = \exp(\mathbf{w})u(r)z',$$

where $\mathbf{w} \in \mathfrak{r}_1 + \mathfrak{r}_2$ with $|p_{\mathbf{v}_1}(\mathbf{w})| \leq e^{120C_1\varepsilon_1 s'}$, and $|p_{\mathbf{v}_2}(\mathbf{w})| \leq e^{-2s'+120C_1\varepsilon_1 s'}$. Then

$$a(-2s')\Omega(y) \subset Q(\mathfrak{r}_1,\ell)B^{A_0H}(\beta)\tilde{z}$$

for $\ell = -s' + 120C_1\varepsilon_1 s'$ and $\tilde{z} = a(-2s')z'$. By Proposition 5.3,

$$\mu_{\mathcal{F}_i}(Q(\mathfrak{r}_1,\ell)B^{A_0H}(\beta)\tilde{z}) \leqslant e^{-d_0\ell/2} < e^{-900C_1\varepsilon_1 s'}.$$

This proves the claim.

We then repeat the above argument for all highly concentrated curves outside \mathcal{N}_{i_1} and get $\Omega(z_2)$ and at least $e^{d_i s' - 2s'} e^{-35C_1 \varepsilon_1 s'}$ different curves contained in \mathcal{N}_{i_2} containing $\Omega(z_2)$ which cover at least $e^{d_i s'} e^{-35C_1 \varepsilon_1 s'}$ different $\Omega(z'') \subset \mathcal{N}_{i_2}$. By the claim, this argument can be repeated for at least $e^{400C_1 \varepsilon_1 s'}$ times and get at least

$$e^{d_i s'} e^{-35C_1 \varepsilon_1 s'} e^{400C_1 \varepsilon_1 s'} \geqslant e^{d_i s' + 20C_1 \varepsilon_1 s'}$$

different $\Omega(z')$'s covered. This completes the proof for this case.

We then turn to the case where the contribution of integrable surfaces can be ignored. Using a dyadic pigeonhole principle, we can only consider highly concentrated curves

$$(X - X_0)(Y - Y_0) + v_1^{-1}(X - X_0) + w_1^{-1}(Y - Y_0) = 0,$$

satisfying that $|k| = |w_1/v_1| \in [e^{\ell}, 2e^{\ell}]$ for some $|\ell| < s' - 120C_1\varepsilon_1 s'$. Without loss of generality, we can assume that $\ell > 0$. By Remark 6.5, we have that every highly concentrated curve can be counted for at most $e^{2s'+2\ell+60C_1\varepsilon_1 s'}$ times. Therefore, there are at least

$$e^{2d_is'}e^{-15C_1\varepsilon_1s'}e^{-2s'-2\ell-60C_1\varepsilon_1s'} = e^{(2d_i-2)s'-2\ell}e^{-75C_1\varepsilon_1s'}$$

different highly concentrated curves in $\Theta(z)$. Every curve covers at least $e^{2s'-7C_1\varepsilon_1s'}$ different $\Omega(z')$'s. Thus, counting multiplicity, they cover at least $e^{2d_is'-2\ell}e^{-90C_1\varepsilon_1s'}$ $\Omega(z')$'s.

Our goal is to show that they cover at least

$$e^{d_i s' + 20C_1 \varepsilon_1 s'}$$

different $\Omega(z')$.

If every $\Omega(z')$ is covered by at most

$$e^{d_i s' - 2\ell} e^{-110C_1 \varepsilon_1 s'}$$

different highly concentrated curves, then we are done. Now let us assume that some $\Omega(z_0)$ is covered by at least

$$e^{d_i s' - 2\ell} e^{-110C_1 \varepsilon_1 s'}$$

different curves. Without loss of generality, let us assume that the coordinate of $\Omega(z_0)$ is **0**. Then every curve passing through $\Omega(z_0)$ is of the form

$$XY + v_1^{-1}X + w_1^{-1}Y = 0.$$

Let \mathfrak{LL} denote the set of highly concentrated curves passing through $\Omega(z_0)$.

Now let us consider the following process:

Step 0: We start with $\mathfrak{L}_0 = \emptyset$ and $\mathfrak{P}_0 = \emptyset$.

Step j+1: Suppose \mathfrak{L}_j and \mathfrak{P}_j have been constructed. We pick a highly concentrated curve $\mathcal{C} \in \mathfrak{L}\mathfrak{L}$ such that the number of points contained in \mathcal{C} but not contained in \mathfrak{P}_j is $\geqslant e^{2s'-7C_1\varepsilon_1s'}$. Then we define $\mathfrak{L}_{j+1}=\mathfrak{L}_j\cup\{\mathcal{C}\}$ and $\mathfrak{P}_{j+1}=\mathfrak{P}_j\cup\mathcal{C}$. If we can not pick such a curve, then we stop the process.

When the process is stopped, say at step n, let us take any curve $\mathcal{C} \in \mathfrak{LL} \setminus \mathfrak{L}_n$. Then by the rule of the process, we have that at least $e^{2s'-7C_1\varepsilon_1s'}$ points in \mathcal{C} are covered by curves from \mathfrak{L}_n . Let the equation of \mathcal{C} be

$$XY + v_1^{-1}X + w_1^{-1}Y = 0.$$

Using the dyadic pigeonhole principle, we have that at least $e^{2s'-7C_1\varepsilon_1s'}$ points in \mathcal{C} are covered by curves from \mathfrak{L}_n whose equation

$$XY + v_1^{'-1}X + w_1^{'-1}Y = 0,$$

satisfies that $|w_1 - w_1'| \in [e^{s'-m}, 2e^{s'-m}]$. Moreover, we can choose m as the minimum of m' satisfying the condition. By direct calculation, we can show that each such curve is contained in the $e^{-s'-m-2\ell}$ neighborhood of \mathcal{C} , and can cover at most $e^{m+2\ell+7C_1\varepsilon_1s'}$ different $\Omega(z')$ of \mathcal{C} . Therefore, there are at least $e^{2s'-m-2\ell-14C_1\varepsilon_1s'}$ such curves. Then these curves cover at least $e^{4s'-m-2\ell-21C_1\varepsilon_1s'}$ different $\Omega(z')$ in the $e^{-s'-m-2\ell}$ neighborhood of \mathcal{C} . Let us call it a (ℓ,m) -neighborhood. It is easy to see that there are at most $e^{4s'-2m-4\ell}$ different curves from \mathfrak{LL} which can be contained in the (ℓ,m) -neighborhood. Then every curve from $\mathfrak{LL} \setminus \mathfrak{L}_n$ is contained in some (ℓ,m') -neighborhood which is $e^{-21C_1\varepsilon_1s'}$ -almost filled (at least $e^{4s'-m'-2\ell-21C_1\varepsilon_1s'}$ different $\Omega(z')$ are covered) by curves from \mathfrak{L}_n . Again by a dyadic pigeonhole principle, we can find a minimal m such that at least β -proportional curves from \mathfrak{LL} is contained in some (ℓ,m) -neighborhood which is $e^{-21C_1\varepsilon_1s'}$ -almost filled by curves from \mathfrak{L}_n . Suppose we have \mathfrak{N} different such (ℓ,m) -neighborhoods. Then they cover at least

$$\mathfrak{N}e^{4s'-m-2\ell-21C_1\varepsilon_1s'}$$

different $\Omega(z')$. If

$$\mathfrak{N} \geqslant e^{d_i s' - 4s' + m + 2\ell + 41C_1 \varepsilon_1 s'},$$

then we are done. Now suppose that

$$\mathfrak{N} < e^{d_i s' - 4s' + m + 2\ell + 41C_1 \varepsilon_1 s'}.$$

Note that every (ℓ, m) -neighborhood can contain at most

$$e^{4s'-2m-4\ell}$$

different curves from \mathfrak{LL} . We have

$$\Re e^{4s'-2m-4\ell} \geqslant e^{d_i s'-2\ell} e^{-110C_1 \varepsilon_1 s'},$$

which implies that

$$\mathfrak{N} \geqslant e^{d_i s' - 4s' + 2m + 2\ell - 110C_1 \varepsilon_1 s'}.$$

This implies that

$$m \leq 151C_1\varepsilon_1s'$$
.

In this case, a (ℓ, m) -neighborhood is contained in the $e^{-s'-2\ell}$ -neighborhood of the straight line

$$v_1^{-1}X + w_1^{-1}Y = 0.$$

Note that every (ℓ, m) -neighborhood contains at least

$$e^{4s'-2\ell-172C_1\varepsilon_1s'}$$

different $\Omega(z')$. If $\ell < 10^{10} C_1 \varepsilon_1 s'$, then it covers at least

$$e^{4s'-3\times 10^{10}C_1\varepsilon_1s'}\geqslant e^{d_is'+20C_1\varepsilon_1s'}$$

different $\Omega(z')$. Then we are done. Now let us assume that $\ell \geqslant 10^{10}C_1\varepsilon_1s'$ These (ℓ, m) -neighborhoods can cover at least

$$e^{d_i s' - 4s' + 2\ell - 110C_1 \varepsilon_1 s'} e^{4s' - 2\ell - 172C_1 \varepsilon_1 s'} = e^{d_i s' - 282C_1 \varepsilon_1 s'}$$

different $\Omega(z')$. Similarly to the case where most curves are integrable, in this case, all $\Omega(z')$ covered by these (ℓ, m) -neighborhoods are contained in

$$\mathcal{N}^{\ell} = (X_0, Y_0) + [e^{-s'-\ell}] \times [\beta e^{-s'}],$$

for some (X_0, Y_0) .

We claim that counting multiplicity, there are at most

$$e^{2d_i s'} e^{-d_0 \ell/2} \le e^{2d_i s'} e^{-1000C_1 \varepsilon_1 s'}$$

highly concentrated curves contained in \mathcal{N}^{ℓ} . In fact, fixing $\Omega(z') \subset \mathcal{N}^{\ell}$, we have that for any $\Omega(y)$, such that $\mathcal{C}(\mathcal{T}(y), z) \subset \mathcal{N}^{\ell}$, after replacing y with another element in $\Omega(y)$, the following holds:

$$y = \exp(\mathbf{w})u(r)z',$$

where $\mathbf{w} \in \mathfrak{r}_1 + \mathfrak{r}_2$ with $|p_{\mathbf{v}_1}(\mathbf{w})| \leq e^{s'-\ell}$, and $|p_{\mathbf{v}_2}(\mathbf{w})| \leq e^{-s'-\ell}$. Then

$$a(-2s')\Omega(y) \subset Q(\mathfrak{r}_1,\ell)B^{A_0H}(\beta)\tilde{z},$$

where $\tilde{z} = a(-2s')z'$. By Proposition 5.3,

$$\mu_{\mathcal{F}_i}(Q(\mathfrak{r}_1,\ell)B^{A_0H}(\beta)\tilde{z})\leqslant e^{-d_0\ell/2}< e^{-1000C_1\varepsilon_1s'}.$$

This implies the claim.

Similarly to the case where most curves are integrable, we can repeat the above argument for curves outside \mathcal{N}^{ℓ} , we can get another

$$\mathcal{N}_2^{\ell} = (X_2, Y_2) + [e^{-s'-\ell}] \times [\beta e^{-s'}]$$

disjoint with \mathcal{N}^{ℓ} which also contains at least

$$e^{d_i s' - 282C_1 \varepsilon_1 s'}$$

different $\Omega(z')$. By our claim, this process can be repeated for at least $e^{800C_1\varepsilon_1 s'}$ times which gives at least

$$e^{d_i s' - 282C_1 \varepsilon_1 s'} e^{800C_1 \varepsilon_1 s'} \geqslant e^{d_i s' + 20C_1 \varepsilon_1 s'}$$

different $\Omega(z')$.

This completes the proof.

7. Proof of the main theorem

We are equipped to prove Theorem 1.3.

Proof of Theorem 1.3. By Proposition 5.2, we have that $\mu_{\mathcal{F}_0}$ is (d_0, s_0) -good. Applying Proposition 5.12 and 6.2, we get that $\mu_{\mathcal{F}_i}$ is (d_i, s_i) -good, where $d_i = d_0 + i\varepsilon_1$ and $s_i = 2^{-i}s_0$. We can proceed this inductive construction until $d_i \geq 4 - \theta$. Then by applying Proposition 3.1, we have that $a(2s_i)_*\mu_{\mathcal{F}_i}$ is effectively equidistributed. Noting that $\mu_{\mathcal{F}_i}$ is a component of $a(t-2s_i)u([1])x$, we complete the proof.

References

- [BKM01] V Bernik, D Kleinbock, and Grigorij A Margulis. Khintchine-type theorems on manifolds: the convergence case for standard and multiplicative versions. *International Mathematics Research Notices*, 2001(9):453–486, 2001.
- [Bur90] Marc Burger. Horocycle flow on geometrically finite surfaces. *Duke Mathematical Journal*, 61(3):779–803, 1990.
- [BV16] Tim Browning and Ilya Vinogradov. Effective Ratner theorem for $SL(2,\mathbb{R}) \ltimes \mathbb{R}^2$ and gaps in \sqrt{n} modulo 1. Journal of the London Mathematical Society, 94(1):61–84, 2016.
- [CY19] Sam Chow and Lei Yang. An effective ratner equidistribution theorem for multiplicative diophantine approximation on planar lines. arXiv preprint arXiv:1902.06081, 2019.
- [Dan81] S. G. Dani. Invariant measures and minimal sets of horospherical flows. *Inventiones mathematicae*, 64(2):357–385, 1981.
- [DS84] Shrikrishna G Dani and John Smillie. Uniform distribution of horocycle orbits for fuchsian groups. *Duke Mathematical Journal*, 51(1):185–194, 1984.
- [Edw21] Samuel Edwards. On the rate of equidistribution of expanding translates of horospheres in $\Gamma \setminus G$. Commentarii Mathematici Helvetici, 96(2):275–337, 2021.
- [EM22] M Einsiedler and A Mohammadi. Effective arguments in unipotent dynamics. Dynamics, geometry, number theory: the impact of margulis on modern mathematics, 2022.
- [EMMV20] Einsiedler, Margulis, Mohammadi, and Venkatesh. Effective equidistribution and property (τ) . Journal of the AMS, 33(1), 2020.
- [EMV09] Manfred Einsiedler, Gregory Margulis, and Akshay Venkatesh. Effective equidistribution for closed orbits of semisimple groups on homogeneous spaces. *Inventiones mathematicae*, 177(1):137–212, 2009.
- [FF03] Livio Flaminio and Giovanni Forni. Invariant distributions and time averages for horocycle flows. *Duke Math. J.*, 119(3):465–526, 2003.
- [Fur73] Harry Furstenberg. The unique ergodigity of the horocycle flow. In *Recent advances in topological dynamics*, pages 95–115. Springer, 1973.
- [GT12] Ben Green and Terence Tao. The quantitative behaviour of polynomial orbits on nilmanifolds. *Annals of Mathematics*, 175(2):465–540, 2012.
- [KdSSY21] Dmitry Kleinbock, Nicolas de Saxcé, Nimish A Shah, and Pengyu Yang. Equidistribution in the space of 3-lattices and dirichlet-improvable vectors on planar lines. arXiv preprint arXiv:2106.08860, 2021.
- [Kim21] Wooyeon Kim. Effective equidistribution of expanding translates in the space of affine lattices. arXiv preprint arXiv:2110.00706, 2021.
- [Kle08] Dmitry Kleinbock. An extension of quantitative nondivergence and applications to diophantine exponents. *Transactions of the American Mathematical Society*, 360(12):6497–6523, 2008.

- [KM98] Dmitry Kleinbock and Gregory Margulis. Flows on homogeneous spaces and diophantine approximation on manifolds. Annals of Mathematics, pages 339– 360, 1998.
- [KM12] Dmitry Kleinbock and Gregory Margulis. On effective equidistribution of expanding translates of certain orbits in the space of lattices, pages 385–396. Springer, Boston, MA, 2012.
- [LM14] Elon Lindenstrauss and Gregory Margulis. Effective estimates on indefinite ternary forms. *Israel Journal of Mathematics*, 203(1):445–499, 2014.
- [LM21] Elon Lindenstrauss and Amir Mohammadi. Polynomial effective density in quotients of H^3 and $H^2 \times H^2$. arXiv preprint arXiv:2112.14562, 2021.
- [LMW22] Elon Lindenstrauss, Amir Mohammadi, and Zhiren Wang. Polynomial effective equidistribution. arXiv preprint arXiv:2202.11815, 2022.
- [Mar04] Gregori Aleksandrovitsch Margulis. On some aspects of the theory of anosov systems. In *On Some Aspects of the Theory of Anosov Systems*, pages 1–71. Springer, 2004.
- [MS95] Shahar Mozes and Nimish Shah. On the space of ergodic invariant measures of unipotent flows. *Ergodic Theory and Dynamical Systems*, 15(01):149–159, 1995.
- [NV21] P. Nelson and A. Venkatesh. The orbit method and analysis of automorphic forms. *Acta Mathematica*, 226(1), 2021.
- [Rat90a] Marina Ratner. On measure rigidity of unipotent subgroups of semisimple groups. *Acta mathematica*, 165:229–309, 1990.
- [Rat90b] Marina Ratner. Strict measure rigidity for unipotent subgroups of solvable groups. *Inventiones mathematicae*, 101(1):449–482, 1990.
- [Rat91a] Marina Ratner. On Raghunathan's measure conjecture. *Annals of Mathematics*, pages 545–607, 1991.
- [Rat91b] Marina Ratner. Raghunathan's topological conjecture and distributions of unipotent flows. *Duke Math. J*, 63(1):235–280, 1991.
- [Sar82] Peter Sarnak. Asymptotic behavior of periodic orbits of the horocycle flow and eisenstein series. Communications on Pure and Applied Mathematics, 34(6):719–739, 1982.
- [Sha09] Nimish A Shah. Equidistribution of expanding translates of curves and Dirichlet's theorem on diophantine approximation. *Inventiones mathematicae*, 177(3):509–532, 2009.
- [Str04] Andreas Strömbergsson. On the uniform equidistribution of long closed horocycles. *Duke Math. J.*, 123(3):507–547, 06 2004.
- [Str13] Andreas Strömbergsson. On the deviation of ergodic averages for horocycle flows. *Journal of Modern Dynamics*, 7:291, 2013.
- [Str15] Andreas Strömbergsson. An effective Ratner equidistribution result for $SL(2, \mathbb{R}) \ltimes \mathbb{R}^2$. Duke Mathematical Journal, 164(5):843–902, 04 2015.
- [SU15] Peter Sarnak and Adrián Ubis. The horocycle flow at prime times. *Journal de Mathématiques Pures et Appliquées*, 103(2):575–618, 2015.
- [Ven10] A. Venkatesh. Sparse equidistribution problems, period bounds and subconvexity. *Annals of Mathematics*, 172:989–1094, 2010.

Lei Yang, College of Mathematics, Sichuan University, Chengdu, Sichuan, 610000, China

Email address: lyang861028@gmail.com