

TWO-TERM SPECTRAL ASYMPTOTICS IN LINEAR ELASTICITY ON A RIEMANNIAN MANIFOLD

GENQIAN LIU

ABSTRACT. In this note, by explaining two key methods that were employed in [Liu-21] and by giving some remarks, we show that the proof of Theorem 1.1 in [Liu-21] is a rigorous proof based on theory of strongly continuous semigroups and pseudodifferential operators. All remarks and comments to paper [Liu-21], which were given by Matteo Capoferri, Leonid Friedlander, Michael Levitin and Dmitri Vassiliev in [CaFrLeVa-22], are incorrect. The so-called “numerical counter-examples” in [CaFrLeVa-22] are useless examples for the two-term asymptotics of the counting functions of the elastic eigenvalues. Clearly, the conclusion and the proof of [Liu-21] are completely correct.

1. TWO KEY METHODS

Let (Ω, g) be a compact smooth Riemannian n -manifold with smooth boundary $\partial\Omega$. Let P_g be the Navier–Lamé operator:

$$(1.1) \quad P_g \mathbf{u} := \mu \nabla^* \nabla \mathbf{u} - (\mu + \lambda) \operatorname{grad} \operatorname{div} \mathbf{u} - \mu \operatorname{Ric}(\mathbf{u}), \quad \mathbf{u} = (u^1, \dots, u^n),$$

where μ and λ are Lamé parameters satisfying $\mu > 0$ and $\mu + \lambda > 0$, $\nabla^* \nabla$ is the Bochner Laplacian (see (2.11) of [Liu-21]), div and grad are the usual divergence and gradient operators, and

$$(1.2) \quad \operatorname{Ric}(\mathbf{u}) = \left(\sum_{l=1}^n R_l^1 u^l, \sum_{l=1}^n R_l^2 u^l, \dots, \sum_{l=1}^n R_l^n u^l \right)$$

denotes the action of Ricci tensor $R_l^j := \sum_{k=1}^n R_{lk}^{kj}$ on \mathbf{u} .

We denote by P_g^- and P_g^+ the Navier–Lamé operators with zero Dirichlet and Neumann boundary conditions, respectively. The Dirichlet boundary condition is $\mathbf{u}|_{\partial\Omega}$, and the Neumann boundary condition is $\frac{\partial \mathbf{u}}{\partial \nu} := 2\mu(\operatorname{Def} \mathbf{u})^\# \nu + \lambda(\operatorname{div} \mathbf{u})\nu$ on $\partial\Omega$, where $\operatorname{Def} \mathbf{u} = \frac{1}{2}(\nabla \mathbf{u} + \nabla \mathbf{u}^T)$, $\#$ is the sharp operator (for a tensor) by raising index, and ν is the unit outer normal to $\partial\Omega$.

Since P_g^- (respectively, P_g^+) is an unbounded, self-adjoint and positive (respectively, nonnegative) operator in $[H_0^1(\Omega)]^n$ (respectively, $[H^1(\Omega)]^n$) with discrete spectrum $0 < \tau_1^- < \tau_2^- \leq \dots \leq \tau_k^- \leq \dots \rightarrow +\infty$ (respectively, $0 \leq \tau_1^+ < \tau_2^+ \leq \dots \leq \tau_k^+ \leq \dots \rightarrow +\infty$), one has

$$(1.3) \quad P_g^\mp \mathbf{u}_k^\mp = \tau_k^\mp \mathbf{u}_k^\mp,$$

where $\mathbf{u}_k^- \in [H_0^1(\Omega)]^n$ (respectively, $\mathbf{u}_k^+ \in [H^1(\Omega)]^n$) is the eigenvector corresponding to eigenvalue τ_k^- (respectively, τ_k^+).

In [Liu-21], we obtained the following result:

Theorem 1.1. *Let (Ω, g) be a compact smooth Riemannian manifold of dimension n with smooth boundary $\partial\Omega$, and let $0 < \tau_1^- < \tau_2^- \leq \dots \leq \tau_k^- \leq \dots$ (respectively, $0 \leq \tau_1^+ < \tau_2^+ \leq \tau_3^+ \leq \dots \leq \tau_k^+ \leq \dots$) be the eigenvalues of the Navier–Lamé operator P_g^- (respectively, P_g^+) with respect to the zero Dirichlet (respectively, Neumann) boundary condition. Then*

$$(1.4) \quad \sum_{k=1}^{\infty} e^{-\tau_k^\mp t} = \operatorname{Tr}(e^{-tP_g^\mp}) = \left[\frac{n-1}{(4\pi\mu t)^{n/2}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}} \right] \operatorname{Vol}(\Omega)$$

2020 *Mathematics Subject Classification.* 74B05, 35K50, 35P20, 35S05.

Key words and phrases. Linear elasticity; Eigenvalues; Spectral asymptotics; Pseudodifferential operators.

$$\mp \frac{1}{4} \left[\frac{n-1}{(4\pi\mu t)^{(n-1)/2}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{(n-1)/2}} \right] \text{Vol}(\partial\Omega) + O(t^{1-n/2}) \quad \text{as } t \rightarrow 0^+.$$

Here $\text{Vol}(\Omega)$ denotes the n -dimensional volume of Ω , $\text{Vol}(\partial\Omega)$ denotes the $(n-1)$ -dimensional volume of $\partial\Omega$.

According to the proof of Theorem 1.1, in particular, by the expressions of \mathbf{q}_{-3} and \mathbf{q}_{-4} in (3.10) of [Liu-21], we actually proved the following result (see, ii) of Remark 4.1 in [Liu-21]):

Theorem 1.2. *Let (Ω, g) be a compact smooth Riemannian manifold of dimension n with smooth boundary $\partial\Omega$. Then*

$$\sum_{k=1}^{\infty} e^{-\tau_k^\mp t} = t^{-n/2} [a_0 + a_1^\mp t^{1/2} + a_2^\mp t + \cdots + a_m^\mp t^{m/2} + O(t^{(m+1)/2})] \quad \text{as } t \rightarrow 0^+,$$

where all coefficients $a_0, a_k^\mp, (k = 1, 2, 3, \dots)$ can be explicitly obtained by an algorithm (see the proof of [Liu-21]). In particular,

$$(1.5) \quad a_0 = \left[\frac{n-1}{(4\pi\mu)^{n/2}} + \frac{1}{(4\pi(2\mu+\lambda))^{n/2}} \right] \text{Vol}(\Omega),$$

$$(1.6) \quad a_1^\mp = \mp \frac{1}{4} \left[\frac{n-1}{(4\pi\mu)^{(n-1)/2}} + \frac{1}{(4\pi(2\mu+\lambda))^{(n-1)/2}} \right] \text{Vol}(\partial\Omega),$$

$$(1.7) \quad a_2^\mp = \left[\frac{n-1}{(4\pi\mu)^{n/2}} + \frac{1}{(4\pi(2\mu+\lambda))^{n/2}} \right] \left(\frac{1}{3} \int_{\Omega} R dV - \frac{1}{6} \int_{\partial\Omega} H ds \right),$$

where R is the scalar curvature of Ω and H is the mean curvature of $\partial\Omega$.

In the proof of Theorem 1.1 in [Liu-21], we have used two key methods: the ‘‘method of image’’ and the ‘‘explicitly calculable method’’ for the full symbol of the resolvent operator $(\tau I - P_g)^{-1}$. In this note, by explaining the two key methods again, and by giving some remarks, we will show that our conclusion and the proof of Theorem 1.1 in [Liu-21] are correct.

In order to explain the ‘‘method of image’’, we first introduce its original idea for calculating the Green’s function of a parabolic (elastic) system with zero Dirichlet (or Neumann) boundary condition in the upper half-space $\mathbb{R}_+^n := \{x = (x', x_n) \in \mathbb{R}^n \mid x_n > 0\}$, where $x' = (x_1, \dots, x_{n-1})$.

Let $\mathbf{K}(t, x, y)$ be the fundamental solution of the parabolic (elastic) system $\frac{\partial \mathbf{u}}{\partial t} - P\mathbf{u} = 0$ in \mathbb{R}^n , where $P = \mu\Delta\mathbf{u} + (\lambda + \mu)\nabla(\nabla \cdot \mathbf{u})$. In fact, $\mathbf{K}(t, x, y)$ in \mathbb{R}^n can be explicitly written out (but we will not need such an explicit expression here). That is, for any $x, y \in \mathbb{R}^n$,

$$\frac{\partial \mathbf{K}(t, x, y)}{\partial t} - P_y \mathbf{K}(t, x, y) = \delta(x - y).$$

Here we write the elastic operator P as P_y to denote the corresponding partial derivatives acting on variable y . Now, let us restrict $x, y \in \mathbb{R}_+^n$. It is very easy to verify that $\mathbf{K}^-(t, x, y) := \mathbf{K}(t, x, y) - \mathbf{K}(t, x, y^*)$ and $\mathbf{K}^+ := \mathbf{K}(t, x, y) + \mathbf{K}(t, x, y^*)$ are the Green’s functions of $\frac{\partial \mathbf{u}}{\partial t} - P_y \mathbf{u} = 0$ in \mathbb{R}_+^n with zero Dirichlet and Neumann boundary conditions, respectively, where $y^* = (y', -y_n)$. In other words,

$$\begin{cases} \frac{\partial \mathbf{K}^-(t, x, y)}{\partial t} - P_y \mathbf{K}^-(t, x, y) = \delta(x - y), & x, y \in \mathbb{R}_+^n, \\ \mathbf{K}^-(t, x, y) = 0, & x \in \mathbb{R}_+^n, y \in \partial\mathbb{R}_+^n, \end{cases}$$

and

$$\begin{cases} \frac{\partial \mathbf{K}^+(t, x, y)}{\partial t} - P_y \mathbf{K}^+(t, x, y) = \delta(x - y), & x, y \in \mathbb{R}_+^n, \\ \frac{\partial \mathbf{K}^+(t, x, y)}{\partial \nu} = 0, & x \in \mathbb{R}_+^n, y \in \partial\mathbb{R}_+^n, \end{cases}$$

where $\frac{\partial \mathbf{K}^+}{\partial \nu} := \mu(\nabla \mathbf{K}^+ + (\nabla \mathbf{K}^+)^T)\nu + \lambda(\text{div } \mathbf{K}^+)\nu$ on $\partial\mathbb{R}_+^n$.

Such a basic “reflection method” for obtaining Green’s functions of parabolic (elastic) system in \mathbb{R}_+^n with zero Dirichlet (or Neumann) boundary condition can be generalized to a smooth Riemannian manifold with smooth boundary. This technique is also called the “method of image”. Let us introduce it again (the details can be seen in [Liu-21]). Let $\mathcal{M} = \Omega \cup (\partial\Omega) \cup \Omega^*$ be the (closed) double of Ω . Note that \mathcal{M} is a closed manifold since we can take the boundary normal coordinates on a neighborhood of $\partial\Omega$, and used the metric (see p. 10169, p. 10183 and p. 10187 of [Liu-21])

$$(1.8) \quad \begin{aligned} g_{jk}(x^*) &= -g_{jk}(x) \quad \text{for } j < k = n \text{ or } k < j = n, \\ &= g_{jk}(x) \quad \text{for } j, k < n \text{ or } j = k = n, \end{aligned}$$

$$(1.9) \quad g_{jk}(x) = 0 \quad \text{for } j < k = n \text{ or } k < j = n \text{ on } \partial\Omega,$$

$$(1.10) \quad \sqrt{|g|/g_{nn}} dx_1 \cdots dx_{n-1} = \text{the element of (Riemannian) surface area on } \partial\Omega,$$

where $x_n(x^*) = -x_n(x)$. For given $\epsilon > 0$, denote by $U_\epsilon(\partial\Omega) = \{z \in \mathcal{M} \mid \text{dist}(z, \partial\Omega) < \epsilon\}$ the ϵ -neighborhood of $\partial\Omega$ in \mathcal{M} . Clearly, for any $W \subset U_\epsilon(\partial\Omega)$ with local coordinates $x = (x', x_n)$ such that $\epsilon > x_n > 0$ in $W \cap \Omega$; $x_n = 0$ on $W \cap \partial\Omega$, and $|\nabla x_n| = 1$ near $\partial\Omega$. It is easy to verify (see [Liu-21]) that such an extend metric g is C^0 -smooth on whole \mathcal{M} and C^∞ -smooth in $\mathcal{M} \setminus \partial\Omega$. Let us point out that obtaining a closed double manifold \mathcal{M} from Ω is a standard and classical method in Riemannian geometry (see, Example 9.32 on p. 226 in [Lee-13] or [MS-67]). Roughly speaking, Ω and Ω^* are C^0 -smoothly glued into \mathcal{M} . Next, let \mathcal{P} be the double to \mathcal{M} of the operator P_g on Ω . Then the coefficients of \mathcal{P} are also C^0 -smooth on \mathcal{M} and C^∞ -smooth in $\mathcal{M} \setminus \partial\Omega$. Of course, \mathcal{P} is an elliptic operator on \mathcal{M} . Thus we immediately get a strongly continuous semigroup $e^{-t\mathcal{P}}$ on \mathcal{M} . Let $\mathbf{K}(t, x, y)$ be the parabolic kernel (i.e., a fundamental solution to parabolic (elastic) system $\frac{\partial \mathbf{u}}{\partial t} - \mathcal{P}\mathbf{u} = 0$ on whole (closed) manifold \mathcal{M}). More precisely,

$$\frac{\partial \mathbf{K}(t, x, y)}{\partial t} - \mathcal{P}_y \mathbf{K}(t, x, y) = \delta(x - y), \quad x, y \in \mathcal{M},$$

where we have written \mathcal{P} as \mathcal{P}_y to denote the corresponding covariant derivatives acting on variable y . Obviously, $\mathbf{K} \in C^1((0, \infty) \times \mathcal{M} \times \mathcal{M}) \cap C^\infty((0, \infty) \times (\mathcal{M} \setminus \partial\Omega) \times (\mathcal{M} \setminus \partial\Omega))$. Then

$$(1.11) \quad \mathbf{K}^-(t, x, y) = \mathbf{K}(t, x, y) - \mathbf{K}(t, x, y^*), \quad x, y \in \Omega$$

and

$$(1.12) \quad \mathbf{K}^+(t, x, y) = \mathbf{K}(t, x, y) + \mathbf{K}(t, x, y^*), \quad x, y \in \Omega$$

are Green’s functions of the parabolic (elastic) system $\frac{\partial \mathbf{u}}{\partial t} - P_g \mathbf{u} = 0$ in Ω with zero Dirichlet and Neumann boundary conditions, respectively. That is,

$$(1.13) \quad \begin{cases} \frac{\partial \mathbf{K}^-(t, x, y)}{\partial t} - P_g \mathbf{K}^-(t, x, y) = \delta(x - y), & x, y \in \Omega, \\ \mathbf{K}^-(t, x, y) = 0, & x \in \Omega, y \in \partial\Omega, \end{cases}$$

$$(1.14) \quad \begin{cases} \frac{\partial \mathbf{K}^+(t, x, y)}{\partial t} - P_g \mathbf{K}^+(t, x, y) = \delta(x - y), & x, y \in \Omega, \\ \frac{\partial \mathbf{K}^+(t, x, y)}{\partial \nu} = 0, & x \in \Omega, y \in \partial\Omega, \end{cases}$$

where $\frac{\partial \mathbf{K}^+}{\partial \nu} := 2\mu(\text{Def } \mathbf{K}^+)^\# \boldsymbol{\nu} + \lambda(\text{div } \mathbf{K}^+) \boldsymbol{\nu}$ is the Neumann boundary condition again. Note that all covariant derivatives act on variable y for any fixed $x \in \Omega$. This method based on theory of parabolic system on Riemannian manifolds. By the “method of image” and (1.11)–(1.14), the studied topic becomes to discuss the differences of two fundamental solutions on \mathcal{M} , further becomes to study such a problem which is restricted on Ω . Of course, one can get all the coefficients a_i^\mp by classical estimates. But we have used the method of pseudodifferential operators in [Liu-21] (which is simpler and more effective) as follows.

It is well-known that

$$(1.15) \quad \int_{\Omega} (\text{Tr } (\mathbf{K}^\mp(t, x, x))) dV = \sum_{k=1}^{\infty} e^{-t\tau_k^\mp}.$$

Since $e^{-t\mathcal{P}}f(x) = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} (\tau I - \mathcal{P})^{-1} f(x) d\tau$, we have

$$e^{-t\mathcal{P}}f(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{ix \cdot \xi} \left(\frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} \iota((\tau I - \mathcal{P})^{-1}) \hat{f}(\xi) d\tau \right) d\xi,$$

so that

$$\begin{aligned} \mathbf{K}(t, x, y) &= e^{-t\mathcal{P}}\delta(x-y) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-y) \cdot \xi} \left(\frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} \iota((\tau I - \mathcal{P})^{-1}) d\tau \right) d\xi \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-y) \cdot \xi} \left(\frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} \left(\sum_{l \geq 0} \mathbf{q}_{-2-l}(x, \xi, \tau) \right) d\tau \right) d\xi, \end{aligned}$$

where \mathcal{C} is a suitable curve in the complex plane in the positive direction around the spectrum of \mathcal{P} , and $\iota((\tau I - \mathcal{P})^{-1}) := \sum_{l \geq 0} \mathbf{q}_{-2-l}(x, \xi, \tau)$ is the full symbol of resolvent operator $(\tau I - \mathcal{P})^{-1}$. This implies that for any $x \in \Omega$,

$$\begin{aligned} \text{Tr}(\mathbf{K}(t, x, x)) &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \left(\frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} \sum_{l \geq 0} \text{Tr}(\mathbf{q}_{-2-l}(x, \xi, \tau)) d\tau \right) d\xi, \\ \text{Tr}(\mathbf{K}(t, x, x^*)) &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-x^*) \cdot \xi} \left(\frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} \sum_{l \geq 0} \text{Tr}(\mathbf{q}_{-2-l}(x, \xi, \tau)) d\tau \right) d\xi. \end{aligned}$$

Therefore, whether one can get all coefficients of the parabolic trace, it depends strongly on whether one can explicitly calculate the full symbol of the resolvent operator $(\tau I - \mathcal{P})^{-1}$. The second key idea (or main contribution) of [Liu-21] is that such kind of full symbol can be explicitly obtained by a surprising new technique [Liu-19] (see below).

Since the full symbol $\sigma((\tau I - P_g)^{-1})$ of $(\tau I - P_g)^{-1}$ is classical, we write the full symbol $\sigma((\tau I - P_g)^{-1}) \sim \mathbf{q}_{-2} + \mathbf{q}_{-3} + \mathbf{q}_{-4} + \dots$ with \mathbf{q}_j being homogeneous of degree j ($j = -2, -3, \dots$) in the variable $(\xi, \tau^{1/2})$. Generally, for an elliptic system, it is not possible to explicitly calculate the full symbol of its resolvent operator. However, for the elastic Lamé system, we discovered that the full symbol is explicitly calculable. In [Liu-21], we have explicitly calculated the full symbol by a method of operator algebra (This is a surprising effective method stemmed from the idea of [Liu-19] for studying the elastic Dirichlet-to-Neumann map). Thus by the parabolic trace method, we can get all coefficients $a_0, a_1^{\mp}, a_2^{\mp}, \dots$ in the expansion of the parabolic trace

$$\sum_{k=1}^{\infty} e^{-t\tau_k^{\mp}} = t^{-n/2} \left[a_0 + a_1^{\mp} t^{1/2} + a_3^{\mp} t + a_4^{\mp} t^{3/2} + \dots \right].$$

In [Liu-21], we had obtained a_0 and a_1^{\mp} by an (algorithm) method. From the expression (3.10) in [Liu-21], we can get \mathbf{q}_{-3} and \mathbf{q}_{-4} , i.e.,

$$(1.16) \quad \mathbf{q}_{-3}(x, \xi, \tau) = -\mathbf{a}_2^{-1} \left(\mathbf{a}_1 \mathbf{a}_2^{-1} - i \sum_{l=1}^n \frac{\partial \mathbf{a}_2}{\partial \xi_l} \frac{\partial \mathbf{a}_2^{-1}}{\partial x_l} \right),$$

$$(1.17) \quad \mathbf{q}_{-4} = -\mathbf{a}_2^{-1} \left(\sum_{\substack{j < 2 \\ j+|\alpha|-k=0}} (\partial_{\xi}^{\alpha} \mathbf{a}_k) (D_x^{\alpha} \mathbf{q}_{-2-j}) / \alpha! \right).$$

Note that \mathbf{q}_{-3} has not any contribution to the coefficients a_l^{\mp} because it is an odd function in variable $\xi \in \mathbb{R}^n$ (the corresponding integral on \mathbb{R}^n will vanish). Further, by inserting \mathbf{q}_{-4} into our calculation procedure for the coefficients a_2^{\mp} (see [Liu-21]), we immediately get the explicit expression of a_2^{\mp} (i.e., (1.7)). To this end, we need to use the expansion of the metric g : we can write

$$g_{jk}(x) = \delta_{jk} + \sum_{l=1}^n \frac{\partial g_{jk}}{\partial x_l}(0) x_l + \frac{1}{2} \sum_{l,m=1}^n \frac{\partial^2 g_{jk}}{\partial x_l \partial x_m}(0) x_l x_m + O(|x|^3) \text{ near } 0.$$

In geodesic normal coordinate system centered at $x = 0$, as Riemann showed, one has

$$\frac{\partial^2 g_{lm}}{\partial x_j \partial x_j} = -\frac{1}{3} R_{l j m k} - \frac{1}{3} R_{l k m j}.$$

Substituting this into \mathbf{q}_{-4} in [Liu-21], and by calculating

$$\int_{W \cap \Omega} \left\{ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \left(\frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} \operatorname{Tr}(\mathbf{q}_{-4}(x, \xi, \tau)) d\tau \right) d\xi \right\} dV,$$

where $W \subset \Omega \setminus U_\epsilon(\partial\Omega)$, we obtain the term that will product the factor $\frac{t}{3} \int_{\Omega} R dV$. Note that under the boundary normal coordinates, $\frac{1}{2} \frac{\partial g_{jk}}{\partial x_n} = \delta_{jk} \tilde{\kappa}_j$ on $\partial\Omega$. By calculating

$$\int_{\tilde{W} \cap \Omega} \left\{ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \left(\frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} \operatorname{Tr}(\mathbf{q}_{-4}(x, \xi, \tau)) d\tau \right) d\xi \right\} dV,$$

we get another term that will product the factor $-\frac{t}{6} \int_{\tilde{W} \cap \partial\Omega} H ds$, where $\tilde{W} \subset U_\epsilon(\partial\Omega)$. In addition, it is easy to verify that

$$\begin{aligned} \int_{W \cap \Omega} \left\{ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-\tilde{x}) \cdot \xi} \left(\frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} \operatorname{Tr}(\mathbf{q}_{-4}(x, \xi, \tau)) d\tau \right) d\xi \right\} dV &= O(t^{2-\frac{n}{2}}) \quad \text{as } t \rightarrow 0^+, \\ \int_{\tilde{W} \cap \Omega} \left\{ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-\tilde{x}) \cdot \xi} \left(\frac{1}{2\pi i} \int_{\mathcal{C}} e^{-t\tau} \operatorname{Tr}(\mathbf{q}_{-4}(x, \xi, \tau)) d\tau \right) d\xi \right\} dV &= O(t^{2-\frac{n}{2}}) \quad \text{as } t \rightarrow 0^+. \end{aligned}$$

Therefore, Theorem 1.2 is proved.

Based on classical theory of strongly continuous semigroups and pseudodifferential operators, we had given a rigorous proof to Theorem 1.1 in [Liu-21]. Here we also give an outline of proof of Theorem 1.2. Clearly, the conclusion and proof of [Liu-21] are correct.

2. SEVERAL REMARKS

Remark 2.1. Besides the elastic Lamé system, by applying our new technique we can also explicitly calculate and obtain the full symbol of the resolvent operator for the thermoelastic system [LiuTan-22] (it is also an elliptic system). Furthermore, the parabolic trace for this elliptic system can also be explicitly obtained.

Generally, for any matrix-valued elliptic operator L , if one can explicitly obtain the inverse matrix for the principal symbol of the operator $(\tau I - L)^{-1}$, then all coefficients in the asymptotic expansion of the parabolic trace will immediately be obtained by our new (algorithm) method.

Remark 2.2. Our paper [Liu-21] only studied the elastic Lamé system (i.e., elliptic system) for given $\mu > 0$ and $\lambda + \mu > 0$. As $\lambda + \mu \rightarrow 0$, our asymptotic expansion tends to the corresponding (well-known) result for the Laplace–Beltrami operator. In fact, all results and discussions in [Liu-21] still hold for $\mu > 0$ and $\lambda + \mu \geq 0$.

Remark 2.3. Two-term asymptotics for the counting function $\mathcal{N}(\Lambda)$ as $\Lambda \rightarrow +\infty$ implies two-term asymptotics for the partition function $\mathcal{Z}(t)$ as $t \rightarrow 0^+$ under “elastic billiards” assumption, however, the converse may not true, where $\mathcal{N}(\Lambda) := \#\{k | \tau_k \leq \Lambda\}$, and $\mathcal{Z}(t) := \sum_{k=1}^{\infty} e^{-t\tau_k}$. The reason is: the famous Tauberian theorem shows that one-term asymptotics for $\mathcal{Z}(t)$ as $t \rightarrow 0^+$ implies one-term asymptotics for $\mathcal{N}(\Lambda)$ as $\Lambda \rightarrow +\infty$. But, the Tauberian theorem is not true for two-term asymptotics. More precisely, let $v(s)$ vanish for $s < 0$, be nondecreasing, continuous from the right, and suppose that for some constant $\alpha > 0$,

$$\int_0^{\infty} e^{-ts} dv(s) \sim \frac{A}{t^\alpha} + \frac{B}{t^{\alpha-1}} + o(t^{1-\alpha}) \quad \text{as } t \rightarrow 0^+,$$

then one can not obtain

$$v(s) \sim \frac{A}{\Gamma(\alpha+1)} s^\alpha + \frac{B}{\Gamma(\alpha)} s^{\alpha-1} + o(s^{\alpha-1}) \quad \text{as } s \rightarrow +\infty.$$

Professor J. Korevaar (the author of “Tauberian Theory: A Century of Developments”, Springer-Verlag, 2004) told the author of [Liu-21] this well-known fact with a detailed example in a private e-mail about sixteen years ago. Clearly, two-term asymptotics as $t \rightarrow 0^+$ for the partition function $\mathcal{Z}(t)$ in [Liu-21] can not go back to two-term asymptotics for $\mathcal{N}(\Lambda)$ as $\Lambda \rightarrow +\infty$. Thus, any numerical (verification) comparison must be done in the setting of $\mathcal{Z}(t)$ as $t \rightarrow 0^+$ (here only an additional “elastic billiards” assumption is added) rather than to consider $\mathcal{N}(\Lambda)$ as $\Lambda \rightarrow +\infty$ as done in [CaFrLeVa-22]. Obviously, putting numerical experimental verification in the setting of $\mathcal{N}(\Lambda)$ with Λ in a finite interval is a fundamental mistake in [CaFrLeVa-22].

In the numerical verification in [CaFrLeVa-22], the number $\Lambda = 1600$ or $\Lambda = 2400$ or $\Lambda = 2800$, or $\Lambda = 2000$ or $\lambda = 3000$ are too small. Even one takes $\Lambda = 1000^{1000}$ or $\Lambda = 10000^{10000}$ or $\Lambda = 100000^{100000}$, it may not describe any true asymptotic case of $\mathcal{N}(\Lambda) - a\Lambda$ as $\Lambda \rightarrow +\infty$, where a is the one-term coefficient of $\mathcal{N}(\Lambda)$. Numerical experimental verification should consider the case after large Λ rather than before. In [CaFrLeVa-22], it seems that the numerical (experimental) data have “satisfying” effect for their second term of the counting function when $0 \leq \Lambda \leq 3000$. However, it is useless to study the asymptotic behavior as $\Lambda \rightarrow +\infty$. Such a numerical experimental verification for small Λ may not provide any valuable information for the future case of $\mathcal{N}(\Lambda) - a\Lambda$ as $\Lambda \rightarrow +\infty$. More precisely, in order to verify

$$\lim_{\Lambda \rightarrow +\infty} \frac{1}{\Lambda^{(n-1)/2}} \left[\mathcal{N}(\Lambda) - a \text{Vol}_n(\Omega) \Lambda^{n/2} \right] = b'$$

for some constant b' , by the definition of limit, one should prove that for every number $\epsilon > 0$ there is a number $M > 0$ which depends on ϵ such that if $\Lambda > M$ then

$$\left| \frac{1}{\Lambda^{(n-1)/2}} \left[\mathcal{N}(\Lambda) - a \text{Vol}_n(\Omega) \Lambda^{n/2} \right] - b' \right| < \epsilon.$$

The authors of [CaFrLeVa-22] claimed that their result was “correct” only by “numerically” verifying the above inequality for some $\epsilon_0 > 0$ when $0 \leq \Lambda \leq 3000$ or $0 \leq \Lambda \leq 2400$ or $0 \leq \Lambda \leq 2600$ for the unit disk and unit square. Such a so-called “numerical verification” method is not believable at all.

Remark 2.4. The authors of [CaFrLeVa-22] essentially discussed the elastic operator in a very special domain (i.e., the Euclidean upper half-space with flat metric $g_{jk} = \delta_{jk}$). Their method is “stretch Ω by a linear factor $\kappa > 0$, note that the eigenvalues then rescale as κ^{-2} , and check the rescaling of the geometric invariants and of (1.20)” (see, from line 1 to line 2 on p. 6 of [CaFrLeVa-22]). However, for the two-term asymptotics, such a method will lead to losing much useful information because the boundary surface “scrambles” the elastic waves. In fact, in a Riemannian manifold (Ω, g) , it follows from Lemma 2.1.1 of [Liu-19] that the elastic operator has the following local representation:

$$\begin{aligned} & \mu \sum_{j=1}^n \left\{ \Delta_g u^j + 2 \sum_{k,s,l=1}^n g^{kl} \Gamma_{sk}^j \frac{\partial u^s}{\partial x_l} + \sum_{k,s,l=1}^n \left(g^{kl} \frac{\partial \Gamma_{sl}^j}{\partial x_k} + \sum_{h=1}^n g^{kl} \Gamma_{hl}^j \Gamma_{sk}^h - \sum_{h=1}^n g^{kl} \Gamma_{sh}^j \Gamma_{kl}^h \right) u^s \right\} \frac{\partial}{\partial x_j} \\ & + (\lambda + \mu) \text{grad div } \mathbf{u} + \mu \text{ Ric}(\mathbf{u}). \end{aligned}$$

In [CaFrLeVa-22], by erroneously thinking of the above elastic operator defined on (Ω, g) as $\mu \sum_{j=1}^n \frac{\partial^2 \mathbf{u}}{\partial x_j^2} + (\lambda + \mu) \nabla(\nabla \cdot \mathbf{u})$ on $\bar{\mathbb{R}}_+^n$, and further by mistaking a neighborhood of the boundary as the Euclidean upper half-space, the authors of [CaFrLeVa-22] had completely changed the original eigenvalues problems into other different spectral problems. The reader can not see where the branching Hamiltonian billiards condition (i.e., the corresponding billiard is neither dead-end nor absolutely periodic) on the Riemannian manifold is used in the proof of [CaFrLeVa-22]. In a Riemannian manifold, in order to get the two-term asymptotics of the counting function, the cut-off function should be introduced. See Response 6 in Section 3 for a correct approach to obtain the two-term asymptotics.

3. SOME RESPONSES

We will give some responses to remarks in [CaFrLeVa-22] that involved the paper [Liu-21]:

Response 1. On p. 27 of [CaFrLeVa-22], from line 6 to line 10, the authors of [CaFrLeVa-22] wrote “Finally, [Liu-21, p. 10166] contains a remarkable claim: “McKean and Singer ...proved the famous Kac conjecture...”. Here the conjecture in question is the “Can one hear the shape of a drum?” one, which was proven to be false in [GoWeWo-92]. The question of whether a similar conjecture is true for the operator of linear elasticity is still fully open, to the best of our knowledge.”

Clearly, the authors of [CaFrLeVa-22] have confused the famous Kac question and Kac conjecture, and given an incorrect comment. Let us write out the original sentences in [Kac-66] and [MS-67] as follows.

The Kac question: On p. 2–p. 3 of [Kac-66], M. Kac wrote “Assume that for each n the eigenvalue λ_n for Ω_1 is equal to the eigenvalue μ_n for Ω_2 . Question: Are the regions Ω_1 and Ω_2 congruent in the sense of Euclidean geometry? ”

The Kac conjecture: On p. 23 of [Kac-66], M. Kac wrote “It is thus natural to conjecture that for a *smooth* drum with r *smooth* holes

$$\sum_{n=1}^{\infty} e^{-\lambda_n t} \sim \frac{|\Omega|}{2\pi t} - \frac{L}{4} \frac{1}{\sqrt{2\pi t}} + \frac{1}{6}(1-r),$$

and that therefore one can “hear” the connectivity of the drum!”

On p. 44 of [MS-67], H. McKean and I. Singer wrote “Kac was led to conjecture

$$(4b) \quad Z = \frac{\text{area}}{4\pi t} - \frac{\text{length}/4}{\sqrt{4\pi t}} + \frac{1}{6}(1-h) + o(1) \quad (t \downarrow 0)$$

for regions D with smooth B and $h < \infty$ holes, and was able to prove the correctness of the first 2 terms. This jibes with an earlier conjecture of A. Pleijel and suggests that you can hear *the number of holes*. (4b) will be proved below in a form applicable both to open manifolds with compact boundary and to closed manifolds.”

In [Liu-21], on p. 10166, from line –14 to line –1, the author of [Liu-21] accurately introduced the Kac question and the Kac conjecture according to the original literatures [Kac-66] and [MS-67]. The comment (in [Liu-21]) “... a celebrated result of the spectral (geometric) invariants had been obtained by McKean and Singer. They proved the famous Kac conjecture and gave an explicit expression to the first three coefficients of asymptotic expansion for the heat trace of the Laplacian on a bounded domain Ω of a Riemannian manifold: ...” is a proper and correct statement.

Response 2. In the footnote 3 on p. 3 of [CaFrLeVa-22], the authors of [CaFrLeVa-22] wrote “We will not call (1.6) the Neumann condition in order to avoid confusion with erroneous “Neumann” condition in [Liu-21].” And for the remark on p. 9 of [CaFrLeVa-22], “**The first fundamental mistake of [Liu-21] is ...** ”

In fact, in [Liu-21] the Neumann boundary condition is $\frac{\partial \mathbf{u}}{\partial \nu} := 2\mu(\text{Def } \mathbf{u})^\# \boldsymbol{\nu} + \lambda(\text{div } \mathbf{u})\boldsymbol{\nu}$, as clearly pointed out from line 18 to line 21 on p. 10166 of [Liu-21], and the author of [Liu-21] wrote “For the derivation of the Navier–Lamé elastic wave equations, its mechanical meaning, and the explanation of the Dirichlet and Neumann boundary conditions, we refer the reader to [Liu-19] for the case of Riemannian manifold and ...”. In [Liu-19], the Neumann boundary condition is $2\mu(\text{Def } \mathbf{u})^\# \boldsymbol{\nu} + \lambda(\text{div } \mathbf{u})\boldsymbol{\nu}$ on $\partial\Omega$. [Liu-19] is an earlier paper which was posed on arXiv on Aug. 14, 2019 by the author (Note that the paper [Liu-21] was submitted to *The Journal Geometric Analysis* on Jul. 21, 2020). In [Liu-21], the Neumann boundary condition is denoted by $\frac{\partial \mathbf{u}}{\partial \nu}$. It is very clear from Lemma 2.1.1 of [Liu-19] that $\frac{\partial \mathbf{u}}{\partial \nu} := 2\mu(\text{Def } \mathbf{u})^\# \boldsymbol{\nu} + \lambda(\text{div } \mathbf{u})\boldsymbol{\nu}$, which equals to $\sum_{j,k=1}^n (\mu(u^j_{;k} + u_k^{;j})\nu^k + \lambda u^k_{;k}\nu^j) \frac{\partial}{\partial x_j}$ (or equivalently, $\lambda n^j \nabla_k u^k + \mu(n^k \nabla_k u^j + n_k \nabla_j u^k)$), where $\{\frac{\partial}{\partial x_j}\}_{j=1}^n$ is the nature coordinate basis. In addition,

by applying the Green's formula to the elastic Lamé system, one can immediately see the corresponding Neumann boundary condition (see also, (12.49) of Exercise 4 on p. 470 in M. Taylor's book [Ta-1]).

Let us point out the following facts (by applying Green's formula to the corresponding partial differential equation (or system of equations)):

i). For the Laplace equation $\Delta u = 0$ in Ω , the Dirichlet boundary condition is $u|_{\partial\Omega}$, and the Neumann boundary condition is $(\boldsymbol{\nu} \cdot \nabla u)|_{\partial\Omega}$, where $\boldsymbol{\nu}$ is the outward normal vector field on $\partial\Omega$;

ii). For the divergence type equation $\nabla \cdot (a(x)\nabla u) = 0$ in Ω , where $a(x) > 0$ on $\bar{\Omega}$, the Dirichlet boundary condition is $u|_{\partial\Omega}$, and the Neumann boundary condition is $(\boldsymbol{\nu} \cdot a(x)\nabla u)|_{\partial\Omega}$;

iii). For the elastic Lamé system $\mu\Delta\mathbf{u} + (\mu + \lambda)\nabla(\nabla \cdot \mathbf{u}) = 0$ in Ω , where Lamé coefficients $\mu > 0$ and $\mu + \lambda \geq 0$ on $\bar{\Omega}$, the Dirichlet boundary condition is $\mathbf{u}|_{\partial\Omega}$, and the Neumann boundary condition is $\mu(\nabla\mathbf{u} + \nabla\mathbf{u}^T)\#\boldsymbol{\nu} + \lambda(\operatorname{div}\mathbf{u})\boldsymbol{\nu}$ on $\partial\Omega$ (see [Liu-19]);

iv). For the Maxwell's equations

$$\begin{cases} \operatorname{curl}\mathbf{E} - i\omega\mu\mathbf{H} = 0 & \text{in } \Omega, \\ \operatorname{curl}\mathbf{H} + i\omega\sigma\mathbf{E} & \text{in } \Omega. \end{cases}$$

where $\omega > 0$ is a fixed frequency; σ is electronic coefficient and μ is the magnetic permeability, the Dirichlet boundary condition is $(\boldsymbol{\nu} \times \mathbf{E})|_{\partial\Omega}$, and the Neumann boundary condition is $(\boldsymbol{\nu} \times \mathbf{H})|_{\partial\Omega}$ (see [Liu-19.2]);

v). For the Stokes flow equations (of a fluid)

$$\begin{cases} \mu\Delta\mathbf{u} + \nabla p = 0 & \text{in } \Omega, \\ \operatorname{div}\mathbf{u} = 0 & \text{in } \Omega, \end{cases}$$

where $\mu > 0$ is the viscosity constant, the Dirichlet boundary condition is $\mathbf{u}|_{\partial\Omega}$, and the Neumann boundary condition is $\mu(\nabla\mathbf{u} + \nabla\mathbf{u}^T)\#\boldsymbol{\nu} + p\boldsymbol{\nu}$ on $\partial\Omega$ (see [Liu-20]).

The boundary conditions depend strongly on the corresponding partial differential equation (or system of equations). They are an indivisible whole. Speaking boundary conditions without any partial differential equation(s) is not a proper statement. In [Liu-21], the author never said $\frac{\partial\mathbf{u}}{\partial\nu}$ to be $\boldsymbol{\nu} \cdot \nabla\mathbf{u}$ as be understood in (1.33) on p. 8 of [CaFrLeVa-22]. Again, in [Liu-21], the Neumann boundary condition for elastic system is $\mu(\nabla\mathbf{u} + \nabla\mathbf{u}^T)\#\boldsymbol{\nu} + \lambda(\operatorname{div}\mathbf{u})\boldsymbol{\nu}$ as pointed out from line 18 to line 21 on p. 10166 (see the paper [Liu-19], from version 1 to version 3).

Response 3. *On p. 26 of [CaFrLeVa-22], from line 20 to line 22, the authors of [CaFrLeVa-22] wrote: "Mathematically the reason for the erroneous expressions comes from the fact that Liu treats the principal symbol of \mathfrak{L} as a diagonal matrix with eigenvalues (1.8) rather than a diagonalisable matrix with these eigenvalues."*

This is not true. Clearly, the authors of [CaFrLeVa-22] gave an irrelevant remark to the proof of [Liu-21]. Obviously, in [Liu-21], the principal symbol of P_g (see, p. 10177–p. 10178 of [Liu-21]) is

$$\mu \left(\sum_{m,l=1}^n g^{ml}\xi_m\xi_l \right) \mathbf{I}_n + (\mu + \lambda) \begin{bmatrix} \sum_{m=1}^n g^{1m}\xi_m\xi_1 & \cdots & \sum_{m=1}^n g^{1m}\xi_m\xi_n \\ \vdots & & \vdots \\ \sum_{m=1}^n g^{nm}\xi_m\xi_1 & \cdots & \sum_{m=1}^n g^{nm}\xi_m\xi_n \end{bmatrix},$$

which is not a diagonal matrix. In [Liu-21], the author of [Liu-21] never treated the principal symbol as being stated by the authors of [CaFrLeVa-22]. Clearly, the authors of [CaFrLeVa-22] misunderstood the main ideas of [Liu-21], and they tried to overthrow Theorem 1.1 of [Liu-21] by their irrelevant remark.

Response 4. From line 3 to line 4 on p. 27 of [CaFrLeVa-22], the authors of [CaFrLeVa-22] wrote “*[Liu-21, formula (1.8)] uses the representation (1.24) in the Riemannian setting in an arbitrary dimension. We recall that the definition of the operator curl is non-trivial in $d > 3$ even in the Euclidean case, and more clarification would be helpful here.*”

In fact, from line 5 to line 17 on p. 10166 of [Liu-21], the author of [Liu-21] wrote “In three spatial dimensions,”. It is clear that equation (1.8) in [Liu-21] is considered in three dimensional case.

Response 5. From line 23 to line 39 on p. 26 of [CaFrLeVa-22], the authors of [CaFrLeVa-22] wrote: “*We note that the expression for \tilde{b}_{Dir} was already found in the 1960 paper by M. Dupuis, R. Mazo, and L. Onsager [DuMaOn-60]. Remarkably, this paper includes the critique of the 1950 paper by E. W. Montroll who presented exactly Liu’s expression (1.34) for the second asymptotic coefficient, modulo some scaling, see [Mo-50, formulae (3)-(5)]. . . . In essence, Liu has rediscovered the seventy-year-old Montroll’s formula which had been known to be incorrect for sixty years.*”

Obviously, the authors of [CaFrLeVa-22] have misunderstood all results in these papers, and have misled the reader by completely wrong statements. In [Mo-50], in order to discuss the effect of the volume and surface area to the heat capacities in low temperature, E. W. Montroll gave the expression for the counting function of eigenvalues for the elastic normal modes of a three-dimensional rectangular solid $\{(x, y, z) \in \mathbb{R}^3 \mid 0 \leq x \leq L_x, 0 \leq y \leq L_y, 0 \leq z \leq L_z\}$ with the boundary condition $\boldsymbol{\nu} \cdot \nabla \mathbf{u}$ (according to the explanation of the boundary condition in [CaFrLeVa-22]). In [DuMaOn-60], M. Dupuis, R. Mazo and L. Onsager investigated an isotropic solid at low temperatures whose model is a rectangular plate of thickness l_3 and other dimensions l_1 and l_2 with realistic boundary conditions. The faces parallel to the plane of the plate are supposed to be free of stresses, whereas the periodic boundary conditions are given on the other faces (see, p. 1453 of [DuMaOn-60]). Clearly, these boundary conditions are completely different from the (whole) Dirichlet or Neumann (i.e., traction) boundary conditions in [Liu-21]. In [Mo-50] and [DuMaOn-60], the authors respectively calculated the counting functions with different boundary conditions for a very special domain (i.e., three-dimensional rectangular solid) by elementary calculations. But, in [Liu-21], we proved the asymptotic formulas for the heat traces of an elastic body for general compact smooth manifold with smooth boundary (for the corresponding Dirichlet and the Neumann (i.e., traction) boundary conditions, respectively). The domains and the boundary conditions in three papers are quite different.

Response 6. On p. 27 of [CaFrLeVa-22], the authors of [CaFrLeVa-22] wrote “*The paper does not refer to previous publications [De-12], [DuMaOn-60] or [SaVa-97].*”

Clearly, the following two conjectures are still open:

Conjecture 1: Let Ω be a compact, connected smooth Riemannian n -manifold with smooth boundary $\partial\Omega$. The following two-term asymptotics hold:

$$(3.1) \quad \begin{aligned} \mathcal{N}_{\mp}(\Lambda) &\sim \frac{1}{\Gamma(1 + \frac{n}{2})} \left(\frac{n-1}{(4\pi\mu)^{n/2}} + \frac{1}{(4\pi(\lambda + 2\mu))^{n/2}} \right) \text{vol}_n(\Omega) \Lambda^{n/2} \\ &\mp \frac{1}{4\Gamma(1 + \frac{n-1}{2})} \left(\frac{n-1}{(4\pi\mu)^{(n-1)/2}} + \frac{1}{(4\pi(\lambda + 2\mu))^{(n-1)/2}} \right) \text{vol}_{n-1}(\partial\Omega) \Lambda^{(n-1)/2} \quad \text{as } \Lambda \rightarrow +\infty, \end{aligned}$$

where $\mathcal{N}_-(\Lambda) := \#\{k \mid \tau_k^- \leq \Lambda\}$ (respectively, $\mathcal{N}_+(\Lambda) := \#\{k \mid \tau_k^+ \leq \Lambda\}$) is the counting function of elastic eigenvalues for the Dirichlet (respectively, Neumann) boundary condition.

When $\lambda + \mu = 0$, Conjecture 1 just is the famous Weyl conjecture, which has about 110-year history and remains open.

Conjecture 2: Let Ω be a compact, connected smooth Riemannian n -manifold with smooth boundary $\partial\Omega$. Suppose that the corresponding elastic billiards is neither dead-end nor absolutely periodic for the Ω . The above two-term asymptotics (3.1) hold.

Obviously, for any given compact smooth Riemannian manifold Ω with smooth boundary, under the elastic billiards condition, an explicit two-term asymptotic expansion of the counting function $\mathcal{N}_{\mp}(\Lambda)$ as $\Lambda \rightarrow +\infty$ had not been given in [SaVa-97]. In [De-12] and [DuMaOn-60], the counting functions of elastic eigenvalues were explicitly calculated only for very special domain (a three-dimensional rectangular solid) with other boundary conditions. However, in Theorem 1.1 of [Liu-21], our two-term asymptotic formulas of the heat trace hold for any given compact connected smooth Riemannian manifold Ω with smooth boundary (do not need any additional assumption).

Similar to Ivrii's method in [Iv-80] (or [Iv-16]) and Hörmander's technique in [Ho-85], it is not difficult to prove the above Conjecture 2. In this new approach which is completely different from [SaVa-97] and [CaFrLeVa-22], the spectral measure and its Fourier transform will be introduced; the fundamental solution $\mathbf{u}(t, x, y)$ of the elastic wave equations with suitable boundary conditions will be considered; the asymptotic expansion of the normal singularity of $\sigma(t) = \int_{\Omega} \text{Tr}(\mathbf{u}(t, x, x)) dx$ as well as $\sigma_{\zeta}(t) = \int_{\Omega} \text{Tr}(\zeta(x)\mathbf{u}(t, x, x)) dx$ will be estimated and explicitly calculated; and the branching Hamiltonian billiards condition on Riemannian manifold will be well used.

However, not every manifold satisfies the billiards condition. For example, when $\lambda + \mu = 0$ (the corresponding elastic operator becomes the Laplacian), the semi-sphere $\mathbb{S}_+^{n-1} := \{(x_1, \dots, x_{n-1}, x_n) \in \mathbb{R}^n | x_1^2 + \dots + x_{n-1}^2 + x_n^2 = 1, x_n \geq 0\}$ does not satisfy the billiards condition because every point $(x, \xi) \in S^*(\mathbb{S}_+^{n-1})$ is periodic (Here $S^*(\mathbb{S}_+^{n-1})$ is the fiber of cotangent unit sphere over \mathbb{S}_+^{n-1}).

ACKNOWLEDGEMENTS

This research was supported by NNSF of China (11671033/A010802) and NNSF of China (11171023/A010801).

REFERENCES

- [Liu-21] G. Q. Liu, *Geometric Invariants of Spectrum of the Navier–Lamé Operator*, J. Geom. Anal. 31 (2021), 10164–10193.
- [Liu-19] G. Q. Liu, *Determination of isometric real-analytic metric and spectral invariants for elastic Dirichlet–Neumann map on Riemannian manifolds*, [arXiv:1908.05096](#).
- [LiuTan-22] G. Q. Liu, X. Tan, *Asymptotic Expansions of the Traces of the Thermoelastic Operators*, [arXiv:2205.13238](#).
- [Liu-19.2] G. Q. Liu, *Determining anisotropic real-analytic metric from boundary electromagnetic information*, [arXiv:1909.12803](#)
- [Liu-20] G. Q. Liu, *Determining the viscosity from the boundary information for incompressible fluid*, [arXiv:2006.04310](#)
- [CaFrLeVa-22] M. Capoferri, L. Friedlander, M. Levitin, D. Vassiliev, *Two-term spectral asymptotics in linear elasticity*, [arXiv:2207.13636](#).
- [Ho-85] L. Hörmander, *The Analysis of Partial Differential Operators IV*, Springer-Verlag, Berlin, Heidelberg, New York (1985, 2009).
- [Kac-66] M. Kac, *Can one hear the shape of a drum?*, Amer. Math. Monthly (Slaught Mem. Papers, no. 11), 73(4) (1966), 1–23.
- [Iv-80] V. Ivrii, *Second term of the spectral asymptotic expansion of the Laplace–Beltrami operator on manifolds with boundary*, Funktsional. Anal. i Prilozhen. 14 (1980), no. 2, 25–34; English transl., Functional Anal. Appl. 14 (1980), 98–106.
- [Iv-16] V. Ivrii, *100 years of Weyl's law*, Bull. Math. Sci. 6 (2016), 379–452.
- [Lee-13] J. M. Lee, *Introduction to smooth manifolds*, Springer Science+Business Media, New York, 2013.
- [MS-67] H. McKean, I. M. Singer, *Curvature and the eigenvalues of the Laplacian*, J. Differential Geometry, 1(1967), 43–69.
- [Ta-1] M. E. Taylor, *Partial Differential Equations I*, 2nd Edition, Appl. Math. Sci., vol. 115, Springer Science+Business Media, LLC 1996, 2011.

- [SaVa-97] Yu. Safarov and D. Vassiliev, *The asymptotic distribution of eigenvalues of partial differential operators*, Amer. Math. Soc., Providence, RI, 1997. DOI: 10.1090/mmono/155. (Chapter VI Mechanical Applications of this book is freely available online at nms.kcl.ac.uk/yuri.safarov/Book/chapter6.pdf).
- [Mo-50] E. W. Montroll, *Size effect in low temperature heat capacities*, J. Chem. Phys. 18:2, 183–185 (1950).
- [DuMaOn-60] M. Dupuis, R. Mazo, and L. Onsager, *Surface specific heat of an isotropic solid at low temperatures*, J. Chem. Phys. 33:5 (1960), 1452–1461.

SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 100081, CHINA
Email address: liugqz@bit.edu.cn