

# Discrete time crystal enabled by Stark many-body localization

Shuo Liu,<sup>1,2</sup> Shi-Xin Zhang,<sup>2,\*</sup> Chang-Yu Hsieh,<sup>2</sup> Shengyu Zhang,<sup>2</sup> and Hong Yao<sup>1,†</sup>

<sup>1</sup>*Institute for Advanced Study, Tsinghua University, Beijing 100084, China*

<sup>2</sup>*Tencent Quantum Laboratory, Tencent, Shenzhen, Guangdong 518057, China*

(Dated: August 8, 2022)

Discrete time crystal (DTC) has recently attracted increasing attention, but most DTC models and their properties are only revealed after disorder average. In this Letter, we propose a simple disorder-free periodically driven model that exhibits nontrivial DTC order stabilized by Stark many-body localization (MBL). We demonstrate the existence of DTC phase by analytical analysis from perturbation theory and convincing numerical evidence from observable dynamics. The new DTC model paves a new promising way for further experiments and deepens our understanding of DTC. Since the DTC order doesn't require special quantum state preparation and the strong disorder average, it can be naturally realized on the noisy intermediate-scale quantum (NISQ) hardware with much fewer resources and repetitions. Moreover, besides the robust subharmonic response, there are other novel robust beating oscillations in Stark-MBL DTC phase which are absent in random or quasi-periodic MBL DTC.

**Introduction:** Spontaneous symmetry breaking (SSB) is one of the most important concepts in modern physics. Phases of matter and phase transitions can be described by SSB mechanism, for example, the formation of crystals is the result of spontaneously breaking continuous spatial translational symmetry. Inspired by this notion, Wilczek proposed the intriguing concept of “time crystal” which spontaneously breaks continuous time translational symmetry [1–3] and various no-go theorems [4–6] have established since then that the continuous time crystal would not exist. However, Floquet systems, quantum systems subject to periodic driving, can indeed exhibit discrete time translational symmetry breaking (DTTSB) [7–10] and have attracted considerable research interest [7, 8, 11–19]. The given observable in the DTC phase can develop persistent oscillations whose period is an integer multiple of the driving period. Recently, DTC has been experimentally realized in programmable quantum devices with periodic driving [20–24].

Due to the existence of periodic driving, energy is no longer conserved in a Floquet system. Thus, in the absence of any other local conservation laws, a generic system will absorb energy from the periodic driving, ultimately heating to infinite temperature. The thermalization of many-body Floquet systems implies that any local physical observable becomes featureless at late times [25–27]. Therefore, strong disorder is required [7–10, 27–30] to realize MBL that exhibits emergent local integrals of motion [31, 32] and prevents absorption of heat from periodic driving. However, to investigate the DTC behavior, we have to average the observable dynamics over a great number of different disorder instances, requiring more quantum resources and severely restricting the efficient experimental study of DTC.

Besides DTC stabilized by MBL phase, so-called prethermal DTC phase exists without the need of MBL. Under some conditions, the dynamics of the many-body

Floquet system can be thought of as being generated by an effective time-independent “prethermal Hamiltonian”,  $H_{\text{eff}}$ . The Floquet system can then display DTC dynamics upon starting from certain low-temperature symmetry-breaking initial states of  $H_{\text{eff}}$  within an exponential heating time window [33–37], realizing prethermal DTC [11, 38–42].

Recently, in kicked PXP model [43, 44], the discrete time crystal order enabled by quantum many-body scars [45–48] with Néel state as the initial state had been identified, which is strongly reminiscent of prethermal DTC. The fidelity  $F_n = |\langle Z_2 | U_F^n | Z_2 \rangle|^2$  is used to characterize the dynamics in this model, where  $|Z_2\rangle$  is the initial Néel state and  $U_F$  is the Floquet evolution unitary. When  $n$  is even,  $F_n > 0$  and when  $n$  is odd,  $F_n = 0$ , this corresponds to the subharmonic response with a timescale  $T_s = 2$ . After a long enough time, the fidelity will decay to zero finally and stay featureless, and this corresponds to the prethermal timescale  $T_p$ . Between the two timescales, there are another two novel timescales that are not reported in DTC or prethermal DTC phases before. One is the emergent beating timescale  $T_b$ , the fidelity at even periods exhibits a beating oscillation which comes from the overlap between the Néel initial state and the lowest lying excited states of  $H_{\text{eff}}$ . The beating timescale  $T_b \propto \Delta^{-1}$  where  $\Delta$  is the gap in the Floquet spectrum. The other is the timescale  $T_g$  that is set by the inverse energy splitting in the ground state manifold and  $T_g \propto e^N$  where  $N$  is the system size. After driving cycles in the order of  $T_g$ , the fidelity at even periods  $F_{2n}$  decreases, while simultaneously the fidelity at odd periods  $F_{2n+1}$  increases. However, different from prethermal DTC phases, these phenomena strongly depend on special Néel initial states where highly accurate quantum state preparation is required.

An extremely important and exciting direction is to identify a clean Floquet system, i.e. without strong disorder, that exhibits a nontrivial DTC phase with no de-

pendence on initial states. To stabilize this intrinsically dynamical phase, MBL is extremely important as discussed above. It has long been established that the quantum systems may enter MBL phases in the presence of sufficiently strong random disorder [31, 32, 49–57], quasi-periodic potential [58–63] or linear Zeeman field [64–67] in one-dimensional (1D) systems. The third one is called Stark MBL [64–67]. By intuition, we may construct a clean many-body Floquet system utilizing Stark MBL to stabilize DTC order.

In this Letter, we propose a clean kicked Floquet model inspired by Stark MBL. It has various nontrivial and interesting properties, including robust subharmonic response as conventional DTC and other novel timescales similar to those reported in kicked PXP model and Rydberg atom experiments [43, 44]. The subharmonic response is robust against imperfection and doesn't depend on special initial states, which is a signal for nontrivial DTC phase. Since the existence of DTC order doesn't depend on the strong disorder, it can be naturally realized on the quantum hardware relying on fewer quantum computational resources and experimental trials.

**Model:** The Floquet unitary for the model considered in the Letter reads:

$$U_F = U_2 U_1 = e^{-iH_2} e^{-i(\frac{\pi}{2} - \epsilon)H_1}, \quad (1)$$

where

$$H_1 = \sum_j X_j, \quad (2)$$

$$H_2 = J_z \sum_j (j+1) Z_j Z_{j+1} + W \sum_j j Z_j. \quad (3)$$

$X$  and  $Z$  are Pauli matrices,  $\epsilon$  is the imperfection in the driving.  $H_1$  is the kicked term, when  $\epsilon = 0$ ,  $U_1 = \prod_j X_j$  and all the spins flip exactly;  $H_2$  has two terms: one is a linear Zeeman field for Stark MBL; the other one is a linear  $zz$  interaction. The linear term for interaction is important to stabilize DTC similar to the case discussed in [22] (see also in the Supplemental Materials). The reason is that the linear Zeeman field, as well as the Stark MBL, will be suppressed by the imperfection  $\epsilon$ . Therefore we need some nonuniform interaction to stabilize the MBL phase. Different from the random MBL DTC case where a strong disorder in  $zz$  interaction is added, we here instead also use a linear  $zz$  interaction.

When  $\epsilon = 0$ , the quasi-eigenstates of  $U_F^0$  can be written as:

$$|\pm\rangle = \frac{1}{\sqrt{2}}(e^{-i\frac{H_2(z)}{2}}|z\rangle \pm e^{-i\frac{H_2(-z)}{2}}|-z\rangle), \quad (4)$$

whose eigenvalues are,

$$U_F^0|\pm\rangle = \pm e^{-i\frac{H_2(z)+H_2(-z)}{2}}|\pm\rangle \quad (5)$$

respectively, where  $|z\rangle$  is the product state and  $|-z\rangle = \prod_j X_j|z\rangle$ .  $|+\rangle$  and  $|-\rangle$  form a so-called  $\pi$ -pair in which

quasi-eigenenergy difference equals  $\pi$ . For simplicity, we use quasi-eigenenergy  $\epsilon_F$  of  $|+\rangle$  to represent this  $\pi$ -pair. And for any product state  $|z\rangle$ , it can be represented by a superposition of a  $\pi$ -pair of quasi-eigenstates,

$$|\pm z\rangle = \frac{1}{\sqrt{2}}e^{i\frac{H_2(\pm z)}{2}}(|+\rangle \pm |-\rangle). \quad (6)$$

Therefore, there is a trivial subharmonic response in  $\epsilon = 0$  limit,

$$U_F^2|z\rangle = U_F| - z\rangle = |z\rangle, \quad (7)$$

where we have ignored the global phase. Accordingly, the local physical observables, such as  $\langle Z(t) \rangle$ , develop persistent oscillations whose periods are twice as the driving period, and the discrete time translational symmetry spontaneously breaks. However, this DTC order depends on fine-tuning of parameters  $\epsilon = 0$ . To establish a nontrivial DTC phase, the subharmonic response must be robust against imperfection  $\epsilon$ . When  $\epsilon \neq 0$ , the quasi-eigenstates can not be analytically exactly tracked, we use perturbation theory and numerical results below to show that the subharmonic response is robust against imperfection  $\epsilon$ .

**Observable:** To describe the dynamics of the kicked model and diagnose the DTC phase, we need to utilize suitable observable. Due to the linear Zeeman field and linear  $zz$  interaction, different spatial sites are not equivalent anymore. Therefore, we don't choose the non-equal time spin-spin correlation on site  $N/2$  commonly used in previous DTC works [12] and instead use more representative site-averaged observables. Spurred by [44], we define two types of fidelity, one is the fidelity for a given initial state:

$$F_n = |\langle \psi_i | U_F^n | \psi_i \rangle|^2, \quad (8)$$

where  $|\psi_i\rangle$  is the initial state. The other is the state-averaged fidelity

$$\bar{F}_n = \frac{1}{2^N} \sum_{\{z\}} |\langle z | U_F^n | z \rangle|^2, \quad (9)$$

where the sum is over all possible product states  $|z\rangle$ . We can also utilize site-averaged spin-spin correlation  $\frac{1}{N} \sum_j \text{Tr}(Z_j(2nT)Z_j(0))$  as the observable, and the conclusions are qualitatively the same.

**Analysis of different timescales:** We observe three different timescales for our Floquet model which are similar to those reported in kicked PXP model. Because the third timescale  $T_g \propto e^N$  is exponential with the system size, we show the dynamics of a small system (3 sites) firstly to demonstrate all the three timescales in the dynamics. The results are summarized in Fig. 1.

To understand these three timescales, we consider a product state  $|z\rangle$  and the corresponding state-dependent fidelity firstly. When  $\epsilon \neq 0$ , although the product state




FIG. 1. There are three timescales for the dynamics of the state-averaged fidelity  $\bar{F}_n$ : when  $n$  is even,  $\bar{F}_n > 0$  and when  $n$  is odd,  $\bar{F}_n = 0$ , which correspond to the subharmonic response  $T_s = 2$ ; besides this subharmonic response, there are a beating timescale  $T_b \approx 66$  and a third timescale  $T_g \approx 16000$  ( $N = 3$ ;  $J_z = \frac{\pi}{2N}$ ;  $W = 5.0$ ;  $\epsilon = 0.05$ ).

$|z\rangle$  can not be represented by a superposition of one  $\pi$ -pair of quasi-eigenstates, we can still do decomposition in the eigenspace of  $U_F$ . Based on the perturbation theory, the quasi-eigenstate of  $U_F$  ( $\epsilon \neq 0$ ) can be written as a superposition of the quasi-eigenstates of  $U_F^0$  ( $\epsilon = 0$ ) with the similar quasi-eigenenergies. For a given product state  $|z\rangle$ , the corresponding original  $\pi$ -pair of quasi-eigenstates and the most related quasi-eigenstates to the first-order perturbation form a Hilbert subspace, and  $|z\rangle$  roughly live in this subspace.

We utilize perturbation theory to locate the subspace where the product state  $|z\rangle$  lives. We use  $|\pm\rangle$  to represent the original  $\pi$ -pair related to  $|z\rangle$  when  $\epsilon = 0$  and the quasi-eigenenergy is  $\varepsilon_F$ , i.e.  $U_F^0|\pm\rangle = \pm e^{-i\varepsilon_F}|\pm\rangle$ . By intuition, the dimension of the subspace is determined by the number of quasi-eigenstates of  $U_F^0$  which have the similar quasi-eigenenergies with  $|\pm\rangle$ . In the Supplemental Material, we show that if there is only one  $\pi$ -pair of quasi-eigenstates  $|\pm'\rangle$  with  $\varepsilon'_F = \varepsilon_F$  or  $\varepsilon_F + \pi \pmod{2\pi}$ ,  $|\pm\rangle$  and  $|\pm'\rangle$  form a subspace with dimension of 4 and  $|z\rangle$  roughly lives in this subspace. Equivalently, if we check the overlaps between  $|z\rangle$  and quasi-eigenstates of  $U_F$ , there is an obvious dominant-subleading  $\pi$ -pair pattern, see Fig. 2(a). Even if there is no exactly matching quasi-eigenstates of  $U_F^0$ , as long as there is one special  $\pi$ -pair of quasi-eigenstates of  $U_F^0$  which has the closer quasi-eigenenergy with  $|\pm\rangle$  than all other eigenstates, the dominant-subleading  $\pi$ -pair pattern for the decomposition of  $|z\rangle$  still exists. On the contrary, if there are several  $\pi$ -pairs of  $U_F^0$  have the same closest quasi-eigenenergy difference with  $|\pm\rangle$  as  $\delta\varepsilon_F = |\varepsilon_F - \varepsilon'_F| = \delta$ , the dominant-subleading  $\pi$ -pair pattern vanishes and we can only see one  $\pi$ -pair (the original one) with dominant overlap, see Fig. 2(b).

When  $\epsilon = 0$ , the product state  $|z\rangle$  can be represented by an equal weight superposition of a  $\pi$ -pair of quasi-eigenstates of  $U_F^0$  and this induces the subharmonic response as discussed above. When  $\epsilon$  is small, as long as the product state  $|z\rangle$  can still be represented by a su-




FIG. 2. Overlaps with quasi-eigenstates of  $U_F$ :  $N = 9$ ;  $J_z = \frac{\pi}{2N}$ ;  $W = 5.0$ ;  $\epsilon = 0.05$ . (a), the product state  $|0000000000\rangle$  roughly lives in a subspace with dimension of 4 and we can see the obvious dominant-subleading  $\pi$ -pair pattern; (b), there is only one obvious  $\pi$ -pair in the decomposition of the product state  $|011001100\rangle$ .

perposition of several  $\pi$ -pairs of quasi-eigenstates of  $U_F$  and the weights of the two quasi-eigenstates in any  $\pi$ -pair have the same absolute value, as guaranteed by the perturbation theory, the  $T_s = 2$  subharmonic response still exists.

The beating timescale  $T_b$  is caused by the quasi-eigenenergy difference between different quasi-eigenstates (see the Supplemental Material for details). Consider a three-site subsystem of a product state  $|z\rangle$ ,  $|S_{j-1}, S_j, S_{j+1}\rangle$ ,  $|z\rangle$  and  $| - z\rangle$  ( $| - S_{j-1}, -S_j, -S_{j+1}\rangle$ ) can be combined into a  $\pi$ -pair with quasi-eigenenergy equals  $\frac{H_2(z)+H_2(-z)}{2}$  when  $\epsilon = 0$ . If we flip spin  $S_j$  of  $|z\rangle$  and  $| - z\rangle$ , there are two new product states  $|z'\rangle$  ( $|S_{j-1}, -S_j, S_{j+1}\rangle$ ) and  $| - z'\rangle$  ( $| - S_{j-1}, S_j, -S_{j+1}\rangle$ ). And they form a new  $\pi$ -pair  $|\pm'\rangle$  with quasi-eigenenergy  $\varepsilon'_F$ . The quasi-eigenenergy difference between the two  $\pi$ -pairs is

$$\delta\varepsilon_F = 2J_z |jS_{j-1} + (j+1)S_{j+1}| \pmod{2\pi}. \quad (10)$$

When  $S_{j-1} = -S_{j+1}$ , the quasi-eigenenergy difference after flipping the spin  $S_j$  equals  $2J_z$ ; when  $S_{j-1} = S_{j+1}$ , the quasi-eigenenergy difference after flipping the spin  $S_j$  equals  $2(2j+1)J_z$ . Suppose  $J_z = \frac{\pi}{2N}$  and assume  $N$  be odd for simplicity, considering the spin configuration around a fixed site  $\frac{N-1}{2}$  (the middle site), the product states can be divided into two parts: when  $S_{\frac{N-1}{2}-1} = S_{\frac{N-1}{2}+1}$ , as discussed above, there are two  $\pi$ -pairs related by a Pauli-X matrix at site  $\frac{N-1}{2}$  and the quasi-eigenenergy difference is equal to  $\pi$ . And the quasi-eigenstates decomposition of the product state includes a dominant  $\pi$ -pair and a subleading  $\pi$ -pair, see Fig. 2(a). We call these product states “good initial states”. The dynamics of these good states have a dominant beating timescale  $T_b$  determined by the quasi-eigenenergy difference between dominant and subleading  $\pi$ -pairs and  $T_b$  fits well with the perturbative predictions (see the Supplemental Material for details), see Fig. 3. When  $S_{\frac{N-1}{2}-1} = -S_{\frac{N-1}{2}+1}$ , there is no  $\pi$ -pair  $|\pm'\rangle$  with quasi-eigenenergy difference  $\delta\varepsilon_F = 0$  or  $\pi$  with  $|\pm\rangle$ . As shown




FIG. 3. Dynamics of the product state  $|000000000\rangle$  which is a “good initial state” ( $N = 9; W = 5.0; \epsilon = 0.05$ ). In (a),(b),  $J_z = \frac{\pi}{2N}$  and in (c),(d),  $J_z = \frac{\pi-0.05}{2N}$ . In both cases, there is a dominant beating timescale.




FIG. 4. (a),(b): dynamics of the product state  $|011001100\rangle$  ( $N = 9; J_z = \frac{\pi}{2N}; W = 5.0; \epsilon = 0.05$ ). It is a bad initial state and there is no dominant beating timescale. (c),(d): dynamics of the state-averaged fidelity  $\bar{F}_n$  ( $N = 9; J_z = \frac{\pi}{2N}; W = 5.0; \epsilon = 0.05$ ). There is a dominant beating timescale  $T_b = \frac{2\pi}{\omega_b}$  which is caused by the “good initial states”. We can also see many other timescales caused by the “bad initial states”.

in Fig. 2(b), there is no dominant-subleading  $\pi$ -pair pattern. So there is no dominant beating oscillation in the dynamics, see Fig. 4(a). We call these product states “bad initial states”. Although there is no obvious subleading  $\pi$ -pair, there are many  $\pi$ -pairs with small overlaps and different quasi-eigenenergy differences with  $|\pm\rangle$ , thus we can see many Fourier peaks in Fig. 4(b).

Now we investigate the robustness of the beating timescale by considering non-perfect  $J_z$ , for example,  $J_z = \frac{\pi-0.05}{2N}$ . For a  $\pi$ -pair of quasi-eigenstates of  $U_F^0$  combined by “good initial states”, although there is no other

$\pi$ -pairs of quasi-eigenstates of  $U_F^0$  with  $\delta\varepsilon_F = 0$  or  $\pi$ , as long as only one  $\pi$ -pair of quasi-eigenstates has the more closer quasi-eigenenergy with  $|\pm\rangle$  than all other quasi-eigenstates, the dominant-subleading  $\pi$ -pair pattern still exists and there is also a dominant beating timescale, see Fig. 3(d).

We can use a more general state-averaged observable, the state-averaged fidelity, to describe the dynamics of the many-body Floquet system. The quasi-eigenenergy corrections due to the first-order perturbation to all “good initial states” are the same, in the case considered here,  $T_b \approx \frac{\pi}{\epsilon \sin(\frac{\pi(N-1)W}{2})}$  (see the Supplemental Material for details), so the dominant beating timescale still exists, see Fig. 4(d). Additionally, there is no scaling relation between  $T_b$  and the system size  $N$ . It needs to be emphasized that the classification of “good initial state” and “bad initial state” is for the beating timescale  $T_b$ , all the initial product states show a robust subharmonic response  $T_s = 2$  and support spontaneously discrete time translational symmetry breaking.

In terms of the timescale  $T_g$ , it is induced by the tiny quasi-energy splitting of a given  $\pi$ -pair, i.e. the quasi-eigenenergy difference between  $|+\rangle$  and  $|-\rangle$  equals  $\pi + \delta$ . This quasi-energy mismatch  $\delta$  due to finite size effect induces the third timescale as  $T_g \propto \frac{1}{\delta}$  and  $\delta \propto e^{-N}$  (see the Supplemental Material for details). As discussed in [68], the existence of the third timescale  $T_g$  depends on the order of the two limits: (a),  $\lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty}$  and (b),  $\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty}$ . (a) characterizes the “intrinsic” quench dynamics of this phase. In (a), we will never reach times of  $O(e^N)$  and the third oscillation timescale  $T_g$  ( $\propto e^N$ ) vanishes. And we can only observe the subharmonic response and beating oscillation out to  $t \rightarrow \infty$ .

**Phase transition:** To stabilize DTC phase, MBL is extremely important as discussed above. We can see the distribution of level spacing ratio gradually crosses from the Poisson limit to the Gaussian orthogonal ensemble (GOE) type with increasing imperfection  $\epsilon$ , see Fig. 5 for level spacing distribution (see the Supplemental Material for details), which means there is a phase transition from MBL phase to trivial thermal phase.

To diagnose the phase transition and discrete time translational symmetry breaking, we utilize two indicators: the magnitude of the subharmonic response and the mutual information (see the Supplemental Material for details). We compute the mutual information between the first and last site of the spin chain as a function of  $\epsilon$  for fixed  $J_z$  and  $W$ . As discussed in [12], we perform a numerical exploration of the discrete time translational symmetry breaking to get the critical  $\epsilon_c$  for fixed  $J_z$  and  $W$ , see Fig. 6 for the schematic phase diagram consisting the new DTC phase and trivial thermal phase.

**Conclusion:** We have demonstrated that the discrete time crystal can be realized in a clean kicked Floquet model stabilized by Stark MBL. The magnitude of the




FIG. 5. Histogram of the distribution of the level spacing ratios:  $N = 13$ ,  $J_z = \pi + 1.5$ ,  $W = 5.0$ . (a),  $\epsilon = 0.05$ ; (b),  $\epsilon = 0.5$ . With increasing the imperfection  $\epsilon$ , the distribution of level spacing ratio gradually crosses from the Poisson limit to the GOE distribution.




FIG. 6. The schematic phase diagram ( $W = 5.0$ ): when  $J_z = n\pi$ , the  $zz$  interaction has no effect on the Floquet evolution and the system enters trivial thermalized phase.

subharmonic oscillation peak and mutual information of quasi-eigenstates provide convincing evidence of the DTTSB. We also utilize the perturbation theory to explain the novel beating timescale absent in conventional DTC. Compared to the conventional DTC stabilized by the strong disorder, the resources required in our model are much fewer and it can be easily realized on the quantum hardware [69] in the NISQ era [70] (see Supplemental Materials for detailed experimental proposals).

**Acknowledgements:** We thank Zhou-Quan Wan for helpful discussions. This work is supported in part by the NSFC under Grant No. 11825404 (SXZ, SL, and HY), the CAS Strategic Priority Research Program under Grant No. XDB28000000 (HY), and Beijing Municipal Science and Technology Commission under Grant No. Z181100004218001 (HY).

\* shixinzhang@tencent.com

† yaohong@tsinghua.edu.cn

[1] F. Wilczek, Quantum time crystals, *Phys. Rev. Lett.* **109**, 160401 (2012).

- [2] T. Li, Z.-X. Gong, Z.-Q. Yin, H. T. Quan, X. Yin, P. Zhang, L.-M. Duan, and X. Zhang, Space-time crystals of trapped ions, *Phys. Rev. Lett.* **109**, 163001 (2012).
- [3] F. Wilczek, Superfluidity and space-time translation symmetry breaking, *Phys. Rev. Lett.* **111**, 250402 (2013).
- [4] P. Bruno, Impossibility of spontaneously rotating time crystals: A no-go theorem, *Phys. Rev. Lett.* **111**, 070402 (2013).
- [5] P. Nozières, Time crystals: can diamagnetic currents drive a charge density wave into rotation?, *EPL (Europhysics Letters)* **103**, 57008 (2013).
- [6] H. Watanabe and M. Oshikawa, Absence of quantum time crystals, *Phys. Rev. Lett.* **114**, 251603 (2015).
- [7] V. Khemani, A. Lazarides, R. Moessner, and S. L. Sondhi, Phase structure of driven quantum systems, *Phys. Rev. Lett.* **116**, 250401 (2016).
- [8] D. V. Else, B. Bauer, and C. Nayak, Floquet time crystals, *Phys. Rev. Lett.* **117**, 090402 (2016).
- [9] C. W. von Keyserlingk and S. L. Sondhi, Phase structure of one-dimensional interacting floquet systems. ii. symmetry-broken phases, *Phys. Rev. B* **93**, 245146 (2016).
- [10] C. W. von Keyserlingk, V. Khemani, and S. L. Sondhi, Absolute stability and spatiotemporal long-range order in floquet systems, *Phys. Rev. B* **94**, 085112 (2016).
- [11] K. Sacha, Modeling spontaneous breaking of time-translation symmetry, *Phys. Rev. A* **91**, 033617 (2015).
- [12] N. Y. Yao, A. C. Potter, I.-D. Potirniche, and A. Vishwanath, Discrete time crystals: Rigidity, criticality, and realizations, *Phys. Rev. Lett.* **118**, 030401 (2017).
- [13] A. Russomanno, F. Iemini, M. Dalmonte, and R. Fazio, Floquet time crystal in the lipkin-meshkov-glick model, *Phys. Rev. B* **95**, 214307 (2017).
- [14] Z. Gong, R. Hamazaki, and M. Ueda, Discrete time-crystalline order in cavity and circuit qed systems, *Phys. Rev. Lett.* **120**, 040404 (2018).
- [15] B. Huang, Y.-H. Wu, and W. V. Liu, Clean floquet time crystals: Models and realizations in cold atoms, *Phys. Rev. Lett.* **120**, 110603 (2018).
- [16] B. Zhu, J. Marino, N. Y. Yao, M. D. Lukin, and E. A. Demler, Dicke time crystals in driven-dissipative quantum many-body systems, *New Journal of Physics* **21**, 073028 (2019).
- [17] V. K. Kozin and O. Kyriienko, Quantum time crystals from hamiltonians with long-range interactions, *Phys. Rev. Lett.* **123**, 210602 (2019).
- [18] K. Chinzei and T. N. Ikeda, Time crystals protected by floquet dynamical symmetry in hubbard models, *Phys. Rev. Lett.* **125**, 060601 (2020).
- [19] N. Y. Yao, C. Nayak, L. Balents, and M. P. Zaletel, Classical discrete time crystals, *Nature Physics* **16**, 438 (2020).
- [20] S. Choi, J. Choi, R. Landig, G. Kucsko, H. Zhou, J. Isoya, F. Jelezko, S. Onoda, H. Sumiya, V. Khemani, C. von Keyserlingk, N. Y. Yao, E. Demler, and M. D. Lukin, Observation of discrete time-crystalline order in a disordered dipolar many-body system, *Nature* **543**, 221 (2017).
- [21] J. Zhang, P. W. Hess, A. Kyprianidis, P. Becker, A. Lee, J. Smith, G. Pagano, I.-D. Potirniche, A. C. Potter, A. Vishwanath, N. Y. Yao, and C. Monroe, Observation of a discrete time crystal, *Nature* **543**, 217 (2017).
- [22] X. Mi, M. Ippoliti, C. Quintana, A. Greene, Z. Chen, J. Gross, F. Arute, K. Arya, J. Atalaya, R. Bab-

- bush, J. C. Bardin, J. Basso, A. Bengtsson, A. Bilmes, A. Bourassa, L. Brill, M. Broughton, B. B. Buckley, D. A. Buell, B. Burkett, N. Bushnell, B. Chiaro, R. Collins, W. Courtney, D. Debroy, S. Demura, A. R. Derk, A. Dunsworth, D. Eppens, C. Erickson, E. Farhi, A. G. Fowler, B. Foxen, C. Gidney, M. Giustina, M. P. Harrigan, S. D. Harrington, J. Hilton, A. Ho, S. Hong, T. Huang, A. Huff, W. J. Huggins, L. B. Ioffe, S. V. Isakov, J. Iveland, E. Jeffrey, Z. Jiang, C. Jones, D. Kafri, T. Khattar, S. Kim, A. Kitaev, P. V. Klimov, A. N. Korotkov, F. Kostritsa, D. Landhuis, P. Laptev, J. Lee, K. Lee, A. Locharla, E. Lucero, O. Martin, J. R. McClean, T. McCourt, M. McEwen, K. C. Miao, M. Mohseni, S. Montazeri, W. Mruczkiewicz, O. Naaman, M. Neeley, C. Neill, M. Newman, M. Y. Niu, T. E. O'Brien, A. Opremcak, E. Ostby, B. Pato, A. Petukhov, N. C. Rubin, D. Sank, K. J. Satzinger, V. Shvarts, Y. Su, D. Strain, M. Szalay, M. D. Trevithick, B. Villalonga, T. White, Z. J. Yao, P. Yeh, J. Yoo, A. Zalcman, H. Neven, S. Boixo, V. Smelyanskiy, A. Megrant, J. Kelly, Y. Chen, S. L. Sondhi, R. Moessner, K. Kechedzhi, V. Khemani, and P. Roushan, Time-Crystalline Eigenstate Order on a Quantum Processor, *Nature* (2021).
- [23] J. Randall, C. E. Bradley, F. V. van der Grienden, A. Galicia, M. H. Abobeih, M. Markham, D. J. Twitchen, F. Machado, N. Y. Yao, and T. H. Taminiau, Many-body-localized discrete time crystal with a programmable spin-based quantum simulator, *Science* **374**, 1474 (2021).
- [24] X. Zhang, W. Jiang, J. Deng, K. Wang, J. Chen, P. Zhang, W. Ren, H. Dong, S. Xu, Y. Gao, F. Jin, X. Zhu, Q. Guo, H. Li, C. Song, A. V. Gorshkov, T. Iadecola, F. Liu, Z.-X. Gong, Z. Wang, D.-L. Deng, and H. Wang, Digital quantum simulation of Floquet symmetry-protected topological phases, *Nature* **607**, 468 (2022).
- [25] L. D'Alessio and M. Rigol, Long-time behavior of isolated periodically driven interacting lattice systems, *Phys. Rev. X* **4**, 041048 (2014).
- [26] A. Lazarides, A. Das, and R. Moessner, Equilibrium states of generic quantum systems subject to periodic driving, *Phys. Rev. E* **90**, 012110 (2014).
- [27] P. Ponte, A. Chandran, Z. Papić, and D. A. Abanin, Periodically driven ergodic and many-body localized quantum systems, *Annals of Physics* **353**, 196 (2015).
- [28] P. Ponte, Z. Papić, F. Huveneers, and D. A. Abanin, Many-body localization in periodically driven systems, *Physical review letters* **114**, 140401 (2015).
- [29] A. Lazarides, A. Das, and R. Moessner, Fate of many-body localization under periodic driving, *Phys. Rev. Lett.* **115**, 030402 (2015).
- [30] D. A. Abanin, W. D. Roeck, and F. Huveneers, Theory of many-body localization in periodically driven systems, *Annals of Physics* **372**, 1 (2016).
- [31] M. Serbyn, Z. Papić, and D. A. Abanin, Local conservation laws and the structure of the many-body localized states, *Phys. Rev. Lett.* **111**, 127201 (2013).
- [32] D. A. Huse, R. Nandkishore, and V. Oganesyan, Phenomenology of fully many-body-localized systems, *Phys. Rev. B* **90**, 174202 (2014).
- [33] D. A. Abanin, W. De Roeck, and F. Huveneers, Exponentially Slow Heating in Periodically Driven Many-Body Systems, *Physical Review Letters* **115**, 256803 (2015).
- [34] T. Mori, T. Kuwahara, and K. Saito, Rigorous Bound on Energy Absorption and Generic Relaxation in Periodically Driven Quantum Systems, *Physical Review Letters* **116**, 120401 (2016).
- [35] T. Kuwahara, T. Mori, and K. Saito, Floquet-Magnus Theory and Generic Transient Dynamics in Periodically Driven Many-Body Quantum Systems, *Annals of Physics* **367**, 96 (2016).
- [36] D. A. Abanin, W. De Roeck, W. W. Ho, and F. Huveneers, Effective Hamiltonians, prethermalization, and slow energy absorption in periodically driven many-body systems, *Physical Review B* **95**, 014112 (2017).
- [37] D. Abanin, W. De Roeck, W. W. Ho, and F. Huveneers, A Rigorous Theory of Many-Body Prethermalization for Periodically Driven and Closed Quantum Systems, *Communications in Mathematical Physics* **354**, 809 (2017).
- [38] D. V. Else, B. Bauer, and C. Nayak, Prethermal phases of matter protected by time-translation symmetry, *Phys. Rev. X* **7**, 011026 (2017).
- [39] T.-S. Zeng and D. N. Sheng, Prethermal time crystals in a one-dimensional periodically driven floquet system, *Phys. Rev. B* **96**, 094202 (2017).
- [40] D. J. Luitz, R. Moessner, S. L. Sondhi, and V. Khemani, Prethermalization without temperature, *Phys. Rev. X* **10**, 021046 (2020).
- [41] A. Kyprianidis, F. Machado, W. Morong, P. Becker, K. S. Collins, D. V. Else, L. Feng, P. W. Hess, C. Nayak, G. Pagano, N. Y. Yao, and C. Monroe, Observation of a prethermal discrete time crystal, *Science* **372**, 1192 (2021).
- [42] M. Natsheh, A. Gambassi, and A. Mitra, Critical properties of the prethermal floquet time crystal, *Phys. Rev. B* **103**, 224311 (2021).
- [43] D. Bluvstein, A. Omran, H. Levine, A. Keesling, G. Semeghini, S. Ebadi, T. T. Wang, A. A. Michailidis, N. Maskara, W. W. Ho, S. Choi, M. Serbyn, M. Greiner, V. Vuletić, and M. D. Lukin, Controlling quantum many-body dynamics in driven rydberg atom arrays, *Science* **371**, 1355 (2021).
- [44] N. Maskara, A. A. Michailidis, W. W. Ho, D. Bluvstein, S. Choi, M. D. Lukin, and M. Serbyn, Discrete time-crystalline order enabled by quantum many-body scars: entanglement steering via periodic driving, *Physical Review Letters* **127**, 090602 (2021).
- [45] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papić, Weak ergodicity breaking from quantum many-body scars, *Nature Physics* **14**, 745 (2018).
- [46] W. W. Ho, S. Choi, H. Pichler, and M. D. Lukin, Periodic orbits, entanglement, and quantum many-body scars in constrained models: Matrix product state approach, *Phys. Rev. Lett.* **122**, 040603 (2019).
- [47] M. Schechter and T. Iadecola, Weak ergodicity breaking and quantum many-body scars in spin-1  $xy$  magnets, *Phys. Rev. Lett.* **123**, 147201 (2019).
- [48] M. Serbyn, D. A. Abanin, and Z. Papić, Quantum many-body scars and weak breaking of ergodicity, *Nature Physics* **17**, 675 (2021).
- [49] I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, Interacting electrons in disordered wires: Anderson localization and low- $t$  transport, *Phys. Rev. Lett.* **95**, 206603 (2005).
- [50] D. M. Basko, I. L. Aleiner, and B. L. Altshuler, Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states, *Annals of Physics* **321**, 1126 (2006).
- [51] V. Oganesyan and D. A. Huse, Localization of interacting fermions at high temperature, *Phys. Rev. B* **75**, 155111

- (2007).
- [52] M. Znidaric, T. Prosen, and P. Prelovsek, Many-body localization in the heisenberg  $xxz$  magnet in a random field, *Physical Review B* **77**, 064426 (2008).
- [53] C. Monthus and T. Garel, Many-body localization transition in a lattice model of interacting fermions: Statistics of renormalized hoppings in configuration space, *Phys. Rev. B* **81**, 134202 (2010).
- [54] E. Cuevas, M. Feigel'man, L. Ioffe, and M. Mezard, Level statistics of disordered spin-1/2 systems and materials with localized Cooper pairs, *Nature Communications* **3**, 1128 (2012).
- [55] J. H. Bardarson, F. Pollmann, and J. E. Moore, Unbounded growth of entanglement in models of many-body localization, *Phys. Rev. Lett.* **109**, 017202 (2012).
- [56] R. Vosk and E. Altman, Many-body localization in one dimension as a dynamical renormalization group fixed point, *Phys. Rev. Lett.* **110**, 067204 (2013).
- [57] M. Serbyn, Z. Papić, and D. A. Abanin, Universal slow growth of entanglement in interacting strongly disordered systems, *Phys. Rev. Lett.* **110**, 260601 (2013).
- [58] S. Iyer, V. Oganesyan, G. Refael, and D. A. Huse, Many-body localization in a quasiperiodic system, *Phys. Rev. B* **87**, 134202 (2013).
- [59] R. Modak and S. Mukerjee, Many-body localization in the presence of a single-particle mobility edge, *Phys. Rev. Lett.* **115**, 230401 (2015).
- [60] S.-X. Zhang and H. Yao, Universal properties of many-body localization transitions in quasiperiodic systems, *Phys. Rev. Lett.* **121**, 206601 (2018).
- [61] T. Kohlert, S. Scherg, X. Li, H. P. Lüschen, S. Das Sarma, I. Bloch, and M. Aidelsburger, Observation of many-body localization in a one-dimensional system with a single-particle mobility edge, *Phys. Rev. Lett.* **122**, 170403 (2019).
- [62] S.-X. Zhang and H. Yao, Strong and Weak Many-Body Localizations, *arXiv:1906.00971* (2019).
- [63] S. Ghosh, J. Gidugu, and S. Mukerjee, Transport in the nonergodic extended phase of interacting quasiperiodic systems, *Phys. Rev. B* **102**, 224203 (2020).
- [64] M. Schulz, C. A. Hooley, R. Moessner, and F. Pollmann, Stark many-body localization, *Phys. Rev. Lett.* **122**, 040606 (2019).
- [65] E. V. H. Doggen, I. V. Gornyi, and D. G. Polyakov, Stark many-body localization: Evidence for hilbert-space shattering, *Phys. Rev. B* **103**, L100202 (2021).
- [66] E. P. L. van Nieuwenburg, Y. Baum, and G. Refael, From Bloch Oscillations to Many Body Localization in Clean Interacting Systems, *Proceedings of the National Academy of Sciences* **116**, 9269 (2019).
- [67] V. Khemani, M. Hermele, and R. M. Nandkishore, Localization from Hilbert space shattering: from theory to physical realizations, *Physical Review B* **101**, 174204 (2020).
- [68] V. Khemani, R. Moessner, and S. L. Sondhi, A Brief History of Time Crystals, *arXiv:1910.10745* (2019).
- [69] F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, R. Barends, R. Biswas, S. Boixo, F. G. S. L. Brandao, D. A. Buell, B. Burkett, Y. Chen, Z. Chen, B. Chiaro, R. Collins, W. Courtney, A. Dunsworth, E. Farhi, B. Foxen, A. Fowler, C. Gidney, M. Giustina, R. Graff, K. Guerin, S. Habegger, M. P. Harrigan, M. J. Hartmann, A. Ho, M. Hoffmann, T. Huang, T. S. Humble, S. V. Isakov, E. Jeffrey, Z. Jiang, D. Kafri, K. Kechedzhi, J. Kelly, P. V. Klimov, S. Knysh, A. Korotkov, F. Kostritsa, D. Landhuis, M. Lindmark, E. Lucero, D. Lyakh, S. Mandrà, J. R. McClean, M. McEwen, A. Megrant, X. Mi, K. Michielsen, M. Mohseni, J. Mutus, O. Naaman, M. Neeley, C. Neill, M. Y. Niu, E. Ostby, A. Petukhov, J. C. Platt, C. Quintana, E. G. Rieffel, P. Roushan, N. C. Rubin, D. Sank, K. J. Satzinger, V. Smelyanskiy, K. J. Sung, M. D. Trevithick, A. Vainsencher, B. Villalonga, T. White, Z. J. Yao, P. Yeh, A. Zalcman, H. Neven, and J. M. Martinis, Quantum supremacy using a programmable superconducting processor, *Nature* **574**, 505 (2019).
- [70] J. Preskill, Quantum Computing in the NISQ era and beyond, *Quantum* **2**, 79 (2018).
- [71] R. Santagati, J. Wang, A. A. Gentile, S. Paesani, N. Wiebe, J. R. McClean, S. Morley-Short, P. J. Shadbolt, D. Bonneau, J. W. Silverstone, D. P. Tew, X. Zhou, J. L. O'Brien, and M. G. Thompson, Witnessing eigenstates for quantum simulation of Hamiltonian spectra, *Science Advances* **4** (2018).
- [72] S. Liu, S.-X. Zhang, C.-Y. Hsieh, S. Zhang, and H. Yao, Probing many-body localization by excited-state VQE, *arXiv:2111.13719* (2021).
- [73] Y. Y. Atas, E. Bogomolny, O. Giraud, and G. Roux, Distribution of the ratio of consecutive level spacings in random matrix ensembles, *Phys. Rev. Lett.* **110**, 084101 (2013).
- [74] M. M. Wolf, F. Verstraete, M. B. Hastings, and J. I. Cirac, Area laws in quantum systems: Mutual information and correlations, *Phys. Rev. Lett.* **100**, 070502 (2008).
- [75] C.-M. Jian, I. H. Kim, and X.-L. Qi, Long-range mutual information and topological uncertainty principle, *arXiv:1508.07006* (2015).

**SUPPLEMENTAL MATERIALS**

**Why is there a dominant-subleading  $\pi$ -pair pattern: A perturbation theory perspective**

In this section, via degenerate perturbation theory, we explain the overlap pattern between the initial product states and the quasi-eigenstates of the Floquet operator  $U_F$ .

The Floquet unitary  $U_F$  reads:

$$U_F = e^{-iH_2} e^{-i(\frac{\pi}{2}-\epsilon)H_1} = e^{-iH_2} e^{-i(\frac{\pi}{2}-\epsilon)\sum_j X_j} = e^{-iH_2} e^{-i\frac{\pi}{2}\sum_j X_j} e^{i\epsilon\sum_j X_j} = U_F^0 e^{i\epsilon\sum_j X_j}, \quad (\text{S1})$$

and by Taylor expansion on small imperfection  $\epsilon$ , we have:

$$\begin{aligned} U_F &= U_F^0 \sum_n \frac{(i\epsilon)^n}{n!} (\sum_j X_j)^n = U_F^0 (I + i\epsilon \sum_j X_j + O(\epsilon^2)) = U_F^0 + i\epsilon U_F^0 \sum_j X_j + O(\epsilon^2) \\ &\approx U_F^0 + U_F^1. \end{aligned} \quad (\text{S2})$$

Assuming  $|\phi\rangle$  and  $|\varphi\rangle$  are two quasi-eigenstates of  $U_F^0$  with quasi-energy difference  $\pi$ , which is approximately the case for ‘‘good initial states’’.

$$U_F^0 |\phi\rangle = e^{-i\varepsilon^0} |\phi\rangle \quad \text{and} \quad U_F^0 |\varphi\rangle = -e^{-i\varepsilon^0} |\varphi\rangle, \quad (\text{S3})$$

then  $U_F^0 |\varphi\rangle = e^{-i\varepsilon^0} |\varphi\rangle$ , where  $\varepsilon^0 = \frac{1}{2}(H_2(z_\phi) + H_2(-z_\phi)) = \frac{1}{2}(H_2(z_\varphi) + H_2(-z_\varphi)) + \pi \pmod{2\pi}$ ,  $z_\phi$  and  $z_\varphi$  are the corresponding computational basis configurations from quasi-eigenstates. And

$$|\phi\rangle = \frac{1}{\sqrt{2}} (e^{-i\frac{H_2(z_\phi)}{2}} |z_\phi\rangle + e^{-i\frac{H_2(-z_\phi)}{2}} | - z_\phi\rangle), \quad (\text{S4})$$

$$|\varphi\rangle = \frac{1}{\sqrt{2}} (e^{-i\frac{H_2(z_\varphi)}{2}} |z_\varphi\rangle - e^{-i\frac{H_2(-z_\varphi)}{2}} | - z_\varphi\rangle), \quad (\text{S5})$$

$|z_\phi\rangle$  and  $|z_\varphi\rangle$  are related by a Pauli- $X$  matrix on site  $j$  (here we assume that the choice of  $J_z$  makes the quasi-energy difference is closest to  $\pi$  with flipping on  $j$  site, in the main text we fix  $j = \frac{N-1}{2}$  for simplicity),

$$|z_\phi\rangle = X_j |z_\varphi\rangle, \quad (\text{S6})$$

and by convention we have,

$$|z_\phi\rangle = \prod_j X_j | - z_\phi\rangle. \quad (\text{S7})$$

The zero order perturbative quasi-eigenstate of  $U_F$  reads

$$|\psi^0\rangle = a|\varphi\rangle + b|\phi\rangle. \quad (\text{S8})$$

Then

$$\begin{aligned} (U_F^0 - e^{-i\varepsilon^0})|\psi^1\rangle &= (E_1 - U_F^1)|\psi^0\rangle \\ &= (E_1 - U_F^1)(a|\varphi\rangle + b|\phi\rangle), \end{aligned} \quad (\text{S9})$$

where  $|\psi^1\rangle$  is the first order perturbative quasi-eigenstate of  $U_F$ . It requires

$$\det \begin{bmatrix} -E_1 & \langle\varphi|U_F^1|\phi\rangle \\ \langle\phi|U_F^1|\varphi\rangle & -E_1 \end{bmatrix} = 0. \quad (\text{S10})$$

According to

$$|z_\phi\rangle = \frac{1}{\sqrt{2}} e^{i\frac{H_2(z_\phi)}{2}} (|\phi\rangle + | - \phi\rangle), \quad (\text{S11})$$

$$| - z_\phi\rangle = \frac{1}{\sqrt{2}} e^{i\frac{H_2(-z_\phi)}{2}} (|\phi\rangle - | - \phi\rangle), \quad (\text{S12})$$



$$|z_\varphi\rangle = \frac{1}{\sqrt{2}} e^{i\frac{H_2(z_\varphi)}{2}} (|\varphi\rangle + |-\varphi\rangle), \quad (\text{S13})$$

$$| - z_\varphi \rangle = \frac{1}{\sqrt{2}} e^{i\frac{H_2(-z_\varphi)}{2}} (|-\varphi\rangle - |\varphi\rangle), \quad (\text{S14})$$

the off-diagonal elements can be calculated:

$$\begin{aligned} \langle \varphi | U_F^1 | \phi \rangle &= \langle \varphi | i\epsilon U_F^0 | \frac{1}{\sqrt{2}} (e^{-i\frac{H_2(z_\phi)}{2}} |z_\phi\rangle + e^{-i\frac{H_2(-z_\phi)}{2}} | - z_\phi \rangle) \\ &= \langle \varphi | i\epsilon U_F^0 | (\frac{1}{2} e^{-i\frac{H_2(z_\phi)}{2}} e^{i\frac{H_2(z_\phi)}{2}} (|\varphi\rangle + |-\varphi\rangle) + \frac{1}{2} e^{-i\frac{H_2(-z_\phi)}{2}} e^{i\frac{H_2(-z_\phi)}{2}} (|-\varphi\rangle - |\varphi\rangle)) \\ &= \frac{i\epsilon}{2} (e^{-i\frac{H_2(z_\phi)}{2}} e^{i\frac{H_2(z_\phi)}{2}} e^{-i\epsilon^0} - e^{-i\frac{H_2(-z_\phi)}{2}} e^{i\frac{H_2(-z_\phi)}{2}} e^{-i\epsilon^0}) \\ &= \frac{i\epsilon}{2} (-e^{-i\frac{H_2(z_\phi)}{2}} e^{-i\frac{H_2(-z_\phi)}{2}} + e^{-i\frac{H_2(-z_\phi)}{2}} e^{-i\frac{H_2(z_\phi)}{2}}), \end{aligned} \quad (\text{S15})$$

$$\begin{aligned} \langle \phi | U_F^1 | \varphi \rangle &= \langle \phi | i\epsilon U_F^0 | \frac{1}{\sqrt{2}} (e^{-i\frac{H_2(z_\varphi)}{2}} |z_\varphi\rangle - e^{-i\frac{H_2(-z_\varphi)}{2}} | - z_\varphi \rangle) \\ &= \langle \phi | i\epsilon U_F^0 | (\frac{1}{2} e^{-i\frac{H_2(z_\varphi)}{2}} e^{i\frac{H_2(z_\varphi)}{2}} (|\phi\rangle + |-\phi\rangle) - \frac{1}{2} e^{-i\frac{H_2(-z_\varphi)}{2}} e^{i\frac{H_2(-z_\varphi)}{2}} (|\phi\rangle - |-\phi\rangle)) \\ &= \frac{i\epsilon}{2} (e^{-i\frac{H_2(z_\varphi)}{2}} e^{i\frac{H_2(z_\varphi)}{2}} e^{-i\epsilon^0} - e^{-i\frac{H_2(-z_\varphi)}{2}} e^{i\frac{H_2(-z_\varphi)}{2}} e^{-i\epsilon^0}) \\ &= \frac{i\epsilon}{2} (e^{-i\frac{H_2(z_\varphi)}{2}} e^{-i\frac{H_2(-z_\varphi)}{2}} - e^{-i\frac{H_2(-z_\varphi)}{2}} e^{-i\frac{H_2(z_\varphi)}{2}}). \end{aligned} \quad (\text{S16})$$

Bring the results of Eq. S15 and Eq. S16 back to Eq. S10, we have

$$E_1^2 + \frac{\epsilon^2}{4} e^{-2i\epsilon^0} (2 - e^{-i(H_{2z}(z_\phi) + H_{2z}(-z_\phi))} - e^{-i(H_{2z}(-z_\phi) + H_{2z}(z_\phi))}) = 0, \quad (\text{S17})$$

$$E_1^2 + \frac{\epsilon^2}{4} e^{-2i\epsilon^0} (2 - e^{-2ijW} - e^{+2ijW}) = 0, \quad (\text{S18})$$

$$E_1^2 + \frac{1}{2} \epsilon^2 e^{-2i\epsilon^0} (1 - \cos(2jW)) = 0, \quad (\text{S19})$$

$$E_1^2 + \epsilon^2 e^{-2i\epsilon^0} \sin^2(jW) = 0, \quad (\text{S20})$$

where  $H_{2z}$  is the linear Zeeman field term. And the first order perturbation correction to quasi-eigenenergy is

$$E_1 = \pm i\epsilon e^{-i\epsilon^0} \sin(jW), \quad (\text{S21})$$

where  $j$  is the site position for the matching flip  $X_j$  and  $W$  is the linear Zeeman field strength. Then the quasi-eigenenergy are

$$\begin{aligned} e^{-i\epsilon'} &= e^{-i\epsilon^0} (1 \pm i\epsilon \sin(jW)), \\ &\approx e^{-i\epsilon^0} e^{\pm i\epsilon \sin(jW)}, \end{aligned} \quad (\text{S22})$$

$$\epsilon' = \epsilon^0 \pm \epsilon \sin(jW) \pmod{2\pi}. \quad (\text{S23})$$

So  $\omega_b = 2\epsilon \sin(jW)$ , i.e. the second timescale  $T_b = \frac{2\pi}{\omega_b} = \frac{\pi}{\epsilon \sin(jW)}$ . And  $T_b$  is the same for all ‘‘good initial states’’ since the derivation above doesn’t depend on special good state configurations (see Fig. S1). This result leads a dominant beating oscillation in the dynamics of the state-averaged fidelity. The predictions from perturbation theory and numerical results are in good agreement, see Fig. S2.

When the quasienergy difference between two quasi-eigenstates of  $U_F^0$  equals to zero instead of  $\pi$ . Then the first order perturbative quasi-eigenenergy are:

$$E_1^2 + \frac{1}{2} \epsilon^2 e^{-2i\epsilon^0} (1 + \cos(2jW)) = 0, \quad (\text{S24})$$

$$E_1^2 + \epsilon^2 e^{-2i\epsilon^0} \cos^2(jW) = 0, \quad (\text{S25})$$




FIG. S1. The quasi-eigenenergy differences between dominant and subleading  $\pi$ -pair  $\delta\varepsilon_F(T_b)$  are the same for all “good initial states”. For “bad initial states”, although there is no obvious subleading  $\pi$ -pair, we can also calculate the  $\delta\varepsilon_F$  according to the magnitude of the overlaps and it is roughly determined by  $2J_z$ . It is consistent with our understanding from perturbation picture. ( $N = 9, J_z = \frac{\pi}{2N}, W = 5.0, \epsilon = 0.05$ )




FIG. S2. The beating oscillation frequency is determined by Fourier transformation on even period of dynamics using state-averaged fidelity mentioned in the main text. As discussed in the text, there are two different frequencies,  $\omega_b$  and  $\omega'_b$ . For “good initial states”,  $\omega_b \approx \frac{2\pi}{T_b} = 2\epsilon \sin(\frac{N-1}{2}W)$ ; for “bad initial states”,  $\omega'_b \approx 2J_z$ . Left:  $\omega_b$ ; Right:  $\omega'_b$ . ( $J_z = \frac{\pi}{2N}; W = 5.0; \epsilon = 0.05$ )

$$\varepsilon' = \varepsilon^0 \pm \epsilon \cos(jW) \pmod{2\pi}. \quad (\text{S26})$$

When we choose a  $J_z$  that is not perfect, i.e.  $\delta_{J_z} > 0$  ( $J_z = \frac{\pi - \delta_{J_z}}{2N}$ ), there are no quasi-eigenstates of  $U_F^0$  with the quasi-eigenenergy difference exactly equals 0 or  $\pi$  besides intrinsic  $\pi$ -pairs. For each  $\pi$ -pair up to one X flip on the target one, the quasi-eigenenergy difference is  $\pi - \delta_{J_z}$ , and the one with the smallest  $\delta_{J_z}$  constitutes the subleading  $\pi$ -pair in the overlap spectrum. Therefore, we can still see a dominant beating timescale  $T_b$  determined by the quasi-eigenenergy difference between dominant and subleading  $\pi$ -pairs.

For “bad initial states”, we show why there is another beating timescale  $T'_b$  different from the  $T_b$  for “good initial state”. For a given bad initial state (assuming matching site  $j = 2$  here), for example,  $|z\rangle = |01100\rangle$ ,  $|z\rangle$  and  $| -z\rangle$  form a  $\pi$ -pair of quasi-eigenstates of  $U_F^0$ , if we flip spin  $S_j$  ( $j = 0, \dots, N$ ) of  $|z\rangle$  and  $| -z\rangle$ , there will be  $N$  new  $\pi$ -pairs of quasi-eigenstates of  $U_F^0$ . In this case, different  $\pi$ -pairs are still related by the Pauli-X matrix and the second timescale  $T'_b$  is roughly determined by the original quasi-eigenenergy difference. For example, we set  $J_z = \frac{\pi}{2N}$  throughout the work, when  $N$  is large,  $J_z \sim 0$ , related quasi-eigenstates with quasi-eigenenergy difference of  $2J_z$  correspond to the  $T'_b$ , see Fig. S2. The difference is that there are more than one subleading  $\pi$ -pairs for a given dominant  $\pi$ -pair. Also from the perturbation theory, we can see many different beating frequencies in such cases.

### The relationship between the second timescale and quasi-energy difference

A random initial product state  $|z\rangle$  can be roughly written as a superposition of a dominant and a subleading  $\pi$  pairs,

$$|z\rangle = \alpha(|m\rangle + |-m\rangle) + \beta(|n\rangle + |-n\rangle), \quad (\text{S27})$$

where  $|\alpha|^2 + |\beta|^2 = \frac{1}{2}$ , and  $U_F|\pm m\rangle = \pm e^{-i\varepsilon_m}|m\rangle$ ,  $U_F|\pm n\rangle = \pm e^{-i\varepsilon_n}|n\rangle$ . At time  $t = 2lT$ .

$$U_F^{2l}|z\rangle = e^{-2li(\varepsilon_m - \varepsilon_n)}\alpha(|m\rangle + |-m\rangle) + \beta(|n\rangle + |-n\rangle), \quad (\text{S28})$$

and

$$F_{2l} = |\langle z|U_F^{2l}|z\rangle|^2 = 4|\alpha|^4 + 4|\beta|^4 + 8|\alpha|^2|\beta|^2 \cos(2l(\varepsilon_m - \varepsilon_n)). \quad (\text{S29})$$

Then the second timescale  $T_b$ , or the frequency is related the quasi-energy difference:

$$\omega_b = \frac{2\pi}{T_b} = \frac{2\pi}{2l} = |\varepsilon_m - \varepsilon_n|. \quad (\text{S30})$$

### The relationship between the third timescale and the quasienergy mismatch

For a finite-size system, there is a tiny quasi-energy mismatch in each  $\pi$ -pair and this is the origin of the third timescale.

$$|z\rangle = \frac{1}{\sqrt{2}}(|m\rangle + |-(m + \delta_m)\rangle), \quad (\text{S31})$$

At time  $t = 2lT$ ,

$$U_F^{2l}|z\rangle = \frac{1}{\sqrt{2}}(|m\rangle + e^{-2li\delta_m}|-(m + \delta_m)\rangle), \quad (\text{S32})$$

then the third timescale for the given initial state  $|z\rangle$  is:

$$T_g = 2l = \frac{2\pi}{\delta_m}. \quad (\text{S33})$$

### The dominant beating timescale in the dynamics of state-averaged fidelity

In Fig. 4 for state-averaged fidelity, we can see a dominant beating timescale  $T_b$ , which is caused by the ‘‘good initial states’’, and many other beating timescales  $T'_b$ , which are caused by ‘‘bad initial states’’. We explain why the  $T_b$  is dominant in state averaged case. Consider the fidelity of a given initial state firstly:

$$F_n = |\langle \psi|U_F^n|\psi\rangle|^2, \quad (\text{S34})$$

where  $|\psi\rangle$  is the initial state. And the initial state can be roughly represented by  $n$   $\pi$ -pairs

$$|\psi\rangle = \sum_{i=0}^{n-1} \alpha_i(|+\rangle_i + |-\rangle_i), \quad (\text{S35})$$

where  $2\sum_{i=0}^{n-1} |\alpha_i|^2 = 1$ . At time  $t = 2lT$ ,

$$U_F^{2l}|\psi\rangle = \sum_{i=0}^{n-1} \alpha_i e^{-2li\delta\varepsilon_F(i,0)}(|+\rangle_i + |-\rangle_i), \quad (\text{S36})$$

where  $\delta\varepsilon_F(i,0)$  is the quasi-eigenenergy difference between  $|+\rangle_i$  and  $|+\rangle_0$ . And

$$F_{2l} = \left| \sum_{i=0}^{n-1} 2|\alpha_i|^2 e^{-2li\delta\varepsilon_F(i,0)} \right|^2 \quad (\text{S37})$$

$$= \sum_{i=0}^{n-1} 4|\alpha_i|^4 + \sum_{i<j}^{n-1} 8|\alpha_i|^2|\alpha_j|^2 \cos(2l(\delta\varepsilon_F(i,0) - \delta\varepsilon_F(j,0))). \quad (\text{S38})$$




FIG. S3. State-averaged fidelity  $\bar{F}_n$  and its Fourier transformation ( $N = 11$ ,  $J_z = \pi + 1.5$ ,  $W = 5.0$ ,  $\epsilon = 0.05$ ).

The total height of the peak after Fourier transformation is:

$$\text{Height} \propto \sum_{i < j}^{n-1} 8|\alpha_i|^2|\alpha_j|^2 = 1 - \sum_{i=0}^{n-1} 4|\alpha_i|^4 = 1 - 4\text{EIPR}, \quad (\text{S39})$$

where  $\text{EIPR} = \sum_i |\alpha_i|^4$  [71, 72]. For “bad initial state”, there is a dominant  $\pi$ -pair and EIPR is large; for “good initial state”, there is a dominant  $\pi$ -pair and a subleading  $\pi$ -pair and EIPR is relatively small. Since the numbers of the “good initial states” and “bad initial states” are both the same:  $2^{N-1}$ , the dynamics of the state-averaged fidelity will exhibit a dominant beating timescale as given by the “good initial states”.

### The Fourier transform on the fidelity dynamics

To see the oscillation period in the dynamics more obviously, we can take Fourier transform of fidelity, which is defined as:

$$f(\omega) = \sum_{l=0}^{n-1} \bar{F}_l e^{-i\omega l}, \quad (\text{S40})$$

where  $\bar{F}_l$  is the state-averaged fidelity in the time step  $l$ . And we take the normalized form of  $f(\omega)$  in this Letter, which reads:

$$\tilde{f}(\omega) = \frac{|f(\omega)|}{n}. \quad (\text{S41})$$

For example, the height of the peak of subharmonic response is  $\tilde{f}(\pi)$ .

### Results for different imperfection $\epsilon$

To distinguish different phases controlled by the imperfection  $\epsilon$ , we can utilize the magnitude,  $\tilde{f}(\pi)$ , of the peak of the subharmonic response. When  $\epsilon$  is small, there is an obvious subharmonic response which means DTTSB, see Fig. S3. And the magnitude of the subharmonic response decreases as the imperfection increases, see Fig. S3 and Fig. S4.

When  $\epsilon$  is large enough, the fidelity at even periods will decay to zero after a few cycles and there is no obvious signal of the subharmonic response, see Fig. S5.

### Level statistics for many-body localization

Level statistics are often used to diagnose MBL phases. The level spacing ratio for quasi-eigenenergies is defined as:

$$r_n = \frac{\min\{\Delta^n, \Delta^{n+1}\}}{\max\{\Delta^n, \Delta^{n+1}\}}, \quad (\text{S42})$$




FIG. S4. State-averaged fidelity  $\bar{F}_n$  and its Fourier transformation ( $N = 11$ ,  $J_z = \pi + 1.5$ ,  $W = 5.0$ ,  $\epsilon = 0.25$ ).




FIG. S5. State-averaged fidelity  $\bar{F}_n$  and its Fourier transformation ( $N = 11$ ,  $J_z = \pi + 1.5$ ,  $W = 5.0$ ,  $\epsilon = 0.5$ ).

where  $\Delta^n = \varepsilon^n - \varepsilon^{n+1}$  is the gap between quasi-eigenenergy levels  $n$  and  $n + 1$ ,  $[r^{(n)}]$  is the level average of this ratio. In the delocalized phase (thermal phase), the level spacings follow a GOE distribution and the level-averaged level spacing ratio is  $[r^{(n)}] \approx 0.53$  [73]. In contrast, in the localized phase, the level spacing follows a Poisson distribution, which gives  $[r^{(n)}] = 2 \ln 2 - 1 \approx 0.39$  [73].

As discussed in the main text, the DTC order is stabilized by Stark MBL. We can see the distribution of level spacing ratio gradually crosses from the Poisson limit to the GOE type with increasing imperfection  $\epsilon$ , see Fig. S6 and Fig. S7 for level spacing distribution.

### Phase transition

To diagnose the phase transition and discrete time translational symmetry breaking, we utilize two indicators: the magnitude of the subharmonic response and the mutual information. The former is related to the definition of *DTTSB-1* in [8]: DTTSB occurs when the observables develop persistent oscillations whose periods are an integer multiple of the driving period; the latter is related to the definition of *DTTSB-2* in [8]: DTTSB occurs if the eigenstates of the Floquet unitary  $U_F$  cannot be short-range correlated.

As we increase the imperfection  $\epsilon$ , the magnitude,  $\tilde{f}(\pi)$ , of the peak of subharmonic response decreases, see Fig. S8, and eventually, becomes completely washed out when the system enters into the trivial thermal phase.

We further use the state-averaged mutual information  $I(A, B) = S_A + S_B - S_{AB}$  [8, 12, 74, 75], to check whether the quasi-eigenstate is short-range entangled. Here  $A$  and  $B$  are spins on opposite ends of a  $N$ -spin chain and  $S$  is the Von Neumann Entropy. For small imperfection  $\epsilon \approx 0$ , any quasi-eigenstate is long-range entangled “cat state”, indicating nearly full  $I(A, B) = \log 2$ , and mutual information drops dramatically upon leaving the DTTSB phase for large  $\epsilon$ , see Fig. S8.

Due to the limitation of computing power and finite size effect, we can not get the accurate  $\epsilon_c$  for fixed  $J_z$  and  $W$  by the numerical exploration of DTTSB, the phase diagram in the main text is schematic. Nonetheless, the distribution of the level spacing ratios, magnitude of the subharmonic oscillation peak, and mutual information of quasi-eigenstates provide convincing evidence of the DTTSB.




FIG. S6. Level spacing ratio:  $J_z = \pi + 1.5$ ,  $W = 5.0$ .




FIG. S7. Histogram of the distribution of the level spacing ratios:  $N = 13$ ,  $J_z = \pi + 1.5$ ,  $W = 5.0$ . (a),  $\epsilon = 0.05$ ; (b),  $\epsilon = 0.5$ . With increasing the imperfection  $\epsilon$ , we find that the distribution of level spacing ratio gradually crosses from the Poisson limit to the GOE distribution.

### Imperfect $J_z$

In the main text, we have shown the robust subharmonic response, i.e. DTC phase, and the novel beating oscillation with  $J_z = \frac{\pi}{2N}$ . This is a perfect  $J_z$  as for a given  $\pi$ -pair  $|\pm\rangle$  of  $U_F^0$ , there is only one  $\pi$ -pair  $|\pm'\rangle$  of  $U_F^0$  with one spin flipped has the quasi-eigenenergy difference exactly 0 or  $\pi$ . As we discussed above, even if we choose a  $J_z$  that is not perfect, the subharmonic response, as well as the beating oscillation, still exists. The dominant beating timescale is shown in Fig. S9 and the phase transition between the DTC phase and trivial thermalized phase is shown in Fig. S10 for imperfect  $J_z$ . Moreover, the classification of perfect  $J_z$  and imperfect  $J_z$  is for the beating timescale  $T_b$ . It is not necessary that  $J_z$  scale with the system size as  $1/N$ . In fact, any  $J_z$  that guarantees the Stark MBL is good enough to stabilize the DTC phase (see the next section).

### Different choices of $J_z$ and $W$

As discussed in [22], the linear Zeeman field, as well as the Stark MBL, will be suppressed by the imperfection  $\epsilon$  and a special term of  $zz$  interaction is important (linear form in our case). In this section, we show the results of different combinations of  $J_z$  and  $W$ . We find that DTC indeed requires the linear  $zz$  interaction and the impact of linear Zeeman field is suppressed. There are four choices of  $\{J_z, W\}$ : (a), {constant, constant}; (b), {constant, linear}; (c), {linear, constant}; (d), {linear, linear}. For example, {constant, linear} stands for

$$H_2 = J_z \sum_j Z_j Z_{j+1} + W \sum_j j Z_j. \quad (\text{S43})$$

The results of the Fourier peak height  $\tilde{f}(\pi)$  with different choices  $\{J_z, W\}$  are shown in Fig. S11. And we choose an arbitrary  $J_z$  with no size dependence on purpose to demonstrate the universality of the DTC phase and the physical




FIG. S8. Left: the height  $\tilde{f}(\pi)$  of the subharmonic response with different  $\epsilon$  ( $J_z = \pi + 1.5, W = 5.0, t = 10000T$ ); right: mutual information  $I(A, B)$  ( $J_z = \pi + 1.5; W = 5.0$ ).




FIG. S9. Dynamics of the state-averaged fidelity  $\bar{F}_n$  ( $N = 9; J_z = \frac{\pi - 0.05}{2N}; W = 5.0; \epsilon = 0.05$ ). The dominant beating timescale  $T_b$  still exists with an imperfect  $J_z$ .

picture in this Letter.

### The realization of DTC on quantum computers

Our model can be easily realized on the NISQ hardware with significantly fewer resources. In this section, we will show the experimental realization and protocol of our model on quantum devices. Our proposed quantum circuit structure for the DTC consists of three parts: the preparation circuit, Floquet unitary evolution circuit, and the measurement part, as shown in Fig. S12.  $R_a(\theta)$  is the rotation gate along  $a$  axis ( $a = x, y, z$ ). For example,  $R_x(\theta) = \exp(-iX\theta/2)$ , where  $X$  is the Pauli matrix. And  $R_{zz}(\theta)$  is parameterized  $zz$  coupling gate defined as  $R_{zz}(\theta) = \exp(-i\frac{\theta}{2}Z \otimes Z)$ . For a given product state  $|z\rangle$ , we can get many computational basis configurations after measuring the output state of the quantum circuit on computational basis and the probability of getting back the original  $|z\rangle$  configuration is the fidelity we focus on. Besides, we can also measure site averaged spin polarization as the dynamics indicator.




FIG. S10. Left: the Fourier spectrum peak height  $\tilde{f}(\pi)$  of the subharmonic response; right: mutual information  $I(A, B)$  ( $J_z = \frac{\pi - 0.05}{2N}$ ;  $W = 5.0$ ). Even if the  $J_z$  is not perfect, the DTC phase is still robust against the imperfection  $\epsilon$ .




FIG. S11. The height  $\tilde{f}(\pi)$  of the subharmonic response with different choices of  $J_z$  and  $W$ : (a), {constant, constant}; (b), {constant, linear}; (c), {linear, constant}; (d), {linear, linear}. With the constant  $zz$  interaction, we find that the subharmonic response is not robust against the imperfection  $\epsilon$  as shown in (a) and (b). And the linear  $zz$  interaction is important to stabilize the DTC as shown in (c) and (d). The difference between constant  $W$  and linear  $W$ , whose impact is suppressed by the imperfection  $\epsilon$ , is not obvious ( $J_z = \pi + 1$ ;  $W = 5.0$ ).






FIG. S12. The circuit structure for the implementation of kicked model in this Letter. In the preparation circuit part, we first prepare a product state  $|z\rangle = |S_0, S_1, \dots, S_{N-1}\rangle$  from the initial state  $|000\dots\rangle$  with  $\theta_j = S_j\pi$  ( $S = 0$ : spin up;  $S = 1$ : spin down). In the Floquet unitary evolution part,  $R_x$  corresponds to the kicked term  $U_1$ ;  $R_z$  is the term of linear Zeeman field and  $R_{zz}$  is the term of linear  $zz$  interaction. In the measurement part, we will get a computational basis configuration  $|z'\rangle$ . The probability of getting  $|z\rangle$  is the fidelity ( $\theta = \pi - 2\epsilon$ ,  $\theta_{z,j}^i = 2jW$ ,  $\theta_{zz,j}^i = 2(j+1)J_z$ ).