

# Molecular states of $D^*D^*\bar{K}^*$ nature

N. Ikeno,<sup>1,\*</sup> M. Bayar,<sup>2,3,†</sup> and E. Oset<sup>3,‡</sup>

<sup>1</sup>*Department of Agricultural, Life and Environmental Sciences,  
Tottori University, Tottori 680-8551, Japan*

<sup>2</sup>*Department of Physics, Kocaeli University, 41380, Izmit, Turkey*

<sup>3</sup>*Departamento de Física Teórica and IFIC,  
Centro Mixto Universidad de Valencia-CSIC Institutos de Investigación de Paterna,  
Aptdo. 22085, 46071 Valencia, Spain*

(Dated: August 9, 2022)

## Abstract

We study the  $D^*D^*\bar{K}^*$  system by using the Fixed Center Approximation to the Faddeev equations. Since the  $D^*D^*$  interaction is attractive and gives a bound state, and so is the case of the  $D^*\bar{K}^*$  interaction, where the  $J^P = 0^+$  bound state is identified with the  $X_0(2900)$ , the  $D^*D^*\bar{K}^*$  system leads to bound states manifestly exotic with *ccs* open quarks. We obtain bound states of isospin  $I = 1/2$ , negative parity and total spin  $J = 0, 1, 2$ . For  $J = 0$  we obtain one state, and for  $J = 1, 2$  we obtain two states in each case. The widths are relatively narrow and the binding energies range from 56 MeV to 158 MeV.

---

\* ikeno@tottori-u.ac.jp

† melahat.bayar@kocaeli.edu.tr

‡ oset@ific.uv.es

## I. INTRODUCTION

While the study of general Few Body Systems has a long tradition, the study of Few Body Systems made of mesons has only a recent history. A review of such systems has been done in Ref. [1]. The study of the meson meson interaction with chiral Lagrangians [2] and the realization that this interaction, properly unitarized, led to meson–meson bound states that could be associated to known mesonic resonances [3–7] (see recent reviews in Refs. [8–11]) prompted the study of three body systems made with mesons, some of which could be associated to known mesonic states [1].

With a few exceptions [12], most of the states found in the past from the meson meson interaction correspond to states which are not exotic in the sense that they could also be formed in principle from a conventional  $q\bar{q}$ . Yet, the recent experimental findings of the  $X_0(2900)$ ,  $X_1(2900)$  [13, 14] in the  $D\bar{K}$  invariant mass and the  $T_{cc}$  state [15, 16] in the  $DD\pi$  spectrum revealed clear exotic mesonic structures, since one has  $cs$  quarks in the first case and  $cc$  quarks in the second one. These findings open the door to the formation of few body systems with several open quarks, having three or more quarks with flavors like  $ccs$ ,  $ccc$ ,  $css$ , etc., after eliminating the  $q\bar{q}$  structures with no flavor, as  $u\bar{u}$ ,  $d\bar{d}$ ,  $s\bar{s}$ ,  $c\bar{c}$ , etc. Since flavor is conserved in strong interactions, these systems can be relatively stable since they cannot decay into lighter mesons with smaller number of flavors. In the present paper we report on calculations for the  $D^*D^*\bar{K}^*$  system, which has open  $ccs$  quarks. The reason to choose this system is because we can establish connection with the experimental finding of Refs. [13–16]. Indeed, the  $X_0(2900)$  was soon identified as a likely  $D^*\bar{K}^*$  molecular state [17–26]. Other works favor compact tetraquarks [27–29], yet, the relativized quark model of Ref. [30] disfavors the compact tetraquark structure and favors the molecular one.

A valuable information concerning the  $D^*\bar{K}^*$  system comes from the work of Ref. [12], where 10 years prior to the observation of the  $X_0(2900)$ , a  $D^*\bar{K}^*$  molecule had been predicted with the same quantum numbers as the  $X_0(2900)$  and a mass and width remarkably close to those of the observed ones. The other part of the interaction is for the  $D^*D^*$  subsystem. Here we rely upon the recent observation of the  $T_{cc}$  state [15, 16]. Once again the  $D^*D$  nature of the state has been advocated from the beginning [16] and supported by many theoretical papers [9, 31–45]. A mapping of the  $D^*D$  interaction to the  $D^*D^*$  system has been done in Ref. [46] using an extrapolation of the local hidden gauge approach [47–50]

to the charm sector, which was already used in Ref. [12] to study the same  $D^*D^*$  exotic system. A similar approach using the boson exchange model is done in Ref. [51]. It was found in Ref. [12] that the  $D^*D^*$  system was bound in  $I = 0$  and  $J^P = 1^+$ . In Ref. [46], the model was refined including decay channels and using the same cut off to regularize the loops that was used in Ref. [32] to describe the  $T_{cc}$  state.

Three body systems containing two charmed quarks (or antiquarks) and a strange quark have been recently studied. In Table 1 of Ref. [1] one finds the  $DDK$  studied in Ref. [52],  $D\bar{D}^*K$  studied in Ref. [53], and  $DD^*K$  studied in Ref. [54]. More recently the  $D\bar{D}^*K$  hexaquark state is also studied in [55] via QCD sum rules. In these systems one has, however,  $cc\bar{s}$  or  $c\bar{c}s$  quarks, but not the  $ccs$  combination that we have in the system that we study, which makes it super exotic. These works are done using different technical approaches. However, for the purpose of justifying the Fixed Center Approximation (FCA) to the Faddeev equations that we follow, we find interesting to mention two recent works on the study of the  $D\bar{D}K$  system. In Ref. [56] it is studied using the Gaussian expansion method, minimizing the energy of the system. The same system is studied in Ref. [57] using the Fixed Center Approximation (FCA) and the results obtained are very similar. The coincidence of two very different methods in a similar system to the one we study gives us confidence in the FCA method that we use in the present approach to study the  $D^*D^*\bar{K}^*$  system. The fact that we use input tuned to the  $T_{cc}$  state in Ref. [32] and the  $D^*\bar{K}^* X_0(2900)$  state in Ref. [22] gives us further confidence, not only on the existence of the bound states that we find, but also on the values of the masses and widths that we predict.

## II. FORMALISM

We will use the FCA to study the  $D^*D^*\bar{K}^*$  system. In this picture there is a cluster of two bound particles and the third one collides with the components of this cluster without modifying its wave function. Certainly, if the third particle is lighter than the constituents of the cluster, the approximation is better, which suggests to consider the  $D^*D^*$  systems as the cluster. This latter system has been shown to be bound in Refs. [46, 58, 59]. In Ref. [46] the local hidden gauge approach extrapolated to the charm sector has been used, and the loops are regularized with the cut off demanded to fit the binding of the  $T_{cc}$  state as a  $D^*D$  molecule in Ref. [32]. A state with isospin  $I = 0$  and  $J^P = 1^+$  was found. The




FIG. 1. Diagrams involved in the FCA for the collision of the  $\bar{K}^*$  with the cluster of  $D^*D^*$ .

predicted binding should be realistic and we take the results of Ref. [46] as input for the  $D^*D^*$  cluster.

We follow the formalism of Ref. [60], according to which, the amplitude for  $\bar{K}^*$  collision with the  $D^*D^*$  cluster is given by the sum of the partition functions  $T_1$ ,  $T_2$ , where  $T_1$  sums all diagrams in Fig. 1 where the  $\bar{K}^*$  collides first with the particle 1 of the cluster, while  $T_2$  sums all diagrams where the  $\bar{K}^*$  collides first with particle 2 of the cluster.

We have for the total amplitude  $T$

$$T \equiv T_1 + T_2$$

and  $T_1$ ,  $T_2$  are coupled through

$$T_1 = t_1 + t_1 G_0 T_2,$$

$$T_2 = t_2 + t_2 G_0 T_1,$$

where  $t_1$  is the scattering amplitude for  $D^*(1)\bar{K}^*$ , and  $t_2$  the corresponding amplitude for  $D^*(2)\bar{K}^*$  scattering and  $G_0$  is the  $\bar{K}^*$  propagator folded with the cluster wave function. However, we should take into account the isospin and spin of the  $D^*(i)\bar{K}^*$  amplitudes.

### A. ISOSPIN CONSIDERATIONS

With the isospin doublet  $(D^+, -D^0)$ , the  $I = 0$   $D^*D^*$  state is given by

$$|D^*D^*, I = 0\rangle = -\frac{1}{\sqrt{2}}(D^{*+}D^{*0} - D^{*0}D^{*+}) \quad (1)$$

and one should also bear in mind that this is accompanied by  $J = 1$  in each of the two components, which we will address below.

When evaluating  $t_1$  of the scattering amplitude for  $D^*(1)\bar{K}^*$  we have the following matrix element, using the notation  $|I_3(D^*(1)), I_3(D^*(2))\rangle |I_3(\bar{K}^*)\rangle$ , where we have chosen  $\bar{K}^*$  to have the  $I_3 = \frac{1}{2}$  component

$$\begin{aligned} & \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} \left( \langle D^*D^*, \frac{1}{2}, -\frac{1}{2} | -\langle D^*D^*, -\frac{1}{2}, \frac{1}{2} | \right) \\ & \times \langle \bar{K}^*, \frac{1}{2} || t_1 | \left( |D^*D^*, \frac{1}{2}, -\frac{1}{2}\rangle - |D^*D^*, -\frac{1}{2}, \frac{1}{2}\rangle \right) | \bar{K}^*, \frac{1}{2} \rangle. \end{aligned} \quad (2)$$

To make connection with the  $D^*\bar{K}^*$  isospin amplitudes of Refs. [12, 22], we combine the third component of  $D^*(1)$  with the one of the  $\bar{K}^*$  to give states of  $D^*\bar{K}^*$  isospin. Hence, we have with the notation  $|I(D^*(1)\bar{K}^*), I_3(D^*(1)\bar{K}^*)\rangle |I_3(D^*(2))\rangle$

$$\begin{aligned} & \frac{1}{2} \left[ \left( \langle 1, 1 | \langle -\frac{1}{2} | -\frac{1}{\sqrt{2}} (\langle 1, 0 | -\langle 0, 0 |) \langle \frac{1}{2} | \right) | t_1 | \left( |1, 1\rangle | -\frac{1}{2}\rangle - \frac{1}{\sqrt{2}} (|10\rangle - |00\rangle) | \frac{1}{2}\rangle \right) \right] \\ & = \frac{1}{2} \left( t^{I=1} + \frac{1}{2} t^{I=1} + \frac{1}{2} t^{I=0} \right) = \frac{3}{4} t^{I=1} + \frac{1}{4} t^{I=0}, \end{aligned} \quad (3)$$

since  $t_1$  only affects  $D^*(1)\bar{K}^*$  and  $D^*(2)$  is a spectator. The amplitude  $t_2$ , for  $D^*(2)\bar{K}^*$  scattering is obviously equal to  $t_1$ .

### B. SPIN CONSIDERATIONS

We start  $D^*D^*$  with  $J = 1$  and the  $\bar{K}^*$  has also  $J = 1$ . Hence, we can have three total spins for  $D^*D^*\bar{K}^*$ ,  $J = 0, 1, 2$ . Let us see each one of the cases.

1. TOTAL  $J = 0$

With the notation  $|j_3(D^*D^*), j_3(\bar{K}^*)\rangle$ , we have

$$|D^*D^*\bar{K}^*, J = 0\rangle = \frac{1}{\sqrt{3}}(|1, -1\rangle - |0, 0\rangle + |-1, 1\rangle)$$

and now we write the state  $|j(D^*D^*) = 1, j_3(D^*D^*)\rangle$  in terms of the  $|j_3(D^*(1))j_3(D^*(2))\rangle$  and we have

$$\frac{1}{\sqrt{3}} \left\{ \frac{1}{\sqrt{2}}(|1, 0\rangle - |0, 1\rangle) |-1\rangle - \frac{1}{\sqrt{2}}(|1, -1\rangle - |-1, 1\rangle) |0\rangle + \frac{1}{\sqrt{2}}(|0, -1\rangle - |-1, 0\rangle) |1\rangle \right\}$$

And now we write  $|j_3(D^*(1))j_3(\bar{K}^*)\rangle$  in terms of  $|j(D^*\bar{K}^*)j_3(D^*\bar{K}^*)\rangle$  and with the notation  $|j(D^*(1)\bar{K}^*), j_3(D^*(1)\bar{K}^*)\rangle |j_3(D^*(2))\rangle$ , we have

$$\begin{aligned} & \frac{1}{\sqrt{3}} \frac{1}{\sqrt{2}} \left\{ \left( \frac{1}{\sqrt{6}} |2, 0\rangle + \frac{1}{\sqrt{2}} |1, 0\rangle + \frac{1}{\sqrt{3}} |0, 0\rangle \right) |0\rangle - \left( \frac{1}{\sqrt{2}} |2, -1\rangle + \frac{1}{\sqrt{2}} |1, -1\rangle \right) |1\rangle \right. \\ & - \left( \frac{1}{\sqrt{2}} |2, 1\rangle + \frac{1}{\sqrt{2}} |1, 1\rangle \right) |-1\rangle + \frac{1}{\sqrt{2}} (|2, -1\rangle - |1, -1\rangle) |1\rangle \\ & \left. + \frac{1}{\sqrt{2}} (|2, 1\rangle - |1, 1\rangle) |-1\rangle - \left( \frac{1}{\sqrt{6}} |2, 0\rangle - \frac{1}{\sqrt{2}} |1, 0\rangle + \frac{1}{\sqrt{3}} |0, 0\rangle \right) |0\rangle \right\} \\ & = \frac{1}{\sqrt{3}} \{ |1, 0\rangle |0\rangle - |1, -1\rangle |1\rangle - |1, 1\rangle |-1\rangle \} \end{aligned}$$

and the matrix element of  $t_1$  sandwiched between this state becomes

$$t_1 = \frac{1}{3} (t^{j=1} + t^{j=1} + t^{j=1}) = t^{j=1}.$$

The result is expected since if we have  $| (D^*(1)\bar{K}^*)D^*(2)\rangle$  with  $j(D^*(2)) = 1$ , the spin of  $D^*(1)\bar{K}^*$  must necessarily be 1 to match total spin zero for the  $(D^*(1)\bar{K}^*)D^*(2)$  system.

2. TOTAL  $J = 1, J_3 = 1$

Once again, with the  $|j_3(D^*D^*), j_3(\bar{K}^*)\rangle$  representation we have the state

$$\frac{1}{\sqrt{2}} \{ |1, 0\rangle - |0, 1\rangle \}$$

which in the  $|j_3(D^*(1))j_3(D^*(2))\rangle |j_3(\bar{K}^*)\rangle$  representation reads

$$\frac{1}{\sqrt{2}} \left\{ \left( \frac{1}{\sqrt{2}} |1, 0\rangle - \frac{1}{\sqrt{2}} |0, 1\rangle \right) |0\rangle - \left( \frac{1}{\sqrt{2}} |1, -1\rangle - \frac{1}{\sqrt{2}} |-1, 1\rangle \right) |1\rangle \right\}$$

and written in terms of  $|j(D^*(1)\bar{K}^*)j_3(D^*(1)\bar{K}^*)\rangle |j_3(D^*(2))\rangle$  is

$$\begin{aligned} & \frac{1}{2} \left\{ \left( \frac{1}{\sqrt{2}} |2, 1\rangle + \frac{1}{\sqrt{2}} |1, 1\rangle \right) |0\rangle + \left( -\sqrt{\frac{2}{3}} |2, 0\rangle + \frac{1}{\sqrt{3}} |0, 0\rangle \right) |1\rangle \right. \\ & \left. - |2, 2\rangle | -1\rangle + \left( \frac{1}{\sqrt{6}} |2, 0\rangle - \frac{1}{\sqrt{2}} |1, 0\rangle + \frac{1}{\sqrt{3}} |0, 0\rangle \right) |1\rangle \right\} \\ & = \frac{1}{2} \left\{ \left( \frac{1}{\sqrt{2}} |2, 1\rangle + \frac{1}{\sqrt{2}} |1, 1\rangle \right) |0\rangle - \frac{1}{\sqrt{6}} |2, 0\rangle |1\rangle + \frac{2}{\sqrt{3}} |0, 0\rangle |1\rangle \right. \\ & \left. - |2, 2\rangle | -1\rangle - \frac{1}{\sqrt{2}} |1, 0\rangle |1\rangle \right\}. \end{aligned}$$

The  $t_1$  matrix sandwiched between this state gives the combination

$$t_1 = \frac{1}{4} \left( \frac{5}{3} t^{j=2} + t^{j=1} + \frac{4}{3} t^{j=0} \right).$$

### 3. TOTAL $J = 2$ , $J_3 = 2$

In the  $|j_3(D^*D^*), j_3(\bar{K}^*)\rangle$  representation the state is

$$|1, 1\rangle.$$

Then with the  $|j_3(D^*(1))j_3(D^*(2))\rangle |j_3(\bar{K}^*)\rangle$  notation we have

$$\left( \frac{1}{\sqrt{2}} |1, 0\rangle - \frac{1}{\sqrt{2}} |0, 1\rangle \right) |1\rangle$$

which in the  $|j(D^*(1)\bar{K}^*)j_3(D^*(1)\bar{K}^*)\rangle |j_3(D^*(2))\rangle$  representation reads

$$\frac{1}{\sqrt{2}} |2, 2\rangle |0\rangle - \frac{1}{\sqrt{2}} \left( \frac{1}{\sqrt{2}} |2, 1\rangle - \frac{1}{\sqrt{2}} |1, 1\rangle \right) |1\rangle.$$

The  $t_1$  matrix sandwiched between this state is then

$$t_1 = \frac{3}{4} t^{j=2} + \frac{1}{4} t^{j=1}.$$

### 4. COMBINED SPIN AND ISOSPIN AMPLITUDE

Combining the isospin and the spin decomposition of the amplitudes in subsections IIA and IIB we find the final contributions

$$\begin{aligned}
t_1 &= \frac{3}{4}t^{I=1, j=1} + \frac{1}{4}t^{I=0, j=1}, \quad \text{for } J = 0 \\
t_1 &= \frac{1}{16} \left\{ 5t^{I=1, j=2} + 3t^{I=1, j=1} + 4t^{I=1, j=0} + \frac{5}{3}t^{I=0, j=2} \right. \\
&\quad \left. + t^{I=0, j=1} + \frac{4}{3}t^{I=0, j=0} \right\}, \quad \text{for } J = 1 \\
t_1 &= \frac{1}{16} \left\{ 9t^{I=1, j=2} + 3t^{I=1, j=1} + 3t^{I=0, j=2} + t^{I=0, j=1} \right\}, \quad \text{for } J = 2.
\end{aligned} \tag{4}$$

### C. NORMALIZATION OF THE AMPLITUDES

By looking at the papers [61, 62] and the diagram of double scattering in Fig. 1 we find for the  $S$  matrix

$$\begin{aligned}
S^{(2)} &= -i(2\pi)^4 \delta^4(p_{fin} - p_{in}) \frac{1}{\mathcal{V}^2} \\
&\times \frac{1}{\sqrt{2\omega_{\bar{K}^*}}} \frac{1}{\sqrt{2\omega_{\bar{K}^*}}} \frac{1}{\sqrt{2\omega_{D^*}}} \frac{1}{\sqrt{2\omega_{D^*}}} \frac{1}{\sqrt{2\omega_{D^*}}} \frac{1}{\sqrt{2\omega_{D^*}}} \\
&\times t_1 t_2 \int \frac{d^3q}{(2\pi)^3} F(\vec{q}) \frac{1}{q^{0^2} - \vec{q}^2 - m_{\bar{K}^*}^2 + i\epsilon},
\end{aligned} \tag{5}$$

where  $\mathcal{V}$  is the volume of the normalization box,  $p_{fin}$  and  $p_{in}$  the final and initial momenta of the three particle system,  $q$  is the momentum of the exchanged  $\bar{K}^*$ ,  $w_i = \sqrt{m_i^2 + \vec{q}_i^2}$  and  $F(\vec{q})$  the form factor of the cluster. If we look at the process from the macroscopic perspective of having  $(D^*(1)D^*(2))_C \bar{K}^*$ , with  $(D^*(1)D^*(2))_C$  meaning the  $D^*D^*$  cluster, the  $S$  matrix reads

$$S^{(2)} = -i(2\pi)^4 \delta^4(p_{fin} - p_{in}) \frac{1}{\mathcal{V}^2} \frac{1}{\sqrt{2\omega_{\bar{K}^*}}} \frac{1}{\sqrt{2\omega_{\bar{K}^*}}} \frac{1}{\sqrt{2\omega_C}} \frac{1}{\sqrt{2\omega_C}} T^{(2)}, \tag{6}$$

Hence

$$\begin{aligned}
T^{(2)} &= 2\omega_C t_1 t_2 \frac{1}{2\omega_{D^*}} \frac{1}{2\omega_{D^*}} \int \frac{d^3q}{(2\pi)^3} F(\vec{q}) \frac{1}{q^{0^2} - \vec{q}^2 - m_{\bar{K}^*}^2 + i\epsilon} \\
&= \frac{2\omega_C}{2\omega_{D^*}} \frac{2\omega_C}{2\omega_{D^*}} \frac{1}{2\omega_C} t_1 t_2 \int \frac{d^3q}{(2\pi)^3} F(\vec{q}) \frac{1}{q^{0^2} - \vec{q}^2 - m_{\bar{K}^*}^2 + i\epsilon}.
\end{aligned} \tag{7}$$

It is thus convenient to write the partition functions suited to the macroscopic formalism as



$$\tilde{T}_1 = \tilde{t}_1 + \tilde{t}_1 \tilde{G}_0 \tilde{T}_2; \quad \tilde{T}_2 = \tilde{t}_2 + \tilde{t}_2 \tilde{G}_0 \tilde{T}_1, \quad (8)$$

where approximating  $\omega_{D^*} = m_{D^*}$ ,  $\omega_C = m_C$

$$\tilde{t}_1 = \frac{2m_C}{2m_{D^*}} t_1; \quad \tilde{t}_2 = \frac{2m_C}{2m_{D^*}} t_2,$$

and

$$\tilde{G}_0 = \frac{1}{2m_C} \int \frac{d^3q}{(2\pi)^3} F(\vec{q}) \frac{1}{q^0 - \vec{q}^2 - m_{\bar{K}^*}^2 + i\epsilon}. \quad (9)$$

The form factor of the cluster is obtained from the wave function of the cluster in momentum space [63] and reads [60, 61]

$$F(q) = \frac{1}{\mathcal{N}} \int_{|\vec{p}-\vec{q}| < q_{max}}^{p < q_{max}} d^3p \frac{1}{m_C - \sqrt{m_{D^*}^2 + \vec{p}^2} - \sqrt{m_{D^*}^2 + \vec{p}^2}} \times \frac{1}{m_C - \sqrt{m_{D^*}^2 + (\vec{p}-\vec{q})^2} - \sqrt{m_{D^*}^2 + (\vec{p}-\vec{q})^2}}, \quad (10)$$

with the normalization constant  $\mathcal{N}$

$$\mathcal{N} = \int_{p < q_{max}} d^3p \left( \frac{1}{m_C - \sqrt{m_{D^*}^2 + \vec{p}^2} - \sqrt{m_{D^*}^2 + \vec{p}^2}} \right)^2 \quad (11)$$

where  $q_{max}$  corresponds to the cut off of the loop function  $G_{D^*D^*}$  in the Bethe Salpeter equation for the  $D^*D^*$  scattering

$$t = V + V G_{D^*D^*} t, \quad (12)$$

for which we use the value  $q_{max} = 420$  MeV used in Ref. [46] deduced to get the  $T_{cc}$  state in Ref. [32]. With this representation of the  $t$  matrix,  $F(\vec{q})$  vanishes for  $|\vec{q}| > 2 q_{max}$ .

Following Refs. [60, 61] we take  $q^0$  in the rest frame of the cluster, which in the present case is

$$q^0 = \frac{s - m_{\bar{K}^*}^2 - m_C^2}{2m_C}$$

with  $s$  the square of the total rest energy of the  $(D^*D^*)_C \bar{K}^*$  system. We also need the argument of the  $t_1$  matrix of the  $D^* \bar{K}^*$  subsystem. This is evaluated taking

$$\begin{aligned} s' &= (p_{\bar{K}^*} + \frac{1}{2} P_C)^2 = m_{\bar{K}^*}^2 + \frac{1}{4} m_C^2 + q^0 m_C \\ &= \frac{1}{2} (s - m_{\bar{K}^*}^2 - m_C^2) + \frac{1}{4} m_C^2 + m_{\bar{K}^*}^2 \simeq \frac{1}{2} (s - m_{\bar{K}^*}^2 - m_C^2) + m_{D^*}^2 + m_{\bar{K}^*}^2, \end{aligned}$$

the last results corresponding to the formula used in Refs. [60, 61]

We should note that, since  $t_1 = t_2$ , then  $T_1 = T_2$  and, thus, Eq. (8) leads to

$$\tilde{T}_1 = \tilde{t}_1 + \tilde{t}_1 \tilde{G}_0 \tilde{T}_1; \quad \tilde{T}_1 = \frac{1}{\tilde{t}_1^{-1} - \tilde{G}_0}; \quad \tilde{T} = \tilde{T}_1 + \tilde{T}_2 = 2\tilde{T}_1 \quad (13)$$

and we represent  $|\tilde{T}|^2$  for real values of  $s$  looking for the peaks, from where we deduce the mass and width of the states that we find.

#### D. ELEMENTARY AMPLITUDES

We need the  $D^* \bar{K}^*$  amplitudes for the different  $I, j$  states. They are obtained in [12] and [22]. The  $t$  matrices are obtained by means of the Bethe Salpeter equations and the potential  $V^{I, j}$  for all the cases that we have in Eq. (4) are given in Tables XI, XII of [12]. To these potentials we add an imaginary part to account for the decay into  $D \bar{K}$  for  $I = 0, j(D^* \bar{K}^*) = 0, 2$  and into  $D^* \bar{K}$  for  $I = 0, j(D^* \bar{K}^*) = 1, 2$ . For the decay into  $D^* \bar{K}$  we find [22]

$$i \text{Im} V^{(D^* \bar{K})} = -i A_j \frac{1}{8\pi} (G' g m_{D^*})^2 q^5 \left( \frac{1}{(\omega_K(q) - p_2^0)^2 - \bar{q}^2 - m_\pi^2} \right)^2 \frac{1}{\sqrt{s}} F'^4(q) \quad (14)$$

where we have used  $q = \frac{\lambda^{1/2}(s', m_{D^*}^2, m_{\bar{K}}^2)}{2\sqrt{s'}}$ ,  $p_2^0 = \frac{s' + m_{\bar{K}^*}^2 - m_{D^*}^2}{2\sqrt{s'}}$ ,  $F'(q) = e^{((\omega_K - p_2^0)^2 - \bar{q}^2)/\Lambda^2}$ ,  $G' = \frac{3g'}{4\pi^2 f}$ ;  $g' = -\frac{G_V m_\rho}{\sqrt{2} f^2}$ ,  $G_V \simeq 55$  MeV,  $f = 93$  MeV. As in [22] we choose  $A_j = 3/2$  for  $j = 1$  and  $A_j = 9/10$  for  $j = 2$ .

The decay into  $D \bar{K}$  is studied in [12] and a long formula is obtained for the box diagrams containing  $D \bar{K}$  intermediate state. The imaginary part alone is easier to evaluate and we sketch its derivation in Appendix A. We consider the imaginary part only for the  $I = 0, D^* \bar{K}^*$  states. The reason is that only the  $I = 0$  states of  $D^* \bar{K}^*$  are bound. The  $D^* \bar{K}^*$  interaction in  $I = 1$  is repulsive. The amplitudes of  $I = 1$  are then small compared to the  $I = 0$  which develop poles in the bound region, and neglecting a small imaginary part of the  $I = 1$  amplitudes has a negligible effect in the final results concerning the three body bound states.

The result for  $I = 0$  of the imaginary part for decay into  $D \bar{K}$  is given by

$$i \text{Im} V^{(D \bar{K})} = -i F_j g^4 \frac{9}{15} \frac{1}{2\pi} \frac{1}{\sqrt{s'}} q^5 \left( \frac{1}{(\omega_K(q) - p_2^0)^2 - \bar{q}^2 - m_\pi^2} \right)^2 F'^4(q) F_{HQ} \quad (15)$$




FIG. 2. Modulus squared of the  $D^*D^*\bar{K}^*$  scattering amplitude with total spin  $J = 0$ . The dashed line ignores the  $\bar{K}^*$  width and the solid one accounts for it via a convolution of the  $G_{D^*\bar{K}^*}$  function.

where  $q = \frac{\lambda^{1/2}(s', m_D^2, m_{\bar{K}}^2)}{2\sqrt{s'}}$ ,  $F_{HQ} = \left(\frac{m_{D^*}}{m_{\bar{K}^*}}\right)^2$  and  $F_j = 5$  for  $j = 0$  and  $F_j = 2$  for  $j = 2$ . Then the  $t_1$  matrix is obtained as

$$t_1 = \frac{1}{V^{-1} - G_{D^*\bar{K}^*}}, \quad (16)$$

and to get the same results as in [22],  $G_{D^*\bar{K}^*}$  is regularized with dimensional regularization with the same input as in [22]. In the  $G_{D^*\bar{K}^*}$  function we also perform a convolution to account for the width of the  $\bar{K}^*$  as done [12, 22].

### III. RESULTS

In Fig. 2 we show the results for  $|\tilde{T}|^2$  as a function of the energy of the  $D^*D^*\bar{K}^*$  system for  $J = 0$ . We find a clear peak around 4845 MeV, about 61 MeV below the  $D^*D^*\bar{K}^*$  threshold. Most of the binding comes from the binding of the  $X_0(2900)$  as a  $D^*\bar{K}^*$  state which is about 30 MeV. The interaction with two  $D^*$  would give rise to a binding about twice as big. The width considering the convolution is about 12 MeV. We see that the consideration of the  $\bar{K}^*$  width by means of a convolution of the  $G_{D^*\bar{K}^*}$  function reduces the strength of  $|\tilde{T}|^2$  and increases a bit the width, but barely changes the mass. As we have seen in Eq. (4), we can




FIG. 3. Modulus squared of the  $D^*D^*\bar{K}^*$  scattering amplitude with total spin  $J = 1$ . Same meaning of the lines as Fig. 2.

see that in this case we have only  $j_{D^*\bar{K}^*} = 1$  and the width comes from the imaginary part of  $t^{I=0, j=1}$ , which has its source in the decay to  $D^*\bar{K}$  (leaving apart the  $\bar{K}^*$  width effect).

In Fig. 3 we show the same picture for total spin  $J = 1$ . Interestingly, we see now two peaks, indicating two states. In this case it is easy to trace the origin of the peaks. As we can see in Eq. (4) for  $J = 1$  we have now contributions from  $j = 0, 1, 2$ . We can see in Table 5 of [22] that the  $1^+$  and  $0^+$  states have about the same mass, 2861 MeV and 2866 MeV respectively, but the  $2^+$  state is more bound, at 2775 MeV. It is then clear that the first peak (higher energy) that we call state I is due to  $t^{I=0, j=0,1}$  while the second peak that we call state II is due to  $t^{I=0, j=2}$ . We also note that the effect of the  $\bar{K}^*$  width is more visible for the first peak, which is understandable because in the second peak there is little phase space for a decay of  $D^*D^*\bar{K}^*$  to  $D^*D^*\bar{K}\pi$  compared to  $D^*D\bar{K}$ .

In Fig. 4 we see a similar pattern for the case of total  $J = 2$ . From Eq. (4) one can also see that now one has  $j = 1, 2$  contribution. The first peak comes from the  $t^{I=0, j=1}$  amplitude while the second peak is due to  $t^{I=0, j=2}$ .




FIG. 4. Modulus squared of the  $D^*D^*\bar{K}^*$  scattering amplitude with total spin  $J = 2$ . Same meaning of the lines as Fig. 2.

Thus, in total we find 5 states which we summarize in Table I. There we also show the main decay channel expected based on the source of imaginary part of the  $t_1$  amplitudes involved in the peak. The binding energies range from 56 MeV to 158 MeV and the widths are relatively small, ranging from 12 MeV to 56 MeV.

TABLE I. Mass, binding, width (including the effects of the  $\bar{K}^*$  width) and decay modes of the states obtained for  $D^*D^*\bar{K}^*$ . The binding is with respect to  $2m_{D^*} + m_{K^*}$ .

$J$		$M$ [MeV]	$B$ [MeV]	$\Gamma$ [MeV]	Main decay mode
0	(State I)	4845	61	12	$D^*D^*\bar{K}$
1	(State I)	4849	57	15	$D^*D\bar{K}, D^*D^*\bar{K}$
1	(State II)	4750	156	55	$D^*D\bar{K}, D^*D^*\bar{K}$
2	(State I)	4850	56	15	$D^*D^*\bar{K}$
2	(State II)	4748	158	56	$D^*D\bar{K}, D^*D^*\bar{K}$

## IV. CONCLUSIONS

We have conducted a search for possible bound states of the three body system  $D^*D^*\bar{K}^*$ . For this we have relied upon the FCA to the Faddeev equations which has proved reliable in the study of related systems compared to the other methods like the variational one using the Gaussian expansion method. For the study of this system we have chosen as the cluster the  $D^*D^*$ , which appears bound in  $I = 0$  and  $J^P = 1^+$ , and the  $\bar{K}^*$  collides with it repeatedly. The FCA method has the virtue of allowing one to trace the structures obtained to the scattering amplitudes of  $D^*\bar{K}^*$ . This latter system has bound states in  $j = 0, 1, 2$ , with the  $j = 0, 1$  states very close in energy and the  $j = 2$  state more bound. In the  $D^*D^*\bar{K}^*$  system we can have total spin  $J = 0, 1, 2$ , and each of these spins the  $D^*\bar{K}^*$  amplitude appears with different combinations of isospin and spin. As a consequence, we obtain two peaks in the three-body partition functions which are tied to the  $j = 0, 1$  and  $j = 2$  of  $D^*\bar{K}^*$ , with the exception of  $J = 0$ , where the  $D^*\bar{K}^*$  system can only be in  $j = 1$ , and here we only get one peak.

In total we obtained 5 three-body states, one for  $J = 0$ , two for  $J = 1$  and two for  $J = 2$  and we evaluate the mass and width of these states. The states obtained, stemming from the cluster of  $D^*D^*$  with  $I = 0$  and  $J^P = 1^+$ , have total isospin  $I = 1/2$ , negative parity and spin  $J = 0, 1, 2$ , the widths are relatively narrow, ranging from 12 MeV to 56 MeV and the bindings range from 56 MeV to 158 MeV. We have also shown which is the main decay channel of each particular state to facilitate its experimental search. We hope the present work stimulates future searches for these super exotic states containing *ccs* quarks. As mentioned in Ref. [64], since the flavor is conserved in the strong interactions, we can expect new states made of many mesons, relatively stable, which cannot decay to mesons with smaller meson number. These would make them similar to nuclei where the baryon number conservation is responsible for the elements of the periodic table, giving rise to a new periodic table of mesons. In coincidence with the point made in Ref. [64] we think that we are just at the beginning of this new era.




FIG. 5. Box diagrams accounting for the decay of  $D^* \bar{K}^*$  in  $I = 0$  into  $D \bar{K}$ .

## ACKNOWLEDGMENTS

The work of N. I. was partly supported by JSPS KAKENHI Grant Number JP19K14709. This work is partly supported by the Spanish Ministerio de Economía y Competitividad (MINECO) and European FEDER funds under Contract No. PID2020-112777GB-I00, and by Generalitat Valenciana under contract PROMETEO/2020/023. This project has received funding from the European Union Horizon 2020 research and innovation programme under the program H2020-INFRAIA- 2018-1, grant agreement No. 824093 of the STRONG-2020 project.

## APPENDIX A

With the  $D^*$ ,  $\bar{K}^*$  doublets  $(D^{*+}, -D^{*0})$   $(\bar{K}^{\bar{*}0}, -K^{*-})$  the  $I = 0$  wave function is given by

$$-\frac{1}{\sqrt{2}}(D^{*+} K^{*-} - D^{*0} \bar{K}^{\bar{*}0})$$

and the box diagrams that account for  $D \bar{K}$  decay are given in Fig. 5

The dynamics comes from the vector  $\rightarrow$  pseudoscalar-pseudoscalar vertex,  $VPP$  given by the Lagrangian

$$\mathcal{L} = -ig \langle [P, \partial_\mu P] V^\mu \rangle, \quad g = \frac{M_V}{2f} \quad (M_V = 800 \text{ MeV}, \quad f = 93 \text{ MeV})$$

where  $P, V$  are  $q\bar{q}$  matrices written in terms of pseudoscalar and vector mesons.

$$P = \begin{pmatrix} \frac{1}{\sqrt{2}}\pi^0 + \frac{1}{\sqrt{3}}\eta + \frac{1}{\sqrt{6}}\eta' & \pi^+ & K^+ & \bar{D}^0 \\ \pi^- & -\frac{1}{\sqrt{2}}\pi^0 + \frac{1}{\sqrt{3}}\eta + \frac{1}{\sqrt{6}}\eta' & K^0 & D^- \\ K^- & \bar{K}^0 & -\frac{1}{\sqrt{3}}\eta + \sqrt{\frac{2}{3}}\eta' & D_s^- \\ D^0 & D^+ & D_s^+ & \eta_c \end{pmatrix}, \quad (17)$$

$$V = \begin{pmatrix} \frac{1}{\sqrt{2}}\rho^0 + \frac{1}{\sqrt{2}}\omega & \rho^+ & K^{*+} & \bar{D}^{*0} \\ \rho^- & -\frac{1}{\sqrt{2}}\rho^0 + \frac{1}{\sqrt{2}}\omega & K^{*0} & D^{*-} \\ K^{*-} & \bar{K}^{*0} & \phi & D_s^{*-} \\ D^{*0} & D^{*+} & D_s^{*+} & J/\psi \end{pmatrix}, \quad (18)$$

To simplify the calculation we evaluate the diagrams at threshold of  $D^*\bar{K}^*$ . The external three momenta are zero and  $\vec{q}$  is the running variable in the loop. In this limit the  $\epsilon^0$  component of the external vectors is zero and all the vertices are of the type of  $\vec{\epsilon} \cdot \vec{q}$ . Since one is concerned about the imaginary part, one realizes that the pions will be off shell because  $D^* \rightarrow D\pi$  is possible but then  $\bar{K}^* + \pi \rightarrow \bar{K}$  does not go. The loop function is then given by

$$i \int \frac{d^4q}{(2\pi)^4} \left( \frac{1}{q^0^2 - \vec{q}^2 - m_\pi^2} \right)^2 \frac{1}{(p_1 - q)^2 - m_D^2 + i\epsilon} \frac{1}{(p_2 + q)^2 - m_K^2 + i\epsilon},$$

and we can perform the  $q^0$  integration analytically using Cauchy's integration. Since we are concerned only about the imaginary part coming from the  $D\bar{K}$  placed on shell we consider only the poles of the  $D$  and  $\bar{K}$  propagators and obtain

$$\int \frac{d^3q}{(2\pi)^3} \left( \frac{1}{(w_K - p_2^0)^2 - \vec{q}^2 - m_\pi^2} \right)^2 \frac{1}{2w_K} \frac{1}{2w_D} \frac{1}{p_1^0 + p_2^0 - w_K - w_D + i\epsilon}.$$

We have integrals of type

$$\epsilon^l \epsilon^m \epsilon^r \epsilon^s \int \frac{d^3q}{(2\pi)^3} f(\vec{q}^2) q^l q^m q^r q^s = \frac{1}{15} \int \frac{d^3q}{(2\pi)^3} f(\vec{q}^2) \vec{q}^4 (\delta_{lm}\delta_{rs} + \delta_{lr}\delta_{ms} + \delta_{ls}\delta_{mr}) \epsilon^l \epsilon^m \epsilon^r \epsilon^s,$$

and considering the  $VV$  spin projectors in  $j = 0, 1, 2$  [65]

$$\begin{aligned} \mathcal{P}^{(0)} &= \frac{1}{3} \epsilon^l \epsilon^l \epsilon^r \epsilon^r \\ \mathcal{P}^{(1)} &= \frac{1}{2} (\epsilon^l \epsilon^m \epsilon^l \epsilon^m - \epsilon^l \epsilon^m \epsilon^m \epsilon^l) \\ \mathcal{P}^{(2)} &= \frac{1}{2} (\epsilon^l \epsilon^m \epsilon^l \epsilon^m + \epsilon^l \epsilon^m \epsilon^m \epsilon^l) - \frac{1}{3} \epsilon^l \epsilon^l \epsilon^m \epsilon^m \end{aligned}$$



and the appropriate factors in the vertices, we obtain from the sum of all the diagrams

$$i \text{Im} V^{(D\bar{K})} = -i F_j g^4 \frac{9}{15} \frac{1}{2\pi} \frac{1}{\sqrt{s'}} q^5 \left( \frac{1}{(\omega_K(q) - p_2^0)^2 - \bar{q}^2 - m_\pi^2} \right)^2 F'^4(q) F_{HQ} \quad (19)$$

where we have also included a factor demanded by heavy quark symmetry  $F_{HQ}$  to get the coupling  $VPP$  in the case of heavy vectors [66] and used  $q = \frac{\lambda^{1/2}(s', m_D^2, m_{\bar{K}}^2)}{2\sqrt{s'}}$ ,  $F_{HQ} = \left( \frac{m_{D^*}}{m_{\bar{K}^*}} \right)^2$ ,  $p_2^0 = \frac{s' + m_{\bar{K}^*}^2 - m_{D^*}^2}{2\sqrt{s'}}$ ,  $F'(q) = e^{((\omega_K - p_2^0)^2 - \bar{q}^2)/\Lambda^2}$ .

Further details can be seen in the evaluation of the width of the analogous  $\bar{B}^* \bar{K}^*$  state in section D of Ref. [67].

- 
- [1] A. Martinez Torres, K. P. Khemchandani, L. Roca and E. Oset, *Few Body Syst.* **61**, 35 (2020).
  - [2] J. Gasser and H. Leutwyler, *Annals Phys.* **158**, 142 (1984).
  - [3] J. A. Oller and E. Oset, *Nucl. Phys. A* **620**, 438-456 (1997) [erratum: *Nucl. Phys. A* **652**, 407-409 (1999)].
  - [4] J. A. Oller, E. Oset and J. R. Pelaez, *Phys. Rev. D* **59**, 074001 (1999) [erratum: *Phys. Rev. D* **60**, 099906 (1999); erratum: *Phys. Rev. D* **75**, 099903 (2007)].
  - [5] N. Kaiser, *Eur. Phys. J. A* **3**, 307-309 (1998).
  - [6] M. P. Locher, V. E. Markushin and H. Q. Zheng, *Eur. Phys. J. C* **4**, 317-326 (1998).
  - [7] J. Nieves and E. Ruiz Arriola, *Nucl. Phys. A* **679**, 57-117 (2000).
  - [8] F. K. Guo, C. Hanhart, U. G. Meißner, Q. Wang, Q. Zhao and B. S. Zou, *Rev. Mod. Phys.* **90**, 015004 (2018) [erratum: *Rev. Mod. Phys.* **94**, 029901 (2022)].
  - [9] X. K. Dong, F. K. Guo and B. S. Zou, *Commun. Theor. Phys.* **73**, 125201 (2021).
  - [10] X. K. Dong, F. K. Guo and B. S. Zou, *Few Body Syst.* **62**, 61 (2021).
  - [11] X. K. Dong, F. K. Guo and B. S. Zou, *Progr. Phys.* **41**, 65-93 (2021).
  - [12] R. Molina, T. Branz and E. Oset, *Phys. Rev. D* **82**, 014010 (2010).
  - [13] R. Aaij *et al.* [LHCb], *Phys. Rev. Lett.* **125**, 242001 (2020).
  - [14] R. Aaij *et al.* [LHCb], *Phys. Rev. D* **102**, 112003 (2020).
  - [15] R. Aaij *et al.* [LHCb], *Nature Phys.* **18**, 751-754 (2022).
  - [16] R. Aaij *et al.* [LHCb], *Nature Commun.* **13**, 3351 (2022).
  - [17] M. Z. Liu, J. J. Xie and L. S. Geng, *Phys. Rev. D* **102**, 091502 (2020).
  - [18] X. G. He, W. Wang and R. Zhu, *Eur. Phys. J. C* **80**, 1026 (2020).

- [19] J. He and D. Y. Chen, *Chin. Phys. C* **45**, 063102 (2021).
- [20] Y. Huang, J. X. Lu, J. J. Xie and L. S. Geng, *Eur. Phys. J. C* **80**, 973 (2020).
- [21] M. W. Hu, X. Y. Lao, P. Ling and Q. Wang, *Chin. Phys. C* **45**, 021003 (2021).
- [22] R. Molina and E. Oset, *Phys. Lett. B* **811**, 135870 (2020).
- [23] S. S. Agaev, K. Azizi and H. Sundu, *J. Phys. G* **48**, 085012 (2021).
- [24] C. J. Xiao, D. Y. Chen, Y. B. Dong and G. W. Meng, *Phys. Rev. D* **103**, 034004 (2021).
- [25] B. Wang and S. L. Zhu, *Eur. Phys. J. C* **82**, 419 (2022).
- [26] H. X. Chen, W. Chen, R. R. Dong and N. Su, *Chin. Phys. Lett.* **37**, 101201 (2020).
- [27] Q. Xin, Z. G. Wang and X. S. Yang, [arXiv:2207.09910 [hep-ph]].
- [28] M. Karliner and J. L. Rosner, *Phys. Rev. D* **102**, 094016 (2020).
- [29] G. J. Wang, L. Meng, L. Y. Xiao, M. Oka and S. L. Zhu, *Eur. Phys. J. C* **81**, 188 (2021).
- [30] Q. Xin and Z. G. Wang, *Eur. Phys. J. A* **58**, 110 (2022).
- [31] X. Z. Ling, M. Z. Liu, L. S. Geng, E. Wang and J. J. Xie, *Phys. Lett. B* **826**, 136897 (2022).
- [32] A. Feijoo, W. H. Liang and E. Oset, *Phys. Rev. D* **104**, 114015 (2021).
- [33] S. Fleming, R. Hodges and T. Mehen, *Phys. Rev. D* **104**, 116010 (2021).
- [34] H. Ren, F. Wu and R. Zhu, *Adv. High Energy Phys.* **2022**, 9103031 (2022).
- [35] K. Chen, R. Chen, L. Meng, B. Wang and S. L. Zhu, *Eur. Phys. J. C* **82**, 581 (2022).
- [36] M. Albaladejo, *Phys. Lett. B* **829**, 137052 (2022).
- [37] M. L. Du, V. Baru, X. K. Dong, A. Filin, F. K. Guo, C. Hanhart, A. Nefediev, J. Nieves and Q. Wang, *Phys. Rev. D* **105**, 014024 (2022).
- [38] N. Santowsky and C. S. Fischer, *Eur. Phys. J. C* **82**, 313 (2022).
- [39] C. Deng and S. L. Zhu, *Phys. Rev. D* **105**, 054015 (2022).
- [40] H. W. Ke, X. H. Liu and X. Q. Li, *Eur. Phys. J. C* **82**, 144 (2022).
- [41] S. S. Agaev, K. Azizi and H. Sundu, *JHEP* **06**, 057 (2022).
- [42] Y. Kamiya, T. Hyodo and A. Ohnishi, *Eur. Phys. J. A* **58**, 131 (2022).
- [43] L. Meng, B. Wang, G. J. Wang and S. L. Zhu, arXiv:2204.08716 [hep-ph].
- [44] L. M. Abreu, arXiv:2206.01166 [hep-ph].
- [45] S. Chen, C. Shi, Y. Chen, M. Gong, Z. Liu, W. Sun and R. Zhang, arXiv:2206.06185 [hep-lat].
- [46] L. R. Dai, R. Molina and E. Oset, *Phys. Rev. D* **105**, 016029 (2022).
- [47] M. Bando, T. Kugo and K. Yamawaki, *Phys. Rept.* **164**, 217-314 (1988).
- [48] M. Harada and K. Yamawaki, *Phys. Rept.* **381**, 1-233 (2003).

- [49] U. G. Meissner, Phys. Rept. **161**, 213 (1988).
- [50] H. Nagahiro, L. Roca, A. Hosaka and E. Oset, Phys. Rev. D **79**, 014015 (2009).
- [51] S. Q. Luo, T. W. Wu, M. Z. Liu, L. S. Geng and X. Liu, Phys. Rev. D **105**, 074033 (2022).
- [52] T. W. Wu, M. Z. Liu, L. S. Geng, E. Hiyama and M. P. Valderrama, Phys. Rev. D **100**, 034029 (2019).
- [53] X. L. Ren, B. B. Malabarba, L. S. Geng, K. P. Khemchandani and A. Martínez Torres, Phys. Lett. B **785**, 112-117 (2018).
- [54] L. Ma, Q. Wang and U. G. Meißner, Chin. Phys. C **43**, 014102 (2019).
- [55] U. Ozdem, arXiv:2207.14549 [hep-ph]
- [56] T. W. Wu, M. Z. Liu and L. S. Geng, Phys. Rev. D **103**, L031501 (2021).
- [57] X. Wei, Q. H. Shen and J. J. Xie, arXiv:2205.12526 [hep-ph].
- [58] M. Z. Liu, T. W. Wu, M. Pavon Valderrama, J. J. Xie and L. S. Geng, Phys. Rev. D **99**, 094018 (2019).
- [59] S. Q. Luo, T. W. Wu, M. Z. Liu, L. S. Geng and X. Liu, Phys. Rev. D **105**, 074033 (2022).
- [60] L. Roca and E. Oset, Phys. Rev. D **82**, 054013 (2010).
- [61] J. Yamagata-Sekihara, L. Roca and E. Oset, Phys. Rev. D **82**, 094017 (2010) [erratum: Phys. Rev. D **85**, 119905 (2012)].
- [62] C. W. Xiao, M. Bayar and E. Oset, Phys. Rev. D **86**, 094019 (2012).
- [63] D. Gamermann, J. Nieves, E. Oset and E. Ruiz Arriola, Phys. Rev. D **81**, 014029 (2010).
- [64] T. W. Wu, Y. W. Pan, M. Z. Liu and L. S. Geng, arXiv:2208.00882 [hep-ph].
- [65] R. Molina, D. Nicmorus and E. Oset, Phys. Rev. D **78**, 114018 (2008).
- [66] W. H. Liang, C. W. Xiao and E. Oset, Phys. Rev. D **89**, 054023 (2014).
- [67] E. Oset and L. Roca, arXiv:2207.08538 [hep-ph].