

Spectroscopy behavior of fully heavy tetraquarks

Hong-Tao An^{1,2,*}, Si-Qiang Luo^{1,2,†}, Zhan-Wei Liu^{1,2,3,‡} and Xiang Liu^{1,2,3,§}

¹*School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China*

²*Research Center for Hadron and CSR Physics, Lanzhou University
and Institute of Modern Physics of CAS, Lanzhou 730000, China*

³*Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou, Gansu 730000, China*

(Dated: August 28, 2023)

Stimulated by the observation of the $X(6900)$ from LHCb in 2020 and the recent results from CMS and ATLAS in the di- J/ψ invariant mass spectrum, in this work we systematically study all possible configurations for the ground fully heavy tetraquark states in the constituent quark model. By our calculation, we present their spectroscopy behaviors such as binding energy, lowest meson-meson thresholds, specific wave function, magnetic moment, transition magnetic moment, radiative decay width, rearrangement strong width ratio, internal mass contributions, relative lengths between (anti)quarks, and the spatial distribution of four valence (anti)quarks. We cannot find a stable S-wave state for the fully heavy tetraquark system. We hope that our results will be valuable to further experimental exploration of fully heavy tetraquark states.

I. INTRODUCTION

At the birth of the quark model [1–3], exotic states beyond conventional hadrons were proposed. The search for exotic hadronic states is full of challenges and opportunities. Since the $X(3872)$ was first reported by the Belle Collaboration in 2003 [4–6], a series of charmonium-like or bottomonium-like exotic states [7–21] and P_c states [22–24] have been observed experimentally, which stimulates extensive discussions of their properties by introducing the assignments of conventional hadron, compact multi-quark states, molecular state, hybrid, glueball, and kinematic effects [7–21].

In 2003, the BaBar Collaboration observed a narrow heavy-light state $D_{s0}^*(2317)$ in the $D_s^+\pi^0$ invariant mass spectrum [25]. However, since the mass of the observed $D_{s0}^*(2317)$ is about 100 MeV below the quark model predictions in Ref.[26], it is difficult to understand the $D_{s0}^*(2317)$ in conventional quark model directly, which is the “low-mass puzzle” of $D_{s0}^*(2317)$. In order to solve the low-mass puzzle, the tetraquark explanation with the $Qq\bar{q}\bar{q}$ configuration was proposed in Refs. [27–29]. Later, the CLEO Collaboration [30] confirmed the $D_{s0}^*(2317)$ and announced another narrow resonance $D_{s1}(2460)$ in the $D_s^+\pi^0$ final states. Low-mass puzzle also happens to the $D_{s1}(2460)$ [26]. The discussion of the $D_{s1}(2460)$ as tetraquark state can be found in Refs. [31–38]. In particular, the LHCb Collaboration reported the discovery of two new exotic structures $X_0(2900)$ and $X_1(2900)$ [39, 40], which inspired the study of exotic charmed tetraquarks [41–50]. In addition, the theorists also began to study the doubly charmed tetraquark states early [51–55]. In 2017, the LHCb Collaboration observed a doubly charmed baryon $\Xi_{cc}^{++}(3620)$ in the $\Lambda_c^+K^-\pi^+\pi^+$

decay mode [56]. Using the $\Xi_{cc}^{++}(3620)$ as the scaling point, the theorists further explored the possible stable doubly charmed tetraquark states with the $QQ\bar{q}\bar{q}$ configuration [57–66]. Surprisingly, as a candidate of doubly charmed tetraquark, the T_{cc}^+ was detected by LHCb in the $D^0D^0\pi^+$ invariant mass spectrum, which has a minimal quark configuration of $cc\bar{u}\bar{d}$ [67]. In addition to these singly and doubly charmed tetraquarks, there should be a triply charmed tetraquark. To our knowledge, the triply charmed tetraquark states with the $QQ\bar{Q}\bar{q}$ configuration have also been studied by various approaches [68–72].

Briefly reviewing the status of heavy flavor tetraquark states, we must mention the fully heavy tetraquark with the $QQ\bar{Q}\bar{Q}$ configuration, which has attracted the attention of both theorists and experimentalists. Chao et al. suggested that the peculiar resonance-like structures of $R(e^+e^- \rightarrow \text{hadrons})$ for $\sqrt{s} = 6 - 7$ GeV may be due to the production of the predicted P-wave (cc)-($\bar{c}\bar{c}$) states in the energy range of 6.4 – 6.8 GeV, which could dominantly decay into charmed mesons [73]. The calculation of the fully heavy tetraquark was then carried out using the potential model [74, 75] and the MIT bag model with the Born-Oppenheimer approximation [76]. This system has also been studied in a non-relativistic potential model, where no $QQ\bar{Q}\bar{Q}$ bound state can be found [77]. However, Lloyd *et al.* adopted a parameterized non-relativistic Hamiltonian to study such system [78], where they found several closely-lying bound states with a large oscillator basis. Later, Karliner *et al.* estimated the masses of the fully heavy tetraquark states by a simple quark model, and obtained $M(X_{cc\bar{c}\bar{c}}) = 6192 \pm 25$ MeV and $M(X_{bb\bar{b}\bar{b}}) = 18826 \pm 25$ MeV for the fully charmed and fully bottom tetraquarks with the $J^{PC} = 0^{++}$ quantum number, respectively [79]. Anwar *et al.* have calculated the ground-state energy of the $bb\bar{b}\bar{b}$ bound state in a nonrelativistic effective field theory with one-gluon-exchange (OGE) color Coulomb interaction, and the ground state $bb\bar{b}\bar{b}$ tetraquark mass is predicted to be (18.72 ± 0.02) GeV [80]. In Ref. [81], Bai *et al.* presented a calculation of the $bb\bar{b}\bar{b}$ tetraquark ground-state

* anht14@lzu.edu.cn

† luosq15@lzu.edu.cn

‡ liuzhanwei@lzu.edu.cn

§ xiangliu@lzu.edu.cn

energy using a diffusion Monte Carlo method to solve the non-relativistic many-body system. Debastiani *et al.* extended the updated Cornell model to study the fully charmed tetraquark in a diquark-antidiquark configuration [82]. Chen *et al.* used a moment QCD sum rule method to give the existence of the exotic states $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ in the compact diquark-antidiquark configuration, where they suggested to search for them in the $J/\psi J/\psi$ and $\eta_c(1S)\eta_{1S}$ channels [83].

With the accumulation of experimental data, many collaborations have tried to search for it. The CMS Collaboration reported the first observation of the $\Upsilon(1S)$ pair production in pp collisions, where there is evidence for a structure around 18.4 GeV with a global significance of 3.6σ exists in the four-lepton channel, which is probably a fully-bottom tetraquark state [84]. However, this structure was not confirmed by the later CMS analysis [85]. Subsequently, the LHCb Collaboration studied the $\Upsilon(1S)_{\mu^+\mu^-}$ invariant mass distribution to search for a possible $bb\bar{b}\bar{b}$ exotic meson, but they did not see any significant excess in the range 17.5 – 20.0 GeV [86]. By 2020, the LHCb Collaboration declared a narrow resonance $X(6900)$ in the di- J/ψ mass spectrum with a significance of more than 5σ [87]. In addition, a broad structure ranging from 6.2 to 6.8 GeV and an underlying peak near 7.3 GeV were also reported at the same time [87]. Recently, the ATLAS and CMS collaborations published their measurements on the di- J/ψ invariant mass spectrum. Here, they not only confirmed the existence of the $X(6900)$, but also found some new peaks [88–90]. There have been extensive discussions about the observed $X(6900)$ from different approaches and with different assignments [91–108].

The problem of the stability of the fully heavy tetraquark state has been debated for a long time. Debastiani *et al.* found that the lowest S-wave $cc\bar{c}\bar{c}$ tetraquarks may be below the dicharmonium thresholds in their updated Cornell model [109]. The 1^+ $bb\bar{b}\bar{c}$ state is thought to be a narrow state in the extended chromomagnetic model [110]. However, many other studies have suggested that the ground state of fully heavy tetraquarks is above the di-meson threshold. Wang *et al.* also calculated the fully-heavy tetraquark state in two nonrelativistic quark models with different OGE Coulomb, linear confinement and hyperfine potentials [111]. Based on the numerical calculations, they suggested that the ground states should be located about 300 – 450 MeV above the lowest scattering states, indicating that there is no bound tetraquark state. The lattice nonrelativistic QCD method was applied to study the lowest energy eigenstate of the $bb\bar{b}\bar{b}$ system, and no state was found below the lowest bottomonium-pair threshold [112]. Furthermore, Richard *et al.* also claimed that the fully heavy configuration $QQ\bar{Q}\bar{Q}$ is not stable if one adopts a standard quark model and treats the four-body problem appropriately [113]. Jin *et al.* studied full-charm and full-bottom tetraquarks using the quark delocalization color screening model and the chiral quark model, respectively, and

the results within the quantum numbers $J^P = 0^+, 1^+$, and 2^+ show that the bound state exists in both models [114]. Frankly speaking, theorists have not come to an agreement on the stability of the fully heavy tetraquark state.

Facing the present status of fully heavy tetraquark, in this work we adopt the variational method to systematically study the fully heavy tetraquark states, where the mass spectrum of the fully heavy tetraquark is given in the framework of nonrelativistic quark model associated with a potential containing Coulomb, linear, and hyperfine terms. The constructed total wave functions involved in these discussed systems satisfy the requirement of the Pauli principle. We should emphasize that we can also reproduce the masses of these conventional hadrons with the same parameters, which is a test of our adopted framework. With this preparation, we calculate the binding energies, the lowest meson-meson thresholds, and the rearrangement strong width ratio, and study the stability of the fully heavy tetraquark states against the decay into two meson states. Furthermore, we discuss whether the discussed tetraquarks have a compact configuration based on the eigenvalue of the hyperfine potential matrix. According to specific wave functions, we could obtain the magnetic moments, transition magnetic moments, and radiative decay widths, which may reflect their electromagnetic properties and internal structures. We also give the size of the tetraquarks, the relative distances between (anti)quarks, and the spatial distribution of the four valence (anti)quarks for each state. Through the present systematic work, we can test whether compact bound fully heavy tetraquarks exist within the given Hamiltonian.

This paper is organized as follows. After the introduction, we present the Hamiltonian of the constituent quark model and list the corresponding parameters in Section II. Then we give the spatial function with a simple Gaussian form and construct the flavor, color, and spin wave functions of the fully heavy tetraquark states (see Section III). In Section IV, we show the numerical results obtained by the variational method and further calculate their magnetic moment, transition magnetic moment, radiative decay width, rearrangement strong width ratio, the internal mass contributions and relative lengths between (anti) quarks. In addition, a comparison of our results with those of other theoretical groups is made in Section V. Finally, the paper ends with a short summary in Section VI.

II. HAMILTONIAN

We choose a nonrelativistic Hamiltonian for the fully heavy tetraquark system, which is written as,

$$H = \sum_{i=1}^4 \left(m_i + \frac{\mathbf{p}_i^2}{2m_i} \right) - \frac{3}{4} \sum_{i<j}^4 \frac{\lambda_i^c}{2} \cdot \frac{\lambda_j^c}{2} (V_{ij}^{Con} + V_{ij}^{SS}). \quad (1)$$

Here, m_i is the (anti)quark mass, λ_i^c is the $SU(3)$ color operator for the i -th quark, and for the antiquark, λ_i^c is replaced by $-\lambda_i^{c*}$. The internal quark potentials V_{ij}^{Con} and V_{ij}^{SS} have the following forms:

$$\begin{aligned} V_{ij}^{Con} &= -\frac{\kappa}{r_{ij}} + \frac{r_{ij}}{a_0^2} - D, \\ V_{ij}^{SS} &= \frac{\kappa'}{m_i m_j} \frac{1}{r_{0ij} r_{ij}} e^{-r_{ij}^2/r_{0ij}^2} \vec{\sigma}_i \cdot \vec{\sigma}_j, \end{aligned} \quad (2)$$

where $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$ is the distance between the i -th (anti)quark and the j -th (anti)quark, and the σ_i is the $SU(2)$ spin operator for the i -th quark. As for the r_{0ij} and κ' , we have

$$\begin{aligned} r_{0ij} &= 1/\left(\alpha + \beta \frac{m_i m_j}{m_i + m_j}\right), \\ \kappa' &= \kappa_0 \left(1 + \gamma \frac{m_i m_j}{m_i + m_j}\right). \end{aligned} \quad (3)$$

The corresponding parameters appearing in Eqs. (2-3) are shown in Table I. Here, κ and κ' are the couplings of the Coulomb and hyperfine potentials, respectively, and they are proportional to the running coupling constant $\alpha_s(r)$ of QCD. The Coulomb and hyperfine interactions can be deduced from the one-gluon-exchange model. $1/a_0^2$ represents the strength of the linear potential. r_{0ij} is the Gaussian-smearing parameter. Furthermore, we introduce κ_0 and γ in κ' to better describe the interaction between different quark pairs [115].

TABLE I. Parameters of the Hamiltonian.

Parameter	κ	a_0	D
Value	120.0 MeV fm	$0.0318119 (\text{MeV}^{-1}\text{fm})^{1/2}$	983 MeV
Parameter	α	β	m_c
Value	1.0499 fm^{-1}	$0.0008314 (\text{MeVfm})^{-1}$	1918 MeV
Parameter	κ_0	γ	m_b
Value	194.144 MeV	0.00088 MeV^{-1}	5343 MeV

III. WAVE FUNCTIONS

Here, we focus on the ground fully heavy tetraquark states. We present the flavor, spatial, and color-spin parts of the total wave function for fully-heavy tetraquark system. In order to consider the constraint by the Pauli principle, we use a diquark-antidiquark picture to analyze this tetraquark system.

A. Flavor Part

First we discuss the flavor part. Here, we list all the possible flavor combinations for the fully-heavy

TABLE II. All possible flavor combinations for the fully-heavy tetraquark system.

System	Flavor combinations		
$QQ\bar{Q}\bar{Q}$	$cc\bar{c}\bar{c}$	$bb\bar{b}\bar{b}$	$cb\bar{c}\bar{b}$
	$cc\bar{b}\bar{b} (bb\bar{c}\bar{c})$	$cc\bar{c}\bar{b} (bc\bar{c}\bar{c})$	$bb\bar{b}\bar{c} (cb\bar{b}\bar{b})$

tetraquark system in Table II.

In Table II, the three flavor combinations in the first row are purely neutral particles and the C parity is a “good” quantum number. For the other six states in the second row, each state has a charge conjugation anti-partner, and their masses, internal mass contributions, relative distances between (anti)quarks are absolutely identical, so we only need to discuss one of the pair.

Furthermore, the $cc\bar{c}\bar{c}$, $bb\bar{b}\bar{b}$, and $cc\bar{b}\bar{b}$ states have the two pairs of (anti)quarks which are identical, but only the first two quarks in the $cc\bar{c}\bar{b}$ and $bb\bar{b}\bar{c}$ states are identical.

B. Spatial Part

In this part, we construct the wave function for the spatial part in a simple Gaussian form. We denote the fully heavy tetraquark state as the $Q(1)Q(2)\bar{Q}(3)\bar{Q}(4)$ configuration, and choose the Jacobian coordinate system as follows:

$$\begin{aligned} \mathbf{x}_1 &= \sqrt{1/2}(\mathbf{r}_1 - \mathbf{r}_2), \\ \mathbf{x}_2 &= \sqrt{1/2}(\mathbf{r}_3 - \mathbf{r}_4), \\ \mathbf{x}_3 &= \left[\left(\frac{m_1 \mathbf{r}_1 + m_2 \mathbf{r}_2}{m_1 + m_2} \right) - \left(\frac{m_3 \mathbf{r}_3 + m_4 \mathbf{r}_4}{m_3 + m_4} \right) \right]. \end{aligned} \quad (4)$$

Here, we set the Jacobi coordinates with the following conditions:

$$\begin{aligned} m_1 = m_2 = m_3 = m_4 = m_c, & \quad \text{for } cc\bar{c}\bar{c}, \\ m_1 = m_2 = m_3 = m_4 = m_b, & \quad \text{for } bb\bar{b}\bar{b}, \\ m_1 = m_2 = m_c, m_3 = m_4 = m_b, & \quad \text{for } cc\bar{b}\bar{b}, \\ m_1 = m_2 = m_3 = m_c, m_4 = m_b, & \quad \text{for } cc\bar{c}\bar{b}, \\ m_1 = m_2 = m_3 = m_b, m_4 = m_c, & \quad \text{for } bb\bar{b}\bar{c}, \\ m_1 = m_c, m_2 = m_b, m_3 = m_c, m_4 = m_b, & \quad \text{for } cb\bar{c}\bar{b}. \end{aligned}$$

Based on this, we construct the spatial wave functions of the $QQ\bar{Q}\bar{Q}$ states in a single Gaussian form. The spatial wave function can satisfy the required symmetry property:

$$R^s = \exp[-C_{11}\mathbf{x}_1^2 - C_{22}\mathbf{x}_2^2 - C_{33}\mathbf{x}_3^2], \quad (5)$$

where C_{11} , C_{22} , and C_{33} are the variational parameters.

It is also useful to introduce the center of mass frame so that the kinetic term in the Hamiltonian of Eq. (1)

can be reduced appropriately for our calculations. The kinetic term, denoted by T_c , is as follows

$$T_c = \sum_{i=1}^4 \frac{\mathbf{p}_i^2}{2m_i} - \frac{\mathbf{p}_{rC}^2}{2M} = \frac{\mathbf{p}_{x_1}^2}{2m'_1} + \frac{\mathbf{p}_{x_2}^2}{2m'_2} + \frac{\mathbf{p}_{x_3}^2}{2m'_3}, \quad (6)$$

where different states have different reduced masses m'_i , which are listed in Table III.

TABLE III. The reduced mass m'_i in different states.

States	m'_1	m'_2	m'_3	States	m'_1	m'_2	m'_3
$cc\bar{c}\bar{c}$	m_c	m_c	m_c	$cc\bar{c}\bar{b}$	m_c	$\frac{2m_cm_b}{m_c+m_b}$	$\frac{(m_c+m_b)m_c}{2(3m_c+m_b)}$
$bb\bar{b}\bar{b}$	m_b	m_b	m_b	$bb\bar{b}\bar{c}$	m_b	$\frac{2m_cm_b}{m_c+m_b}$	$\frac{(m_c+m_b)m_b}{2(3m_b+m_c)}$
$cc\bar{b}\bar{b}$	m_c	m_b	$\frac{2m_cm_b}{m_c+m_b}$	$cb\bar{c}\bar{b}$	$\frac{2m_cm_b}{m_c+m_b}$	$\frac{2m_cm_b}{m_c+m_b}$	$\frac{m_c+m_b}{2}$

C. Color-spin Part

In the color space, the color wave functions can be analyzed using the SU(3) group theory, where the direct product of the diquark and antidiquark components reads

$$(3_c \otimes 3_c) \otimes (\bar{3}_c \otimes \bar{3}_c) = (6_c \oplus \bar{3}_c) \otimes (\bar{6}_c \oplus 3_c). \quad (7)$$

Based on this, we get two types of color-singlet states:

$$\phi_1 = |(Q_1 Q_2)^{\bar{3}} (\bar{Q}_3 \bar{Q}_4)^3\rangle, \phi_2 = |(Q_1 Q_2)^6 (\bar{Q}_3 \bar{Q}_4)^{\bar{6}}\rangle. \quad (8)$$

In the spin space, the allowed wave functions are in the diquark-antidiquark picture:

$$\begin{aligned} \chi_1 &= |(Q_1 Q_2)_1 (\bar{Q}_3 \bar{Q}_4)_1\rangle_2, \chi_2 = |(Q_1 Q_2)_1 (\bar{Q}_3 \bar{Q}_4)_1\rangle_1, \\ \chi_3 &= |(Q_1 Q_2)_1 (\bar{Q}_3 \bar{Q}_4)_0\rangle_1, \chi_4 = |(Q_1 Q_2)_0 (\bar{Q}_3 \bar{Q}_4)_1\rangle_1, \\ \chi_5 &= |(Q_1 Q_2)_1 (\bar{Q}_3 \bar{Q}_4)_1\rangle_0, \chi_6 = |(Q_1 Q_2)_0 (\bar{Q}_3 \bar{Q}_4)_0\rangle_0. \end{aligned} \quad (9)$$

In the notation $|(Q_1 Q_2)_{\text{spin}1} (\bar{Q}_3 \bar{Q}_4)_{\text{spin}2}\rangle_{\text{spin}3}$, the spin1, spin2, and spin3 represent the spin of the diquark, the spin of the antidiquark, and the total spin of the tetraquark state, respectively.

Since the flavor part and spatial parts are chosen to be fully symmetric for the (anti)diquark, the color-spin part of the total wave function should be fully antisymmetric. Combining the flavor part, we show all possible color-spin part satisfying the Pauli principle with J^{PC} in Table IV.

In addition, it is convenient to consider the strong decay properties, we use the meson-meson configuration to represent color-singlet and spin wave functions, again. The color wave functions in the meson-meson configuration can be derived from the following direct product:

$$(3_c \otimes \bar{3}_c) \otimes (3_c \otimes \bar{3}_c) = (1_c \oplus 8_c) \otimes (1_c \oplus 8_c). \quad (10)$$

Based on Eq. (10), they can be expressed as

$$\psi_1 = |(Q_1 \bar{Q}_3)^1 (Q_2 \bar{Q}_4)^1\rangle, \psi_2 = |(Q_1 \bar{Q}_3)^8 (Q_2 \bar{Q}_4)^8\rangle. \quad (11)$$

Similarly, the spin wave functions in the meson-meson configuration read as

$$\begin{aligned} \zeta_1 &= |(Q_1 \bar{Q}_3)_1 (Q_2 \bar{Q}_4)_1\rangle_2, \zeta_2 = |(Q_1 \bar{Q}_3)_0 (Q_2 \bar{Q}_4)_1\rangle_1, \\ \zeta_3 &= |(Q_1 \bar{Q}_3)_1 (Q_2 \bar{Q}_4)_0\rangle_1, \zeta_4 = |(Q_1 \bar{Q}_3)_1 (Q_2 \bar{Q}_4)_1\rangle_1, \\ \zeta_5 &= |(Q_1 \bar{Q}_3)_1 (Q_2 \bar{Q}_4)_1\rangle_0, \zeta_6 = |(Q_1 \bar{Q}_3)_0 (Q_2 \bar{Q}_4)_0\rangle_0. \end{aligned} \quad (12)$$

TABLE IV. The allowed color-spin parts for each flavor configuration.

Type	$J^{P(C)}$	Color-spin Part			
$cc\bar{c}\bar{c}$	$2^{+(+)}$	$\phi_1 \chi_1$			
$bb\bar{b}\bar{b}$	$1^{+(-)}$	$\phi_1 \chi_2$			
$cc\bar{b}\bar{b}$	$0^{+(+)}$	$\phi_1 \chi_5$	$\phi_2 \chi_6$		
$cc\bar{c}\bar{b}$	2^+	$\phi_1 \chi_1$			
$bb\bar{b}\bar{c}$	1^+	$\phi_1 \chi_2$	$\phi_1 \chi_3$	$\phi_2 \chi_4$	
	0^+	$\phi_1 \chi_5$	$\phi_2 \chi_6$		
$cb\bar{c}\bar{b}$	2^{++}	$\phi_1 \chi_1$	$\phi_2 \chi_1$		
	1^{+-}	$\phi_1 \chi_2$	$\phi_2 \chi_2$	$\frac{1}{\sqrt{2}}(\phi_1 \chi_3 + \phi_1 \chi_4)$	
		$\frac{1}{\sqrt{2}}(\phi_2 \chi_3 + \phi_2 \chi_4)$			
	1^{++}	$\frac{1}{\sqrt{2}}(\phi_1 \chi_3 - \phi_1 \chi_4)$	$\frac{1}{\sqrt{2}}(\phi_2 \chi_3 - \phi_2 \chi_4)$		
0^{++}	$\phi_1 \chi_5$	$\phi_2 \chi_5$	$\phi_1 \chi_6$	$\phi_2 \chi_6$	

IV. NUMERICAL ANALYSIS

1. Mass spectrum, internal contribution, and spatial size

In this subsection, we check the consistency between the experimental masses and the obtained masses of traditional hadrons using the variational method based on the Hamiltonian of Eq. (1) and the parameters in Table I. We show the results in Table VI and note that our values are relatively reliable since the deviations for most states are less than 20 MeV.

In addition, we have systematically constructed the total wave function satisfied by the Pauli principle in the previous section. The corresponding total wave function could be expanded as follows:

$$|\Psi_\alpha\rangle = \sum_{ij} C_{ij}^\alpha |F\rangle |R^s\rangle |\phi_i \chi_j\rangle. \quad (13)$$

To study the mass of the fully heavy tetraquarks with the variational method, we calculate the Schrödinger

equation $H|\Psi_\alpha\rangle = E_\alpha|\Psi_\alpha\rangle$, diagonalize the corresponding matrix, and then determine the ground state masses for the fully heavy tetraquarks. According to the corresponding variational parameters, we also give the internal mass contributions, including the quark mass part, the kinetic energy part, the confinement potential part, and the hyperfine potential part. For comparison, we also show the lowest meson-meson thresholds for the tetraquarks with different quantum numbers and their internal contributions. This is how we define the binding energy:

$$B_T = M_{tetraquark} - M_{meson1} - M_{meson2}, \quad (14)$$

where $M_{tetraquark}$, M_{meson1} , and M_{meson2} are the masses of the tetraquark and the two mesons at the lowest threshold allowed in the rearrangement decay of the tetraquark, respectively. To facilitate the discussion in the next subsection, we also define the V^C , which is the sum of the Coulomb potential and the linear potential.

Here, it is also useful to investigate the spatial size of the tetraquarks, which is strongly related to the magnitude of the various kinetic energies and the potential energies between the quarks. It is also important to understand the relative lengths between the quarks in the tetraquarks and their lowest thresholds, and the relative distance between the heavier quarks is generally shorter than that between the lighter quarks [61]. This tendency is also maintained in each tetraquark state according to the corresponding tables.

2. Magnetic moments, transition magnetic moments, and radiative decay widths

The magnetic moment of hadrons is a physical quantity that reflects their internal structures [121]. The total magnetic moment $\vec{\mu}_{total}$ of a compound system contains the spin magnetic moment $\vec{\mu}_{spin}$ and the orbital magnetic moment $\vec{\mu}_{orbital}$ from all of its constituent quarks. For ground hadron states, their contribution of the orbital magnetic moment $\vec{\mu}_{orbital}$ is zero, and so we only concentrate on the spin magnetic moment $\vec{\mu}_{spin}$. The explicit expression for the spin magnetic moment $\vec{\mu}_{spin}$ is written as

$$\vec{\mu}_{spin} = \sum_i \mu_i \vec{\sigma}_i = \sum_i \frac{Q_i^{eff}}{2M_i^{eff}} \vec{\sigma}_i, \quad (15)$$

where Q_i^{eff} and M_i^{eff} are the effective charge and effective mass of the i -th constituent quark, respectively. The $\vec{\sigma}_i$ denotes the Pauli's spin matrix of the i -th constituent quark. According to Ref. [122], the effective charge of the quark is affected by other quarks in the inner hadron. We now assume that the effective charge is linearly dependent on the charge of the shielding quarks. So the effective charge Q_i^{eff} is defined as

$$Q_i^{eff} = Q_i + \sum_{i \neq j} \alpha_{ij} Q_j, \quad (16)$$

where Q_i is the bare charge of the i -th constituent quark, the α_{ij} is a corrected parameter that reflects how much the charge of other quarks affects the charge of the i -th quark. To simplify the calculation, we also set α_{ij} always equal to 0.033 according to the Ref. [122]. The effective quark masses M_i^{eff} contain the contributions from both the bare quark mass terms and the interaction terms in the chromomagnetic model, and their values are taken from Ref. [123].

To obtain the magnetic moment of the discussed hadron, we calculate the z -component of the magnetic moment operator $\hat{\mu}^z$ sandwiched by the corresponding total wave function Ψ_α (Eq.[9]). Now, only the spin part of the total wave function is involved. The total spin wave functions of the discussed hadrons are written as

$$|\chi_{total}\rangle = C_1\chi_1 + C_2\chi_2 + \dots \quad (17)$$

Based on this, we can quantitatively obtain the magnetic moment of the discussed hadron

$$\begin{aligned} \mu &= \langle \Psi_\alpha | \hat{\mu}^z | \Psi_\alpha \rangle = \langle \chi_\alpha | \hat{\mu}^z | \chi_\alpha \rangle \\ &= C_1^2 \mu(\chi_1) + C_2^2 \mu(\chi_2) + \dots + 2C_1 C_2 \mu^{tr}(\chi_1, \chi_2) + \dots \end{aligned}$$

where μ^{tr} is the cross-term representing the transition moment, and C_1, C_2 are the eigenvectors of the given mixing state [124]. Similarly, the transition magnetic moments between the hadrons can be obtained as $\mu_{H' \rightarrow H\gamma} = \langle \Psi_{H'} | \hat{\mu}^z | \Psi_H \rangle$.

According to Eq. (18), the numerical values for the magnetic moments of the traditional hadrons have been listed in Table VI. Here, $\mu_N = e/2m_N$ is the nuclear magnetic moment with $m_N = 938$ MeV as the nuclear mass, which is the unit of the magnetic moment. For comparison, we also show the experimental values and other theoretical results from Refs. [121, 122, 124–127]. Because of the $\mu_Q = -\mu_{\bar{Q}}$, the magnetic moment of all of the $J^P = 0^+$ ground mesons and tetraquarks and the ground states with certain C -parity is 0.

The decay property is another important aspect to investigate the nature of the exotic hadron. According to the transition magnetic moments in the above subsection, we can further obtain the radiative decay widths around fully heavy tetraquarks [128–137].

$$\Gamma = \frac{|\mathbf{k}|^2}{\pi} \frac{2}{2J_i + 1} \frac{M_f}{M_i} \sum_{M_{J_f}, M_{J_i}} |\mathcal{M}_{M_{J_f}, M_{J_i}}|^2, \quad (18)$$

where the J_i and J_f are the total angular momenta of the initial and final hadrons, respectively. The M_i and M_f in Eq. (18) represent initial and final hadron masses, respectively.

3. Relative decay widths of tetraquarks

In addition to radiative decay, we also consider the rearrangement strong decay properties for fully heavy

tetraquarks. Based on Eqs. (10-12), the color wave function also falls into two categories: the color-singlet $\psi_1 = |(Q_1\bar{Q}_3)^1(Q_2\bar{Q}_4)^1\rangle$ which can easily decay into two S-wave mesons, and the color-octet $\psi_2 = |(Q_1\bar{Q}_3)^8(Q_2\bar{Q}_4)^8\rangle$ which can only fall apart by the gluon exchange. Thus we transform the total wave functions Ψ_α into the new configuration,

$$|\Psi_\alpha\rangle = \sum_{ij} C'_{ij}{}^\alpha |F\rangle |R^s\rangle |[\psi_i\zeta_j]\rangle. \quad (19)$$

Among the decay behaviors of the tetraquarks, one decay mode is that the quarks simply fall apart into the final decay channels without quark pair creations or annihilations, which is donated as ‘‘Okubo-Zweig-Iizuka (OZI)-superallowed’’ decays. In this part, we will only focus this type of decay channels. For two body decay by L-wave, the partial decay width reads as [72, 110, 138–140]:

$$\Gamma_i = \gamma_i \alpha \frac{k^{2L+1}}{m^{2L}} |c_i|^2, \quad (20)$$

where α is an effective coupling constant, c_i is the overlap corresponding exactly to $C'_{ij}{}^\alpha$ of Eq. (19), m is the mass of the initial state, k is the momentum of the final state in the rest frame of the initial state. For the decays of the S-wave tetraquarks, $(k/m)^{-2}$ is of order $\mathcal{O}(10^{-2})$ or even smaller, so all higher-wave decays are suppressed. So we only need to consider the S-wave decays. As for γ_i , it is determined by the spatial wave functions of the initial and final states, which are different for each decay process. In the quark model in the heavy quark limit, the spatial wave functions of the ground S-wave pseudoscalar and the vector meson are the same. The relations of γ_i for fully heavy tetraquarks are given in Table V. Based on this, the branching fraction is proportional to the square of the coefficient of the corresponding component in the eigenvectors, and the strong decay phase space, i.e., $k \cdot |c_i|^2$, for each decay mode. From the value of $k \cdot |c_i|^2$, one can roughly estimate the ratios of the relative decay widths between different decay processes of different initial tetraquarks.

In the following subsections, we concretely discuss all possible configurations for fully heavy tetraquarks.

A. $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ states

First we investigate the $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ systems. There are two $J^{PC} = 0^{++}$ states, one $J^{PC} = 1^{+-}$ state, and one $J^{PC} = 2^{++}$ state according to Table IV. We show the masses of the ground states, the variational parameters, the internal mass contributions, the relative lengths between the quarks, their lowest meson-meson thresholds, the specific wave function, the magnetic moments, the transition magnetic moments, the radiative decay widths, and the rearrangement strong width ratios in Tables VII-IX, respectively.

TABLE V. The approximate relation about γ_i for the $QQ\bar{Q}\bar{Q}$ system.

States	γ_i
$cc\bar{c}\bar{c}$	$\gamma_{J/\psi J/\psi} = \gamma_{\eta_c J/\psi} = \gamma_{\eta_c \eta_c}$
$bb\bar{b}\bar{b}$	$\gamma_{\Upsilon\Upsilon} = \gamma_{\eta_b \Upsilon} = \gamma_{\eta_b \eta_b}$
$cc\bar{b}\bar{b}$	$\gamma_{B_c^* B_c^*} = \gamma_{B_c B_c^*} = \gamma_{B_c B_c}$
$cc\bar{c}\bar{b}$	$\gamma_{J/\psi B_c^*} = \gamma_{J/\psi B_c} = \gamma_{\eta_c B_c^*} = \gamma_{\eta_c B_c}$
$bb\bar{b}\bar{c}$	$\gamma_{\Upsilon \bar{B}_c^*} = \gamma_{\Upsilon \bar{B}_c} = \gamma_{\eta_b \bar{B}_c^*} = \gamma_{\eta_b \bar{B}_c}$
$cb\bar{c}\bar{b}$	$\gamma_{J/\psi \Upsilon} = \gamma_{J/\psi \eta_b} = \gamma_{\eta_c \Upsilon} = \gamma_{\eta_c \eta_b}$ $\gamma_{B_c^* \bar{B}_c^*} = \gamma_{B_c^* \bar{B}_c} = \gamma_{B_c \bar{B}_c^*} = \gamma_{B_c \bar{B}_c}$

Here, we take the $J^{PC} = 0^{++}$ $bb\bar{b}\bar{b}$ ground state as an example, and others have similar discussions according to Tables VII-IX. We now analyze the numerical results obtained from the variational method. For the $J^{PC} = 0^{++}$ $bb\bar{b}\bar{b}$ ground state, its mass is 19240.0 MeV and the corresponding binding energy B_T is +461.9 MeV. Its variational parameters are given as $C_{11} = 7.7 \text{ fm}^{-2}$, $C_{22} = 7.7 \text{ fm}^{-2}$, and $C_{33} = 11.4 \text{ fm}^{-2}$, giving roughly the inverse ratios of the size for the diquark, the antidiquark, and between the center of the diquark and the antidiquark, respectively. We naturally find that the C_{11} is equal to C_{22} , so the distance of $(b - b)$ would be equal to that of $(\bar{b} - \bar{b})$, and the reason is that the $bb\bar{b}\bar{b}$ system is a neutral system.

The total wave function in the diquark-antidiquark configuration is given by

$$|\Psi_{tot}\rangle = -0.936|F\rangle |R^s\rangle |[\phi_2\chi_6]\rangle + 0.352|F\rangle |R^s\rangle |[\phi_1\chi_5]\rangle. \quad (21)$$

The meson-meson configuration is connected to the diquark-antidiquark configuration by a linear transformation. We then obtain the total wave function in the meson-meson configuration:

$$|\Psi_{tot}\rangle = 0.558|F\rangle |R^s\rangle |[\psi_1\zeta_5]\rangle + 0.560|F\rangle |R^s\rangle |[\psi_1\zeta_6]\rangle + 0.021|F\rangle |R^s\rangle |[\psi_2\zeta_5]\rangle + 0.612|F\rangle |R^s\rangle |[\psi_2\zeta_6]\rangle. \quad (22)$$

According to Eq. (22), we are sure that the overlaps c_i of $\eta_b\eta_b$ and $\Upsilon\Upsilon$ are 0.560 and 0.558, respectively. Then, based on Eq. (20), the rearrangement strong width ratios are

$$\frac{\Gamma_{T_{b^2\bar{b}^2}(19240.0,0^{++}) \rightarrow \Upsilon\Upsilon}}{\Gamma_{T_{b^2\bar{b}^2}(19240.0,0^{++}) \rightarrow \eta_b\eta_b}} = 1 : 1.2, \quad (23)$$

i.e., both the $\Upsilon\Upsilon$ and $\eta_b\eta_b$ are dominant decay channels for the $T_{b^2\bar{b}^2}(19240.0, \Upsilon\Upsilon)$ state.

TABLE VI. Masses and magnetic moments of some ground hadrons obtained from the theoretical calculations. M_{result} , μ_{result} , μ_{bag} , $\mu_{the(1)}$, and $\mu_{the(2)}$ are theoretical masses and magnetic moments for Eq. (1), Eq. (15), and Refs. [121, 122, 124], respectively. M_{exp} and μ_{exp} are the observed values of masses and magnetic moments. The masses and errors are in units of MeV. The magnetic moment is in units of the nuclear magnetic moment μ_N . The variational parameter is in units of fm^{-2} .

Hadron	Σ^+	Σ^0	Σ^-	Ξ^0	Ξ^-	Σ_c^{++}	Σ_c^+	Σ_c^0	Σ_c^{*++}	Σ_c^{*+}	Σ_c^{*0}	Σ_b^+	Σ_b^0	Σ_b^-	Σ_b^{*+}	Σ_b^{*0}	Σ_b^{*-}
M_{result}	1187.7		1295.4			2445.2			2518.3			5832.1			5860.8		
Parameters	2.1		3.3			2.1			2.0			2.1			2.0		
	3.1		2.9			3.7			3.4			4.0			3.4		
M_{exp}	1189.4		1314.9			2454.0			2851.4			5811.3			5832.1		
Error	-1.7		-19.5			-8.8			-0.1			20.8			28.7		
μ_{result}	2.53	0.75	-1.04	-1.31	-0.52	2.34	0.55	-1.24	4.09	1.38	-1.32	2.37	0.58	-1.21	3.48	0.78	-1.92
μ_{bag} [124]	2.72	0.86	-1.01	-1.58	-0.64	2.13	0.41	-1.31	4.07	1.39	-1.29	2.23	0.58	-1.07	3.29	0.76	-1.77
$\mu_{the(1)}$ [121]	2.74	0.84	-1.06	-1.47	-0.52	2.36	0.50	-1.37	4.09	1.30	-1.49						
$\mu_{the(2)}$ [122]	2.46	0.47	-1.10	-1.61	-0.65	3.57	1.96	0.04									
μ_{exp}	2.46		-1.16	-1.25	-0.65							6.14	2.70			-2.02	
Hadron	D^{*0}	D^{*+}	D_s^{*+}	B^{*0}	B^{*+}	B_s^{*0}	J/ψ	η_c	Υ	η_b	B_c	B_c^*	Δ^{++}	Δ^+	Δ^0	Δ^0	Ω^-
M_{result}	1996.9	2093.3	5363.6	5434.7	3092.2	2998.5	9468.9	9389.0	6287.9	6350.5	1245.6			1675.8			
Parameters	3.8	6.2	4.2	7.5	12.5	15.0	49.7	57.4	22.9	20.2	1.8			3.3			
M_{exp}	2010.3	2112.2	5324.7	5415.4	3096.9	2983.9	9460.3	9399.0	6274.9	(6332)	1232.0			1672.5			
Error	-13.4	-18.9	38.8	19.3	-4.7	14.6	8.6	10.0	13.0	(17.5)	13.6			3.1			
μ_{result}	-1.37	1.24	1.00	-0.78	1.83	0.51	0	-	0	-	0.44	5.57	-2.78	0	-2.78	-1.86	
μ_{bag} [124]	-0.98	1.21	1.08	-0.53	1.21	1.01	0	-	0	-	0.52	5.70	2.85	0	-2.85	-2.20	
$\mu_{the(1)}$ [121]	-1.49	1.30	1.07				0	-				5.58	2.79	0	-2.79	-1.88	

As for the magnetic moments of the $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ ground states, their values are all 0, because the same quark and antiquark have exactly opposite magnetic moments, which cancel each other out. We also discuss the transition magnetic moment of the $T_{b_2\bar{b}_2}(19303.9, 1^{+-}) \rightarrow T_{b_2\bar{b}_2}(19240.0, 0^{++})\gamma$ process. We construct their flavor \otimes spin wave functions as

$$\begin{aligned}
|\Psi\rangle_{T_{b_2\bar{b}_2}(1^{+-})}^{S=1;S_s=0} &= |R^s\rangle|\psi\rangle|bb\bar{b}\bar{b}\rangle\frac{1}{\sqrt{2}}(\uparrow\uparrow\downarrow\downarrow - \downarrow\downarrow\uparrow\uparrow) \\
|\Psi\rangle_{T_{b_2\bar{b}_2}(0^{++})}^{S=0;S_s=0} &= |R^s\rangle|\psi\rangle|bb\bar{b}\bar{b}\rangle|0.352\frac{1}{\sqrt{3}}(\uparrow\uparrow\downarrow\downarrow + \downarrow\downarrow\uparrow\uparrow) + \dots.
\end{aligned} \tag{24}$$

And then, the transition magnetic momentum of the $T_{b_2\bar{b}_2}(19303.9, 1^{+-}) \rightarrow T_{b_2\bar{b}_2}(19240.0, 0^{++})\gamma$ process can be given by the z -component of the magnetic moment operator $\hat{\mu}^z$ sandwiched by the flavor-spin wave functions of the $T_{b_2\bar{b}_2}(19303.9, 1^{+-})$ and $T_{b_2\bar{b}_2}(19240.0, 0^{++})$. So, the corresponding transition magnetic momentum is

$$\begin{aligned}
\mu_{T_{b_2\bar{b}_2}(1^{+-}) \rightarrow T_{b_2\bar{b}_2}(0^{++})} &= \langle \Psi_{tot}^{1^{+-}} | \hat{\mu}^z | \Psi_{tot}^{0^{++}} \rangle \\
&= 0.352 \times \frac{1}{\sqrt{6}}(4\mu_b - 4\mu_{\bar{b}}) = -0.072 \mu_N.
\end{aligned} \tag{25}$$

As for the transition magnetic moment of the $T_{b_2\bar{b}_2}(19327.9, 2^{++}) \rightarrow T_{b_2\bar{b}_2}(19240.0, 0^{++})\gamma$ process, its value is 0 due to the C parity conservation restriction.

Furthermore, according to Eq. (18) and Eq. (25), we also obtain the corresponding radiative decay widths

$$\Gamma_{T_{b_2\bar{b}_2}(19303.9, 1^{+-}) \rightarrow T_{b_2\bar{b}_2}(19240.0, 0^{++})\gamma} = 2.8 \text{ keV}, \tag{26}$$

$$\Gamma_{T_{b_2\bar{b}_2}(19327.9, 2^{++}) \rightarrow T_{b_2\bar{b}_2}(19240.0, 0^{++})\gamma} = 0 \text{ keV}. \tag{27}$$

1. Relative distances and symmetry

Here, we concentrate on the the relative distances between the (anti)quarks in tetraquarks. Looking at the relative distances in Table IX, we find that the relative distances of (1,2) and (3,4) pairs are the same, and other relative distances are the same in all the $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ states This is due to the permutation symmetry for the ground state wave function in each tetraquark [65]. For the $c_1c_2\bar{c}_3\bar{c}_4$ and $b_1b_2\bar{b}_3\bar{b}_4$ states, they need to satisfy the Pauli principle for identical particles are as follows:

$$A_{12}|\Psi_{tot}\rangle = A_{34}|\Psi_{tot}\rangle = -|\Psi_{tot}\rangle, \tag{28}$$

where the operator A_{ij} means exchanging the coordinate of Q_i (\bar{Q}_i) and Q_j (\bar{Q}_j).

Meanwhile, they are pure neutral particles with definite C-parity, so the permutation symmetries for total wave functions are as follows:

$$A_{12-34}|\Psi_{tot}\rangle = \pm|\Psi_{tot}\rangle, \quad (29)$$

where A_{12-34} means that the coordinates of the diquark and the antidiquark are exchanged.

Based on this, the relationship of the relative distances for all the $c_1c_2\bar{c}_3\bar{c}_4$ and $b_1b_2\bar{b}_3\bar{b}_4$ states can be obtained as follows:

$$\begin{aligned} & \langle\Psi_{tot}|\mathbf{r}_1 - \mathbf{r}_3|\Psi_{tot}\rangle \\ &= \langle\Psi_{tot}|A_{12}^{-1}A_{12}|\mathbf{r}_1 - \mathbf{r}_3|A_{12}^{-1}A_{12}|\Psi_{tot}\rangle = \langle\Psi_{tot}|\mathbf{r}_2 - \mathbf{r}_3|\Psi_{tot}\rangle \\ &= \langle\Psi_{tot}|A_{34}^{-1}A_{34}|\mathbf{r}_2 - \mathbf{r}_3|A_{34}^{-1}A_{34}|\Psi_{tot}\rangle = \langle\Psi_{tot}|\mathbf{r}_2 - \mathbf{r}_4|\Psi_{tot}\rangle \\ &= \langle\Psi_{tot}|A_{12}^{-1}A_{12}|\mathbf{r}_2 - \mathbf{r}_4|A_{12}^{-1}A_{12}|\Psi_{tot}\rangle = \langle\Psi_{tot}|\mathbf{r}_1 - \mathbf{r}_4|\Psi_{tot}\rangle, \end{aligned} \quad (30)$$

and

$$\begin{aligned} & \langle\Psi_{tot}|\mathbf{r}_1 - \mathbf{r}_2|\Psi_{tot}\rangle \\ &= \langle\Psi_{tot}|A_{12-34}^{-1}A_{12-34}|\mathbf{r}_1 - \mathbf{r}_2|A_{12-34}^{-1}A_{12-34}|\Psi_{tot}\rangle \\ &= \langle\Psi_{tot}|\mathbf{r}_3 - \mathbf{r}_4|\Psi_{tot}\rangle. \end{aligned} \quad (31)$$

Obviously, our theoretical derivations are in perfect agreement with the calculated results in Table IX.

We can also prove three Jacobi coordinates, $\mathbf{R}_{1,2} = \mathbf{r}_1 - \mathbf{r}_2$, $\mathbf{R}_{3,4} = \mathbf{r}_3 - \mathbf{r}_4$, and $\mathbf{R}' = 1/2(\mathbf{r}_1 + \mathbf{r}_2 - \mathbf{r}_3 - \mathbf{r}_4)$, are orthogonal to each other for all the $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ states:

$$\begin{aligned} & \langle\Psi_{tot}|\mathbf{R}_{1,2} \cdot \mathbf{R}_{3,4}|\Psi_{tot}\rangle \\ &= \langle\Psi_{tot}|(34)^{-1}(34)|\mathbf{R}_{1,2} \cdot \mathbf{R}_{3,4}|(34)^{-1}(34)|\Psi_{tot}\rangle \\ &= -\langle\Psi_{tot}|\mathbf{R}_{1,2} \cdot \mathbf{R}_{3,4}|\Psi_{tot}\rangle = 0, \end{aligned} \quad (32)$$

$$\begin{aligned} & \langle\Psi_{tot}|\mathbf{R}_{1,2} \cdot \mathbf{R}'|\Psi_{tot}\rangle \\ &= \langle\Psi_{tot}|(12)^{-1}(12)|\mathbf{R}_{1,2} \cdot \mathbf{R}'|(12)^{-1}(12)|\Psi_{tot}\rangle \\ &= -\langle\Psi_{tot}|\mathbf{R}_{1,2} \cdot \mathbf{R}'|\Psi_{tot}\rangle = 0, \end{aligned} \quad (33)$$

and

$$\begin{aligned} & \langle\Psi_{tot}|\mathbf{R}_{3,4} \cdot \mathbf{R}'|\Psi_{tot}\rangle \\ &= \langle\Psi_{tot}|(34)^{-1}(34)|\mathbf{R}_{3,4} \cdot \mathbf{R}'|(34)^{-1}(34)|\Psi_{tot}\rangle \\ &= -\langle\Psi_{tot}|\mathbf{R}_{3,4} \cdot \mathbf{R}'|\Psi_{tot}\rangle = 0. \end{aligned} \quad (34)$$

According to the relative distances in Table IX and the relationship of Eqs. (28-34), we can well describe the relative positions of the four valence quarks for all the $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ states. Meanwhile, using the relative distances between (anti)quarks and the orthogonal relation, one can also determine the relative distance of (12) – (34), which is consistent with our results in Table IX. We can also give the relative position of R_c and the spherical radius of the tetraquarks. Here, we define R_c to be the geometric center of the four quarks (the center of the




FIG. 1. Relative positions for four valence quarks and R_c in the $J^{PC} = 0^{++} bb\bar{b}\bar{b}$ ground state. Meanwhile, we label the relative distances of $R_{b,b}$, $R_{b,\bar{b}}$, $R_{\bar{b},\bar{b}}$, R' , and the radius (units: fm).

sphere). Based on these results, we show the spatial distribution of the four valence quarks for the $J^{PC} = 0^{++} bb\bar{b}\bar{b}$ ground state in Fig. 1.

In quark model, a compact tetraquark state has no color-singlet substructure, while a hadronic molecule is a loosely bound state which contains several color-singlet hadrons. According to Table IX, we easily find the relative distances of (1,2), (1,3), (1,4), (2,3), (2,4), and (3,4) quark pairs are all 0.227 or 0.204 fm. Meanwhile, the radius of the state is only 0.130 fm. Thus, in this state, all the distances between the quark pairs are roughly the same order of magnitude apart. If it is a molecular configuration, the distances between two quarks and two antiquarks should be much greater than the distances in the compact multiquark scheme. And the radius of molecular configuration can reach several femtometers. So, our calculations are consistent with the compact tetraquark expectations.

2. The internal contribution

Let us now turn our discussion to the internal mass contribution for the $J^{PC} = 0^{++} bb\bar{b}\bar{b}$ ground state.

First, for the kinetic energy, this $bb\bar{b}\bar{b}$ state has 814.0 MeV, which can be understood as the sum of three internal kinetic energies: kinetic energies of two pairs of the $b - \bar{b}$, and the $(bb) - (\bar{b}\bar{b})$ pair. Accordingly, the sum of the internal kinetic energies of the $\eta_b\eta_b$ state only comes from the two pairs of the $b - \bar{b}$. Therefore, this $bb\bar{b}\bar{b}$ state has an additional kinetic energy needed to bring the $\eta_b\eta_b$ into a compact configuration. The actual kinetic energies of two pairs of the $b - \bar{b}$ in the the $J^{PC} = 0^{++} bb\bar{b}\bar{b}$ ground state are smaller than those in the $\eta_b\eta_b$ state. This is because, as can be seen in Table IX, the distance of $b - \bar{b}$ is larger in the tetraquark state than in the meson: the

distance of $b - \bar{b}$ is 0.204 fm in this $b\bar{b}\bar{b}\bar{b}$ state while it is 0.148 fm in η_b . Meanwhile, we find that even if we consider the additional kinetic energy between the $(b\bar{b}) - (b\bar{b})$ pair, the total kinetic energies in this $b\bar{b}\bar{b}\bar{b}$ state are still smaller than that in the $\eta_b\eta_b$ state. However, this does not lead the ground $J^{PC} = 0^{++} b\bar{b}\bar{b}\bar{b}$ state to a stable state due to the confinement potential part.

As for the confinement potential part, the contributions from V^C for the $J^{PC} = 0^{++} b\bar{b}\bar{b}\bar{b}$ ground state in Table IX are all attractive. Thus, this state has a large positive binding energy. However, it is still above the meson-meson threshold because the $V^C(b\bar{b})$ in η_b is very attractive. As for the other internal contributions, the quark contents of this state are the same as the corresponding rearrangement decay threshold. Moreover, the mass contribution from the hyperfine potential term is negligible compared to the contributions from other terms.

3. Comparison with two models of chromomagnetic interaction

Now, we compare the numerical values for the $cc\bar{c}\bar{c}$ and $b\bar{b}\bar{b}\bar{b}$ systems between the constituent quark model and two CMI models [110, 123] in Tables VII-IX. The comparisons of the values for $cc\bar{c}\bar{c}$ and $b\bar{b}\bar{b}\bar{b}$ states, which are in the last three columns of Tables VII-IX, can be summarized in the following important conclusions.

First, we find that there is no stable state below the lowest heavy quarkonium pair thresholds in any of the three different models. In all three models, we consider two possible color configurations, the color-sextet $|(QQ)^{3_c}(\bar{Q}\bar{Q})^{3_c}\rangle$ and the color-triplet $|(QQ)^{6_c}(\bar{Q}\bar{Q})^{6_c}\rangle$. According to the extended chromomagnetic model [110], the ground state is always dominated by the color-sextet configuration. This view is consistent with the specific wave function of the ground state in Eq.(21) given by the constituent quark model.

In contrast, the obtained masses from the constituent quark model are systematically larger than those from the extended CMI model [110] according to Tables VII-IX. Meanwhile, the obtained masses from the the CMI model [123] are obviously larger than those of the constituent quark model. Their mass differences are mainly due to the effective quark masses as given in the last three columns of Tables VII-IX. The effective quark masses are the sum of the quark mass, the relevant kinetic term, and all the relevant interaction terms in the constituent quark model, which indeed seems to approximately reproduce the effective quark mass from two CMI model [110, 123]. We compare the subtotal values of c and \bar{c} quark part in Table VIII. The c effective quark mass in constituent quark model is 3225 MeV, which is about 100 MeV larger than that of the extended CMI model in the $J^{PC} = 0^{++} cc\bar{c}\bar{c}$ state. Correspondingly, we also find that the c effective quark mass in the CMI model [123] is 3450 MeV and about 200 MeV larger than that

of constituent quark model. The effective quark masses in extended CMI model depend on the parameters of the traditional hadron. However, the effective quark masses should be different depending on whether they are inside a meson, a baryon, or a tetraquark. The effective quark masses trend to be large when they are inside configurations with larger constituents in the extended CMI model, as can be seen from the comparisons of the last three columns in Tables VII-IX. Moreover, we notice the similar situation also occurs in the $ud\bar{c}\bar{c}$ state in Table X of Ref. [61].

B. $cc\bar{b}\bar{b}$ state

Here, we will concentrate on the $cc\bar{b}\bar{b}$ system. Similar to the $cc\bar{c}\bar{c}$ and $b\bar{b}\bar{b}\bar{b}$ systems, the $cc\bar{b}\bar{b}$ system is also satisfied with fully antisymmetric for diquarks and anti-quarks. There are two $J^P = 0^+$ states, one $J^P = 1^+$ state, and one $J^P = 2^+$ state in the $cc\bar{b}\bar{b}$ system. We show the masses of the ground states, the variational parameters, the internal mass contributions, the relative lengths between the quarks, their lowest meson-meson thresholds, the specific wave function, the magnetic moments, the transition magnetic moments, the radiative decay widths, and the rearrangement strong width ratios in Tables XI and XII.

First, we take the $J^P = 0^+$ $cc\bar{b}\bar{b}$ ground state as an example to discuss its properties with the variational method. A similar situation occurs in the other two quantum numbers according to Tables XI, and XII. The mass of the lowest $J^P = 0^+$ $cc\bar{b}\bar{b}$ state is 12920.0 MeV, and the corresponding binding energy B_T is +344.2 MeV according to Table XI. Thus, the state is obviously higher than the corresponding rearrangement meson-meson thresholds. The wave function is given by

$$|\Psi_{tot}\rangle = -0.966|F\rangle|R^s\rangle|[\phi_1\chi_5]\rangle + 0.259|F\rangle|R^s\rangle|[\phi_2\chi_6]\rangle. \quad (35)$$

Here, we see that the mass contribution of ground state mainly comes from the $|(Q_1Q_2)_1^3(\bar{Q}_3\bar{Q}_4)_1^3\rangle_0$ component, and the $|(Q_1Q_2)_0^6(\bar{Q}_3\bar{Q}_4)_0^6\rangle_0$ component is negligible. Its variational parameters are given as $C_{11} = 23.9\text{fm}^{-2}$, $C_{22} = 10.5\text{fm}^{-2}$, and $C_{33} = 12.3\text{fm}^{-2}$.

The meson-meson configuration is connected to the diquark-antidiquark configuration by a linear transformation. Then, we obtain the total wave function in the meson-meson configuration:

$$|\Psi_{tot}\rangle = -0.589|F\rangle|R^s\rangle|[\psi_1\zeta_5]\rangle + 0.095|F\rangle|R^s\rangle|[\psi_1\zeta_6]\rangle + 0.608|F\rangle|R^s\rangle|[\psi_2\zeta_5]\rangle + 0.524|F\rangle|R^s\rangle|[\psi_2\zeta_6]\rangle. \quad (36)$$

According to Eq. (36), we are sure that the overlaps c_i of B_cB_c and $B_c^*B_c^*$ are 0.095 and 0.589, respectively. Then, based on Eq. (20), the rearrangement strong width

TABLE VII. The masses, binding energies, variational parameters, the internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between the quarks for the $J^{PC} = 0^{++}, 1^{+-}$ $cc\bar{c}\bar{c}$ states and their lowest meson-meson thresholds. Here, (i, j) denotes the contribution of the i -th and j -th quarks. The number is given as $i=1$ and 2 for the quarks, 3 and 4 for the antiquark. The masses and corresponding contributions are given in units of MeV, and the relative lengths (variational parameters) are in units of fm (fm^{-2}). Meanwhile, we give a comparison with the other two CMI models [110, 123], to further secure the effective quark mass.

$cc\bar{c}\bar{c}$				The contribution from each term		Relative Lengths (fm)		Overall	Present Work		CMI Model				
$J^{PC} = 0^{++}$		Value	$\eta_c\eta_c$	Difference	(i, j)	Value	$\eta_c\eta_c$		Contribution	Value	Ref. [110]	Ref. [123]			
Mass/ B_T		6384.4	5997.0	387.4	(1,2)	0.406		$2m_c$	3836.0						
Variational Parameters (fm^{-2})	C_{11}	7.7	15.0		(1,3)	0.371	0.290(η_c)	$\frac{\mathbf{p}_{x1}^2}{2m_1^2}$	233.9						
	C_{22}	7.7	15.9		(2,3)	0.371		$\frac{m_{\bar{c}}}{m_c+m_{\bar{c}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$	123.1						
	C_{33}	11.4	-		(1,4)	0.371		$V^C(12)$	-6.8	$-\frac{1}{4}m_{cc}$					
Quark Mass		7672.0	7672.0	0.0	(2,4)	0.371	0.290(η_c)	$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	-52.1	-792.9	$2m_c$				
Confinement Potential		-2083.8	-2440.4	356.6	(3,4)	0.406		$-D$	-983.0	$\frac{5}{4}m_{c\bar{c}}$	3449.6				
Kinetic Energy		814.0	915.1	-101.1	(1,2)-(3,4):	0.235 fm									
CS Interaction		22.7	-150.0	172.7	Radius:		0.235 fm								
V^C	(1,2)	-6.8			(1,3)	-26.1	$-237.2(\eta_c)$								
	(2,3)	-26.1			(3,4)	-6.8									
	(1,4)	-26.1			(2,4)	-26.1	$-237.2(\eta_c)$								
Subtotal		-117.8	-474.4	356.6				Subtotal	3151.1	3042.7	3449.6				
Total Contribution		718.9	291.0	427.9											
Total Wave function:															
$\Psi_{tot} = 0.535 F\rangle R^s\rangle[\phi_1\chi_5\rangle] - 0.845 F\rangle R^s\rangle[\phi_2\chi_6\rangle] = 0.612 F\rangle R^s\rangle[\psi_1\zeta_5\rangle] + 0.443 F\rangle R^s\rangle[\psi_1\zeta_6\rangle] = 0.612 F\rangle R^s\rangle[\psi_1\zeta_5\rangle] + 0.443 F\rangle R^s\rangle[\psi_1\zeta_6\rangle]$															
The rearrangement strong width ratios:															
$\Gamma_{T_{c_2\bar{c}2}(6384.3,0^{++}) \rightarrow J/\psi J/\psi} : \Gamma_{T_{c_2\bar{c}2}(6384.3,0^{++}) \rightarrow \eta_c \eta_c} = 1 : 2.8$															
The radiative decay widths: $\Gamma_{T_{c_2\bar{c}2}(6482.7,2^{++}) \rightarrow T_{c_2\bar{c}2}(6384.3,0^{++})\gamma} = 0$ keV															
$\Gamma_{T_{c_2\bar{c}2}(6451.5,1^{+-}) \rightarrow T_{c_2\bar{c}2}(6384.3,0^{++})\gamma} = 238.1$ keV															
The magnetic moments: $\mu_{T_{c_2\bar{c}2}(6384.3,0^{++})} = \langle \Psi_{tot}^{0^{++}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0$															
The transition magnetic moments:															
$\mu_{T_{c_2\bar{c}2}(6451.5,1^{+-}) \rightarrow T_{c_2\bar{c}2}(6384.3,0^{++})\gamma} = \langle \Psi_{tot}^{1^{+-}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0.671 \mu_N$															
$\mu_{T_{c_2\bar{c}2}(6482.7,2^{++}) \rightarrow T_{c_2\bar{c}2}(6384.3,0^{++})\gamma} = \langle \Psi_{tot}^{2^{++}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0$															
$J^{PC} = 1^{+-}$				Value		$J/\psi\eta_c$		Relative Lengths (fm)		Overall	Contribution		Value	Ref. [110]	Ref. [123]
Mass/ B_T		6451.5	6090.7	360.8	(i,j)	Value	$J/\psi\eta_c$								
Variational Parameters (fm^{-2})	C_{11}	9.1	15.0		(1,2)	0.373		$2m_c$	3836.0						
	C_{22}	9.1	12.5		(1,3)	0.395	0.290(η_c)	$\frac{\mathbf{p}_{x1}^2}{2m_1^2}$	277.2						
	C_{33}	7.3	-		(2,3)	0.395		$\frac{m_{\bar{c}}}{m_c+m_{\bar{c}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$	111.2	$\frac{1}{2}m_{cc}$					
Quark Mass		7672.0	7672.0	0.0	(1,4)	0.395		$V^C(12)$	-19.4	1585.8					
Confinement Potential		-1998.8	-2367.4	368.6	(2,4)	0.395	0.318(J/ψ)	$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	3.0	$\frac{1}{2}m_{c\bar{c}}$	$2m_c$				
Kinetic Energy		767.2	839.0	-71.8	(3,4)	0.373		$-D$	-983.0	1534.3	3449.6				
CS Interaction		1.5	-53.9	54.4				Subtotal	3225.0	3120.0	3449.6				
V^C	(1,2)	-19.4			(1,2)-(3,4):	0.294 fm									
	(2,3)	1.5			Radius:		0.235 fm								
	(1,4)	1.5			(1,3)	1.5	$-237.2(\eta_c)$								
Subtotal		-32.8	-401.4	368.6	(3,4)	-19.4									
Total Contribution		718.9	291.0	427.9	(2,4)	-1.5	$-164.2(J/\psi)$								
Total Wave function: $\Psi_{tot} = F\rangle R^s\rangle[\phi_1\chi_2\rangle] = -0.408 F\rangle R^s\rangle[\psi_1\zeta_2\rangle] - 0.408 F\rangle R^s\rangle[\psi_1\zeta_3\rangle] + 0.577 F\rangle R^s\rangle[\psi_2\zeta_5\rangle] + 0.577 F\rangle R^s\rangle[\psi_2\zeta_6\rangle]$															
The rearrangement strong decay channel: $J/\psi\eta_c$															
The radiative decay widths:															
$\Gamma_{T_{c_2\bar{c}2}(6482.7,2^{++}) \rightarrow T_{c_2\bar{c}2}(6451.5,1^{+-})\gamma} = 70.4$ keV															
$\Gamma_{T_{c_2\bar{c}2}(6451.5,1^{+-}) \rightarrow T_{c_2\bar{c}2}(6384.3,0^{++})\gamma} = 238.1$ keV															
The magnetic moments: $\mu_{T_{c_2\bar{c}2}(6451.5,1^{+-})} = \langle \Psi_{tot}^{1^{+-}} \hat{\mu}^z \Psi_{tot}^{1^{+-}} \rangle = 0$															
The transition magnetic moments:															
$\mu_{T_{c_2\bar{c}2}(6482.7,2^{++}) \rightarrow T_{c_2\bar{c}2}(6451.5,1^{+-})\gamma} = \langle \Psi_{tot}^{2^{++}} \hat{\mu}^z \Psi_{tot}^{1^{+-}} \rangle = 0.750 \mu_N$															
$\mu_{T_{c_2\bar{c}2}(6451.5,1^{+-}) \rightarrow T_{c_2\bar{c}2}(6384.3,0^{++})\gamma} = \langle \Psi_{tot}^{1^{+-}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0.335 \mu_N$															
c-quark:	m_c^{eff}	$2m_c$	3836.0												
		$\frac{\mathbf{p}_{x1}^2}{2m_1^2}$	277.2												
		$\frac{m_{\bar{c}}}{m_c+m_{\bar{c}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$	111.2	$\frac{1}{2}m_{cc}$											
\bar{c} -quark:	$m_{\bar{c}}^{eff}$	$V^C(12)$	-19.4	1585.8											
		$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	3.0	$\frac{1}{2}m_{c\bar{c}}$	$2m_c$										
		$-D$	-983.0	1534.3	3449.6										
Subtotal															
Subtotal		3225.0	3120.0	3449.6											
Total contribution		6451.5	6231.0	6899.0											
CS Interaction															
$\frac{1}{2}V^{SS}(12)$															
$\frac{1}{2}V^{SS}(34)$															
$-\frac{1}{4}(V^{SS}(13) + V^{SS}(14) + V^{SS}(23) + V^{SS}(24))$															
Subtotal		1.5	-9.5	0.0											
Total contribution		6451.5	6231.0	6899.0											

TABLE VIII. The masses, binding energies, variational parameters, the internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between quarks for the $J^{PC} = 2^{++} cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ states and their lowest meson-meson thresholds. The notation is the same as that in Table VII.

$cc\bar{c}\bar{c}$				The contribution from each term		Relative Lengths (fm)		Overall	Present Work		CMI Model		
$J^{PC} = 2^{++}$		Value	$J/\psi J/\psi$	Difference	(i, j)	Value	$J/\psi J/\psi$		Contribution	Value	Ref. [110]	Ref. [123]	
Mass/ B_T		6482.7	6184.5	298.2	(1,2)	0.377		$2m_c$	3836.0				
Variational Parameters (fm $^{-2}$)	C_{11}	8.9	12.5		(1,3)	0.403	0.318(J/ψ)	$\frac{p_{x1}^2}{2m_1^2}$	270.4				
	C_{22}	8.9	12.5		(2,3)	0.403		$\frac{m_{\bar{c}}}{m_c+m_{\bar{c}}}\frac{p_{x3}^2}{2m_3^2}$	105.6	$\frac{1}{2}m_{cc}$			
	C_{33}	6.9	-		(1,4)	0.403		$V^C(12)$	-14.6	1585.8			
Quark Mass		7672.0	7672.0	0.0	(1,4)	0.403		$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	10.8	$\frac{1}{2}m_{c\bar{c}}$	$2m_c$		
Confinement Potential		-1973.6	-2294.4	320.8	(2,4)	0.403	0.318(J/ψ)	$-D$	-983.0	1534.3	3449.6		
Kinetic Energy		752.0	769.9	-10.9	(3,4)	0.377		Subtotal	3225.2	3120.0	3449.6		
CS Interaction		32.3	43.9	-11.6	(1,2)-(3,4):	0.302 fm							
					Radius:	0.241 fm							
V^C	(1,2)	-14.6			(1,3)	5.4	-164.2(J/ψ)	$2m_c$	3836.0				
	(2,3)	5.4			(3,4)	-14.6		$\frac{p_{x2}^2}{2m_2^2}$	270.4				
	(1,4)	5.5			(1,3)	5.4	-164.2(J/ψ)	$\frac{m_{\bar{c}}}{m_c+m_{\bar{c}}}\frac{p_{x3}^2}{2m_3^2}$	105.6	$\frac{1}{2}m_{cc}$			
Subtotal		-7.6	-328.4	320.8	(3,4)	-14.6		$V^C(12)$	-14.6	1585.8			
Total Contribution		776.7	478.5	298.2	(2,4)	5.4	-164.2(J/ψ)	$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	10.8	$\frac{1}{2}m_{c\bar{c}}$	$2m_c$		
					(2,4)	5.4	-164.2(J/ψ)	$-D$	-983.0	1534.3	3449.6		
Total Wave function:													
$\Psi_{tot} = F\rangle R^s\rangle \phi_1\chi_1\rangle = 0.577 F\rangle R^s\rangle \psi_1\zeta_1\rangle - 0.816 F\rangle R^s\rangle \psi_2\zeta_1\rangle$													
The rearrangement strong decay channel: $J/\psi J/\psi$													
The radiative decay widths:													
$\Gamma_{T_{c2\bar{c}2}(6482.7,2^{++}) \rightarrow T_{c2\bar{c}2}(6384.3,0^{++})\gamma} = 0$ keV													
$\Gamma_{T_{c2\bar{c}2}(6482.7,2^{++}) \rightarrow T_{c2\bar{c}2}(6451.5,1^{+-})\gamma} = 70.4$ keV													
The magnetic moments: $\mu_{T_{c2\bar{c}2}(6451.5,1^{+-})} = \langle \Psi_{tot}^{1+-} \hat{\mu}^z \Psi_{tot}^{1+-} \rangle = 0$.													
The transition magnetic moments:													
$\mu_{T_{c2\bar{c}2}(6482.7,2^{++}) \rightarrow T_{c2\bar{c}2}(6384.3,0^{++})\gamma} = \langle \Psi_{tot}^{2++} \hat{\mu}^z \Psi_{tot}^{0++} \rangle = 0$													
$\mu_{T_{c2\bar{c}2}(6482.7,2^{++}) \rightarrow T_{c2\bar{c}2}(6451.4,1^{+-})\gamma} = \langle \Psi_{tot}^{2++} \hat{\mu}^z \Psi_{tot}^{1+-} \rangle = \mu_c - \mu_{\bar{c}} = 0.750\mu_N$													
Subtotal													
Total contribution									6482.7	6287.3	6956.0		
$bb\bar{b}\bar{b}$ $J^{PC} = 2^{++}$				Value		$\Upsilon\Upsilon$	Difference	Relative Lengths (fm)		Contribution	Value	Ref. [110]	Ref. [123]
Mass/ B_T		19327.9	18938.8	390.1	(i, j)	Value	$\Upsilon\Upsilon$						
Variational Parameters (fm $^{-2}$)	C_{11}	30.0	49.4		(1,2)	0.205		$2m_b$	10686.0				
	C_{22}	30.0	49.4		(1,3)	0.220	0.160(Υ)	$\frac{p_{x1}^2}{2m_1^2}$	328.0				
	C_{33}	23.0	-		(1,3)	0.220	0.160(Υ)	$\frac{m_{\bar{b}}}{m_b+m_{\bar{b}}}\frac{p_{x3}^2}{2m_3^2}$	126.0	$\frac{1}{2}m_{bb}$			
Quark Mass		21372.0	21372.0	0.0	(2,3)	0.220		$V^C(12)$	-269.1	4764.8			
Confinement Potential		-2977.3	-3559.5	582.2	(1,4)	0.220		$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	-236.6	$\frac{1}{2}m_{b\bar{b}}$	$2m_b$		
Kinetic Energy		908.1	1087.3	-179.2	(2,4)	0.220	0.160(Υ)	$-D$	-983.0	4722.5	10105.8		
CS Interaction		25.1	38.0	-12.9	(3,4)	0.205		Subtotal	9651.3	9487.3	10105.8		
V^C	(1,2)	-269.1			(1,2)-(3,4):	0.165 fm		$2m_b$	10686.0				
	(2,3)	-118.3			Radius:	0.132 fm		$\frac{p_{x2}^2}{2m_2^2}$	328.0				
	(1,4)	-118.3			(1,3)	-118.3	-796.7(Υ)	$\frac{m_{\bar{b}}}{m_b+m_{\bar{b}}}\frac{p_{x3}^2}{2m_3^2}$	126.0	$\frac{1}{2}m_{bb}$			
Subtotal		-1011.3	-1593.5	582.2	(3,4)	-269.1		$V^C(12)$	-269.1	4764.8			
Total Contribution		-78.1	-468.2	390.1	(2,4)	-118.3	-796.7(Υ)	$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	-236.6	$\frac{1}{2}m_{b\bar{b}}$	$2m_b$		
					(2,4)	-118.3	-796.7(Υ)	$-D$	-983.0	4722.5	10105.8		
Total Wave function:													
$\Psi_{tot} = F\rangle R^s\rangle \phi_1\chi_2\rangle = 0.577 F\rangle R^s\rangle \psi_1\zeta_1\rangle - 0.816 F\rangle R^s\rangle \psi_2\zeta_1\rangle$													
The rearrangement strong decay channel: $B_c^* B_c^*$													
The radiative decay widths:													
$\Gamma_{T_{b2\bar{b}2}(19327.9,2^{++}) \rightarrow T_{b2\bar{b}2}(19240.0,0^{++})\gamma} = 0$ keV													
$\Gamma_{T_{b2\bar{b}2}(19327.9,2^{++}) \rightarrow T_{b2\bar{b}2}(19303.9,1^{+-})\gamma} = 1.0$ keV													
The magnetic moments: $\mu_{T_{b2\bar{b}2}(19303.9,1^{+-})} = \langle \Psi_{tot}^{1+-} \hat{\mu}^z \Psi_{tot}^{1+-} \rangle = 0$													
The transition magnetic moments:													
$\mu_{T_{b2\bar{b}2}(19327.9,2^{++}) \rightarrow T_{b2\bar{b}2}(19240.0,0^{++})\gamma} = \langle \Psi_{tot}^{2++} \hat{\mu}^z \Psi_{tot}^{0++} \rangle = 0$													
$\mu_{T_{b2\bar{b}2}(19327.9,2^{++}) \rightarrow T_{b2\bar{b}2}(19303.9,1^{+-})\gamma} = \langle \Psi_{tot}^{2++} \hat{\mu}^z \Psi_{tot}^{1+-} \rangle = \mu_b - \mu_{\bar{b}} = -0.125\mu_N$													
Total contribution									19327.9	19000.1	20243.0		

TABLE IX. The masses, binding energies, variational parameters, the internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between quarks for the $J^{PC} = 0^{++}$, 1^{+-} $b\bar{b}b\bar{b}$ states and their lowest meson-meson thresholds. The notation is the same as that in Table VII.

$b\bar{b}b\bar{b}$		The contribution from each term			Relative Lengths (fm)		Overall	Present Work		CMI Model		
$J^{PC} = 0^{++}$	Value	$\eta_b\eta_{\bar{b}}$	Difference	(i, j)	Vaule	$\eta_b\eta_{\bar{b}}$		Contribution	Value	Ref. [110]	Ref. [123]	
Mass/ B_T	19240.0	18778.1	461.9	(1,2)	0.227		b -quark: m_b^{eff}	$2m_b$	10686.0			
Variational Parameters (fm $^{-2}$)	C_{11}	24.6	57.4	(1,3)	0.204	0.148(η_b)		$\frac{\mathbf{p}_{x1}^2}{2m_1^2}$	269.2			
	C_{22}	24.6	57.4	(2,3)	0.204			$\frac{m_{\bar{b}}}{m_b+m_{\bar{b}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$	216.0			
	C_{33}	39.5	-	(1,4)	0.204			$V^C(12)$	111.8	$-\frac{1}{4}m_{bb}$		
Quark Mass	21372.0	21372.0	0.0	(2,4)	0.204	0.148(η_b)		$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	-697.4	-2382.4	$2m_b$	
Confinement Potential	-3101.0	-3724.2	623.2	(3,4)	0.227			$-D$	-983.0	$\frac{5}{4}m_{b\bar{b}}$	11806.2	10105.8
Kinetic Energy	970.4	1255.9	-285.5	(1,2)-(3,4):		0.126 fm		Subtotal	9602.6	9423.8	10105.8	
CS Interaction	17.0	-125.5	142.5	Radius:		0.130 fm						
V^C	(1,2)	-111.8		(1,3)	-339.7	-879.1(η_b)		\bar{b} -quark: $m_{\bar{b}}^{eff}$	$2m_b$	10686.0		
	(2,3)	-339.7		(3,4)	-111.8				$\frac{\mathbf{p}_{x2}^2}{2m_2^2}$	216.0		
	(1,4)	-339.7		(3,4)	-111.8		$\frac{m_{\bar{b}}}{m_b+m_{\bar{b}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$		216.0			
Subtotal	-1135.0	-1758.2	623.2	(2,4)	-339.7	-879.1(η_b)	$V^C(34)$		111.8	$-\frac{1}{4}m_{bb}$		
Total Contribution	-147.6	-627.9	480.3				$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$		-697.4	-2382.4	$2m_b$	
Total Wave function:							CS Interaction		$-D$	-983.0	11806.2	10105.8
$\Psi_{tot} = 0.352 F\rangle R^s\rangle[\phi_1\chi_5\rangle] - 0.936 F\rangle R^s\rangle[\phi_2\chi_6\rangle] = 0.558 F\rangle R^s\rangle[\psi_1\zeta_5\rangle] + 0.560 F\rangle R^s\rangle[\psi_1\zeta_6\rangle] + 0.021 F\rangle R^s\rangle[\psi_2\zeta_5\rangle] + 0.612 F\rangle R^s\rangle[\psi_2\zeta_6\rangle]$									Subtotal	9602.6	9423.8	10105.8
The rearrangement strong width ratios:									$\frac{3}{4}V^{SS}(12)$	8.5	$4v_{bb}$	$4C_{bb}$
$\Gamma_{T_{c2,c2}(19240.0,0^{++}) \rightarrow \Upsilon\Upsilon} : \Gamma_{T_{c2,c2}(19240.0,0^{++}) \rightarrow \eta_b\eta_{\bar{b}}} = 1 : 1.2$									$\frac{3}{4}V^{SS}(34)$	8.5	$4v_{\bar{b}\bar{b}}$	$4C_{\bar{b}\bar{b}}$
The radiative decay widths:									Subtotal	17.0	15.4	23.2
$\Gamma_{T_{b2,\bar{b}2}(19327.9,2^{++}) \rightarrow T_{b2,\bar{b}2}(19240.0,0^{++})\gamma} = 0 \text{ keV}$								Matrix nondiagonal element	17.8	-27.0	-40.2	
$\Gamma_{T_{b2,\bar{b}2}(19303.9,1^{+-}) \rightarrow T_{b2,\bar{b}2}(19240.0,0^{++})\gamma} = 2.8 \text{ keV}$								Total contribution	19240.0	18836.0	20275.0	
The magnetic moments: $\mu_{T_{b2,\bar{b}2}(19240.0,0^{++})} = \langle \Psi_{tot}^{0^{++}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0$												
The transition magnetic moments: $\mu_{T_{b2,\bar{b}2}(19327.9,2^{++}) \rightarrow T_{b2,\bar{b}2}(19240.0,0^{++})\gamma} = \langle \Psi_{tot}^{2^{++}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0$												
$\mu_{T_{b2,\bar{b}2}(19393.9,1^{+-}) \rightarrow T_{b2,\bar{b}2}(19240.0,0^{++})\gamma} = \langle \Psi_{tot}^{1^{+-}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0.352 \times \frac{4}{\sqrt{6}}(\mu_b - \mu_{\bar{b}}) = -0.072\mu_N$												
$J^{PC} = 1^{+-}$	Value	$\Upsilon\eta_b$	Difference	Relative Lengths (fm)			Contribution	Value	Ref. [110]	Ref. [123]		
Mass/ B_T	19303.9	18857.9	446.0	(i, j)	Value	$\Upsilon\eta_b$	b -quark: m_b^{eff}	$2m_b$	10686.0			
Variational Parameters (fm $^{-2}$)	C_{11}	30.7	57.4	(1,2)	0.203			$\frac{\mathbf{p}_{x1}^2}{2m_1^2}$	335.6			
	C_{22}	30.7	49.4	(1,3)	0.217	0.148(η_b)		$\frac{m_{\bar{b}}}{m_b+m_{\bar{b}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$	131.4	$\frac{1}{2}m_{bb}$		
	C_{33}	24.0	-	(2,3)	0.217			$V^C(12)$	-274.5	4764.8		
Quark Mass	21372.0	21372.0	0.0	(1,4)	0.217			$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	-244.2	$\frac{1}{2}m_{b\bar{b}}$	$2m_b$	
Confinement Potential	-3003.4	-3641.9	638.4	(2,4)	0.217	Υ		$-D$	-983.0	4722.5	10105.8	
Kinetic Energy	934.0	1171.6	-237.6	(3,4)	0.203			Subtotal	9653.3	9487.3	10105.8	
CS Interaction	1.3	-43.8	45.1	(1,2)-(3,4):		0.162 fm						
V^C	(1,2)	-274.5		(1,3)	-122.1	-879.1(η_b)		\bar{b} -quark: $m_{\bar{b}}^{eff}$	$2m_b$	10686.0		
	(2,3)	-122.1		(3,4)	-274.5				$\frac{\mathbf{p}_{x2}^2}{2m_2^2}$	335.6		
	(1,4)	-122.1		(3,4)	-274.5		$\frac{m_{\bar{b}}}{m_b+m_{\bar{b}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$		131.4	$\frac{1}{2}m_{bb}$		
Subtotal	-1037.4	-1675.9	638.5	(2,4)	-122.1	-796.7(Υ)	$V^C(12)$		-274.5	4764.8		
Total Contribution	-102.1	-548.1	446.0				$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$		-244.2	$\frac{1}{2}m_{b\bar{b}}$	$2m_b$	
Total Wave function: $\Psi_{tot} = F\rangle R^s\rangle[\phi_1\chi_2\rangle] = -0.408 F\rangle R^s\rangle[\psi_1\zeta_2\rangle] - 0.408 F\rangle R^s\rangle[\psi_1\zeta_3\rangle] + 0.577 F\rangle R^s\rangle[\psi_2\zeta_5\rangle] + 0.577 F\rangle R^s\rangle[\psi_2\zeta_6\rangle]$							$-D$		-983.0	4722.5	10105.8	
The rearrangement strong decay channel: $J/\psi\eta_c$							CS Interaction		Subtotal	9653.3	9487.3	10105.8
The radiative decay widths:									$\frac{1}{2}V^{SS}(12)$	6.7	$\frac{8}{3}v_{bb}$	$\frac{8}{3}C_{bb}$
$\Gamma_{T_{b2,\bar{b}2}(19327.9,2^{++}) \rightarrow T_{b2,\bar{b}2}(19303.9,1^{+-})\gamma} = 1.0 \text{ keV}$									$\frac{1}{2}V^{SS}(34)$	6.7	$\frac{8}{3}v_{\bar{b}\bar{b}}$	$\frac{8}{3}C_{\bar{b}\bar{b}}$
$\Gamma_{T_{b2,\bar{b}2}(19303.9,1^{+-}) \rightarrow T_{b2,\bar{b}2}(19240.0,0^{++})\gamma} = 2.8 \text{ keV}$									$-\frac{1}{4}(V^{SS}(13) + V^{SS}(14) + V^{SS}(23) + V^{SS}(24))$	-12.2	$-\frac{16}{3}v_{b\bar{b}}$	$-\frac{16}{3}C_{b\bar{b}}$
The magnetic moments: $\mu_{T_{b2,\bar{b}2}(19303.9,1^{+-})} = \langle \Psi_{tot}^{1^{+-}} \hat{\mu}^z \Psi_{tot}^{1^{+-}} \rangle = 0$								Subtotal	1.3	-5.1	0.0	
The transition magnetic moments: $\mu_{T_{b2,\bar{b}2}(19327.9,2^{++}) \rightarrow T_{b2,\bar{b}2}(19303.9,1^{+-})\gamma} = \langle \Psi_{tot}^{2^{++}} \hat{\mu}^z \Psi_{tot}^{1^{+-}} \rangle = -0.125\mu_N$												
$\mu_{T_{b2,\bar{b}2}(19393.9,1^{+-}) \rightarrow T_{b2,\bar{b}2}(19240.0,0^{++})\gamma} = \langle \Psi_{tot}^{1^{+-}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = -0.072\mu_N$												
								Total contribution	19303.9	18696.4	20211.6	

ratios is

$$\frac{\Gamma_{T_{c^2\bar{b}^2}(12920.0,0^+) \rightarrow B_c^* B_c^*}}{\Gamma_{T_{c^2\bar{b}^2}(12920.0,0^+) \rightarrow B_c B_c}} = 1 : 50, \quad (37)$$

i.e., $B_c B_c$ is the dominant rearrangement decay channel for the $T_{c^2\bar{b}^2}(12920.0,0^+)$ state.

As for the magnetic moment of the $J^P = 0^+ c\bar{c}b\bar{b}$ ground state, its value is 0, while the magnetic moment of all $J^P = 0^+$ tetraquark states is 0. As for the $J^P = 1^+ c\bar{c}b\bar{b}$ state, we construct its flavor \otimes spin wave functions as

$$\begin{aligned} & |\Psi\rangle_{T_{c^2\bar{b}^2}(12939.9,1^+)}^{S=1;S_s=1} \\ &= |R^s\rangle|\psi\rangle|c\bar{c}b\bar{b}\rangle \frac{1}{2}(\uparrow\uparrow\downarrow\downarrow + \uparrow\downarrow\uparrow\downarrow - \downarrow\uparrow\uparrow\downarrow - \downarrow\downarrow\uparrow\uparrow) \end{aligned} \quad (38)$$

So the corresponding transition magnetic momentum is

$$\mu_{T_{c^2\bar{b}^2}(12939.9,1^+)} = \langle \Psi_{tot}^{1^+} | \hat{\mu}^z | \Psi_{tot}^{1^+} \rangle = \mu_c + \mu_{\bar{b}} = 0.490\mu_N. \quad (39)$$

We also discuss the transition magnetic moment of the $T_{c^2\bar{b}^2}(12939.9,1^+) \rightarrow T_{c^2\bar{b}^2}(12920.0,0^+)\gamma$ process. We still construct their flavor \otimes spin wave functions as

$$\begin{aligned} |\Psi\rangle_{T_{c^2\bar{b}^2}(1^+)}^{S=1;S_s=0} &= |R^s\rangle|\psi\rangle|c\bar{c}b\bar{b}\rangle \frac{1}{\sqrt{2}}(\uparrow\uparrow\downarrow\downarrow - \downarrow\downarrow\uparrow\uparrow), \\ |\Psi\rangle_{T_{c^2\bar{b}^2}(0^+)}^{S=0;S_s=0} &= |R^s\rangle|\psi\rangle|c\bar{c}b\bar{b}\rangle 0.966 \frac{1}{\sqrt{3}}(\uparrow\uparrow\downarrow\downarrow + \downarrow\downarrow\uparrow\uparrow) + \dots \end{aligned} \quad (40)$$

And then, the transition magnetic momentum of the $T_{c^2\bar{b}^2}(12939.9,1^+) \rightarrow T_{c^2\bar{b}^2}(12920.0,0^+)\gamma$ process can be described by the z -component of the magnetic moment operator $\hat{\mu}^z$ sandwiched by the flavor-spin wave functions of the $T_{c^2\bar{b}^2}(12939.9,1^+)$ and $T_{c^2\bar{b}^2}(12920.0,0^+)$. So the corresponding transition magnetic momentum is

$$\begin{aligned} \mu_{T_{c^2\bar{b}^2}(12939.9,1^+) \rightarrow T_{c^2\bar{b}^2}(12920.0,0^+)\gamma} &= \langle \Psi_{tot}^{1^+} | \hat{\mu}^z | \Psi_{tot}^{0^+} \rangle \\ &= 0.966 \times \frac{1}{\sqrt{6}}(4\mu_c - 4\mu_{\bar{b}}) = 0.534\mu_N. \end{aligned} \quad (41)$$

Further, according to Eq. (18) and Eq. (41), we also obtain the radiative decay widths

$$\Gamma_{T_{c^2\bar{b}^2}(12960.9,2^+) \rightarrow T_{c^2\bar{b}^2}(12939.9,1^+)\gamma} = 3.6 \text{ keV}. \quad (42)$$

Finally, we turn to the internal contribution for the $c\bar{c}b\bar{b}$ ground state. For the kinetic energy part, the $J^P = 0^+ c\bar{c}b\bar{b}$ state receives 835.9 MeV, which is smaller than that of the meson-meson threshold $B_c B_c$. The potential part of this state is much smaller than that of the lowest meson-meson threshold. Furthermore, we find that all the V^C for this state are attractive. However, compared to the V^C of $B_c B_c$, these attractive values seem to trivial. This is because the length between $c - \bar{b}$ in tetraquarks is longer than that in B_c according to Table XI and XII. In summary, we tend to think that these $c\bar{c}b\bar{b}$ states are unstable compact states.

C. $c\bar{c}b\bar{b}$ and $b\bar{b}c\bar{c}$ states

Here, we discuss the $c\bar{c}b\bar{b}$ and $b\bar{b}c\bar{c}$ systems. For these two systems, they only need to satisfy the antisymmetry for the diquark. Thus, compared to the above three systems, the $c\bar{c}b\bar{b}$ and $b\bar{b}c\bar{c}$ systems have more allowed states. There are two $J^P = 0^+$ states, three $J^P = 1^+$ states, one $J^P = 2^+$ state in the $c\bar{c}b\bar{b}$ and $b\bar{b}c\bar{c}$ systems. We calculate the masses of the ground states, the corresponding variational parameters, the various internal contributions, the relative lengths between the quarks, and their lowest meson-meson thresholds, their lowest meson-meson thresholds, specific wave functions, magnetic moments, transition magnetic moments, radiative decay widths, and rearrangement strong width ratios in Tables XIII, XIV, and XV, respectively.

We now analyse the numerical results of the $J^P = 1^+$ ground $b\bar{b}c\bar{c}$ state obtained from the variational method according to Table XIV. Other states would have similar discussions from Tables XIII-XV. The mass of the lowest $J^P = 1^+ b\bar{b}c\bar{c}$ state is 16043.2 MeV, and the corresponding binding energy B_T is +303.7 MeV. Thus, the state is obviously above the lowest rearrangement meson-meson decay channel $B_c^* \eta_b$, and it is an unstable tetraquark state. Its variational parameters are given as $C_{11} = 12.4\text{fm}^{-2}$, $C_{22} = 21.0\text{fm}^{-2}$, and $C_{33} = 28.9\text{fm}^{-2}$. The corresponding wave function is given by

$$\begin{aligned} |\Psi_{tot}\rangle &= 0.984|F\rangle|R^s\rangle|[\phi_2\chi_4]\rangle + 0.171|F\rangle|R^s\rangle|[\phi_1\chi_3]\rangle \\ &\quad - 0.044|F\rangle|R^s\rangle|[\phi_1\chi_2]\rangle. \end{aligned} \quad (43)$$

Here, we notice that the mass contribution of ground state mainly comes from the $|(Q_1 Q_2)_0^6 (\bar{Q}_3 \bar{Q}_4)_1^6\rangle$ component, and the other two components are negligible. Then we transform Eq. (43) to the meson-meson configuration via a linear transformation, and the corresponding wave function is given as:

$$\begin{aligned} |\Psi_{tot}\rangle &= 0.494|F\rangle|R^s\rangle|[\psi_1\zeta_2]\rangle - 0.396|F\rangle|R^s\rangle|[\psi_1\zeta_3]\rangle \\ &\quad - 0.487|F\rangle|R^s\rangle|[\psi_1\zeta_4]\rangle + 0.111|F\rangle|R^s\rangle|[\psi_2\zeta_2]\rangle \\ &\quad - 0.246|F\rangle|R^s\rangle|[\psi_2\zeta_3]\rangle - 0.537|F\rangle|R^s\rangle|[\psi_2\zeta_4]\rangle. \end{aligned} \quad (44)$$

Furthermore, we can sure that its rearrangement strong width ratios is:

$$\begin{aligned} \Gamma_{T_{b^2\bar{c}^2}(16043.2,1^+) \rightarrow B_c^* \Upsilon} : \Gamma_{T_{b^2\bar{c}^2}(16043.2,1^+) \rightarrow B_c \Upsilon} : \\ \Gamma_{T_{b^2\bar{c}^2}(16043.2,1^+) \rightarrow B_c^* \eta_b} = 1 : 1.3 : 1.5. \end{aligned} \quad (45)$$

And its radiative decay widths are:

$$\begin{aligned} \Gamma_{T_{b^2\bar{c}^2}(16149.2,2^+) \rightarrow T_{c^2\bar{c}^2}(16043.2,1^+)\gamma} &= 435.0 \text{ keV}, \\ \Gamma_{T_{c^2\bar{c}^2}(16043.9,0^+) \rightarrow T_{c^2\bar{c}^2}(16043.2,1^+)\gamma} &= 10^{-6} \text{ keV}. \end{aligned}$$

Let us now focus on the internal contributions for this state and the relative lengths between the quarks. For the kinetic energy part, the state gets 876.1 MeV, which is obviously smaller than that of the lowest meson-meson

threshold $B_c\eta_b$. The actual kinetic energy of the $b - \bar{b}$ ($b - \bar{c}$) in the $J^P = 1^+ bb\bar{b}\bar{c}$ state is smaller than that in the η_b (B_c^*) meson. The reason for this can be seen in Table XIV. The size of this pair is larger in the $J^P = 1^+ bb\bar{b}\bar{c}$ state than in the meson: the distance (3,4) is 0.245 fm in this tetraquark while it is 0.148 fm in η_b .

Here, let us turn our discussion to the potential parts. The potential part of this state is much smaller than that of its lowest meson-meson threshold. Although the V^C between quark and antiquark are attractive, the V^C in the diquark and antiquark are repulsive. However, relative to the η_b and B_c mesons, the V^C in the tetraquark are less attractive. Therefore, they still have a relatively large positive binding energy in this state.

D. $cb\bar{c}\bar{b}$ state

Finally, we investigate the $cb\bar{c}\bar{b}$ system. Similar to the $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ systems, the $cb\bar{c}\bar{b}$ system is also a pure neutral system and has a certain C-parity. Thus the corresponding magnetic moment is $0\mu_N$ for all the ground $cb\bar{c}\bar{b}$ states. Moreover, the Pauli principle does not impose any constraints on the wave functions of the $cb\bar{c}\bar{b}$ system. Thus, compared to other discussed tetraquark systems, the $cb\bar{c}\bar{b}$ system has more allowed states. There are four $J^{PC} = 0^{++}$ states, four $J^{PC} = 1^{+-}$ states, two $J^{PC} = 1^{++}$ states, two $J^{PC} = 2^{++}$ states in the $cb\bar{c}\bar{b}$ system.

Here, we now analyze the numerical results about the $cb\bar{c}\bar{b}$ system obtained from the variational method. Here, we take the $J^{PC} = 0^{++}$ $cb\bar{c}\bar{b}$ ground state as an example to discuss specifically, and others would have similar discussions. The mass of the lowest $J^{PC} = 0^{++}$ $cb\bar{c}\bar{b}$ state is 12759.3 MeV, and the corresponding binding energy B_T is +371.8 MeV. Thus, the state obviously has a larger mass than the lowest rearrangement meson-meson decay channel $\eta_b\eta_c$, and it should be an unstable compact tetraquark state. Its variational parameters are given as $C_{11} = 11.9 \text{ fm}^{-2}$, $C_{22} = 11.9 \text{ fm}^{-2}$, and $C_{33} = 22.9 \text{ fm}^{-2}$. Since this state is a pure neutral state, we naturally notice that the value of C_{11} is equal to C_{22} , which means that the distance of ($b - b$) is equal to ($\bar{b} - \bar{b}$). Our results also reflect these properties according to Table XVI. The corresponding wave function is given as:

$$\begin{aligned} \Psi_{tot} = & 0.961|F\rangle|R^s\rangle[|\phi_2\chi_5\rangle] + 0.114|F\rangle|R^s\rangle[|\phi_2\chi_6\rangle] \\ & - 0.069|F\rangle|R^s\rangle[|\phi_1\chi_5\rangle] - 0.241|F\rangle|R^s\rangle[|\phi_1\chi_6\rangle]. \end{aligned} \quad (46)$$

Based on Eq. (46), we find that its mass contribution to the ground state mainly comes from the $6 \otimes \bar{6}$ component, the corresponding $3 \otimes \bar{3}$ component being negligible. Then we transform Eq. (46) to $c\bar{c} - b\bar{b}$ and $c\bar{b} - b\bar{c}$ configurations via a linear transformation, and the corresponding two

wave functions are given as:

$$\begin{aligned} |\Psi_{tot}\rangle = & -0.830|F\rangle|R^s\rangle[|\psi_1\zeta_5\rangle] + 0.211|F\rangle|R^s\rangle[|\psi_1\zeta_6\rangle] \\ & - 0.367|F\rangle|R^s\rangle[|\psi_2\zeta_5\rangle] + 0.363|F\rangle|R^s\rangle[|\psi_2\zeta_6\rangle] \\ = & -0.668|F\rangle|R^s\rangle[|\psi'_1\zeta'_5\rangle] + 0.333|F\rangle|R^s\rangle[|\psi'_1\zeta'_6\rangle] \\ & - 0.398|F\rangle|R^s\rangle[|\psi'_2\zeta'_5\rangle] + 0.533|F\rangle|R^s\rangle[|\psi'_2\zeta'_6\rangle]. \end{aligned} \quad (47)$$

Further, we can certain its rearrangement strong width ratios. For the $c\bar{b} - b\bar{c}$ decay mode

$$\begin{aligned} \Gamma_{T_{cb\bar{c}\bar{b}}(12796.9,1^{+-}) \rightarrow B_c^* \bar{B}_c^*} : \Gamma_{T_{cb\bar{c}\bar{b}}(12796.9,1^{+-}) \rightarrow B_c^* \bar{B}_c} : \\ \Gamma_{T_{cb\bar{c}\bar{b}}(12796.9,1^{+-}) \rightarrow B_c \bar{B}_c^*} = 1 : 3.9 : 3.9, \end{aligned} \quad (48)$$

where both the $B_c^* \bar{B}_c$ and $B_c \bar{B}_c^*$ channels are the dominant decay modes for the $T_{cb\bar{c}\bar{b}}(12796.9,1^{+-})$ tetraquark state.

$$\begin{aligned} \Gamma_{T_{cb\bar{c}\bar{b}}(12796.9,1^{+-}) \rightarrow \eta_b J/\psi} : \Gamma_{T_{cb\bar{c}\bar{b}}(12796.9,1^{+-}) \rightarrow \Upsilon \eta_c} \\ = 1 : 18.4. \end{aligned} \quad (49)$$

The dominant decay channel is the $\Upsilon\eta_c$ final states in the $c\bar{c} - b\bar{b}$ decay mode.

We also calculate the transition magnetic moments for this state:

$$\begin{aligned} \mu_{T_{cb\bar{c}\bar{b}}(12882.4,2^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12759.6,0^{++})\gamma} &= 0, \\ \mu_{T_{cb\bar{c}\bar{b}}(12856.6,1^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12759.6,0^{++})\gamma} &= 0, \\ \mu_{T_{cb\bar{c}\bar{b}}(12796.9,1^{+-}) \rightarrow T_{cb\bar{c}\bar{b}}(12759.6,0^{++})\gamma} &= 0.081, \end{aligned} \quad (50)$$

which are in units of μ_N . Furthermore, we can obtain its radiative decay widths are:

$$\begin{aligned} \Gamma_{T_{cb\bar{c}\bar{b}}(12882.4,2^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12759.6,0^{++})\gamma} &= 0, \\ \Gamma_{T_{cb\bar{c}\bar{b}}(12856.6,1^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12759.6,0^{++})\gamma} &= 0, \\ \Gamma_{T_{cb\bar{c}\bar{b}}(12797.3,1^{+-}) \rightarrow T_{cb\bar{c}\bar{b}}(12759.6,0^{++})\gamma} &= 33.1, \end{aligned} \quad (51)$$

which are in units of keV.

Let us now turn our discussion to the internal contribution for the $J^{PC} = 1^{+-}$ $cb\bar{c}\bar{b}$ ground state. For the kinetic energy part, the state obtains 858.5 MeV, which is smaller than the 1001.2 MeV of the lowest meson-meson threshold $B_c\eta_b$ according to Table XVI. As for the potential part, although the V^C between quark and antiquark are attractive, the V^C in the diquark and antiquark are repulsive. However, relative to the lowest meson-meson threshold $B_c\eta_b$, the total V^C is not attractive than the $B_c\eta_b$, which leads to this state having a relatively larger mass.

We also notice that the $V^C(1,3)$, $V^C(2,3)$, $V^C(1,4)$, and $V^C(2,4)$ are absolutely the same, and meanwhile the distances of (1,3), (1,4), (2,3), and (2,4) are also the same. These actually reflect $\langle \Psi_{tot} | (\mathbf{R}_{1,2} \cdot \mathbf{R}_{3,4}) | \Psi_{tot} \rangle = \langle \Psi_{tot} | (\mathbf{R}_{1,2} \cdot \mathbf{R}') | \Psi_{tot} \rangle = \langle \Psi_{tot} | (\mathbf{R}_{3,4} \cdot \mathbf{R}') | \Psi_{tot} \rangle = 0$. Obviously, it is unreasonable that the distance of $c\bar{c}$ is exactly the same as that of the $c\bar{b}$ and $b\bar{b}$. According to Sec IV of Ref. [65], we only consider single Gaussian

form which the $l_1 = l_2 = l_3 = 0$ in spatial part of the total wave function is not sufficient. These lead to the $cb\bar{c}\bar{b}$ state, which is far away from the real structures in nature. We have reason enough to believe that the $\langle \Psi_{tot} | (\mathbf{R}_{1,2} \cdot \mathbf{R}_{3,4}) | \Psi_{tot} \rangle$ should not be zero. Meanwhile, considering other spatial basis would reduce the corresponding to the binding energy B_T [65]. But these corrections would be powerless against the higher binding energy B_T of the ground $J^{PC} = 1^{+-} cb\bar{c}\bar{b}$. In conclusion, we tend to think that the $J^{PC} = 1^{+-} cb\bar{c}\bar{b}$ ground state should be an unstable compact state.

V. COMPARISON WITH OTHER WORK

The mass spectra have been studied with different approaches such as different nonrelativistic constituent quark models, different chromomagnetic models, the relativistic quark models, the nonrelativistic chiral quark model, the diquark models, the diffusion Monte Carlo calculation, and the QCD sum rule. In addition, these fully heavy tetraquark systems have been discussed with different color structures such as the $8_{Q\bar{Q}} \otimes 8_{Q\bar{Q}}$ configuration, the diquark-antiquark configuration ($3 \otimes \bar{3}$ and the $6 \otimes \bar{6}$) and the couplings between the above color configurations. For comparison, we briefly list our results and other theoretical results in Table X.

Compared to other systems, there is the most extensive discussion about the $cc\bar{c}\bar{c}$ system. So we will concentrate on the $cc\bar{c}\bar{c}$ system, but other systems can be discussed in a similar way. After comparing our results with other researches, we can see that the most theoretical masses of $cc\bar{c}\bar{c}$ in ground states lie in a wide range of 6.0 – 6.8 GeV in Table X. Our results are 6.38, 6.45, and 6.48 GeV for the 0^{++} , 1^{+-} , and 2^{++} $cc\bar{c}\bar{c}$ ground states, respectively. These three ground states are expected to be broad because they can all decay to charmonium pairs: $\eta_c\eta_c$, $\eta_c J/\psi$, or $J/\psi J/\psi$ through the quark (antiquark) rearrangements. Therefore, these types of decays are favored both dynamically and kinematically. According to Table X, we can conclude that the obtained masses of the ground states are obviously smaller than the $X(6900)$ observed by the LHCb collaboration. The observed $X(6900)$ is less likely to be the ground compact tetraquark state and could be a first or second radial excited $cc\bar{c}\bar{c}$ state.

Although we all use a similar Hamiltonian expression as in the nonrelativistic constituent quark model [111, 141–143], the spatial wave function is mostly expanded in the Gaussian basis according to Ref. [144], while we treat the spatial function as a Gaussian function, which is convenient for use in further variational methods to handle calculations in the four-body problem. Our results for the $cc\bar{c}\bar{c}$ system are roughly compatible with other nonrelativistic constituent quark models, although different papers have chosen different potential forms.

It is also interesting to note that relatively larger results are also given by the QCD sum rules [145], the Monte Carlo method [146], the diquark model [147], and the chiral quark model [148]. However, the results given by the QCD sum rules [145] are about 1 GeV below those of the constituent quark models for the $bb\bar{b}\bar{b}$ system. In contrast, our results are obviously larger than the chromomagnetic models [79, 107, 110, 123], and the diquark models [100, 149], where these models usually neglect the kinematic term and explicitly include confining potential contributions or adopt a diquark picture.

VI. SUMMARY

The discovery of exotic structures in the di- J/ψ invariant mass spectrum from the LHCb, CMS, and ATLAS collaborations gives us strong confidence to investigate the fully heavy tetraquark system. Thus, we use the variational method to systematically calculate the masses of all possible configurations for fully heavy tetraquarks within the framework of the constituent quark model. Meanwhile, we also give the corresponding internal mass contributions, the relative lengths between (anti)quarks, their lowest meson-meson thresholds, the specific wave function, the magnetic moments, the transition magnetic moments, the radiative decay widths, the rearrangement strong width ratios, and the comparisons with the two different CMI models.

To obtain the above results, we need to construct the total wave functions of the tetraquark states, including the flavor part, the color part, the spin part, and the spatial part, which is chosen to be a simple Gaussian form. Here, we first estimate the theoretical values of traditional hadrons, which are used to compare the experimental values to prove the reliability of this model. Before the discussing the numerical analysis, we analyze the stability condition by using only the color-spin interaction. Then, we obtain the specific numerical values and show them in corresponding Tables and the spatial distribution of valence quarks for the $J^{PC} = 0^{++} bb\bar{b}\bar{b}$ ground state in Fig. 1.

For the $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ systems, there are two pure neutral systems with definite C-parity. There are only two $J^{PC} = 0^{++}$ states, one $J^{PC} = 1^{+-}$ state, and one $J^{PC} = 2^{++}$ state, due to the Pauli principle. We also find that these states with different quantum numbers are all above the lowest thresholds, and have larger masses. Since these states are pure neutral particles, the corresponding magnetic moments are all 0 for the ground $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ states. Meanwhile, of course, the variational parameters C_{11} and C_{22} are the same, so the distances of the diquark and antidiquark are also the same. Moreover, the distances between quark and antiquark are all the same according to the symmetry analysis of Eqs. (30-31). Furthermore, three Jacobi coordinates are orthogonal to each other according to Eqs. (32-34). Based on this, we take the $J^{PC} = 0^{++} bb\bar{b}\bar{b}$ ground state as an example to

TABLE X. Comparison of the results of different methods for the $QQ\bar{Q}\bar{Q}$ tetraquark states.

Systems $J^{P(C)}$		$cc\bar{c}\bar{c}$				$bb\bar{b}\bar{b}$				$cc\bar{b}\bar{b}$			
		0^{++}	1^{+-}	2^{++}		0^{++}	1^{+-}	2^{++}		0^+	1^+	2^+	
The nonrelativistic constituent quark models	Our result	6384	6512	6452	6483	19352	19240	19304	19328	12920	13008	12940	12961
	Ref. [111]	6377	6425	6425	6432	19215	19247	19247	19249	12847	12866	12864	12868
		6371	6483	6450	6479	19243	19305	19311	19325	12886	12946	12924	12940
	Ref. [142]	6487	6518	6500	6524	19332	19338	19329	19341	12947	13039	12960	12972
	Ref. [143]	6500	6411	6453	6475	19200	19235	19216	19225	12880	12981	12890	12902
Ref. [78]	6477	6695	6528	6573	-	-	-	-	-	-	-	-	
Multiquark color flux-tube model	Ref. [141]	6491	6580	6607		19357	19413	19429		12963	13024	13041	
		6407	6463	6486		19329	19373	19387		12906	12946	12960	
The chromo-magnetic model	Ref. [79]	6035	6139	6194		18834	18890	18921		12597	12660	12695	
		6192	-	-		18826	-	-		-	-	-	
	Ref. [123]	6899	7016	6899	6956	20155	20275	20212	20243	13496	13634	13560	13595
		6035	6253	6137	6194	18834	18954	18890	18921	12597	12734	12660	12695
Ref. [110]	6045	6271	6231	6287	18836	18981	18969	19000	12596	12712	12672	12703	
Ref. [107]	6034	6254	6137	6194	18834	18953	18890	18921	-	-	-	-	
The Bethe-Salpeter equations	Ref. [150]	6419	6456	6516		19205	19221	19253		-	-	-	
The relativistic quark model	Ref. [100]	6190	6271	6367		19314	19320	19330		12846	12859	12883	
	Ref. [94]	6435	6542	6515	6543	19201	19255	19251	19262	-	-	-	-
Monte Carlo method	Ref. [146]	6351	6441	6471		19199	19276	19289		12865	12908	12926	
The diquark model	Ref. [149]	5966	6051	6223		18754	18808	18916		-	-	-	
	Ref. [147, 151]	6322	6354	6385		19666	19673	19680		12401	12409	12427	
The QCD sum rule method	Ref. [145]	6360	6540	6470	6520	18130	18150	18140	18150	-	-	-	-
	Ref. [152, 153]	5990	6050	6090		18840	18840	18850		-	-	-	-
The nonrelativistic chiral quark model	Ref. [148, 154]	6510	6600	6708		-	-	-		12684	12737	12791	
An effective potential model	Ref. [91]	6346	6476	6441	6475	19154	19226	19214	19232	-	-	-	-

Systems $J^{P(C)}$		$cb\bar{c}\bar{b}$				$cc\bar{c}\bar{b}$				$bb\bar{b}\bar{c}$						
		0^{++}	1^{+-}	1^{++}	2^{++}	0^+	1^+	2^+		0^+	1^+	2^+				
Our result		12760	12851	12797	12856	12857	128824		9621	96246	9706	9731	16044	16043	16125	16149
		12989	13008	12999	13056	12960	12971		9766	9729			16163	16144		
Ref. [143]		12783	12850	12802	12835	12851	12852		9665	9676	9699	9713	16061	16046	16079	16089
		12966	13035	12949	12964	12938	12964		9732	9718			16100	16089		
Ref. [142]		12835	12864	12852	12864	12870	12864		9740	9746	9749	9768	16158	16157	16164	16176
		12864	13050	13047	13052	13056	13070		9763	9757			16158	16167		
Ref. [141]		12894		12955		13000		9735	9766		9839	16175	16179	16274		
		12829		12881		12925		9670	9683		9732	16126	16130	16182		
Ref. [110]		12354		12436		12548		9705	9705		9732	15713	15729	15806		
		12363	12509	12425	12477	12524	12537		9318	9335	9384	9526	15712	15719	15851	15882
		12682	12747	12720	12744	12703	12755		9506	9499			15862	15854		
		13396	13634	13478	13592	13510	13590		10144	10174	10231	10273	16832	16840	16884	16917
Ref. [123]		13483	13553	13520	13555	13599	13599		10322	10282			16952	16915		
		12354	12592	12436	12550	12468	12548		9313	9343	9400	9442	15713	15729	15773	15806
		12441	12511	12478	12513	12557	12557		9491	9451			15841	15804		
		12578	12620	12496	12583	12611	12690									
	12656	12693	12653	12735	12700	12700										
Ref. [100]		12813	12824	12826	12831	12831	12849		9572	9602	9619	9647	16109	16117	16117	16132
Ref. [146]		12534		12510		12569	12582		9615	9610		9719	16040	16013	16129	
Ref. [149]		12359	12471	12424	12488	12485	12566		-	-	-	-	-	-	-	-
Ref. [154]		-		-		-	-		9579	9590	9613		16060	16062	16068	

show the spatial distribution of four valence quarks. As for the internal contribution, although the kinetic energy part is smaller than that of the $\eta_b\eta_b$ state, the V^C in η_b is much more attractive relative to the $J^{PC} = 0^{++} b\bar{b}b\bar{b}$ ground state, which is the main reason why this state has a larger mass than the meson-meson threshold. Similar situations also occur in other systems.

Similar to the $cc\bar{c}\bar{c}$ and $bb\bar{b}\bar{b}$ systems, the $cb\bar{c}\bar{b}$ system has the same number of the allowed ground states. According to the specific function, their mass contribution mainly comes from the $\bar{3} \otimes 3$ component within the diquark-antiquark configuration. Furthermore, we get the relevant the values of the magnetic moments, the transition magnetic moments, and the radiative decay widths. We also obtain the rearrangement strong width ratios within the meson-meson configuration.

As for the $cc\bar{c}\bar{b}$ and $bb\bar{b}\bar{c}$ systems, there are more allowed states due to fewer symmetry restrictions. Considering only the hyperfine potential, we can expect to have a compact stable state for $J^P = 1^+ bb\bar{b}\bar{c}$ configuration. However, since the V^C of the tetraquark are less attractive than the corresponding mesons, this state still has a mass larger than the meson-meson threshold.

In the $cb\bar{c}\bar{b}$ system, these states are also pure neutral particles, and we naturally obtain that their variational parameters C_{11} and C_{22} are the same. There is no constraint from the Pauli principle, so there are four $J^{PC} = 0^{++}$ states, four $J^{PC} = 1^{+-}$ states, two $J^{PC} = 1^{++}$ states, two $J^{PC} = 2^{++}$ states. All of the $cb\bar{c}\bar{b}$ states have larger masses relative to the lowest thresholds. Moreover, they all have two different rearrangement strong decay modes: $c\bar{c} - b\bar{b}$ and $c\bar{b} - b\bar{c}$.

Then we compare our results with other theoretical work. Our results are roughly compatible with other non-relativistic constituent quark models, although different papers have chosen different potential forms. Meanwhile, it is also interesting to find that similar mass ranges are given by the QCD sum rules, the Monte Carlo method, and the chiral quark model. This shows that our results are quite reasonable.

In summary, our theoretical calculations show that

the masses of the $cc\bar{c}\bar{c}$ ground states are around 6.45 GeV, which is obviously lower than 6.9 GeV. Thus, the experimentally observed $X(6900)$ state does not seem to be a ground $cc\bar{c}\bar{c}$ tetraquark state, but could be a radially or orbitally excited state. We also find that these lowest states all have a large positive binding energy B_T . In other words, all these states are found to have masses above the corresponding two meson decay thresholds via the quark rearrangement. Hence, we conclude that there is no compact bound ground fully heavy tetraquark state which is stable against the strong decay into two mesons within the constituent quark model. Finally, we hope that more relevant experimental analyses will be able to focus on this system in the near future.

VII. ACKNOWLEDGMENTS

This work is supported by the China National Funds for Distinguished Young Scientists under Grant No. 11825503, National Key Research and Development Program of China under Contract No. 2020YFA0406400, the 111 Project under Grant No. B20063, the National Natural Science Foundation of China under Grant No. 12247101, and the project for top-notch innovative talents of Gansu province. Z.W.L. would like to thank the support from the National Natural Science Foundation of China under Grants No. 12175091, and 11965016, and CAS Interdisciplinary Innovation Team.

VIII. APPENDIX

In this appendix, we show the masses, binding energies, variational parameters, internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between the quarks for the $cc\bar{b}\bar{b}$, $cc\bar{c}\bar{b}$, $bb\bar{b}\bar{c}$, and $cb\bar{c}\bar{b}$ states with different $J^{P(C)}$ quantum numbers and their lowest meson-meson thresholds.

-
- [1] M. Gell-Mann, A Schematic Model of Baryons and Mesons, *Phys. Lett.* **8** (1964), 214-215.
 - [2] G. Zweig, An SU(3) model for strong interaction symmetry and its breaking. Version 1, CERN-TH-401.
 - [3] G. Zweig, An SU(3) model for strong interaction symmetry and its breaking. Version 2, CERN-TH-412.
 - [4] S. K. Choi *et al.* [Belle Collaboration], Observation of a narrow charmonium - like state in exclusive $B^{+-} \rightarrow K^{+-}\pi^+\pi^- J/\psi$ decays, *Phys. Rev. Lett.* **91**, 262001 (2003).
 - [5] D. Acosta *et al.* [CDF Collaboration], Observation of the narrow state $X(3872) \rightarrow J/\psi\pi^+\pi^-$ in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV, *Phys. Rev. Lett.* **93**, 072001 (2004).
 - [6] V. M. Abazov *et al.* [D0 Collaboration], Observation and properties of the $X(3872)$ decaying to $J/\psi\pi^+\pi^-$ in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV, *Phys. Rev. Lett.* **93**, 162002 (2004).
 - [7] E. S. Swanson, The New heavy mesons: A Status report, *Phys. Rept.* **429**, 243 (2006).
 - [8] S. L. Zhu, New hadron states, *Int. J. Mod. Phys. E* **17** (2008) 283.
 - [9] M. B. Voloshin, Charmonium, *Prog. Part. Nucl. Phys.* **61** (2008) 455.
 - [10] N. Drenska, R. Faccini, F. Piccinini, A. Polosa, F. Renga and C. Sabelli, New Hadronic Spectroscopy, *Riv. Nuovo Cim.* **33** (2010) 633.

TABLE XI. The masses, binding energies, variational parameters, the internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between quarks for the $J^P = 0^+, 1^+$ $c\bar{c}b\bar{b}$ states and their lowest meson-meson thresholds. The notation is the same as that of Table VII.

$c\bar{c}b\bar{b}$		The contribution from each term			Relative Lengths (fm)		Overall	Present Work		CMI Model	
$J^P = 0^+$		Value	$B_c B_c$	Difference	(i, j)	Value		$B_c B_c$	Contribution	Value	Ref. [110]
Mass/ B_T		12920.0	12575.8	344.2	(1,2)	0.348		$2m_c$	3836.0		
Variational Parameters (fm ⁻²)	C_{11}	23.9	22.9		(1,3)	0.308	0.235(B_c)	$\frac{p_{x1}^2}{2m_1^2}$	319.0		
	C_{22}	10.5	22.9		(2,3)	0.308		$\frac{m_{\bar{b}}}{m_c+m_{\bar{b}}}\frac{p_{x3}^2}{2m_3^2}$	188.0		
	C_{33}	12.3	-		(1,4)	0.308		$V^C(12)$	-46.9	$\frac{1}{2}m_{cc}$	
Quark Mass		14522.0	14522.0	0.0	(2,4)	0.308	0.235(B_c)	$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	-95.0	1585.8	$2m_c$
Confinement Potential		-2420.1	-2795.5	375.4	(1,2)-(3,4):	0.226 fm		$-D$	-983.0	1578.7	3449.6
Kinetic Energy		835.9	947.3	-111.4	Radius:	0.151 fm		Subtotal	3218.1	3164.5	3449.6
CS Interaction		-7.0	-98.0	91.0							
V^C	(1,2)	-46.9			(1,3)	-47.5	-414.8(B_c)	$2m_b$	10686.0		
	(2,3)	-47.5			(3,4)	-217.1		$\frac{p_{x2}^2}{2m_2^2}$	216.4		
	(1,4)	-47.5			(2,4)	-47.5	-414.8(B_c)	$\frac{m_c}{m_c+m_{\bar{b}}}\frac{p_{x3}^2}{2m_3^2}$	67.5	$\frac{1}{2}m_{bb}$	
Subtotal	-454.0	-829.5	375.3				$V^C(34)$	-217.1	4764.8		
Total Contribution	-374.9	19.8	355.1				$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	-95.0	$\frac{m_{\bar{b}}}{m_{\bar{b}}+m_c}m_{c\bar{b}}$	$2m_b$	
Total Wave function:											
$\Psi_{tot} = 0.259 F\rangle R^s\rangle[\phi_2\chi_6\rangle] - 0.966 F\rangle R^s\rangle[\phi_1\chi_5\rangle] = -0.589 F\rangle R^s\rangle[\psi_1\zeta_5\rangle] + 0.095 F\rangle R^s\rangle[\psi_1\zeta_6\rangle] + 0.608 F\rangle R^s\rangle[\psi_2\zeta_5\rangle] + 0.524 F\rangle R^s\rangle[\psi_2\zeta_6\rangle]$											
The rearrangement strong width ratios :											
$\Gamma_{T_{c2\bar{b}2}(12920.0,0^{++})\rightarrow B_c^*B_c^+} : \Gamma_{T_{c2\bar{b}2}(12920.0,0,0^{++})\rightarrow B_c B_c} = 1 : 50$											
The radiative decay widths :											
$\Gamma_{T_{c2\bar{b}2}(12960.9,2^+)\rightarrow T_{c2\bar{b}2}(12920.0,0^+)\gamma} = 0 \text{ keV} ;$											
$\Gamma_{T_{c2\bar{b}2}(12939.9,1^+)\rightarrow T_{c2\bar{b}2}(12920.0,0^+)\gamma} = 3.8 \text{ keV}$											
The magnetic moments : $\mu_{T_{c2\bar{b}2}(12920.0,0^+)} = \langle \Psi_{tot}^0 \hat{\mu}^z \Psi_{tot}^0 \rangle = 0$											
The transition magnetic moments:											
$\mu_{T_{c2\bar{b}2}(12960.9,2^+)\rightarrow T_{c2\bar{b}2}(12920.0,0^+)\gamma} = \langle \Psi_{tot}^{2^+} \hat{\mu}^z \Psi_{tot}^0 \rangle = 0$											
$\mu_{T_{c2\bar{b}2}(12939.9,1^+)\rightarrow T_{c2\bar{b}2}(12920.0,0^+)\gamma} = \langle \Psi_{tot}^{1^+} \hat{\mu}^z \Psi_{tot}^0 \rangle = 0.966 \times \frac{4}{\sqrt{6}}(\mu_c - \mu_{\bar{b}}) = 0.534\mu_N$											
$J^P = 1^+$		Value	$B_c^* B_c$	Difference	Relative Lengths (fm)			Contribution	Value	Ref. [110]	Ref. [123]
Mass/ B_T		12939.9	12638.4	301.5	(i, j)	Value	$B_c^* B_c$	$2m_c$	3836.0		
Variational Parameters (fm ⁻²)	C_{11}	24.8	20.2		(1,2)	0.351		$\frac{p_{x1}^2}{2m_1^2}$	312.7		
	C_{22}	10.3	22.9		(1,3)	0.317	0.250(B_c^*)	$\frac{m_{\bar{b}}}{m_c+m_{\bar{b}}}\frac{p_{x3}^2}{2m_3^2}$	169.1	$\frac{1}{2}m_{cc}$	
	C_{33}	11.1	-		(2,3)	0.317		$V^C(12)$	-43.5	1585.8	
Quark Mass		14522.0	14522.0	0.0	(1,4)	0.317		$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	-83.2	$\frac{m_c}{m_{\bar{b}}+m_c}m_{c\bar{b}}$	$2m_c$
Confinement Potential		-2400.7	-2741.1	340.4	(2,4)	0.317	0.235(B_c)	$-D$	-983.0	1587.7	3449.6
Kinetic Energy		814.0	891.5	-77.5	(3,4)	0.226		Subtotal	3208.1	3164.5	3449.6
CS Interaction		4.6	-34.0	38.6							
V^C	(1,2)	-43.5			(1,2)-(3,4):	0.238 fm		$2m_b$	10686.0		
	(2,3)	-41.6			Radius:	0.157 fm		$\frac{p_{x2}^2}{2m_2^2}$	271.5		
	(1,4)	-41.6			(1,3)	-41.6	-360.4(B_c^*)	$\frac{m_c}{m_c+m_{\bar{b}}}\frac{p_{x3}^2}{2m_3^2}$	60.7	$\frac{1}{2}m_{bb}$	
Subtotal	-454.0	-829.5	375.3	(3,4)	-225.5		$V^C(34)$	-225.5	4764.8		
Total Contribution	383.9	82.4	301.5	(2,4)	-41.6	-414.8(B_c)	$\frac{1}{2}[V^C(13) + V^C(14) + V^C(23) + V^C(24)]$	-83.2	$\frac{m_{\bar{b}}}{m_{\bar{b}}+m_c}m_{c\bar{b}}$	$2m_b$	
Total Wave function: $\Psi_{tot} = F\rangle R^s\rangle[\phi_1\chi_2\rangle] = -0.419 F\rangle R^s\rangle[\psi_1\zeta_2\rangle] - 0.393 F\rangle R^s\rangle[\psi_1\zeta_3\rangle] - 0.066 F\rangle R^s\rangle[\psi_1\zeta_4\rangle] + 0.587 F\rangle R^s\rangle[\psi_2\zeta_2\rangle] + 0.557 F\rangle R^s\rangle[\psi_2\zeta_3\rangle] + 0.105 F\rangle R^s\rangle[\psi_2\zeta_4\rangle]$											
The rearrangement strong decay channel: $B^* B_c$											
The radiative decay widths: $\Gamma_{T_{b2\bar{b}2}(12960.9,2^+)\rightarrow T_{b2\bar{b}2}(12939.9,1^+)\gamma} = 3.6 \text{ keV}$											
$\Gamma_{T_{c2\bar{b}2}(12939.9,1^+)\rightarrow T_{c2\bar{b}2}(12920.0,0^+)\gamma} = 3.8 \text{ keV}$											
The magnetic moments:											
$\mu_{T_{c2\bar{b}2}(12939.9,1^+)} = \langle \Psi_{tot}^{1^+} \hat{\mu}^z \Psi_{tot}^{1^+} \rangle = \mu_c + \mu_{\bar{b}} = 0.490\mu_N$											
The transition magnetic moments: $\mu_{T_{b2\bar{b}2}(12960.9,2^+)\rightarrow T_{b2\bar{b}2}(12939.9,1^+)\gamma} = \langle \Psi_{tot}^{2^+} \hat{\mu}^z \Psi_{tot}^{1^+} \rangle = \mu_c - \mu_{\bar{b}} = 0.342\mu_N$											
$\mu_{T_{c2\bar{b}2}(12939.9,1^+)\rightarrow T_{c2\bar{b}2}(12920.0,0^+)\gamma} = \langle \Psi_{tot}^{1^+} \hat{\mu}^z \Psi_{tot}^0 \rangle = 0.966 \times \frac{4}{\sqrt{6}}(\mu_c - \mu_{\bar{b}}) = 0.534\mu_N$											
								Subtotal	9726.5	9508.4	10105.8
								$\frac{1}{2}V^{SS}(12)$	5.7	$\frac{8}{3}v_{cc} + \frac{8}{3}v_{bb}$	$\frac{8}{3}C_{cc} + \frac{8}{3}C_{bb}$
								$\frac{1}{2}V^{SS}(34)$	4.6	9.5+5.1	14.1+7.7
								$-\frac{1}{4}(V^{SS}(13) + V^{SS}(14) + V^{SS}(23) + V^{SS}(24))$	-10.6	$-\frac{16}{3}v_{c\bar{b}}$	$-\frac{16}{3}C_{c\bar{b}}$
								Subtotal	4.6	-1.2	4.3
								Total contribution	12939.9	12671.6	13560.0

TABLE XIII. The masses, binding energy, variational parameters, the internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between quarks for the $J^P = 0^+, 1^+ cc\bar{c}\bar{b}$ states and their lowest meson-meson thresholds. The notation is the same as that in Table VII.

$cc\bar{c}\bar{b}$		The contribution from each term			Relative Lengths (fm)		Overall	Present Work		CMI Model	
$J^P = 0^+$		Value	$B_c\eta_c$	Difference	(i, j)	Value		$B_c\eta_c$	Contribution	Value	Ref. [110]
Mass/ B_T		9620.5	9286.4	332.1	(1,2)	0.418		$2m_c$	3836.0	$-\frac{1}{4}m_{cc}$	
Variational Parameters (fm ⁻²)	C_{11}	11.4	22.9		(1,3)	0.325	0.290(η_c)	$\frac{\mathbf{p}_{x1}^2}{2m_1^2} + \frac{m_c+m_{\bar{b}}}{3m_c+m_{\bar{b}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$	451.6	$-\frac{1}{4}m_{cc}$	
	C_{22}	7.2	15.0		(2,3)	0.325		$V^C(12)$	-12.6	$\frac{5m_c}{4(m_c+m_{\bar{c}})}m_{c\bar{c}}$	
	C_{33}	15.2	-		(1,4)	0.336		$\frac{1}{2}[V^C(13)+V^C(23)]$	-91.8	1917.8	
Quark Mass		11097.0	11097.0	0.0	(1,4)	0.336		$\frac{1}{2}[V^C(14)+V^C(24)]$	-74.8	$\frac{5m_c}{4(m_c+m_{\bar{b}})}m_{c\bar{b}}$	$2m_c$
Confinement Potential		-2280.0	-2618.0	338.0	(2,4)	0.336	0.235(B_c)	$-D$	-983.0	1973.7	3449.6
Kinetic Energy		810.3	931.2	-120.9	(3,4)	0.333					
CS Interaction		18.2	-123.8	142.0	(1,2)-(3,4):	0.204 fm					
V^C	(1,2)	-12.6			(1,3)	-91.9	-237.2(η_c)				
	(1,4)	-74.8			(2,3)	-91.9					
	Subtotal	-314.0	-652.0	338.0	(2,4)	-74.8	-414.8(B_c)				
Total Contribution		514.4	155.4	359.0	(3,4)	32.0					
Total Wave function:											
$\Psi_{tot} = 0.401 F\rangle R^s\rangle[\phi_1\chi_5\rangle] - 0.916 F\rangle R^s\rangle[\phi_2\chi_6\rangle] = -0.574 F\rangle R^s\rangle[\psi_1\zeta_5\rangle] - 0.532 F\rangle R^s\rangle[\psi_1\zeta_6\rangle] + 0.019 F\rangle R^s\rangle[\psi_2\zeta_5\rangle] - 0.622 F\rangle R^s\rangle[\psi_2\zeta_6\rangle]$											
The rearrangement strong decay channel:											
$\Gamma_{T_{c2\bar{b}2}(9620.5,0^+) \rightarrow B_c^* J/\psi} : \Gamma_{T_{c2\bar{b}2}(9620.5,0^+) \rightarrow B_c\eta_c} = 1 : 1.2$											
The radiative decay widths: $\Gamma_{T_{c2\bar{c}\bar{b}}(9730.5,2^+) \rightarrow T_{c2\bar{c}\bar{b}}(9620.5,0^+)\gamma} = 0\text{keV}$ $\Gamma_{T_{c2\bar{c}\bar{b}}(9624.6,1^+) \rightarrow T_{c2\bar{c}\bar{b}}(9620.5,0^+)\gamma} = 0.007\text{keV}$											
The magnetic moments: $\mu_{T_{c2\bar{c}\bar{b}}(9620.5,0^+)} = \langle \Psi_{tot}^0 \hat{\mu}^z \Psi_{tot}^0 \rangle = 0$											
The transition magnetic moments:											
$\mu_{T_{c2\bar{c}\bar{b}}(9730.5,2^+) \rightarrow T_{c2\bar{c}\bar{b}}(9620.5,0^+)\gamma} = \langle \Psi_{tot}^2 \hat{\mu}^z \Psi_{tot}^0 \rangle = 0$											
$\mu_{T_{c2\bar{c}\bar{b}}(9624.6,1^+) \rightarrow T_{c2\bar{c}\bar{b}}(9620.5,0^+)\gamma} = \langle \Psi_{tot}^1 \hat{\mu}^z \Psi_{tot}^0 \rangle = -0.096\mu_N$											
$J^P = 1^+$		Value	$B_c^*\eta_c$	Difference	Relative Lengths (fm)			Contribution	Value	Ref. [110]	Ref. [123]
Mass/ B_T		9624.6	9349.0	275.6	(i, j)	$B_c^*\eta_c$					
Variational Parameters (fm ⁻²)	C_{11}	11.1	20.2		(1,2)	0.429		$2m_c$	3836.0	$-\frac{1}{4}m_{cc}$	
	C_{22}	6.9	15.0		(1,3)	0.328	0.290(η_c)	$\frac{\mathbf{p}_{x1}^2}{2m_1^2} + \frac{m_c+m_{\bar{b}}}{3m_c+m_{\bar{b}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$	442.8	$-\frac{1}{4}m_{cc}$	
	C_{33}	15.3	-		(2,3)	0.328		$V^C(12)$	-17.4	$\frac{5m_c}{4(m_c+m_{\bar{c}})}m_{c\bar{c}}$	
Quark Mass		11097.0	11097.0	0.0	(2,3)	0.328		$\frac{1}{2}[V^C(13)+V^C(23)]$	-87.0	1917.8	
Confinement Potential		-2266.2	-2563.6	297.4	(1,4)	0.340		$\frac{1}{2}[V^C(14)+V^C(24)]$	-68.9	$\frac{5m_c}{4(m_c+m_{\bar{b}})}m_{c\bar{b}}$	$2m_c$
Kinetic Energy		795.2	875.4	-80.2	(2,4)	0.340	0.250(B_c^*)	$-D$	-983.0	1973.7	3449.6
CS Interaction		8.0	-59.9	67.9	(3,4)	0.338					
V^C	(1,2)	-17.4			(1,2)-(3,4):	0.204 fm					
	(2,3)	-87.0			(2,4)	-68.9	-360.4(B_c^*)				
	(1,4)	-68.9			(3,4)	29.0					
Subtotal		-300.2	-597.6	297.4	(3,4)	29.0					
Total Contribution		503.0	218.0	285.0	(1,3)	-87.0	-237.2(η_c)				
Total Wave function:											
$\Psi_{tot} = 0.220 F\rangle R^s\rangle[\phi_1\chi_3\rangle] + 0.968 F\rangle R^s\rangle[\phi_2\chi_4\rangle] + 0.118 F\rangle R^s\rangle[\phi_1\chi_2\rangle] = 0.494 F\rangle R^s\rangle[\psi_1\zeta_2\rangle] - 0.396 F\rangle R^s\rangle[\psi_1\zeta_3\rangle] - 0.487 F\rangle R^s\rangle[\psi_1\zeta_4\rangle] + 0.111 F\rangle R^s\rangle[\psi_2\zeta_2\rangle] - 0.246 F\rangle R^s\rangle[\psi_2\zeta_3\rangle] - 0.537 F\rangle R^s\rangle[\psi_2\zeta_4\rangle]$											
The rearrangement strong decay channel: $\Gamma_{T_{c2\bar{c}\bar{b}}(9624.6,1^+) \rightarrow B_c^* J/\psi}$:											
$\Gamma_{T_{c2\bar{c}\bar{b}}(9624.6,1^+) \rightarrow B_c J/\psi} : \Gamma_{T_{c2\bar{c}\bar{b}}(9624.6,1^+) \rightarrow B_c^*\eta_c} = 1 : 0.8 : 1.2$											
The radiative decay widths: $\Gamma_{T_{c2\bar{c}\bar{b}}(9730.5,2^+) \rightarrow T_{c2\bar{c}\bar{b}}(9624.6,1^+)\gamma} = 145.0\text{keV}$ $\Gamma_{T_{c2\bar{c}\bar{b}}(9624.6,1^+) \rightarrow T_{c2\bar{c}\bar{b}}(9620.5,0^+)\gamma} = 0.007\text{keV}$											
The magnetic moments: $\mu_{T_{c2\bar{c}\bar{b}}(9624.6,1^+)} = \langle \Psi_{tot}^1 \hat{\mu}^z \Psi_{tot}^1 \rangle = -0.233\mu_N$											
The transition magnetic moments:											
$\mu_{T_{c2\bar{c}\bar{b}}(9624.6,1^+) \rightarrow T_{c2\bar{c}\bar{b}}(9620.5,0^+)\gamma} = \langle \Psi_{tot}^1 \hat{\mu}^z \Psi_{tot}^0 \rangle = -0.096\mu_N$ $\mu_{T_{c2\bar{c}\bar{b}}(9730.5,2^+) \rightarrow T_{c2\bar{c}\bar{b}}(9624.6,1^+)\gamma} = \langle \Psi_{tot}^2 \hat{\mu}^z \Psi_{tot}^1 \rangle = -0.294\mu_N$											
\bar{c} -quark:								$m_{\bar{c}}$	1918.0	$\frac{-m_{\bar{c}}}{4(m_{\bar{b}}+m_{\bar{c}})}m_{\bar{c}\bar{b}}$	
								$\frac{m_{\bar{b}}}{m_{\bar{c}}+m_{\bar{b}}}\frac{\mathbf{p}_{x2}^2}{2m_2^2} + \frac{m_c}{3m_c+m_{\bar{b}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$	230.3	-400.5	
								$\frac{1}{2}V^C(34)$	14.5	$\frac{5m_{\bar{b}}}{4(m_{\bar{c}}+m_{\bar{b}})}m_{\bar{c}\bar{b}}$	$m_{\bar{c}}$
$m_{\bar{c}}^{eff}$								$\frac{1}{2}[V^C(13)+V^C(23)]$	-87.0	1917.8	
								$-\frac{1}{2}D$	-491.5	1917.8	1724.8
								Subtotal	1584.3	1517.3	1724.8
\bar{b} -quark:								$m_{\bar{b}}$	5343.0	$\frac{-m_{\bar{b}}}{4(m_{\bar{b}}+m_{\bar{c}})}m_{\bar{c}\bar{b}}$	
								$\frac{m_{\bar{c}}}{m_{\bar{c}}+m_{\bar{b}}}\frac{\mathbf{p}_{x2}^2}{2m_2^2} + \frac{m_c}{3m_c+m_{\bar{b}}}\frac{\mathbf{p}_{x3}^2}{2m_3^2}$	122.1	-1203.5	
								$\frac{1}{2}V^C(34)$	14.5	$\frac{5m_{\bar{b}}}{4(m_{\bar{c}}+m_{\bar{b}})}m_{\bar{c}\bar{b}}$	$m_{\bar{b}}$
$m_{\bar{b}}^{eff}$								$\frac{1}{2}[V^C(14)+V^C(24)]$	-68.9	5929.5	5052.9
								$-\frac{1}{2}D$	-491.5	5929.5	5052.9
								Subtotal	4919.2	4726.0	5052.9
CS Interaction								$-\frac{1}{4}V^{SS}(12)$	-2.3	$-\frac{4}{3}v_{c\bar{b}}$	$-\frac{4}{3}C_{\bar{c}\bar{b}}$
								$\frac{3}{4}V^{SS}(34)$	10.4	-2.6	-4.4
								Subtotal	8.0	11.6	16.8
Matrix nondiagonal element									-9.3	-18.4	-100.0
Total contribution									9624.6	9335.1	10144

TABLE XIV. The masses, binding energy, variational parameters, the internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between quarks for the $J^P = 0^+, 1^+$ $bbb\bar{c}$ states and their lowest meson-meson thresholds. The notation is the same as that in Table VII.

$bbb\bar{c}$		The contribution from each term			Relative Lengths (fm)		Overall	Present Work		CMI Model		
$J^P = 0^+$	Value	$B_c\eta_b$	Difference	(i, j)	Vaule	$B_c\eta_b$		Contribution	Value	Ref. [110]	Ref. [123]	
Mass/ B_T	16043.9	15676.9	367.0	(1,2)	0.242		$2m_b$	10686.0	$-\frac{1}{4}m_{bb}$			
Variational Parameters (fm ⁻²)	C_{11} C_{22} C_{33}	12.5 21.7 28.7	22.9 58.8 -	(1,3) (2,3) (1,4)	0.239 0.239 0.249	0.148(η_b)	$\frac{p_1^2}{2m_1^2} + \frac{m_c+m_b}{3m_c+m_b} \frac{p_3^2}{2m_3^2}$	393.8	$-\frac{2382.4}{5m_b}$	$\frac{5903.1}{4(m_b+m_b)}$	$m_{b\bar{b}}$	
Quark Mass	17947.0	17947.0	0.0	(2,3)	0.239		$V^C(12)$	97.7	5903.1		$2m_b$	
Confinement Potential	-2786.7	-3259.9	473.2	(1,4)	0.249		$\frac{1}{2}[V^C(13) + V^C(23)]$	-251.3	$\frac{5m_b}{4(m_b+m_b)}$	$m_{b\bar{c}}$		
Kinetic Energy	883.3	1101.6	-218.3	(2,4)	0.249	0.235(B_c)	$\frac{1}{2}[V^C(14) + V^C(24)]$	-228.3	5929.5		10105.8	
CS Interaction	15.5	-111.8	127.3	(3,4)	0.318		$-D$	-983.0			10105.8	
V^C	(1,2) (1,4) Subtotal	97.7 -225.8 -820.7	-1293.9 473.2	(1,2)-(3,4) (1,3) (2,3) (2,4)	0.148 fm -251.3 -251.3 -228.3	0.148 fm (η_b) (B_c)	$\frac{m_b}{m_b+m_c} \frac{p_2^2}{2m_2^2} + \frac{m_b}{3m_b+m_c} \frac{p_3^2}{2m_3^2}$	5343.0 183.8 20.4	$\frac{-m_b}{4(m_c+m_b)}$	$m_{c\bar{b}}$		
Total Contribution	78.1	-304.1	382.2	(3,4)	40.8		$\frac{1}{2}[V^C(13) + V^C(23)]$	-491.5	5903.1		5052.9	
Total Wave function:								Subtotal	4804.4	4699.6	5052.9	
$\Psi_{tot} = 0.308 F\rangle R^s\rangle \phi_1\chi_5\rangle - 0.951 F\rangle R^s\rangle \phi_2\chi_6\rangle - 0.543 F\rangle R^s\rangle \psi_1\zeta_5\rangle - 0.584 F\rangle R^s\rangle \psi_1\zeta_6\rangle - 0.056 F\rangle R^s\rangle \psi_2\zeta_5\rangle - 0.602 F\rangle R^s\rangle \psi_2\zeta_6\rangle$							\bar{c} -quark:	$\frac{m_c}{m_b+m_c} \frac{p_2^2}{2m_2^2} + \frac{m_b}{3m_b+m_c} \frac{p_3^2}{2m_3^2}$	1918.0	$\frac{-m_c}{4(m_c+m_b)}$	$m_{b\bar{c}}$	
The rearrangement strong decay channel:								$\frac{1}{2}[V^C(14) + V^C(24)]$	-228.3	$\frac{5m_c}{4(m_b+m_c)}$	$m_{c\bar{b}}$	
$\Gamma_{T_{b_2\bar{b}\bar{c}}(16043.9,0^+) \rightarrow B_c^*\bar{c}} : \Gamma_{T_{b_2\bar{b}\bar{c}}(16043.9,0^+) \rightarrow B_c\eta_b} = 1 : 1.4$								$-\frac{1}{2}D$	-491.5	1973.8.8	1724.8	
The radiative decay widths: $\Gamma_{T_{b_2\bar{b}\bar{c}}(16149.2,2^+) \rightarrow T_{b_2\bar{b}\bar{c}}(16043.9,0^+)\gamma} = 0$ keV								Subtotal	1524.3	1573.3	1724.8	
$\Gamma_{T_{c_2\bar{c}\bar{b}}(16043.9,0^+) \rightarrow T_{c_2\bar{c}\bar{b}}(16043.2,1^+)\gamma} = 10^{-6}$ keV							CS Interaction	$\frac{3}{4}V^{SS}(12)$	7.9	4 v_{bb}	4 C_{bb}	
The magnetic moments: $\mu_{T_{b_2\bar{b}\bar{c}}(16043.9,0^+)} = \langle \Psi_{tot}^0 \hat{\mu}^z \Psi_{tot}^0 \rangle = 0$								$\frac{3}{4}V^{SS}(34)$	7.7	7.7	11.6	
The transition magnetic moments:								Subtotal	15.5	15.5	24.8	
$\mu_{T_{b_2\bar{b}\bar{c}}(16149.2,2^+) \rightarrow T_{b_2\bar{b}\bar{c}}(16043.9,0^+)\gamma} = \langle \Psi_{tot}^2 \hat{\mu}^z \Psi_{tot}^0 \rangle = 0$								Matrix nondiagonal element	-15.2	-26.7	43.7	
$\mu_{T_{b_2\bar{b}\bar{c}}(16043.9,0^+) \rightarrow T_{b_2\bar{b}\bar{c}}(16043.2,1^+)\gamma} = \langle \Psi_{tot}^0 \hat{\mu}^z \Psi_{tot}^1 \rangle = 0.096\mu_N$								Total contribution	16043.9	15711.9	16932.0	
$J^P = 1^+$	Value	$B_c^*\eta_b$	Difference	Relative Lengths (fm)			Contribution	Value	Ref. [110]	Ref. [123]		
Mass/ B_T	16043.2	15739.5	303.7	(i, j)	Value	$B_c^*\eta_b$	$2m_b$	10686.0	$-\frac{1}{4}m_{bb}$			
Variational Parameters (fm ⁻²)	C_{11} C_{22} C_{33}	12.4 21.0 28.9	20.2 57.4 -	(1,2) (1,3) (2,3)	0.245 0.240 0.240	0.148(η_b)	$\frac{p_1^2}{2m_1^2} + \frac{m_b+m_c}{3m_b+m_c} \frac{p_3^2}{2m_3^2}$	387.9	$-\frac{2382.4}{5m_b}$	$\frac{5903.1}{4(m_b+m_b)}$	$m_{b\bar{b}}$	
Quark Mass	17947.0	17947.0	0.0	(2,3)	0.240		$V^C(12)$	94.5	5903.1		$2m_b$	
Confinement Potential	-2779.8	-3205.5	425.7	(1,4)	0.250		$\frac{1}{2}[V^C(13) + V^C(23)]$	-248.2	$\frac{5m_b}{4(m_b+m_b)}$	$m_{b\bar{c}}$		
Kinetic Energy	876.1	961.5	-123.3	(2,4)	0.250	0.250(B_c^*)	$\frac{1}{2}[V^C(14) + V^C(24)]$	-225.8	5929.5		10105.8	
CS Interaction	4.8	-47.8	52.6	(3,4)	0.320		$-D$	-983.0			10105.8	
V^C	(1,2) (2,3) (1,4) Subtotal	94.5 -248.2 -225.8 -813.7	-1239.5 425.7	(1,2)-(3,4) (1,2) (2,3) (2,4)	0.148 fm 0.245 0.240 0.250	0.148 fm (B_c^*)	$\frac{m_b}{m_b+m_c} \frac{p_2^2}{2m_2^2} + \frac{m_b}{3m_b+m_c} \frac{p_3^2}{2m_3^2}$	5342.0 183.8 19.9	$\frac{-m_b}{4(m_c+m_b)}$	$m_{c\bar{b}}$		
Total Contribution	67.1	-241.5	308.6	(3,4)	39.7		$\frac{1}{2}[V^C(13) + V^C(23)]$	-248.2	5903.1		5052.9	
Total Wave function:								Subtotal	4807.0	4699.6	5052.9	
$\Psi_{tot} = 0.171 F\rangle R^s\rangle \phi_1\chi_3\rangle + 0.984 F\rangle R^s\rangle \phi_2\chi_4\rangle + 0.044 F\rangle R^s\rangle \phi_1\chi_2\rangle = 0.431 F\rangle R^s\rangle \psi_1\zeta_2\rangle - 0.467 F\rangle R^s\rangle \psi_1\zeta_3\rangle - 0.500 F\rangle R^s\rangle \psi_1\zeta_4\rangle + 0.240 F\rangle R^s\rangle \psi_2\zeta_2\rangle - 0.188 F\rangle R^s\rangle \psi_2\zeta_3\rangle - 0.502 F\rangle R^s\rangle \psi_2\zeta_4\rangle$							\bar{c} -quark:	$\frac{m_c}{m_b+m_c} \frac{p_2^2}{2m_2^2} + \frac{m_b}{3m_b+m_c} \frac{p_3^2}{2m_3^2}$	1918.0	$\frac{-m_c}{4(m_c+m_b)}$	$m_{b\bar{c}}$	
The rearrangement strong decay channel: $\Gamma_{T_{b_2\bar{b}\bar{c}}(16043.2,1^+) \rightarrow B_c^*\bar{c}} :$								$\frac{1}{2}[V^C(14) + V^C(24)]$	-225.8	$\frac{5m_c}{4(m_b+m_c)}$	$m_{c\bar{b}}$	
$\Gamma_{T_{b_2\bar{b}\bar{c}}(16043.2,1^+) \rightarrow B_c^*\eta_b} = 1 : 1.3 : 1.5$								$-\frac{1}{2}D$	-491.5	1973.8	1724.8	
The radiative decay widths: $\Gamma_{T_{b_2\bar{b}\bar{c}}(16149.2,2^+) \rightarrow T_{c_2\bar{c}\bar{b}}(16043.2,1^+)\gamma} = 435.0$ keV								Subtotal	1525.0	1573.3	1724.8	
$\Gamma_{T_{c_2\bar{c}\bar{b}}(16043.9,0^+) \rightarrow T_{c_2\bar{c}\bar{b}}(16043.2,1^+)\gamma} = 10^{-6}$ keV							CS Interaction	$-\frac{1}{4}V^{SS}(12)$	-2.6	$-\frac{4}{3}v_{b\bar{c}}$	$-\frac{4}{3}C_{b\bar{c}}$	
The magnetic moments: $\mu_{T_{b_2\bar{b}\bar{c}}(16043.2,1^+)} = \langle \Psi_{tot}^1 \hat{\mu}^z \Psi_{tot}^1 \rangle = -0.346\mu_N$								$\frac{3}{4}V^{SS}(34)$	7.5	7.7	11.6	
The transition magnetic moments:								Subtotal	4.8	5.0	7.2	
$\mu_{T_{b_2\bar{b}\bar{c}}(16043.9,0^+) \rightarrow T_{b_2\bar{b}\bar{c}}(16043.2,1^+)\gamma} = \langle \Psi_{tot}^0 \hat{\mu}^z \Psi_{tot}^1 \rangle = 0.201\mu_N$								Matrix nondiagonal element	-5.0	-8.6	24.3	
$\mu_{T_{b_2\bar{b}\bar{c}}(16149.2,2^+) \rightarrow T_{b_2\bar{b}\bar{c}}(16043.2,1^+)\gamma} = \langle \Psi_{tot}^2 \hat{\mu}^z \Psi_{tot}^1 \rangle = 0.329\mu_N$								Total contribution	16048.1	15719.1	16915.0	

TABLE XV. The masses, binding energy, variational parameters, the internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between quarks for the $J^P = 2^+ cc\bar{c}\bar{b}$ and $bb\bar{b}\bar{c}$ states and their lowest meson-meson thresholds. The notation is the same as that in Table VII.

$cc\bar{c}\bar{b}$		The contribution from each term			Relative Lengths (fm)		Overall	Present Work		CMI Model	
$J^P = 2^+$		Value	$B_c^* J/\psi$	Difference	(i, j)	Value		$B_c J/\psi$	Contribution	Value	Ref. [110]
Mass/ B_T		9730.5	9442.7	287.8	(1,2)	0.378		$2m_c$	3836.0	$\frac{1}{2}m_{cc}$	
Variational Parameters (fm ⁻²)	C_{11}	13.7	20.2		(1,3)	0.350	0.318(J/ψ)	$\frac{p_1^2}{2m_1^2} + \frac{m_c+m_b}{3m_c+m_b} \frac{p_3^2}{2m_3^2}$	408.0	$\frac{1585.8}{4(m_c+m_b)} m_{c\bar{c}}$	
	C_{22}	8.9	12.5		(2,3)	0.350		$V^C(12)$	-14.3	$\frac{767.1}{4(m_c+m_b)} m_{c\bar{c}}$	
	C_{33}	9.1	-		(1,4)	0.359		$\frac{1}{2}[V^C(13) + V^C(23)]$	-21.9	$\frac{m_c}{4(m_c+m_b)} m_{c\bar{c}}$	$2m_c$
Quark Mass		11097.0	11097.0	0.0	(1,4)	0.359		$\frac{1}{2}[V^C(14) + V^C(24)]$	-17.3	$\frac{m_c}{4(m_c+m_b)} m_{c\bar{c}}$	
Confinement Potential		-2158.4	-2490.6	332.2	(2,4)	0.359	0.250(B_c^*)	$-D$	-983.0		3449.6
Kinetic Energy		763.9	799.3	-35.4	(3,4)	0.304					3449.6
CS Interaction		28.0	36.9	-8.9	(1,2)-(3,4):		0.264 fm				
V^C	(1,2)	-14.3			(1,3)	-21.9	-164.2(J/ψ)	$\frac{m_b}{m_c+m_b} \frac{p_2^2}{2m_2^2} + \frac{m_c}{3m_c+m_b} \frac{p_3^2}{2m_3^2}$	1918.0	$\frac{m_c}{4(m_c+m_b)} m_{c\bar{c}}$	
	(1,4)	-17.3			(2,3)	-21.9		$\frac{1}{2}V^C(34)$	244.7	767.1	
	Subtotal	-192.4	-524.6	332.2	(2,4)	-17.3	-360.4(B_c^*)	$\frac{1}{2}[V^C(13) + V^C(23)]$	-21.9	$\frac{m_c}{4(m_c+m_b)} m_{c\bar{c}}$	m_c
Total Contribution		599.5	311.7	287.8	(3,4)	-99.6		$-\frac{1}{2}D$	-491.5	801.1	1724.8
Total Wave function:								Subtotal	1599.5	1568.2	1724.8
$\Psi_{tot} = F\rangle R^s\rangle[\phi_1\chi_1\rangle] = 0.577 F\rangle R^s\rangle[\psi_1\zeta_1\rangle] - 0.816 F\rangle R^s\rangle[\psi_2\zeta_1\rangle]$								$\frac{m_b}{m_c+m_b} \frac{p_2^2}{2m_2^2} + \frac{m_c}{3m_c+m_b} \frac{p_3^2}{2m_3^2}$	5343.0	$\frac{m_b}{4(m_b+m_c)} m_{c\bar{b}}$	
The rearrangement strong decay channel: $B_c^* J/\psi$.								$\frac{1}{2}V^C(34)$	111.2		2407.0
The radiative decay widths:								$\frac{1}{2}[V^C(14) + V^C(24)]$	-49.8	$\frac{m_b}{4(m_c+m_b)} m_{c\bar{b}}$	m_b
$\Gamma_{T_{c_2\bar{c}\bar{b}}(9730.5,2^+) \rightarrow T_{c_2\bar{c}\bar{b}}(9620.5,0^+)\gamma} = 0 \text{ keV}$								$-\frac{1}{2}D$	-491.5	2371.8	5052.9
$\Gamma_{T_{c_2\bar{c}\bar{b}}(9730.5,2^+) \rightarrow T_{c_2\bar{c}\bar{b}}(9624.6,1^+)\gamma} = 145.0 \text{ keV}$								Subtotal	4895.5	4778.8	5052.9
The magnetic moments:								$\frac{1}{2}V^{SS}(12)$	5.6	$\frac{8}{3}v_{cc} + \frac{8}{3}v_{c\bar{b}}$	$\frac{8}{3}C_{cc} + \frac{8}{3}C_{c\bar{b}}$
$\mu_{T_{c_2\bar{c}\bar{b}}(9730.5,2^+)} = \langle \Psi_{tot}^{2^+} \hat{\mu}^z \Psi_{tot}^{2^+} \rangle = 2\mu_c + \mu_{\bar{c}} + \mu_{\bar{b}} = 0.464\mu_N$								$\frac{1}{2}V^{SS}(34)$	8.5	$9.5+5.2$	$14.1+8.8$
The transition magnetic moments:								$\frac{1}{4}(V^{SS}(13) + V^{SS}(23))$	9.5	$\frac{8}{3}v_{c\bar{c}} + \frac{8}{3}v_{c\bar{b}}$	$\frac{8}{3}C_{c\bar{c}} + \frac{8}{3}C_{c\bar{b}}$
$\mu_{T_{c_2\bar{c}\bar{b}}(9730.5,2^+) \rightarrow T_{c_2\bar{c}\bar{b}}(9620.5,0^+)\gamma} = \langle \Psi_{tot}^{2^+} \hat{\mu}^z \Psi_{tot}^{0^+} \rangle = 0$								$\frac{1}{4}(V^{SS}(14) + V^{SS}(24))$	4.3	$14.2+7.9$	$14.1+8.8$
$\mu_{T_{c_2\bar{c}\bar{b}}(9730.5,2^+) \rightarrow T_{c_2\bar{c}\bar{b}}(9624.6,1^+)\gamma} = \langle \Psi_{tot}^{2^+} \hat{\mu}^z \Psi_{tot}^{1^+} \rangle = -0.294\mu_N$								Subtotal	28.0	36.8	45.9
Total contribution									9730.5	9526.0	10273.2
$bb\bar{b}\bar{c}$ $J^P = 2^+$		Value	$B_c^* \Upsilon$	Difference	Relative Lengths (fm)			Contribution	Value	Ref. [110]	Ref. [123]
Mass/ B_T		16149.2	15819.4	329.8	(i, j)	Value	$B_c^* \Upsilon$	$2m_c$	10686.0	$\frac{1}{2}m_{bb}$	
Variational Parameters (fm ⁻²)	C_{11}	14.4	20.2		(1,2)	0.210		$\frac{p_1^2}{2m_1^2} + \frac{m_b+m_{\bar{c}}}{3m_b+m_{\bar{c}}} \frac{p_3^2}{2m_3^2}$	404.4	$\frac{4764.8}{4(m_b+m_{\bar{c}})} m_{bb}$	
	C_{22}	28.6	49.7		(1,3)	0.256	0.160(Υ)	$V^C(12)$	-257.5	$\frac{2361.2}{4(m_b+m_{\bar{c}})} m_{bb}$	
	C_{33}	16.9	-		(2,3)	0.256		$\frac{1}{2}[V^C(13) + V^C(23)]$	-85.5	$\frac{m_b}{4(m_b+m_{\bar{c}})} m_{b\bar{c}}$	$2m_b$
Quark Mass		17947.0	17497.0	0.0	(1,4)	0.266		$\frac{1}{2}[V^C(14) + V^C(24)]$	-77.6	$\frac{m_b}{4(m_b+m_{\bar{c}})} m_{b\bar{c}}$	
Confinement Potential		-2659.7	-3123.1	463.4	(2,4)	0.266	0.250(B_c^*)	$-D$	-983.0		10105.8
Kinetic Energy		838.2	961.5	-123.3	(3,4)	0.296					10105.8
CS Interaction		23.8	34.0	-10.2	(1,2)-(3,4):		0.194 fm				
V^C	(1,2)	-257.5			(1,2)	-257.5		$\frac{m_b}{m_c+m_b} \frac{p_2^2}{2m_2^2} + \frac{m_b}{3m_b+m_{\bar{c}}} \frac{p_3^2}{2m_3^2}$	5343.0	$\frac{m_b}{4(m_b+m_{\bar{c}})} m_{bb}$	
	(2,3)	-85.5			(2,4)	-77.6	-360.4(B_c^*)	$\frac{1}{2}V^C(34)$	146.6	2382.4	
	(1,4)	-77.6			(2,4)	-77.6	-360.4(B_c^*)	$\frac{1}{2}[V^C(13) + V^C(23)]$	-55.1	$\frac{m_b}{4(m_b+m_{\bar{c}})} m_{b\bar{c}}$	m_b
Subtotal		-693.7	-1157.1	463.4	(3,4)	-110.1		$-\frac{1}{2}D$	-491.5	2360.6	5052.9
Total Contribution		168.2	-161.6	329.8	(1,3)	-85.5	-796.7(Υ)	Subtotal	4857.5	4743.0	5052.9
Total Wave function:								$\frac{m_b}{m_c+m_b} \frac{p_2^2}{2m_2^2} + \frac{m_b}{3m_b+m_{\bar{c}}} \frac{p_3^2}{2m_3^2}$	1918.0	$\frac{m_c}{4(m_b+m_{\bar{c}})} m_{c\bar{b}}$	
$\Psi_{tot} = F\rangle R^s\rangle[\phi_1\chi_1\rangle] = 0.577 F\rangle R^s\rangle[\psi_1\zeta_1\rangle] - 0.816 F\rangle R^s\rangle[\psi_2\zeta_1\rangle]$								$\frac{1}{2}V^C(34)$	287.2		801.3
The rearrangement strong decay channel: $B_c^* \Upsilon$.								$\frac{1}{2}[V^C(14) + V^C(24)]$	-55.1	$\frac{m_c}{4(m_b+m_{\bar{c}})} m_{b\bar{c}}$	m_c
The radiative decay widths:								$-\frac{1}{2}D$	-491.5	789.3	1724.8
$\Gamma_{T_{b_2\bar{b}\bar{c}}(16149.2,2^+) \rightarrow T_{b_2\bar{b}\bar{c}}(16043.9,0^+)\gamma} = 0 \text{ keV}$								Subtotal	1581.2	1590.6	1724.8
$\Gamma_{T_{b_2\bar{b}\bar{c}}(16149.2,2^+) \rightarrow T_{c_2\bar{c}\bar{b}}(16043.2,1^+)\gamma} = 435.0 \text{ keV}$								$\frac{1}{2}V^{SS}(12)$	5.9	$\frac{8}{3}v_{bb} + \frac{8}{3}v_{b\bar{c}}$	$\frac{8}{3}C_{bb} + \frac{8}{3}C_{c\bar{b}}$
The magnetic moments:								$\frac{1}{2}V^{SS}(34)$	6.4	$5.1+5.2$	$7.7+8.8$
$\mu_{T_{b_2\bar{b}\bar{c}}(16149.2,2^+)} = \langle \Psi_{tot}^{2^+} \hat{\mu}^z \Psi_{tot}^{2^+} \rangle = 2\mu_b + \mu_{\bar{b}} + \mu_{\bar{c}} = -0.472\mu_N$								$\frac{1}{4}(V^{SS}(13) + V^{SS}(23))$	4.6	$\frac{8}{3}v_{b\bar{b}} + \frac{8}{3}v_{b\bar{c}}$	$\frac{8}{3}C_{b\bar{b}} + \frac{8}{3}C_{b\bar{c}}$
The transition magnetic moments:								$\frac{1}{4}(V^{SS}(14) + V^{SS}(24))$	6.9	$7.7+7.9$	$7.7+8.8$
$\mu_{T_{b_2\bar{b}\bar{c}}(16149.2,2^+) \rightarrow T_{b_2\bar{b}\bar{c}}(16043.2,1^+)\gamma} = \langle \Psi_{tot}^{2^+} \hat{\mu}^z \Psi_{tot}^{1^+} \rangle = 0.329\mu_N$								Subtotal	23.8	25.9	33.1
$\mu_{T_{b_2\bar{b}\bar{c}}(16149.2,2^+) \rightarrow T_{b_2\bar{b}\bar{c}}(16043.9,0^+)\gamma} = \langle \Psi_{tot}^{2^+} \hat{\mu}^z \Psi_{tot}^{0^+} \rangle = 0$								Total contribution	16149.2	15882.3	16917.0

TABLE XVI. The masses, binding energies, variational parameters, the internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between quarks for the the $J^{PC} = 0^{++}$, 2^{++} $cb\bar{c}\bar{b}$ states and their lowest meson-meson thresholds. The notation is the same as that in Table VII.

$cb\bar{c}\bar{b}$		The contribution from each term			Relative Lengths (fm)		Overall	Present Work		CMI Model	
$J^{PC} = 0^{++}$	Value	$\eta_b\eta_c$	Difference	(i, j)	Value	$\eta_b\eta_c$		Contribution	Value	Ref. [110]	Ref. [123]
Mass/ B_T	12759.6	12387.5	372.1	(1,2)	0.315		m_c	1918.0	$\frac{-m_c}{4(m_c+m_b)}m_{cb}$		
Variational Parameters (fm ⁻²)	C_{11}	12.7	15.0	(1,3)	0.277	0.290(η_c)	$\frac{m_b}{m_c+m_b}\frac{p_{x1}^2}{2m_1^2}$	193.7	$\frac{-m_c}{4(m_c+m_b)}m_{cb}$		
	C_{22}	12.7	57.4	(2,3)	0.277		$\frac{m_b}{2m_c+2m_b}\frac{p_{x3}^2}{2m_3^2}$	139.6	$\frac{5m_c}{8(m_c+m_b)}m_{c\bar{b}}$		
	C_{33}	23.6	-	(1,4)	0.277		$\frac{1}{2}V^C(12)$	21.3	986.7		
Quark Mass	14522.0	14522.0	0.0	(2,4)	0.277	0.148(η_b)	$\frac{1}{2}[V^C(13) + V^C(14)]$	-172.6	$\frac{5m_c}{8(m_c+m_b)}m_{c\bar{e}}$	m_c	
Confinement Potential	-2571.3	-3082.3	511.0	(3,4)	0.315		$-1/2D$	-491.5	958.9	1724.8	
Kinetic Energy	905.8	1085.5	-179.7	(1,2)-(3,4):	0.164 fm		Subtotal	1608.5	1545.1	1724.8	
CS Interaction	-81.1	-137.6	56.6								
V^C	(1,2)	42.6		(1,3)	-172.6	-237.2(η_c)	$\frac{m_b}{m_c+m_b}\frac{p_{x1}^2}{2m_1^2}$	5343.0	$\frac{-m_b}{4(m_c+m_b)}m_{cb}$		
	(1,4)	-172.6		(2,3)	-172.6		$\frac{m_c}{2m_c+2m_b}\frac{p_{x3}^2}{2m_3^2}$	69.5	$\frac{-m_b}{4(m_c+m_b)}m_{cb}$		
	Subtotal	-605.3	-1116.3	511.0	(2,4)	-172.6	-879.1(η_b)	$\frac{1}{2}V^C(12)$	50.1	$\frac{5m_b}{8(m_c+m_b)}m_{c\bar{b}}$	
Total Contribution	219.3	-168.5	387.8	(3,4)	42.6		$\frac{1}{2}[V^C(23) + V^C(24)]$	21.3	2965.5		
							$-1/2D$	-491.5	$\frac{5m_b}{8(m_b+m_b)}m_{b\bar{b}}$	m_b	
							Subtotal	4819.8	4713.3	5052.9	
Total Wave function:							CS Interaction	$-\frac{1}{4}(V^{SS}(12) + V^{SS}(34))$	-5.3	$-\frac{8}{3}v_{cb} - \frac{20}{3}v_{c\bar{e}}$	$-\frac{8}{3}C_{cb} - \frac{20}{3}C_{c\bar{e}}$
$\Psi_{tot} = 0.961 F\rangle R^s\rangle[\phi_2\chi_5\rangle + 0.114 F\rangle R^s\rangle[\phi_2\chi_6\rangle - 0.069 F\rangle R^s\rangle[\phi_1\chi_5\rangle$								$-\frac{5}{4}V^{SS}(13)$	-33.3	-5.1-35.5	-8.8-35.3
$-0.241 F\rangle R^s\rangle[\phi_1\chi_6\rangle = 0.211 F\rangle R^s\rangle[\psi_1\zeta_6\rangle - 0.830 F\rangle R^s\rangle[\psi_1\zeta_5\rangle$								$-\frac{5}{4}V^{SS}(24)$	-10.1	$-\frac{20}{3}v_{b\bar{b}} - \frac{40}{3}v_{b\bar{c}}$	$-\frac{20}{3}C_{b\bar{b}} - \frac{40}{3}C_{b\bar{c}}$
$-0.367 F\rangle R^s\rangle[\psi_2\zeta_5\rangle + 0.363 F\rangle R^s\rangle[\psi_2\zeta_6\rangle = 0.333 F\rangle R^s\rangle[\psi_1'\zeta_6\rangle$								$-\frac{5}{4}(V^{SS}(14) + V^{SS}(23))$	-32.5	-19.2-39.3	-19.3-44.3
$-0.668 F\rangle R^s\rangle[\psi_1'\zeta_5\rangle - 0.398 F\rangle R^s\rangle[\psi_2'\zeta_5\rangle + 0.533 F\rangle R^s\rangle[\psi_2'\zeta_6\rangle$								Subtotal	-81.1	-99.2	-107.5
The rearrangement strong decay channel:							Matrix nondiagonal element		-15.8	-80.7	-51.9
$\Gamma_{T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\rightarrow\eta_c\eta_b} : \Gamma_{T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\rightarrow J/\psi\Upsilon} = 21 : 1$							Total contribution		12759.6	12336.1	13396.0
$\Gamma_{T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\rightarrow B_c B_c} : \Gamma_{T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\rightarrow B_c^* B_c^*} = 7.6 : 1$							The radiative decay widths:				
$\Gamma_{T_{cb\bar{e}\bar{b}}(12797.3,1^{+-})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\gamma} = 33.1$ keV							The magnetic moments: $T_{cb\bar{e}\bar{b}}(12759.6,0^{++}) = \langle \Psi_{tot}^{0^{++}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0$				
$\Gamma_{T_{cb\bar{e}\bar{b}}(12856.6,1^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\gamma} = 0$							The transition magnetic moments: $\mu_{T_{cb\bar{e}\bar{b}}(12882.4,2^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\gamma} = \langle \Psi_{tot}^{2^{++}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0$				
$\Gamma_{T_{cb\bar{e}\bar{b}}(12882.4,2^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\gamma} = 0$							$\mu_{T_{cb\bar{e}\bar{b}}(12856.6,1^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\gamma} = \langle \Psi_{tot}^{1^{++}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0$				
$\Gamma_{T_{cb\bar{e}\bar{b}}(12882.4,2^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\gamma} = 0$							$\mu_{T_{cb\bar{e}\bar{b}}(12796.9,1^{+-})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\gamma} = \langle \Psi_{tot}^{1^{+-}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0.226 \times 2\sqrt{\frac{1}{6}}(\mu_c - \mu_b) = 0.081\mu_N$				
$J^{PC} = 2^{++}$	Value	$\Upsilon J/\psi$	Difference	Relative Lengths (fm)			Contribution	Value	Ref. [110]	Ref. [123]	
Mass/ B_T	12882.4	12561.1	321.3	(i, j)	Value	$\Upsilon J/\psi$	m_c	1918.0	$\frac{-m_c}{4(m_c+m_b)}m_{cb}$		
Variational Parameters (fm ⁻²)	C_{11}	11.0	12.5	(1,3)	0.340		$\frac{m_b}{m_c+m_b}\frac{p_{x1}^2}{2m_1^2}$	167.1	$\frac{-m_c}{4(m_c+m_b)}m_{cb}$		
	C_{22}	11.0	49.7	(1,3)	0.296	0.318(J/ψ)	$\frac{m_b}{2m_c+2m_b}\frac{p_{x3}^2}{2m_3^2}$	124.2	$\frac{5m_c}{8(m_c+m_b)}m_{c\bar{b}}$		
	C_{33}	21.0	-	(2,3)	0.296		$\frac{1}{2}V^C(12)$	14.0	986.7		
Quark Mass	14522.0	14522.0	0.0	(2,4)	0.296		$\frac{1}{2}[V^C(13) + V^C(14)]$	-137.7	$\frac{5m_c}{8(m_c+m_b)}m_{c\bar{e}}$	m_c	
Confinement Potential	-2460.7	-2926.9	466.2	(1,4)	0.296		$-1/2D$	-491.5	958.9	1724.8	
Kinetic Energy	791.7	925.1	-133.4	(2,4)	0.296	0.160(Υ)	Subtotal	1594.1	1545.1	1724.8	
CS Interaction	29.5	41.0	-11.5	(3,4)	0.340						
V^C	(1,2)	28.0		(1,2)-(3,4):	0.174 fm		$\frac{m_b}{m_c+m_b}\frac{p_{x1}^2}{2m_1^2}$	5343.0	$\frac{-m_b}{4(m_c+m_b)}m_{cb}$		
	(2,3)	-137.7		(2,4)	-137.7	-796.7(Υ)	$\frac{m_c}{2m_c+2m_b}\frac{p_{x3}^2}{2m_3^2}$	60.0	$\frac{-m_b}{4(m_c+m_b)}m_{cb}$		
	(1,4)	-137.7		(3,4)	28.0		$\frac{1}{2}V^C(12)$	44.6	$\frac{5m_b}{8(m_c+m_b)}m_{c\bar{b}}$		
Subtotal	-494.7	-960.9	466.2				$\frac{1}{2}[V^C(23) + V^C(24)]$	14.0	2965.5		
Total Contribution	326.5	5.1	321.4	(1,3)	-137.7	-164.2(J/ψ)	$-1/2D$	-491.5	2951.6	5052.9	
							Subtotal	4832.4	4713.3	5052.9	
Total Wave function:							CS Interaction	$-\frac{1}{4}(V^{SS}(12) + V^{SS}(34))$	-4.7	$-\frac{8}{3}v_{cb} + \frac{10}{3}v_{c\bar{e}}$	$-\frac{8}{3}C_{cb} + \frac{10}{3}C_{c\bar{e}}$
$\Psi_{tot} = 0.999 F\rangle R^s\rangle[\phi_2\chi_1\rangle - 0.027 F\rangle R^s\rangle[\phi_1\chi_1\rangle = -0.801 F\rangle R^s\rangle[\psi_1\zeta_1\rangle$								$\frac{5}{8}V^{SS}(13)$	15.2	-5.1+17.7	-8.8+17.7
$-0.599 F\rangle R^s\rangle[\psi_2\zeta_1\rangle = -0.832 F\rangle R^s\rangle[\psi_1'\zeta_1\rangle + 0.555 F\rangle R^s\rangle[\psi_2'\zeta_1\rangle$								$\frac{5}{8}V^{SS}(24)$	4.5	$\frac{10}{3}v_{b\bar{b}} + \frac{20}{3}v_{b\bar{c}}$	$\frac{10}{3}C_{b\bar{b}} + \frac{20}{3}C_{b\bar{c}}$
The rearrangement strong decay channel: $\Upsilon J/\psi$ and $B_c^* B_c^*$								$\frac{5}{8}(V^{SS}(14) + V^{SS}(23))$	14.7	9.6+19.7	9.6+22.0
The magnetic moments: $T_{cb\bar{e}\bar{b}}(12882.4,2^{++}) = \langle \Psi_{tot}^{2^{++}} \hat{\mu}^z \Psi_{tot}^{2^{++}} \rangle = 0$							Subtotal	29.5	41.7	40.5	
The radiative decay widths:							Matrix nondiagonal element		0.1	-29.1	3.2
$\Gamma_{T_{cb\bar{e}\bar{b}}(12882.4,2^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12797.3,1^{+-})\gamma} = 4.9$ keV							Total contribution		12882.4	12529.4	13599.0
$\Gamma_{T_{cb\bar{e}\bar{b}}(12882.4,2^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\gamma} = 0$							The transition magnetic moments: $\mu_{T_{cb\bar{e}\bar{b}}(12882.4,2^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,1^{+-})\gamma} = \langle \Psi_{tot}^{2^{++}} \hat{\mu}^z \Psi_{tot}^{1^{+-}} \rangle = 0.774 \times (\mu_c + \mu_b) + 0.226 \times (\mu_c - \mu_b) = 0.345\mu_N$				
$\Gamma_{T_{cb\bar{e}\bar{b}}(12882.4,2^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12759.6,0^{++})\gamma} = 0$							$\mu_{T_{cb\bar{e}\bar{b}}(12882.4,2^{++})\rightarrow T_{cb\bar{e}\bar{b}}(12759.3,0^{++})\gamma} = \langle \Psi_{tot}^{2^{++}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0$				

TABLE XVII. The masses, binding energies, variational parameters, the internal contribution, total wave functions, magnetic moments, transition magnetic moments, radiative decay widths, rearrangement strong width ratios, and the relative lengths between quarks for the the $J^{PC} = 1^{+-}, 1^{++} c\bar{c}\bar{b}$ states and their lowest meson-meson thresholds. The notation it the same as that in Table VII.

$c\bar{c}\bar{b}$		The contribution from each term			Relative Length (fm)		Overall	Present Work		CMI Model	
$J^{PC} = 1^{+-}$	Value	$\Upsilon\eta_c$	Difference	(i, j)	Value	$\Upsilon\eta_c$		Contribution	Value	Ref. [110]	Ref. [123]
Mass/ B_T	12796.9	12467.4	329.5	(1,2)	0.331		m_c	1918.0	$\frac{-m_c}{4(m_c+m_b)}m_{cb}$		
Variational Parameters (fm ⁻²)	C_{11}	11.9	15.0	(1,3)	0.289	0.290(η_c)	$\frac{m_b}{m_c+m_b}\frac{p_{x1}^2}{2m_1^2}$	180.6	-400.5		
	C_{22}	11.9	49.7	(2,3)	0.289		$\frac{m_b}{2m_c+2m_b}\frac{p_{x3}^2}{2m_3^2}$	135.3	$\frac{5m_c}{8(m_c+m_b)}m_{c\bar{b}}$		
	C_{33}	22.9	-	(1,4)	0.289		$\frac{1}{2}V^C(12)$	17.8	986.7		
Quark Mass	14522.0	14522.0	0.0	(2,4)	0.289	0.160(Υ)	$\frac{1}{2}[V^C(13) + V^C(14)]$	-158.1	$\frac{5m_c}{8(m_c+m_c)}m_{c\bar{c}}$	m_c	
Confinement Potential	-2527.1	-3000.0	472.9	(3,4)	0.331		$-\frac{1}{2}D$	-491.5	958.9	1724.8	
Kinetic Energy	858.5	1001.2	-142.7	(1,2)-(3,4):	0.168 fm		Subtotal	1602.1	1545.1	1724.8	
CS Interaction	-41.4	-55.8	14.4	(1,3)	-158.1	-237.2(η_c)	$\frac{m_b}{m_c+m_b}\frac{p_{x1}^2}{2m_1^2}$	5343.0	$\frac{-m_b}{4(m_c+m_b)}m_{cb}$		
V^C	(1,2)	35.6		(2,3)	-158.1		$\frac{m_c}{2m_c+2m_b}\frac{p_{x3}^2}{2m_3^2}$	64.8	-1203.8		
	(1,4)	-158.1		(2,4)	-158.1	-796.7(Υ)	$\frac{1}{2}V^C(12)$	48.6	$\frac{5m_b}{8(m_c+m_b)}m_{c\bar{b}}$		
	Subtotal	-561.1	-1034.0	472.9	(3,4)	35.6	$\frac{1}{2}[V^C(23) + V^C(24)]$	17.8	2965.5	m_b	
Total Contribution	256.0	-88.6	344.6	(3,4)	35.6		$-\frac{1}{2}D$	-491.5	2951.6	5052.9	
Total Wave function:							CS Interaction	Subtotal	4824.7	4713.3	5052.9
$\Psi_{tot} = 0.877 F\rangle R^s\rangle[\phi_1\chi_2\rangle - 0.064 F\rangle R^s\rangle[\phi_2\chi_2\rangle + 0.320 F\rangle R^s\rangle[\phi_1\chi_3\rangle + 0.320 F\rangle R^s\rangle[\phi_1\chi_4\rangle + 0.105 F\rangle R^s\rangle[\phi_2\chi_3\rangle + 0.105 F\rangle R^s\rangle[\phi_2\chi_4\rangle + 0.211 F\rangle R^s\rangle[\psi_1\zeta_2\rangle + 0.854 F\rangle R^s\rangle[\psi_1\zeta_3\rangle + 0.223 F\rangle R^s\rangle[\psi_2\zeta_2\rangle + 0.420 F\rangle R^s\rangle[\psi_2\zeta_3\rangle - 0.481 F\rangle R^s\rangle[\psi'_1\zeta'_2\rangle - 0.481 F\rangle R^s\rangle[\psi'_1\zeta'_3\rangle + 0.283 F\rangle R^s\rangle[\psi'_1\zeta'_4\rangle - 0.395 F\rangle R^s\rangle[\psi'_2\zeta'_2\rangle - 0.395 F\rangle R^s\rangle[\psi'_2\zeta'_3\rangle + 0.382 F\rangle R^s\rangle[\psi'_2\zeta'_4\rangle]$								$-\frac{1}{4}[V^{SS}(12) + V^{SS}(34)]$	-5.0	$-\frac{8}{3}v_{cb} - \frac{10}{3}v_{c\bar{c}}$	$-\frac{8}{3}C_{cb} - \frac{10}{3}C_{c\bar{c}}$
								$-\frac{5}{8}V^{SS}(13)$	-16.0	-5.2-17.7	-8.8-17.7
								$-\frac{5}{8}V^{SS}(24)$	-4.8	$-\frac{10}{3}v_{b\bar{b}} - \frac{20}{3}v_{b\bar{c}}$	$-\frac{10}{3}C_{b\bar{b}} - \frac{20}{3}C_{b\bar{c}}$
								$-\frac{5}{8}[V^{SS}(14) + V^{SS}(23)]$	-15.6	-9.6-19.7	-9.7-22.0
							Subtotal	-41.4	-52.2	-58.1	
The rearrangement strong decay channel:							Matrix nondiagonal element	-15.3	-54.7	-19.6	
							Total contribution	12796.9	12409.9	13478.0	
$\Gamma_{T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-}) \rightarrow \eta_b J/\psi} : \Gamma_{T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-}) \rightarrow \Upsilon\eta_c} = 1 : 18.4$							The magnetic moments: $T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-}) = \langle \Psi_{tot}^{1+-} \hat{\mu}^z \Psi_{tot}^{1+-} \rangle = 0$				
$\Gamma_{T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = 33.1$ keV							The transition magnetic moments: $\mu_{T_{cb\bar{c}\bar{b}}(12882.4, 2^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = 0.345\mu_N$				
$\Gamma_{T_{cb\bar{c}\bar{b}}(12856.6, 1^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = 0.1$ keV							$\mu_{T_{cb\bar{c}\bar{b}}(12856.6, 1^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = \langle \Psi_{tot}^{1++} \hat{\mu}^z \Psi_{tot}^{1+-} \rangle = 0.113(\mu_c + \mu_b) = 0.036\mu_N$				
$\Gamma_{T_{cb\bar{c}\bar{b}}(12882.4, 2^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = 4.9$ keV							$\mu_{T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = \langle \Psi_{tot}^{1+-} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0.226 \times 2\sqrt{\frac{1}{6}}(\mu_c - \mu_b) = 0.081\mu_N$				
$J^{PC} = 1^{++}$		Value	$\Upsilon J/\psi$	Difference	Relative Lengths (fm)		Overall	Contribution	Value	Ref. [110]	Ref. [123]
Mass/ B_T	12856.6	12561.1	295.5	(i, j)	Value	$\Upsilon J/\psi$		m_c	1918.0	$\frac{-m_c}{4(m_c+m_b)}m_{cb}$	
Variational Parameters (fm ⁻²)	C_{11}	11.4	12.5	(1,2)	0.333		$\frac{m_b}{m_c+m_b}\frac{p_{x1}^2}{2m_1^2}$	174.0	-400.5		
	C_{22}	11.4	49.7	(1,3)	0.291	0.318(J/ψ)	$\frac{m_b}{2m_c+2m_b}\frac{p_{x3}^2}{2m_3^2}$	127.2	$\frac{5m_c}{8(m_c+m_b)}m_{c\bar{b}}$		
	C_{33}	21.5	-	(2,3)	0.291		$\frac{1}{2}V^C(12)$	16.0	986.7		
Quark Mass	14522.0	14522.0	0.0	(2,4)	0.291	0.160(Υ)	$\frac{1}{2}[V^C(13) + V^C(14)]$	-146.5	$\frac{5m_c}{8(m_c+m_c)}m_{c\bar{c}}$	m_c	
Confinement Potential	-2488.0	-2926.9	438.9	(1,4)	0.291		$-\frac{1}{2}D$	-491.5	958.9	1724.8	
Kinetic Energy	818.6	925.1	-106.5	(2,4)	0.291	0.160(Υ)	Subtotal	1597.2	1545.1	1724.8	
CS Interaction	10.0	41.0	-31.0	(3,4)	0.333		$\frac{m_b}{m_c+m_b}\frac{p_{x1}^2}{2m_1^2}$	5343.0	$\frac{-m_b}{4(m_c+m_b)}m_{cb}$		
V^C	(1,2)	32.0		(1,2)-(3,4):	0.172 fm		$\frac{m_c}{2m_c+2m_b}\frac{p_{x3}^2}{2m_3^2}$	62.5	-1203.8		
	(2,3)	-146.5		(2,4)	-146.5	-796.7(Υ)	$\frac{1}{2}V^C(12)$	45.7	$\frac{5m_b}{8(m_c+m_b)}m_{c\bar{b}}$		
	(1,4)	-146.5		(3,4)	32.0		$\frac{1}{2}[V^C(23) + V^C(24)]$	16.0	2965.5	m_b	
Subtotal	-522.0	-960.9	438.9	(3,4)	32.0		$-\frac{1}{2}D$	-491.5	2951.6	5052.9	
Total Contribution	306.6	5.1	301.5	(1,3)	-146.5	-164.2(J/ψ)	Subtotal	4829.2	4713.3	5052.9	
Total Wave function:							CS Interaction	$\frac{1}{4}[V^{SS}(12) + V^{SS}(34)]$	4.9	$\frac{8}{3}v_{cb} + \frac{10}{3}v_{c\bar{c}}$	$\frac{8}{3}C_{cb} + \frac{10}{3}C_{c\bar{c}}$
$\Psi_{tot} = 0.693 F\rangle R^s\rangle[\phi_2\chi_3\rangle - 0.693 F\rangle R^s\rangle[\phi_2\chi_4\rangle + 0.139 F\rangle R^s\rangle[\phi_1\chi_3\rangle - 0.139 F\rangle R^s\rangle[\phi_1\chi_4\rangle + 0.686 F\rangle R^s\rangle[\psi_1\zeta_4\rangle + 0.727 F\rangle R^s\rangle[\psi_2\zeta_4\rangle - 0.480 F\rangle R^s\rangle[\psi'_1\zeta'_2\rangle - 0.480 F\rangle R^s\rangle[\psi'_1\zeta'_3\rangle - 0.395 F\rangle R^s\rangle[\psi'_2\zeta'_2\rangle - 0.395 F\rangle R^s\rangle[\psi'_2\zeta'_3\rangle]$								$\frac{5}{8}V^{SS}(13)$	15.5	5.2+17.7	8.8+17.7
								$\frac{5}{8}V^{SS}(24)$	4.6	$\frac{10}{3}v_{b\bar{b}} - \frac{20}{3}v_{b\bar{c}}$	$\frac{10}{3}C_{b\bar{b}} - \frac{20}{3}C_{b\bar{c}}$
								$-\frac{5}{8}[V^{SS}(14) + V^{SS}(23)]$	-15.0	9.6-19.7	9.7-22.0
								Subtotal	10.0	12.9	14.1
The magnetic moments: $T_{cb\bar{c}\bar{b}}(12856.6, 1^{++}) = \langle \Psi_{tot}^{1++} \hat{\mu}^z \Psi_{tot}^{1++} \rangle = 0$							Matrix nondiagonal element	-6.3	-6.1	-59.5	
The radiative decay widths:							Total contribution	12856.6	12523.6	13510.0	
$\Gamma_{T_{cb\bar{c}\bar{b}}(12856.6, 1^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = 0$ keV							The transition magnetic moments: $\mu_{T_{cb\bar{c}\bar{b}}(12882.4, 2^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12856.6, 1^{++})\gamma} = \langle \Psi_{tot}^{2^{++}} \hat{\mu}^z \Psi_{tot}^{1^{++}} \rangle = 0$				
$\Gamma_{T_{cb\bar{c}\bar{b}}(12856.6, 1^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = 0.1$ keV							$\mu_{T_{cb\bar{c}\bar{b}}(12856.6, 1^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = \langle \Psi_{tot}^{1^{++}} \hat{\mu}^z \Psi_{tot}^{0^{++}} \rangle = 0$				
$\Gamma_{T_{cb\bar{c}\bar{b}}(12882.4, 2^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12856.6, 1^{++})\gamma} = 0$ keV							$\mu_{T_{cb\bar{c}\bar{b}}(12856.6, 1^{++}) \rightarrow T_{cb\bar{c}\bar{b}}(12796.9, 1^{+-})\gamma} = \langle \Psi_{tot}^{1^{++}} \hat{\mu}^z \Psi_{tot}^{1^{+-}} \rangle = 0.113(\mu_c + \mu_b) = 0.036\mu_N$				
The radiative decay widths: $J/\psi\Upsilon$							$\Gamma_{T_{cb\bar{c}\bar{b}}(12856.6, 1^{++}) \rightarrow B_c^* \bar{B}_c} : \Gamma_{T_{cb\bar{c}\bar{b}}(12856.6, 1^{++}) \rightarrow B_c \bar{B}_c^*} = 1 : 1$				

- [11] A. Esposito, A. L. Guerrieri, F. Piccinini, A. Pilloni and A. D. Polosa, Four-Quark Hadrons: an Updated Review, *Int. J. Mod. Phys. A* **30** (2015) 1530002.
- [12] H. X. Chen, W. Chen, X. Liu and S. L. Zhu, The hidden-charm pentaquark and tetraquark states, *Phys. Rept.* **639** (2016) 1.
- [13] R. Chen, X. Liu and S. L. Zhu, Hidden-charm molecular pentaquarks and their charm-strange partners, *Nucl. Phys. A* **954** (2016) 406.
- [14] A. Hosaka, T. Iijima, K. Miyabayashi, Y. Sakai and S. Yasui, Exotic hadrons with heavy flavors: X, Y, Z, and related states, *PTEP* **2016** (2016) no.6, 062C01.
- [15] J. M. Richard, Exotic hadrons: review and perspectives, *Few Body Syst.* **57** (2016) no.12, 1185.
- [16] R. F. Lebed, R. E. Mitchell and E. S. Swanson, Heavy-Quark QCD Exotica, *Prog. Part. Nucl. Phys.* **93**, 143-194 (2017).
- [17] A. Esposito, A. Pilloni and A. D. Polosa, Multiquark Resonances, *Phys. Rept.* **668**, 1-97 (2017).
- [18] A. Ali, J. S. Lange and S. Stone, Exotics: Heavy Pentaquarks and Tetraquarks, *Prog. Part. Nucl. Phys.* **97**, 123-198 (2017).
- [19] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C. P. Shen, C. E. Thomas, A. Vairo and C. Z. Yuan, The *XYZ* states: experimental and theoretical status and perspectives, *Phys. Rept.* **873**, 1-154 (2020).
- [20] Y. R. Liu, H. X. Chen, W. Chen, X. Liu and S. L. Zhu, Pentaquark and Tetraquark states, *Prog. Part. Nucl. Phys.* **107** (2019), 237-320.
- [21] H. X. Chen, W. Chen, X. Liu, Y. R. Liu and S. L. Zhu, An updated review of the new hadron states, *Rept. Prog. Phys.* **86** (2023) no.2, 026201
- [22] R. Aaij *et al.* [LHCb Collaboration], Observation of $J/\psi p$ Resonances Consistent with Pentaquark States in $\Lambda_b^0 \rightarrow J/\psi K^- p$ Decays, *Phys. Rev. Lett.* **115**, 072001 (2015).
- [23] R. Aaij *et al.* [LHCb Collaboration], Model-independent evidence for $J/\psi p$ contributions to $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays, *Phys. Rev. Lett.* **117**, no. 8, 082002 (2016).
- [24] R. Aaij *et al.* [LHCb Collaboration], Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$, *Phys. Rev. Lett.* **122**, no. 22, 222001 (2019).
- [25] B. Aubert *et al.* [BaBar Collaboration], Observation of a narrow meson decaying to $D_s^+ \pi^0$ at a mass of 2.32 GeV/ c^2 , *Phys. Rev. Lett.* **90**, 242001 (2003).
- [26] S. Godfrey and N. Isgur, Mesons in a Relativized Quark Model with Chromodynamics, *Phys. Rev. D* **32**, 189 (1985).
- [27] T. Barnes, F. E. Close and H. J. Lipkin, Implications of a DK molecule at 2.32-GeV, *Phys. Rev. D* **68** (2003), 054006.
- [28] H. Y. Cheng and W. S. Hou, B decays as spectroscopy for charmed four quark states, *Phys. Lett. B* **566** (2003), 193-200.
- [29] A. P. Szczepaniak, Description of the $D^*(s)(2320)$ resonance as the D pi atom, *Phys. Lett. B* **567** (2003), 23-26.
- [30] D. Besson *et al.* [CLEO Collaboration], Observation of a narrow resonance of mass 2.46 GeV/ c^2 decaying to $D_s^{*+} \pi^0$ and confirmation of the $D_{sJ}^*(2317)$ state, *Phys. Rev. D* **68**, 032002 (2003).
- [31] V. Dmitrasinovic, D_s^{*+} (2317) and D_s^{*+} (2460): Tetraquarks bound by the t Hooft instanton-induced interaction?, *Phys. Rev. D* **70** (2004), 096011.
- [32] Y. Q. Chen and X. Q. Li, A Comprehensive four-quark interpretation of $D(s)(2317)$, $D(s)(2457)$ and $D(s)(2632)$, *Phys. Rev. Lett.* **93** (2004), 232001.
- [33] Y. R. Liu, S. L. Zhu, Y. B. Dai and C. Liu, $D+(sJ)(2632)$: An Excellent candidate of tetraquarks, *Phys. Rev. D* **70** (2004), 094009.
- [34] A. Faessler, T. Gutsche, V. E. Lyubovitskij and Y. L. Ma, $D^* K$ molecular structure of the $D(s1)(2460)$ meson, *Phys. Rev. D* **76** (2007), 114008.
- [35] D. Ebert, R. N. Faustov and V. O. Galkin, Masses of tetraquarks with open charm and bottom, *Phys. Lett. B* **696** (2011), 241-245.
- [36] c. J. Xiao, D. Y. Chen and Y. L. Ma, Radiative and pionic transitions from the $D_{s1}(2460)$ to the $D_{s0}^*(2317)$, *Phys. Rev. D* **93** (2016) no.9, 094011.
- [37] S. Y. Kong, J. T. Zhu, D. Song and J. He, Heavy-strange meson molecules and possible candidates $Ds0^*(2317)$, $Ds1(2460)$, and $X0(2900)$, *Phys. Rev. D* **104** (2021) no.9, 094012.
- [38] H. L. Fu, H. W. Grieffhammer, F. K. Guo, C. Hanhart and U. G. Meißner, Update on strong and radiative decays of the $D_{s0}^*(2317)$ and $D_{s1}(2460)$ and their bottom cousins, *Eur. Phys. J. A* **58** (2022) no.4, 70.
- [39] R. Aaij *et al.* [LHCb], A model-independent study of resonant structure in $B^+ \rightarrow D^+ D^- K^+$ decays, *Phys. Rev. Lett.* **125**, 242001 (2020).
- [40] R. Aaij *et al.* [LHCb], Amplitude analysis of the $B^+ \rightarrow D^+ D^- K^+$ decay, *Phys. Rev. D* **102** (2020), 112003.
- [41] X. H. Liu, M. J. Yan, H. W. Ke, G. Li and J. J. Xie, Triangle singularity as the origin of $X_0(2900)$ and $X_1(2900)$ observed in $B^+ \rightarrow D^+ D^- K^+$, *Eur. Phys. J. C* **80** (2020) no.12, 1178.
- [42] M. Karliner and J. L. Rosner, First exotic hadron with open heavy flavor: $cs\bar{u}\bar{d}$ tetraquark, *Phys. Rev. D* **102** (2020) no.9, 094016.
- [43] Z. G. Wang, Analysis of the $X_0(2900)$ as the scalar tetraquark state via the QCD sum rules, *Int. J. Mod. Phys. A* **35** (2020) no.30, 2050187.
- [44] Y. Huang, J. X. Lu, J. J. Xie and L. S. Geng, Strong decays of $\bar{D}^* K^*$ molecules and the newly observed $X_{0,1}$ states, *Eur. Phys. J. C* **80** (2020) no.10, 973.
- [45] T. J. Burns and E. S. Swanson, Kinematical cusp and resonance interpretations of the $X(2900)$, *Phys. Lett. B* **813** (2021), 136057.
- [46] M. W. Hu, X. Y. Lao, P. Ling and Q. Wang, $X_0(2900)$ and its heavy quark spin partners in molecular picture, *Chin. Phys. C* **45** (2021) no.2, 021003.
- [47] G. J. Wang, L. Meng, L. Y. Xiao, M. Oka and S. L. Zhu, Mass spectrum and strong decays of tetraquark $\bar{c}\bar{s}q\bar{q}$ states, *Eur. Phys. J. C* **81** (2021) no.2, 188.
- [48] Y. Xue, X. Jin, H. Huang and J. Ping, Tetraquarks with open charm flavor, *Phys. Rev. D* **103** (2021) no.5, 054010.
- [49] S. S. Agaev, K. Azizi and H. Sundu, Vector resonance $X_1(2900)$ and its structure, *Nucl. Phys. A* **1011** (2021), 122202.
- [50] U. Özdem and K. Azizi, Magnetic moment of the $X_1(2900)$ state in the diquark-antidiquark picture, [arXiv:2202.11466 [hep-ph]].
- [51] A. Del Fabbro, D. Janc, M. Rosina and D. Treleani, Production and detection of doubly charmed tetraquarks, *Phys. Rev. D* **71** (2005), 014008.

- [52] D. Ebert, R. N. Faustov, V. O. Galkin and W. Lucha, Masses of tetraquarks with two heavy quarks in the relativistic quark model, *Phys. Rev. D* **76** (2007), 114015.
- [53] T. Hyodo, Y. R. Liu, M. Oka, K. Sudoh and S. Yasui, Production of doubly charmed tetraquarks with exotic color configurations in electron-positron collisions, *Phys. Lett. B* **721** (2013), 56-60.
- [54] Y. Ikeda, B. Charron, S. Aoki, T. Doi, T. Hatsuda, T. Inoue, N. Ishii, K. Murano, H. Nemura and K. Sasaki, Charmed tetraquarks T_{cc} and T_{cs} from dynamical lattice QCD simulations, *Phys. Lett. B* **729** (2014), 85-90.
- [55] A. Esposito, M. Papinutto, A. Pilloni, A. D. Polosa and N. Tantalo, Doubly charmed tetraquarks in B_c and Ξ_{bc} decays, *Phys. Rev. D* **88** (2013) no.5, 054029.
- [56] R. Aaij *et al.* [LHCb Collaboration], Observation of the doubly charmed baryon Ξ_{cc}^{++} , *Phys. Rev. Lett.* **119**, no. 11, 112001 (2017).
- [57] S. Q. Luo, K. Chen, X. Liu, Y. R. Liu and S. L. Zhu, Exotic tetraquark states with the $qq\bar{Q}\bar{Q}$ configuration, *Eur. Phys. J. C* **77**, no. 10, 709 (2017).
- [58] M. Karliner and J. L. Rosner, Discovery of doubly-charmed Ξ_{cc} baryon implies a stable ($bb\bar{u}\bar{d}$) tetraquark, *Phys. Rev. Lett.* **119**, no. 20, 202001 (2017).
- [59] E. J. Eichten and C. Quigg, Heavy-quark symmetry implies stable heavy tetraquark mesons $Q_i Q_j \bar{q}_k \bar{q}_l$, *Phys. Rev. Lett.* **119**, no. 20, 202002 (2017).
- [60] A. Ali, Q. Qin and W. Wang, Discovery potential of stable and near-threshold doubly heavy tetraquarks at the LHC, *Phys. Lett. B* **785**, 605-609 (2018).
- [61] W. Park, S. Noh and S. H. Lee, Masses of the doubly heavy tetraquarks in a constituent quark model, *Acta Phys. Polon. B* **50**, 1151-1157 (2019).
- [62] T. Guo, J. Li, J. Zhao and L. He, Mass spectra of doubly heavy tetraquarks in an improved chromomagnetic interaction model, *Phys. Rev. D* **105** (2022) no.1, 014021.
- [63] Q. F. Lü, D. Y. Chen and Y. B. Dong, Masses of doubly heavy tetraquarks $T_{QQ'}$ in a relativized quark model, *Phys. Rev. D* **102** (2020) no.3, 034012.
- [64] W. Park and S. H. Lee, Color spin wave functions of heavy tetraquark states, *Nucl. Phys. A* **925** (2014), 161-184.
- [65] S. Noh, W. Park and S. H. Lee, The Doubly-heavy Tetraquarks ($qq'\bar{Q}\bar{Q}'$) in a Constituent Quark Model with a Complete Set of Harmonic Oscillator Bases, *Phys. Rev. D* **103** (2021), 114009.
- [66] C. Deng, H. Chen and J. Ping, Systematical investigation on the stability of doubly heavy tetraquark states, *Eur. Phys. J. A* **56** (2020) no.1, 9.
- [67] R. Aaij *et al.* [LHCb], Observation of an exotic narrow doubly charmed tetraquark, [arXiv:2109.01038].
- [68] Q. F. Lü, D. Y. Chen, Y. B. Dong and E. Santopinto, Triply-heavy tetraquarks in an extended relativized quark model, *Phys. Rev. D* **104** (2021) no.5, 054026.
- [69] K. Chen, X. Liu, J. Wu, Y. R. Liu and S. L. Zhu, Triply heavy tetraquark states with the $QQ\bar{Q}\bar{q}$ configuration, *Eur. Phys. J. A* **53**, no. 1, 5 (2017).
- [70] Y. Xing, Weak decays of triply heavy tetraquarks $b\bar{c}b\bar{q}$, *Eur. Phys. J. C* **80** (2020) no.1, 57.
- [71] J. F. Jiang, W. Chen and S. L. Zhu, Triply heavy $QQ\bar{Q}\bar{q}$ tetraquark states, *Phys. Rev. D* **96** (2017) no.9, 094022.
- [72] X. Z. Weng, W. Z. Deng and S. L. Zhu, Triply heavy tetraquark states, *Phys. Rev. D* **105** (2022) no.3, 034026.
- [73] K. T. Chao, The $(cc) - (\bar{c}\bar{c})$ (Diquark - Anti-Diquark) States in e^+e^- Annihilation, *Z. Phys. C* **7** (1981), 317.
- [74] J. P. Ader, J. M. Richard and P. Taxil, DO NARROW HEAVY MULTI - QUARK STATES EXIST?, *Phys. Rev. D* **25** (1982), 2370.
- [75] A. M. Badalian, B. L. Ioffe and A. V. Smilga, FOUR QUARK STATES IN THE HEAVY QUARK SYSTEM, *Nucl. Phys. B* **281** (1987), 85
- [76] L. Heller and J. A. Tjon, On Bound States of Heavy $Q^2\bar{Q}^2$ Systems, *Phys. Rev. D* **32**, 755 (1985).
- [77] S. Zouzou, B. Silvestre-Brac, C. Gignoux and J. M. Richard, FOUR QUARK BOUND STATES, *Z. Phys. C* **30** (1986), 457
- [78] R. J. Lloyd and J. P. Vary, All charm tetraquarks, *Phys. Rev. D* **70**, 014009 (2004).
- [79] M. Karliner, S. Nussinov and J. L. Rosner, $QQ\bar{Q}\bar{Q}$ states: masses, production, and decays, *Phys. Rev. D* **95**, no.3, 034011 (2017).
- [80] M. N. Anwar, J. Ferretti, F. K. Guo, E. Santopinto and B. S. Zou, Spectroscopy and decays of the fully-heavy tetraquarks, *Eur. Phys. J. C* **78**, no.8, 647 (2018).
- [81] Y. Bai, S. Lu and J. Osborne, Beauty-full Tetraquarks, *Phys. Lett. B* **798**, 134930 (2019).
- [82] V. R. Debastiani and F. S. Navarra, Spectroscopy of the All-Charmed Tetraquark, *PoS Hadron2017*, 238 (2018).
- [83] W. Chen, H. X. Chen, X. Liu, T. G. Steele and S. L. Zhu, Hunting for exotic doubly hidden-charm/bottom tetraquark states, *Phys. Lett. B* **773** (2017), 247-251.
- [84] V. Khachatryan *et al.* [CMS], Observation of $\Upsilon(1S)$ pair production in proton-proton collisions at $\sqrt{s} = 8$ TeV, *JHEP* **05** (2017), 013.
- [85] A. M. Sirunyan *et al.* [CMS], Measurement of the $\Upsilon(1S)$ pair production cross section and search for resonances decaying to $\Upsilon(1S)\mu^+\mu^-$ in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Lett. B* **808** (2020), 135578.
- [86] R. Aaij *et al.* [LHCb], Search for beautiful tetraquarks in the $\Upsilon(1S)\mu^+\mu^-$ invariant-mass spectrum, *JHEP* **10** (2018), 086.
- [87] R. Aaij *et al.* [LHCb], Observation of structure in the J/ψ -pair mass spectrum, *Sci. Bull.* **65** (2020) no.23, 1983-1993.
- [88] A. Hayrapetyan *et al.* [CMS], Observation of new structure in the $J/\psi J/\psi$ mass spectrum in proton-proton collisions at $\sqrt{s} = 13$ TeV, [arXiv:2306.07164 [hep-ex]].
- [89] J. Zhang *et al.* [CMS], Recent CMS results on exotic resonances, *PoS ICHEP2022*, 775 [arXiv:2212.00504 [hep-ex]].
- [90] Y. Xu [ATLAS], ATLAS Results on Exotic Hadronic Resonances, *Acta Phys. Polon. Supp.* **16** (2023) no.3, 21 [arXiv:2209.12173 [hep-ex]].
- [91] J. Zhao, S. Shi and P. Zhuang, Fully-heavy tetraquarks in a strongly interacting medium, *Phys. Rev. D* **102** (2020) no.11, 114001.
- [92] H. X. Chen, W. Chen, X. Liu and S. L. Zhu, Strong decays of fully-charm tetraquarks into di-charmonia, *Sci. Bull.* **65** (2020), 1994-2000.
- [93] M. S. Liu, F. X. Liu, X. H. Zhong and Q. Zhao, Full-heavy tetraquark states and their evidences in the LHCb di- J/ψ spectrum, [arXiv:2006.11952 [hep-ph]].
- [94] Q. F. Lü, D. Y. Chen and Y. B. Dong, Masses of fully heavy tetraquarks $QQ\bar{Q}\bar{Q}$ in an extended relativized quark model, *Eur. Phys. J. C* **80** (2020) no.9, 871.

- [95] Z. Zhao, K. Xu, A. Kaewsnod, X. Liu, A. Limphirat and Y. Yan, Study of charmoniumlike and fully-charm tetraquark spectroscopy, *Phys. Rev. D* **103** (2021) no.11, 116027.
- [96] M. A. Bedolla, J. Ferretti, C. D. Roberts and E. Santopinto, Spectrum of fully-heavy tetraquarks from a diquark+antidiquark perspective, *Eur. Phys. J. C* **80** (2020) no.11, 1004
- [97] F. X. Liu, M. S. Liu, X. H. Zhong and Q. Zhao, Higher mass spectra of the fully-charmed and fully-bottom tetraquarks, *Phys. Rev. D* **104** (2021) no.11, 116029.
- [98] H. W. Ke, X. Han, X. H. Liu and Y. L. Shi, Tetraquark state $X(6900)$ and the interaction between diquark and antidiquark, *Eur. Phys. J. C* **81** (2021) no.5, 427.
- [99] M. Karliner and J. L. Rosner, Interpretation of structure in the di- J/ψ spectrum, *Phys. Rev. D* **102** (2020) no.11, 114039.
- [100] R. N. Faustov, V. O. Galkin and E. M. Savchenko, Masses of the $QQ\bar{Q}\bar{Q}$ tetraquarks in the relativistic diquark-antidiquark picture, *Phys. Rev. D* **102** (2020), 114030.
- [101] B. D. Wan and C. F. Qiao, Gluonic tetracharm configuration of $X(6900)$, *Phys. Lett. B* **817** (2021), 136339.
- [102] C. Gong, M. C. Du, Q. Zhao, X. H. Zhong and B. Zhou, Nature of $X(6900)$ and its production mechanism at LHCb, *Phys. Lett. B* **824** (2022), 136794.
- [103] J. W. Zhu, X. D. Guo, R. Y. Zhang, W. G. Ma and X. Q. Li, A possible interpretation for $X(6900)$ observed in four-muon final state by LHCb – A light Higgs-like boson?, [arXiv:2011.07799 [hep-ph]].
- [104] J. Z. Wang, D. Y. Chen, X. Liu and T. Matsuki, Producing fully charm structures in the J/ψ -pair invariant mass spectrum, *Phys. Rev. D* **103** (2021) no.7, 071503.
- [105] J. Z. Wang and X. Liu, Improved understanding of the peaking phenomenon existing in the new di- J/ψ invariant mass spectrum from the CMS Collaboration, *Phys. Rev. D* **106** (2022) no.5, 054015.
- [106] N. Santowsky and C. S. Fischer, Four-quark states with charm quarks in a two-body Bethe–Salpeter approach, *Eur. Phys. J. C* **82** (2022) no.4, 313.
- [107] Z. Zhuang, Y. Zhang, Y. Ma and Q. Wang, Lineshape of the compact fully heavy tetraquark, *Phys. Rev. D* **105** (2022) no.5, 054026.
- [108] E. Lötstedt, L. Wang, R. Yoshida, Y. Zhang and K. Yamanouchi, Error-mitigated quantum computing of Heisenberg spin chain dynamics, *Phys. Scripta* **98** (2023) no.3, 035111.
- [109] V. R. Debastiani and F. S. Navarra, A non-relativistic model for the $[cc][\bar{c}\bar{c}]$ tetraquark, *Chin. Phys. C* **43** (2019) no.1, 013105.
- [110] X. Z. Weng, X. L. Chen, W. Z. Deng and S. L. Zhu, Systematics of fully heavy tetraquarks, *Phys. Rev. D* **103** (2021) no.3, 034001.
- [111] G. J. Wang, L. Meng and S. L. Zhu, Spectrum of the fully-heavy tetraquark state $QQ\bar{Q}'\bar{Q}'$, *Phys. Rev. D* **100** (2019) no.9, 096013.
- [112] C. Hughes, E. Eichten and C. T. H. Davies, Searching for beauty-fully bound tetraquarks using lattice nonrelativistic QCD, *Phys. Rev. D* **97** (2018) no.5, 054505.
- [113] J. M. Richard, A. Valcarce and J. Vijande, Few-body quark dynamics for doubly heavy baryons and tetraquarks, *Phys. Rev. C* **97** (2018) no.3, 035211.
- [114] X. Jin, Y. Xue, H. Huang and J. Ping, Full-heavy tetraquarks in constituent quark models, *Eur. Phys. J. C* **80**, no.11, 1083 (2020).
- [115] H. T. An, S. Q. Luo, Z. W. Liu and X. Liu, Fully heavy pentaquark states in constituent quark model, *Phys. Rev. D* **105** (2022) no.7, 7.
- [116] W. Park, A. Park and S. H. Lee, Dibaryons in a constituent quark model, *Phys. Rev. D* **92** (2015) no.1, 014037.
- [117] W. Park, A. Park and S. H. Lee, Dibaryons with two strange quarks and total spin zero in a constituent quark model, *Phys. Rev. D* **93** (2016) no.7, 074007.
- [118] A. Park, W. Park and S. H. Lee, Dibaryons with two strange quarks and one heavy flavor in a constituent quark model, *Phys. Rev. D* **94** (2016) no.5, 054027.
- [119] W. Park, A. Park, S. Cho and S. H. Lee, $P_c(4380)$ in a constituent quark model, *Phys. Rev. D* **95** (2017) no.5, 054027.
- [120] W. Park, S. Cho and S. H. Lee, Where is the stable pentaquark?, *Phys. Rev. D* **99** (2019) no.9, 094023.
- [121] H. Y. Zhou, F. L. Wang, Z. W. Liu and X. Liu, Probing the electromagnetic properties of the $\Sigma_c^{(*)}D^{(*)}$ -type doubly charmed molecular pentaquarks, *Phys. Rev. D* **106** (2022) no.3, 034034.
- [122] S. Kumar, R. Dhir and R. C. Verma, Magnetic moments of charm baryons using effective mass and screened charge of quarks, *J. Phys. G* **31** (2005) no.2, 141-147
- [123] J. Wu, Y. R. Liu, K. Chen, X. Liu and S. L. Zhu, Heavy-flavored tetraquark states with the $QQ\bar{Q}\bar{Q}$ configuration, *Phys. Rev. D* **97** (2018) no.9, 094015.
- [124] W. X. Zhang, H. Xu and D. Jia, Masses and magnetic moments of hadrons with one and two open heavy quarks: Heavy baryons and tetraquarks, *Phys. Rev. D* **104** (2021) no.11, 114011.
- [125] G. J. Wang, R. Chen, L. Ma, X. Liu and S. L. Zhu, Magnetic moments of the hidden-charm pentaquark states, *Phys. Rev. D* **94** (2016) no.9, 094018.
- [126] F. L. Wang, H. Y. Zhou, Z. W. Liu and X. Liu, What can we learn from the electromagnetic properties of hidden-charm molecular pentaquarks with single strangeness?, *Phys. Rev. D* **106** (2022) no.5, 054020.
- [127] F. L. Wang, H. Y. Zhou, Z. W. Liu and X. Liu, Exploring the electromagnetic properties of the $\Xi_c^{(\prime)*}\bar{D}_s^*$ and $\Omega_c^{(*)}\bar{D}_s^*$ molecular states, [arXiv:2210.02809 [hep-ph]].
- [128] Z. P. Li, The Threshold pion photoproduction of nucleons in the chiral quark model, *Phys. Rev. D* **50** (1994), 5639-5646.
- [129] W. J. Deng, H. Liu, L. C. Gui and X. H. Zhong, Spectrum and electromagnetic transitions of bottomonium, *Phys. Rev. D* **95** (2017) no.7, 074002.
- [130] W. J. Deng, H. Liu, L. C. Gui and X. H. Zhong, Charmonium spectrum and their electromagnetic transitions with higher multipole contributions, *Phys. Rev. D* **95** (2017) no.3, 034026.
- [131] L. Y. Xiao, K. L. Wang, Q. f. Lu, X. H. Zhong and S. L. Zhu, Strong and radiative decays of the doubly charmed baryons, *Phys. Rev. D* **96**, no.9, 094005 (2017).
- [132] Q. F. Lü, K. L. Wang, L. Y. Xiao and X. H. Zhong, Mass spectra and radiative transitions of doubly heavy baryons in a relativized quark model, *Phys. Rev. D* **96**, no.11, 114006 (2017).
- [133] K. L. Wang, Y. X. Yao, X. H. Zhong and Q. Zhao, Strong and radiative decays of the low-lying S - and P -wave singly heavy baryons, *Phys. Rev. D* **96**, no.11, 116016 (2017).

- [134] Y. X. Yao, K. L. Wang and X. H. Zhong, Strong and radiative decays of the low-lying D -wave singly heavy baryons, *Phys. Rev. D* **98**, no.7, 076015 (2018).
- [135] Z. p. Li, H. x. Ye and M. h. Lu, An Unified approach to pseudoscalar meson photoproductions off nucleons in the quark model, *Phys. Rev. C* **56** (1997), 1099-1113.
- [136] S. J. Brodsky and J. R. Primack, The Electromagnetic Interactions of Composite Systems, *Annals Phys.* **52** (1969), 315-365.
- [137] Q. Zhao, J. S. Al-Khalili, Z. P. Li and R. L. Workman, Pion photoproduction on the nucleon in the quark model, *Phys. Rev. C* **65** (2002), 065204.
- [138] H. T. An, K. Chen and X. Liu, Manifestly exotic pentaquarks with a single heavy quark, *Phys. Rev. D* **105** (2022) no.3, 034018.
- [139] T. Guo, J. Li, J. Zhao and L. He, Mass spectra and decays of open-heavy tetraquark states, *Phys. Rev. D* **105** (2022) no.5, 054018.
- [140] X. Z. Weng, W. Z. Deng and S. L. Zhu, Doubly heavy tetraquarks in an extended chromomagnetic model, *Chin. Phys. C* **46** (2022) no.1, 013102.
- [141] C. Deng, H. Chen and J. Ping, Towards the understanding of fully-heavy tetraquark states from various models, *Phys. Rev. D* **103** (2021) no.1, 014001.
- [142] M. S. Liu, Q. F. Lü, X. H. Zhong and Q. Zhao, All-heavy tetraquarks, *Phys. Rev. D* **100** (2019) no.1, 016006.
- [143] J. Zhang, J. B. Wang, G. Li, C. S. An, C. R. Deng and J. J. Xie, Spectrum of the S-wave fully-heavy tetraquark states, [arXiv:2209.13856 [hep-ph]].
- [144] E. Hiyama, Y. Kino and M. Kamimura, Gaussian expansion method for few-body systems, *Prog. Part. Nucl. Phys.* **51** (2003), 223-307.
- [145] Q. N. Wang, Z. Y. Yang and W. Chen, Exotic fully-heavy $QQ\bar{Q}\bar{Q}$ tetraquark states in $\mathbf{8}_{[Q\bar{Q}]} \otimes \mathbf{8}_{[Q\bar{Q}]}$ color configuration, *Phys. Rev. D* **104** (2021) no.11, 114037.
- [146] M. C. Gordillo, F. De Soto and J. Segovia, Diffusion Monte Carlo calculations of fully-heavy multi-quark bound states, *Phys. Rev. D* **102** (2020) no.11, 114007.
- [147] H. Mutuk, Nonrelativistic treatment of fully-heavy tetraquarks as diquark-antidiquark states, *Eur. Phys. J. C* **81** (2021) no.4, 367.
- [148] X. Chen, Fully-heavy tetraquarks: $bb\bar{c}\bar{c}$ and $bc\bar{b}\bar{c}$, *Phys. Rev. D* **100** (2019) no.9, 094009.
- [149] A. V. Berezhnoy, A. V. Luchinsky and A. A. Novoselov, Tetraquarks Composed of 4 Heavy Quarks, *Phys. Rev. D* **86** (2012), 034004.
- [150] Q. Li, C. H. Chang, G. L. Wang and T. Wang, Mass spectra and wave functions of $T_{QQ\bar{Q}\bar{Q}}$ tetraquarks, *Phys. Rev. D* **104** (2021) no.1, 014018.
- [151] H. Mutuk, Spectrum of $cc\bar{b}\bar{b}$, $bc\bar{c}\bar{c}$, and $bc\bar{b}\bar{b}$ tetraquark states in the dynamical diquark model, *Phys. Lett. B* **834** (2022), 137404.
- [152] Z. G. Wang, Analysis of the $QQ\bar{Q}\bar{Q}$ tetraquark states with QCD sum rules, *Eur. Phys. J. C* **77** (2017) no.7, 432.
- [153] Z. G. Wang and Z. Y. Di, Analysis of the vector and axialvector $QQ\bar{Q}\bar{Q}$ tetraquark states with QCD sum rules, *Acta Phys. Polon. B* **50** (2019), 1335.
- [154] X. Chen, Fully-charm tetraquarks: $cc\bar{c}\bar{c}$, [arXiv:2001.06755 [hep-ph]].