

# Sensitivities on dark photon from the Forward Physics Experiments

Kingman Cheung<sup>a,b,c</sup>, C.J. Ouseph<sup>a,b</sup>

<sup>a</sup> *Department of Physics, National Tsing Hua University, Hsinchu 30013, Taiwan*

<sup>b</sup> *Center for Theory and Computation,*

*National Tsing Hua University, Hsinchu 30013, Taiwan*

<sup>c</sup> *Division of Quantum Phases and Devices, School of Physics,*

*Konkuk University, Seoul 143-701, Republic of Korea*

(Dated: August 10, 2022)

## Abstract

Neutrino-electron scattering experiments can explore the potential presence of a light gauge boson  $A'$  which arises from an additional  $U(1)_{B-L}$  group, or a dark photon  $A'$  which arises from a dark sector and has kinetic mixing with the SM hypercharge gauge field. We generically call it a dark photon. In this study, we investigate the effect of the dark photon on neutrino-electron scattering  $\nu e^- \rightarrow \nu e^-$  at the newly proposed forward physics experiments such as FASER $\nu$ , FASER $\nu$ 2, SND@LHC and FLArE(10 tons). We estimate the anticipated sensitivities to the  $U(1)_{B-L}$  gauge coupling in a wide range of the dark photon mass  $M_{A'}$ . We compare the sensitivities of the proposed forward physics experiments with the current limits from TEXONO, GEMMA, BOREXINO, and LSND, as well as CHARM II experiments. We also interpret the anticipated sensitivities in terms of the kinetic mixing parameter  $\epsilon$  in a wide range of dark photon mass  $M_{A'}$ .

## I. INTRODUCTION

Neutrino-charged-lepton scattering and neutrino-nucleon scattering have a long history of establishing the standard model (SM) [1], e.g., the discovery of neutral current in  $\nu N$  scattering in 1973 [2], the measurement of the weak-mixing angle, as well as constraining new physics that violates parity [1]. In the last two decades, neutrino physics mainly concerned with the oscillations of flavors of neutrinos and now the oscillation parameters are reasonably well determined, except for the CP-violation parameter, the mass hierarchy of neutrino masses and  $\theta_{23}$ -octant degeneracy. A number of new experiments are proposed to solve these subtleties, such as DUNE, JUNO, Hyper-K, etc.

On the other hand, neutrino scattering received renewed interests in a number of proposals, including the near detector of DUNE making use of an off-axis neutrino beam, and FASER $\nu$  making use of the neutrinos coming from the interaction point (IP) of the ATLAS experiment. FASER $\nu$  is indeed running and taking data [3, 4]. The most distinct feature of the neutrinos coming off the IP of ATLAS is that the energy range can be as high as TeV, which provides an unprecedented energy scale of studying neutrino-nucleon and neutrino-electron scattering. The FASER $\nu$  experiment can cover a unique energy range that other experiments cannot cover. The ICECUBE focuses on very high-energy neutrinos with energy 10 TeV to 1 PeV. On the other hand, the short- and long-baseline experiments cover mostly around MeV up to a few GeV. There are no precise measurements of neutrino scattering in a few tens of GeV to a few hundreds of GeV region. FASER $\nu$  based on the neutrino flux coming off the LHC opens such a unique window in this energy range. A further proposal is to build the Forward Physics Facilities (FPF) [12] that can house a number of proposed experiments within the same location, including FASER2, Advanced SND, and FORMOSA. Thus, such experiments open an avenue to search for new physics associated with the neutrinos, which are not easily tested at the  $pp$  collisions of the LHC experiments.

In this work, we investigate the potential of detecting a light  $U(1)_{B-L}$  gauge boson or a hidden  $U(1)$  gauge boson (aka. dark photon) at such proposed neutrino-scattering experiments. Generically, a hidden  $U(1)$  does not couple to the SM particles, but it can mix with the SM  $U(1)_Y$  gauge field through the  $B^{\mu\nu}F'_{\mu\nu}$  term. Throughout the work we use the  $U(1)_{B-L}$  gauge boson as illustration and at the end we also translate the results to the dark photon model.

The organization is as follows. In the next section, we describe the  $U(1)_{B-L}$  model and the relation to the hidden  $U(1)$  model (dark photon). In Sec. III, we calculate the neutrino-electron scattering in the SM and the  $U(1)_{B-L}$  model. In Sec. IV, we estimate the sensitivity reach at FASER $\nu$ , FASER $\nu$ 2, SND@LHC, and FLArE. We compare the results of Sec. IV to other experiments in Sec. V. We conclude in Sec. VI.

## II. THE $U(1)_{B-L}$ MODEL

We first introduce the  $U(1)_{B-L}$  gauge interactions of a new gauge boson, denoted by  $A'$ , to the SM fermions with the charge proportional to quantum number  $B-L$  of the fermions, where  $B$  and  $L$  are the baryon and lepton number, respectively. The interaction Lagrangian is given by

$$\mathcal{L}_{B-L} \supset g_{B-L} \left[ -\bar{l}\gamma^\mu A'_\mu l - \bar{\nu}_\alpha \gamma^\mu A'_\mu \nu_\alpha + \frac{1}{3} \bar{q}\gamma^\mu A'_\mu q \right], \quad (1)$$

where  $g_{B-L}$  is the  $U(1)_{B-L}$  coupling strength and  $q, l$  and  $\nu$  are quark, charged lepton and neutrino fields, respectively. Here we assume that  $A'$  couples vectorially to the SM fermions without loss of generality and the gauge boson  $A'$  receives the mass by either a spontaneous breaking of the  $U(1)_{B-L}$  or Stueckelberg mechanism, which is not our concern here. Thus, the model is characterized by the mass  $M_{A'}$  and the gauge coupling strength  $g_{B-L}$ .

Hidden-sector  $U(1)'$  models belong to a class of hidden-valley models [5]. It is often used to build models of dark matter. Such  $U(1)'$  models can be connected to the SM via a small kinetic mixing though it is hidden from our world. The mass of  $A'$  and mixing parameter  $\epsilon$  can be tuned to satisfy the relic density of the Universe. The Lagrangian describing the kinetic mixing with the SM hypercharge field is given by

$$\mathcal{L} = -\frac{1}{4} B'^2_{\mu\nu} - \frac{1}{4} F''^2_{\mu\nu} + \frac{1}{2} \epsilon B'_{\mu\nu} F''^{\mu\nu} + \frac{1}{2} M_{A'}^2 A''^2_\mu - g_Y \frac{Y}{2} B'_\mu \bar{f} \gamma^\mu f, \quad (2)$$

where  $B'_{\mu\nu} = \partial_\mu B'_\nu - \partial_\nu B'_\mu$  and  $F''_{\mu\nu} = \partial_\mu A''_\nu - \partial_\nu A''_\mu$ , and here the primed fields  $B'$  and  $A''$  denote the fields before rotating into physical fields. The last term denotes the interaction of  $B'_\mu$  with SM fermions. The kinetic mixing term can be eliminated by rotating  $(B'_\mu, A''_\mu)$  into physical fields  $(B_\mu, A'_\mu)$  by a field redefinition:

$$\begin{aligned} B'_\mu &\simeq B_\mu + \epsilon A'_\mu \\ A''_\mu &\simeq A'_\mu. \end{aligned}$$

Then the Lagrangian becomes

$$\mathcal{L} = -\frac{1}{4}B_{\mu\nu}^2 - \frac{1}{4}F_{\mu\nu}^{\prime 2} + \frac{1}{2}M_{A'}^2 A_{\mu}^{\prime 2} - g_Y \frac{Y}{2}(B_{\mu} + \epsilon A'_{\mu}) \bar{f} \gamma^{\mu} f. \quad (3)$$

Thus, the hidden  $U(1)'$  gauge boson  $A'$  can now couple to the SM fermion with an interaction strength  $\epsilon g_Y(Y/2)$ .

Conventionally, the term "dark photon" is used to refer to a  $U(1)'$  gauge boson that is coupled to the SM only through kinetic mixing with the photon. Strictly speaking such a dark photon does not couple to neutrinos because of zero electric charge of the neutrino. However, as shown above the kinetic mixing with the  $B_{\mu}$  does create an interaction of  $A'$  with neutrinos proportional to the hypercharge of the neutrinos. Simultaneously, it also mixes with the SM  $Z$  boson and thus the mixing parameter  $\epsilon$  is constrained to be small by the LEP data [1]. Nevertheless, we will continue to use the term dark photon in a more general sense for the gauge bosons coupled to SM particles only through the kinetic mixing.

The  $U(1)_{B-L}$  gauge boson couples to the left- and right-handed fermions with the same quantum number, and thus the interaction is vector-like. On the other hand, the hidden  $U(1)'$  gauge boson discussed above couples to SM fermions with quantum number equal to the hypercharge of the fermion. For simplicity we assume the hidden  $U(1)'$  gauge boson also has the vector-like couplings to neutrinos and electrons with the hypercharge  $Y/2 = -1/2$ . We can easily translate between the two coupling strengths:

$$|g_{B-L}| = |\epsilon g_Y(Y/2)|, \quad (4)$$

where we take the hypercharge for electrons and neutrinos to be  $Y/2 = -1/2$ .

### III. NEUTRINO-ELECTRON SCATTERING

Neutrino-electron scattering is a viable alternative to collider searches for physics beyond the SM [12]. In this section, we discuss the neutrino-electron scattering in the SM and with the presence of a dark photon.

#### A. Standard Model cross section

In SM, the  $\nu_e$ - $e$  scattering takes place via the charged and neutral current interactions. However, the  $\nu_{\mu}$ - $e$  and  $\nu_{\tau}$ - $e$  scatterings are mediated only by the neutral  $Z$  boson. The

Table I. Valued for  $a$  and  $b$  in the expression of Eq. (5), where  $\alpha = \mu, \tau$

Process	$a$	$b$
$\nu_e e^- \rightarrow \nu_e e^-$	$\sin^2 \theta_w + \frac{1}{2}$	$\sin^2 \theta_w$
$\bar{\nu}_e e^- \rightarrow \bar{\nu}_e e^-$	$\sin^2 \theta_w$	$\sin^2 \theta_w + \frac{1}{2}$
$\nu_\alpha e^- \rightarrow \nu_\alpha e^-$	$\sin^2 \theta_w - \frac{1}{2}$	$\sin^2 \theta_w$
$\bar{\nu}_\alpha e^- \rightarrow \bar{\nu}_\alpha e^-$	$\sin^2 \theta_w$	$\sin^2 \theta_w - \frac{1}{2}$

differential cross-section in the rest frame of the electron can be expressed as [7]

$$\left[ \frac{d\sigma}{dT}(\nu e^- \rightarrow \nu e^-) \right]_{SM} = \frac{2G_F^2 m_e}{\pi E_\nu^2} (a^2 E_\nu^2 + b^2 (E_\nu - T)^2 - ab m_e T), \quad (5)$$

where  $G_F$  denotes the Fermi constant,  $T$  indicates the recoil energy of the electron,  $E_\nu$  is the energy of the incoming neutrino, and  $m_e$  is the mass of electron. The values for  $a$  and  $b$  depend on the neutrino flavor, which are given in Table. I. The contributing Feynman diagrams of the SM neutrino-electron scattering in the SM are shown in Fig. 1(a).

## B. Dark photon contribution to the cross-section

The scattering cross section of a neutrino off an electron or nucleus depends mildly on the chiral structure of the  $A'$  coupling. Spectra would look similar for other chiral structures, but here, for the sake of concreteness, we assume that  $A'$  has pure vector couplings of the form  $\bar{f}\gamma^\mu f A'_\mu$ . With this assumption we obtain the differential cross section for neutrino-electron scattering as a function of the recoil energy. The effective renormalizable Lagrangian given in Eq. (3), where the SM photon and dark photon mixed via a kinetic term. This mixing has been studied widely in literature [9–11]. The  $U(1)_{B-L}$  gauging induces a new coupling  $g_{B-L}$  in addition to its mass  $M_{A'}$  and  $\epsilon$ . The parameter space of the current dark photon model contains a set of three variables  $(M_{A'}, \epsilon, g_{B-L})$ . In this study we focus on the dark model with two parameters,  $M_{A'}$  and  $g_{B-L}$ . Figure 1(b) shows the Feynman diagram of neutrino-electron scattering mediated by the dark photon.

The scattering cross section from the pure contribution of  $A'$  is expressed as follows [7].

$$\left[ \frac{d\sigma}{dT}(\nu e^- \rightarrow \nu e^-) \right]_{DP} = \frac{g_{B-L}^4 m_e}{4\pi E_\nu^2 (M_{A'}^2 + 2m_e T)^2} (2E_\nu^2 + T^2 - 2TE_\nu - m_e T) \quad (6)$$




Figure 1. (a) Contributing Feynman diagrams for anti-electron-neutrino and electron scattering via the  $W$  and  $Z$  boson exchanges. Note that muon- and tau-neutrinos (including anti-muon- and anti-tau-neutrinos) scatter with electron only via the  $Z$  boson exchange. (b) The Feynman diagram showing the scattering of (anti-)neutrino with electron via the  $t$ -channel dark photon  $A'$  exchange.

where the cross section is flavor blind. On the other hand, the interference term in the differential cross-section depends on the neutrino flavor and whether it is the neutrino or anti-neutrino. The interference terms are given by [7].,

$$\frac{d\sigma_{\text{INT}}(\nu_e e^-)}{dT} = \frac{g_{B-L}^2 G_F m_e}{2\sqrt{2}\pi E_\nu^2 (M_{A'}^2 + 2m_e T)} (2E_\nu^2 - m_e T + \beta) \quad (7)$$

$$\frac{d\sigma_{\text{INT}}(\bar{\nu}_e e^-)}{dT} = \frac{g_{B-L}^2 G_F m_e}{2\sqrt{2}\pi E_\nu^2 (M_{A'}^2 + 2m_e T)} (2E_\nu^2 + 2T^2 - T(4E_\nu + m_e) + \beta) \quad (8)$$

$$\frac{d\sigma_{\text{INT}}(\nu_\alpha e^-)}{dT} = \frac{g_{B-L}^2 G_F m_e}{2\sqrt{2}\pi E_\nu^2 (M_{A'}^2 + 2m_e T)} (-2E_\nu^2 + m_e T + \beta) \quad (9)$$

$$\frac{d\sigma_{\text{INT}}(\bar{\nu}_\alpha e^-)}{dT} = \frac{g_{B-L}^2 G_F m_e}{2\sqrt{2}\pi E_\nu^2 (M_{A'}^2 + 2m_e T)} (-2E_\nu^2 - 2T^2 - T(4E_\nu + m_e) + \beta) \quad (10)$$

where  $\beta$  is given by

$$\beta = \sin^2 \theta_w (8E_\nu^2 - 8E_\nu T - 4m_e T + 4T^2).$$

Here  $\alpha$  in  $\nu_\alpha$  denotes either  $\mu$  or  $\tau$ . The  $\nu_\mu e^-$  and  $\nu_\tau e^-$  scatterings can occur through the SM  $Z$  and the dark photon  $A'$  mediated processes, while the  $\nu_e e^-$  scattering is mediated by the SM  $Z, W$  bosons and dark photon  $A'$ . The pure contribution of the dark photon cross section is proportional to the fourth power of  $g_{B-L}$  while the interference term is proportional to the second power of  $g_{B-L}$ . The total cross-section  $\sigma_T$  is then given by  $\sigma_T = \sigma_{DP} + \sigma_{INT} + \sigma_{SM}$ .

#### IV. SENSITIVITY REACH ON PARAMETER SPACE OF THE DARK PHOTON MODEL AT THE FORWARD PHYSICS EXPERIMENTS

Proton-proton collisions at the Large Hadron Collider (LHC) produce many particles along the beam axis. Forward Physics Facilities (FPF) [12–21], situated several hundred meters away from the ATLAS interaction point and protected by concrete and rock, will host a wide variety of experiments probing the SM processes and also searching for new physics beyond the standard model (BSM) [12, 22–39]. Each experiment in the FPF is uniquely optimized for its own physics goals.

In this section, we study the neutrino-electron scattering mediated by the  $t$ -channel dark photon exchange at the running FASER $\nu$  and the proposed FASER $\nu$ 2, FLArE (Forward Liquid Argon Experiment), and SND@LHC, where FASER $\nu$ 2 and FLArE are parts of the proposed Forward Physics Facility (FPF). FPF also contains other detectors called the FASER2, Advanced SND, and FORMOSA. The dark photon sensitivity at FASER2 was discussed in the literature [12, 35, 40, 41].

The FASER $\nu$  and SND@LHC detectors are made of tungsten with masses of 1.1 tons and 830 kg respectively. FASER $\nu$  is made of 1000 layers of emulsion films interleaved with tungsten plates of thickness of 1 mm. The size of FASER $\nu$  is 25 cm  $\times$  30 cm  $\times$  1.1 m [3, 4]. That of SND@LHC is 39 cm  $\times$  39 cm [4]. The upgraded version of FASER $\nu$  – FASER $\nu$ 2 will have a size 50 cm  $\times$  50 cm  $\times$  5 m and will weigh 10 tons. FLArE is composed of 10 ton-scale liquid argon time projection chamber (LArTPC) of the type being employed in several modern neutrino experiments. The size of the FLArE detector is 1 m  $\times$  1 m  $\times$  7 m and will weigh 10 tons.

We are interested in the neutrino-electron scattering at the aforementioned detectors. It is well known that huge number of hadrons, such as pions, kaons and other hadrons, are produced along the beam direction. These hadrons will decay during the flight, thus producing a lot of neutrinos of all three flavors at very high energy up to a few TeV[19]. We find that muon neutrinos are mostly produced from charged-pion decays, electron neutrinos from hyperon, kaon and  $D$ -meson decays, and tau neutrinos from  $D_s$  meson decays. With average energies ranging from 600 GeV to 1 TeV, the spectra of the three neutrino flavors cover a broad energy range.

### A. Dark photon interactions at FASER $\nu$ and FASER $\nu$ 2

In this subsection we discuss the neutrino-electron  $\nu e^- \rightarrow \nu e^-$  scattering at FASER $\nu$  and FASER $\nu$ 2 mediated by a new neutral  $U(1)_{B-L}$  gauge field  $A'$ . The cross-section of the fixed target neutrino-electron scattering is given in Sec. III. We consider the dark photon of mass  $M_{A'}$  ranging from  $10^{-6}$  GeV to 10 GeV with an incoming neutrino beam energy  $E_\nu$  from 10 GeV to 10 TeV. The cross section is calculated by setting the value of  $g_{B-L} = 10^{-2}$ . We use the neutrino fluxes and energy spectra obtained in [19, 20] to study the neutrinos that pass through FASER $\nu$  and FASER $\nu$ 2. We estimate the 90% C.L. sensitivity reach on the parameter space of the dark photon model. We show that the best sensitivity can be achieved in the small  $M_{A'}$  region.

#### 1. FASER $\nu$

The FASER $\nu$  neutrino detector is located in front of the FASER main detector [18, 20], which is aligned along the collision axis to maximize the neutrino interaction. FASER $\nu$ [20] is a 25 cm  $\times$  30 cm  $\times$  1.1 m emulsion detector, consisting of 770 layers of emulsion films interleaved with 1-mm-thick tungsten plates with mass 1.2 tons [3]. The features of high density and shorten radiation length of tungsten are advocated as the material for the detector, because such features help keeping the detector small in size and localizing electromagnetic showers in a small volume.

The expected numbers of neutrino events from the three flavors of active neutrinos with different dark photon masses  $M_{A'}$  in FASER $\nu$  are shown in Fig. 2. The solid lines represent the number of neutrino events as a function of dark photon mass  $M_{A'}$  while the dashed lines represent the SM event rates. The number of events associated with dark photon overlaps with the SM value as the dark photon mass  $M_{A'}$  increases. The highest number of events comes in with the muon neutrino  $\nu_\mu$  scattering with electron ( $\nu_\mu e^- \rightarrow \nu_\mu e^-$ ), while the lowest one is from the tau neutrino  $\nu_\tau$  scattering ( $\nu_\tau e^- \rightarrow \nu_\tau e^-$ ), and that from the electron neutrino  $\nu_e$  scattering is in between. Total number of events follows the same behavior as the scattering cross section. We can summarize the results in Fig. 2 as follows,  $N_{\nu_\tau}(g_{B-L}, M_{A'}) < N_{\nu_e}(g_{B-L}, M_{A'}) < N_{\nu_\mu}(g_{B-L}, M_{A'})$ .

To estimate the sensitivity reach in the parameter space ( $g_{B-L}$  vs  $M_{A'}$ ) of the dark photon






Figure 2. Total number of active neutrinos produced at FASER $\nu$  from neutrino-electron scattering ( $\nu_\alpha e^- \rightarrow \nu_\alpha e^-$ ), where  $\alpha = e, \mu, \tau$ . Contributions from neutrinos and anti-neutrinos are included.

model, we first calculate the predicted number of events  $N_{BSM}$  for the dark photon model and the SM number of events  $N_{SM}$ , and treat the statistical error as  $\sqrt{N_{BSM}}$  and systematic uncertainty  $\sigma$  as a fraction ( $\sigma = 20\%, 5\%$ ) of the normalization of the SM predictions. We then define the measure of  $\chi^2$  as a function of  $(g_{B-L})$  and a nuisance parameter  $\alpha$  as follows [28, 42]

$$\chi^2(g_{B-L}, \alpha) = \min_{\alpha} \left[ \frac{(N_{BSM}^{\nu_e} - (1 + \alpha)N_{SM}^{\nu_e})^2}{N_{BSM}^{\nu_e}} + \frac{(N_{BSM}^{\nu_\mu} - (1 + \alpha)N_{SM}^{\nu_\mu})^2}{N_{BSM}^{\nu_\mu}} + \frac{(N_{BSM}^{\nu_\tau} - (1 + \alpha)N_{SM}^{\nu_\tau})^2}{N_{BSM}^{\nu_\tau}} + \left(\frac{\alpha}{\sigma}\right)^2 \right], \quad (11)$$

where  $N_{BSM} = N_{DP} + N_{INT} + N_{SM}$  and the minimization is over the nuisance parameter  $\alpha$ . Here  $N_{DP}$  is the number of events from the dark photon mediated process only,  $N_{INT}$  is the interference term. Here we have treated the systematic uncertainties in each neutrino flavor to be the same and use only one nuisance parameter  $\alpha$ . Physics-wise the systematic uncertainties come from theoretical calculations, the flux of neutrinos from the ATLAS IP, detector response, etc. We show in the left panel of Fig. 3 the 90% C.L. sensitivity reach




Figure 3. Left Panel: Sensitivity reach on the coupling  $g_{B-L}$  versus the dark photon mass  $M_{A'}$  achieved at FASER $\nu$ . Systematic uncertainties  $\sigma = 5\%$ ,  $20\%$  and without systematic uncertainties are shown. Right Panel: sensitivity reach on the coupling  $g_{B-L}^\alpha$  for each neutrino flavor  $\alpha = e, \mu, \tau$  versus the dark photon mass  $M_{A'}$  achieved at FASER $\nu$  (without systematic uncertainties) are shown.

(corresponding to  $\chi^2 = 2.71$ ) of the coupling strength  $g_{B-L}$  versus the dark photon mass  $M_{A'}$  at FASER $\nu$ . The higher the systematic uncertainty the weaker the limit on  $g_{B-L}$  will be. Nevertheless, the differences among  $\sigma = 5\%$ ,  $20\%$  and without systematic uncertainties are relatively small. The sensitivity reach on  $g_{B-L}$  is the best at very small  $M_{A'}$ , around  $2 \times 10^{-6}$  at  $M_{A'} = 10^{-6}$  GeV and reduces to about  $2 \times 10^{-2}$  at  $M_{A'} = 10$  GeV.

FASER $\nu$  is primarily designed for the purpose of identifying the flavors of neutrinos [20]. The expected sensitivity for each flavor at FASER $\nu$  is shown in the right panel of Fig. 3. We only show the curves with no systematic uncertainty included. The curves can be compared to the corresponding one “Without Systematic” of the left panel of Fig. 3. For the evaluation of  $\chi^2$ , we consider the special case of Eq. 11 with  $\alpha = 0$  and it reduces to

$$\chi^2(g_{B-L}^\alpha) = \left[ \frac{(N_{BSM}^{\nu_\alpha} - N_{SM}^{\nu_\alpha})^2}{N_{BSM}^{\nu_\alpha}} \right] \quad (12)$$

where  $\alpha = e, \mu, \tau$ . The green and orange curves of Fig. 3 depict the sensitivity reach of  $g_{B-L}^\tau$  and  $g_{B-L}^e$  ( $\chi^2 = 2.71$ ) versus  $M_{A'}$ , respectively, while the blue curve represents the sensitivity reach of  $g_{B-L}^\mu$  versus  $M_{A'}$ . It is clear from the right and left panel of Fig. 3 that

the overall sensitivity reach of  $g_{B-L}$  is dominated by  $g_{B-L}^\mu$ . The sensitivity reach by the neutrino flavors is in the order of  $\nu_\tau < \nu_e < \nu_\mu$ .

## 2. FASER $\nu$ 2

FASER $\nu$ 2 is a much more advanced version of FASER $\nu$ [13]. FASER $\nu$ 2 detector is currently proposed to be composed of 3330 emulsion layers interleaved with 2 mm thick tungsten plates. It will also accommodate a veto detector and an interface detector to the FASER2 spectrometer, with one detector in the middle of the emulsion modules and the other detector downstream of the emulsion modules to make the global analysis and muon charge measurement possible. Both the emulsion modules and interface detectors will be put in a cooling system. The total volume of the tungsten target is  $50 \text{ cm} \times 50 \text{ cm} \times 5 \text{ m}$ , and the mass is 10 tons. The detector length, including the emulsion films and interface detectors, will be about 8 m.




Figure 4. Total number of active neutrinos produced at FASER $\nu$ 2 from neutrino-electron scattering ( $\nu_\alpha e^- \rightarrow \nu_\alpha e^-$ ), where  $\alpha = e, \mu, \tau$ . Contributions from neutrinos and anti-neutrinos are included.

In this subsection we calculate the sensitivity reach on the parameter space of the dark

photon model at FASER $\nu$ 2. We follow the same procedures as the last subsection for the estimation of the total number of neutrino events at FASER $\nu$ 2. The total numbers of neutrino events at FASER $\nu$ 2 are shown in Fig. 4, which follows the same notation as Fig. 2. Comparing with FASER $\nu$ , FASER $\nu$ 2 could collect a larger number of neutrino events. There are sizable increments in both the BSM neutrino events and SM neutrino events. In the heavy  $M_{A'}$  mass regime ( $M_{A'} \gtrsim 5$  GeV), the  $N_{\nu_\mu}$  curve overlaps with the  $N_{\nu_e}$  curve. This behavior arises from the large interference effect associated with the muon neutrinos. Among the three neutrino flavors, the tau neutrino is the least in event rates.




Figure 5. Left Panel: Sensitivity reach on the coupling  $g_{B-L}$  versus the dark photon mass  $M_{A'}$  achieved at FASER $\nu$ 2. Systematic uncertainties  $\sigma = 5\%$ ,  $20\%$  and without systematic uncertainties are shown. Right Panel: sensitivity reach on the coupling  $g_{B-L}^\alpha$  for each neutrino flavor  $\alpha = e, \mu, \tau$  versus the dark photon mass  $M_{A'}$  achieved at FASER $\nu$ 2 (without systematic uncertainties) are shown.

Sensitivity reach at FASER $\nu$ 2 is carried out by a similar  $\chi^2$  analysis in Eq. (11). We estimate the 90% C.L. sensitivity reach (corresponding to  $\chi^2 = 2.71$ ) of the coupling strength  $g_{B-L}$  versus dark photon mass  $M_{A'}$  at FASER $\nu$ 2, as shown in Fig. 5. The dark photon sensitivity reach at FASER $\nu$ 2 is shown in the left panel of Fig. 5, in which the curves with  $\sigma = 5\%$  and  $20\%$  are overlapping with each other. FASER $\nu$ 2 has better sensitivity compared to FASER $\nu$ . In the very small dark photon mass region,  $M_{A'} = 10^{-6}$  GeV,

FASER $\nu$ 2 can reach as low as  $10^{-6}$  while FASER $\nu$  only reaches around  $2 \times 10^{-6}$ . The sensitivity devalues with the increment of dark photon mass.

The expected sensitivity reach of FASER $\nu$ 2 for each neutrino flavor is given in the right panel of Fig. 5. The effect of interference is clearly visible from the cross-over of  $g_{B-L}^e$  curve over the  $g_{B-L}^\mu$  curve. In the higher  $M_{A'}$  mass region, the  $g_{B-L}^e$  curve has better sensitivity compared with the  $g_{B-L}^\mu$  curve. The total sensitivity curve of  $g_{B-L}$  in the left panel of Fig. 5 resembles the  $g_{B-L}^\mu$  curve of the right panel in the low  $M_{A'}$  region and switches to  $g_{B-L}^e$  in the heavier  $M_{A'}$  region.

## B. Dark photon interactions at SND@LHC




Figure 6. Total number of active neutrinos produced at SND@LHC from neutrino-electron scattering ( $\nu_\alpha e^- \rightarrow \nu_\alpha e^-$ ), where  $\alpha = e, \mu, \tau$ . Contributions from neutrinos and anti-neutrinos are included.

SND@LHC is a compact detector of neutrinos at the LHC and it consists of a neutrino target followed downstream by a device for detecting muons, which are produced when neutrinos interact with the target. It is especially complementary to the FASER $\nu$ . Targets are made from tungsten plates interlaced with emulsion films and electronic tracking devices.

Emulsion films display the tracks of particles produced in neutrino interactions, while electronic tracking devices provide the time stamps for these tracks. The tracking devices also measure the energy of the neutrinos along with the muon detector.

The predicted number of neutrino events mediated by the dark photon of mass ranging from  $M_{A'} \sim 10^{-6} - 10$  GeV is shown in Fig. 6. Comparing with FASER $\nu$  and FASER $\nu$ 2, SND@LHC has fewer neutrino events. We summarize the predicted numbers of neutrino events for these three forward-physics experiments as follows,  $N_{\text{BSM}}^{\text{SND@LHC}}(g_{B-L}, M_{A'}) < N_{\text{BSM}}^{\text{FASER}\nu}(g_{B-L}, M_{A'}) < N_{\text{BSM}}^{\text{FASER}\nu 2}(g_{B-L}, M_{A'})$ . The difference in the event rate arises from the neutrino flux at each detector volume.




Figure 7. Left Panel: Sensitivity reach on the coupling  $g_{B-L}$  versus the dark photon mass  $M_{A'}$  achieved at SND@LHC. Systematic uncertainties  $\sigma = 5\%$ ,  $20\%$  and without systematic uncertainties are shown. Right Panel: sensitivity reach on the coupling  $g_{B-L}^\alpha$  for each neutrino flavor  $\alpha = e, \mu, \tau$  versus the dark photon mass  $M_{A'}$  achieved at SND@LHC (without systematic uncertainties) are shown.

A similar  $\chi^2$  analysis (Eq. 11) is carried out at SND@LHC to assess dark photon sensitivity. Accordingly, a 90% C.L. sensitivity reach is estimated (corresponding to  $\chi^2 = 2.71$ ) for the coupling strength  $g_{B-L}$  versus the dark photon mass  $M_{A'}$  at the SND@LHC, as shown in the left panel of Fig. 7. Here we have assumed a benchmark detector made of tungsten with dimensions  $39 \text{ cm} \times 39 \text{ cm}$  at the 14 TeV LHC with an integrated luminosity



of  $L = 150 \text{ fb}^{-1}$ . The higher the systematic uncertainty the weaker the limit on  $g_{B-L}$  will be. Nevertheless, the differences among  $\sigma = 5\%$ ,  $20\%$  and without systematic uncertainties are relatively small. The sensitivity reach on  $g_{B-L}$  is the best at very small  $M_{A'}$  as low as  $4 \times 10^{-6}$  at  $M_{A'} = 10^{-6}$  GeV and devalues to about  $8 \times 10^{-2}$  at  $M_{A'} = 10$  GeV. The reach of  $g_{B-L}^\alpha$  for each neutrino flavor as a function of dark photon mass  $M_{A'}$  is shown in the right panel of Fig. 7.




Figure 8. Total number of active neutrinos produced at FLArE (10 tons) from neutrino-electron scattering ( $\nu_\alpha e^- \rightarrow \nu_\alpha e^-$ ), where  $\alpha = e, \mu, \tau$ . Contributions from neutrinos and anti-neutrinos are included.

### C. Dark photon interactions at FLArE (10 tons)

As a part of the suite of detectors for the FPF, the liquid argon time projection chamber (LArTPC) is being considered. This type of detector offers the capability to determine particle identification, track angle, and kinetic energy over a wide range of energies. Dark matter (DM) direct detection searches [39] and neutrino experiments have successfully used liquid argon as an active detector. Detecting neutrino energy [13] and searching for dark matter are therefore well suited to be put together to utilize such a liquid-argon detector.

There is an imperative motivation to detect and measure TeV-scale neutrino events from a laboratory-generated and well-characterized source. Moreover, the LHC could be the only source of high-energy intense tau neutrinos on Earth. Here we have assumed a benchmark detector made of tungsten with dimensions  $1\text{ m} \times 1\text{ m} \times 8\text{ m}$  at the 14 TeV LHC with an integrated luminosity of  $L = 3000\text{ fb}^{-1}$ .

The expected number of neutrino events at FLArE(10 tons) is presented in Fig. 8. Comparing with the aforementioned forward physics experiments, the expected number of neutrino events at the FLArE as a function of dark photon mass  $M_{A'}$  and coupling  $g_{B-L}$  are in the following ordering:  $N_{\text{BSM}}^{\text{SND@LHC}}(g_{B-L}, M_{A'}) < N_{\text{BSM}}^{\text{FASER}\nu}(g_{B-L}, M_{A'}) < N_{\text{BSM}}^{\text{FLArE}}(g_{B-L}, M_{A'}) < N_{\text{BSM}}^{\text{FASER}\nu^2}(g_{B-L}, M_{A'})$ . This ordering will reflect the bound on the dark photon coupling.




Figure 9. Left Panel: Sensitivity reach on the coupling  $g_{B-L}$  versus the dark photon mass  $M_{A'}$  achieved at FLArE (10 tons). Systematic uncertainties  $\sigma = 5\%$ ,  $20\%$  and without systematic uncertainties are shown. Right Panel: sensitivity reach on the coupling  $g_{B-L}^\alpha$  for each neutrino flavor  $\alpha = e, \mu, \tau$  versus the dark photon mass  $M_{A'}$  achieved at FLArE(10 tons) (without systematic uncertainties) are shown.

The bound on the dark-photon coupling is obtained through the  $\chi^2$  defined in Eq. (11). We obtain the 90% C.L. dark photon sensitivity at FLArE (10 tons), as shown in the left panel of the Fig. 9, which shows the coupling  $g_{B-L}$  versus the dark photon mass  $M_{A'}$  with



systematic uncertainties  $\sigma = 5\%$ ,  $20\%$  and without systematic uncertainties. The right panel of Fig. 9 obtained by Eq. (12) shows the sensitivity reach on the coupling  $g_{B-L}^\alpha$  for each neutrino flavor  $\alpha = e, \mu, \tau$  versus the dark photon mass  $M_{A'}$  achieved at FLArE (10 tons) (without systematic uncertainties). The effect of interference is visible from the right panel of the Fig. 9, in which the  $g_{B-L}^e$  curve overlaps with the  $g_{B-L}^\mu$  curve for  $M_{A'} \gtrsim 0.1$  GeV.

## V. COMPLEMENTARITY OF FASER $\nu/2\nu$ , SND@LHC AND FLArE RESULTS

We have obtained the future sensitivity reach in the parameter space ( $g_{B-L}, M_{A'}$ ) of the dark photon model at FASER $\nu$ , FASER $2\nu$ , SND@LHC, and FLArE(10 tons). In this section, we compare these forward physics experiments with existing neutrino-electron scattering experiments. While comparing the Forward Physics Experiments to one another, FASER $\nu 2$  has the best sensitivity among the four experiments while SND@LHC is the least sensitive one. For the case of a very small dark photon mass  $M_{A'} = 10^{-6}$  GeV the sensitivity reach of  $g_{B-L}$  is around  $4 \times 10^{-6}$  at SND@LHC,  $2 \times 10^{-6}$  at FASER $\nu$ ,  $1.6 \times 10^{-6}$  at FLArE(10 tons), and  $10^{-6}$  at FASER $\nu 2$ . FASER $\nu 2$  has the best sensitivity for the whole range of dark photon mass considered in this work.

The GEMMA Collaboration [43] measured the  $\bar{\nu}_e - e$  scattering cross section, which can be used to put a bound on neutrino magnetic moments. The Gemma (dashed magenta curve) in Fig. 10 has the best sensitivity among all the considered forward Physics experiments at the low dark photon mass region. The Gemma sensitivity gets weakened with increment of dark photon mass, after  $M_{A'} > 0.01$  GeV there is a considerable reduction in the Gemma sensitivity to the dark photon coupling. In the heavier mass region the forward physics experiments estimated in this work in general have better sensitivity.

The TEXONO-NPCGe[44] estimated the constraints on millicharged neutrinos via analysis of data from atomic ionizations. NPCGe (dashed orange curve) also has better sensitivity in the low dark photon mass region, and the sensitivity gets weakened with increases in dark photon mass. The sensitivity curve of NPCGe follows the similar pattern of Gemma. The TEXONO-HPGe [45] estimated the limit on the electron-neutrino magnetic moment. FASER $\nu 2$  (blue curve) has better sensitivity than that of HPGe (dashed blue) in the very small dark photon mass  $M_{A'} \sim 10^{-6}$  GeV. However, in the intermediate mass




Figure 10. Future sensitivity reach of FASER $\nu$  (dark green), FASER2 $\nu$  (red), SND@LHC (violet), and FLArE (10 tons)(blue) at 90% C.L.(without systematic uncertainties) on the gauge coupling  $g_{B-L}$  of the  $U_{B-L}(1)$  group as the function of dark photon mass  $M_{A'}$ . Other existing constraints at 90% C.L. shown include (i)  $\bar{\nu}_e - e$  scattering measurement at Gemma (dashed magenta curve), (ii) NPCGe (dashed orange curve), (iii) HPGe (dashed blue curve), (iv) elastic scattering of neutrinos using a liquid scintillator at Borexino (dashed black curve), (v) CsI (dashed red curve) measurement of  $\bar{\nu}_e - e^-$ , (vi) LSND (dashed light green curve) measurement of  $\nu_e - e^-$ , (vii) CHARM-II- $\nu_\mu - e$  (dashed light blue), and (viii) CHARM-II- $\bar{\nu}_\mu - e$  (dashed light yellow).

region  $10^{-6} \text{ GeV} < M_{A'} < 0.01 \text{ GeV}$ , HPGe has better sensitivity than all the forward physics experiments considered in this work.

The Borexino Collaboration [46] measured the spectrum of the  ${}^7\text{Be}$  solar neutrino (with 862 keV energy) via the elastic scattering of neutrinos using a liquid scintillator. The sensitivity curve of Borexino (dashed black curve) also followed the similar pattern of TEXONO. However, both FLArE (red curve) and FASER $\nu$ 2 (blue) have better sensitivity at  $M_{A'} = 10^{-6} \text{ GeV}$  compared with Borexino.

TEXONO-CsI [47] measured the  $\bar{\nu}_e - e^-$  scattering. FASER $\nu$  (dark green), FASER $\nu$ 2




Figure 11. Future sensitivity reach of FASER $\nu$  (dark green), FASER2 $\nu$  (red), SND@LHC (violet), and FLArE (10 tons)(blue) at 90% C.L.(without systematic uncertainties) on the kinetic mixing parameter  $\epsilon$  as a function of dark photon mass  $M_{A'}$ . Other existing constraints at 90% C.L. shown include (i)  $\bar{\nu}_e - e$  scattering measurement at Gemma (dashed magenta curve), (ii) NPCGe (dashed orange curve), (iii) HPGe (dashed blue curve), (iv) elastic scattering of neutrinos using a liquid scintillator at Borexino (dashed black curve), (v) CsI (dashed red curve) measurement of  $\bar{\nu}_e e^-$ , (vi) LSND (dashed light green curve) measurement of  $\nu_e - e^-$ , (vii) CHARM-II- $\nu_\mu - e$  (dashed light blue), and (viii) CHARM-II- $\bar{\nu}_\mu - e$  (dashed light yellow).

(blue), and FLArE (red curve) all have better sensitivity in the very low  $M_{A'}$  mass region ( $M_{A'} < 2 \times 10^{-6}$  GeV). However, in the heavier dark photon mass region ( $M_{A'} < 1$  GeV) CsI has better sensitivity than any of the forward physics experiments. LSND [48] also followed the similar pattern as CsI. However SND@LHC(violet curve) has better sensitivity compared with LSND (dashed light green) in the low  $M_{A'}$  mass region. In all the above, except CsI, the forward physics experiments have better sensitivity in the heavy dark photon mass region. CHARM II [49] also measured the neutrino-electron scattering in the lower dark photon mass region. The forward physics experiments have better sensitivity compared

to CHARM II (CHARM-II  $\nu_\mu - e$  (dashed light blue) and CHARM-II  $\bar{\nu}_\mu - e$  (dashed light yellow)).

The bounds on the kinetic mixing parameter  $\epsilon$  as a function of dark photon mass  $M_{A'}$  are shown in Fig. 11. We have translated the limits on kinetic mixing from the limits on the gauge coupling  $g_{B-L}$  by  $|g_{B-L}| = |\epsilon g_Y(Y/2)|$  in Eq. 4. The sensitivity reach on the dark photon kinetic mixing  $\epsilon$  follows the same pattern as  $g_{B-L}$  coupling. While comparing the bounds on  $\epsilon$  achieved in the forward physics experiments with existing neutrino-electron scattering experiments, the forward physics experiments have better bounds in the very low dark photon mass region ( $M_{A'} \sim 10^{-6}$  GeV) and in the heavier dark photon mass region  $M_{A'} > 0.1$  GeV. For the case of a very small dark photon mass  $M_{A'} = 10^{-6}$  GeV the sensitivity reach of  $\epsilon$  is around  $10^{-5}$  for SND@LHC,  $5 \times 10^{-5}$  for FASER $\nu$ ,  $4 \times 10^{-5}$  for FLArE(10 tons), and  $3 \times 10^{-5}$  for FASER $\nu$ 2. FASER $\nu$ 2 has the best sensitivity for all dark photon mass range considered in this work. The sensitivity on  $\epsilon$  gets weakened with increment of dark photon mass.

## VI. CONCLUSIONS

Forward Physics Facilities (FPF) can host a number of experiments that make use of the unique opportunities of neutrino beams in the energy range of a few hundred GeV to TeV in exploring physics beyond the SM. We have investigated sensitivity reach on a couple of dark photon models at a number of experiments, including FASER $\nu$ , FASER $\nu$ 2, SND@LHC, and FLArE(10 tons). We employed two models: (i)  $U(1)_{B-L}$  and (ii) a hidden  $U(1)'$  that mixes with the SM gauge fields via the kinetic mixing (vector portal). The presence of such a  $U(1)_{B-L}$  gauge field or a dark photon gives rise to a detectable signal in neutrino-electron scattering. We have studied neutrino-electron scattering mediated via  $t$ -channel exchange of the  $U(1)_{B-L}$  gauge field at the FPE experiments such as FASER $\nu$ , FASER $\nu$ 2, SND@LHC and FLArE (10 tons), and calculated the expected sensitivity reach on the parameter space of the  $U(1)_{B-L}$  model. We investigated the advantage of FPF in a wide mass range of the  $U(1)_{B-L}$  gauge field and to determine the flavor dependence of the coupling between neutrino and this new gauge boson. We found that both FASER $\nu$  and SND@LHC are more sensitive to  $g_{B-L}^\mu$ . On the other hand, FASER $\nu$ 2 is more sensitive to  $g_{B-L}^\mu$  in the lower dark photon mass region, but shifts to  $g_{B-L}^e$  in the heavier dark photon mass region due to

the interference effect. A similar pattern happens in FLArE, however, in the heavier dark photon mass region both  $g_{B-L}^\mu$  and  $g_{B-L}^e$  receive similar sensitivity.

Once we obtained the sensitivities on the coupling  $g_{B-L}$  we translated the results into the sensitivities on the kinetic mixing parameter  $\epsilon$  via Eq. (4) as a function of dark photon mass  $M_{A'}$ . The bound of kinetic mixing also follows the same pattern of the gauge coupling  $g_{B-L}$ .

We have covered a wide mass range of dark photon  $10^{-6} \text{ GeV} \leq M_{A'} \leq 10 \text{ GeV}$  in our study. Among all the proposed forward physics experiments, FASER $\nu$ 2 has the best sensitivity to the  $U(1)_{B-L}$  and the dark photon model, whereas the SND@LHC has the least. The sensitivity curves for all the experiments follow a similar pattern, in which the sensitivity gets weakened with the increment of dark photon mass. We also compared the 90% C.L. sensitivity reach on the parameter space of the models at the FPF experiments with the other existing bounds from the experiments such as TEXONO, GEMMA, BOREXINO, LSND, and CHARM II.

In summary, the FPF experiments considered in this work can achieve better sensitivities at higher  $M_{A'}$  mass region because of the higher energy range of the neutrino flux coming from the ATLAS IP.

## ACKNOWLEDGEMENT

This work was supported in parts by the Ministry of Science and Technology with the grant number MoST-110-2112-M-007-017-MY3.

- 
- [1] R. L. Workman *et al.* [Particle Data Group], PTEP **2022**, 083C01 (2022)
  - [2] D. Z. Freedman, Phys. Rev. D **9**, 1389-1392 (1974) doi:10.1103/PhysRevD.9.1389
  - [3] Poster by D. Hayakawa in NEUTRINO 2022, held in Seoul Korea, May 30 – June 4, 2022.
  - [4] Talk by A.Di Crescenzo and Felix Kling in NEUTRINO 2022, held in Seoul Korea, May 30 – June 4, 2022.
  - [5] M. J. Strassler and K. M. Zurek, Phys. Lett. B **651**, 374-379 (2007) doi:10.1016/j.physletb.2007.06.055 [arXiv:hep-ph/0604261 [hep-ph]].

- [6] M. Fabbrichesi, E. Gabrielli and G. Lanfranchi, “The Dark Photon,” doi:10.1007/978-3-030-62519-1 [arXiv:2005.01515 [hep-ph]].
- [7] S. Bilmis, I. Turan, T. M. Aliev, M. Deniz, L. Singh and H. T. Wong, “Constraints on Dark Photon from Neutrino-Electron Scattering Experiments,” Phys. Rev. D **92**, no.3, 033009 (2015) doi:10.1103/PhysRevD.92.033009 [arXiv:1502.07763 [hep-ph]].
- [8] R. Harnik, J. Kopp and P. A. N. Machado, “Exploring nu Signals in Dark Matter Detectors,” JCAP **07**, 026 (2012) doi:10.1088/1475-7516/2012/07/026 [arXiv:1202.6073 [hep-ph]].
- [9] R. Essig, J. A. Jaros, W. Wester, P. Hansson Adrian, S. Andreas, T. Averett, O. Baker, B. Batell, M. Battaglieri and J. Beacham, *et al.* “Working Group Report: New Light Weakly Coupled Particles,” [arXiv:1311.0029 [hep-ph]].
- [10] R. Essig, R. Harnik, J. Kaplan and N. Toro, “Discovering New Light States at Neutrino Experiments,” Phys. Rev. D **82** (2010), 113008 doi:10.1103/PhysRevD.82.113008 [arXiv:1008.0636 [hep-ph]].
- [11] R. Essig, P. Schuster and N. Toro, “Probing Dark Forces and Light Hidden Sectors at Low-Energy e+e- Colliders,” Phys. Rev. D **80** (2009), 015003 doi:10.1103/PhysRevD.80.015003 [arXiv:0903.3941 [hep-ph]].
- [12] J. L. Feng, F. Kling, M. H. Reno, J. Rojo, D. Soldin, L. A. Anchordoqui, J. Boyd, A. Ismail, L. Harland-Lang and K. J. Kelly, *et al.* “The Forward Physics Facility at the High-Luminosity LHC,” [arXiv:2203.05090 [hep-ex]].
- [13] L. A. Anchordoqui, A. Ariga, T. Ariga, W. Bai, K. Balazs, B. Batell, J. Boyd, J. Bramante, M. Campanelli and A. Carmona, *et al.* “The Forward Physics Facility: Sites, experiments, and physics potential,” Phys. Rept. **968** (2022), 1-50 doi:10.1016/j.physrep.2022.04.004 [arXiv:2109.10905 [hep-ph]].
- [14] A. Ariga *et al.* [FASER], “FASER: ForwArd Search ExpeRiment at the LHC,” [arXiv:1901.04468 [hep-ex]].
- [15] H. Abreu *et al.* [FASER], “The tracking detector of the FASER experiment,” [arXiv:2112.01116 [physics.ins-det]].
- [16] H. Abreu *et al.* [FASER], “The trigger and data acquisition system of the FASER experiment,” JINST **16** (2021) no.12, P12028 doi:10.1088/1748-0221/16/12/P12028 [arXiv:2110.15186 [physics.ins-det]].

- [17] H. Abreu *et al.* [FASER], “Technical Proposal: FASERnu,” [arXiv:2001.03073 [physics.ins-det]].
- [18] A. Ariga *et al.* [FASER], “Technical Proposal for FASER: ForwArd Search ExpeRiment at the LHC,” [arXiv:1812.09139 [physics.ins-det]].
- [19] F. Kling and L. J. Nevay, “Forward neutrino fluxes at the LHC,” *Phys. Rev. D* **104** (2021) no.11, 113008 doi:10.1103/PhysRevD.104.113008 [arXiv:2105.08270 [hep-ph]].
- [20] H. Abreu *et al.* [FASER], “Detecting and Studying High-Energy Collider Neutrinos with FASER at the LHC,” *Eur. Phys. J. C* **80** (2020) no.1, 61 doi:10.1140/epjc/s10052-020-7631-5 [arXiv:1908.02310 [hep-ex]].
- [21] A. Ismail, R. Mammen Abraham and F. Kling, “Neutral current neutrino interactions at FASER $\nu$ ,” *Phys. Rev. D* **103** (2021) no.5, 056014 doi:10.1103/PhysRevD.103.056014 [arXiv:2012.10500 [hep-ph]].
- [22] S. Ansarifard and Y. Farzan, “Neutral exotica at FASER $\nu$  and SND@LHC,” *JHEP* **02** (2022), 049 doi:10.1007/JHEP02(2022)049 [arXiv:2109.13962 [hep-ph]].
- [23] K. Jodowski and S. Trojanowski, “Neutrino beam-dump experiment with FASER at the LHC,” *JHEP* **05**, 191 (2021) doi:10.1007/JHEP05(2021)191 [arXiv:2011.04751 [hep-ph]].
- [24] J. L. Feng, I. Galon, F. Kling and S. Trojanowski, “Dark Higgs bosons at the ForwArd Search ExpeRiment,” *Phys. Rev. D* **97** (2018) no.5, 055034 doi:10.1103/PhysRevD.97.055034 [arXiv:1710.09387 [hep-ph]].
- [25] F. Kling and S. Trojanowski, “Heavy Neutral Leptons at FASER,” *Phys. Rev. D* **97** (2018) no.9, 095016 doi:10.1103/PhysRevD.97.095016 [arXiv:1801.08947 [hep-ph]].
- [26] J. L. Feng, I. Galon, F. Kling and S. Trojanowski, “Axionlike particles at FASER: The LHC as a photon beam dump,” *Phys. Rev. D* **98** (2018) no.5, 055021 doi:10.1103/PhysRevD.98.055021 [arXiv:1806.02348 [hep-ph]].
- [27] F. Deppisch, S. Kulkarni and W. Liu, “Heavy neutrino production via  $Z'$  at the lifetime frontier,” *Phys. Rev. D* **100** (2019) no.3, 035005 doi:10.1103/PhysRevD.100.035005 [arXiv:1905.11889 [hep-ph]].
- [28] K. Cheung, C. J. Ouseph and T. Wang, “Non-standard neutrino and  $Z'$  interactions at the FASER $\nu$  and the LHC,” *JHEP* **12** (2021), 209 doi:10.1007/JHEP12(2021)209 [arXiv:2111.08375 [hep-ph]].

- [29] P. Bakhti, Y. Farzan and S. Pascoli, “Discovery potential of FASER $\nu$  with contained vertex and through-going events,” *JHEP* **04** (2021), 075 doi:10.1007/JHEP04(2021)075 [arXiv:2010.16312 [hep-ph]].
- [30] Y. Jho, J. Kim, P. Ko and S. C. Park, “Search for sterile neutrino with light gauge interactions: recasting collider, beam-dump, and neutrino telescope searches,” [arXiv:2008.12598 [hep-ph]].
- [31] N. Okada, S. Okada and Q. Shafi, “Light Z and dark matter from U(1) $X$  gauge symmetry,” *Phys. Lett. B* **810**, 135845 (2020) doi:10.1016/j.physletb.2020.135845 [arXiv:2003.02667 [hep-ph]].
- [32] M. Bahraminasr, P. Bakhti and M. Rajaei, “Sensitivities to secret neutrino interaction at FASER $\nu$ ,” *J. Phys. G* **48** (2021) no.9, 095001 doi:10.1088/1361-6471/ac11c2 [arXiv:2003.09985 [hep-ph]].
- [33] K. J. Kelly, M. Sen, W. Tangarife and Y. Zhang, “Origin of sterile neutrino dark matter via secret neutrino interactions with vector bosons,” *Phys. Rev. D* **101** (2020) no.11, 115031 doi:10.1103/PhysRevD.101.115031 [arXiv:2005.03681 [hep-ph]].
- [34] A. Falkowski, M. González-Alonso, J. Kopp, Y. Soreq and Z. Tabrizi, “EFT at FASER $\nu$ ,” *JHEP* **10** (2021), 086 doi:10.1007/JHEP10(2021)086 [arXiv:2105.12136 [hep-ph]].
- [35] A. Ariga *et al.* [FASER], “FASER’s physics reach for long-lived particles,” *Phys. Rev. D* **99** (2019) no.9, 095011 doi:10.1103/PhysRevD.99.095011 [arXiv:1811.12522 [hep-ph]].
- [36] G. Cottin, J. C. Helo, M. Hirsch, A. Titov and Z. S. Wang, “Heavy neutral leptons in effective field theory and the high-luminosity LHC,” *JHEP* **09**, 039 (2021) doi:10.1007/JHEP09(2021)039 [arXiv:2105.13851 [hep-ph]].
- [37] A. Ismail, S. Jana and R. M. Abraham, “Neutrino up-scattering via the dipole portal at forward LHC detectors,” *Phys. Rev. D* **105**, no.5, 055008 (2022) doi:10.1103/PhysRevD.105.055008 [arXiv:2109.05032 [hep-ph]].
- [38] K. Cheung and C. J. Ouseph, “Constraining the Active-to-Heavy-Neutrino transitional magnetic moments associated with the  $Z'$  interactions at FASER $\nu$ ,” [arXiv:2205.11077 [hep-ph]].
- [39] B. Batell, J. L. Feng and S. Trojanowski, “Detecting Dark Matter with Far-Forward Emulsion and Liquid Argon Detectors at the LHC,” *Phys. Rev. D* **103**, no.7, 075023 (2021) doi:10.1103/PhysRevD.103.075023 [arXiv:2101.10338 [hep-ph]].
- [40] M. Bauer, P. Foldenauer and J. Jaeckel, “Hunting All the Hidden Photons,” *JHEP* **07**, 094 (2018) doi:10.1007/JHEP07(2018)094 [arXiv:1803.05466 [hep-ph]].



- [41] B. Dutta, S. Ghosh, P. Huang and J. Kumar, “Explaining  $g\mu-2$  and  $RK(*)$  using the light mediators of  $U(1)T3R$ ,” *Phys. Rev. D* **105**, no.1, 015011 (2022) doi:10.1103/PhysRevD.105.015011 [arXiv:2105.07655 [hep-ph]].
- [42] P. Ballett, M. Hostert, S. Pascoli, Y. F. Perez-Gonzalez, Z. Tabrizi and R. Zukanovich Funchal, “ $Z$ 's in neutrino scattering at DUNE,” *Phys. Rev. D* **100**, no.5, 055012 (2019) doi:10.1103/PhysRevD.100.055012 [arXiv:1902.08579 [hep-ph]].
- [43] A. G. Beda, E. V. Demidova, A. S. Starostin, V. B. Brudanin, V. G. Egorov, D. V. Medvedev, M. V. Shirchenko and T. Vylov, “GEMMA experiment: Three years of the search for the neutrino magnetic moment,” *Phys. Part. Nucl. Lett.* **7** (2010), 406-409 doi:10.1134/S1547477110060063 [arXiv:0906.1926 [hep-ex]].
- [44] J. W. Chen, H. C. Chi, H. B. Li, C. P. Liu, L. Singh, H. T. Wong, C. L. Wu and C. P. Wu, “Constraints on millicharged neutrinos via analysis of data from atomic ionizations with germanium detectors at sub-keV sensitivities,” *Phys. Rev. D* **90** (2014) no.1, 011301 doi:10.1103/PhysRevD.90.011301 [arXiv:1405.7168 [hep-ph]].
- [45] H. B. Li *et al.* [TEXONO], “Limit on the electron neutrino magnetic moment from the Kuo-Sheng reactor neutrino experiment,” *Phys. Rev. Lett.* **90** (2003), 131802 doi:10.1103/PhysRevLett.90.131802 [arXiv:hep-ex/0212003 [hep-ex]].
- [46] G. Bellini, J. Benziger, D. Bick, S. Bonetti, G. Bonfini, M. Buizza Avanzini, B. Caccianiga, L. Cadonati, F. Calaprice and C. Carraro, *et al.* “Precision measurement of the  $^7\text{Be}$  solar neutrino interaction rate in Borexino,” *Phys. Rev. Lett.* **107** (2011), 141302 doi:10.1103/PhysRevLett.107.141302 [arXiv:1104.1816 [hep-ex]].
- [47] M. Deniz *et al.* [TEXONO], “Measurement of  $\text{Nu}(e)\text{-bar}$  -Electron Scattering Cross-Section with a CsI(Tl) Scintillating Crystal Array at the Kuo-Sheng Nuclear Power Reactor,” *Phys. Rev. D* **81** (2010), 072001 doi:10.1103/PhysRevD.81.072001 [arXiv:0911.1597 [hep-ex]].
- [48] L. B. Auerbach *et al.* [LSND], “Measurement of electron - neutrino - electron elastic scattering,” *Phys. Rev. D* **63** (2001), 112001 doi:10.1103/PhysRevD.63.112001 [arXiv:hep-ex/0101039 [hep-ex]].
- [49] P. Vilain *et al.* [CHARM-II], “Measurement of differential cross-sections for muon-neutrino electron scattering,” *Phys. Lett. B* **302** (1993), 351-355 doi:10.1016/0370-2693(93)90408-A