

Atmospheric axion-like particles at Super-Kamiokande

Kingman Cheung,^{1,2,3,*} Jui-Lin Kuo,^{4,†} Po-Yan Tseng,^{1,2,‡} and Zeren Simon Wang^{1,2,§}

¹*Department of Physics, National Tsing Hua University, Hsinchu 300, Taiwan*

²*Center for Theory and Computation, National Tsing Hua University, Hsinchu 300, Taiwan*

³*Division of Quantum Phases and Devices, School of Physics,*

Konkuk University, Seoul 143-701, Republic of Korea

⁴*Department of Physics and Astronomy, University of California, Irvine, CA 92697-4575, USA*

We consider a muonphilic axion-like-particle (ALP), denoted as a , lighter than twice the muon mass. ALPs of this mass range dominantly decay into a pair of photons, induced by a triangular muon loop. Such light muonphilic ALPs are naturally long-lived. At the atmosphere, the ALPs are copiously produced from charged-meson decays in air showers, such as $\pi^\pm \rightarrow \mu^\pm \nu a$, via the ALP-muon coupling $g_{a\mu\mu}$. After propagating tens of kilometers, the ALPs decay with $a \rightarrow \gamma\gamma$ inside large-volume Cherenkov detectors near the Earth surface, such as Super-Kamiokande (SK). We find the present SK observation constrains on muonphilic ALPs of mass range [1 MeV, 30 MeV] and ALP-muon coupling $[10^{-3}, 10^2]$, assuming the proper decay length $c\tau_a$ in $[10^{-3} \text{ km}, 10^6 \text{ km}]$ either correlated or uncorrelated with $g_{a\mu\mu}$. We conclude that atmospheric searches of such exotic states can be complementary to collider and beam-dump experiments as well as astrophysical probes.

I. INTRODUCTION

The strong CP problem [1–4] in the Standard Model (SM) can be solved by introducing a global $U(1)_{\text{PQ}}$ symmetry which was spontaneously broken down by a dynamical CP-conserving axion field. The corresponding pseudo-Nambu-Goldstone-boson of the broken symmetry is called the QCD axion, which in addition serves as a dark matter candidate [5–7]. The breaking scale of the new symmetry should be high: $f_a \gtrsim 10^9 \text{ GeV}$ [8], demanding tiny masses of the QCD axion and their couplings to the SM particles, since the latter two are inversely proportional to f_a . This results in a very long lifetime of the QCD axions.

A closely-related hypothetical particle is known as axion-like particle (ALP), which, like the QCD axion, is also a pseudoscalar boson. Unlike the QCD axion, the ALP mass is not linearly proportional to the couplings to the SM particles, and the ALP hence does not necessarily fix the strong CP problem. However, the ALP remains one of the possible dark matter candidates, and its mass could possibly range across more than 20 orders of magnitude [9–11]. Further, such ALPs appear in various theoretical models beyond the SM [12–14].

In general, the ALPs can couple to photons, leptons, quarks, as well as gauge bosons at either tree level or loop level. The phenomenology with only ALP-photon interaction $g_{a\gamma\gamma}$ has been vastly investigated (see Ref. [15] and the references therein). In particular, for sub-GeV ALPs, PRIMEX [16] and Belle II [17] provide the most stringent upper bounds on $|g_{a\gamma\gamma}|$, and ALPs of mass $m_a \lesssim 30 \text{ MeV}$ [15, 18] are disfavored by beam-dump experiments. However, for ALP-muon interactions, only BaBar [19]

gives constraints, for ALPs heavier than twice the muon mass [18]. As far as we know, muonphilic ALPs lighter than twice the muon mass have not been directly constrained. Therefore, we choose to focus on this scenario in the present work.

When cosmic rays reach the Earth’s atmosphere, large atmospheric air showers are produced including copious production of pseudoscalar mesons. Such mesons can decay to light long-lived particles (LLPs) (see Refs. [20–22] for reviews on LLPs), which travel macroscopic distances before decaying potentially in the large-volume neutrino experiments at the Earth surface. This allows to probe various models predicting such LLPs including heavy neutral leptons [23–28], the lightest neutralinos in the R-parity-violating supersymmetry [29], the hadrophilic light dark matter [30], and milli-charged particles [31]. Similarly, the muonphilic ALPs can be abundantly produced via charged-meson decays from the atmosphere air showers. Such ALPs should be long-lived, because both they are very light and their decay channels are radiatively suppressed if their mass is below twice the muon mass. After traveling tens of kilometers across the atmosphere, these ALPs may subsequently decay into two photons inside the detectors of neutrino experiments, such as Super-Kamiokande (SK). With the tool MCEq [32], we numerically compute the ALPs’ flux from the air showers including the propagation through dense medium. We then estimate the signal event rates at the SK detector, which is sensitive to events of energy below $\mathcal{O}(100) \text{ GeV}$ [33]. After discussing the background events, we obtain SK bounds on both physical observables and model parameters.

This article is organized as follows. In Sec. II, we introduce the theoretical scenario we investigate in this work. The estimation of the ALP flux from the air showers is detailed in Sec. III, followed by Sec. IV and Sec. V explaining the ALP detection on the Earth and introducing the SK experiment, respectively. The final numerical results are presented and discussed in Sec. VI. At the end,

* cheung@phys.nthu.edu.tw

† juilink1@uci.edu

‡ tpoyan1209@gmail.com

§ wzs@mx.nthu.edu.tw

Sec. VII provides a summary and outlook of this work.

II. ALP-MUON INTERACTION

In this work, we consider the interaction between the ALP and muon with an effective Lagrangian expressed as

$$\mathcal{L} \supset -ig_{a\mu\mu}a\bar{\mu}\gamma_5\mu, \quad (1)$$

where $g_{a\mu\mu}$ is a dimensionless coupling constant. For ALP mass m_a larger than twice the muon mass m_μ , the ALP can decay into a pair of muons, while for a lighter ALP only decays into a SM photon pair induced by a triangular muon loop. The loop-induced interaction between ALP and photons can be described by

$$\mathcal{L}_{\text{loop}} \supset -\frac{1}{4}g_{a\gamma\gamma}^{\text{eff}}aF^{\mu\nu}\tilde{F}_{\mu\nu}, \quad (2)$$

with the effective coupling being [34]

$$g_{a\gamma\gamma}^{\text{eff}} = \frac{g_{a\mu\mu}\alpha}{m_\mu\pi} \left[1 - \frac{4m_\mu^2}{m_a^2} \arcsin^2\left(\frac{m_a}{2m_\mu}\right) \right], \quad (3)$$

which is valid for $m_a \leq 2m_\mu$. The lifetime of ALP with $m_a < 2m_\mu$ then reads

$$\tau_a = \Gamma_{a \rightarrow \gamma\gamma}^{-1} = \frac{64\pi}{(g_{a\gamma\gamma}^{\text{eff}})^2 m_a^3}. \quad (4)$$

With the ALP-muon coupling, ALPs can be produced from charged-meson decays in air shower, dominated by the decay of charged pions $\pi^\pm \rightarrow \mu^\pm\nu a$; therefore, the kinematically allowed ALP mass range is $0 \leq m_a \leq m_\pi - m_\mu$ assuming massless SM neutrinos.

In Fig. 1, we present two Feynman diagrams for the production and decay of the ALPs, respectively.

The ALP-muon and ALP-photon couplings both contribute to the muon magnetic dipole moment, $a_\mu = (g-2)_\mu/2$ [18]. The one-loop result of $g_{a\mu\mu}$ leads to negative contributions to a_μ . However, if we also include the ALP-photon coupling, $g_{a\gamma\gamma}$, the two couplings will induce two-loop light-by-light and Barr-Zee diagrams [35, 36], which, in combination, provide positive contributions to a_μ [37]. On the experimental side, the updated combined results of Fermilab [38] and BNL [39] measurements indicate a 4.25σ positive deviation from the SM theoretical prediction:

$$\Delta a_\mu^{\text{BNL}} = a_\mu^{\text{BNL}} - a_\mu^{\text{SM}} = (251 \pm 59) \times 10^{-11}. \quad (5)$$

However, theoretical uncertainties arising from hadronic vacuum polarization may alleviate the tension between these measurements and the SM [40]. Given the large uncertainties within the SM computation, we do not take into account $(g-2)_\mu$ in our analysis.

III. ALP FLUX FROM AIR SHOWER

We utilize the numerical code MCEq [32] to compute the ALP flux at the Earth surface. MCEq numerically solves cascade equations of particles propagating in a dense medium; in this work, we use it to study the ALP production throughout the cascade of secondary cosmic rays. We adopt the parameterization of the cosmic ray flux at the top of atmosphere provided in [41] and take the hadronic interaction model in [42]. The atmosphere is modeled by the CORSIKA parameterizations [43].

To implement the process $\pi^\pm \rightarrow \mu^\pm\nu a$ in MCEq, we compute the corresponding decay matrix

$$D_{\pi^\pm \rightarrow a}^{ij} = \Delta T_{\pi^\pm}^i \frac{dN_a}{dT_a}(T_{\pi^\pm}^i, T_a^j), \quad (6)$$

where T_{π^\pm} and T_a are the kinetic energy of the pion and the ALP in the lab frame, with i, j and $\Delta T_{\pi^\pm}^i$ being the kinetic energy bin indices and width, respectively. The ALP energy spectrum dN_a/dT_a in the lab frame is obtained by applying a Lorentz boost to the energy spectrum in the pion rest frame

$$\frac{dN_a}{dT_a} = \int \frac{d\Omega}{4\pi} \frac{dN_a}{dE_a^*} \left| \frac{\partial E_a^*}{\partial T_a} \right|, \quad (7)$$

where E_a^* is the energy of the ALP in the pion rest frame and $|\partial E_a^*/\partial T_a|$ is the Jacobian between E_a^* and T_a .

After the decay matrix is tabulated, we augment the decay channels of π^\pm with $\pi^\pm \rightarrow \mu\nu a$. We first consider the case that production and the subsequent decay of the ALP into two photons are uncorrelated for simplicity, *i.e.*, the ALP flux at production is proportional to $g_{a\mu\mu}^2$ while the decay is determined by the decay length $c\tau_a$ in the ALP rest frame. The results are model-independent in the sense that they can be mapped onto another model with similar decay topologies. Then, we study the case that both production and decay depend on the coupling constant $g_{a\mu\mu}$.

IV. ALP DETECTION ON THE EARTH

After arriving at the Earth, the ALP can decay into two photons through a muon loop, with a lifetime given in Eq. (4). The photons so-produced can then be detected by the Cherenkov detector in neutrino experiments. Given zenith angle θ , detector geometry, and data-taking time Δt , the event rate can be calculated by

$$\frac{dN_{\text{event}}}{dT_a} = \epsilon \Delta t A_{\text{eff}} \frac{d\Phi_a}{dT_a}, \quad (8)$$

where ϵ is the detection efficiency and we use the output of MCEq for the differential flux $d\Phi_a/dT_a$. The effective decay area A_{eff} , depending on T_a , θ and $c\tau_a$, is given in the appendix of Ref. [26]. The main SM background of such a two-photon signal from the ALP decay




FIG. 1. *Left panel:* Feynman diagram for production of the muonphilic ALP from the charged-pion decay. The decay vertex factor $g_{\pi\mu\nu}$ denotes the effective coupling for the charged pion decay $\pi^+ \rightarrow \mu^+\nu_\mu$. Its conjugated diagram with π^- decays is not shown here. *Right panel:* Feynman diagram for the decay of the muonphilic ALP into a pair of photons, via a triangular muon loop.

consists of neutral pions decaying into two photons, and neutrino-induced electron-like events that create multiple Cherenkov rings in the electromagnetic shower. The number of these events have been studied in Ref. [33], with the best-fit values being 1727 and 797, respectively. These background events will be taken into account in Sec. VI when we estimate the sensitivity reach of SK.

In addition to the signals from ALP decaying into a γ -pair, ALP can interact with atoms in the detector to create mono- γ signal with an energy similar to the energy of ALP, the so-called inverse-Primakoff process. The cross section of inverse-Primakoff process was studied in details in [44], which can be expressed as

$$\sigma_{\text{IP}} \simeq \left(\frac{g_{a\gamma\gamma}^{\text{eff}}}{1 \text{ GeV}^{-1}} \right)^2 \times 2 \text{ GeV}^{-2}. \quad (9)$$

However, since A_{eff} of decay is larger by orders of magnitude than the effective cross section of inverse-Primakoff process $N_T \sigma_{\text{IP}}$ with N_T being the total number of target atoms inside the fiducial volume of the detector, we can infer that the event rate from the ALP decay dominates over that from the inverse-Primakoff process; therefore, we will not consider this possibility further in this work.

V. SUPER-KAMIOKANDE

We note that only when the charged pions have a kinetic energy below $\mathcal{O}(\text{TeV})$ and hence a small Lorentz boost, do they essentially all decay well before reaching the Earth surface; therefore, ALP flux at $T_a > \mathcal{O}(\text{TeV})$ is strongly suppressed. In order to maximize the sensitivity, we focus on the water-based Cherenkov detector of Super-Kamiokande (SK), which has good energy resolution in the sub- and multi-GeV ranges [33].

Following the analysis in Ref. [33], the geometry of the SK detector is assumed to be a cylinder with radius

$R_{\text{SK}} = 20 \text{ m}$ and height $H_{\text{SK}} = 40 \text{ m}$. The lifetime of SK is taken to be 5326 days with a flat detection efficiency of 0.75. Fully-contained events in SK can be grouped into different categories according to the energy and configuration of observed Cherenkov rings. Since the signal from ALP decay constitutes two electron-like Cherenkov rings, we consider data of π^0 -like two-ring events in 5 energy-bins for sub-GeV T_a and electron-like multi-ring events in 5 $\cos\theta$ -bins for multi-GeV T_a provided in Ref. [33].

VI. CONSTRAINTS ON THE PARAMETER SPACE

We perform χ^2 -fit to the SK data mentioned in Sec. V using [26]

$$\chi_i^2 = 2 \left\{ N_{\text{sig}}^i + N_{\text{bkg}}^i - N_{\text{obs}}^i \left[1 - \log \left(\frac{N_{\text{obs}}^i}{N_{\text{sig}}^i + N_{\text{bkg}}^i} \right) \right] \right\}, \quad (10)$$

where N_{sig}^i , N_{bkg}^i , and N_{obs}^i are numbers of expected signal events of ALP, background events, and observed events in each data bin, respectively. The background and observed events are extracted from Ref. [33]. For a total of 10 bins considered in this work, we derive the 90% C.L. constraint by requiring $\Delta\chi^2 \equiv \chi^2 - \chi_0^2 \leq 4.865$ with $\chi^2 = \sum_i \chi_i^2$ and χ_0^2 being the case without ALP contribution.

We show the sensitivity reach of SK in Fig. 2 for $g_{a\mu\mu}$ - $c\tau_a$ uncorrelated (solid lines in the left panel) and correlated (blue area in the right panel) cases. In the left panel, we observe that for $m_a = 1 \text{ MeV}$, the best sensitivity of SK lies at $c\tau_a \sim 5 \times 10^{-2} \text{ km}$, while for larger m_a , the best sensitivity of SK is reached at larger $c\tau_a$. In the same plot we also overlap the uncorrelated sensitivity curves with dashed lines depicting $c\tau_a$ as a function of $g_{a\mu\mu}$ using Eq. (4). This allows us to crosscheck with the




FIG. 2. *Left panel*: sensitivity reach of SK to the muonphilic ALPs for uncorrelated $g_{a\mu\mu}$ and $c\tau_a$ (solid curves), and $c\tau_a$ as a function of $g_{a\mu\mu}$ according to Eq. (4) (dashed lines) in the $(c\tau_a, g_{a\mu\mu})$ plane, for three benchmark values of m_a . *Right panel*: constraints on $(m_a, g_{a\mu\mu})$ assuming $c\tau_a$ is correlated with $g_{a\mu\mu}$. For comparison, we also include the constraint from BaBar which holds only for larger m_a [19], and the bounds from SN1987A which cover $g_{a\mu\mu} \sim [10^{-10}, 5 \times 10^{-3}]$ for sub-GeV m_a [45].

right panel where we assume both production and decay are mediated by $g_{a\mu\mu}$; the intersection of the solid and dashed lines in the left plot for each fixed mass should coincide with the corresponding parameter point on the outer edge of the blue area in the $g_{a\mu\mu}$ vs. m_a plane in the right panel. We find in the right plot that for $m_a = [0.1, 33]$ MeV, SK can exclude $g_{a\mu\mu} = [5 \times 10^{-3}, 50]$, comparable to the BaBar exclusion limits, which, however, only hold for larger m_a [19]. Below $g_{a\mu\mu} \sim 5 \times 10^{-3}$ and for sub-GeV m_a , we note the parameter space is covered by the SN1987A constraint [45, 46]. For each mass value within the sensitive range, the SK exclusion limits are bounded from both top and bottom. This is because when $g_{a\mu\mu}$ is too small, the production rate of the ALPs is insufficient and the decay length is too long. On the other hand, with a too large $g_{a\mu\mu}$, despite the enhanced production rate, the decay length is so short that the ALPs decay before reaching the SK detector. We note that future muon beam-dump experiments can improve the sensitivity at the ALP mass range of interest in this work down to around $g_{a\mu\mu} \sim 6 \times 10^{-6}$; see Ref. [47] for details. The results presented here are only based on charged pion flux. One can in principle extend the constraint to $m_a \geq m_{\pi^\pm} - m_\mu$ by further including heavier mesons such as kaons that have a similar decay channel. However, once $m_a \geq 2m_\mu$ is fulfilled, the ALPs can decay into two muons, resulting in further complication during the air shower; we reserve this possibility for future work.

In addition to ALP-muon interaction, we have checked the case that the ALP has a direct coupling to the SM photons. In this case, the main ALP production chan-

nel in air shower is π^0 decay, with a smaller branching ratio compared to charged pion decay as a result of the much shorter lifetime of π^0 . Therefore, we infer that given a fixed decay rate of $\pi \rightarrow a$, for $T_a < \mathcal{O}(\text{TeV})$ the ALP flux from π^0 is less intense than that from π^\pm , but for higher masses the ALP flux from π^0 dominates, as π^\pm 's will not all decay well before reaching the Earth surface. We find that this excludes $g_{a\gamma\gamma} \gtrsim 10^{-2} \text{ GeV}^{-1}$ for $m_a \sim \mathcal{O}(1\text{--}100 \text{ MeV})$, assuming the decay of ALP is independently determined by $c\tau_a$. Since $g_{a\gamma\gamma} \gtrsim 10^{-2} \text{ GeV}^{-1}$ is already excluded by accelerator and collider constraints (see Ref. [48] for a recent summary), we do not demonstrate the result in this work. In addition, we note that in this scenario the case that the production and the decay are correlated cannot be probed by SK, since tree-level decay results in a much shorter $c\tau_a$ for the given couplings; see also other scenarios with tree-level decays discussed in Ref. [26].

VII. CONCLUSIONS

Similar to QCD axions, axion-like-particles (ALPs), denoted as a in this work, are also among the most plausible candidates of dark matter. With its mass and couplings to the SM particles uncorrelated, such exotic particles are being searched for at various experimental facilities across a wide range of masses. While the ALPs can in theory couple to various types of particles, here, we have chosen to focus on the case that the ALPs are dom-

inantly or solely interacting with the SM muons at tree level. In addition, we have restricted ourselves to ALP mass below the muon-pair threshold so that the ALPs only decay radiatively into a pair of photons. Such ALPs can be produced from charged pion decays, $\pi^\pm \rightarrow \mu^\pm \nu a$ via the ALP-muon coupling $g_{a\mu\mu}$.

Large numbers of mesons including charged pions are produced in the atmospheric air showers resulting from cosmic rays. Once ALPs are produced from these charged-pion decays, if long-lived, they can travel tens of kilometers downwards to the Earth surface thanks to the large Lorentz boost, and decay in large-volume neutrino experiments such as Super-Kamiokande (SK), leading to Cherenkov signal events.

We make use of the numerical tool MCEq in order to estimate the ALP flux at the Earth surface stemming from cosmic-ray-induced atmospheric air showers of the charged pions. We further compute the signal event rates, taking into account the differential ALP flux, detector efficiency, data-taking time, etc., at SK. In addition, there are background events mainly stemming from neutral pion decays into two photons as well as neutrino-induced electron-like events that lead to multiple Cherenkov rings in the electromagnetic shower. We have extracted the level of these background events from Ref. [33], and evaluated the sensitivity reach of the SK experiments to such muonphilic ALPs. Results are presented for both production-decay correlated and uncorrelated cases, shown in Fig. 2. In particular, we find that if both production and decay are mediated by the coupling $g_{a\mu\mu}$, SK can probe $g_{a\mu\mu}$ down to 5×10^{-3} at $m_a \sim 20$ MeV, complementary to the exclusion limits obtained at BaBar which is sensitive to larger masses, as well as SN1987A constraint which covers $g_{a\mu\mu} \lesssim 5 \times 10^{-3}$.

Additionally, we have commented on further possibilities such as the inverse-Primakoff process, and the case that the ALP is coupled to the SM photons at tree level. While the former is expected to be dominated over by the main process considered in this work, the latter is checked to give only rather weak limits that have already been excluded by past experiments.

We have also briefly discussed the implication of the anomalous muon magnetic moment measurements on our model. Based on the current sensitivity of Fermilab $(g - 2)_\mu$ measurement and once theoretical uncertainties are clarified, $|g_{a\mu\mu}| \gtrsim 5 \times 10^{-4}$ can be probed for correlated $g_{a\gamma\gamma} - g_{a\gamma\gamma}$ and $m_a \leq 2m_\mu$. However, with this coupling correlation, the ALPs always contribute negatively to a_μ . Therefore, our model is unable to alleviate the present tension between the SM prediction and observation results.

Before closing, we comment on the sensitivities of other present and future neutrino telescopes to the scenario considered in this work. For the future Hyper-Kamiokande experiment (HK) [49, 50], its fiducial volume is increased by a factor 25 compared to SK. Therefore, we naively estimate that it can improve the sensitivity reach to $g_{a\mu\mu}$ by a factor $\sqrt{5}$ ($5^{1/4}$) for the uncorre-

lated (correlated) case. We note that the precise sensitivity reach ultimately depends on the detector configuration of HK. Moreover, if the signal discrimination rate is also enhanced, we expect the sensitivity reach could potentially enclose a large portion of the parameter space currently covered by the SN1987A constraint. Finally, we note that in principle IceCube [51] is also capable of probing the atmospheric ALP flux considered in this work. However, since IceCube focuses on the ultrahigh energy range, the best sensitivity lies at $c\tau_a \sim 5 \times 10^{-5}$ km for $m_a \sim 10$ MeV [26]; therefore, with a smaller $c\tau_a$ needed, we can expect that the constraint on $g_{a\mu\mu}$ will be much weaker, despite a much larger fiducial volume at IceCube.

ACKNOWLEDGMENT

We thank Giovanna Cottin, Felix Kling, and Víctor Muñoz for useful discussions. Z.S.W. is supported by the Ministry of Science and Technology (MoST) of Taiwan with grant number MoST-110-2811-M-007-542-MY3. J.L.K. is supported by the U.S. National Science Foundation (NSF) Theoretical Physics Program, Grant PHY-1915005. P.Y.T. and K.C. are supported in part by MoST with grant numbers MoST-111-2112-M-007-012-MY3 and MoST-110-2112-M-007-017-MY3, respectively.

-
- [1] R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38**, 1440 (1977).
- [2] R. D. Peccei and H. R. Quinn, Phys. Rev. D **16**, 1791 (1977).
- [3] S. Weinberg, Phys. Rev. Lett. **40**, 223 (1978).
- [4] F. Wilczek, Phys. Rev. Lett. **40**, 279 (1978).
- [5] J. Preskill, M. B. Wise, and F. Wilczek, Phys. Lett. B **120**, 127 (1983).
- [6] L. F. Abbott and P. Sikivie, Phys. Lett. B **120**, 133 (1983).
- [7] M. Dine and W. Fischler, Phys. Lett. B **120**, 137 (1983).
- [8] J. L. Feng, T. Moroi, H. Murayama, and E. Schnapka, Phys. Rev. D **57**, 5875 (1998), arXiv:hep-ph/9709411.
- [9] J. E. Kim and D. J. E. Marsh, Phys. Rev. D **93**, 025027 (2016), arXiv:1510.01701 [hep-ph].
- [10] I. De Martino, T. Broadhurst, S. H. Henry Tye, T. Chiueh, H.-Y. Schive, and R. Lazkoz, Phys. Rev. Lett. **119**, 221103 (2017), arXiv:1705.04367 [astro-ph.CO].
- [11] V. A. Rubakov, JETP Lett. **65**, 621 (1997), arXiv:hep-ph/9703409.
- [12] P. Svrcek and E. Witten, JHEP **06**, 051 (2006), arXiv:hep-th/0605206.
- [13] G. B. Gelmini and M. Roncadelli, Phys. Lett. B **99**, 411 (1981).
- [14] F. Wilczek, Phys. Rev. Lett. **49**, 1549 (1982).
- [15] P. Agrawal *et al.*, Eur. Phys. J. C **81**, 1015 (2021), arXiv:2102.12143 [hep-ph].
- [16] D. Aloni, C. Fanelli, Y. Soreq, and M. Williams, Phys. Rev. Lett. **123**, 071801 (2019), arXiv:1903.03586 [hep-ph].
- [17] F. Abudinén *et al.* (Belle-II), Phys. Rev. Lett. **125**, 161806 (2020), arXiv:2007.13071 [hep-ex].
- [18] M. A. Buen-Abad, J. Fan, M. Reece, and C. Sun, JHEP **09**, 101 (2021), arXiv:2104.03267 [hep-ph].
- [19] J. P. Lees *et al.* (BaBar), Phys. Rev. D **94**, 011102 (2016), arXiv:1606.03501 [hep-ex].
- [20] D. Curtin *et al.*, Rept. Prog. Phys. **82**, 116201 (2019), arXiv:1806.07396 [hep-ph].
- [21] L. Lee, C. Ohm, A. Soffer, and T.-T. Yu, Prog. Part. Nucl. Phys. **106**, 210 (2019), [Erratum: Prog.Part.Nucl.Phys. 122, 103912 (2022)], arXiv:1810.12602 [hep-ph].
- [22] J. Alimena *et al.*, J. Phys. G **47**, 090501 (2020), arXiv:1903.04497 [hep-ex].
- [23] A. Kusenko, S. Pascoli, and D. Semikoz, JHEP **11**, 028 (2005), arXiv:hep-ph/0405198.
- [24] T. Asaka and A. Watanabe, JHEP **07**, 112 (2012), arXiv:1202.0725 [hep-ph].
- [25] M. Masip, AIP Conf. Proc. **1606**, 59 (2015), arXiv:1402.0665 [hep-ph].
- [26] C. Argüelles, P. Coloma, P. Hernández, and V. Muñoz, JHEP **02**, 190 (2020), arXiv:1910.12839 [hep-ph].
- [27] P. Coloma, P. Hernández, V. Muñoz, and I. M. Shoemaker, Eur. Phys. J. C **80**, 235 (2020), arXiv:1911.09129 [hep-ph].
- [28] S. Meighen-Berger, M. Agostini, A. Ibarra, K. Krings, H. Niederhausen, A. Rappelt, E. Resconi, and A. Turcati, Phys. Lett. B **811**, 135929 (2020), arXiv:2005.07523 [hep-ph].
- [29] P. Candia, G. Cottin, A. Méndez, and V. Muñoz, Phys. Rev. D **104**, 055024 (2021), arXiv:2107.02804 [hep-ph].
- [30] C. A. Argüelles, V. Muñoz, I. M. Shoemaker, and V. Takhistov, (2022), arXiv:2203.12630 [hep-ph].
- [31] C. A. Argüelles Delgado, K. J. Kelly, and V. Muñoz Albornoz, (2021), arXiv:2104.13924 [hep-ph].
- [32] A. Fedynitch, R. Engel, T. K. Gaisser, F. Riehn, and T. Stanev, EPJ Web Conf. **99**, 08001 (2015), arXiv:1503.00544 [hep-ph].
- [33] K. Abe *et al.* (Super-Kamiokande), Phys. Rev. D **97**, 072001 (2018), arXiv:1710.09126 [hep-ex].
- [34] M. Bauer, M. Neubert, and A. Thamm, JHEP **12**, 044 (2017), arXiv:1708.00443 [hep-ph].
- [35] E. J. Chun, EPJ Web Conf. **118**, 01006 (2016), arXiv:1511.05225 [hep-ph].
- [36] W. J. Marciano, A. Masiero, P. Paradisi, and M. Passera, Phys. Rev. D **94**, 115033 (2016), arXiv:1607.01022 [hep-ph].
- [37] W.-Y. Keung, D. Marfatia, and P.-Y. Tseng, LHEP **2021**, 209 (2021), arXiv:2104.03341 [hep-ph].
- [38] B. Abi *et al.* (Muon $g-2$), Phys. Rev. Lett. **126**, 141801 (2021), arXiv:2104.03281 [hep-ex].
- [39] G. W. Bennett *et al.* (Muon $g-2$), Phys. Rev. D **73**, 072003 (2006), arXiv:hep-ex/0602035.
- [40] S. Borsanyi *et al.*, Nature **593**, 51 (2021), arXiv:2002.12347 [hep-lat].
- [41] T. K. Gaisser, Astropart. Phys. **35**, 801 (2012), arXiv:1111.6675 [astro-ph.HE].
- [42] F. Riehn, H. P. Dembinski, R. Engel, A. Fedynitch, T. K. Gaisser, and T. Stanev, PoS **ICRC2017**, 301 (2018), arXiv:1709.07227 [hep-ph].
- [43] D. Heck, J. Knapp, J. N. Capdevielle, G. Schatz, and T. Thouw, (1998).
- [44] T. Abe, K. Hamaguchi, and N. Nagata, Phys. Lett. B **815**, 136174 (2021), arXiv:2012.02508 [hep-ph].
- [45] D. Croon, G. Elor, R. K. Leane, and S. D. McDermott, JHEP **01**, 107 (2021), arXiv:2006.13942 [hep-ph].
- [46] R. Bollig, W. DeRocco, P. W. Graham, and H.-T. Janka, Phys. Rev. Lett. **125**, 051104 (2020), [Erratum: Phys.Rev.Lett. 126, 189901 (2021)], arXiv:2005.07141 [hep-ph].
- [47] C.-Y. Chen, M. Pospelov, and Y.-M. Zhong, Phys. Rev. D **95**, 115005 (2017), arXiv:1701.07437 [hep-ph].
- [48] D. d’Enterria, in *Workshop on Feebly Interacting Particles* (2021) arXiv:2102.08971 [hep-ex].
- [49] K. Abe *et al.*, (2011), arXiv:1109.3262 [hep-ex].
- [50] E. Kearns *et al.* (Hyper-Kamiokande Working Group), in *Community Summer Study 2013: Snowmass on the Mississippi* (2013) arXiv:1309.0184 [hep-ex].
- [51] M. G. Aartsen *et al.* (IceCube), JINST **12**, P03012 (2017), arXiv:1612.05093 [astro-ph.IM].