

## Van der Waals type PV diagrams of PNJL matter

Manjeet Seth<sup>1,\*</sup>, Kinkar Saha<sup>2,†</sup>, Sudipa Upadhaya<sup>3</sup>, Soumitra Maity<sup>4</sup>, and Sabyasachi Ghosh<sup>1</sup>

<sup>1</sup>Indian Institute of Technology Bhilai, GEC Campus,  
Sejbahar, Raipur - 492015, Chhattisgarh, India

<sup>2</sup>Uluberia College, Howrah, West Bengal, India, 711315

<sup>3</sup>Ramsaday College, Howrah, West Bengal, India, 711401 and

<sup>4</sup>Bose Institute, Block-EN, Sector-V, Salt Lake City, Kolkata, West Bengal, India, 700091

Study of the systems produced in high-energy heavy ion collision experiments are of immense interest for various reasons. Here we intend to take an attempt to discuss the presence of interactions in such systems and their effects to distinguish them from the ideal scenario under the framework of Polyakov–Nambu–Jona-Lasinio model. This model encapsulates two very important aspects of Quantum Chromo Dynamics (QCD), viz. the chiral and the confinement-deconfinement phase transitions. With the help of  $T$  and  $\mu$  dependent constituent quark mass  $M(T, \mu)$ , one can calculate net quark number density  $n(T, \mu)$ , pressure  $P(T, \mu)$ :

$$n(T, \mu) = g \int \frac{d^3p}{(2\pi)^3} (f_0^+ - f_0^-)$$

$$P(T, \mu) = g \int \frac{d^3p}{(2\pi)^3} \left( \frac{p^2}{3E} \right) (f_0^+ + f_0^-), \quad (1)$$

where  $g = 3 \times 3 \times 2 = 18$  is degeneracy factor of quark or anti-quark with energy  $E = \sqrt{p^2 + M^2(T, \mu)}$  and  $f_0^\pm = 1/\{e^{\beta(E \mp \mu)} + 1\}$  for NJL model case and for PNJL model, we have to replaced by modified distribution function [1].

Now let us go for rough estimation number of hadrons or quarks, which can be produced in heavy ion collision experiments with large range of  $\sqrt{s} = 1 - 200$  GeV. Considering only pion production from this energy, one can get rough no  $\frac{(1-200)}{0.14} \approx 7 - 1, 428$  pions and  $14 - 2, 856 \approx 10^{1-3}$ . So if we want to restrict 100




FIG. 1: Pressure as a function of volume at various characteristic temperatures by using PNJL model.

net quark by varying volume from  $V = 1 \text{ fm}^3$  to  $100 \text{ fm}^3$  with number density  $n = 100 - 1/\text{fm}^3$ . Now for 3 temperatures  $T = 0.400$  GeV,  $0.200$  GeV,  $0.120$  GeV, fixing net quark number as

$$N = n(T, \mu) \times V = 100$$

$$Vg \int \frac{d^3p}{(2\pi)^3} (f_0^+ - f_0^-) = 100, \quad (2)$$

Fig.(1) shows the dependence of Pressure,  $P$  on the volume,  $V$  of the system, where  $V$  is obtained from Eqn.(2) as function of chemical potential,  $\mu$ . In order to obtain the con-

\*Electronic address: manjeetseth28@gmail.com

†Electronic address: saha.k.09@gmail.com




FIG. 2: Van Der Waal like Plots

stituent masses from PNJL model [5, 6, 7] and thereby the volume,  $V$  from Eqn.(2) at various characteristic temperatures, we have varied  $\mu$ . The Pressure,  $P$  in Fig.(1) show non-monotonic behaviour with  $V$ . The results are portrayers of the quantum level interactions present in the system among the associated degrees of freedom. Also the modified potential in Polyakov loop part, mimicking to some extent the interactions present among gluonic degrees of freedom in Lattice QCD [2], also has a considerable effect [comparison with NJL]. From any one isotherm below  $T=160$  MeV, we can get the idea of the corresponding  $\mu$  where a transition happens indicating a critical point on the phase diagram, using the  $V$ - $\mu$  relation. The transition point is signified by the minima present in the corresponding isotherms. For  $T>160$  MeV, any such minima are absent as the system then resides entirely in the QGP region no matter what the chemical potential is. This result is quite commensurate with the QCD phase diagram and hence  $T\sim 160$ MeV can be considered as the cross-over transition temperature at  $\mu = 0$ , as extracted from the model study. This agrees quite satisfactorily with Lattice QCD results at continuum limit [3, 4] and as predicted by our earlier study [2, 5, 6, 7].

The results appear simpler to explain if we look for isotherms with reduced Pressure,  $P_r$

and reduced Volume,  $V_r$  in Fig.(2) as was done by Van Der Waal. The presence of non-monotonicities below  $T_r=1$  signifies the interactions present causing transitions occurring in the system at particular chemical potential for one characteristic temperature. These critical temperatures and chemical potentials fall on the phase diagram shown in Ref. [2] for six-quark interactions. Below  $T_r=1$ , the isotherms therefore represent various states which could be unstable or metastable. Above  $T_r=1$ , the system comprises of quarks only and hence possibility of phase transition does not occur. Similar results had also been shown in Ref. [8], for normal nuclear matter with corrections from Quantum Statistical effects.

## References

- [1] Sanjay K. Ghosh, Tamal K. Mukherjee, Munshi Golam Mustafa, and Rajarshi Ray, Phys. Rev. D77, 094024 (2008).
- [2] Abhijit Bhattacharyya, Sanjay K. Ghosh, Soumitra Maity, Sibaji Raha, Rajarshi Ray, Kinkar Saha, and Sudipa Upadhaya, Phys. Rev. D95, 054005 (2017).
- [3] S. Borsanyi et al., Phys. Lett. B730, 99 (2014).
- [4] A. Bazavov et al., Phys. Rev. D90, 094503 (2014).
- [5] Sanjay K. Ghosh, Sibaji Raha, Rajarshi Ray, Kinkar Saha, and Sudipa Upadhaya, Phys. Rev. D91, 054005 (2015).
- [6] Kinkar Saha, Sudipa Upadhaya, and Sabyasachi Ghosh, Mod. Phys. Lett. A32 05, 1750018 (2016).
- [7] Abhijit Bhattacharyya, Sanjay K. Ghosh, Soumitra Maity, Sibaji Raha, Rajarshi Ray, Kinkar Saha, Subhasis Samanta, and Sudipa Upadhaya, Phys. Rev. C99, 045207 (2019).
- [8] S.N. Fedotkin, A.G. Magner, and U.V. Grygoriev, arXiv:2012.09695 [nucl-th]