Heterogeneity assessment in causal data fusion problems

Tat-Thang Vo¹, Kara E. Rudolph², and Iván Díaz³

¹Department of Statistics and Data Science, The Wharton School, University of Pennsylvania, USA

²Department of Epidemiology, Mailman School of Public Health, Columbia University, USA

³Division of Biostatistics, Department of Population Health Sciences, Weill Cornell Medicine, USA

August 12, 2022

Abstract

Previous works have formalized the conditions under which findings from a source population could be reasonably extrapolated to another target population, the so-called "transportability" problem. While most of these works focus on a setting with two populations, many recent works have also provided the identifiability of a causal parameter when multiple data sources are available, under certain homogeneity assumptions. However, we know of little work examining transportability when data sources are possibly heterogeneous, e.g. in the distribution of mediators of the exposure-outcome relation. The presence of such heterogeneity generally invalidates the transportability assumption required in most of the literature. In this paper, we will propose a general approach for heterogeneity assessment when estimating the average exposure effect in a target population, with mediator and outcome data obtained from multiple external sources. To account for heterogeneity, we define different effect estimands when the mediator and outcome information is transported from different sources. We discuss the causal assumptions to identify these estimands, then propose efficient semi-parametric estimation strategies that allow the use of flexible data-adaptive machine learning methods to estimate the nuisance parameters. We also propose two new methods to investigate sources of heterogeneity in the transported estimates. These methods will inform users about how much of the observed statistical heterogeneity in the transported effects is due to the differences across data sources in: 1) conditional distribution of mediator variables, and/or 2) conditional distribution of the outcome. We illustrate the proposed methods using four sites that were part of the Moving to Opportunity Study, which was an experiment that randomized housing voucher receipt to participating families living in public housing.

1 Introduction

The integration of data and knowledge from several data sources is often known as data fusion (Bareinboim and Pearl, 2016). Across different research fields, data fusion methods are increasingly needed to satisfy the requirement of combining information from multiple databases, so as to improve our ability to perform inferences from data. Although the ideas behind data fusion have been around for quite long, the first formal frameworks for this concept have just been recently developed. Bareinboim and Pearl (2016), for instance, formalize the conditions under which causal findings from a source population could be reasonably extrapolated to another target population. This framework serves as a theoretical base for many data fusion methods such as covariate shift (Uehara *and others*, 2020), selection bias adjustment (Ferri-García and Rueda, 2020), external validity (Stuart *and others*, 2011), randomized and observational data combination (Colnet *and others*, 2020), transported indirect effects (Rudolph *and others*, 2021), causally interpretable evidence synthesis and heterogeneity assessment (Vo *and others*, 2019; Dahabreh *and others*, 2020). While most of these methods focus on a setting with two populations, one source and one target,

many recent works have also provided the identifiability of a causal parameter when multiple data sources are available. See for instance, the work by Sobel *and others* (2017); Vo *and others* (2019, 2021); Bareinboim and Pearl (2016); Dahabreh *and others* (2019); Li and Luedtke (2021).

Existing data fusion approaches require assuming that certain portion of the data generating mechanism (e.g. the outcome model) is homogeneous across populations. This assumption is often used to justify merging all individual-level data across data sources, then using these to provide an estimate of the missing distribution in the target population (Dahabreh *and others*, 2019; Li and Luedtke, 2021). However, important heterogeneity may exist among data sources, for instance, due to the difference in settings or background conditions, in versions of interventions and controls being assessed, in the mechanisms by which the interventions exert effects, or in outcome measurement. Depending on the data setting/structural causal model and estimand of interest, the presence of such heterogeneity may violate the transportability assumption(s) required for identification, and may induce bias if estimation proceeds anyway. Assessing and incorporating this heterogeneity is therefore of critical importance in causal data fusion problems.

In this paper, we propose a general approach for heterogeneity assessment when combining data from multiple data sources. We assume the presence of heterogeneity across databases. Instead of directly fusing data from different sources, we use data from each source separately to estimate the causal parameter of interest in the target population. We then develop two methods to quantify how much of the variability observed among different transported estimates can be explained by each source of heterogeneity across databases. For instance, when estimating the average exposure effect in a target population by using the intermediate variables and outcome data from elsewhere, we measure how much of the variability among the transported estimates can be explained by the difference between populations in conditional intermediate variables distribution and in conditional outcome distribution. Such a heterogeneity decomposition allows one to assess the extent to which the average exposure effect estimates obtained by using data from different sources are homogeneous (so that they can be combined to derive a single summary estimate for the target parameter). We show that transporting effects separately and then fusing them, or a subset of then, when warranted, instead of fusing the individual-level data avoids possible identification biases at the expense of estimation variance.

The paper is structured as follow. In Section 2, we discuss the importance of heterogeneity assessment when using data from multiple sources to estimate the average exposure effect in a target population. In Section 3, we propose two novel approaches to investigate the impact of heterogeneity between these data sources on the transported effect estimates. We evaluate the finite-sample performance of these approaches via a simulation study in Section 4, and apply them to analyze the data from four sites that were part of the Moving to Opportunity Study in Section 5. Specifically, we decompose heterogeneity in estimating the average causal effect of moving with a Section 8 housing voucher on risk of children subsequently developing a psychiatric disorder, with neighborhood poverty being the mediator. In section 6, we briefly discuss the generalization of the proposed framework to a multiple mediator setting. We end this paper by some final discussions and suggestions for future research.

2 Data fusion in the context of multiple heterogeneous data sources

2.1 Setting and notations

Suppose that the aim of the analysis is to investigate the causal effect of a binary exposure X on a binary outcome Y in a target population S = j, where S is a categorical variable denoting the population. In S = j, we have data on a vector of baseline covariates L, but no data on X and Y because there are not yet studies conducted to assess the $X \to Y$ relationship. The information on (L, X, Y) is then obtained from a set of external sources S, where $j \notin S$. In these external sources, we further observe a vector of intermediate variables $M = (M_1, \ldots, M_Z)$ that partially mediates the effect of X on Y. Toward the end of this Section, we will also discuss the setting in which only data on (X, M) are available in some additional external data sources. Note that data on M are not available in the target population S = j. For each individual *i*, we thus observe $O_i = (S_i, L_i, X_i I(S_i \in S), M_i I(S_i \in S), Y_i I(S_i \in S))$. We assume that (O_1, \ldots, O_n) are independent and identically distributed (i.i.d.) realizations of $O = (S, L, XI(S \in S), MI(S \in S), YI(S \in S))$. The causal diagram describing the data generating mechanism is depicted in figure 1. This causal diagram also represents a nonparametric structural equation model (NPSEM) satisfying:

$$L = f_L(U_l, \epsilon_L); \quad S = f_S(L, \epsilon_S); \quad X = f_X(L, S, \epsilon_X); \quad M = f_M(X, L, S, \epsilon_M); \quad Y = f_Y(L, X, M, S, U_l, \epsilon_Y)$$

where U_l denotes unobserved common causes of L and Y and $(\epsilon_L, \epsilon_S, \epsilon_X, \epsilon_M, \epsilon_Y)$ denotes a vector of exogenous factors. Here, the functions f are assumed to be deterministic but unknown.




Figure 1: A causal diagram illustrating the setting of interest

Some remarks are noteworthy here. First, the populations $S \in \{j\} \cup S$ are subgroups of a super-population obtained by combining all hypothetical populations in the analysis. The chance of an individual being in each population depends on his/her baseline covariate profile, but such inclusion status does not change or affect patient characteristics. In figure 1, we thus let *L* cause *S* but not vice-versa, as in other related works. However, the proposal discussed below also works under an alternative causal diagram with an arrow $S \rightarrow L$ instead of $L \rightarrow S$.

Another remark is that we allow S to have a causal impact on X and on post-baseline variables such as M and Y. This will reflect the heterogeneity across datasets that can influence the outcome (see more discussion below). Finally, by considering all mediators M_1, \ldots, M_Z as a bloc, we avoid making assumptions on the causal structure among these mediators. This will simplify the proposal that we discuss below. In appendix A3, we will extend it to the setting in which the causal structure between mediators is known.

2.2 Addressing heterogeneity: lessons learned from the meta-analysis literature

On the log relative risk scale, the average exposure effect in the target population S = j can be expressed as:

$$\lambda(j) = \log \theta(x = 1, j) - \log \theta(x = 0, j)$$

where $\theta(x, j) = P\{Y(x) = 1 | S = j\}$, and Y(x) denotes the counterfactual outcome that could be observed in a patient if this patient were exposed to X = x. We focus on the log relative risk scale as it is frequently used to measure the treatment effect for a binary outcome. However, the discussion below can be generalized to other scales (such as risk difference or odds ratio), and other outcome types.

Assume consistency (i.e. Y(x) = Y among those with X = x), and ignorability of the exposure assignment in the target population j (i.e. $Y(x) \perp X | L, S = j$), one then has:

$$\theta(x,j) = \int P(Y=1|x,l,S=j) \cdot f_L(l|S=j)dl$$

where P(Y = 1 | x, l, S = j) is the short-hand notation for P(Y = 1 | X = x, L = l, S = j) and so forth. To borrow the information of the exposure and outcome from other populations in S, the following assumption is often made:

(A1) Common outcome distribution given L, i.e. $P(Y = 1|x, l, S = j) = P(Y = 1|x, l, S = j') \forall j' \in S$

for then $\theta(x, j)$ can be reexpressed as:

$$\theta(x,j) = \int P(Y=1|x,l,S\in\mathcal{S}) \cdot f_L(l|S=j)dl$$

Assumption (A1) requires that there is no arrow going from *S* to *M* and *Y* in the causal diagram. In that case, one possible approach to estimate $\theta(x, j)$ is to combine individual-level data from different sources in *S* to construct a conditional outcome model (given *X* and *L*), then using this model to predict the chance of experiencing the outcome under exposure level *x* of patients in population S = j, and averaging these predicted values.

Concerns arise when heterogeneity exists across data sources such that there are arrows going from S to M or Y, as (A1) may be violated in those situations. For example, consider our illustrative Moving to Opportunity (MTO) study, which aims to assess the effect of a housing voucher on psychiatric outcomes among children in various US cities. A mediator of interest in this case is the type of school (high-poverty vs not) that the child attends. If the city has a direct impact on the type of school, then assumption (A1) will be violated even when there is no arrow $S \rightarrow Y$, since:

$$P(Y = 1|x, l, S = j) = \sum_{m} P(Y = 1|x, l, m, S = j) \times P(m \mid x, l, S = j)$$

=
$$\sum_{m} P(Y = 1|x, l, m, S = j') \times P(m \mid x, l, S = j)$$

\neq
$$\sum_{m} P(Y = 1|x, l, m, S = j') \times P(m \mid x, l, S = j')$$

=
$$P(Y = 1|x, l, S = j')$$

where the mediator mechanism $P(m \mid x, l, S = j)$ varies across *j*.

To weaken assumption (A1), one can choose to borrow the information on the mediator and the outcome generating mechanism from separate sources. More precisely, if one believes that the mediator generating mechanism in the target population S = j is closer to the one observed in data sources $S \in S_2$, and the outcome generating mechanism is closer to the one observed in data sources $S \in S_3$, where $S_2, S_3 \subseteq S$, then the following (weaker) assumptions can be considered:

(A2) Common mediator distribution given L, i.e.:

$$f_M(m|x, l, S = j) = f_M(m|x, l, S = j_1) = \dots = f_M(m|x, l, S = j_{|S_2|}) \quad \forall j_1, \dots, j_{|S_2|} \in S_2$$

(A3) Common outcome distribution given L and M, i.e.:

$$P(Y = 1 | x, m, l, S = j) = P(Y = 1 | x, m, l, S = j_1) = \dots = P(Y = 1 | x, m, l, S = j_{|S_3|}) \quad \forall j_1, \dots, j_{|S_3|} \in S_3$$

Under these assumptions, $\theta(x, j)$ can be reexpressed as:

$$\theta(x,j) = \int P(Y=1|x,m,l,S\in\mathcal{S}_2) \cdot f_M(m|x,l,S\in\mathcal{S}_3) \cdot f_L(l|S=j)dl \tag{1}$$

Despite the weaker assumptions, combining data from different sources as in (1) may still suffer from residual bias when distributions within S_2 (or within S_3) are not entirely homogeneous. Note that in assumption (A2) and (A3), the first equality is untestable as we do not observe (X, M, Y) in the target population S = j. Nonetheless, other equalities in these assumptions (which assume data sources in S_2 (or S_3) are homogeneous in the mediator distribution (or the outcome distribution)) can be assessed by using statistical tests (Racine *and others*, 2006; Luedtke *and others*, 2019). The absence of heterogeneity across distributions in S_2 (or S_3) could also informally suggests whether the first equality in (A2) and (A3) is realistic. Assume that New York City is the target population and we borrow data from similar East Coast US cities such as Philadelphia, DC, Boston. If we see that the mediator (or outcome) distributions from these cities are homogeneous, it is reasonable to believe that the mediator (or outcome) distribution in New York City will be the same as in those cities, since they are all from the same region and have likely similar clinical standard. Statistical testing, however, are subject to the lack of power, let alone other risks such as model mispecification in some approaches. Besides, simply testing the equality of mediator (or outcome) distribution in S_2 (or S_3) do not inform us about the magnitude of heterogeneity across these data sources if it exists, or how we can adjust our analysis plan to account for it. The fact that statistical heterogeneity is inevitable (regardless of whether or not we happen to be able to detect it) also motivates the need for methods that can quantify heterogeneity across data sources, and shift focus from testing the presence of heterogeneity, to assessing the impact of it on the transported estimates (Higgins *and others*, 2003, 2019).

The challenge of heterogeneity also arises in other research areas such as evidence synthesis and meta-analysis. In this literature, the slightly different focus is on combining multiple exposure effect estimates $\hat{\lambda}(1), \ldots, \hat{\lambda}(K)$ that are obtained from different populations $1, \ldots, K$. The heterogeneity across these populations is taken into account in multiple ways. For instance, studies are only meta-analyzed when they are deemed as sufficiently similar. And even when this condition holds, a meta-analysis will not naively combine individual-level data from different studies (whenever such data are available). Instead, it assumes that $\lambda(1), \ldots, \lambda(K)$ are heterogeneous but all coming from an underlying distribution. The target of the analysis is to obtain the mean of such distribution as a summary, and the variance as a measurement of heterogeneity (Higgins *and others*, 2009).

The above principle of meta-analysis seems to be particularly relevant to a data fusion setting. According to this principle, data from different sources should be independently transported to the target population S = j, and the transported findings should be summarized (or meta-analyzed) in a way that takes into account the possible heterogeneity between different data sources. To formalize this, we denote:

$$\theta(x, j, k, p) = \int E(Y|L = l, M = m, X = x, S = k) \cdot f_M(m|X = x, L = l, S = p) \cdot f_L(l|S = j) \, dm dl$$
(2)

for $k \in S_2$ and $p \in S_3$. When data sources are homogeneous (hence the two assumptions (A2) and (A3) are satisfied), we have $\theta(x, j, k, p) = \theta(x, j, j, j) = \theta(x, j)$. However, such equality may not hold in general. Intuitively, $\theta(x, j, k, p)$ describes the potential result that can be observed in population S = j under exposure level x, if this population has the same mediator and outcome generating mechanism as in data sources p and k, respectively. In section 2.2 and 2.3, we moreover show that $\theta(x, j, k, p)$ can have a causal interpretation when certain assumptions on the data structure are satisfied. Based on $\theta(x, j, k, p)$, we can define:

$$\lambda(j,k,p) = \log \theta(x=1,j,k,p) - \log \theta(x=0,j,k,p)$$

Estimating and comparing different $\lambda(j, k, p)$ will tell us to what extent the transported effect estimates will be consistent when using data from different sources. This is extensively discussed in section 3, but we briefly provide here some intuitions. First, we summarize all $\lambda(j, k, p)$ by calculating the expectation $E\{\lambda(j, k, p)|j\}$. We then compute the corresponding variance, $V\{\lambda(j, k, p)|j\}$, to quantify the total heterogeneity in the (conditional) mediator and outcome distribution between data sources. These expectations and variances are computed across a uniform distribution on (k, p) and should not be confused with expectations and variances computed with respect to the distribution of the observed data. When $V\{\lambda(j, k, p)|j\} = 0$, data sources in S_2 are homogeneous in the mediator distribution, and data sources in S_3 are homogeneous in the outcome distribution. Under the assumption that these (homogeneous) data sources are similar to the target population j in the mediator distribution (for S_2) and the outcome distribution (for S_3), averaging $\hat{\lambda}(j, k, p)$ will provide a consistent estimate for $\lambda(j)$.

When $V\{\lambda(j, k, p)|j\} \neq 0$, either data sources in S_2 are heterogeneous in the mediator distribution, or data sources in S_3 are heterogeneous in the outcome distribution, or both. We can investigate the exact source of heterogeneity by estimating $V\{\lambda(j, k, p)|j, p\}$ for some $p \in S_2$ fixed, and $V\{\lambda(j, k, p)|j, k\}$ for some $k \in S_3$ fixed. By construction, $V\{\lambda(j, k, p)|j, p\}$ describes the impact of the heterogeneity in the outcome distribution across data sources in S_3 on the variability of $\lambda(j, k, p)$. Similarly, $V\{\lambda(j, k, p)|j, k\}$ describes the impact of the heterogeneity in the mediator distribution across data sources in S_2 on the variability of $\lambda(j, k, p)$. The magnitude of these variances might also shed insights on the magnitude of bias due to heterogeneity. Intuitively, if the amount of heterogeneity (measured by the above variance components) is smaller than the sampling error (measured by the variance of $\hat{\lambda}(j, k, p)$), the bias due to heterogeneity might probably be small and negligible. This evaluation of magnitude of bias is not possible if one only uses statistical tests to assess the presence of heterogeneity. In the next section we discuss optimal and robust estimation of $\lambda(j, k, p)$. These estimators will then be used in section 3 to study heterogeneity with the tools discussed in this paragraph.

2.3 Estimand and identification

Before discussing the estimation strategies for $\lambda(j, k, p)$, we will show that $\lambda(j, k, p)$ can also have a causal interpretation under certain assumptions on the data structure. Note that regardless of whether these causal assumptions are satisfied, $\lambda(j, k, p)$ is still well-defined as a population parameter by formula (2).

We denote (i) M(x, k) the potential value of M if a patient were in population k and exposed to x; (ii) Y(x, m, k) the potential value of Y if a patient were in population k, exposed to x and having the mediator level fixed at m; (iii) Y(x, k, M(x, p)) the potential value of Y that could be observed if an individual were in population k, exposed to x and having his/her mediator level fixed at the value potentially observed if (s)he were instead in population p and exposed to x. Under the causal assumptions that we provide below, one then has:

$$\theta(x, j, k, p) := P\{Y(x, k, M(x, p)) = 1 | S = j\}$$

Likewise,

$$\lambda(j,k,p) := \log \frac{\theta(x=1,j,k,p)}{\theta(x=0,j,k,p)} = \log \frac{P\{Y(1,k,M(1,p)) = 1 | S = j\}}{P\{Y(0,k,M(0,p)) = 1 | S = j\}}$$

where $x = 0, 1, k \in S_3$ and $p \in S_2$. Hence, $\lambda(j, k, p)$ describes the average exposure effect that could have been observed in population S = j, had individuals in this population been treated (versus not treated) as in population S = k, and the mediator level of each subject been fixed at the value potentially observed if being treated (versus not treated) as in population S = p. The intuition behind fixing M and Y at the values counterfactually observed in a different population, is to isolate the heterogeneity due to the mediator distribution from the heterogeneity due to the outcome distribution, thereby making the impact of each source of heterogeneity on the transported effect estimates quantifiable (see section 3).

For $\theta(x, j, k, p)$ to have the above causal interpretation, the following assumptions are needed (Appendix A1):

- (B1) **Consistency**, i.e. (X, S) = (x, k) implies M(x, k) = M and (X, M, S) = (x, m, k) implies Y(x, k, m) = YImplicit in this assumption is that an individual's observed mediator and outcome is not affected by others' exposure level or study membership; which is also known as the Stable Unit Treatment Value Assumption (Rubin, 1978, 1990).
- (B2) **Positivity**, i.e. $P(0 < P(S = p|L) < 1) = P(0 < P(S = p|M, L) < 1) = P(0 < P(X = 1|M, L, S) < 1) = P(0 < P(X = 1|L, S) < 1) = 1 \forall p$ This postulates that the support of *L* and of (*L*, *M*) is the same for each level of *S*, and the support of (*L*, *M*, *S*) and of (*L*, *S*) is the same for each level of *X*, to guarantee an adequate overlap between different population in terms of case-mix and of the mediator distribution. Such an assumption is important to learn about the treatment effect in one population by using external data.
- (B3) Within-trial ignorability, i.e. $M(x, k) \perp X | L, S = k$ and $Y(x, k, m) \perp X | L, S = k \forall x, k, m$ This states that in each population, the treatment is independent of all counterfactual mediators and outcomes given L, which is often guaranteed to hold in a trial setting because of treatment randomization.
- (B4) **Between-trial ignorability**, i.e. $Y(x, k, m) \perp S|L$ and $M(x, k) \perp S|L \forall x, k, m$. This states that the population indicator S is independent of all counterfactual mediators outcomes, conditioning on L. Thus, the set L must contain all prognostic factors of the mediator and outcome that are differentially distributed across population.
- (B5) No unmeasured mediator-outcome confounders, i.e. $Y(x, k, m) \perp M | X = x, L, S = k$ As we are interested in the effects of the mediator on the outcome, mediator-outcome confounders must be properly taken into account.

(B6) **Cross-world assumption**, i.e. $Y(x, k, m) \perp M(x, p)|L, S = j$

This assumption requires conditional independence between Y and M when the individual is recruited in two different studies. Such an assumption is satisfied under the non-parametric structural equation model associated with the causal diagram depicted in figure 1.

2.4 Estimation and inference

In the discussion below, we will introduce the vector of nuisance parameters $\eta = (a, b, g, e, r, h)$, where a = a(x, m, l, k) denotes E(Y|x, m, l, S = k), b = b(x, l, p) denotes E(a(x, M, L, k)|x, l, S = p), g = g(x, m, l, p) denotes P(X = x|m, l, S = p), e = e(p, m, l) denotes P(S = p|m, l), r = r(p, l) denotes P(S = p|l), t(x, l, p) denotes P(X = x|l, S = p) and h = h(p) denotes P(S = p).

2.4.1 G-computation

By construction, a simple g-computation approach could be used to estimate $\theta(x, j, k, p)$, where $x = 0, 1, k \in S_3$ and $p \in S_2$. In step 1, a regression model for a(X, M, L, S) is postulated and fitted by using the data on Y, X, M, L and S of subjects in S_3 . This model is then used to compute the predicted values $\hat{a}(x, M_i, L_i, k)$ for each subject i in S_2 . In step 2, a model for b(X, L, S) is postulated and fitted by using the estimates $\hat{a}(x, M, L, k)$ obtained from step 1 and the data on X, L, S of subjects in S_2 . The predicted values $\hat{b}(x, L_{i'}, p)$ are computed for each individual i' with $S_{i'} = j$. These predicted values are then averaged to obtain an estimator $\hat{\theta}_0(x, j, k, p)$ for $\theta(x, j, k, p)$.

Note that $\hat{\theta}_0(x, j, k, p) \xrightarrow{P} \theta(x, j, k, p)$ (where \xrightarrow{P} denotes convergence in probability) if one can correctly specify a parametric model for a(X, M, L, S) and b(X, L, S), in the sense that $a = a(X, M, L, S, \psi_a)$ and $b = b(X, L, S, \psi_b)$ for some finite-dimensional parameter ψ_0 and ψ_1 , respectively. In that case, an estimator $\hat{\lambda}_0(j, k, p)$ for $\lambda(j, k, p)$ can be obtained by:

$$\hat{\lambda}(j,k,p) := \log \hat{\theta}_0(x=1,j,k,p) - \log \hat{\theta}_0(x=0,j,k,p)$$

The Delta method or parametric bootstrap sampling then allows one to derive the asymptotic variance of $\hat{\lambda}_0(j, k, p)$.

The consistency of $\hat{\theta}_0(x, j, k, p)$ with respect to $\theta(x, j, k, p)$, however, is not ensured if the nuisance parameters a(X, M, L, S) and b(X, L, S) cannot be described by parametric models, or if at least one of these models is incorrectly specified. In the next section, we will construct alternative approaches to estimate $\theta(x, j, k, p)$, which will rely on models of different aspects of the data generating mechanism other than the outcome, and will allow for the use of flexible, data-adaptive regression methods to prevent bias due to (nuisance) model mispecifications.

2.4.2 More robust estimators

In this section, we consider using semi-parametric theory to construct an efficient, non-parametric estimation approach for $\theta(x, j, k, p)$. This approach can achieve root *n* rate of convergence to the parameter of interest even when the nuisance functions in η are estimated at slower rates. To achieve this, one first need to characterize the efficient influence function (EIF) $IF(O, \eta)$ of $\theta(x, j, k, p)$. For a fixed value η_1 of the nuisance parameter (corresponding, e.g., to a preliminary estimator or its probability limit), the EIF generally allows us to write an expansion

$$\theta(x, j, k, p, \boldsymbol{\eta}_1) - \theta(x, j, k, p, \boldsymbol{\eta}) \coloneqq -E\{IF(O, \boldsymbol{\eta}_1)\} + R(\boldsymbol{\eta}, \boldsymbol{\eta}_1)$$

where $R(\eta, \eta_1)$ is a second-order term that can be written as sums of terms of the type $E[c(\eta, \eta_1) \{ f(\eta_1) - f(\eta) \} \{ g(\eta_1) - g(\eta) \} \}$. The above expression then means that $E\{IF(O, \eta_1)\}$ represents the first-order bias of a plug in estimator given by $\theta(x, j, k, p, \eta_1)$. This motivates the construction of de-biased estimators by first computing the plug-in estimator, then subtracting the estimate of its first order bias. If $R(\eta, \eta_1)$ can be assumed to be small (see below), de-biased estimators can be proven to have optimal properties. We start our proposal by giving the efficient influence function for $\theta(x, j, k, p)$ in the non-parametric model (Appendix A2).

Proposition 2.1 By using the Delta method for the functionals, one can derive the following efficient influence function for $\theta(x, j, k, p)$:

$$\begin{split} IF(O, \eta) &= \frac{g(x, M, L, p)}{g(x, M, L, k)} \cdot \frac{e(p, M, L)}{e(k, M, L)} \cdot \frac{r(j, L)}{r(p, L)} \cdot \frac{I(x, k)}{t(x, L, p) \cdot h(j)} \{Y - a(M, x, L, k)\} \\ &+ \frac{r(j, L)}{r(p, L)} \cdot \frac{I(x, p)}{t(x, L, p) \cdot h(j)} \{a(M, x, L, k) - b(x, L, p)\} + \frac{I(j)}{h(j)} \{b(x, L, p) - \theta(x, j, k, p, \eta)\} \end{split}$$

where I(x, p) = I(X = x, S = p), I(j) = I(S = j) are indicator functions.

Some remarks are noteworthy here. The above EIF for $\theta(x, j, k, p)$ is closely related to the EIF proposed recently by Rudolph and van der Laan (2017), or by Li and Luedtke (2021) for the intention-to-treat treatment effect under non-adherence. The adherence status in such a setting can be viewed as a single binary mediator which entirely mediates the causal effect of X on Y. Our setting here is more general. We allow for M to be a vector of multiple mediators of any nature, and these meditors might only partially mediates the causal effect of X on Y. The above EIF can thus be seen as an extension, or a reparameterization of the EIF in previous works to enable the construction of efficient estimators in more general settings.

Let $\hat{\eta} = (\hat{a}, \hat{b}, \hat{g}, \hat{e}, \hat{r}, \hat{h}, \hat{t})$ denote an estimate for η . Assume that $\hat{\eta}$ converges in probability to some η_1 that might be potentially different from the true value of the nuisance parameter η . In appendix 1, we prove that the remainder term:

$$R(\boldsymbol{\eta}, \boldsymbol{\eta}_1) := \theta(x, j, k, p, \boldsymbol{\eta}_1) - \theta(x, j, k, p, \boldsymbol{\eta}) + E\{IF(O, \boldsymbol{\eta}_1)\}$$

is indeed a second order term. This then leads to the following lemma (Appendix A2):

Lemma 2.2 Let $\eta_1 = (a_1, b_1, g_1, e_1, r_1, h_1, t_1)$ be such that one of the following conditions hold:

- (*i*) $a = a_1$ and $b = b_1$
- (*ii*) $r = r_1$ and $t = t_1$ and either $(g, e) = (g_1, e_1)$ or $a = a_1$

then $E\{IF(O, \eta_1)\} = R(\eta, \eta_1) = 0.$

Lemma 2.2 implies that it is possible to construct consistent estimators for $\theta(x, j, k, p)$ under consistent estimation of a subset of the nuisance parameters in η as specified above. Note that condition (i) in lemma 2.2 corresponds to the requirement under which the previously discussed g-computation estimator is consistent. To construct alternative estimators for $\theta(x, j, k, p)$ that are consistent under condition (ii), $\theta(x, j, k, p)$ is reexpressed as:

$$\begin{aligned} \theta(x,j,k,p) &= E\left\{Y \cdot \frac{t(x,k)}{t(x,p)} \cdot \frac{g(x,M,L,p)}{g(x,M,L,k)} \cdot \frac{e(p,M,L)}{e(k,M,L)} \cdot \frac{r(j,L)}{r(p,L)} \cdot \frac{h(k)}{h(j)}\right\} \\ &= E\left\{a(x,M,L,k) \cdot \frac{I(x,p)}{t(x,p)} \cdot \frac{r(j,L)}{r(p,L)} \cdot \frac{1}{h(j)}\right\}\end{aligned}$$

The first equality hence suggests a weighting-based approach that consistently estimates $\theta(x, j, k, p)$ if g, e and r can be correctly specified as $g = g(x, M, L, p, \psi_g)$, $r = r(j, L, \psi_r)$ and $e = e(p, M, L, \psi_e)$ for some finite-dimensional parameter ψ_g, ψ_r and ψ_e , respectively. In contrast, the second equality suggests a weighting-regression approach that consistently estimate $\theta(x, j, k, p)$ if the nuisance parameters a and r can be correctly specified as $a(x, M, L, k, \psi_a)$ and $r(j, L, \psi_r)$ for some finite-dimensional parameter ψ_a and ψ_r , respectively. Depending on the specific applications, one approach may be more preferable than the other two when knowledge about certain nuisance parameters is available. However, similar to g-computation, both approaches are asymptotically biased if the corresponding nuisance models are incorrectly specified or if they cannot be described by a finite-dimension vector of parameters. Furthermore, if model selection or regularization techniques must be employed to estimate the nuisance parameters, then the estimator distribution is generally unknown, making it very difficult to construct valid methods for confidence intervals and hypothesis tests. To overcome the limitations of the aforementioned approaches, we propose below an alternative estimator that does not require parametric assumptions. This, often referred to as the *one-step* estimator, can be expressed as:

$$\hat{\theta}_1(x, j, k, p) = \hat{\theta}_0(x, j, k, p) + \frac{1}{N} \sum_{i=1}^N IF(O_i, \hat{\eta}_1)$$
(3)

where $\hat{\theta}_0(x, j, k, p)$ is computed as in section 3.1, but with the two nuisance parameters a(x, M, L, k)and b(x, L, p) possibly estimated by a flexible data-adaptive algorithm. Likewise, $IF(O_i, \hat{\eta}_1)$ is computed by plugging into $IF(O_i, \eta_1)$ an estimate $\hat{\eta}_1$ of η_1 that is possibly obtained by using a flexible data-adaptive algorithm. In the theorem below, we clarify the conditions on $\hat{\eta}_1$ for $\hat{\theta}_1(x, j, k, p)$ to be asymptotically normal and efficient.

Theorem 2.3 (Asymptotic normality and efficiency) Assume

- (i) Positivity, described as identification assumption (A2) in Section 3.2, and
- (ii) The second-order term $R(\hat{\eta}, \eta)$ is $o_P(n^{-1/2})$ and
- (iii) The class of functions $\{IF(\eta, \theta') : |\theta' \theta| < \delta, ||\eta \eta_1|| < \delta\}$ is Donsker for some $\delta > 0$ and such that $P\{IF(\eta, \theta') - IF(\eta_1, \theta)\}^2 \rightarrow 0$ as $(\eta, \theta') \rightarrow (\eta_1, \theta)$

then:

$$\hat{\theta}_1(x, j, k, p) = \theta_1(x, j, k, p) + \frac{1}{N} \sum_{i=1}^N IF(O_i, \eta) + o_P(1),$$

due to which $\sqrt{N}(\hat{\theta}_1 - \theta) \xrightarrow{D} N(0, \zeta^2)$, where $\zeta^2 = V\{IF(O, \eta)\}$ is the non parametric efficiency bound.

Note that condition (ii) for asymptotic normality in theorem 2.3 is satisfied if all components of $\hat{\eta}$ converges in $L_2(P)$ norm to their true counterparts in η at $n^{-1/4}$ -rate or faster. This is the case for many data-adaptive algorithms such as LASSO or highly adaptive LASSO, under certain conditions (Belloni *and others*, 2015; Benkeser and Van Der Laan, 2016). In contrast, condition (iii) (i.e. Donsker condition) may be avoided by using cross-fitting in the estimation procedure. To achieve this, the dataset is randomly partitioned into Q sets of approximately equal size, namely V_1, \ldots, V_Q . On each sample $T_q = \{1, \ldots, N\} \setminus V_q$, the data-adaptive algorithm will be trained and then used to produce a prediction $\hat{\eta}^*$ of η for each patient in the validation set V_q . The one-step estimator is finally adapted to cross-fitting by substituting all occurrences of $\hat{\eta}(O_i)$ by $\hat{\eta}^*(O_i)$ in the estimation procedure.

As a direct consequence of theorem 2.3, the variance of the plug-in estimator $\hat{\theta}_1(x, j, k, p)$ can be estimated by the sample variance of the efficient influence function, i.e. $\hat{\zeta}^2 = \hat{V}(IF(O, \hat{\eta}))$, with $\hat{\theta}_1(x, j, k, p)$ and the nuisance parameter vector $\boldsymbol{\eta}$ estimated as described above. In what follows, a plug-in estimator for $\lambda(j, k, p)$ and its variance can be constructed from $\hat{\theta}_1(x, j, k, p)$ and $\hat{\zeta}^2$ by applying the Delta method, as is done for the simple g-computation approach.

3 Heterogeneity assessment among transported estimates

In this section, we propose simple methods to summarize the estimates $\hat{\lambda}(j, k, p)$ ($p \in S_2, k \in S_3$) obtained from the previously suggested procedures, with aim to investigate the impact of heterogeneity across data sources in S_2 and S_3 on the variability observed among $\lambda(j, k, p)$. Note that by construction, the difference between various $\lambda(j, k, p)$ can be explained by two possible reasons, which are:

- (i) *M*-related heterogeneity, i.e. the difference across studies in S_2 in the mediator distribution P(M|X = x, L, S = p)
- (ii) *Y*-related heterogeneity, i.e. the difference across studies in S_3 in the outcome distribution E(Y|L, M, X = x, S = k).

One can however quantify the impact of one heterogeneity source on the estimands by keeping the other source fixed. For instance, different $\lambda(j, k_0, p)$ for some k_0 fixed and p varying are different due to the differential distribution of M across studies, since P(L|j) and $E(Y|L, M, X = x, S = k_0)$ are unchanged among these estimands. Likewise, $\lambda(j, k, p_0)$ for some p_0 fixed and k varying are heterogeneous due to the differential outcome distribution across studies, since $P(M|X = x, L, S = p_0)$ and P(L|S = j) are unchanged among these estimands.

3.1 Parametric random-effect models

We start the discussion with a common parametric model to summarize $\hat{\lambda}(j, k, p)$, and then introduce our proposal based on a non-parametric generalization of the parametric random-effect model. This model involves two different random effects to capture the two aforementioned sources of heterogeneity, that is:

$$\hat{\lambda}(j,k,p) = \Lambda + \gamma_k + \delta_p + \epsilon_{jkp} \tag{4}$$

where Λ denotes the summary effect, $\boldsymbol{\gamma} = (\gamma_k \ \delta_p)^T \sim N[\mathbf{0}, \operatorname{diag}(\omega^2, \zeta^2)]$ denotes the randomeffect vector and $\epsilon_{jkp} \sim N(0, \sigma_{jkp}^2)$ denotes the residual random error. In this model, we assume that the random effects and the residual random error are pairwise independent, for the sake of simplicity. The covariance matrix of the random errors ϵ_{jkp} , however, is left unstructured, i.e. $\operatorname{cov}(\epsilon_{jkp}, \epsilon_{jk'p'}) \neq 0$ when (k - k')(p - p') = 0. A direct consequence of model (4) is that:

$$V\{\lambda(j,k,p)|j,p\} = V(\gamma_k) = \omega^2 \quad \forall p \in S_2$$
$$V\{\lambda(j,k,p)|j,k\} = V(\delta_p) = \zeta^2 \quad \forall k \in S_3$$

 ω^2 thus captures the so-called *Y*-related heterogeneity, while ζ^2 captures the *M*-related heterogeneity. As $V(\lambda(j, k, p)|j) = \omega^2 + \zeta^2$, the contribution of each source of heterogeneity to the total variability observed among all $\lambda(j, k, p)$ can then be expressed on the relative scale as:

$$I_Y^2 = \frac{\hat{\omega}^2}{\hat{\omega}^2 + \hat{\zeta}^2} \qquad \qquad I_M^2 = \frac{\hat{\zeta}^2}{\hat{\omega}^2 + \hat{\zeta}^2}$$

where $\hat{\omega}^2$ and $\hat{\zeta}^2$ are consistent estimates of the corresponding variance components.

Model (4) can be fitted by using restricted maximum likelihood. Alternatively, a Bayesian approach such as Markov chain Monte Carlo can be used to avoid convergence issues. From this, we obtain $\hat{\Lambda}$ which is a weighted average across all $\hat{\lambda}(j, k, p)$. When $\omega^2 = \zeta^2 = 0$, $\hat{\Lambda}$ will be a valid estimate for the average exposure effect in population *j*, given that the (homogeneous) mediator and outcome distributions across all data sources in S_2 and S_3 are similar to their analogues in population *j*.

Meanwhile, if $\zeta^2 \neq 0$, $\omega^2 = 0$, and we believe that the mediator distribution in the target population *j* is closest to its analogue in data source $p_0 \in S_2$, while the (homogeneous) outcome distribution across all data sources in S_3 are similar to their analogue in population *j*, then $\hat{\Lambda} + \hat{\delta}_{p_0}$ is a valid estimate for the average exposure effect in population *j*. Here, $\hat{\delta}_{p_0}$ denotes a valid estimate (e.g. the Best Linear Unbiased Prediction) for δ_{p_0} . When $\zeta = 0$, $\omega^2 \neq 0$ and we believe that the outcome distribution in the target population *j* is closest to its analogue in data source $k_0 \in S_3$, while the (homogeneous) mediator distribution across all data sources in S_2 are similar to their analogue in population *j*, then $\hat{\Lambda} + \hat{\gamma}_{k_0}$ is a valid estimate for the average exposure effect $\lambda(j)$ in population *j*. Here, $\hat{\gamma}_{k_0}$ denotes a valid estimate (e.g. the Best Linear Unbiased Prediction) for γ_{k_0} . Depending on the context, users might report the most appropriate summary estimates.

The situation, however, is more complicated when $\omega^2 \neq 0$ and $\zeta^2 \neq 0$. In that case, it is likely that combining different data sources to estimate $\lambda(j)$ will imply bias. However, the magnitude of ω^2 and ζ^2 might not be larger than the sampling variance of the estimators. If so, the importance of inconsistency across data sources might arguably be low, and the residual bias due to heterogeneity when averaging different $\lambda(j, k, p)$ might still be acceptable.

Model (4), however, can provide invalid findings when the normality and independence assumptions made on the random effects are violated. Indeed, these assumptions are purely made for statistical convenience and cannot be verified by empirical evidence. In the next section, we will thus provide an alternative approach to summarize $\hat{\lambda}(j, k, p)$ that does not rely on parametric assumptions. This approach will also allow one to decompose the heterogeneity across data sources into different components as described above, but in a non-parametric manner.

3.2 Non-parametric heterogeneity assessment

As before, one can obtain different summary effects by averaging different sets of $\lambda(j, k, p)$. For instance, the mean of $\lambda(j, k, p)$ that share the same outcome distribution E(Y|X, L, M, S = k), where $k \in S_3$, can be expressed as:

$$\Lambda(j,k,\bullet) = \frac{1}{|\mathcal{S}_2|} \sum_{p \in \mathcal{S}_2} \lambda(j,k,p)$$

 $\Lambda(j, \bullet, p)$ can also be defined in a similar way. Marginalizing $\Lambda(j, k, \bullet)$. over k (or $\Lambda(j, \bullet, p)$ over p), we then obtain the mean of all $\Lambda(j, k, p)$, i.e.:

$$\Lambda(j,\bullet,\bullet) = \frac{1}{|\mathcal{S}_2| \cdot |\mathcal{S}_3|} \sum_{k \in \mathcal{S}_3} \sum_{p \in \mathcal{S}_2} \lambda(j,k,p)$$

The total variance across studies can then be defined as:

$$\tau^{2'} := V\{\lambda(j,k,p)|j\} = \frac{1}{|\mathcal{S}_2| \cdot |\mathcal{S}_3|} \sum_{\substack{p \in \mathcal{S}_2; \\ k \in \mathcal{S}_3}} \left\{\lambda(j,k,p) - \Lambda(j,\bullet,\bullet)\right\}^2$$

By considering k, p as two independent variables that are discrete and uniformly distributed, one can apply the law of total variance to decompose $\tau^{2'}$ into $\tau^{2'} = \omega^{2'} + \zeta^{2'}$, where:

$$\begin{split} \zeta^{2'} &:= E\left\{V\big(\lambda(j,k,p)|j,k\big)|j\right\} = \frac{1}{|\mathcal{S}_2| \cdot |\mathcal{S}_3|} \sum_{\substack{p \in \mathcal{S}_2;\\k \in \mathcal{S}_3}} \left\{\lambda(j,k,p) - \Lambda(j,k,\bullet)\right\}^2 \\ \omega^{2'} &:= V\left\{E(\lambda(j,k,p)|j,k)|j\right\} = \frac{1}{|\mathcal{S}_3|} \sum_{k \in \mathcal{S}_3} \left\{\Lambda(j,k,\bullet) - \Lambda(j,\bullet,\bullet)\right\}^2 \end{split}$$

The first component $\zeta^{2'}$ describes the (averaged) variability of $\Lambda(j, k, p)$ when p varies and (j, k) are fixed. It hence reflects the impact of M-related heterogeneity across data sources in S_2 on the inconsistency observed among $\lambda(j, k, p)$. The second component $\omega_{2'}$ describes the variability of $\Lambda(j, k, \bullet) = E(\lambda(j, k, p)|j, k)$ when j is fixed and k varies. It hence reflects the impact of Y-related heterogeneity across data sources in S_3 on the inconsistency observed among $\lambda(j, k, p)$. In practice, each component can be estimated by plugging in the estimates $\hat{\lambda}(j, k, p)$.

Note that when the random-effect model (4) holds, the different variance components discussed in this section will converge (almost surely) to the corresponding analogues in model (4) when the number of studies $K \to \infty$. The advantage, however, is that $\tau^{2'}, \omega^{2'}, \zeta^{2'}$ and $\xi^{2'}$ still provide valid measurements of the impact of different sources of heterogeneity on $\lambda(j, k, p)$ when model (4) is incorrectly specified.

Finally, weights can be used when calculating the above means and variances to make the contribution of different data sources proportional to their sizes. For instance, one can consider the indexes k, p as independent random variables following the distributions:

$$P(k = p_0) = \frac{\sum_i I(S_i = p_0)}{\sum_i I(S_i \in S_2)}; \qquad P(k = k_0) = \frac{\sum_i I(S_i = k_0)}{\sum_i I(S_i \in S_2)}$$

for all $p_0 \in S_2$ and $k_0 \in S_3$. Using deterministic (a priori known) weights as these can prevent complications due to having to estimate the unknown weights, such as the inverse of the variance as in standard meta-analysis. In many recent works, complications due to weight estimation have been shown to be important when the studies is of small sample size (Bakbergenuly *and others*, 2020; Walter and Balakrishnan, 2022).

4 A simulation study

In this section, we assess the finite-sample performance of the proposed methods when being used to estimate the average treatment effect in a target population S = 1, by using the treatment, mediator and outcome data from 4 simulated randomized controlled trials conducted in 4 other populations, i.e. S = 2, ..., 5. We thus have $S_2 = S_3 = \{2, ..., 5\}$. The data-generating mechanism is as follow, with all variables (except for S) being Bernouilli distributed:

$$\begin{split} P(L_1 = 1) &= P(L_2 = 1) = 0.5, \quad L_1 \perp L_2 \\ P(S = 2|L) &= P(S = 1|L) \cdot \exp(0.01 + 0.45L_1 + 0.3L_2 + 0.5L_1L_2) \\ P(S = 3|L) &= P(S = 1|L) \cdot \exp(0.01 - 0.2L_1 + 0.2L_2 - L_1L_2) \\ P(S = 4|L) &= P(S = 1|L) \cdot \exp(0.01 + 0.45L_1 + 0.1L_2) \\ P(S = 5|L) &= P(S = 1|L) \cdot \exp(-0.25 + 0.55L_1 - 0.25L_2 - L_1L_2) \\ P(X = 1|L, S) &= 0.5 \\ P(M = 1|X, L, S) &= \exp((1.37 - 0.5X - 0.5L_1 + XL_1 - 0.5L_2) \\ P(Y = 1|X, L, S) &= \exp((-1 - X + 0.5 \cdot I(S \in \{2, 4\}) + 0.75M + 0.65XM \cdot I(S \in \{1, 3\}) \\ &+ 0.5L_1 + 0.5ML_1 - 0.5L_2 + XL_1) \end{split}$$

We consider three sample size, namely 1000, 5000 and 10000. For each sample size, we first estimate $\lambda(j, k, p)$ by the (non-parametric) plug-in approach proposed in the previous section. The nuisance parameters in η are estimated by super learner (SL) algorithms, whose library includes the sample mean, the generalized linear model with a logistic link function and the highly adaptive LASSO. Although cross fitting helps ensure asymptotic normality of the proposed estimator without relying on the Donsker condition, we will not consider it here due to the prohibitive computational times of the resulting simulation study. For each $\lambda(j, k, p)$, we calculate (i) the root-*N* bias, (ii) the mean squared error (scaled by the sample size) and (iii) the coverage of the 95% confidence interval (CI). In every simulation, we summarize each quantity (i–iii) by computing the mean across 25 parameters $\lambda(j, k, p)$.

In what follows, the different estimates $\hat{\lambda}(j, k, p)$ are summarized by using the random-effect model (4) and the non-parametric approach proposed in section 4.2. Note that the mediator distribution P(M = 1|X, L, S) is the same across data sources, hence there is no *M*-related heterogeneity. In contrast, there is heterogeneity in the outcome distribution across data sources, and only one of these (S = 3) has the outcome distribution similar to that in the target population S = 1. The magnitude of *Y*-related heterogeneity variance is 0.013, which is calculated by applying the non-parametric approach proposed in section 4.2 on the true values of $\lambda(j, k, p)$.

Model (4) with two random effects γ_k and δ_p is fitted by using a fully Bayesian approach based on Markov Chain Monte Carlo (MCMC) simulation. We consider weakly informative, independent prior distributions for the unknown parameters Λ , ω^2 and ζ^2 . The prior distribution for Λ is N(0, 1000), and for ω^2 and ζ^2 it is U(0, 100). For the sake of simplicity, we assume that the residual covariance matrix is known a priori. This assumption is also commonly adopted in the random-effect meta-analysis literature. The empirical (posterior) distributions of ω^2 and ζ^2 are summarized by the posterior median.

When $\hat{\lambda}(j, k, p)$ are summarized non-parametrically, we consider the weight of 1 for each data source for the sake of simplicity.

Table 1: Heterogeneity assessment by the random-effect model and the non-parametric approach

Heterogeneity	True value	Random-effect model			Non-parametric approach		
		1000	5000	10000	1000	5000	10000
<i>M</i> -related ($\times 10^{-3}$)	0.00	3.29	0.47	0.12	1.53	0.26	0.13
		(1.14, 9.41)	(0.19, 1.02)	(0.05, 0.29)	(1.03, 2.26)	(0.15, 0.41)	(0.09, 0.21)
<i>Y</i> -related ($\times 10^{-3}$)	12.5	30.9	21.5	26.7	30.2	16.2	14.5
		(1.22, 112)	(9.51, 42.0)	(16.6, 45.5)	(17.5, 48.0)	(11.4, 22.3)	(10.6, 18.7)

The root-N bias, the scaled MSE and the 95%CI coverage of the plug-in approach when estimating 25 parameters $\lambda(j, k, p)$ are summarized in figure 2. As the sample size increases,




Figure 2: Root-N bias, Scaled MSE and the coverage of the confidence interval across simulations for all $\lambda(j, k, p)$

the root-*N* bias distribution tends to approach the expected value of zero, while the scale MSE distribution remains stable. The coverage of the 95%CI also gets better when then sample size is larger. These findings indicate that the proposed plug-in approach performs sufficiently well in estimating $\lambda(j, k, p)$.

Results of the two heterogeneity assessment approaches are provided in table 1. As can be seen from this table, both approaches correctly identified the absence of *M*-related heterogeneity across data sources when the sample size is sufficiently large. However, the random-effect model (4) tends to overestimate the impact of *Y*-related heterogeneity on the variability of $\lambda(j, k, p)$, due to the violation of the parametric assumptions in model (4). In contrast, the non-parametric approach correctly estimates the amount of *Y*-related heterogeneity across data sources when the sample size is sufficiently large.

5 Illustrative data analysis

We now apply our proposed methods to data from the Moving to Opportunity study (MTO). MTO was a large-scale housing voucher experiment conducted by the US Department of Housing and Urban Development in the 1990s-2000s (Sanbonmatsu *and others*, 2011; Kling *and others*, 2007). In this experiment, families living in high-rise public housing developments in five US cities could sign up to be randomized to receive a Section 8 housing voucher. These vouchers, which continue to be the primary form of federal housing assistance today (now called Housing Choice vouchers), can be used by families to move out of public housing and into a rental on the private market by subsidizing their rent to be approximately one-third of their income (Development, 2022). The participating MTO families (adults and their children) were followed up for 10-15 years after randomization and economic, educational, and health-related outcomes were assessed (Sanbonmatsu *and others*, 2011).

As stated above, there were five MTO cities/sites, but we exclude one of those, Baltimore, because of evidence that the housing voucher intervention did not work the same there due to a concurrent housing voucher program (Rudolph *and others*, 2018). There is ample evidence of effect heterogeneity across MTO sites, even qualitative effect heterogeneity in which some sites exhibited negative effect estimates but others exhibited positive effect estimates (Rudolph and others, 2018). This evidence of effect heterogeneity is seen not only for total effects, but also for indirect effects (Rudolph *and others*, 2020, 2021; Rudolph and Díaz, 2022).

When such effect heterogeneity exists, it is natural to want to understand why: what components of the causal process are contributing to such heterogeneity? We address such a question here. Specifically, in this analysis, we are interested in understanding contributions to site-level heterogeneity in the effect of moving with the voucher on risk of developing a psychiatric disorder in adolescence (10-15 years after randomization), using attendance of a lower-poverty school during follow-up as an intermediate variable/ mediator, among boys who were younger than 5 years old when their family enrolled in MTO. Using the proposed methods, we decompose the site-level heterogeneity into heterogeneity in the mediation model or heterogeneity in the outcome model.

Baseline covariates, L, included family-level, child-level, and neighborhood characteristics at time of randomization. Child variables included: race/ethnicity, age, history of behavioral problems, and whether the child was ever enrolled in special class for gifted and talented students. Adult variables included: whether the adult: was a high school graduate or had their GED, never married, was under 18 when child was born, was currently working, was currently receiving welfare, perceived the baseline neighborhood as being unsafe at night, was very dissatisfied with neighborhood, had moved 3 times or more, signed up to participate in MTO so that their child(ren) could attend better quality schools, had a Section 8 voucher previously; whether a household member had a disability, size of household (2, 3, or \geq 4), and the poverty rate of the baseline neighborhood. Site, S, included Boston, Chicago, Los Angeles (LA), and New York City (NYC). Exposure, X, was moving with the voucher out of public housing, and was a binary variable. The mediator, M, was a binary variable indicating whether the child had ever attended a school that was not high-poverty (a non-Title I school). The outcome, Y, was a binary variable indicating whether the child had any psychiatric disorder in the past year, as defined by the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV).

Our aim here is to estimate the average treatment effect in the target MTO population of LA, using mediator and outcome data from other sites, i.e. Boston, Chicago and NYC. We first transport the data from these sites to the target population by using the one-step estimator proposed in section 2.4.2. The nuisance parameters in this approach are estimated by using the Super Learner algorithm (Van der Laan *and others*, 2007), whose library includes an intercept only model, a main terms general linear model, and extreme gradient boosting. We then use the non-parametric heterogeneity assessment approach proposed in section 3.2 to investigate the impact of mediator- and outcome-related heterogeneity on the variability of the transported effect estimates.

Figure 3 gives the log relative risks of developing a psychiatric disorder comparing moving with the voucher versus not, predicted for the target MTO population in LA. The predicted effects transported to LA vary depending on which other MTO sites are used as the source populations for the mediation model and the outcome model. Specifically, when the outcome model is fixed at one given site, the predicted effects transported to LA seem to be quite homogeneous regardless of which site is used for the mediation model. In contrast, when the MGM is fixed at one given site, using Boston as the source site for the outcome model generates very different predicted effect for LA, as compared to using either Chicago or NYC as the source site for the outcome model. This suggests that important heterogeneity might exist in the outcome model, but not in the mediation model across data sources.

The above observation is indeed confirmed by results of the heterogeneity assessment. Using the proposed non-parametric approach, we obtain an outcome-related heterogeneity variance of 0.354, much larger than the noise, 0.195, which is defined as the average of the variances of all transported effect estimates. In contrast, the mediator-related heterogeneity variance is 0.057, about 4 times smaller than the noise. Based on these findings, we suggest that one needs to be cautious about what site to use for the outcome source model as there might be important heterogeneity between the outcome model for Boston compared to other sites. Simply merging data from all sites and use these to estimate the treatment effect in the target population, LA, will overlook such heterogeneity.




Figure 3: Predicted log relative risks of developing a psychiatric disorder in adolescence among boys comparing moving with a Section 8 voucher versus not in LA (i.e. the target population), using varying MTO sites as the source population for the mediation model and for the outcome model. In each line, the first site denotes the source population for the outcome model, and the second site denotes the source population for the mediation model. All results were approved for release by the U.S. Census Bureau, authorization number CBDRB-FY22-CES018-013.

6 Conclusion

Heterogeneity across data sources is an important challenge in data fusion. In this paper, we have proposed novel methods to identify and assess the impact of heterogeneity when the mediator and outcome data from multiple sources are transported to a target population, with aim to estimate the average treatment effect in such population. Instead of combining data from these external sources (and overlooking the potential heterogeneity between populations), we here suggest that data from each source should be transported separately to the target population. Evaluating the variability across obtained effect estimates can then shed insights on the amount and source of heterogeneity between populations, and whether combining (or averaging) all transported effect estimates is a valid option to obtain a final estimate for the ATE in the target population. While in this paper, we specifically focus on the ATE, the proposed framework can be easily extended to other estimands of the target population, such as those discussed by Li and Luedtke (2021). One can also decompose the *M*-related heterogeneity into different components when there are multiple mediators, and the causal structure among these mediators are specified (see appendix A3 for more details).

One challenge, however, is that the proposed approaches require a stronger positivity assumption to hold than when all data sources are combined into one. For instance, there must be sufficient overlap in the baseline covariate distribution (i.e. case-mix) between the target population and each data source. When all data sources are merged, the case-mix overlap between the resulting dataset and the target population might be better. In practice, it might be the case that positivity is violated for separate data sources but not when all sources are combined. This combination, however, implicitly assumes no heterogeneity across data sources. As we argued throughout this paper, such an assumption is quite unlikely to hold in practice, especially when the number of external data sources is high.

Acknowledgement

This research was conducted as a part of the U.S. Census Bureau's Evidence Building Project Series. The U.S. Census Bureau has not reviewed the paper for accuracy or reliability and does not endorse its contents. Any conclusions expressed herein are those of the authors and do not necessarily represent the views of the U.S. Census Bureau. All results were approved for release by the U.S. Census Bureau, authorization numbers CBDRB-FY22-CES018-013. KER time was supported by R01DA053243, R00DA042127.

References

- BAKBERGENULY, ILYAS, HOAGLIN, DAVID C AND KULINSKAYA, ELENA. (2020). Methods for estimating between-study variance and overall effect in meta-analysis of odds ratios. *Research Synthesis Methods* **11**(3), 426–442.
- BAREINBOIM, ELIAS AND PEARL, JUDEA. (2016). Causal inference and the data-fusion problem. *Proceedings of the National Academy of Sciences* **113**(27), 7345–7352.
- BELLONI, ALEXANDRE, CHERNOZHUKOV, VICTOR AND KATO, KENGO. (2015). Uniform postselection inference for least absolute deviation regression and other z-estimation problems. *Biometrika* **102**(1), 77–94.
- BENKESER, DAVID AND VAN DER LAAN, MARK. (2016). The highly adaptive lasso estimator. In: 2016 IEEE international conference on data science and advanced analytics (DSAA). IEEE. pp. 689–696.
- COLNET, BÉNÉDICTE, MAYER, IMKE, CHEN, GUANHUA, DIENG, AWA, LI, RUOHONG, VAROQUAUX, GAËL, VERT, JEAN-PHILIPPE, JOSSE, JULIE AND YANG, SHU. (2020). Causal inference methods for combining randomized trials and observational studies: a review. *arXiv preprint arXiv:2011.08047*.
- DAHABREH, ISSA J, PETITO, LUCIA C, ROBERTSON, SARAH E, HERNÁN, MIGUEL A AND STEINGRIMS-SON, JON A. (2020). Towards causally interpretable meta-analysis: transporting inferences from multiple randomized trials to a new target population. *Epidemiology (Cambridge, Mass.)* **31**(3), 334.
- DAHABREH, ISSA J, ROBERTSON, SARAH E, PETITO, LUCIA C, HERNÁN, MIGUEL A AND STEIN-GRIMSSON, JON A. (2019). Efficient and robust methods for causally interpretable meta-analysis: Transporting inferences from multiple randomized trials to a target population. *Biometrics*.
- DEVELOPMENT, US DEPARTMENT OF HOUSING URBAN. (2022, Jan). Housing choice voucher program section 8.
- FERRI-GARCÍA, RAMÓN AND RUEDA, MARÍA DEL MAR. (2020). Propensity score adjustment using machine learning classification algorithms to control selection bias in online surveys. *PloS* one 15(4), e0231500.

- HIGGINS, JULIAN PT, THOMAS, JAMES, CHANDLER, JACQUELINE, CUMPSTON, MIRANDA, LI, TIAN-JING, PAGE, MATTHEW J AND WELCH, VIVIAN A. (2019). *Cochrane handbook for systematic reviews of interventions*. John Wiley & Sons.
- HIGGINS, JULIAN PT, THOMPSON, SIMON G, DEEKS, JONATHAN J AND ALTMAN, DOUGLAS G. (2003). Measuring inconsistency in meta-analyses. *Bmj* **327**(7414), 557–560.
- HIGGINS, JULIAN PT, THOMPSON, SIMON G AND SPIEGELHALTER, DAVID J. (2009). A re-evaluation of random-effects meta-analysis. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* **172**(1), 137–159.
- KLING, JEFFREY R, LIEBMAN, JEFFREY B AND KATZ, LAWRENCE F. (2007). Experimental analysis of neighborhood effects. *Econometrica* **75**(1), 83–119.
- LI, SIJIA AND LUEDTKE, ALEX. (2021). Efficient estimation under data fusion. arXiv preprint arXiv:2111.14945.
- LUEDTKE, ALEX, CARONE, MARCO AND VAN DER LAAN, MARK J. (2019). An omnibus nonparametric test of equality in distribution for unknown functions. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **81**(1), 75–99.
- RACINE, JEFFERY S, HART, JEFFREY AND LI, QI. (2006). Testing the significance of categorical predictor variables in nonparametric regression models. *Econometric Reviews* **25**(4), 523–544.
- RUDOLPH, KARA E AND DÍAZ, IVÁN. (2022). Efficiently transporting causal direct and indirect effects to new populations under intermediate confounding and with multiple mediators. *Biostatistics* **23**(3), 789–806.
- RUDOLPH, KARA E, LEVY, JONATHAN, SCHMIDT, NICOLE M, STUART, ELIZABETH AND AHERN, JENNIFER. (2020). Using transportability to understand differences in mediation mechanisms across trial sites of a housing voucher experiment. *Epidemiology (Cambridge, Mass.)* **31**(4), 523.
- RUDOLPH, KARA E, LEVY, JONATHAN AND VAN DER LAAN, MARK J. (2021). Transporting stochastic direct and indirect effects to new populations. *Biometrics* **77**(1), 197–211.
- RUDOLPH, KARA E, SCHMIDT, NICOLE M, GLYMOUR, M MARIA, CROWDER, REBECCA, GALIN, JESSICA, AHERN, JENNIFER AND OSYPUK, THERESA L. (2018*a*). Composition or context: using transportability to understand drivers of site differences in a large-scale housing experiment. *Epidemiology (Cambridge, Mass.)* 29(2), 199.
- RUDOLPH, KARA E, SOFRYGIN, OLEG, SCHMIDT, NICOLE M, CROWDER, REBECCA, GLYMOUR, M MARIA, AHERN, JENNIFER AND OSYPUK, THERESA L. (2018*b*). Mediation of neighborhood effects on adolescent substance use by the school and peer environments. *Epidemiology* **29**(4), 590–598.
- RUDOLPH, KARA E AND VAN DER LAAN, MARK J. (2017). Robust estimation of encouragement design intervention effects transported across sites. *Journal of the Royal Statistical Society:* Series B (Statistical Methodology) **79**(5), 1509–1525.
- SANBONMATSU, LISA, LUDWIG, JENS, KATZ, LAWRENCE F, GENNETIAN, LISA A, DUNCAN, GREG J, KESSLER, RONALD C, ADAM, EMMA, MCDADE, THOMAS W AND LINDAU, STACY TESSLER. (2011). Moving to Opportunity for Fair Housing Demonstration Program-Final Impacts Evaluation. Washington, DC: US Department of Housing and Urban Development, Office of Policy Development and Research.
- SOBEL, MICHAEL, MADIGAN, DAVID AND WANG, WEI. (2017). Causal inference for meta-analysis and multi-level data structures, with application to randomized studies of vioxx. *Psychometrika* **82**(2), 459–474.
- STUART, ELIZABETH A, COLE, STEPHEN R, BRADSHAW, CATHERINE P AND LEAF, PHILIP J. (2011). The use of propensity scores to assess the generalizability of results from randomized trials. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* 174(2), 369–386.

- UEHARA, MASATOSHI, KATO, MASAHIRO AND YASUI, SHOTA. (2020). Off-policy evaluation and learning for external validity under a covariate shift. *Advances in Neural Information Processing Systems* **33**, 49–61.
- VAN DER LAAN, MARK J, POLLEY, ERIC C AND HUBBARD, ALAN E. (2007). Super learner. *Statistical applications in genetics and molecular biology* **6**(1).
- Vo, TAT-THANG, PORCHER, RAPHAEL, CHAIMANI, ANNA AND VANSTEELANDT, STIJN. (2019). A novel approach for identifying and addressing case-mix heterogeneity in individual participant data meta-analysis. *Research synthesis methods* **10**(4), 582–596.
- Vo, TAT-THANG, PORCHER, RAPHAËL AND VANSTEELANDT, STIJN. (2021). Assessing the impact of case-mix heterogeneity in individual participant data meta-analysis: Novel use of i 2 statistic and prediction interval. *Research Methods in Medicine & Health Sciences* **2**(1), 12–30.
- WALTER, STEPHEN AND BALAKRISHNAN, NARAYANASWAMY. (2022). The usual method of estimating weights for studies in meta-analyses is biased. *Available at SSRN 4104102*.