

Three-loop master integrals for the Higgs boson self-energy with internal top-quarks and W-bosons

Ekta Chaubey ^a, Ina Hönemann ^b and Stefan Weinzierl ^b

^a *Physics Department, University of Turin and INFN Turin,
Via Pietro Giuria 1, I-10125, Turin, Italy*

^b *PRISMA Cluster of Excellence, Institut für Physik,
Johannes Gutenberg-Universität Mainz,
D - 55099 Mainz, Germany*

Abstract

We consider the full set of master integrals with internal top- and W -propagators contributing to the three-loop Higgs self-energy diagrams of order $O(\alpha^2\alpha_s)$. We split the master integrals into a system relevant to the Feynman diagrams proportional to the product of Yukawa couplings $y_b y_t$ and the complement. For both systems we define master integrals of uniform weight, such that the associated differential equation is in ϵ -factorised form. The occurring square roots are rationalised and all master integrals are expressible in multiple polylogarithms.

1 Introduction

Higgs precision physics is at the core of the experimental LHC program. On the theoretical side this requires higher-order calculations in perturbation theory [1–12]. The Higgs boson self-energy is an essential building block for many observables. Through renormalisation it enters the Higgs mass definition. Furthermore, the imaginary part of the Higgs boson self-energy is directly related through the optical theorem to the Higgs decay rate [13–29].

In this article we consider three-loop contributions with internal top- and W -propagators to the Higgs boson self-energy. Due to the presence of the internal top- and W -propagators the corresponding Feynman diagrams are among the more complicated Feynman integrals contributing at this order in perturbation theory. These integrals have been computed for $p^2 = m_H^2$ approximately as an expansion in m_H^2/m_t^2 in ref. [23].

In this paper we present the analytic calculation of the relevant master integrals. We keep the full dependence on the external momentum squared p^2 and the heavy particle masses (m_t and m_W), but neglect the b -quark mass. The Feynman integrals depend on two dimensionless variables, which we may take initially as

$$v = \frac{p^2}{m_t^2}, \quad w = \frac{m_W^2}{m_t^2}. \quad (1)$$

It is convenient to divide the Feynman integrals into two sets: A set of 105 master integrals corresponding to the set of Feynman diagrams being proportional to the product of Yukawa couplings $y_b y_t$ and the complement, consisting of 33 additional master integrals. We treat the first set of 105 master integrals in the main part of the paper and the 33 additional integrals in appendix A. The reason is that the sets of occurring square roots differ. In total we encounter four different square roots r_1 – r_4 . While we are not able to rationalise simultaneously all four square roots, we may set up three independent systems of differential equations, such that within one system all square roots can be rationalised.

In the main part of the paper we treat the set of 105 master integrals corresponding to the set of Feynman diagrams being proportional to the product of Yukawa couplings $y_b y_t$. We define a basis of master integrals of uniform weight. In this basis, the associated differential equation is ε -factorised and all occurring differential one-forms are dlog-forms. In this system we encounter two square roots, which can be rationalised simultaneously. This is done by a change of variables from (v, w) to (x, y) . The variables x and y will be defined in the main part of this paper. This change of variables allows us to express all master integrals to any order in the dimensional regularisation parameter ε in terms of multiple polylogarithms. The alphabet for the multiple polylogarithms consists of 21 letters.

The remaining master integrals, relevant to Feynman diagrams with internal top- and W -propagators, where the Higgs boson couples either to a W -boson and a b -quark or to two b -quarks are discussed in appendix A. The calculation proceeds exactly along the same lines as for the first set. The only difference is that additional square roots occur, which require different rationalisations.

This paper is organised as follows: In section 2 we introduce our notation for the three-loop integrals. The master integrals of uniform weight contributing to Feynman diagrams proportional




Figure 1: Examples of three-loop Higgs self-energy diagrams containing a top propagator and a W -propagator and being proportional to the product of Yukawa couplings $y_b y_t$. The internal masses of the propagators are encoded by the colour of the propagators: massless (black), m_t (green), m_W (red).

to the Yukawa couplings $y_b y_t$ are presented in section 3. The differential equation for these master integrals is in ϵ -factorised form. The differential equation is discussed in section 4. The solution of the differential equation requires boundary values, which we give in section 5. Analytical results for the master integrals are presented in section 6. Numerical results are presented in section 7. Finally, our conclusions are given in section 8. Appendix A presents the additional master integrals needed for the contributions not proportional to the Yukawa couplings $y_b y_t$. In appendix B we describe the content of the supplementary electronic file attached to the arxiv version of this article.

2 Notation

We consider three-loop Higgs self-energy diagrams of order $O(\alpha^2 \alpha_s)$ proportional to the product of Yukawa couplings $y_b y_t$. Examples of diagrams are shown in fig. 1. The diagrams shown in fig. 1 have all eight propagators and define the top sectors. Simpler diagrams, obtained from the ones shown in fig. 1 by pinching a subset of the propagators, are not shown.

In this paper we calculate all relevant master integrals. As we have three independent loop momenta and one independent external momentum, we need nine propagators in order to be

able to express any scalar product involving at least one loop momentum in terms of inverse propagators. We therefore consider auxiliary graphs with nine propagators, shown in fig. 2. We label these auxiliary graphs topology A, B, C and D in correspondence with the diagrams shown in fig. 1.

As a consequence we consider the integrals

$$I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X = e^{3\gamma_E \varepsilon} (\mu^2)^{v - \frac{3}{2}D} \int \frac{d^D k_1}{i\pi^{\frac{D}{2}}} \frac{d^D k_2}{i\pi^{\frac{D}{2}}} \frac{d^D k_3}{i\pi^{\frac{D}{2}}} \prod_{j=1}^9 \frac{1}{(P_j^X)^{v_j}},$$

$$X \in \{A, B, C, D\}, \quad (2)$$

where $D = 4 - 2\varepsilon$ denotes the number of space-time dimensions, γ_E denotes the Euler-Mascheroni constant, μ is an arbitrary scale introduced to render the Feynman integral dimensionless, and the quantity v is defined by

$$v = \sum_{j=1}^9 v_j. \quad (3)$$

The inverse propagators P_j^X are defined as follows:

Topology A :

$$\begin{aligned} P_1^A &= -k_1^2 + m_t^2, & P_2^A &= -(k_1 - p)^2 + m_t^2, & P_3^A &= -(k_1 + k_2)^2, \\ P_4^A &= -k_2^2 + m_t^2, & P_5^A &= -(k_2 + k_3)^2 + m_W^2, & P_6^A &= -(k_2 + p)^2 + m_t^2, \\ P_7^A &= -k_3^2, & P_8^A &= -(k_3 - p)^2, & P_9^A &= -(k_1 - k_3)^2 + m_t^2. \end{aligned} \quad (4)$$

Topology B :

$$\begin{aligned} P_1^B &= -k_1^2 + m_t^2, & P_2^B &= -(k_1 - p)^2 + m_t^2, & P_3^B &= -(k_1 + k_2)^2 + m_W^2, \\ P_4^B &= -k_2^2, & P_5^B &= -(k_2 + k_3)^2, & P_6^B &= -(k_2 + p)^2, \\ P_7^B &= -k_3^2, & P_8^B &= -(k_3 - p)^2, & P_9^B &= -(k_1 - k_3)^2 + m_W^2. \end{aligned} \quad (5)$$

Topology C :

$$\begin{aligned} P_1^C &= -k_1^2 + m_t^2, & P_2^C &= -(k_1 - p)^2 + m_t^2, & P_3^C &= -(k_1 + k_2)^2, \\ P_4^C &= -(k_1 + k_2 - k_3)^2, & P_5^C &= -k_2^2 + m_W^2, & P_6^C &= -(k_2 - k_3 + p)^2 + m_t^2, \\ P_7^C &= -k_3^2, & P_8^C &= -(k_3 - p)^2, & P_9^C &= -(k_2 + p)^2 + m_t^2. \end{aligned} \quad (6)$$

Topology D :

$$\begin{aligned} P_1^D &= -k_1^2 + m_t^2, & P_2^D &= -(k_1 - p)^2 + m_t^2, & P_3^D &= -(k_1 - k_3)^2 + m_W^2, \\ P_4^D &= -(k_1 + k_2)^2 + m_t^2, & P_5^D &= -k_2^2, & P_6^D &= -(k_2 + k_3)^2, \\ P_7^D &= -k_3^2, & P_8^D &= -(k_3 - p)^2, & P_9^D &= -(k_2 + p)^2. \end{aligned} \quad (7)$$




Figure 2: Graphs for the four topologies. The internal masses of the propagators are encoded by the colour of the propagators: massless (black), m_t (green), m_W (red).

Topology and sector id	N_{master}
A_{255}	50
$A_{255} + B_{255}$	67
$A_{255} + B_{255} + C_{255}$	104
$A_{255} + B_{255} + C_{255} + D_{255}$	105

(9)

Table 1: The number of master integrals. The table gives the cumulative sum, as for example topology B contains master integrals common to topology A and B .

Topology and sector id	N_{master}
A_{255}	50
B_{255}	32
C_{255}	95
D_{255}	67

(10)

Table 2: The number of master integrals for a given top sector, if we treat each top sector individually.

We introduce the sector id for the Feynman integral $I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X$ by

$$N_{\text{id}}^X = \sum_{j=1}^9 2^{j-1} \Theta(v_j > 0). \quad (8)$$

With the help of the sector id we may specify all the reductions which we will need: We have to reduce in every topology $A - D$ the top sector 255. These top sectors correspond to the diagrams shown in fig. 1. The integral reduction gives us in total 105 master integrals. A detailed list is given in table 1. Note that this list gives the cumulative number of master integrals. This takes into account that certain master integral occur in more than one topology or sector. These master integrals are only counted once. If on the other hand we treat each top sector individually, we find the number of master integrals per top sector as shown in table 2. In this case the total number of master integrals is higher, as we recompute master integrals, which are identical or are related by symmetries. The integral reduction can be carried out with public available computer programs like FIRE/Litered [30–33], Reduze [34, 35], Kira [36, 37] or FiniteFlow [38, 39].

Let us point out, that it is non-trivial that the total number of master integrals is 105 and not 106. There is one relation not easily found by standard integration-by-parts reduction programs¹:

$$I_{011011100}^B = \frac{1}{(m_t^2 - m_W^2)} (I_{001011100}^B - I_{100011100}^B)$$

¹We thank Johann Usovitsch for contributing this relation.

$$\begin{aligned}
& -\frac{1}{s} \left(I_{01111(-1)100}^A - I_{011010100}^A - I_{(-1)11110100}^A \right) \\
& -\frac{3}{s(m_t^2 - m_W^2)} \left(I_{011(-1)10100}^A - I_{01101(-1)100}^A \right). \tag{11}
\end{aligned}$$

Without this relation we may end up with a non-integrable system.

The Feynman parameter representations for the four topologies are given by

$$I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X = e^{3\gamma_E \varepsilon} \frac{\Gamma(v - \frac{3}{2}D)}{9 \prod_{j=1}^9 \Gamma(v_j)} \int_{x_i \geq 0} d^9 x \delta \left(1 - \sum_{j=1}^9 x_j \right) \left(\prod_{j=1}^9 x_j^{v_j - 1} \right) \frac{\mathcal{U}_X^{v-2D}}{\mathcal{F}_X^{v-\frac{3}{2}D}}, \tag{12}$$

where \mathcal{U}_X and \mathcal{F}_X denote the first and second graph polynomial, respectively. The first graph polynomials for the individual topologies are given by

$$\begin{aligned}
\mathcal{U}_A &= \mathcal{U}_B = x_{12} x_3 x_5 + x_{12} x_3 x_7 + x_{12} x_3 x_9 + x_{12} x_4 x_5 + x_{12} x_4 x_7 + x_{12} x_4 x_9 + x_{12} x_5 x_7 \\
&+ x_{12} x_5 x_9 + x_3 x_4 x_5 + x_3 x_4 x_7 + x_3 x_4 x_9 + x_3 x_5 x_7 + x_3 x_7 x_9 + x_4 x_5 x_9 + x_4 x_7 x_9 \\
&+ x_5 x_7 x_9, \\
\mathcal{U}_C &= x_{12} x_3 x_4 + x_{12} x_3 x_6 + x_{12} x_3 x_7 + x_{12} x_4 x_5 + x_{12} x_4 x_7 + x_{12} x_5 x_6 + x_{12} x_5 x_7 \\
&+ x_{12} x_6 x_7 + x_3 x_4 x_5 + x_3 x_4 x_6 + x_3 x_5 x_6 + x_3 x_5 x_7 + x_3 x_6 x_7 + x_4 x_5 x_6 + x_4 x_5 x_7 \\
&+ x_4 x_6 x_7, \\
\mathcal{U}_D &= x_{12} x_3 x_4 + x_{12} x_3 x_5 + x_{12} x_3 x_6 + x_{12} x_4 x_6 + x_{12} x_4 x_7 + x_{12} x_5 x_6 + x_{12} x_5 x_7 \\
&+ x_{12} x_6 x_7 + x_3 x_4 x_5 + x_3 x_4 x_7 + x_3 x_5 x_6 + x_3 x_5 x_7 + x_3 x_6 x_7 + x_4 x_5 x_6 + x_4 x_5 x_7 \\
&+ x_4 x_6 x_7, \tag{13}
\end{aligned}$$

where we used the abbreviation $x_{ij} = x_i + x_j$.

2.1 Dimensional shift relations and differential equations

Let us introduce an operator \mathbf{i}^+ , which raises the power of the propagator i by one and multiplies by v_i , e.g.

$$\mathbf{1}^+ I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X = v_1 \cdot I_{(v_1+1)V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X. \tag{14}$$

In addition we define two operators \mathbf{D}^\pm , which shift the dimension of space-time by two through

$$\mathbf{D}^\pm I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X(D) = I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X(D \pm 2). \tag{15}$$

The dimensional shift relations read [40, 41]

$$\begin{aligned}
\mathbf{D}^- I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X(D) &= \\
& \mathcal{U}_X(\mathbf{1}^+, \mathbf{2}^+, \mathbf{3}^+, \mathbf{4}^+, \mathbf{5}^+, \mathbf{6}^+, \mathbf{7}^+, \mathbf{8}^+, \mathbf{9}^+) I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X(D). \tag{16}
\end{aligned}$$

We denote by $J_{m_i}^X$ the set of internal edge indices, which correspond to the propagation of a particle with mass m_i . For example

$$J_{m_t}^A = \{1, 2, 4, 6, 9\}. \quad (17)$$

With this notation the differential equations read

$$\begin{aligned} \mu^2 \frac{\partial}{\partial m_t^2} I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X &= - \sum_{j \in J_{m_t}^X} \mathbf{j}^+ I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X, \\ \mu^2 \frac{\partial}{\partial m_W^2} I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X &= - \sum_{j \in J_{m_W}^X} \mathbf{j}^+ I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X, \\ \mu^2 \frac{\partial}{\partial s} I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X &= \left(\frac{3}{2} D - v \right) \frac{\mu^2}{s} I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X + \frac{m_t^2}{s} \sum_{j \in J_{m_t}^X} \mathbf{j}^+ I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X \\ &\quad + \frac{m_W^2}{s} \sum_{j \in J_{m_W}^X} \mathbf{j}^+ I_{V_1 V_2 V_3 V_4 V_5 V_6 V_7 V_8 V_9}^X. \end{aligned} \quad (18)$$

For the derivative with respect to s we used the scaling relation. This has the advantage that on the right-hand side at most one raising operator occurs. This minimises the integral reductions we have to compute.

2.2 Variables

In the following we will set

$$\mu^2 = m_t^2. \quad (19)$$

The master integrals depend then kinematically on two dimensionless quantities. We set

$$v = \frac{p^2}{m_t^2}, \quad w = \frac{m_W^2}{m_t^2}. \quad (20)$$

We will encounter the following two square roots

$$r_1 = \sqrt{-v(4-v)}, \quad r_2 = \sqrt{\lambda(v, w, 1)}. \quad (21)$$

The Källén function is defined by

$$\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2yz - 2zx. \quad (22)$$

The variable transformation from (v, w) to (x, y) , where the latter variables are defined by [42–44]

$$v = -\frac{(1-x)^2}{x}, \quad w = \frac{(1-y+2xy)(x-2y+xy)}{x(1-y^2)}, \quad (23)$$

rationalises the square roots r_1 and r_2 :

$$r_1 = \frac{1-x^2}{x}, \quad r_2 = \frac{(1-x) \left[(1-y)^2 + x(1+y)^2 \right]}{x(1-y^2)}. \quad (24)$$

The inverse transformation is given by

$$x = \frac{1}{2}(2-v-r_1), \quad y = \frac{r_2-r_1}{1-w+2v}, \quad (25)$$

such that $x = 0$ corresponds to $v = \infty$ and $y = 0$ corresponds to $w = 1$. We note that the point $(v, w) = (0, 1)$ is blown up in (x, y) -space to the hypersurface $x = 1$. This is convenient, as it allows us to obtain the results for the master integrals by integrating the differential equation in x with constant y . We make a final change of coordinates and we introduce

$$x' = 1-x. \quad (26)$$

This transforms the starting point of the integration from $x = 1$ to $x' = 0$ and is helpful for the conversion to multiple polylogarithms.

3 Master integrals

In this section we define master integrals of uniform weight. This is the main result of this paper.

We have 105 master integrals and 59 master topologies (or sectors). These are summarised in table 3 and 4. We denote the master integrals of uniform weight by J_1 - J_{105} .

Existing deterministic algorithms [45–50] for the construction of master integrals of uniform weight cannot be used for problems of this complexity. Instead, we revert to heuristic methods. We first note that given an educated guess for a basis of master integrals of uniform weight, it is easy to verify/falsify if the proposed basis has the desired properties: One simply computes the differential equation in this basis. For the educated guess we study the maximal cuts in the loop-by-loop Baikov representation [51, 52] and construct integrands, which give to leading order in ϵ constants when integrated over a basis of independent contours [53, 54]. In a second step, possible contributions from sub-topologies are fixed by an ansatz. A pedagogical description of this method can be found in chapter 16 of [55].

For all these master integrals we find compact expressions in terms of pre-canonical Feynman integrals. It is worth noting that all prefactors are polynomials in ϵ, v, w, r_1, r_2 , i.e. no denominators occur. The master integrals J_1 - J_{105} are defined by

$$\begin{aligned} J_1 &= \epsilon^3 \mathbf{D}^- I_{100110000}^A, \\ J_2 &= \epsilon^3 r_1 \mathbf{D}^- I_{110110000}^A, \\ J_3 &= \epsilon^3 r_1 \mathbf{D}^- I_{011110000}^A, \\ J_4 &= \epsilon^3 \mathbf{D}^- I_{(-1)11110000}^A, \\ J_5 &= \epsilon^3 (1-w) \mathbf{D}^- I_{101010100}^A, \end{aligned}$$

$$\begin{aligned}
J_6 &= \varepsilon^3 \mathbf{D}^- I_{101(-1)10100}^A, \\
J_7 &= \varepsilon^3 r_2 \mathbf{D}^- I_{011010100}^A, \\
J_8 &= \varepsilon^3 \left[\mathbf{D}^- I_{011(-1)10100}^A - (1-w) \mathbf{D}^- I_{011010100}^A \right], \\
J_9 &= \varepsilon^3 \mathbf{D}^- I_{01101(-1)100}^A, \\
J_{10} &= \varepsilon^3 \left[\mathbf{D}^- I_{0110101(-1)0}^A + \nu \mathbf{D}^- I_{011010100}^A \right], \\
J_{11} &= \varepsilon^3 (1-w) \mathbf{D}^- I_{100110100}^A, \\
J_{12} &= \varepsilon^3 r_2 \mathbf{D}^- I_{100011100}^A, \\
J_{13} &= \varepsilon^3 \left[\mathbf{D}^- I_{100(-1)11100}^A - (1-w) \mathbf{D}^- I_{100011100}^A \right], \\
J_{14} &= \varepsilon^3 \left[\mathbf{D}^- I_{1000111(-1)0}^A + \nu \mathbf{D}^- I_{100011100}^A \right], \\
J_{15} &= \varepsilon^3 \nu \mathbf{D}^- I_{100100110}^A, \\
J_{16} &= \varepsilon^3 \nu \mathbf{D}^- I_{100010110}^A, \\
J_{17} &= \varepsilon^3 \nu(4-\nu) \mathbf{D}^- I_{110111000}^A, \\
J_{18} &= \varepsilon^3 r_1 \left[\mathbf{D}^- I_{111(-1)10100}^A - (1-w) \mathbf{D}^- I_{111010100}^A \right], \\
J_{19} &= \varepsilon^3 r_1 \left[\mathbf{D}^- I_{111(-1)10100}^A - \mathbf{D}^- I_{011010100}^A \right], \\
J_{20} &= \varepsilon^3 (1-w) r_1 \mathbf{D}^- I_{110110100}^A, \\
J_{21} &= \varepsilon^3 r_1 \left[(1-w) \mathbf{D}^- I_{011110100}^A - \mathbf{D}^- I_{011010100}^A \right], \\
J_{22} &= \varepsilon^3 (1-w) \left[\mathbf{D}^- I_{(-1)11110100}^A - \mathbf{D}^- I_{011010100}^A \right], \\
J_{23} &= \varepsilon^3 (1-w) \mathbf{D}^- I_{01111(-1)100}^A, \\
J_{24} &= \varepsilon^3 r_1 r_2 \mathbf{D}^- I_{110011100}^A, \\
J_{25} &= \varepsilon^3 r_1 \left[\mathbf{D}^- I_{110(-1)11100}^A - (1-w) \mathbf{D}^- I_{110011100}^A \right], \\
J_{26} &= \varepsilon^3 r_1 \left[\mathbf{D}^- I_{1100111(-1)0}^A + \nu \mathbf{D}^- I_{110011100}^A \right], \\
J_{27} &= \varepsilon^3 r_2 \mathbf{D}^- I_{101(-1)11100}^A, \\
J_{28} &= \varepsilon^3 r_1 \left[\mathbf{D}^- I_{101(-1)11100}^A - (1-w) \mathbf{D}^- I_{101011100}^A \right], \\
J_{29} &= \varepsilon^3 \left[\mathbf{D}^- I_{101(-2)11100}^A - (1-w) \mathbf{D}^- I_{101(-1)11100}^A \right], \\
J_{30} &= \varepsilon^3 (1-\nu) \left[\mathbf{D}^- I_{101(-1)11100}^A + \mathbf{D}^- I_{10101110(-1)}^A - (1-w) \mathbf{D}^- I_{101011100}^A \right], \\
J_{31} &= \varepsilon^3 r_1 \left[(1-w) \mathbf{D}^- I_{100111100}^A - \mathbf{D}^- I_{100011100}^A \right], \\
J_{32} &= \varepsilon^3 \nu r_1 \mathbf{D}^- I_{110100110}^A, \\
J_{33} &= \varepsilon^3 \nu r_1 \mathbf{D}^- I_{011100110}^A,
\end{aligned}$$

$$\begin{aligned}
J_{34} &= \varepsilon^3 v \mathbf{D}^- I_{(-1)11100110}^A, \\
J_{35} &= \varepsilon^3 v r_1 \mathbf{D}^- I_{110010110}^A, \\
J_{36} &= \varepsilon^3 (1-w) \left[v \mathbf{D}^- I_{101010110}^A + \mathbf{D}^- I_{011010100}^A \right], \\
J_{37} &= \varepsilon^4 (1-2\varepsilon) v I_{111021100}^A, \\
J_{38} &= \varepsilon^4 (1-2\varepsilon) v I_{111011200}^A, \\
J_{39} &= \varepsilon^3 v (4-v) \left[(1-w) \mathbf{D}^- I_{110111100}^A - \mathbf{D}^- I_{110011100}^A \right], \\
J_{40} &= \varepsilon^3 v r_1 \left[\mathbf{D}^- I_{11101011(-1)}^A - (1-w) \mathbf{D}^- I_{111010110}^A \right], \\
J_{41} &= \varepsilon^4 (1-2\varepsilon) v I_{112010110}^A, \\
J_{42} &= \varepsilon^4 (1-2\varepsilon) v I_{111020110}^A, \\
J_{43} &= \varepsilon^4 (1-2\varepsilon) v I_{021110110}^A, \\
J_{44} &= \varepsilon^4 (1-2\varepsilon) v I_{012110110}^A, \\
J_{45} &= \varepsilon^3 v^2 (4-v) \mathbf{D}^- I_{110101110}^A, \\
J_{46} &= \varepsilon^4 (1-\varepsilon) (1-2\varepsilon) v I_{100111110}^A, \\
J_{47} &= \varepsilon^5 (1-2\varepsilon) v I_{111110110}^A, \\
J_{48} &= \varepsilon^4 (1-2\varepsilon) v (1-w) I_{111120110}^A, \\
J_{49} &= \varepsilon^4 (1-2\varepsilon) v \left[I_{211110110}^A + \frac{1}{2} (2-v+r_1) I_{121110110}^A \right], \\
J_{50} &= \varepsilon^4 (1-2\varepsilon) v r_1 I_{210111110}^A, \\
J_{51} &= \varepsilon^3 v \mathbf{D}^- I_{100011100}^B, \\
J_{52} &= \varepsilon^3 v \mathbf{D}^- I_{001011100}^B, \\
J_{53} &= \varepsilon^3 v r_1 \mathbf{D}^- I_{110011100}^B, \\
J_{54} &= \varepsilon^3 r_2 \mathbf{D}^- I_{101(-1)11100}^B, \\
J_{55} &= \varepsilon^3 (1-w) \left[\mathbf{D}^- I_{101(-1)11100}^B + v \mathbf{D}^- I_{101011100}^B \right], \\
J_{56} &= \varepsilon^3 \left[\mathbf{D}^- I_{101(-2)11100}^B + v \mathbf{D}^- I_{101(-1)11100}^B \right], \\
J_{57} &= \varepsilon^3 v r_2 \mathbf{D}^- I_{011100110}^B, \\
J_{58} &= \varepsilon^3 v \left[\mathbf{D}^- I_{(-1)11100110}^B - (1-w) \mathbf{D}^- I_{011100110}^B \right], \\
J_{59} &= \varepsilon^3 v \mathbf{D}^- I_{0111(-1)0110}^B, \\
J_{60} &= \varepsilon^3 v^2 \mathbf{D}^- I_{100101110}^B, \\
J_{61} &= \varepsilon^3 v^2 \mathbf{D}^- I_{001101110}^B, \\
J_{62} &= \varepsilon^4 (1-2\varepsilon) v I_{111021100}^B, \\
J_{63} &= \varepsilon^3 r_1 \left\{ (1-w) \mathbf{D}^- I_{111(-1)11100}^B - \mathbf{D}^- I_{101(-1)11100}^B \right\},
\end{aligned}$$

$$\begin{aligned}
J_{64} &= \varepsilon^3 v r_1 \{ (1-w) \mathbf{D}^- I_{111100110}^B - \mathbf{D}^- I_{011100110}^B \}, \\
J_{65} &= \varepsilon^3 v^2 r_1 \mathbf{D}^- I_{110101110}^B, \\
J_{66} &= \varepsilon^5 (1-2\varepsilon) v I_{111110110}^B, \\
J_{67} &= \varepsilon^4 (1-2\varepsilon)^2 v I_{111101110}^B, \\
J_{68} &= \varepsilon^3 r_1 \mathbf{D}^- I_{101001100}^C, \\
J_{69} &= \varepsilon^3 \mathbf{D}^- I_{1(-1)1001100}^C, \\
J_{70} &= \varepsilon^3 \mathbf{D}^- I_{10100110(-1)}^C, \\
J_{71} &= \varepsilon^2 (1+4\varepsilon) \mathbf{D}^- I_{011001100}^C, \\
J_{72} &= \varepsilon^3 r_1 \mathbf{D}^- I_{11100110(-1)}^C, \\
J_{73} &= \varepsilon^3 r_2 \left[(1-w) \mathbf{D}^- I_{101011100}^C + \mathbf{D}^- I_{101001100}^C \right], \\
J_{74} &= \varepsilon^3 (1-w) \mathbf{D}^- I_{10101110(-1)}^C, \\
J_{75} &= \varepsilon^3 (1-w) \left[\mathbf{D}^- I_{1010111(-1)0}^C + v \mathbf{D}^- I_{101011100}^C \right], \\
J_{76} &= \varepsilon^3 r_2 \mathbf{D}^- I_{01101110(-1)}^C, \\
J_{77} &= \varepsilon^3 \mathbf{D}^- I_{01101110(-2)}^C, \\
J_{78} &= \varepsilon^3 (w-v) \left\{ \mathbf{D}^- I_{011(-1)11100}^C - 2 \mathbf{D}^- I_{100011100}^A \right\}, \\
J_{79} &= \varepsilon^3 r_1 \left\{ 2 \mathbf{D}^- I_{1(-1)1101010}^C + v \mathbf{D}^- I_{101101010}^C - \mathbf{D}^- I_{011001100}^C - 2 \mathbf{D}^- I_{101100010}^C \right\}, \\
J_{80} &= \varepsilon^3 v \left[2(1-3\varepsilon) \left(I_{202101010}^C - \varepsilon I_{102101010}^C \right) + \varepsilon(1-2\varepsilon) I_{101201010}^C \right], \\
J_{81} &= \varepsilon^3 (1-w) \left[(1-w) \mathbf{D}^- I_{101011010}^C + \mathbf{D}^- I_{011001100}^C \right], \\
J_{82} &= \varepsilon^3 r_1 \left\{ (1-w) \mathbf{D}^- I_{11101110(-1)}^C - \mathbf{D}^- I_{01101110(-1)}^C \right\}, \\
J_{83} &= \varepsilon^4 (1-2\varepsilon) v I_{111011200}^C, \\
J_{84} &= \varepsilon^4 (1-2\varepsilon) v I_{111012100}^C, \\
J_{85} &= \varepsilon^4 (1-2\varepsilon) v I_{112101010}^C, \\
J_{86} &= \varepsilon^3 (1-w) r_1 \left[(1-w) \mathbf{D}^- I_{111011010}^C - \mathbf{D}^- I_{101011100}^C + \mathbf{D}^- I_{111001100}^C \right], \\
J_{87} &= \varepsilon^4 (1-2\varepsilon) \left[I_{101111010}^C - \frac{1}{2} (1-w) I_{201111010}^C - \frac{3}{2} I_{101211(-1)10}^C + \frac{1}{4} v I_{101201010}^C \right. \\
&\quad \left. - \frac{1}{2} I_{201101010}^C + \frac{1}{2} I_{200111010}^C + \frac{3}{2} I_{100211010}^C \right] + \frac{\varepsilon^3}{2} (1-3\varepsilon) v \left[I_{202101010}^C - \varepsilon I_{102101010}^C \right], \\
J_{88} &= \varepsilon^4 (1-2\varepsilon) v I_{101211010}^C, \\
J_{89} &= \varepsilon^4 (1-2\varepsilon) v I_{012111010}^C, \\
J_{90} &= \varepsilon^4 (1-2\varepsilon) v I_{112001110}^C,
\end{aligned}$$

$$\begin{aligned}
J_{91} &= \varepsilon^4 (1 - 2\varepsilon) v I_{111002110}^C, \\
J_{92} &= \varepsilon^2 r_1 \left[(1 - 2\varepsilon)^2 v I_{211001210}^C - \varepsilon (1 - 2\varepsilon) I_{111002110}^C - I_{202002100}^C - v I_{210200210}^A \right], \\
J_{93} &= \varepsilon^4 (1 - 2\varepsilon) v I_{021011110}^C, \\
J_{94} &= \varepsilon^4 (1 - 2\varepsilon) v I_{012011110}^C, \\
J_{95} &= \varepsilon^5 (1 - 2\varepsilon) v I_{111111010}^C, \\
J_{96} &= \varepsilon^4 (1 - 2\varepsilon) vw I_{111121010}^C, \\
J_{97} &= \varepsilon^4 (1 - 2\varepsilon) v(1 - w) I_{111111020}^C, \\
J_{98} &= \varepsilon^4 (1 - 2\varepsilon) r_1 \left(I_{211111010}^C + I_{121111010}^C + I_{111112010}^C + w I_{111121010}^C - 4\varepsilon I_{111111010}^C \right), \\
J_{99} &= \varepsilon^5 (1 - 2\varepsilon) v I_{111011110}^C, \\
J_{100} &= \varepsilon^4 (1 - 2\varepsilon) v(1 - w) I_{111012110}^C, \\
J_{101} &= \varepsilon^4 (1 - 2\varepsilon) v(1 - w) I_{112011110}^C, \\
J_{102} &= \varepsilon^4 (1 - 2\varepsilon) vr_1 I_{121011110}^C, \\
J_{103} &= \varepsilon^5 (1 - 2\varepsilon) v I_{101111110}^C, \\
J_{104} &= \varepsilon^5 (1 - 2\varepsilon) v(1 - v - w) I_{111111110}^C, \\
J_{105} &= \varepsilon^4 (1 - 2\varepsilon) vw I_{111201110}^D. \tag{27}
\end{aligned}$$

4 The differential equation and differential forms

We write the differential equation for the master integrals as

$$dJ = AJ, \tag{28}$$

with

$$A = A_v dv + A_w dw. \tag{29}$$

The matrix-valued one-form A satisfies the integrability condition

$$dA - A \wedge A = 0. \tag{30}$$

For the choice of master integrals as in eq. (27), the differential equation is in ε -form [56] and we have

$$A = \varepsilon \sum_{k=1}^{19} M_k \omega_k, \tag{31}$$

where the M_k 's are 105×105 -matrices, whose entries are rational numbers. The ω_k 's are differential one-forms. We have 19 differential one-forms. All are dlog-forms. The differential one-forms are

$$\omega_1 = d \ln(w),$$

topology	block	sector	propagators	master integrals
<i>A</i>	1	25	1, 4, 5	J_1
	2	27	1, 2, 4, 5	J_2
	3	30	2, 3, 4, 5	J_3, J_4
	4	85	1, 3, 5, 7	J_5, J_6
	5	86	2, 3, 5, 7	J_7, J_8, J_9, J_{10}
	6	89	1, 4, 5, 7	J_{11}
	7	113	1, 5, 6, 7	J_{12}, J_{13}, J_{14}
	8	201	1, 4, 7, 8	J_{15}
	9	209	1, 5, 7, 8	J_{16}
	10	59	1, 2, 4, 5, 6	J_{17}
	11	87	1, 2, 3, 5, 7	J_{18}, J_{19}
	12	91	1, 2, 4, 5, 7	J_{20}
	13	94	2, 3, 4, 5, 7	J_{21}, J_{22}, J_{23}
	14	115	1, 2, 5, 6, 7	J_{24}, J_{25}, J_{26}
	15	117	1, 3, 5, 6, 7	$J_{27}, J_{28}, J_{29}, J_{30}$
	16	121	1, 4, 5, 6, 7	J_{31}
	17	203	1, 2, 4, 7, 8	J_{32}
	18	206	2, 3, 4, 7, 8	J_{33}, J_{34}
	19	211	1, 2, 5, 7, 8	J_{35}
	20	213	1, 3, 5, 7, 8	J_{36}
	21	119	1, 2, 3, 5, 6, 7	J_{37}, J_{38}
	22	123	1, 2, 4, 5, 6, 7	J_{39}
	23	215	1, 2, 3, 5, 7, 8	J_{40}, J_{41}, J_{42}
	24	222	2, 3, 4, 5, 7, 8	J_{43}, J_{44}
	25	235	1, 2, 4, 6, 7, 8	J_{45}
	26	249	1, 4, 5, 6, 7, 8	J_{46}
	27	223	1, 2, 3, 4, 5, 7, 8	J_{47}, J_{48}, J_{49}
	28	251	1, 2, 4, 5, 6, 7, 8	J_{50}

Table 3: Overview of the set of master integrals. The first column denotes the topology, the second column labels consecutively the sectors, the third column gives the sector id (defined in eq. (8)), the fourth column lists the propagators with positive exponent, the fifth column lists the master integrals in the basis \vec{J} .

topology	block	sector	propagators	master integrals
<i>B</i>	29	113	1, 5, 6, 7	J_{51}
	30	116	3, 5, 6, 7	J_{52}
	31	115	1, 2, 5, 6, 7	J_{53}
	32	117	1, 3, 5, 6, 7	J_{54}, J_{55}, J_{56}
	33	206	2, 3, 4, 7, 8	J_{57}, J_{58}, J_{59}
	34	233	1, 4, 6, 7, 8	J_{60}
	35	236	3, 4, 6, 7, 8	J_{61}
	36	119	1, 2, 3, 5, 6, 7	J_{62}, J_{63}
	37	207	1, 2, 3, 4, 7, 8	J_{64}
	38	235	1, 2, 4, 6, 7, 8	J_{65}
	39	223	1, 2, 3, 4, 5, 7, 8	J_{66}
40	239	1, 2, 3, 4, 6, 7, 8	J_{67}	
<i>C</i>	41	101	1, 3, 6, 7	J_{68}, J_{69}, J_{70}
	42	102	2, 3, 6, 7	J_{71}
	43	103	1, 2, 3, 6, 7	J_{72}
	44	117	1, 3, 5, 6, 7	J_{73}, J_{74}, J_{75}
	45	118	2, 3, 5, 6, 7	J_{76}, J_{77}, J_{78}
	46	173	1, 3, 4, 6, 8	J_{79}, J_{80}
	47	181	1, 3, 5, 6, 8	J_{81}
	48	119	1, 2, 3, 5, 6, 7	J_{82}, J_{83}, J_{84}
	49	175	1, 2, 3, 4, 6, 8	J_{85}
	50	183	1, 2, 3, 5, 6, 8	J_{86}
	51	189	1, 3, 4, 5, 6, 8	J_{87}, J_{88}
	52	190	2, 3, 4, 5, 6, 8	J_{89}
	53	231	1, 2, 3, 6, 7, 8	J_{90}, J_{91}, J_{92}
	54	246	2, 3, 5, 6, 7, 8	J_{93}, J_{94}
	55	191	1, 2, 3, 4, 5, 6, 8	$J_{95}, J_{96}, J_{97}, J_{98}$
	56	247	1, 2, 3, 5, 6, 7, 8	$J_{99}, J_{100}, J_{101}, J_{102}$
	57	253	1, 3, 4, 5, 6, 7, 8	J_{103}
	58	255	1, 2, 3, 4, 5, 6, 7, 8	J_{104}
<i>D</i>	59	239	1, 2, 3, 4, 6, 7, 8	J_{105}

Table 4: Overview of the set of master integrals (continued).

$$\begin{aligned}
\omega_2 &= d \ln(1-w), \\
\omega_3 &= d \ln(-v), \\
\omega_4 &= d \ln(1-v), \\
\omega_5 &= d \ln(4-v), \\
\omega_6 &= d \ln(w-v), \\
\omega_7 &= d \ln(1-w-v), \\
\omega_8 &= d \ln(1-w+v), \\
\omega_9 &= d \ln(w^2+v(1-w)), \\
\omega_{10} &= d \ln\left((1-w)^2+vw\right), \\
\omega_{11} &= d \ln\left((1-w)^2+v(2-w)\right), \\
\omega_{12} &= d \ln(\lambda(v,w,1)), \\
\omega_{13} &= d \ln\left(\frac{2-v-r_1}{2-v+r_1}\right), \\
\omega_{14} &= d \ln\left(\frac{2w+v(1-w)-(1-w)r_1}{2w+v(1-w)+(1-w)r_1}\right), \\
\omega_{15} &= d \ln\left(\frac{2(1-w)+vw-wr_1}{2(1-w)+vw+wr_1}\right), \\
\omega_{16} &= d \ln\left(\frac{wv^2+(1-2w-w^2)v+2(1-w)^2-(1-w^2+vw)r_1}{wv^2+(1-2w-w^2)v+2(1-w)^2+(1-w^2+vw)r_1}\right), \\
\omega_{17} &= d \ln\left(\frac{1+w-v-r_2}{1+w-v+r_2}\right), \\
\omega_{18} &= d \ln\left(\frac{(1-w)^2-v(1+w)-(1-w)r_2}{(1-w)^2-v(1+w)+(1-w)r_2}\right), \\
\omega_{19} &= d \ln\left(\frac{-v(3-v+w)-r_1r_2}{-v(3-v+w)+r_1r_2}\right). \tag{32}
\end{aligned}$$

We write

$$\omega_k = d \ln f_k. \tag{33}$$

The variable transformation from (v, w) to (x, y) rationalises all square roots. The f_k 's are then rational functions in x and y . In detail we have

$$\begin{aligned}
f_1 &= -p_1^{-1} p_5^{-1} p_6^{-1} p_8 p_9, & f_2 &= 2p_1^{-1} p_2 p_4 p_5^{-1} p_6^{-1} p_7, & f_3 &= p_1^{-1} p_2^2, \\
f_4 &= p_1^{-1} p_{17}, & f_5 &= p_1^{-1} p_3^2, & f_6 &= -p_1^{-1} p_5^{-1} p_6^{-1} p_{18}, \\
f_7 &= p_1^{-1} p_2 p_5^{-1} p_6^{-1} p_{16}, & f_8 &= p_1^{-1} p_2 p_5^{-1} p_6^{-1} p_{12}, & f_9 &= p_1^{-2} p_5^{-2} p_6^{-2} p_{19} p_{20}, \\
f_{10} &= p_1^{-2} p_2^2 p_5^{-2} p_6^{-2} p_{14} p_{15}, & f_{11} &= p_1^{-2} p_2^2 p_5^{-2} p_6^{-2} p_{10} p_{13}, & f_{12} &= p_1^{-2} p_2^2 p_5^{-2} p_6^{-2} p_{11}^2,
\end{aligned}$$

$$\begin{aligned}
f_{13} &= p_1^2, & f_{14} &= p_1^{-2} p_{19} p_{20}^{-1}, & f_{15} &= p_1^{-2} p_{14}^{-1} p_{15}, \\
f_{16} &= p_1^{-2} p_{10}^{-1} p_{13} p_{14}^{-1} p_{15}, & f_{17} &= -p_1 p_5^{-1} p_6 p_8 p_9^{-1}, & f_{18} &= -p_1^{-1} p_5^3 p_6^{-3} p_8 p_9^{-1}, \\
f_{19} &= -4 p_1 p_4^2 p_7^{-2}. & & & &
\end{aligned} \tag{34}$$

The p_j 's are polynomials in x and y and given by

$$\begin{aligned}
p_1 &= x, \\
p_2 &= x - 1, \\
p_3 &= x + 1, \\
p_4 &= y, \\
p_5 &= y - 1, \\
p_6 &= y + 1, \\
p_7 &= xy + x - y + 1, \\
p_8 &= xy + x - 2y, \\
p_9 &= 2xy - y + 1, \\
p_{10} &= xy^2 + 2xy - 2y^2 + x + 2y, \\
p_{11} &= xy^2 + 2xy + y^2 + x - 2y + 1, \\
p_{12} &= xy^2 + 2xy - y^2 + x + 2y - 1, \\
p_{13} &= 2xy^2 + 2xy - y^2 + 2y - 1, \\
p_{14} &= 2xy^2 + 2xy - 3y^2 + 2y + 1, \\
p_{15} &= 3xy^2 + 2xy - 2y^2 - x + 2y, \\
p_{16} &= 3xy^2 + 2xy - 3y^2 - x + 2y + 1, \\
p_{17} &= x^2 - x + 1, \\
p_{18} &= x^2 y^2 + 2x^2 y - 3xy^2 + x^2 + y^2 - x - 2y + 1, \\
p_{19} &= x^2 y^2 + 2x^2 y - 4xy^2 + x^2 + 2y^2 - 2y, \\
p_{20} &= 2x^2 y^2 + 2x^2 y - 4xy^2 + y^2 - 2y + 1.
\end{aligned} \tag{35}$$

The polynomials p_1 - p_{16} are linear in x and appeared previously in the two-loop mixed QCD-electroweak corrections to $H \rightarrow b\bar{b}$ through a $Ht\bar{t}$ -coupling [42]. The polynomials p_{17} - p_{20} are quadratic in x .

5 Boundary values

In order to solve the differential equation we need boundary values. As boundary point we take the point $(v, w) = (0, 1)$, which corresponds in (x, y) -space to the line $x = 1$. The map from (v, w) -space to (x, y) -space blows up the point $(v, w) = (0, 1)$.

Most master integrals vanish at the boundary point. A few boundary values we have to

compute explicitly. They are obtained from rather simple integrals. We have

$$J_1 = C_1 w^{-\varepsilon}, \quad J_{15} = C_{15} (-v)^{-\varepsilon}, \quad J_{51} = C_{51} (-v)^{-2\varepsilon} w^{-\varepsilon}, \quad J_{60} = C_{60} (-v)^{-2\varepsilon}, \quad J_{71} = C_{71}, \quad (36)$$

with boundary values

$$\begin{aligned} C_1 &= e^{3\gamma_E \varepsilon} (\Gamma(1+\varepsilon))^3, \\ C_{15} &= 2e^{3\gamma_E \varepsilon} \frac{(\Gamma(1+\varepsilon))^3 (\Gamma(1-\varepsilon))^2}{\Gamma(1-2\varepsilon)}, \\ C_{51} &= -3e^{3\gamma_E \varepsilon} \frac{\Gamma(1+\varepsilon) (\Gamma(1-\varepsilon))^3 \Gamma(1+2\varepsilon)}{\Gamma(1-3\varepsilon)}, \\ C_{60} &= 4e^{3\gamma_E \varepsilon} \frac{(\Gamma(1+\varepsilon))^3 (\Gamma(1-\varepsilon))^4}{\Gamma(1-2\varepsilon)^2}, \\ C_{71} &= e^{3\gamma_E \varepsilon} \frac{\Gamma(1-\varepsilon) \Gamma(1+\varepsilon) \Gamma(1+2\varepsilon)^2 \Gamma(1+3\varepsilon)}{\Gamma(1+4\varepsilon)}. \end{aligned} \quad (37)$$

Note that J_{15} , J_{51} and J_{60} have logarithmic singularities at $v = 0$ and C_{15} , C_{51} , C_{60} denote ‘‘regularised’’ boundary values: They are given by the finite values obtained after removing all powers of $\ln(-v)$. The remaining non-zero boundary values are

$$\begin{aligned} C_{14} &= C_1, \\ C_{16} &= C_{59} = C_{15}, \\ C_{52} &= C_{51}, \\ C_{61} &= C_{60}, \\ C_6 &= C_8 = C_9 = C_{29} = C_{70} = C_{77} = \frac{1}{3} C_{71}, \\ C_{56} &= -\frac{2}{3} C_{71}, \\ C_{30} &= -\frac{1}{2} C_1 + \frac{1}{2} C_{71}, \\ C_{78} &= \frac{1}{2} C_1 - \frac{1}{2} C_{71}, \\ C_{87} &= -\frac{1}{4} C_1 + \frac{1}{4} C_{71}. \end{aligned} \quad (38)$$

All other boundary values are zero.

All non-trivial boundary values follow from (i) the explicit computations of eq. 37, (ii) a dedicated integral reduction for $s = 0$ and $m_W^2 = m_t^2$ and (iii) the requirement that the master integrals J_1 - J_{105} are constant on the hypersurface $x = 1$. Note that this does not require that A vanishes when restricted to $x = 1$, it only requires that the vector J is in the kernel of A on the hypersurface $x = 1$.

6 Analytical results

For the multiple polylogarithms [57–59] we use the notation $G(z_1, \dots, z_k; y)$, which is associated with the iterated integral representation. One defines $G(0, \dots, 0; y)$ with k zeros to be

$$G(0, \dots, 0; y) = \frac{1}{k!} (\ln y)^k. \quad (39)$$

This includes the trivial case $G(; y) = 1$. Multiple polylogarithms are then defined recursively by

$$G(z_1, z_2, \dots, z_k; y) = \int_0^y \frac{dy_1}{y_1 - z_1} G(z_2, \dots, z_k; y_1). \quad (40)$$

A discussion of the basic properties of multiple polylogarithms can be found in ref. [55].

We take the point $(v, w) = (0, 1)$ in (v, w) -space as our boundary point. In (x, y) -space this point corresponds to the hyperplane $x = 1$. The master integrals are constant on this hyperplane. We therefore have to integrate only along x in (x, y) -space. It is advantageous to change variables from x to $x' = 1 - x$. Doing so, the starting point of the integration will be $x' = 0$. The integration gives multiple polylogarithms of the form

$$G(l'_1, \dots, l'_k; x'), \quad (41)$$

where the letters l'_1, \dots, l'_k are from the alphabet

$$\mathcal{A} = \{0, 1, 2, x'_7, x'_8, x'_9, x'_{10}, x'_{11}, x'_{12}, x'_{13}, x'_{14}, x'_{15}, x'_{16}, x'_{17,a}, x'_{17,b}, x'_{18,a}, x'_{18,b}, x'_{19,a}, x'_{19,b}, x'_{20,a}, x'_{20,b}\}. \quad (42)$$

This is an alphabet with 21 letters. The non-trivial letters are given by

$$\begin{aligned} x'_7 &= \frac{2}{1+y}, & x'_8 &= \frac{1-y}{1+y}, & x'_9 &= \frac{1+y}{2y}, \\ x'_{10} &= \frac{1+4y-y^2}{(1+y)^2}, & x'_{11} &= \frac{2(1+y^2)}{(1+y)^2}, & x'_{12} &= \frac{4y}{(1+y)^2}, \\ x'_{13} &= -\frac{1-4y-y^2}{2y(1+y)}, & x'_{14} &= \frac{1+4y-y^2}{2y(1+y)}, & x'_{15} &= \frac{1-4y-y^2}{(1+y)(1-3y)}, \\ x'_{16} &= -\frac{4y}{(1+y)(1-3y)} \end{aligned} \quad (43)$$

and

$$\begin{aligned} x'_{17,a} &= e^{\frac{\pi i}{3}}, & x'_{17,b} &= e^{-\frac{\pi i}{3}}, \\ x'_{18,a} &= \frac{1+4y-y^2+i\sqrt{(3+y^2)(1-5y^2)}}{2(1+y)^2}, & x'_{18,b} &= \frac{1+4y-y^2-i\sqrt{(3+y^2)(1-5y^2)}}{2(1+y)^2}, \end{aligned}$$

$$\begin{aligned}
x'_{19,a} &= \frac{1 + 2y - y^2 + \sqrt{2y(1+y-y^2+y^3)}}{(1+y)^2}, & x'_{19,b} &= \frac{1 + 2y - y^2 - \sqrt{2y(1+y-y^2+y^3)}}{(1+y)^2}, \\
x'_{20,a} &= \frac{2y + i\sqrt{2y(1-y-y^2-y^3)}}{2y(1+y)}, & x'_{20,b} &= \frac{2y - i\sqrt{2y(1-y-y^2-y^3)}}{2y(1+y)}.
\end{aligned} \tag{44}$$

The letters are the roots of the polynomials p_1 - p_{20} in x' with x substituted by $1 - x'$.

For all master integrals we write

$$J_k = \sum_{j=0}^{\infty} \varepsilon^j J_k^{(j)}. \tag{45}$$

The ε^j -term $J_k^{(j)}$ is of uniform weight j . To give an example, we have

$$\begin{aligned}
J_{67} &= 2 \left[G(0, x'_8, x'_9; x') + G(0, x'_9, x'_8; x') - G(x'_{15}, x'_8, 0; x') - G(x'_{14}, x'_9, 0; x') \right. \\
&\quad - G(x'_{14}, x'_8, 0; x') - G(x'_{15}, x'_9, 0; x') + G(x'_{14}, x'_8, x'_9; x') + G(x'_{14}, x'_9, x'_8; x') \\
&\quad + G(x'_{15}, x'_8, x'_9; x') + G(x'_{15}, x'_9, x'_8; x') - G(1, x'_9, x'_8; x') + G(x'_{15}, x'_9, 1; x') \\
&\quad - G(x'_7, x'_9, 1; x') + G(x'_7, 1, x'_9; x') - G(0, x'_8, 1; x') - G(1, x'_9, 1; x') \\
&\quad - G(0, 1, x'_8; x') - 2G(x'_7, 1, 0; x') + G(1, 0, 1; x') - G(x'_{14}, x'_8, 1; x') \\
&\quad + 2G(x'_{15}, 1, 0; x') + 2G(x'_7, x'_8, 0; x') + 2G(x'_7, x'_9, 0; x') - G(x'_{15}, x'_7, 1; x') \\
&\quad - G(x'_7, x'_9, x'_8; x') - G(x'_{15}, 1, x'_9; x') - G(x'_7, x'_8, x'_9; x') - G(x'_{14}, x'_7, x'_9; x') \\
&\quad - G(x'_{15}, x'_7, x'_8; x') - G(1, x'_8, x'_9; x') + 2G(x'_{14}, x'_7, 1; x') + G(1, x'_7, 1; x') \\
&\quad \left. + G(1, x'_7, x'_8; x') \right] \varepsilon^3 + O(\varepsilon^4). \tag{46}
\end{aligned}$$

The analytical results for the master integrals are given up to the order ε^5 in a supplementary electronic file attached to the arxiv version of this article.

7 Numerical results

As a reference value we give numerical results for

$$p^2 = m_H^2. \tag{47}$$

Since $p^2 > 0$, we are not in the Euclidean region. Feynman's $i0$ -prescription instructs us to take a small imaginary part into account: $p^2 \rightarrow p^2 + i0$. This selects the correct branches for the two square roots $\sqrt{-v(4-v)}$ and $\sqrt{\lambda(v, w, 1)}$. With

$$m_W = 80.38 \text{ GeV}, \quad m_H = 125.2 \text{ GeV}, \quad m_t = 173.1 \text{ GeV} \tag{48}$$

we obtain for the variables x and y

$$x = 0.7384 + 0.6743i, \quad y = 0.3987i. \tag{49}$$

The values of the master integrals at this point are given to 8 digits in tables 5 and 6. They are easily computed to arbitrary precision by evaluating the multiple polylogarithms with the help of GiNaC [60, 61]. In addition we verified the correctness of our results at several kinematic points with the help of the programs AMFlow [62–64] and sector_decomposition [65, 66].

	ϵ^0	ϵ^1	ϵ^2	ϵ^3	ϵ^4	ϵ^5
J_1	1	1.5342081	3.6442984	3.1853184	5.1462966	2.4686377
J_2	0	-1.4801083 <i>i</i>	-2.4097495 <i>i</i>	-5.6149488 <i>i</i>	-5.2333001 <i>i</i>	-8.0886313 <i>i</i>
J_3	0	1.4801083 <i>i</i>	-0.22727279 <i>i</i>	7.1496293 <i>i</i>	-6.383137 <i>i</i>	25.607456 <i>i</i>
J_4	0	0	-0.54768013	-0.36476197	-2.891637	0.046729212
J_5	0	1.5342081	-2.9983034	19.075365	-53.393746	195.47467
J_6	0.33333333	1.5342081	-0.088235985	15.450021	-32.95737	146.32479
J_7	0	-1.4585842 <i>i</i>	1.294182 <i>i</i>	-18.724204 <i>i</i>	33.200239 <i>i</i>	-173.64768 <i>i</i>
J_8	0.33333333	0	3.7496407	-3.6797304	30.246157	-60.506131
J_9	0.33333333	1.5342081	1.3128219	16.578415	-17.750719	142.4298
J_{10}	0	0	-2.2406312	1.0647958	-26.559377	40.135978
J_{11}	0	-1.5342081	0.91070302	-7.2586694	10.224944	-28.168258
J_{12}	0	1.4585842 <i>i</i>	0.050868134 <i>i</i>	7.6533633 <i>i</i>	-4.6991476 <i>i</i>	27.583113 <i>i</i>
J_{13}	0	1.5342081	-1.7502763	7.0599975	-14.845047	30.115459
J_{14}	1	1.5342081	5.8849296	4.6861424	17.388435	2.5361954
J_{15}	2	1.295828 + 6.2831853 <i>i</i>	-7.8048778 + 4.0709638 <i>i</i>	-12.450557 - 3.8488956 <i>i</i>	-11.660945 - 25.721646 <i>i</i>	10.668736 - 35.695423 <i>i</i>
J_{16}	2	4.3642443 + 6.2831853 <i>i</i>	-3.4630132 + 13.710678 <i>i</i>	-21.696071 + 9.7914742 <i>i</i>	-38.706617 - 23.053895 <i>i</i>	-26.131312 - 75.786891 <i>i</i>
J_{17}	0	0	2.1907205	3.772359	8.6571337	8.5453519
J_{18}	0	-1.4801083 <i>i</i>	0.5445262 <i>i</i>	-15.995422 <i>i</i>	23.86858 <i>i</i>	-132.81753 <i>i</i>
J_{19}	0	0	-5.2250698 <i>i</i>	10.295242 <i>i</i>	-65.298537 <i>i</i>	183.26131 <i>i</i>
J_{20}	0	0	2.2707942 <i>i</i>	-1.1347528 <i>i</i>	10.629046 <i>i</i>	-14.132023 <i>i</i>
J_{21}	0	1.4801083 <i>i</i>	-5.4523426 <i>i</i>	28.891559 <i>i</i>	-88.331183 <i>i</i>	331.78433 <i>i</i>
J_{22}	0	-1.5342081	3.2901965	-18.78899	54.646442	-195.09957
J_{23}	0	1.5342081	-2.1449256	21.075678	-46.255399	207.64266
J_{24}	0	0	2.1588625	0.27796832	11.346261	-5.8909184
J_{25}	0	0	-2.2707942 <i>i</i>	2.3774122 <i>i</i>	-10.218327 <i>i</i>	21.004436 <i>i</i>
J_{26}	0	-1.4801083 <i>i</i>	-2.4097495 <i>i</i>	-8.9313256 <i>i</i>	-7.7660295 <i>i</i>	-26.43436 <i>i</i>
J_{27}	0	-1.4585842 <i>i</i>	1.5911765 <i>i</i>	-16.821522 <i>i</i>	38.929474 <i>i</i>	-158.55603 <i>i</i>
J_{28}	0	-1.4801083 <i>i</i>	5.4523426 <i>i</i>	-27.832513 <i>i</i>	90.872416 <i>i</i>	-322.88758 <i>i</i>
J_{29}	0.33333333	0	4.2973208	-7.8296127	39.354989	-110.36825
J_{30}	0	0	0	3.9327313	-9.7902493	50.090814
J_{31}	0	-1.4801083 <i>i</i>	2.8153204 <i>i</i>	-9.1045495 <i>i</i>	18.752518 <i>i</i>	-45.336109 <i>i</i>
J_{32}	0	-2.9602166 <i>i</i>	9.2997946 - 2.1958763 <i>i</i>	6.898549 + 11.35639 <i>i</i>	-5.0820522 + 19.501961 <i>i</i>	-38.5719 + 19.050034 <i>i</i>
J_{33}	0	2.9602166 <i>i</i>	-9.2997946 - 3.0781682 <i>i</i>	9.6703505 - 3.6126738 <i>i</i>	-19.245548 - 12.824796 <i>i</i>	72.104463 + 0.12405401 <i>i</i>
J_{34}	0	0	-1.0953603	0.24128744 - 3.4411758 <i>i</i>	1.6402683 + 0.75802684 <i>i</i>	6.1054746 - 6.1679597 <i>i</i>
J_{35}	0	-2.9602166 <i>i</i>	9.2997946 - 6.7374647 <i>i</i>	21.16637 + 4.5035873 <i>i</i>	16.446661 + 32.559044 <i>i</i>	-32.652689 + 60.330416 <i>i</i>
J_{36}	0	-1.5342081	-2.6034827 - 9.6397141 <i>i</i>	26.9974 - 0.52358996 <i>i</i>	-27.747558 - 12.210078 <i>i</i>	223.69832 + 60.357017 <i>i</i>
J_{37}	0	0	0	0	2.4095606	-7.4050612
J_{38}	0	0	0	-0.80345343	2.3774716	-10.59662
J_{39}	0	0	2.1907205	-3.9613101	13.09607	-26.512146
J_{40}	0	2.9602166 <i>i</i>	-9.2997946 - 3.712675 <i>i</i>	14.988491 - 6.3598329 <i>i</i>	-8.0684303 - 1.80014 <i>i</i>	80.59546 + 18.841397 <i>i</i>
J_{41}	0	0	0	-1.2199963 - 1.2715067 <i>i</i>	5.3143264 - 0.86415471 <i>i</i>	-13.01594 - 2.7052663 <i>i</i>
J_{42}	0	0	0	0	4.2051087 + 3.3549988 <i>i</i>	-14.946099 + 3.7606676 <i>i</i>
J_{43}	0	0	0	0	2.6414875 + 2.2409565 <i>i</i>	-11.227012 + 0.75555211 <i>i</i>
J_{44}	0	0	0	-1.2199963 - 1.2715067 <i>i</i>	4.98217 - 0.70082991 <i>i</i>	-11.56359 - 2.4458449 <i>i</i>
J_{45}	0	0	4.381441	3.6614725 + 13.764703 <i>i</i>	-16.48045 + 11.502855 <i>i</i>	-30.424976 - 6.4908014 <i>i</i>
J_{46}	0	0	0	-1.2199963 - 1.2715067 <i>i</i>	3.4643984 - 1.4351478 <i>i</i>	-2.3605461 + 0.17411836 <i>i</i>
J_{47}	0	0	0	0	0	2.7105126 + 2.3055744 <i>i</i>
J_{48}	0	0	0	0	0.65371112 + 1.0931833 <i>i</i>	-1.0109578 + 3.0469453 <i>i</i>
J_{49}	0	0	0	0	1.6251574 + 0.75070903 <i>i</i>	-1.068945 + 3.6561297 <i>i</i>
J_{50}	0	0	0	0	0.94098377 - 0.90286329 <i>i</i>	1.1504284 + 2.4970979 <i>i</i>
J_{51}	-3	-3.887484 - 18.849556 <i>i</i>	59.166272 - 24.425783 <i>i</i>	118.51311 + 123.70244 <i>i</i>	-92.180991 + 423.20939 <i>i</i>	-842.91965 + 395.8266 <i>i</i>
J_{52}	-3	-8.4901085 - 18.849556 <i>i</i>	49.671371 - 53.344925 <i>i</i>	202.90571 + 64.044212 <i>i</i>	156.24311 + 572.90309 <i>i</i>	-810.36632 + 1171.6505 <i>i</i>
J_{53}	0	4.4403248 <i>i</i>	-27.899384 + 6.1707632 <i>i</i>	-38.772048 - 87.008884 <i>i</i>	179.55177 - 183.60241 <i>i</i>	643.38826 + 119.50921 <i>i</i>
J_{54}	0	-2.9171683 <i>i</i>	9.1645545 - 1.9918105 <i>i</i>	24.127791 + 5.3212259 <i>i</i>	-21.804448 + 120.95424 <i>i</i>	-196.92433 - 102.13413 <i>i</i>
J_{55}	0	-1.5342081	-8.2052687 - 19.279428 <i>i</i>	79.333963 - 36.244239 <i>i</i>	118.23648 + 121.5975 <i>i</i>	181.95398 + 573.21464 <i>i</i>
J_{56}	-0.66666667	-1.5342081	-7.3030938	-22.007405 - 14.078301 <i>i</i>	-7.7964625 - 73.29467 <i>i</i>	88.637395 - 153.09033 <i>i</i>
J_{57}	0	2.9171683 <i>i</i>	-9.1645545 + 1.9918105 <i>i</i>	-6.2574572 - 3.2092524 <i>i</i>	-20.068012 - 19.446051 <i>i</i>	40.505362 - 29.233598 <i>i</i>
J_{58}	0	3.0684163	-1.5124828 + 9.6397141 <i>i</i>	-7.6933829 - 4.7516047 <i>i</i>	-18.562293 + 7.543913 <i>i</i>	-4.8730848 - 73.947317 <i>i</i>
J_{59}	2	4.3642443 + 6.2831853 <i>i</i>	-1.2223821 + 13.710678 <i>i</i>	-18.743511 + 16.830625 <i>i</i>	-39.764537 - 13.778153 <i>i</i>	-42.529113 - 55.952566 <i>i</i>
J_{60}	4	5.1833121 + 25.132741 <i>i</i>	-78.888363 + 32.56771 <i>i</i>	-129.16811 - 164.93658 <i>i</i>	199.25544 - 383.01328 <i>i</i>	779.6171 - 48.063607 <i>i</i>
J_{61}	4	11.320145 + 25.132741 <i>i</i>	-66.228494 + 71.126566 <i>i</i>	-241.69158 - 85.392283 <i>i</i>	-87.715765 - 582.60488 <i>i</i>	887.29829 - 804.39581 <i>i</i>
J_{62}	0	0	0	-1.2199963 - 1.2715067 <i>i</i>	5.9914141 - 3.4779304 <i>i</i>	-0.18199463 + 1.5542242 <i>i</i>
J_{63}	0	2.9602166 <i>i</i>	-9.2997946 - 3.712675 <i>i</i>	-16.248795 - 0.64764184 <i>i</i>	40.454671 - 155.25155 <i>i</i>	168.63449 + 314.27487 <i>i</i>
J_{64}	0	-2.9602166 <i>i</i>	9.2997946 + 3.712675 <i>i</i>	-11.663712 + 4.2951778 <i>i</i>	17.101399 + 9.4430398 <i>i</i>	-68.03826 - 7.8262762 <i>i</i>
J_{65}	0	-5.9204331 <i>i</i>	37.199178 - 8.2276842 <i>i</i>	51.696065 + 116.01185 <i>i</i>	-239.40236 + 202.10303 <i>i</i>	-589.55784 - 276.35688 <i>i</i>
J_{66}	0	0	0	0	0	2.6632592 + 5.3571394 <i>i</i>
J_{67}	0	0	0	1.2199963 + 1.2715067 <i>i</i>	-6.6685018 + 6.0917061 <i>i</i>	-14.752046 - 15.744008 <i>i</i>

Table 5: Numerical results for the first six terms of the ϵ -expansion of the master integrals J_1 - J_{67} at the kinematic point $p^2 = m_H^2$.

	ϵ^0	ϵ^1	ϵ^2	ϵ^3	ϵ^4	ϵ^5
J_{68}	0	-1.4801083i	5.1350892i	-25.837256i	86.938604i	-308.88849i
J_{69}	0	0	1.0953603	-3.0231166	16.571142	-51.663494
J_{70}	0.33333333	0	1.3701472	1.8283464	-4.959024	38.837723
J_{71}	1	0	2.4674011	8.4143983	-35.10786	169.45714
J_{72}	0	-1.4801083i	2.498067i	-15.01875i	37.560716i	-132.37051i
J_{73}	0	-1.4585842i	1.5911765i	-15.779761i	37.376156i	-152.62396i
J_{74}	0	1.5342081	-2.1449256	18.664175	-47.494038	189.76691
J_{75}	0	-1.5342081	-0.2345679	-8.9971424	3.8407769	-34.623785
J_{76}	0	-1.4585842i	1.294182i	-16.104568i	36.151551i	-153.04501i
J_{77}	0.33333333	1.5342081	0.62627943	18.031643	-37.794484	187.99495
J_{78}	0	0	0	1.5692999	-6.5217998	25.959061
J_{79}	0	2.9602166i	-9.2997946 - 3.0781682i	-14.67018 - 1.6806018i	50.404394 - 150.6488i	194.75027 + 331.14331i
J_{80}	0	0	-0.54768013	-0.41446154 - 1.7205879i	-2.3612249 - 7.5128487i	29.563404 - 11.514285i
J_{81}	0	1.5342081	-2.9983034	17.247662	-52.439144	184.22811
J_{82}	0	1.4801083i	-5.4523426i	23.141893i	-87.484869i	292.60701i
J_{83}	0	0	0	-0.80345343	2.4731997	-10.583527
J_{84}	0	0	0	0	1.6505684	-6.5723197
J_{85}	0	0	0	-0.9800827 - 0.86029394i	4.9546534 - 1.9953368i	-4.239423 + 1.9524939i
J_{86}	0	0	-0.31725341i	-1.4970783i	1.498074i	-9.4006924i
J_{87}	0	0	-0.13692003	2.3004984 - 0.43014697i	-6.0933617 - 0.63742354i	39.061373 + 3.0381966i
J_{88}	0	0	0	-1.2199963 - 1.2715067i	5.1907304 - 3.6242119i	2.1054694 + 0.72653192i
J_{89}	0	0	0	-1.3411296 - 1.3187987i	5.076922 - 3.9113363i	2.0464347 - 0.00022367332i
J_{90}	0	0	0	-0.9800827 - 0.86029394i	4.8178105 + 0.24314247i	-14.201072 - 1.4422657i
J_{91}	0	0	0	0	2.4985099 + 1.7074518i	-12.225556 - 0.48055535i
J_{92}	0	1.4801083i	-9.2997946 + 2.0569211i	2.2589826 - 23.245834i	-5.6384268 + 23.974917i	56.355336 - 142.57165i
J_{93}	0	0	0	0	2.7500878 + 2.2730554i	-11.719892 + 0.89738075i
J_{94}	0	0	0	-1.3411296 - 1.3187987i	4.9222387 - 0.91431241i	-11.726836 - 2.8899212i
J_{95}	0	0	0	0	0	1.8404286 + 0.14449436i
J_{96}	0	0	0	0	0.21789937 + 0.042131655i	0.50386273 + 0.33655445i
J_{97}	0	0	0	-0.23991356 - 0.41121273i	0.85908359 - 1.5044267i	3.8455066 - 0.56878995i
J_{98}	0	0	0	0	0.78680569 - 5.3224605i	4.5796027 + 13.058166i
J_{99}	0	0	0	0	0	2.8101582 + 2.3220016i
J_{100}	0	0	0	0	0.39649658 + 0.58554103i	-0.70924343 + 1.3293236i
J_{101}	0	0	0	-0.36104693 - 0.45850473i	0.35197787 - 1.3297874i	0.96759852 - 1.7438646i
J_{102}	0	0	0	0	0.96352362 - 0.98327656i	2.3321043 + 1.5761209i
J_{103}	0	0	0	0	0	1.9661517 + 5.3108372i
J_{104}	0	0	0	0	0	0.022784811 + 0.43623285i
J_{105}	0	0	0	0	0.10737705 + 0.15759037i	-0.197588 + 0.3448854i

Table 6: Numerical results for the first six terms of the ϵ -expansion of the master integrals J_{68} - J_{105} at the kinematic point $p^2 = m_H^2$.

8 Conclusions

In this paper we considered all master integrals with internal top- and W -propagators contributing to the three-loop Higgs self-energy diagrams of order $O(\alpha^2\alpha_s)$. We obtained analytic results for all master integrals, keeping the full dependence on p^2 , m_W^2 and m_t^2 . We presented a basis of master integrals of uniform weight. In this basis the associated differential equation is ϵ -factorised. We set up three independent systems of differential equations, such that for each system all occurring square roots can be rationalised simultaneously. This allows us to express all master integrals to any order in the dimensional regularisation parameter ϵ in terms of multiple polylogarithms. Our results are a building block for precision Higgs physics.

Acknowledgements

We thank Johann Usovitsch for support with the computer program `Kira`. E.C. receives funding from the European Union’s Horizon 2020 research and innovation programme “High precision multi-jet dynamics at the LHC” (consolidator grant agreement No. 772099). S.W. would like to thank the Munich Institute for Astro-, Particle and BioPhysics (MIAPbP) for hospitality, where this work was finalised.

A The additional topologies

In this appendix we consider the additional topologies containing a top propagator and a W -propagator and not being proportional to the product of Yukawa couplings $y_b y_t$. There are three top sector diagrams which we have to consider. These are shown in fig. 3. The diagrams shown in fig. 3 have all eight propagators and define the top sectors. Simpler diagrams, obtained from the ones shown in fig. 3 by pinching a subset of the propagators, are not shown. The three top sectors can be obtained from one auxiliary graph with nine propagators, shown in fig. 4. The auxiliary graph B' is obtained from the auxiliary graph B through the exchange $m_W \leftrightarrow m_t$. The three diagrams in fig. 3 correspond to the sectors 255, 479 and 509 of the auxiliary graph B' . The auxiliary graph B' introduces additional square roots

$$r_3 = \sqrt{-v(4w-v)}, \quad r_4 = \sqrt{-v(-4-v)}. \quad (50)$$

For the sectors B'_{255} and B'_{479} we encounter the set of square roots (r_1, r_2, r_3) , while for the sector B'_{509} we encounter the set of square roots (r_1, r_2, r_4) . We do not know how to rationalise all four square roots simultaneously. But this is not needed: It is sufficient being able to rationalise the set of square roots (r_1, r_2, r_3) and independently the set of square roots (r_1, r_2, r_4) . This can be done. We are therefore led to two systems of differential equations, one for the sectors B'_{255} and B'_{479} with 61 master integrals K_1 - K_{61} and another system for the sector B'_{509} with 43 master integrals L_1 - L_{43} . In both cases we are able to transform the differential equation into an ϵ -factorised form

$$dK = BK, \quad B = \epsilon \sum_{k=1}^{28} M'_k \omega_k,$$




Figure 3: Examples of additional three-loop Higgs self-energy diagrams containing a top propagator and a W -propagator and not being proportional to the product of Yukawa couplings $y_b y_t$. The internal masses of the propagators are encoded by the colour of the propagators: massless (black), m_t (green), m_W (red).




Figure 4: The topology B' . This topology is related by the exchange $m_W \leftrightarrow m_t$ to topology B .

$$dL = CL, \quad C = \varepsilon \sum_{k=1}^{28} M_k'' \omega_k. \quad (51)$$

The entries of the 61×61 matrices M_k' and of the 43×43 matrices M_k'' are rational numbers. The differential one-forms ω_1 - ω_{19} are given in eq. (32), the additional differential one-forms are given by

$$\begin{aligned} \omega_{20} &= d \ln(4w - v), \\ \omega_{21} &= d \ln\left((1-w)^2 + v\right), \\ \omega_{22} &= d \ln\left(\frac{2w - v - r_3}{2w - v + r_3}\right), \\ \omega_{23} &= d \ln\left(\frac{2w(1-w) - v - r_3}{2w(1-w) - v + r_3}\right), \\ \omega_{24} &= d \ln\left(\frac{-v(1-v+3w) - r_2 r_3}{-v(1-v+3w) + r_2 r_3}\right), \\ \omega_{25} &= d \ln(4+v), \\ \omega_{26} &= d \ln\left((1-w)^2 - vw\right), \\ \omega_{27} &= d \ln\left(\frac{2+v-r_4}{2+v+r_4}\right), \\ \omega_{28} &= d \ln\left(\frac{-v(1+w) - (1-w)r_4}{-v(1+w) + (1-w)r_4}\right). \end{aligned} \quad (52)$$

Not all differential one-forms do occur in the two differential equation in eq. (51). The occurring ones are

$$\begin{aligned} K: & \quad \{\omega_1, \omega_2, \omega_3, \omega_5, \omega_6, \omega_8, \omega_{12}, \omega_{13}, \omega_{17}, \omega_{18}, \omega_{19}, \omega_{20}, \omega_{21}, \omega_{22}, \omega_{23}, \omega_{24}\}, \\ L: & \quad \{\omega_1, \omega_2, \omega_3, \omega_5, \omega_6, \omega_8, \omega_{12}, \omega_{13}, \omega_{17}, \omega_{18}, \omega_{19}, \omega_{25}, \omega_{26}, \omega_{27}, \omega_{28}\}. \end{aligned} \quad (53)$$

We remark that the three systems of master integrals $(A_{255}, B_{255}, C_{255}, D_{255})$, (B'_{255}, B'_{479}) and (B'_{509}) are redundant: The master integrals of B'_{255} are related through the exchange $m_W \leftrightarrow m_t$ to the ones of B_{255} . The master integrals of B'_{255} are K_1 - K_{32} , hence only K_{33} - K_{61} are “new”. The same situation appears in the system B'_{509} : The master integrals L_1 - L_{39} appear already in the system (B'_{255}, B'_{479}) , hence only L_{40} - L_{43} are “new”. This redundancy is the price to pay for having individual systems which can be solved independently with a single rationalisation.

A.1 The sectors B'_{255} and B'_{479}

The sector B'_{479} involves the square roots (r_1, r_2, r_3) . (The sector B'_{255} alone involves only the square roots (r_2, r_3) .) The square roots (r_1, r_2, r_3) are rationalised by the variables (\hat{x}, y) . The variable \hat{x} is defined by

$$x = \frac{(1-y) [(1+y)(1-4\hat{x}) + (3+11y)\hat{x}^2]}{(1+y) [1-y - (1+7y)\hat{x}^2]},$$

$$\hat{x} = \frac{(1-y^2)}{(1-x)} \frac{[2(1-x) - xr_3]}{[3+x+8y-11y^2+8xy+7xy^2]}. \quad (54)$$

This rationalisation is easily obtained with the help of the computer program `RationalizeRoots` [43, 67]. The value $x = 1$ corresponds to $\hat{x} = 0$. The root r_3 is given in the variables (\hat{x}, y) by

$$r_3 = \frac{8\hat{x}[1-y^2-(1+4y-y^2)\hat{x}][1-y^2-2(1+4y-y^2)\hat{x}+(1+y)(1+7y)\hat{x}^2]}{(1-y^2)[1-y-(1+7y)\hat{x}^2][(1+y)(1-4\hat{x})+(3+11y)\hat{x}^2]}. \quad (55)$$

With this rationalisation the differential one-forms translate to

$$d \ln(\hat{p}_1), \dots, d \ln(\hat{p}_{20}), \quad (56)$$

where $\hat{p}_1-\hat{p}_{20}$ are polynomials in (\hat{x}, y) . We note that the differential one-form $\omega_6 = d \ln(w - v)$ leads to a quartic polynomial in \hat{x} :

$$\begin{aligned} \hat{p}_{20} = & 1 - 4\hat{x} - 2y^2 - 16y\hat{x} + 18\hat{x}^2 + 8y^2\hat{x} + 48\hat{x}^2y - 28\hat{x}^3 + y^4 + 16y^3\hat{x} + 44\hat{x}^2y^2 - 144\hat{x}^3y \\ & + 13\hat{x}^4 - 4y^4\hat{x} - 48\hat{x}^2y^3 - 104y^2\hat{x}^3 + 112\hat{x}^4y + 2y^4\hat{x}^2 + 16y^3\hat{x}^3 + 246\hat{x}^4y^2 + 4y^4\hat{x}^3 \\ & + 16y^3\hat{x}^4 + 61y^4\hat{x}^4, \end{aligned} \quad (57)$$

the remaining 19 polynomials $\hat{p}_1-\hat{p}_{19}$ are at most quadratic in \hat{x} . The polynomial \hat{p}_{20} will lead to roots of a quartic polynomial as arguments of the multiple polylogarithms if the differential equation is integrated along \hat{x} for generic and constant y .

The master integrals for the sectors B'_{255} and B'_{479} are

$$\begin{aligned} K_1 &= \varepsilon^3 (1-w) \mathbf{D}^- I_{101010100}^{B'}, \\ K_2 &= \varepsilon^3 \mathbf{D}^- I_{101(-1)10100}^{B'}, \\ K_3 &= \varepsilon^3 r_2 \mathbf{D}^- I_{011010100}^{B'}, \\ K_4 &= \varepsilon^3 \left[\mathbf{D}^- I_{(-1)11010100}^{B'} + (1-w) \mathbf{D}^- I_{011010100}^{B'} \right], \\ K_5 &= \varepsilon^3 \left[\mathbf{D}^- I_{01101(-1)100}^{B'} + v \mathbf{D}^- I_{011010100}^{B'} \right], \\ K_6 &= \varepsilon^3 \left[\mathbf{D}^- I_{0110101(-1)0}^{B'} + v \mathbf{D}^- I_{011010100}^{B'} \right], \\ K_7 &= \varepsilon^3 v \mathbf{D}^- I_{100011100}^{B'}, \\ K_8 &= \varepsilon^3 v \mathbf{D}^- I_{001011100}^{B'}, \\ K_9 &= \varepsilon^3 v \mathbf{D}^- I_{101000110}^{B'}, \\ K_{10} &= \varepsilon^3 r_3 \left[\mathbf{D}^- I_{011010100}^{B'} + \mathbf{D}^- I_{111(-1)10100}^{B'} + (1-w) \mathbf{D}^- I_{111010100}^{B'} \right], \\ K_{11} &= \varepsilon^3 r_3 \mathbf{D}^- I_{111(-1)10100}^{B'}, \\ K_{12} &= \varepsilon^3 v r_3 \mathbf{D}^- I_{110011100}^{B'}, \\ K_{13} &= \varepsilon^3 r_2 \mathbf{D}^- I_{101(-1)11100}^{B'}, \end{aligned}$$

$$\begin{aligned}
K_{14} &= \varepsilon^3 (1-w) \left[\mathbf{D}^- I_{101(-1)11100}^{B'} + v \mathbf{D}^- I_{101011100}^{B'} \right], \\
K_{15} &= \varepsilon^3 \left[\mathbf{D}^- I_{101(-2)11100}^{B'} + v \mathbf{D}^- I_{101(-1)11100}^{B'} \right], \\
K_{16} &= \varepsilon^3 (1-w) \left[v \mathbf{D}^- I_{011011100}^{B'} + \mathbf{D}^- I_{011010100}^{B'} \right], \\
K_{17} &= \varepsilon^3 v r_3 \mathbf{D}^- I_{111000110}^{B'}, \\
K_{18} &= \varepsilon^3 v (1-w) \mathbf{D}^- I_{101100110}^{B'}, \\
K_{19} &= \varepsilon^3 v r_2 \mathbf{D}^- I_{011100110}^{B'}, \\
K_{20} &= \varepsilon^3 v \left[\mathbf{D}^- I_{(-1)11100110}^{B'} + (1-w) \mathbf{D}^- I_{011100110}^{B'} \right], \\
K_{21} &= \varepsilon^3 v \mathbf{D}^- I_{0111(-1)0110}^{B'}, \\
K_{22} &= \varepsilon^3 v^2 \mathbf{D}^- I_{100101110}^{B'}, \\
K_{23} &= \varepsilon^3 v^2 \mathbf{D}^- I_{001101110}^{B'}, \\
K_{24} &= \varepsilon^4 (1-2\varepsilon) v I_{111021100}^{B'}, \\
K_{25} &= \varepsilon^3 r_3 \left[(1-w) \mathbf{D}^- I_{111(-1)11100}^{B'} + \mathbf{D}^- I_{101(-1)11100}^{B'} \right], \\
K_{26} &= \varepsilon^3 v r_3 \left[(1-w) \mathbf{D}^- I_{111100110}^{B'} + \mathbf{D}^- I_{011100110}^{B'} \right], \\
K_{27} &= \varepsilon^3 v r_3 \mathbf{D}^- I_{11101011(-1)}^{B'}, \\
K_{28} &= \varepsilon^4 (1-2\varepsilon) v I_{112010110}^{B'}, \\
K_{29} &= \varepsilon^4 (1-2\varepsilon) v I_{111020110}^{B'}, \\
K_{30} &= \varepsilon^3 v^2 r_3 \mathbf{D}^- I_{110101110}^{B'}, \\
K_{31} &= \varepsilon^5 (1-2\varepsilon) v I_{111110110}^{B'}, \\
K_{32} &= \varepsilon^4 (1-2\varepsilon)^2 v I_{111101110}^{B'}, \\
K_{33} &= \varepsilon^3 \mathbf{D}^- I_{101000001}^{B'}, \\
K_{34} &= \varepsilon^3 r_3 \mathbf{D}^- I_{111000001}^{B'}, \\
K_{35} &= \varepsilon^3 (1-w) \mathbf{D}^- I_{101100001}^{B'}, \\
K_{36} &= \varepsilon^3 r_2 \mathbf{D}^- I_{011100001}^{B'}, \\
K_{37} &= \varepsilon^3 \left[\mathbf{D}^- I_{(-1)11100001}^{B'} + (1-w) \mathbf{D}^- I_{011100001}^{B'} \right], \\
K_{38} &= \varepsilon^3 \left[\mathbf{D}^- I_{01110(-1)001}^{B'} + v \mathbf{D}^- I_{011100001}^{B'} \right], \\
K_{39} &= \varepsilon^2 (1+4\varepsilon) \mathbf{D}^- I_{001100101}^{B'}, \\
K_{40} &= \varepsilon^3 r_1 \mathbf{D}^- I_{001100011}^{B'}, \\
K_{41} &= \varepsilon^3 \left[\mathbf{D}^- I_{(-1)01100011}^{B'} + (1-w) \mathbf{D}^- I_{001100011}^{B'} \right], \\
K_{42} &= \varepsilon^3 \left[\mathbf{D}^- I_{0011(-1)0011}^{B'} + v \mathbf{D}^- I_{001100011}^{B'} \right],
\end{aligned}$$

$$\begin{aligned}
K_{43} &= \varepsilon^3 v \mathbf{D}^- I_{001000111}^{B'}, \\
K_{44} &= \varepsilon^3 r_3 \left[(1-w) \mathbf{D}^- I_{111100001}^{B'} + \mathbf{D}^- I_{011100001}^{B'} \right], \\
K_{45} &= \varepsilon^3 (1-w) \left[(1-w) \mathbf{D}^- I_{101100101}^{B'} + \mathbf{D}^- I_{001100101}^{B'} \right], \\
K_{46} &= \varepsilon^3 r_2 \left[\mathbf{D}^- I_{(-1)11100101}^{B'} + (1-w) \mathbf{D}^- I_{011100101}^{B'} \right], \\
K_{47} &= \varepsilon^3 \left[\mathbf{D}^- I_{(-2)11100101}^{B'} + 2(1-w) \mathbf{D}^- I_{(-1)11100101}^{B'} + (1-w)^2 \mathbf{D}^- I_{011100101}^{B'} \right], \\
K_{48} &= \varepsilon^3 (w-v) \left[\mathbf{D}^- I_{0111(-1)0101}^{B'} - 2 \mathbf{D}^- I_{011000101}^{B'} \right], \\
K_{49} &= \varepsilon^3 r_2 \left[(1-w) \mathbf{D}^- I_{101100011}^{B'} + \mathbf{D}^- I_{001100011}^{B'} \right], \\
K_{50} &= \varepsilon^3 (1-w) \left[\mathbf{D}^- I_{1(-1)1100011}^{B'} + (1-w) \mathbf{D}^- I_{101100011}^{B'} \right], \\
K_{51} &= \varepsilon^3 (1-w) \left[\mathbf{D}^- I_{101100(-1)11}^{B'} + v \mathbf{D}^- I_{101100011}^{B'} \right], \\
K_{52} &= \varepsilon^3 r_3 \left[(1-w)^2 \mathbf{D}^- I_{111100101}^{B'} + 2(1-w) \mathbf{D}^- I_{011100101}^{B'} + \mathbf{D}^- I_{(-1)11100101}^{B'} \right], \\
K_{53} &= \varepsilon^3 r_3 \left[(1-w)^2 \mathbf{D}^- I_{111100011}^{B'} + (1-w) \left(\mathbf{D}^- I_{011100011}^{B'} + \mathbf{D}^- I_{101100011}^{B'} \right) + \mathbf{D}^- I_{001100011}^{B'} \right], \\
K_{54} &= \varepsilon^4 (1-2\varepsilon) v I_{101110021}^{B'}, \\
K_{55} &= \varepsilon^4 (1-2\varepsilon) v I_{112000111}^{B'}, \\
K_{56} &= \varepsilon^4 (1-2\varepsilon) v I_{012100111}^{B'}, \\
K_{57} &= \varepsilon^4 (1-2\varepsilon) v I_{011200111}^{B'}, \\
K_{58} &= \varepsilon^5 (1-2\varepsilon) v I_{111110011}^{B'}, \\
K_{59} &= \varepsilon^4 (1-2\varepsilon) v I_{112110011}^{B'}, \\
K_{60} &= \varepsilon^4 (1-2\varepsilon) r_3 \left[w I_{211110011}^{B'} + I_{112110011}^{B'} - 2\varepsilon I_{111110011}^{B'} \right], \\
K_{61} &= \varepsilon^4 (1-2\varepsilon) vw I_{112100111}^{B'}. \tag{58}
\end{aligned}$$

As boundary point we take again the point $(v, w) = (0, 1)$. We denote the boundary values for this system by C' . The non-zero boundary values are

$$\begin{aligned}
C'_{33} &= C'_{38} = C_1, \\
C'_9 &= C'_{21} = C'_{43} = C_{15}, \\
C'_7 &= C'_8 = C_{51}, \\
C'_{22} &= C'_{23} = C_{60}, \\
C'_{39} &= C_{71}, \\
C'_{41} &= C'_{47} = \frac{1}{3}C_{71}, \\
C'_2 &= C'_5 = C'_{15} = C'_{42} = -\frac{2}{3}C_{71},
\end{aligned}$$

$$C'_{48} = \frac{1}{2}C_1 - \frac{1}{2}C_{71}. \quad (59)$$

A.2 The sector B'_{509}

The sector B'_{509} involves the square roots (r_1, r_2, r_4) . These are rationalised by the variables (\tilde{x}, y) . The variable \tilde{x} is defined by

$$x = \tilde{x} \frac{(1 - \tilde{x})}{(1 + \tilde{x})}, \quad \tilde{x} = \frac{1}{2} \left(1 - x - \sqrt{x^2 - 6x + 1} \right). \quad (60)$$

The variables x and y are defined in eq. (25). The variable \tilde{x} rationalises simultaneously r_1 and r_3 and has already appeared in [68]. The root r_4 is given in the variable \tilde{x} by

$$r_4 = \frac{(1 + \tilde{x}^2)(1 - 2\tilde{x} - \tilde{x}^2)}{\tilde{x}(1 - \tilde{x}^2)}. \quad (61)$$

The value $x = 1 + i\delta$ (with δ being an infinitesimal small positive number) corresponds to $\tilde{x} = i$. We set

$$\tilde{x}' = i - \tilde{x}. \quad (62)$$

With this definition the value $x = 1 + i\delta$ corresponds to $\tilde{x}' = 0$.

With this rationalisation the differential one-forms translate to

$$d \ln(\tilde{p}_1), \dots, d \ln(\tilde{p}_{17}), \quad (63)$$

where $\tilde{p}_1 - \tilde{p}_{17}$ are polynomials in (\tilde{x}, y) . The differential one-form $\omega_6 = d \ln(w - v)$ leads also with this rationalisation to a quartic polynomial in \tilde{x} :

$$\tilde{p}_{17} = 1 - 2y + \tilde{x} + y^2 - 4y\tilde{x} + 2\tilde{x}^2 - y^2\tilde{x} - \tilde{x}^3 + 2y^2\tilde{x}^2 - 4\tilde{x}^3y + \tilde{x}^4 + \tilde{x}^3y^2 + 2\tilde{x}^4y + \tilde{x}^4y^2, \quad (64)$$

the remaining 16 polynomials $\tilde{p}_1 - \tilde{p}_{16}$ are at most quadratic in \tilde{x} . We note that \tilde{p}_{17} (and all other polynomials $\tilde{p}_1 - \tilde{p}_{16}$) are quadratic in y .

The master integrals for the sector B'_{509} are

$$\begin{aligned} L_1 &= K_1 = \varepsilon^3 (1 - w) \mathbf{D}^- I_{101010100}^{B'}, \\ L_2 &= K_2 = \varepsilon^3 \mathbf{D}^- I_{101(-1)10100}^{B'}, \\ L_3 &= K_3 = \varepsilon^3 r_2 \mathbf{D}^- I_{011010100}^{B'}, \\ L_4 &= K_4 = \varepsilon^3 \left[\mathbf{D}^- I_{(-1)11010100}^{B'} + (1 - w) \mathbf{D}^- I_{011010100}^{B'} \right], \\ L_5 &= K_5 = \varepsilon^3 \left[\mathbf{D}^- I_{01101(-1)100}^{B'} + v \mathbf{D}^- I_{011010100}^{B'} \right], \\ L_6 &= K_6 = \varepsilon^3 \left[\mathbf{D}^- I_{0110101(-1)0}^{B'} + v \mathbf{D}^- I_{011010100}^{B'} \right], \end{aligned}$$

$$\begin{aligned}
L_7 &= K_7 = \varepsilon^3 v \mathbf{D}^- I_{100011100}^{B'}, \\
L_8 &= K_8 = \varepsilon^3 v \mathbf{D}^- I_{001011100}^{B'}, \\
L_9 &= K_9 = \varepsilon^3 v \mathbf{D}^- I_{101000110}^{B'}, \\
L_{10} &= K_{13} = \varepsilon^3 r_2 \mathbf{D}^- I_{101(-1)11100}^{B'}, \\
L_{11} &= K_{14} = \varepsilon^3 (1-w) \left[\mathbf{D}^- I_{101(-1)11100}^{B'} + v \mathbf{D}^- I_{101011100}^{B'} \right], \\
L_{12} &= K_{15} = \varepsilon^3 \left[\mathbf{D}^- I_{101(-2)11100}^{B'} + v \mathbf{D}^- I_{101(-1)11100}^{B'} \right], \\
L_{13} &= K_{16} = \varepsilon^3 (1-w) \left[v \mathbf{D}^- I_{011011100}^{B'} + \mathbf{D}^- I_{011010100}^{B'} \right], \\
L_{14} &= K_{18} = \varepsilon^3 v (1-w) \mathbf{D}^- I_{101100110}^{B'}, \\
L_{15} &= K_{19} = \varepsilon^3 v r_2 \mathbf{D}^- I_{011100110}^{B'}, \\
L_{16} &= K_{20} = \varepsilon^3 v \left[\mathbf{D}^- I_{(-1)11100110}^{B'} + (1-w) \mathbf{D}^- I_{011100110}^{B'} \right], \\
L_{17} &= K_{21} = \varepsilon^3 v \mathbf{D}^- I_{0111(-1)0110}^{B'}, \\
L_{18} &= K_{22} = \varepsilon^3 v^2 \mathbf{D}^- I_{100101110}^{B'}, \\
L_{19} &= K_{23} = \varepsilon^3 v^2 \mathbf{D}^- I_{001101110}^{B'}, \\
L_{20} &= K_{33} = \varepsilon^3 \mathbf{D}^- I_{101000001}^{B'}, \\
L_{21} &= K_{35} = \varepsilon^3 (1-w) \mathbf{D}^- I_{101100001}^{B'}, \\
L_{22} &= K_{36} = \varepsilon^3 r_2 \mathbf{D}^- I_{011100001}^{B'}, \\
L_{23} &= K_{37} = \varepsilon^3 \left[\mathbf{D}^- I_{(-1)11100001}^{B'} + (1-w) \mathbf{D}^- I_{011100001}^{B'} \right], \\
L_{24} &= K_{38} = \varepsilon^3 \left[\mathbf{D}^- I_{01110(-1)001}^{B'} + v \mathbf{D}^- I_{011100001}^{B'} \right], \\
L_{25} &= K_{39} = \varepsilon^2 (1+4\varepsilon) \mathbf{D}^- I_{001100101}^{B'}, \\
L_{26} &= K_{40} = \varepsilon^3 r_1 \mathbf{D}^- I_{001100011}^{B'}, \\
L_{27} &= K_{41} = \varepsilon^3 \left[\mathbf{D}^- I_{(-1)01100011}^{B'} + (1-w) \mathbf{D}^- I_{001100011}^{B'} \right], \\
L_{28} &= K_{42} = \varepsilon^3 \left[\mathbf{D}^- I_{0011(-1)0011}^{B'} + v \mathbf{D}^- I_{001100011}^{B'} \right], \\
L_{29} &= K_{43} = \varepsilon^3 v \mathbf{D}^- I_{001000111}^{B'}, \\
L_{30} &= K_{45} = \varepsilon^3 (1-w) \left[(1-w) \mathbf{D}^- I_{101100101}^{B'} + \mathbf{D}^- I_{001100101}^{B'} \right], \\
L_{31} &= K_{46} = \varepsilon^3 r_2 \left[\mathbf{D}^- I_{(-1)11100101}^{B'} + (1-w) \mathbf{D}^- I_{011100101}^{B'} \right], \\
L_{32} &= K_{47} = \varepsilon^3 \left[\mathbf{D}^- I_{(-2)11100101}^{B'} + 2(1-w) \mathbf{D}^- I_{(-1)11100101}^{B'} + (1-w)^2 \mathbf{D}^- I_{011100101}^{B'} \right], \\
L_{33} &= K_{48} = \varepsilon^3 (w-v) \left[\mathbf{D}^- I_{0111(-1)0101}^{B'} - 2 \mathbf{D}^- I_{011000101}^{B'} \right], \\
L_{34} &= K_{49} = \varepsilon^3 r_2 \left[(1-w) \mathbf{D}^- I_{101100011}^{B'} + \mathbf{D}^- I_{001100011}^{B'} \right],
\end{aligned}$$

$$\begin{aligned}
L_{35} &= K_{50} = \varepsilon^3 (1-w) \left[\mathbf{D}^- I_{1(-1)1100011}^{B'} + (1-w) \mathbf{D}^- I_{101100011}^{B'} \right], \\
L_{36} &= K_{51} = \varepsilon^3 (1-w) \left[\mathbf{D}^- I_{101100(-1)11}^{B'} + v \mathbf{D}^- I_{101100011}^{B'} \right], \\
L_{37} &= K_{54} = \varepsilon^4 (1-2\varepsilon) v I_{101110021}^{B'}, \\
L_{38} &= K_{56} = \varepsilon^4 (1-2\varepsilon) v I_{012100111}^{B'}, \\
L_{39} &= K_{57} = \varepsilon^4 (1-2\varepsilon) v I_{011200111}^{B'}, \\
L_{40} &= \varepsilon^3 r_4 \left[v \mathbf{D}^- I_{0011(-1)1111}^{B'} + 2 \mathbf{D}^- I_{001100011}^{B'} \right], \\
L_{41} &= \varepsilon^4 (1-2\varepsilon) v I_{002101111}^{B'}, \\
L_{42} &= \varepsilon^5 (1-2\varepsilon) v I_{101101111}^{B'}, \\
L_{43} &= \varepsilon^4 (1-2\varepsilon) v (1-w) I_{102101111}^{B'}.
\end{aligned} \tag{65}$$

As boundary point we take again the point $(v, w) = (0, 1)$. We denote the boundary values for this system by C'' . The non-zero boundary values are

$$\begin{aligned}
C_{20}'' &= C_{24}'' = C_1, \\
C_9'' &= C_{17}'' = C_{29}'' = C_{15}, \\
C_7'' &= C_8'' = C_{51}, \\
C_{18}'' &= C_{19}'' = C_{60}, \\
C_{25}'' &= C_{71}, \\
C_{27}'' &= C_{32}'' = \frac{1}{3}C_{71}, \\
C_2'' &= C_5'' = C_{12}'' = C_{28}'' = -\frac{2}{3}C_{71}, \\
C_{33}'' &= \frac{1}{2}C_1 - \frac{1}{2}C_{71}.
\end{aligned} \tag{66}$$

B Supplementary material

Attached to the arxiv version of this article is an electronic file in ASCII format with Maple syntax, defining the quantities

$$A, B, C, J.$$

The matrix A appears in the differential equation

$$d\vec{J} = A\vec{J}. \tag{67}$$

The entries of the matrix A are linear combinations of $\omega_1, \dots, \omega_{19}$, defined in eq. (32). These differential forms are denoted by

$$\omega_1, \dots, \omega_{19}.$$

The matrices B and C appear in the differential equations in eq. (51).

The vector \mathbb{J} contains the results for the master integrals up to order ϵ^5 in terms of multiple polylogarithms. The variable ϵ is denoted by `eps`, ζ_2 , ζ_3 , ζ_5 by

`zeta_2, zeta_3, zeta_5,`

respectively. For the notation of multiple polylogarithms we give an example: $G(x'_7, x'_8, 1; x')$ is denoted by

`Glog([xp7, xp8, 1], xp).`

The file size is 10MB, roughly 90% of the file size is used for the ϵ^5 -term of J . The file size increases approximately by a factor of 10 for every additional order. This is expected: With an alphabet of

$$N_{\text{letter}} = 21 \tag{68}$$

letters we have

$$(N_{\text{letter}})^w \tag{69}$$

possible multiple polylogarithms at weight w . As the matrix A is sparse, the actual increase factor is of the order $O(10)$ and roughly a factor of two smaller than $N_{\text{letter}} = 21$. Note that the explicit expressions for J are convenient, but not essential: All information to any order in ϵ is stored compactly in the matrix A and the boundary values given in section 5.

References

- [1] G. P. Lepage, P. B. Mackenzie, and M. E. Peskin, (2014), arXiv:1404.0319.
- [2] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog, and B. Mistlberger, Phys. Rev. Lett. **114**, 212001 (2015), arXiv:1503.06056.
- [3] R. Bonciani *et al.*, JHEP **12**, 096 (2016), arXiv:1609.06685.
- [4] B. Mistlberger, JHEP **05**, 028 (2018), arXiv:1802.00833.
- [5] D. Kara, *Higher-Order Corrections to Higgs Boson Amplitudes with Full Quark Mass Dependence in Quantum Chromodynamics*, PhD thesis, Zurich U., 2018, arXiv:1908.09932.
- [6] C. Duhr, F. Dulat, and B. Mistlberger, Phys. Rev. Lett. **125**, 051804 (2020), arXiv:1904.09990.
- [7] A. H. Ajjath *et al.*, Phys. Rev. D **100**, 114016 (2019), arXiv:1906.09028.
- [8] J. Aparisi *et al.*, Phys. Rev. Lett. **128**, 122001 (2022), arXiv:2110.10202.
- [9] S. P. Martin, Phys. Rev. D **106**, 013007 (2022), arXiv:2203.05042.
- [10] M. Spira, CERN Yellow Reports: Monographs **3**, 123 (2020).
- [11] A. Chakraborty *et al.*, (2022), arXiv:2204.02422.
- [12] A. L. Kataev and V. S. Molokoedov, (2022), arXiv:2201.12073.
- [13] A. L. Kataev and V. T. Kim, Mod. Phys. Lett. A **9**, 1309 (1994).

- [14] K. G. Chetyrkin and A. Kwiatkowski, Nucl. Phys. **B461**, 3 (1996), arXiv:hep-ph/9505358.
- [15] K. G. Chetyrkin, J. H. Kühn, and M. Steinhauser, Nucl. Phys. **B505**, 40 (1997), arXiv:hep-ph/9705254.
- [16] K. G. Chetyrkin and M. Steinhauser, Phys. Lett. **B408**, 320 (1997), arXiv:hep-ph/9706462.
- [17] P. A. Baikov, K. G. Chetyrkin, and J. H. Kühn, Phys. Rev. Lett. **96**, 012003 (2006), arXiv:hep-ph/0511063.
- [18] W. Bernreuther, L. Chen, and Z.-G. Si, JHEP **07**, 159 (2018), arXiv:1805.06658.
- [19] A. Primo, G. Sasso, G. Somogyi, and F. Tramontano, Phys. Rev. **D99**, 054013 (2019), arXiv:1812.07811.
- [20] A. L. Kataev, JETP Lett. **66**, 327 (1997), arXiv:hep-ph/9708292.
- [21] A. Kwiatkowski and M. Steinhauser, Phys. Lett. **B338**, 66 (1994), arXiv:hep-ph/9405308, [Erratum: Phys. Lett.**B342**,455(1995)].
- [22] B. A. Kniehl and M. Spira, Nucl. Phys. **B432**, 39 (1994), arXiv:hep-ph/9410319.
- [23] L. Mihaila, B. Schmidt, and M. Steinhauser, Phys. Lett. **B751**, 442 (2015), arXiv:1509.02294.
- [24] M. Butenschön, F. Fugel, and B. A. Kniehl, Phys. Rev. Lett. **98**, 071602 (2007), arXiv:hep-ph/0612184.
- [25] M. Butenschön, F. Fugel, and B. A. Kniehl, Nucl. Phys. **B772**, 25 (2007), arXiv:hep-ph/0702215.
- [26] C. Anastasiou, F. Herzog, and A. Lazopoulos, JHEP **03**, 035 (2012), arXiv:1110.2368.
- [27] R. Mondini, M. Schiavi, and C. Williams, JHEP **06**, 079 (2019), arXiv:1904.08960.
- [28] R. Mondini, U. Schubert, and C. Williams, JHEP **12**, 058 (2020), arXiv:2006.03563.
- [29] A. Behring and W. Bizoń, JHEP **01**, 189 (2020), arXiv:1911.11524.
- [30] A. Smirnov, JHEP **10**, 107 (2008), arXiv:0807.3243.
- [31] A. V. Smirnov and F. S. Chuharev, Comput. Phys. Commun. **247**, 106877 (2020), arXiv:1901.07808.
- [32] R. N. Lee, (2012), arXiv:1212.2685.
- [33] R. N. Lee, J. Phys. Conf. Ser. **523**, 012059 (2014), arXiv:1310.1145.
- [34] C. Studerus, Comput. Phys. Commun. **181**, 1293 (2010), arXiv:0912.2546.
- [35] A. von Manteuffel and C. Studerus, (2012), arXiv:1201.4330.
- [36] P. Maierhöfer, J. Usovitsch, and P. Uwer, Comput. Phys. Commun. **230**, 99 (2018), arXiv:1705.05610.
- [37] J. Klappert, F. Lange, P. Maierhöfer, and J. Usovitsch, Comput. Phys. Commun. **266**, 108024 (2021), arXiv:2008.06494.
- [38] T. Peraro, JHEP **12**, 030 (2016), arXiv:1608.01902.
- [39] T. Peraro, JHEP **07**, 031 (2019), arXiv:1905.08019.
- [40] O. V. Tarasov, Phys. Rev. **D54**, 6479 (1996), hep-th/9606018.
- [41] O. V. Tarasov, Nucl. Phys. **B502**, 455 (1997), hep-ph/9703319.
- [42] E. Chaubey and S. Weinzierl, JHEP **05**, 185 (2019), arXiv:1904.00382.
- [43] M. Besier, D. Van Straten, and S. Weinzierl, Commun. Num. Theor. Phys. **13**, 253 (2019), arXiv:1809.10983.
- [44] M. Besier, P. Wasser, and S. Weinzierl, Comput. Phys. Commun. **253**, 107197 (2020), arXiv:1910.13251.
- [45] J. Moser, Mathematische Zeitschrift **1**, 379 (1959).
- [46] R. N. Lee, JHEP **04**, 108 (2015), arXiv:1411.0911.
- [47] R. N. Lee and A. A. Pomeransky, (2017), arXiv:1707.07856.

- [48] M. Prausa, *Comput. Phys. Commun.* **219**, 361 (2017), arXiv:1701.00725.
- [49] O. Gituliar and V. Magerya, *Comput. Phys. Commun.* **219**, 329 (2017), arXiv:1701.04269.
- [50] R. N. Lee, *Comput. Phys. Commun.* **267**, 108058 (2021), arXiv:2012.00279.
- [51] P. A. Baikov, *Nucl. Instrum. Meth.* **A389**, 347 (1997), arXiv:hep-ph/9611449.
- [52] H. Frellesvig and C. G. Papadopoulos, *JHEP* **04**, 083 (2017), arXiv:1701.07356.
- [53] F. Cachazo, (2008), arXiv:0803.1988.
- [54] N. Arkani-Hamed, J. L. Bourjaily, F. Cachazo, and J. Trnka, *JHEP* **06**, 125 (2012), arXiv:1012.6032.
- [55] S. Weinzierl, *Feynman Integrals* (Springer, 2022), arXiv:2201.03593.
- [56] J. M. Henn, *Phys. Rev. Lett.* **110**, 251601 (2013), arXiv:1304.1806.
- [57] A. B. Goncharov, *Math. Res. Lett.* **5**, 497 (1998).
- [58] A. B. Goncharov, (2001), math.AG/0103059.
- [59] J. M. Borwein, D. M. Bradley, D. J. Broadhurst, and P. Lisonek, *Trans. Amer. Math. Soc.* **353:3**, 907 (2001), math.CA/9910045.
- [60] C. Bauer, A. Frink, and R. Kreckel, *J. Symbolic Computation* **33**, 1 (2002), cs.sc/0004015.
- [61] J. Vollinga and S. Weinzierl, *Comput. Phys. Commun.* **167**, 177 (2005), hep-ph/0410259.
- [62] X. Liu and Y.-Q. Ma, (2022), arXiv:2201.11669.
- [63] X. Liu, Y.-Q. Ma, and C.-Y. Wang, *Phys. Lett. B* **779**, 353 (2018), arXiv:1711.09572.
- [64] Z.-F. Liu and Y.-Q. Ma, (2022), arXiv:2201.11637.
- [65] C. Bogner and S. Weinzierl, *Comput. Phys. Commun.* **178**, 596 (2008), arXiv:0709.4092.
- [66] T. Binoth and G. Heinrich, *Nucl. Phys.* **B585**, 741 (2000), hep-ph/0004013.
- [67] M. Besier, P. Wasser, and S. Weinzierl, (2019), arXiv:1910.13251.
- [68] L. Adams, E. Chaubey, and S. Weinzierl, *JHEP* **10**, 206 (2018), arXiv:1806.04981.