

# Three-loop master integrals for the Higgs boson self-energy proportional to the top and bottom Yukawa couplings

Ekta Chaubey <sup>a</sup>, Ina Hönemann <sup>b</sup> and Stefan Weinzierl <sup>b</sup>

<sup>a</sup> *Physics Department, University of Turin and INFN Turin,  
Via Pietro Giuria 1, I-10125, Turin, Italy*

<sup>b</sup> *PRISMA Cluster of Excellence, Institut für Physik,  
Johannes Gutenberg-Universität Mainz,  
D - 55099 Mainz, Germany*

## Abstract

We present analytic results for all master integrals contributing to the three-loop Higgs self-energy diagrams of order  $O(\alpha^2\alpha_s)$  proportional to the product of Yukawa couplings  $y_b y_t$ . These diagrams contain internal top- and  $W$ -propagators. We define master integrals of uniform weight, such that the associated differential equation is in  $\epsilon$ -factorised form. The occurring square roots are rationalised and all master integrals are expressed in multiple polylogarithms.

# 1 Introduction

Higgs precision physics is at the core of the experimental LHC program. On the theoretical side this requires higher-order calculations in perturbation theory [1–12]. The Higgs boson self-energy is an essential building block for many observables. Through renormalisation it enters the Higgs mass definition. Furthermore, the imaginary part of the Higgs boson self-energy is directly related through the optical theorem to the Higgs decay rate [13–29].

In this article we consider three-loop contributions to the Higgs boson self-energy, which are of order  $O(\alpha^2\alpha_s)$  proportional to the product of Yukawa couplings  $y_b y_t$ . The corresponding Feynman diagrams contain internal top- and  $W$ -propagators and are therefore among the more complicated Feynman integrals contributing at this order in perturbation theory. These integrals have been computed for  $p^2 = m_H^2$  approximately as an expansion in  $m_H^2/m_t^2$  in ref. [23].

In this paper we present the analytic calculation of the relevant master integrals. We keep the full dependence on the external momentum squared  $p^2$  and the heavy particle masses ( $m_t$  and  $m_W$ ), but neglect the  $b$ -quark mass. The Feynman integrals depend on two dimensionless variables, which we may take initially as

$$v = \frac{p^2}{m_t^2}, \quad w = \frac{m_W^2}{m_t^2}. \quad (1)$$

There are 105 master integrals. We define a basis of master integrals of uniform weight. In this basis, the associated differential equation is  $\epsilon$ -factorised and all occurring differential one-forms are dlog-forms. We encounter two square roots, which can be rationalised simultaneously. This is done by a change of variables from  $(v, w)$  to  $(x, y)$ . The variables  $x$  and  $y$  will be defined in the main part of this paper. This change of variables allows us to express all master integrals to any order in the dimensional regularisation parameter  $\epsilon$  in terms of multiple polylogarithms. The alphabet for the multiple polylogarithms consists of 21 letters.

As a side remark we mention that at three-loop order there are additional diagrams containing internal top- and  $W$ -propagators. In these diagrams the Higgs boson couples either to a  $W$ -boson and a  $b$ -quark or to two  $b$ -quarks. The corresponding Feynman integrals have different/additional square roots. As they are different from an analytical point of view, they are not included in this paper, but will be dealt with in a separate publication.

This paper is organised as follows: In section 2 we introduce our notation for the three-loop integrals. The master integrals of uniform weight are presented in section 3. The differential equation for these master integrals is in  $\epsilon$ -factorised form. The differential equation is discussed in section 4. The solution of the differential equation requires boundary values, which we give in section 5. Analytical results for the master integrals are presented in section 6. Numerical results are presented in section 7. Finally, our conclusions are given in section 8. In appendix A we describe the content of the supplementary electronic file attached to the arxiv version of this article.




Figure 1: Examples of three-loop Higgs self-energy diagrams containing a top propagator and a  $W$ -propagator and being proportional to the product of Yukawa couplings  $y_b y_t$ . The internal masses of the propagators are encoded by the colour of the propagators: massless (black),  $m_t$  (green),  $m_W$  (red).

## 2 Notation

We consider three-loop Higgs self-energy diagrams of order  $O(\alpha^2 \alpha_s)$  proportional to the product of Yukawa couplings  $y_b y_t$ . Examples of diagrams are shown in fig. 1. The diagrams shown in fig. 1 have all eight propagators and define the top sectors. Simpler diagrams, obtained from the ones shown in fig. 1 by pinching a subset of the propagators, are not shown.

In this paper we calculate all relevant master integrals. As we have three independent loop momenta and one independent external momentum, we need nine propagators in order to be able to express any scalar product involving at least one loop momentum in terms of inverse propagators. We therefore consider auxiliary graphs with nine propagators, shown in fig. 2. We label these auxiliary graphs topology  $A$ ,  $B$ ,  $C$  and  $D$  in correspondence with the diagrams shown in fig. 1.

As a consequence we consider the integrals

$$I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X = e^{3\gamma_E \epsilon} (\mu^2)^{v - \frac{3}{2}D} \int \frac{d^D k_1}{i\pi^{\frac{D}{2}}} \frac{d^D k_2}{i\pi^{\frac{D}{2}}} \frac{d^D k_3}{i\pi^{\frac{D}{2}}} \prod_{j=1}^9 \frac{1}{(P_j^X)^{v_j}},$$

$$X \in \{A, B, C, D\}, \quad (2)$$

where  $D = 4 - 2\epsilon$  denotes the number of space-time dimensions,  $\gamma_E$  denotes the Euler-Mascheroni constant,  $\mu$  is an arbitrary scale introduced to render the Feynman integral dimensionless, and the quantity  $\mathbf{v}$  is defined by

$$\mathbf{v} = \sum_{j=1}^9 v_j. \quad (3)$$

The inverse propagators  $P_j^X$  are defined as follows:

Topology A:

$$\begin{aligned} P_1^A &= -k_1^2 + m_t^2, & P_2^A &= -(k_1 - p)^2 + m_t^2, & P_3^A &= -(k_1 + k_2)^2, \\ P_4^A &= -k_2^2 + m_t^2, & P_5^A &= -(k_2 + k_3)^2 + m_W^2, & P_6^A &= -(k_2 + p)^2 + m_t^2, \\ P_7^A &= -k_3^2, & P_8^A &= -(k_3 - p)^2, & P_9^A &= -(k_1 - k_3)^2 + m_t^2. \end{aligned} \quad (4)$$

Topology B:

$$\begin{aligned} P_1^B &= -k_1^2 + m_t^2, & P_2^B &= -(k_1 - p)^2 + m_t^2, & P_3^B &= -(k_1 + k_2)^2 + m_W^2, \\ P_4^B &= -k_2^2, & P_5^B &= -(k_2 + k_3)^2, & P_6^B &= -(k_2 + p)^2, \\ P_7^B &= -k_3^2, & P_8^B &= -(k_3 - p)^2, & P_9^B &= -(k_1 - k_3)^2 + m_W^2. \end{aligned} \quad (5)$$

Topology C:

$$\begin{aligned} P_1^C &= -k_1^2 + m_t^2, & P_2^C &= -(k_1 - p)^2 + m_t^2, & P_3^C &= -(k_1 + k_2)^2, \\ P_4^C &= -(k_1 + k_2 - k_3)^2, & P_5^C &= -k_2^2 + m_W^2, & P_6^C &= -(k_2 - k_3 + p)^2 + m_t^2, \\ P_7^C &= -k_3^2, & P_8^C &= -(k_3 - p)^2, & P_9^C &= -(k_2 + p)^2 + m_t^2. \end{aligned} \quad (6)$$

Topology D:

$$\begin{aligned} P_1^D &= -k_1^2 + m_t^2, & P_2^D &= -(k_1 - p)^2 + m_t^2, & P_3^D &= -(k_1 - k_3)^2 + m_W^2, \\ P_4^D &= -(k_1 + k_2)^2 + m_t^2, & P_5^D &= -k_2^2, & P_6^D &= -(k_2 + k_3)^2, \\ P_7^D &= -k_3^2, & P_8^D &= -(k_3 - p)^2, & P_9^D &= -(k_2 + p)^2. \end{aligned} \quad (7)$$

We introduce the sector id for the Feynman integral  $I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X$  by

$$N_{\text{id}}^X = \sum_{j=1}^9 2^{j-1} \Theta(v_j > 0). \quad (8)$$

With the help of the sector id we may specify all the reductions which we will need: We have to reduce in every topology  $A - D$  the top sector 255. These top sectors correspond to the diagrams shown in fig. 1. The integral reduction gives us in total 105 master integrals. A detailed list is




Figure 2: Graphs for the four topologies. The internal masses of the propagators are encoded by the colour of the propagators: massless (black),  $m_t$  (green),  $m_W$  (red).

Topology and sector id	$N_{\text{master}}$
$A_{255}$	50
$A_{255} + B_{255}$	67
$A_{255} + B_{255} + C_{255}$	104
$A_{255} + B_{255} + C_{255} + D_{255}$	105

(9)

Table 1: The number of master integrals. The table gives the cumulative sum, as for example topology  $B$  contains master integrals common to topology  $A$  and  $B$ .

Topology and sector id	$N_{\text{master}}$
$A_{255}$	50
$B_{255}$	32
$C_{255}$	95
$D_{255}$	67

(10)

Table 2: The number of master integrals for a given top sector, if we treat each top sector individually.

given in table 1. Note that this list gives the cumulative number of master integrals. This takes into account that certain master integral occur in more than one topology or sector. These master integrals are only counted once. If on the other hand we treat each top sector individually, we find the number of master integrals per top sector as shown in table 2. In this case the total number of master integrals is higher, as we recompute master integrals, which are identical or are related by symmetries. The integral reduction can be carried out with public available computer programs like FIRE/Litered [30–33], Reduze [34, 35], Kira [36, 37] or FiniteFlow [38, 39].

Let us point out, that it is non-trivial that the total number of master integrals is 105 and not 106. There is one relation not easily found by standard integration-by-parts reduction programs<sup>1</sup>:

$$\begin{aligned}
I_{011011100}^B &= \frac{1}{(m_t^2 - m_W^2)} (I_{001011100}^B - I_{100011100}^B) \\
&\quad - \frac{1}{s} \left( I_{01111(-1)100}^A - I_{011010100}^A - I_{(-1)11110100}^A \right) \\
&\quad - \frac{3}{s(m_t^2 - m_W^2)} \left( I_{011(-1)10100}^A - I_{01101(-1)100}^A \right). \tag{11}
\end{aligned}$$

Without this relation we may end up with a non-integrable system.

The Feynman parameter representations for the four topologies are given by

$$I_{\mathbf{v}_1 \mathbf{v}_2 \mathbf{v}_3 \mathbf{v}_4 \mathbf{v}_5 \mathbf{v}_6 \mathbf{v}_7 \mathbf{v}_8 \mathbf{v}_9}^X = e^{3\gamma_E \epsilon} \frac{\Gamma(\mathbf{v} - \frac{3}{2}D)}{9 \prod_{j=1}^9 \Gamma(\mathbf{v}_j)} \int_{x_i \geq 0} d^9 x \delta \left( 1 - \sum_{j=1}^9 x_j \right) \left( \prod_{j=1}^9 x_j^{\mathbf{v}_j - 1} \right) \frac{\mathcal{U}_X^{\mathbf{v} - 2D}}{\mathcal{F}_X^{\mathbf{v} - \frac{3}{2}D}}, \tag{12}$$

where  $\mathcal{U}_X$  and  $\mathcal{F}_X$  denote the first and second graph polynomial, respectively. The first graph polynomials for the individual topologies are given by

$$\begin{aligned}
\mathcal{U}_A &= \mathcal{U}_B = x_{12} x_3 x_5 + x_{12} x_3 x_7 + x_{12} x_3 x_9 + x_{12} x_4 x_5 + x_{12} x_4 x_7 + x_{12} x_4 x_9 + x_{12} x_5 x_7 \\
&\quad + x_{12} x_5 x_9 + x_3 x_4 x_5 + x_3 x_4 x_7 + x_3 x_4 x_9 + x_3 x_5 x_7 + x_3 x_7 x_9 + x_4 x_5 x_9 + x_4 x_7 x_9 \\
&\quad + x_5 x_7 x_9,
\end{aligned}$$

$$\mathcal{U}_C = x_{12} x_3 x_4 + x_{12} x_3 x_6 + x_{12} x_3 x_7 + x_{12} x_4 x_5 + x_{12} x_4 x_7 + x_{12} x_5 x_6 + x_{12} x_5 x_7$$

---

<sup>1</sup>We thank Johann Usovitsch for contributing this relation.

$$\begin{aligned}
& +x_{12}x_6x_{78} + x_3x_4x_{59} + x_3x_4x_6 + x_3x_{59}x_6 + x_3x_{59}x_{78} + x_3x_6x_{78} + x_4x_{59}x_6 + x_4x_{59}x_{78} \\
& +x_4x_6x_{78}, \\
\mathcal{U}_D = & x_{12}x_3x_4 + x_{12}x_3x_{59} + x_{12}x_3x_6 + x_{12}x_4x_6 + x_{12}x_4x_{78} + x_{12}x_{59}x_6 + x_{12}x_{59}x_{78} \\
& +x_{12}x_6x_{78} + x_3x_4x_{59} + x_3x_4x_{78} + x_3x_{59}x_6 + x_3x_{59}x_{78} + x_3x_6x_{78} + x_4x_{59}x_6 + x_4x_{59}x_{78} \\
& +x_4x_6x_{78}, \tag{13}
\end{aligned}$$

where we used the abbreviation  $x_{ij} = x_i + x_j$ .

## 2.1 Dimensional shift relations and differential equations

Let us introduce an operator  $\mathbf{i}^+$ , which raises the power of the propagator  $i$  by one and multiplies by  $v_i$ , e.g.

$$\mathbf{1}^+ I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X = v_1 \cdot I_{(v_1+1)v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X. \tag{14}$$

In addition we define two operators  $\mathbf{D}^\pm$ , which shift the dimension of space-time by two through

$$\mathbf{D}^\pm I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X(D) = I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X(D \pm 2). \tag{15}$$

The dimensional shift relations read [40, 41]

$$\begin{aligned}
\mathbf{D}^- I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X(D) = & \tag{16} \\
\mathcal{U}_X(\mathbf{1}^+, \mathbf{2}^+, \mathbf{3}^+, \mathbf{4}^+, \mathbf{5}^+, \mathbf{6}^+, \mathbf{7}^+, \mathbf{8}^+, \mathbf{9}^+) I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X(D). &
\end{aligned}$$

We denote by  $J_{m_i}^X$  the set of internal edge indices, which correspond to the propagation of a particle with mass  $m_i$ . For example

$$J_{m_t}^A = \{1, 2, 4, 6, 9\}. \tag{17}$$

With this notation the differential equations read

$$\begin{aligned}
\mu^2 \frac{\partial}{\partial m_t^2} I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X &= - \sum_{j \in J_{m_t}^X} \mathbf{j}^+ I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X, \\
\mu^2 \frac{\partial}{\partial m_W^2} I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X &= - \sum_{j \in J_{m_W}^X} \mathbf{j}^+ I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X, \\
\mu^2 \frac{\partial}{\partial s} I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X &= \left( \frac{3}{2}D - v \right) \frac{\mu^2}{s} I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X + \frac{m_t^2}{s} \sum_{j \in J_{m_t}^X} \mathbf{j}^+ I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X \\
&+ \frac{m_W^2}{s} \sum_{j \in J_{m_W}^X} \mathbf{j}^+ I_{v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_8 v_9}^X. \tag{18}
\end{aligned}$$

For the derivative with respect to  $s$  we used the scaling relation. This has the advantage that on the right-hand side at most one raising operator occurs. This minimises the integral reductions we have to compute.

## 2.2 Variables

In the following we will set

$$\mu^2 = m_t^2. \quad (19)$$

The master integrals depend then kinematically on two dimensionless quantities. We set

$$v = \frac{p^2}{m_t^2}, \quad w = \frac{m_W^2}{m_t^2}. \quad (20)$$

We will encounter the following two square roots

$$r_1 = \sqrt{-v(4-v)}, \quad r_2 = \sqrt{\lambda(v, w, 1)}. \quad (21)$$

The Källén function is defined by

$$\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2yz - 2zx. \quad (22)$$

The variable transformation from  $(v, w)$  to  $(x, y)$ , where the latter variables are defined by [42–44]

$$v = -\frac{(1-x)^2}{x}, \quad w = \frac{(1-y+2xy)(x-2y+xy)}{x(1-y^2)}, \quad (23)$$

rationalises the square roots  $r_1$  and  $r_2$ :

$$r_1 = \frac{1-x^2}{x}, \quad r_2 = \frac{(1-x) \left[ (1-y)^2 + x(1+y)^2 \right]}{x(1-y^2)}. \quad (24)$$

The inverse transformation is given by

$$x = \frac{1}{2}(2-v-r_1), \quad y = \frac{r_2-r_1}{1-w+2v}, \quad (25)$$

such that  $x = 0$  corresponds to  $v = \infty$  and  $y = 0$  corresponds to  $w = 1$ . We note that the point  $(v, w) = (0, 1)$  is blown up in  $(x, y)$ -space to the hypersurface  $x = 1$ . This is convenient, as it allows us to obtain the results for the master integrals by integrating the differential equation in  $x$  with constant  $y$ . We make a final change of coordinates and we introduce

$$x' = 1-x. \quad (26)$$

This transforms the starting point of the integration from  $x = 1$  to  $x' = 0$  and is helpful for the conversion to multiple polylogarithms.



### 3 Master integrals

In this section we define master integrals of uniform weight. We have 105 master integrals and 59 master topologies (or sectors). These are summarised in table 3 and 4. We denote the master integrals of uniform weight by  $J_1$ - $J_{105}$ . For all these master integrals we find compact expressions in terms of pre-canonical Feynman integrals. The master integrals  $J_1$ - $J_{105}$  are defined by

$$\begin{aligned}
J_1 &= \varepsilon^3 \mathbf{D}^- I_{100110000}^A, \\
J_2 &= \varepsilon^3 r_1 \mathbf{D}^- I_{110110000}^A, \\
J_3 &= \varepsilon^3 r_1 \mathbf{D}^- I_{011110000}^A, \\
J_4 &= \varepsilon^3 \mathbf{D}^- I_{(-1)11110000}^A, \\
J_5 &= \varepsilon^3 (1-w) \mathbf{D}^- I_{101010100}^A, \\
J_6 &= \varepsilon^3 \mathbf{D}^- I_{101(-1)10100}^A, \\
J_7 &= \varepsilon^3 r_2 \mathbf{D}^- I_{011010100}^A, \\
J_8 &= \varepsilon^3 \left[ \mathbf{D}^- I_{011(-1)10100}^A - (1-w) \mathbf{D}^- I_{011010100}^A \right], \\
J_9 &= \varepsilon^3 \mathbf{D}^- I_{01101(-1)100}^A, \\
J_{10} &= \varepsilon^3 \left[ \mathbf{D}^- I_{0110101(-1)0}^A + v \mathbf{D}^- I_{011010100}^A \right], \\
J_{11} &= \varepsilon^3 (1-w) \mathbf{D}^- I_{100110100}^A, \\
J_{12} &= \varepsilon^3 r_2 \mathbf{D}^- I_{100011100}^A, \\
J_{13} &= \varepsilon^3 \left[ \mathbf{D}^- I_{100(-1)11100}^A - (1-w) \mathbf{D}^- I_{100011100}^A \right], \\
J_{14} &= \varepsilon^3 \left[ \mathbf{D}^- I_{1000111(-1)0}^A + v \mathbf{D}^- I_{100011100}^A \right], \\
J_{15} &= \varepsilon^3 v \mathbf{D}^- I_{100100110}^A, \\
J_{16} &= \varepsilon^3 v \mathbf{D}^- I_{100010110}^A, \\
J_{17} &= \varepsilon^3 v(4-v) \mathbf{D}^- I_{110111000}^A, \\
J_{18} &= \varepsilon^3 r_1 \left[ \mathbf{D}^- I_{111(-1)10100}^A - (1-w) \mathbf{D}^- I_{111010100}^A \right], \\
J_{19} &= \varepsilon^3 r_1 \left[ \mathbf{D}^- I_{111(-1)10100}^A - \mathbf{D}^- I_{011010100}^A \right], \\
J_{20} &= \varepsilon^3 (1-w) r_1 \mathbf{D}^- I_{110110100}^A, \\
J_{21} &= \varepsilon^3 r_1 \left[ (1-w) \mathbf{D}^- I_{011110100}^A - \mathbf{D}^- I_{011010100}^A \right], \\
J_{22} &= \varepsilon^3 (1-w) \left[ \mathbf{D}^- I_{(-1)11110100}^A - \mathbf{D}^- I_{011010100}^A \right], \\
J_{23} &= \varepsilon^3 (1-w) \mathbf{D}^- I_{01111(-1)100}^A, \\
J_{24} &= \varepsilon^3 r_1 r_2 \mathbf{D}^- I_{110011100}^A, \\
J_{25} &= \varepsilon^3 r_1 \left[ \mathbf{D}^- I_{110(-1)11100}^A - (1-w) \mathbf{D}^- I_{110011100}^A \right],
\end{aligned}$$

$$\begin{aligned}
J_{26} &= \varepsilon^3 r_1 \left[ \mathbf{D}^- I_{1100111(-1)0}^A + v \mathbf{D}^- I_{110011100}^A \right], \\
J_{27} &= \varepsilon^3 r_2 \mathbf{D}^- I_{101(-1)11100}^A, \\
J_{28} &= \varepsilon^3 r_1 \left[ \mathbf{D}^- I_{101(-1)11100}^A - (1-w) \mathbf{D}^- I_{101011100}^A \right], \\
J_{29} &= \varepsilon^3 \left[ \mathbf{D}^- I_{101(-2)11100}^A - (1-w) \mathbf{D}^- I_{101(-1)11100}^A \right], \\
J_{30} &= \varepsilon^3 (1-v) \left[ \mathbf{D}^- I_{101(-1)11100}^A + \mathbf{D}^- I_{10101110(-1)}^A - (1-w) \mathbf{D}^- I_{101011100}^A \right], \\
J_{31} &= \varepsilon^3 r_1 \left[ (1-w) \mathbf{D}^- I_{100111100}^A - \mathbf{D}^- I_{100011100}^A \right], \\
J_{32} &= \varepsilon^3 v r_1 \mathbf{D}^- I_{110100110}^A, \\
J_{33} &= \varepsilon^3 v r_1 \mathbf{D}^- I_{011100110}^A, \\
J_{34} &= \varepsilon^3 v \mathbf{D}^- I_{(-1)11100110}^A, \\
J_{35} &= \varepsilon^3 v r_1 \mathbf{D}^- I_{110010110}^A, \\
J_{36} &= \varepsilon^3 (1-w) \left[ v \mathbf{D}^- I_{101010110}^A + \mathbf{D}^- I_{011010100}^A \right], \\
J_{37} &= \varepsilon^4 (1-2\varepsilon) v I_{111021100}^A, \\
J_{38} &= \varepsilon^4 (1-2\varepsilon) v I_{111011200}^A, \\
J_{39} &= \varepsilon^3 v (4-v) \left[ (1-w) \mathbf{D}^- I_{110111100}^A - \mathbf{D}^- I_{110011100}^A \right], \\
J_{40} &= \varepsilon^3 v r_1 \left[ \mathbf{D}^- I_{11101011(-1)}^A - (1-w) \mathbf{D}^- I_{111010110}^A \right], \\
J_{41} &= \varepsilon^4 (1-2\varepsilon) v I_{112010110}^A, \\
J_{42} &= \varepsilon^4 (1-2\varepsilon) v I_{111020110}^A, \\
J_{43} &= \varepsilon^4 (1-2\varepsilon) v I_{021110110}^A, \\
J_{44} &= \varepsilon^4 (1-2\varepsilon) v I_{012110110}^A, \\
J_{45} &= \varepsilon^3 v^2 (4-v) \mathbf{D}^- I_{110101110}^A, \\
J_{46} &= \varepsilon^4 (1-\varepsilon) (1-2\varepsilon) v I_{100111110}^A, \\
J_{47} &= \varepsilon^5 (1-2\varepsilon) v I_{111110110}^A, \\
J_{48} &= \varepsilon^4 (1-2\varepsilon) v (1-w) I_{111120110}^A, \\
J_{49} &= \varepsilon^4 (1-2\varepsilon) v \left[ I_{211110110}^A + \frac{1}{2} (2-v+r_1) I_{121110110}^A \right], \\
J_{50} &= \varepsilon^4 (1-2\varepsilon) v r_1 I_{210111110}^A, \\
J_{51} &= \varepsilon^3 v \mathbf{D}^- I_{100011100}^B, \\
J_{52} &= \varepsilon^3 v \mathbf{D}^- I_{001011100}^B, \\
J_{53} &= \varepsilon^3 v r_1 \mathbf{D}^- I_{110011100}^B, \\
J_{54} &= \varepsilon^3 r_2 \mathbf{D}^- I_{101(-1)11100}^B, \\
J_{55} &= \varepsilon^3 (1-w) \left[ \mathbf{D}^- I_{101(-1)11100}^B + v \mathbf{D}^- I_{101011100}^B \right],
\end{aligned}$$

$$\begin{aligned}
J_{56} &= \varepsilon^3 \left[ \mathbf{D}^- I_{101(-2)11100}^B + \nu \mathbf{D}^- I_{101(-1)11100}^B \right], \\
J_{57} &= \varepsilon^3 \nu r_2 \mathbf{D}^- I_{011100110}^B, \\
J_{58} &= \varepsilon^3 \nu \left[ \mathbf{D}^- I_{(-1)11100110}^B - (1-w) \mathbf{D}^- I_{011100110}^B \right], \\
J_{59} &= \varepsilon^3 \nu \mathbf{D}^- I_{0111(-1)0110}^B, \\
J_{60} &= \varepsilon^3 \nu^2 \mathbf{D}^- I_{100101110}^B, \\
J_{61} &= \varepsilon^3 \nu^2 \mathbf{D}^- I_{001101110}^B, \\
J_{62} &= \varepsilon^4 (1-2\varepsilon) \nu I_{111021100}^B, \\
J_{63} &= \varepsilon^3 r_1 \left\{ (1-w) \mathbf{D}^- I_{111(-1)11100}^B - \mathbf{D}^- I_{101(-1)11100}^B \right\}, \\
J_{64} &= \varepsilon^3 \nu r_1 \left\{ (1-w) \mathbf{D}^- I_{111100110}^B - \mathbf{D}^- I_{011100110}^B \right\}, \\
J_{65} &= \varepsilon^3 \nu^2 r_1 \mathbf{D}^- I_{110101110}^B, \\
J_{66} &= \varepsilon^5 (1-2\varepsilon) \nu I_{111110110}^B, \\
J_{67} &= \varepsilon^4 (1-2\varepsilon)^2 \nu I_{111101110}^B, \\
J_{68} &= \varepsilon^3 r_1 \mathbf{D}^- I_{101001100}^C, \\
J_{69} &= \varepsilon^3 \mathbf{D}^- I_{1(-1)1001100}^C, \\
J_{70} &= \varepsilon^3 \mathbf{D}^- I_{10100110(-1)}^C, \\
J_{71} &= \varepsilon^2 (1+4\varepsilon) \mathbf{D}^- I_{011001100}^C, \\
J_{72} &= \varepsilon^3 r_1 \mathbf{D}^- I_{11100110(-1)}^C, \\
J_{73} &= \varepsilon^3 r_2 \left[ (1-w) \mathbf{D}^- I_{101011100}^C + \mathbf{D}^- I_{101001100}^C \right], \\
J_{74} &= \varepsilon^3 (1-w) \mathbf{D}^- I_{10101110(-1)}^C, \\
J_{75} &= \varepsilon^3 (1-w) \left[ \mathbf{D}^- I_{1010111(-1)0}^C + \nu \mathbf{D}^- I_{101011100}^C \right], \\
J_{76} &= \varepsilon^3 r_2 \mathbf{D}^- I_{01101110(-1)}^C, \\
J_{77} &= \varepsilon^3 \mathbf{D}^- I_{01101110(-2)}^C, \\
J_{78} &= \varepsilon^3 (w-\nu) \left\{ \mathbf{D}^- I_{011(-1)11100}^C - 2 \mathbf{D}^- I_{100011100}^A \right\}, \\
J_{79} &= \varepsilon^3 r_1 \left\{ 2 \mathbf{D}^- I_{1(-1)1101010}^C + \nu \mathbf{D}^- I_{101101010}^C - \mathbf{D}^- I_{011001100}^C - 2 \mathbf{D}^- I_{101100010}^C \right\}, \\
J_{80} &= \varepsilon^3 \nu \left[ 2(1-3\varepsilon) \left( I_{202101010}^C - \varepsilon I_{102101010}^C \right) + \varepsilon(1-2\varepsilon) I_{101201010}^C \right], \\
J_{81} &= \varepsilon^3 (1-w) \left[ (1-w) \mathbf{D}^- I_{101011010}^C + \mathbf{D}^- I_{011001100}^C \right], \\
J_{82} &= \varepsilon^3 r_1 \left\{ (1-w) \mathbf{D}^- I_{11101110(-1)}^C - \mathbf{D}^- I_{01101110(-1)}^C \right\}, \\
J_{83} &= \varepsilon^4 (1-2\varepsilon) \nu I_{1111011200}^C, \\
J_{84} &= \varepsilon^4 (1-2\varepsilon) \nu I_{111012100}^C,
\end{aligned}$$

$$\begin{aligned}
J_{85} &= \varepsilon^4 (1 - 2\varepsilon) v I_{112101010}^C, \\
J_{86} &= \varepsilon^3 (1 - w) r_1 \left[ (1 - w) \mathbf{D}^- I_{111011010}^C - \mathbf{D}^- I_{101011100}^C + \mathbf{D}^- I_{111001100}^C \right], \\
J_{87} &= \varepsilon^4 (1 - 2\varepsilon) \left[ I_{101111010}^C - \frac{1}{2} (1 - w) I_{201111010}^C - \frac{3}{2} I_{101211(-1)10}^C + \frac{1}{4} v I_{101201010}^C \right. \\
&\quad \left. - \frac{1}{2} I_{201101010}^C + \frac{1}{2} I_{200111010}^C + \frac{3}{2} I_{100211010}^C \right] + \frac{\varepsilon^3}{2} (1 - 3\varepsilon) v \left[ I_{202101010}^C - \varepsilon I_{102101010}^C \right], \\
J_{88} &= \varepsilon^4 (1 - 2\varepsilon) v I_{101211010}^C, \\
J_{89} &= \varepsilon^4 (1 - 2\varepsilon) v I_{012111010}^C, \\
J_{90} &= \varepsilon^4 (1 - 2\varepsilon) v I_{112001110}^C, \\
J_{91} &= \varepsilon^4 (1 - 2\varepsilon) v I_{111002110}^C, \\
J_{92} &= \varepsilon^2 r_1 \left[ (1 - 2\varepsilon)^2 v I_{211001210}^C - \varepsilon (1 - 2\varepsilon) I_{111002110}^C - I_{202002100}^C - v I_{210200210}^A \right], \\
J_{93} &= \varepsilon^4 (1 - 2\varepsilon) v I_{021011110}^C, \\
J_{94} &= \varepsilon^4 (1 - 2\varepsilon) v I_{012011110}^C, \\
J_{95} &= \varepsilon^5 (1 - 2\varepsilon) v I_{111111010}^C, \\
J_{96} &= \varepsilon^4 (1 - 2\varepsilon) vw I_{111121010}^C, \\
J_{97} &= \varepsilon^4 (1 - 2\varepsilon) v (1 - w) I_{111111020}^C, \\
J_{98} &= \varepsilon^4 (1 - 2\varepsilon) r_1 \left( I_{211111010}^C + I_{121111010}^C + I_{111112010}^C + w I_{111121010}^C - 4\varepsilon I_{111111010}^C \right), \\
J_{99} &= \varepsilon^5 (1 - 2\varepsilon) v I_{111011110}^C, \\
J_{100} &= \varepsilon^4 (1 - 2\varepsilon) v (1 - w) I_{111012110}^C, \\
J_{101} &= \varepsilon^4 (1 - 2\varepsilon) v (1 - w) I_{112011110}^C, \\
J_{102} &= \varepsilon^4 (1 - 2\varepsilon) vr_1 I_{121011110}^C, \\
J_{103} &= \varepsilon^5 (1 - 2\varepsilon) v I_{101111110}^C, \\
J_{104} &= \varepsilon^5 (1 - 2\varepsilon) v (1 - v - w) I_{111111110}^C, \\
J_{105} &= \varepsilon^4 (1 - 2\varepsilon) vw I_{112011110}^D. \tag{27}
\end{aligned}$$

## 4 The differential equation and differential forms

We write the differential equation for the master integrals as

$$dJ = AJ, \tag{28}$$

with

$$A = A_v dv + A_w dw. \tag{29}$$

topology	block	sector	propagators	master integrals
<i>A</i>	1	25	1, 4, 5	$J_1$
	2	27	1, 2, 4, 5	$J_2$
	3	30	2, 3, 4, 5	$J_3, J_4$
	4	85	1, 3, 5, 7	$J_5, J_6$
	5	86	2, 3, 5, 7	$J_7, J_8, J_9, J_{10}$
	6	89	1, 4, 5, 7	$J_{11}$
	7	113	1, 5, 6, 7	$J_{12}, J_{13}, J_{14}$
	8	201	1, 4, 7, 8	$J_{15}$
	9	209	1, 5, 7, 8	$J_{16}$
	10	59	1, 2, 4, 5, 6	$J_{17}$
	11	87	1, 2, 3, 5, 7	$J_{18}, J_{19}$
	12	91	1, 2, 4, 5, 7	$J_{20}$
	13	94	2, 3, 4, 5, 7	$J_{21}, J_{22}, J_{23}$
	14	115	1, 2, 5, 6, 7	$J_{24}, J_{25}, J_{26}$
	15	117	1, 3, 5, 6, 7	$J_{27}, J_{28}, J_{29}, J_{30}$
	16	121	1, 4, 5, 6, 7	$J_{31}$
	17	203	1, 2, 4, 7, 8	$J_{32}$
	18	206	2, 3, 4, 7, 8	$J_{33}, J_{34}$
	19	211	1, 2, 5, 7, 8	$J_{35}$
	20	213	1, 3, 5, 7, 8	$J_{36}$
	21	119	1, 2, 3, 5, 6, 7	$J_{37}, J_{38}$
	22	123	1, 2, 4, 5, 6, 7	$J_{39}$
	23	215	1, 2, 3, 5, 7, 8	$J_{40}, J_{41}, J_{42}$
	24	222	2, 3, 4, 5, 7, 8	$J_{43}, J_{44}$
	25	235	1, 2, 4, 6, 7, 8	$J_{45}$
	26	249	1, 4, 5, 6, 7, 8	$J_{46}$
	27	223	1, 2, 3, 4, 5, 7, 8	$J_{47}, J_{48}, J_{49}$
	28	251	1, 2, 4, 5, 6, 7, 8	$J_{50}$

Table 3: Overview of the set of master integrals. The first column denotes the topology, the second column labels consecutively the sectors, the third column gives the sector id (defined in eq. (8)), the fourth column lists the propagators with positive exponent, the fifth column lists the master integrals in the basis  $\vec{J}$ .

topology	block	sector	propagators	master integrals
<i>B</i>	29	113	1, 5, 6, 7	$J_{51}$
	30	116	3, 5, 6, 7	$J_{52}$
	31	115	1, 2, 5, 6, 7	$J_{53}$
	32	117	1, 3, 5, 6, 7	$J_{54}, J_{55}, J_{56}$
	33	206	2, 3, 4, 7, 8	$J_{57}, J_{58}, J_{59}$
	34	233	1, 4, 6, 7, 8	$J_{60}$
	35	236	3, 4, 6, 7, 8	$J_{61}$
	36	119	1, 2, 3, 5, 6, 7	$J_{62}, J_{63}$
	37	207	1, 2, 3, 4, 7, 8	$J_{64}$
	38	235	1, 2, 4, 6, 7, 8	$J_{65}$
	39	223	1, 2, 3, 4, 5, 7, 8	$J_{66}$
40	239	1, 2, 3, 4, 6, 7, 8	$J_{67}$	
<i>C</i>	41	101	1, 3, 6, 7	$J_{68}, J_{69}, J_{70}$
	42	102	2, 3, 6, 7	$J_{71}$
	43	103	1, 2, 3, 6, 7	$J_{72}$
	44	117	1, 3, 5, 6, 7	$J_{73}, J_{74}, J_{75}$
	45	118	2, 3, 5, 6, 7	$J_{76}, J_{77}, J_{78}$
	46	173	1, 3, 4, 6, 8	$J_{79}, J_{80}$
	47	181	1, 3, 5, 6, 8	$J_{81}$
	48	119	1, 2, 3, 5, 6, 7	$J_{82}, J_{83}, J_{84}$
	49	175	1, 2, 3, 4, 6, 8	$J_{85}$
	50	183	1, 2, 3, 5, 6, 8	$J_{86}$
	51	189	1, 3, 4, 5, 6, 8	$J_{87}, J_{88}$
	52	190	2, 3, 4, 5, 6, 8	$J_{89}$
	53	231	1, 2, 3, 6, 7, 8	$J_{90}, J_{91}, J_{92}$
	54	246	2, 3, 5, 6, 7, 8	$J_{93}, J_{94}$
	55	191	1, 2, 3, 4, 5, 6, 8	$J_{95}, J_{96}, J_{97}, J_{98}$
	56	247	1, 2, 3, 5, 6, 7, 8	$J_{99}, J_{100}, J_{101}, J_{102}$
	57	253	1, 3, 4, 5, 6, 7, 8	$J_{103}$
	58	255	1, 2, 3, 4, 5, 6, 7, 8	$J_{104}$
<i>D</i>	59	239	1, 2, 3, 4, 6, 7, 8	$J_{105}$

Table 4: Overview of the set of master integrals (continued).

The matrix-valued one-form  $A$  satisfies the integrability condition

$$dA - A \wedge A = 0. \quad (30)$$

For the choice of master integrals as in eq. (27), the differential equation is in  $\varepsilon$ -form [45] and we have

$$A = \varepsilon \sum_{k=1}^{19} C_k \omega_k, \quad (31)$$

where the  $C_k$ 's are  $105 \times 105$ -matrices, whose entries are rational numbers. The  $\omega_k$ 's are differential one-forms. We have 19 differential one-forms. All are dlog-forms. The differential one-forms are

$$\begin{aligned} \omega_1 &= d \ln(w), \\ \omega_2 &= d \ln(1-w), \\ \omega_3 &= d \ln(-v), \\ \omega_4 &= d \ln(1-v), \\ \omega_5 &= d \ln(4-v), \\ \omega_6 &= d \ln(w-v), \\ \omega_7 &= d \ln(1-w-v), \\ \omega_8 &= d \ln(1-w+v), \\ \omega_9 &= d \ln(w^2 + v(1-w)), \\ \omega_{10} &= d \ln\left((1-w)^2 + vw\right), \\ \omega_{11} &= d \ln\left((1-w)^2 + v(2-w)\right), \\ \omega_{12} &= d \ln(\lambda(v, w, 1)), \\ \omega_{13} &= d \ln\left(\frac{2-v-r_1}{2-v+r_1}\right), \\ \omega_{14} &= d \ln\left(\frac{2w+v(1-w)-(1-w)r_1}{2w+v(1-w)+(1-w)r_1}\right), \\ \omega_{15} &= d \ln\left(\frac{2(1-w)+vw-wr_1}{2(1-w)+vw+wr_1}\right), \\ \omega_{16} &= d \ln\left(\frac{wv^2 + (1-2w-w^2)v + 2(1-w)^2 - (1-w^2+vw)r_1}{wv^2 + (1-2w-w^2)v + 2(1-w)^2 + (1-w^2+vw)r_1}\right), \\ \omega_{17} &= d \ln\left(\frac{1+w-v-r_2}{1+w-v+r_2}\right), \\ \omega_{18} &= d \ln\left(\frac{(1-w)^2 - v(1+w) - (1-w)r_2}{(1-w)^2 - v(1+w) + (1-w)r_2}\right), \end{aligned}$$

$$\omega_{19} = d \ln \left( \frac{-v(3-v+w) - r_1 r_2}{-v(3-v+w) + r_1 r_2} \right). \quad (32)$$

We write

$$\omega_k = d \ln f_k. \quad (33)$$

The variable transformation from  $(v, w)$  to  $(x, y)$  rationalises all square roots. The  $f_k$ 's are then rational functions in  $x$  and  $y$ . In detail we have

$$\begin{aligned} f_1 &= -p_1^{-1} p_5^{-1} p_6^{-1} p_8 p_9, & f_2 &= 2p_1^{-1} p_2 p_4 p_5^{-1} p_6^{-1} p_7, & f_3 &= p_1^{-1} p_2^2, \\ f_4 &= p_1^{-1} p_{17}, & f_5 &= p_1^{-1} p_3^2, & f_6 &= -p_1^{-1} p_5^{-1} p_6^{-1} p_{18}, \\ f_7 &= p_1^{-1} p_2 p_5^{-1} p_6^{-1} p_{16}, & f_8 &= p_1^{-1} p_2 p_5^{-1} p_6^{-1} p_{12}, & f_9 &= p_1^{-2} p_5^{-2} p_6^{-2} p_{19} p_{20}, \\ f_{10} &= p_1^{-2} p_2^2 p_5^{-2} p_6^{-2} p_{14} p_{15}, & f_{11} &= p_1^{-2} p_2^2 p_5^{-2} p_6^{-2} p_{10} p_{13}, & f_{12} &= p_1^{-2} p_2^2 p_5^{-2} p_6^{-2} p_{11}^2, \\ f_{13} &= p_1^2, & f_{14} &= p_1^{-2} p_{19} p_{20}^{-1}, & f_{15} &= p_1^{-2} p_{14}^{-1} p_{15}, \\ f_{16} &= p_1^{-2} p_{10}^{-1} p_{13} p_{14}^{-1} p_{15}, & f_{17} &= -p_1 p_5^{-1} p_6 p_8 p_9^{-1}, & f_{18} &= -p_1^{-1} p_5^3 p_6^{-3} p_8 p_9^{-1}, \\ f_{19} &= -4p_1 p_4^2 p_7^{-2}. \end{aligned} \quad (34)$$

The  $p_j$ 's are polynomials in  $x$  and  $y$  and given by

$$\begin{aligned} p_1 &= x, \\ p_2 &= x - 1, \\ p_3 &= x + 1, \\ p_4 &= y, \\ p_5 &= y - 1, \\ p_6 &= y + 1, \\ p_7 &= xy + x - y + 1, \\ p_8 &= xy + x - 2y, \\ p_9 &= 2xy - y + 1, \\ p_{10} &= xy^2 + 2xy - 2y^2 + x + 2y, \\ p_{11} &= xy^2 + 2xy + y^2 + x - 2y + 1, \\ p_{12} &= xy^2 + 2xy - y^2 + x + 2y - 1, \\ p_{13} &= 2xy^2 + 2xy - y^2 + 2y - 1, \\ p_{14} &= 2xy^2 + 2xy - 3y^2 + 2y + 1, \\ p_{15} &= 3xy^2 + 2xy - 2y^2 - x + 2y, \\ p_{16} &= 3xy^2 + 2xy - 3y^2 - x + 2y + 1, \\ p_{17} &= x^2 - x + 1, \\ p_{18} &= x^2 y^2 + 2x^2 y - 3xy^2 + x^2 + y^2 - x - 2y + 1, \\ p_{19} &= x^2 y^2 + 2x^2 y - 4xy^2 + x^2 + 2y^2 - 2y, \end{aligned}$$



$$p_{20} = 2x^2y^2 + 2x^2y - 4xy^2 + y^2 - 2y + 1. \quad (35)$$

The polynomials  $p_1$ - $p_{16}$  are linear in  $x$  and appeared previously in the two-loop mixed QCD-electroweak corrections to  $H \rightarrow b\bar{b}$  through a  $Ht\bar{t}$ -coupling [42]. The polynomials  $p_{17}$ - $p_{20}$  are quadratic in  $x$ .

## 5 Boundary values

In order to solve the differential equation we need boundary values. As boundary point we take the point  $(v, w) = (0, 1)$ , which corresponds in  $(x, y)$ -space to the line  $x = 1$ . The map from  $(v, w)$ -space to  $(x, y)$ -space blows up the point  $(v, w) = (0, 1)$ .

Most master integrals vanish at the boundary point. A few boundary values we have to compute explicitly. They are obtained from rather simple integrals. We have

$$J_1 = C_1 w^{-\varepsilon}, \quad J_{15} = C_{15} (-v)^{-\varepsilon}, \quad J_{51} = C_{51} (-v)^{-2\varepsilon} w^{-\varepsilon}, \quad J_{60} = C_{60} (-v)^{-2\varepsilon}, \quad J_{71} = C_{71}, \quad (36)$$

with boundary values

$$\begin{aligned} C_1 &= e^{3\gamma_E \varepsilon} (\Gamma(1 + \varepsilon))^3, \\ C_{15} &= 2e^{3\gamma_E \varepsilon} \frac{(\Gamma(1 + \varepsilon))^3 (\Gamma(1 - \varepsilon))^2}{\Gamma(1 - 2\varepsilon)}, \\ C_{51} &= -3e^{3\gamma_E \varepsilon} \frac{\Gamma(1 + \varepsilon) (\Gamma(1 - \varepsilon))^3 \Gamma(1 + 2\varepsilon)}{\Gamma(1 - 3\varepsilon)}, \\ C_{60} &= 4e^{3\gamma_E \varepsilon} \frac{(\Gamma(1 + \varepsilon))^3 (\Gamma(1 - \varepsilon))^4}{\Gamma(1 - 2\varepsilon)^2}, \\ C_{71} &= e^{3\gamma_E \varepsilon} \frac{\Gamma(1 - \varepsilon) \Gamma(1 + \varepsilon) \Gamma(1 + 2\varepsilon)^2 \Gamma(1 + 3\varepsilon)}{\Gamma(1 + 4\varepsilon)}. \end{aligned} \quad (37)$$

Note that  $J_{15}$ ,  $J_{51}$  and  $J_{60}$  have logarithmic singularities at  $v = 0$ . The remaining non-zero boundary values are

$$\begin{aligned} C_{14} &= C_1, \\ C_{16} &= C_{59} = C_{15}, \\ C_{52} &= C_{51}, \\ C_{61} &= C_{60}, \\ C_6 &= C_8 = C_9 = C_{29} = C_{70} = C_{77} = \frac{1}{3} C_{71}, \\ C_{56} &= -\frac{2}{3} C_{71}, \\ C_{30} &= -\frac{1}{2} C_1 + \frac{1}{2} C_{71}, \end{aligned}$$

$$\begin{aligned}
C_{78} &= \frac{1}{2}C_1 - \frac{1}{2}C_{71}, \\
C_{87} &= -\frac{1}{4}C_1 + \frac{1}{4}C_{71}.
\end{aligned} \tag{38}$$

All other boundary values are zero.

All non-trivial boundary values follow from (i) the explicit computations of eq. 37, (ii) a dedicated integral reduction for  $s = 0$  and  $m_W^2 = m_t^2$  and (iii) the requirement that the master integrals  $J_1$ - $J_{105}$  are constant on the hypersurface  $x = 1$ . Note that this does not require that  $A$  vanishes when restricted to  $x = 1$ , it only requires that the vector  $J$  is in the kernel of  $A$  on the hypersurface  $x = 1$ .

## 6 Analytical results

For the multiple polylogarithms [46–48] we use the notation  $G(z_1, \dots, z_k; y)$ , which is associated with the iterated integral representation. One defines  $G(0, \dots, 0; y)$  with  $k$  zeros to be

$$G(0, \dots, 0; y) = \frac{1}{k!} (\ln y)^k. \tag{39}$$

This includes the trivial case  $G(; y) = 1$ . Multiple polylogarithms are then defined recursively by

$$G(z_1, z_2, \dots, z_k; y) = \int_0^y \frac{dy_1}{y_1 - z_1} G(z_2, \dots, z_k; y_1). \tag{40}$$

A discussion of the basic properties of multiple polylogarithms can be found in ref. [49].

We take the point  $(v, w) = (0, 1)$  in  $(v, w)$ -space as our boundary point. In  $(x, y)$ -space this point corresponds to the hyperplane  $x = 1$ . The master integrals are constant on this hyperplane. We therefore have to integrate only along  $x$  in  $(x, y)$ -space. It is advantageous to change variables from  $x$  to  $x' = 1 - x$ . Doing so, the starting point of the integration will be  $x' = 0$ . The integration gives multiple polylogarithms of the form

$$G(l'_1, \dots, l'_k; x'), \tag{41}$$

where the letters  $l'_1, \dots, l'_k$  are from the alphabet

$$\begin{aligned}
\mathcal{A} = \{ &0, 1, 2, x'_7, x'_8, x'_9, x'_{10}, x'_{11}, x'_{12}, x'_{13}, x'_{14}, x'_{15}, x'_{16}, \\
&x'_{17,a}, x'_{17,b}, x'_{18,a}, x'_{18,b}, x'_{19,a}, x'_{19,b}, x'_{20,a}, x'_{20,b} \}.
\end{aligned} \tag{42}$$

This is an alphabet with 21 letters. The non-trivial letters are given by

$$\begin{aligned}
x'_7 &= \frac{2}{1+y}, & x'_8 &= \frac{1-y}{1+y}, & x'_9 &= \frac{1+y}{2y}, \\
x'_{10} &= \frac{1+4y-y^2}{(1+y)^2}, & x'_{11} &= \frac{2(1+y^2)}{(1+y)^2}, & x'_{12} &= \frac{4y}{(1+y)^2},
\end{aligned}$$

$$\begin{aligned}
x'_{13} &= -\frac{1-4y-y^2}{2y(1+y)}, & x'_{14} &= \frac{1+4y-y^2}{2y(1+y)}, & x'_{15} &= \frac{1-4y-y^2}{(1+y)(1-3y)}, \\
x'_{16} &= -\frac{4y}{(1+y)(1-3y)}
\end{aligned} \tag{43}$$

and

$$\begin{aligned}
x'_{17,a} &= e^{\frac{\pi i}{3}}, & x'_{17,b} &= e^{-\frac{\pi i}{3}}, \\
x'_{18,a} &= \frac{1+4y-y^2+i\sqrt{(3+y^2)(1-5y^2)}}{2(1+y)^2}, & x'_{18,b} &= \frac{1+4y-y^2-i\sqrt{(3+y^2)(1-5y^2)}}{2(1+y)^2}, \\
x'_{19,a} &= \frac{1+2y-y^2+\sqrt{2y(1+y-y^2+y^3)}}{(1+y)^2}, & x'_{19,b} &= \frac{1+2y-y^2-\sqrt{2y(1+y-y^2+y^3)}}{(1+y)^2}, \\
x'_{20,a} &= \frac{2y+i\sqrt{2y(1-y-y^2-y^3)}}{2y(1+y)}, & x'_{20,b} &= \frac{2y-i\sqrt{2y(1-y-y^2-y^3)}}{2y(1+y)}.
\end{aligned} \tag{44}$$

The letters are the roots of the polynomials  $p_1$ - $p_{20}$  in  $x'$  with  $x$  substituted by  $1-x'$ .

For all master integrals we write

$$J_k = \sum_{j=0}^{\infty} \varepsilon^j J_k^{(j)}. \tag{45}$$

The  $\varepsilon^j$ -term  $J_k^{(j)}$  is of uniform weight  $j$ . To give an example, we have

$$\begin{aligned}
J_{67} &= 2 \left[ G(0, x'_8, x'_9; x') + G(0, x'_9, x'_8; x') - G(x'_{15}, x'_8, 0; x') - G(x'_{14}, x'_9, 0; x') \right. \\
&\quad - G(x'_{14}, x'_8, 0; x') - G(x'_{15}, x'_9, 0; x') + G(x'_{14}, x'_8, x'_9; x') + G(x'_{14}, x'_9, x'_8; x') \\
&\quad + G(x'_{15}, x'_8, x'_9; x') + G(x'_{15}, x'_9, x'_8; x') - G(1, x'_9, x'_8; x') + G(x'_{15}, x'_9, 1; x') \\
&\quad - G(x'_7, x'_9, 1; x') + G(x'_7, 1, x'_9; x') - G(0, x'_8, 1; x') - G(1, x'_9, 1; x') \\
&\quad - G(0, 1, x'_8; x') - 2G(x'_7, 1, 0; x') + G(1, 0, 1; x') - G(x'_{14}, x'_8, 1; x') \\
&\quad + 2G(x'_{15}, 1, 0; x') + 2G(x'_7, x'_8, 0; x') + 2G(x'_7, x'_9, 0; x') - G(x'_{15}, x'_7, 1; x') \\
&\quad - G(x'_7, x'_9, x'_8; x') - G(x'_{15}, 1, x'_9; x') - G(x'_7, x'_8, x'_9; x') - G(x'_{14}, x'_7, x'_9; x') \\
&\quad - G(x'_{15}, x'_7, x'_8; x') - G(1, x'_8, x'_9; x') + 2G(x'_{14}, x'_7, 1; x') + G(1, x'_7, 1; x') \\
&\quad \left. + G(1, x'_7, x'_8; x') \right] \varepsilon^3 + O(\varepsilon^4).
\end{aligned} \tag{46}$$

The analytical results for the master integrals are given up to the order  $\varepsilon^5$  in a supplementary electronic file attached to the arxiv version of this article.

## 7 Numerical results

As a reference value we give numerical results for

$$p^2 = m_H^2. \tag{47}$$

Since  $p^2 > 0$ , we are not in the Euclidean region. Feynman’s  $i0$ -prescription instructs us to take a small imaginary part into account:  $p^2 \rightarrow p^2 + i0$ . This selects the correct branches for the two square roots  $\sqrt{-v(4-v)}$  and  $\sqrt{\lambda(v, w, 1)}$ . With

$$m_W = 80.38 \text{ GeV}, \quad m_H = 125.2 \text{ GeV}, \quad m_t = 173.1 \text{ GeV} \quad (48)$$

we obtain for the variables  $x$  and  $y$

$$x = 0.7384 + 0.6743i, \quad y = 0.3987i. \quad (49)$$

The values of the master integrals at this point are given to 8 digits in tables 5 and 6. They are easily computed to arbitrary precision by evaluating the multiple polylogarithms with the help of GiNaC [50, 51]. In addition we verified the correctness of our results at several kinematic points with the help of the programs AMFlow [52–54] and `sector_decomposition` [55, 56].

## 8 Conclusions

In this paper we considered all master integrals contributing to the three-loop Higgs self-energy diagrams of order  $O(\alpha^2 \alpha_s)$  proportional to the product of Yukawa couplings  $y_b y_t$ . These Feynman integrals contain internal top- and  $W$ -propagators. We obtained analytic results for all master integrals, keeping the full dependence on  $p^2$ ,  $m_W^2$  and  $m_t^2$ . We presented a basis of master integrals of uniform weight. In this basis the associated differential equation is  $\epsilon$ -factorised. Two square roots occur, which can be rationalised simultaneously. This allows us to express all master integrals to any order in the dimensional regularisation parameter  $\epsilon$  in terms of multiple polylogarithms. Our results are a building block for precision Higgs physics.

## Acknowledgements

We thank Johann Usovitsch for support with the computer program `Kira`. E.C. receives funding from the European Union’s Horizon 2020 research and innovation programme “High precision multi-jet dynamics at the LHC” (consolidator grant agreement No. 772009). S.W. would like to thank the Munich Institute for Astro-, Particle and BioPhysics (MIAPbP) for hospitality, where this work was finalised.

## A Supplementary material

Attached to the arxiv version of this article is an electronic file in ASCII format with `Maple` syntax, defining the quantities

$$A, \quad J.$$

The matrix  $A$  appears in the differential equation

$$d\vec{J} = A\vec{J}. \quad (50)$$

	$\epsilon^0$	$\epsilon^1$	$\epsilon^2$	$\epsilon^3$	$\epsilon^4$	$\epsilon^5$
$J_1$	1	1.5342081	3.6442984	3.1853184	5.1462966	2.4686377
$J_2$	0	-1.4801083 <i>i</i>	-2.4097495 <i>i</i>	-5.6149488 <i>i</i>	-5.2333001 <i>i</i>	-8.0886313 <i>i</i>
$J_3$	0	1.4801083 <i>i</i>	-0.22727279 <i>i</i>	7.1496293 <i>i</i>	-6.383137 <i>i</i>	25.607456 <i>i</i>
$J_4$	0	0	-0.54768013	-0.36476197	-2.891637	0.046729212
$J_5$	0	1.5342081	-2.9983034	19.075365	-53.393746	195.47467
$J_6$	0.33333333	1.5342081	-0.088235985	15.450021	-32.95737	146.32479
$J_7$	0	-1.4585842 <i>i</i>	1.294182 <i>i</i>	-18.724204 <i>i</i>	33.200239 <i>i</i>	-173.64768 <i>i</i>
$J_8$	0.33333333	0	3.7496407	-3.6797304	30.246157	-60.506131
$J_9$	0.33333333	1.5342081	1.3128219	16.578415	-17.750719	142.4298
$J_{10}$	0	0	-2.2406312	1.0647958	-26.559377	40.135978
$J_{11}$	0	-1.5342081	0.91070302	-7.2586694	10.224944	-28.168258
$J_{12}$	0	1.4585842 <i>i</i>	0.050868134 <i>i</i>	7.6533633 <i>i</i>	-4.6991476 <i>i</i>	27.583113 <i>i</i>
$J_{13}$	0	1.5342081	-1.7502763	7.0599975	-14.845047	30.115459
$J_{14}$	1	1.5342081	5.8849296	4.6861424	17.388435	2.5361954
$J_{15}$	2	1.295828 + 6.2831853 <i>i</i>	-7.8048778 + 4.0709638 <i>i</i>	-12.450557 - 3.8488956 <i>i</i>	-11.660945 - 25.721646 <i>i</i>	10.668736 - 35.695423 <i>i</i>
$J_{16}$	2	4.3642443 + 6.2831853 <i>i</i>	-3.4630132 + 13.710678 <i>i</i>	-21.696071 + 9.7914742 <i>i</i>	-38.706617 - 23.053895 <i>i</i>	-26.131312 - 75.786891 <i>i</i>
$J_{17}$	0	0	2.1907205	3.772359	8.6571337	8.5453519
$J_{18}$	0	-1.4801083 <i>i</i>	0.5445262 <i>i</i>	-15.995422 <i>i</i>	23.86858 <i>i</i>	-132.81753 <i>i</i>
$J_{19}$	0	0	-5.2250698 <i>i</i>	10.295242 <i>i</i>	-65.298537 <i>i</i>	183.26131 <i>i</i>
$J_{20}$	0	0	2.2707942 <i>i</i>	-1.1347528 <i>i</i>	10.629046 <i>i</i>	-14.132023 <i>i</i>
$J_{21}$	0	1.4801083 <i>i</i>	-5.4523426 <i>i</i>	28.891559 <i>i</i>	-88.331183 <i>i</i>	331.78433 <i>i</i>
$J_{22}$	0	-1.5342081	3.2901965	-18.78899	54.646442	-195.09957
$J_{23}$	0	1.5342081	-2.1449256	21.075678	-46.255399	207.64266
$J_{24}$	0	0	2.1588625	0.27796832	11.346261	-5.8909184
$J_{25}$	0	0	-2.2707942 <i>i</i>	2.3774122 <i>i</i>	-10.218327 <i>i</i>	21.004436 <i>i</i>
$J_{26}$	0	-1.4801083 <i>i</i>	-2.4097495 <i>i</i>	-8.9313256 <i>i</i>	-7.7660295 <i>i</i>	-26.43436 <i>i</i>
$J_{27}$	0	-1.4585842 <i>i</i>	1.5911765 <i>i</i>	-16.821522 <i>i</i>	38.929474 <i>i</i>	-158.55603 <i>i</i>
$J_{28}$	0	-1.4801083 <i>i</i>	5.4523426 <i>i</i>	-27.832513 <i>i</i>	90.872416 <i>i</i>	-322.88758 <i>i</i>
$J_{29}$	0.33333333	0	4.2973208	-7.8296127	39.354989	-110.36825
$J_{30}$	0	0	0	3.9327313	-9.7902493	50.090814
$J_{31}$	0	-1.4801083 <i>i</i>	2.8153204 <i>i</i>	-9.1045495 <i>i</i>	18.752518 <i>i</i>	-45.336109 <i>i</i>
$J_{32}$	0	-2.9602166 <i>i</i>	9.2997946 - 2.1958763 <i>i</i>	6.898549 + 11.35639 <i>i</i>	-5.0820522 + 19.501961 <i>i</i>	-38.5719 + 19.050034 <i>i</i>
$J_{33}$	0	2.9602166 <i>i</i>	-9.2997946 - 3.0781682 <i>i</i>	9.6703505 - 3.6126738 <i>i</i>	-19.245548 - 12.824796 <i>i</i>	72.104463 + 0.12405401 <i>i</i>
$J_{34}$	0	0	-1.0953603	0.24128744 - 3.4411758 <i>i</i>	1.6402683 + 0.75802684 <i>i</i>	6.1054746 - 6.1679597 <i>i</i>
$J_{35}$	0	-2.9602166 <i>i</i>	9.2997946 - 6.7374647 <i>i</i>	21.16637 + 4.5035873 <i>i</i>	16.446661 + 32.559044 <i>i</i>	-32.652689 + 60.330416 <i>i</i>
$J_{36}$	0	-1.5342081	-2.6034827 - 9.6397141 <i>i</i>	26.9974 - 0.52358996 <i>i</i>	-27.747558 - 12.210078 <i>i</i>	223.69832 + 60.357017 <i>i</i>
$J_{37}$	0	0	0	0	2.4095606	-7.4050612
$J_{38}$	0	0	0	-0.80345343	2.3774716	-10.59662
$J_{39}$	0	0	2.1907205	-3.9613101	13.09607	-26.512146
$J_{40}$	0	2.9602166 <i>i</i>	-9.2997946 - 3.712675 <i>i</i>	14.988491 - 6.3598329 <i>i</i>	-8.0684303 - 1.80014 <i>i</i>	80.59546 + 18.841397 <i>i</i>
$J_{41}$	0	0	0	-1.2199963 - 1.2715067 <i>i</i>	5.3143264 - 0.86415471 <i>i</i>	-13.01594 - 2.7052663 <i>i</i>
$J_{42}$	0	0	0	0	4.2051087 + 3.3549988 <i>i</i>	-14.946099 + 3.7606676 <i>i</i>
$J_{43}$	0	0	0	0	2.6414875 + 2.2409565 <i>i</i>	-11.227012 + 0.75555211 <i>i</i>
$J_{44}$	0	0	0	-1.2199963 - 1.2715067 <i>i</i>	4.98217 - 0.70082991 <i>i</i>	-11.56359 - 2.4458449 <i>i</i>
$J_{45}$	0	0	4.381441	3.6614725 + 13.764703 <i>i</i>	-16.48045 + 11.502855 <i>i</i>	-30.424976 - 6.4908014 <i>i</i>
$J_{46}$	0	0	0	-1.2199963 - 1.2715067 <i>i</i>	3.4643984 - 1.4351478 <i>i</i>	-2.3605461 + 0.17411836 <i>i</i>
$J_{47}$	0	0	0	0	0	2.7105126 + 2.3055744 <i>i</i>
$J_{48}$	0	0	0	0	0.65371112 + 1.0931833 <i>i</i>	-1.0109578 + 3.0469453 <i>i</i>
$J_{49}$	0	0	0	0	1.6251574 + 0.75070903 <i>i</i>	-1.068945 + 3.6561297 <i>i</i>
$J_{50}$	0	0	0	0	0.94098377 - 0.90286329 <i>i</i>	1.1504284 + 2.4970979 <i>i</i>
$J_{51}$	-3	-3.887484 - 18.849556 <i>i</i>	59.166272 - 24.425783 <i>i</i>	118.51311 + 123.70244 <i>i</i>	-92.180991 + 423.20939 <i>i</i>	-842.91965 + 395.8266 <i>i</i>
$J_{52}$	-3	-8.4901085 - 18.849556 <i>i</i>	49.671371 - 53.344925 <i>i</i>	202.90571 + 64.044212 <i>i</i>	156.24311 + 572.90309 <i>i</i>	-810.36632 + 1171.6505 <i>i</i>
$J_{53}$	0	4.4403248 <i>i</i>	-27.899384 + 6.1707632 <i>i</i>	-38.772048 - 87.008884 <i>i</i>	179.55177 - 183.60241 <i>i</i>	643.38826 + 119.50921 <i>i</i>
$J_{54}$	0	-2.9171683 <i>i</i>	9.1645545 - 1.9918105 <i>i</i>	24.127791 + 5.3212259 <i>i</i>	-21.804448 + 120.95424 <i>i</i>	-196.92433 - 102.13413 <i>i</i>
$J_{55}$	0	-1.5342081	-8.2052687 - 19.279428 <i>i</i>	79.333963 - 36.244239 <i>i</i>	118.23648 + 121.5975 <i>i</i>	181.95398 + 573.21464 <i>i</i>
$J_{56}$	-0.66666667	-1.5342081	-7.3030938	-22.007405 - 14.078301 <i>i</i>	-7.7964625 - 73.29467 <i>i</i>	88.637395 - 153.09033 <i>i</i>
$J_{57}$	0	2.9171683 <i>i</i>	-9.1645545 + 1.9918105 <i>i</i>	-6.2574572 - 3.2092524 <i>i</i>	-20.068012 - 19.446051 <i>i</i>	40.505362 - 29.233598 <i>i</i>
$J_{58}$	0	3.0684163	-1.5124828 + 9.6397141 <i>i</i>	-7.6933829 - 4.7516047 <i>i</i>	-18.562293 + 7.543913 <i>i</i>	-4.8730848 - 73.947317 <i>i</i>
$J_{59}$	2	4.3642443 + 6.2831853 <i>i</i>	-1.2223821 + 13.710678 <i>i</i>	-18.743511 + 16.830625 <i>i</i>	-39.764537 - 13.778153 <i>i</i>	-42.529113 - 55.952566 <i>i</i>
$J_{60}$	4	5.1833121 + 25.132741 <i>i</i>	-78.888363 + 32.56771 <i>i</i>	-129.16811 - 164.93658 <i>i</i>	199.25544 - 383.01328 <i>i</i>	779.6171 - 48.063607 <i>i</i>
$J_{61}$	4	11.320145 + 25.132741 <i>i</i>	-66.228494 + 71.126566 <i>i</i>	-241.69158 - 85.392283 <i>i</i>	-87.715765 - 582.60488 <i>i</i>	887.29829 - 804.39581 <i>i</i>
$J_{62}$	0	0	0	-1.2199963 - 1.2715067 <i>i</i>	5.9914141 - 3.4779304 <i>i</i>	-0.18199463 + 1.5542242 <i>i</i>
$J_{63}$	0	2.9602166 <i>i</i>	-9.2997946 - 3.712675 <i>i</i>	-16.248795 - 0.64764184 <i>i</i>	40.454671 - 155.25155 <i>i</i>	168.63449 + 314.27487 <i>i</i>
$J_{64}$	0	-2.9602166 <i>i</i>	9.2997946 + 3.712675 <i>i</i>	-11.663712 + 4.2951778 <i>i</i>	17.101399 + 9.4430398 <i>i</i>	-68.03826 - 7.8262762 <i>i</i>
$J_{65}$	0	-5.9204331 <i>i</i>	37.199178 - 8.2276842 <i>i</i>	51.696065 + 116.01185 <i>i</i>	-239.40236 + 202.10303 <i>i</i>	-589.55784 - 276.35688 <i>i</i>
$J_{66}$	0	0	0	0	0	2.6632592 + 5.3571394 <i>i</i>
$J_{67}$	0	0	0	1.2199963 + 1.2715067 <i>i</i>	-6.6685018 + 6.0917061 <i>i</i>	-14.752046 - 15.744008 <i>i</i>

Table 5: Numerical results for the first six terms of the  $\epsilon$ -expansion of the master integrals  $J_1$ - $J_{67}$  at the kinematic point  $p^2 = m_H^2$ .

	$\epsilon^0$	$\epsilon^1$	$\epsilon^2$	$\epsilon^3$	$\epsilon^4$	$\epsilon^5$
$J_{68}$	0	-1.4801083i	5.1350892i	-25.837256i	86.938604i	-308.88849i
$J_{69}$	0	0	1.0953603	-3.0231166	16.571142	-51.663494
$J_{70}$	0.33333333	0	1.3701472	1.8283464	-4.959024	38.837723
$J_{71}$	1	0	2.4674011	8.4143983	-35.10786	169.45714
$J_{72}$	0	-1.4801083i	2.498067i	-15.01875i	37.560716i	-132.37051i
$J_{73}$	0	-1.4585842i	1.5911765i	-15.779761i	37.376156i	-152.62396i
$J_{74}$	0	1.5342081	-2.1449256	18.664175	-47.494038	189.76691
$J_{75}$	0	-1.5342081	-0.2345679	-8.9971424	3.8407769	-34.623785
$J_{76}$	0	-1.4585842i	1.294182i	-16.104568i	36.151551i	-153.04501i
$J_{77}$	0.33333333	1.5342081	0.62627943	18.031643	-37.794484	187.99495
$J_{78}$	0	0	0	1.5692999	-6.5217998	25.959061
$J_{79}$	0	2.9602166i	-9.2997946 - 3.0781682i	-14.67018 - 1.6806018i	50.404394 - 150.6488i	194.75027 + 331.14331i
$J_{80}$	0	0	-0.54768013	-0.41446154 - 1.7205879i	-2.3612249 - 7.5128487i	29.563404 - 11.514285i
$J_{81}$	0	1.5342081	-2.9983034	17.247662	-52.439144	184.22811
$J_{82}$	0	1.4801083i	-5.4523426i	23.141893i	-87.484869i	292.60701i
$J_{83}$	0	0	0	-0.80345343	2.4731997	-10.583527
$J_{84}$	0	0	0	0	1.6505684	-6.5723197
$J_{85}$	0	0	0	-0.9800827 - 0.86029394i	4.9546534 - 1.9953368i	-4.239423 + 1.9524939i
$J_{86}$	0	0	-0.31725341i	-1.4970783i	1.498074i	-9.4006924i
$J_{87}$	0	0	-0.13692003	2.3004984 - 0.43014697i	-6.0933617 - 0.63742354i	39.061373 + 3.0381966i
$J_{88}$	0	0	0	-1.2199963 - 1.2715067i	5.1907304 - 3.6242119i	2.1054694 + 0.72653192i
$J_{89}$	0	0	0	-1.3411296 - 1.3187987i	5.076922 - 3.9113363i	2.0464347 - 0.00022367332i
$J_{90}$	0	0	0	-0.9800827 - 0.86029394i	4.8178105 + 0.24314247i	-14.201072 - 1.4422657i
$J_{91}$	0	0	0	0	2.4985099 + 1.7074518i	-12.225556 - 0.48055535i
$J_{92}$	0	1.4801083i	-9.2997946 + 2.0569211i	2.2589826 - 23.245834i	-5.6384268 + 23.974917i	56.355336 - 142.57165i
$J_{93}$	0	0	0	0	2.7500878 + 2.2730554i	-11.219892 + 0.89738075i
$J_{94}$	0	0	0	-1.3411296 - 1.3187987i	4.9222387 - 0.91431241i	-11.726836 - 2.8899212i
$J_{95}$	0	0	0	0	0	1.8404286 + 0.14449436i
$J_{96}$	0	0	0	0	0.21789937 + 0.042131655i	0.50386273 + 0.33655445i
$J_{97}$	0	0	0	-0.23991356 - 0.41121273i	0.85908359 - 1.5044267i	3.8455066 - 0.56878995i
$J_{98}$	0	0	0	0	0.78680569 - 5.3224605i	4.5796027 + 13.058166i
$J_{99}$	0	0	0	0	0	2.8101582 + 2.3220016i
$J_{100}$	0	0	0	0	0.39649658 + 0.58554103i	-0.70924343 + 1.3293236i
$J_{101}$	0	0	0	-0.36104693 - 0.45850473i	0.35197787 - 1.3297874i	0.96759852 - 1.7438646i
$J_{102}$	0	0	0	0	0.96352362 - 0.98327656i	2.3321043 + 1.5761209i
$J_{103}$	0	0	0	0	0	1.9661517 + 5.3108372i
$J_{104}$	0	0	0	0	0	0.022784811 + 0.43623285i
$J_{105}$	0	0	0	0	0.10737705 + 0.15759037i	-0.197588 + 0.3448854i

Table 6: Numerical results for the first six terms of the  $\epsilon$ -expansion of the master integrals  $J_{68}$ - $J_{105}$  at the kinematic point  $p^2 = m_H^2$ .

The entries of the matrix  $A$  are linear combinations of  $\omega_1, \dots, \omega_{19}$ , defined in eq. (32). These differential forms are denoted by

$$\text{omega}_1, \dots, \text{omega}_{19}.$$

The vector  $J$  contains the results for the master integrals up to order  $\epsilon^5$  in terms of multiple polylogarithms. The variable  $\epsilon$  is denoted by `eps`,  $\zeta_2, \zeta_3, \zeta_5$  by

$$\text{zeta}_2, \text{zeta}_3, \text{zeta}_5,$$

respectively. For the notation of multiple polylogarithms we give an example:  $G(x'_7, x'_8, 1; x')$  is denoted by

$$\text{Glog}([\text{xp7}, \text{xp8}, 1], \text{xp}).$$

The file size is 10MB, roughly 90% of the file size is used for the  $\epsilon^5$ -term of  $J$ . The file size increases approximately by a factor of 10 for every additional order. This is expected: With an alphabet of

$$N_{\text{letter}} = 21 \tag{51}$$

letters we have

$$(N_{\text{letter}})^w \tag{52}$$

possible multiple polylogarithms at weight  $w$ . As the matrix  $A$  is sparse, the actual increase factor is of the order  $O(10)$  and roughly a factor of two smaller than  $N_{\text{letter}} = 21$ . Note that the explicit expressions for  $J$  are convenient, but not essential: All information to any order in  $\epsilon$  is stored compactly in the matrix  $A$  and the boundary values given in section 5.

## References

- [1] G. P. Lepage, P. B. Mackenzie, and M. E. Peskin, (2014), arXiv:1404.0319.
- [2] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog, and B. Mistlberger, *Phys. Rev. Lett.* **114**, 212001 (2015), arXiv:1503.06056.
- [3] R. Bonciani *et al.*, *JHEP* **12**, 096 (2016), arXiv:1609.06685.
- [4] B. Mistlberger, *JHEP* **05**, 028 (2018), arXiv:1802.00833.
- [5] D. Kara, *Higher-Order Corrections to Higgs Boson Amplitudes with Full Quark Mass Dependence in Quantum Chromodynamics*, PhD thesis, Zurich U., 2018, arXiv:1908.09932.
- [6] C. Duhr, F. Dulat, and B. Mistlberger, *Phys. Rev. Lett.* **125**, 051804 (2020), arXiv:1904.09990.
- [7] A. H. Ajjath *et al.*, *Phys. Rev. D* **100**, 114016 (2019), arXiv:1906.09028.
- [8] J. Aparisi *et al.*, *Phys. Rev. Lett.* **128**, 122001 (2022), arXiv:2110.10202.
- [9] S. P. Martin, *Phys. Rev. D* **106**, 013007 (2022), arXiv:2203.05042.
- [10] M. Spira, CERN Yellow Reports: Monographs **3**, 123 (2020).

- [11] A. Chakraborty *et al.*, (2022), arXiv:2204.02422.
- [12] A. L. Kataev and V. S. Molokoedov, (2022), arXiv:2201.12073.
- [13] A. L. Kataev and V. T. Kim, *Mod. Phys. Lett. A* **9**, 1309 (1994).
- [14] K. G. Chetyrkin and A. Kwiatkowski, *Nucl. Phys.* **B461**, 3 (1996), arXiv:hep-ph/9505358.
- [15] K. G. Chetyrkin, J. H. Kühn, and M. Steinhauser, *Nucl. Phys.* **B505**, 40 (1997), arXiv:hep-ph/9705254.
- [16] K. G. Chetyrkin and M. Steinhauser, *Phys. Lett.* **B408**, 320 (1997), arXiv:hep-ph/9706462.
- [17] P. A. Baikov, K. G. Chetyrkin, and J. H. Kühn, *Phys. Rev. Lett.* **96**, 012003 (2006), arXiv:hep-ph/0511063.
- [18] W. Bernreuther, L. Chen, and Z.-G. Si, *JHEP* **07**, 159 (2018), arXiv:1805.06658.
- [19] A. Primo, G. Sasso, G. Somogyi, and F. Tramontano, *Phys. Rev.* **D99**, 054013 (2019), arXiv:1812.07811.
- [20] A. L. Kataev, *JETP Lett.* **66**, 327 (1997), arXiv:hep-ph/9708292.
- [21] A. Kwiatkowski and M. Steinhauser, *Phys. Lett.* **B338**, 66 (1994), arXiv:hep-ph/9405308, [Erratum: *Phys. Lett.*B342,455(1995)].
- [22] B. A. Kniehl and M. Spira, *Nucl. Phys.* **B432**, 39 (1994), arXiv:hep-ph/9410319.
- [23] L. Mihaila, B. Schmidt, and M. Steinhauser, *Phys. Lett.* **B751**, 442 (2015), arXiv:1509.02294.
- [24] M. Butenschön, F. Fugel, and B. A. Kniehl, *Phys. Rev. Lett.* **98**, 071602 (2007), arXiv:hep-ph/0612184.
- [25] M. Butenschön, F. Fugel, and B. A. Kniehl, *Nucl. Phys.* **B772**, 25 (2007), arXiv:hep-ph/0702215.
- [26] C. Anastasiou, F. Herzog, and A. Lazopoulos, *JHEP* **03**, 035 (2012), arXiv:1110.2368.
- [27] R. Mondini, M. Schiavi, and C. Williams, *JHEP* **06**, 079 (2019), arXiv:1904.08960.
- [28] R. Mondini, U. Schubert, and C. Williams, *JHEP* **12**, 058 (2020), arXiv:2006.03563.
- [29] A. Behring and W. Bizoń, *JHEP* **01**, 189 (2020), arXiv:1911.11524.
- [30] A. Smirnov, *JHEP* **10**, 107 (2008), arXiv:0807.3243.
- [31] A. V. Smirnov and F. S. Chuharev, *Comput. Phys. Commun.* **247**, 106877 (2020), arXiv:1901.07808.
- [32] R. N. Lee, (2012), arXiv:1212.2685.
- [33] R. N. Lee, *J. Phys. Conf. Ser.* **523**, 012059 (2014), arXiv:1310.1145.
- [34] C. Studerus, *Comput. Phys. Commun.* **181**, 1293 (2010), arXiv:0912.2546.
- [35] A. von Manteuffel and C. Studerus, (2012), arXiv:1201.4330.
- [36] P. Maierhöfer, J. Usovitsch, and P. Uwer, *Comput. Phys. Commun.* **230**, 99 (2018), arXiv:1705.05610.
- [37] J. Klappert, F. Lange, P. Maierhöfer, and J. Usovitsch, *Comput. Phys. Commun.* **266**, 108024 (2021), arXiv:2008.06494.
- [38] T. Peraro, *JHEP* **12**, 030 (2016), arXiv:1608.01902.
- [39] T. Peraro, *JHEP* **07**, 031 (2019), arXiv:1905.08019.
- [40] O. V. Tarasov, *Phys. Rev.* **D54**, 6479 (1996), hep-th/9606018.
- [41] O. V. Tarasov, *Nucl. Phys.* **B502**, 455 (1997), hep-ph/9703319.
- [42] E. Chaubey and S. Weinzierl, *JHEP* **05**, 185 (2019), arXiv:1904.00382.
- [43] M. Besier, D. Van Straten, and S. Weinzierl, *Commun. Num. Theor. Phys.* **13**, 253 (2019), arXiv:1809.10983.
- [44] M. Besier, P. Wasser, and S. Weinzierl, *Comput. Phys. Commun.* **253**, 107197 (2020), arXiv:1910.13251.



- [45] J. M. Henn, Phys. Rev. Lett. **110**, 251601 (2013), arXiv:1304.1806.
- [46] A. B. Goncharov, Math. Res. Lett. **5**, 497 (1998).
- [47] A. B. Goncharov, (2001), math.AG/0103059.
- [48] J. M. Borwein, D. M. Bradley, D. J. Broadhurst, and P. Lisonek, Trans. Amer. Math. Soc. **353:3**, 907 (2001), math.CA/9910045.
- [49] S. Weinzierl, *Feynman Integrals* (Springer, 2022), arXiv:2201.03593.
- [50] C. Bauer, A. Frink, and R. Kreckel, J. Symbolic Computation **33**, 1 (2002), cs.sc/0004015.
- [51] J. Vollinga and S. Weinzierl, Comput. Phys. Commun. **167**, 177 (2005), hep-ph/0410259.
- [52] X. Liu and Y.-Q. Ma, (2022), arXiv:2201.11669.
- [53] X. Liu, Y.-Q. Ma, and C.-Y. Wang, Phys. Lett. B **779**, 353 (2018), arXiv:1711.09572.
- [54] Z.-F. Liu and Y.-Q. Ma, (2022), arXiv:2201.11637.
- [55] C. Bogner and S. Weinzierl, Comput. Phys. Commun. **178**, 596 (2008), arXiv:0709.4092.
- [56] T. Binoth and G. Heinrich, Nucl. Phys. **B585**, 741 (2000), hep-ph/0004013.