Simple restricted modules over the N = 1 Ramond algebra as weak modules for vertex operator superalgebras

Haibo Chen

Dedicated to my supervisor Professor Yucai Su on his 60th birthday

Abstract: In the present paper, a class of new simple modules over the N = 1 Ramond algebra are constructed, which are induced from simple modules over some finite dimensional solvable Lie superalgebras. These new modules are simple restricted modules over the N = 1 Ramond algebra. Combined with the result in [22], a classification of simple weak ψ -twisted $\overline{W}(0, c)$ -modules under certain conditions is also given. At last, some examples of simple restricted N = 1 Ramond modules as various versions of Whittaker modules are presented (classical Whittaker modules were studied in [17]).

Key words: N = 1 Ramond algebra, restricted module, simple module, vertex operator superalgebra.

Mathematics Subject Classification (2020): 17B10, 17B65, 17B68, 17B69.

1 Introduction

The N = 1 Ramond algebra is an infinite dimensional Lie superalgebra

$$\mathcal{R} = \bigoplus_{m \in \mathbb{Z}} \mathbb{C}L_m \oplus \bigoplus_{m \in \mathbb{Z}} \mathbb{C}G_m \oplus \mathbb{C}C,$$

which satisfies the following Lie super-brackets

$$[L_m, L_n] = (n - m)L_{m+n} + \frac{n^3 - n}{12}\delta_{m+n,0}C,$$

$$[G_m, G_n] = -2L_{m+n} + \frac{4m^2 - 1}{12}\delta_{m+n,0}C,$$

$$[L_m, G_n] = (n - \frac{m}{2})G_{m+n}, \ [\mathcal{R}, C] = 0,$$

(1.1)

where $m, n \in \mathbb{Z}$. By definition, we have the following decomposition:

$$\mathcal{R}=\mathcal{R}_{ar{0}}\oplus\mathcal{R}_{ar{1}},$$

where $\mathcal{R}_{\bar{0}} = \operatorname{span}\{L_m, C \mid m \in \mathbb{Z}\}, \mathcal{R}_{\bar{1}} = \operatorname{span}\{G_m \mid m \in \mathbb{Z}\}$. Notice that the even part $\mathcal{R}_{\bar{0}}$ is isomorphic to the classical Virasoro algebra. Clearly, $\mathbb{C}C$ is the center of \mathcal{R} . Let $\mathcal{R}_m = \operatorname{span}\{L_m, G_m, \delta_{m,0}C\}$ for all $m \in \mathbb{Z}$. Then $[\mathcal{R}_m, \mathcal{R}_n] \subset \mathcal{R}_{m+n}$, and $\mathcal{R}, \mathcal{U}(\mathcal{R})$ are \mathbb{Z} -graded. It is easy to see that \mathcal{R} has the following triangular decomposition:

$$\mathcal{R} = \mathcal{R}_+ \oplus \mathcal{R}_0 \oplus \mathcal{R}_-,$$

H. Chen (hypo1025@163.com).

where $\mathcal{R}_+ = \operatorname{span}\{L_m, G_m \mid m \in \mathbb{Z}_+\}$ and $\mathcal{R}_- = \operatorname{span}\{L_m, G_m \mid -m \in \mathbb{Z}_+\}.$

The representation theory of the N = 1 Ramond algebra has attracted a lot of attention from many researchers. The structure of Verma modules over the N = 1 Ramond algebra was investigated by some physicists and mathematicians (see, e.g., [8–10,13]). The structures of Fock modules and pre-Verma modules for the N = 1 Ramond algebra were respectively studied in [14,15]. All simple Harish-Chandra modules over the N = 1 Ramond algebra were classified in [26] (also see [2]). Recently, some non-weight modules as Whittaker modules and $\mathcal{U}(\mathbb{C}L_0 \oplus \mathbb{C}G_0)$ -free modules of rank 1 over the N = 1 Ramond algebra were studied in [17,27], respectively. Moreover, a class of non-weight modules over the N = 1 Ramond algebra were developed in [4], which include super intermediate series modules, $\mathcal{U}(\mathbb{C}L_0)$ -free modules of rank 2 and so on. Clearly, the N = 1 Ramond algebra can be seen as a certain supersymmetric extension of the Virasoro algebra. However, the representation theory of the N = 1 Ramond algebra is far less abundant than the Virasoro algebra.

Highest weight modules and Whittaker modules are two classes of classical representation of Lie (super)algebra, which both belong to the category of restricted modules. In [25], a generalized construction for simple Virasoro modules was given by Mazorchuk and Zhao, which included highest weight modules and various version Whittaker modules. These simple Virasoro modules were restricted modules. In [3, 20], a class of weak modules over the Virasoro vertex operator algebra V(c, 0) were classified by the restricted Virasoro modules of the same level. From then on, to enrich the representation theory of infinite dimensional Lie (super)algebras, the restricted modules over some other Lie (super)algebras such as twisted (mirror) Heisenberg-Virasoro algebras, N = 1 Neveu-Schwarz algebras, gap-p Virasoro algebras, affine Lie algebras $A_1^{(1)}$ were investigated (see, e.g., [1, 5, 6, 12, 18, 19, 21, 24]). In Proposition 4.1 of [22], they showed that any restricted module for the N = 1 Ramond algebra of central charge c is a weak ψ -twisted $\overline{W}(0, c)$ -module. In order to determine those weak modules, we need to give a characterization for restricted modules over the N = 1Ramond algebra. Motivated by this, we consider writing this paper.

The rest of this paper is organized as follows. In Section 2, some definitions and notations of restricted modules are presented for later use. In Section 3, a class of new simple restricted modules over the N = 1 Ramond algebra are constructed in Theorem 3.4. In Section 4, we give a characterization of simple restricted \mathcal{R} -modules under certain conditions in Theorem 4.2, which reduces the problem of classification of simple restricted \mathcal{R} -modules to classification of simple modules over a class of finite-dimensional solvable Lie superalgebras. In Section 5, we see that those induced simple \mathcal{R} -modules defined in Theorem 3.4 are weak ψ -twisted $\overline{W}(0, c)$ -modules. Finally, some examples of restricted \mathcal{R} -modules are given, such as Whittaker modules and high order Whittaker modules.

Throughout this paper, we denote by \mathbb{C} , \mathbb{Z} , \mathbb{N} and \mathbb{Z}_+ the sets of complex numbers, integers, nonnegative integers and positive integers, respectively. All vector superspaces (resp. superalgebras, supermodules) and spaces (resp. algebras, modules) are considered to

be over \mathbb{C} . We use $\mathcal{U}(\mathfrak{a})$ to denote the universal enveloping algebra for a Lie (super)algebra \mathfrak{a} .

2 Preliminaries

Let $M = M_{\bar{0}} \oplus M_{\bar{1}}$ be a \mathbb{Z}_2 -graded vector space. We say that any element $v \in M_{\bar{0}}$ (resp. $v \in M_{\bar{1}}$) is even (resp. odd). Set |v| = 0 if $v \in M_{\bar{0}}$ and |v| = 1 if $v \in M_{\bar{1}}$. We call that all elements in $M_{\bar{0}}$ or $M_{\bar{1}}$ are homogeneous. Throughout this paper, all elements in Lie superalgebras and modules are homogenous. All modules for Lie superalgebras are \mathbb{Z}_2 -graded. All simple modules over Lie (super)algebras are non-trivial unless specified.

Let \mathcal{G} be a Lie superalgebra. A \mathcal{G} -module is a \mathbb{Z}_2 -graded vector space M together with a bilinear map $\mathcal{G} \times M \to M$, denoted $(x, v) \mapsto xv$ satisfying the following conditions

$$x(yv) - (-1)^{|x||y|} y(xv) = [x, y]v, \ \mathcal{G}_{\bar{i}}M_{\bar{j}} \subseteq M_{\bar{i}+\bar{j}}$$

for all $x, y \in \mathcal{G}, v \in M$. It is clear that there is a parity-change functor Π on the category of \mathcal{G} -modules to itself.

Definition 2.1. Assume that M is a \mathcal{G} -module and $x \in \mathcal{G}$.

- (i) If for any $v \in M$ there exists $m \in \mathbb{Z}_+$ such that $x^m v = 0$, we say that the action of x on M is *locally nilpotent*. Similarly, if for any $v \in M$ there exists $m \in \mathbb{Z}_+$ such that $\mathcal{G}^m v = 0$, we say that the action of \mathcal{G} on M is *locally nilpotent*.
- (ii) If for any $v \in M$ we have $\dim(\sum_{m \in \mathbb{Z}_+} \mathbb{C}x^m v) < +\infty$, we call that the $x \in \mathcal{G}$ acts *locally* finite on M. Similarly, if for any $v \in M$ we get $\dim(\sum_{m \in \mathbb{Z}_+} \mathcal{G}^m v) < +\infty$, we call that the \mathcal{G} acts *locally finite* on M.

The action of x on M is locally nilpotent, this implies that the action of x on M is locally finite. Generally speaking, it is not true for any Lie (super)algebra \mathcal{G} . But if \mathcal{G} is a finitely generated Lie (super)algebra, that the action of \mathcal{G} on M is locally nilpotent implies that the action of \mathcal{G} on M is locally finite.

Definition 2.2. Assume that $\mathcal{G} = \bigoplus_{m \in \mathbb{Z}} \mathcal{G}_m$ is a \mathbb{Z} -graded Lie superalgebra. A \mathcal{G} -module M is called the *restricted* module if for any $v \in M$ there exists $k \in \mathbb{N}$ such that $\mathcal{G}_m v = 0$ for m > k.

For simplicity, write $\mathbb{X} = \{0, 1\}$. We denote by \mathcal{S} the set of all infinite vectors of the form $\mathbf{i} := (\dots, i_2, i_1)$ with $i_{2m-1} \in \mathbb{N}, i_{2m} \in \mathbb{X}, m \in \mathbb{Z}_+$, satisfying the condition that the number of nonzero entries is finite. For $k \in \mathbb{Z}_+$, write $\epsilon_k = (\dots, \delta_{k,3}, \delta_{k,2}, \delta_{k,1})$ and $\mathbf{0} = (\dots, 0, 0)$. Denote

$$\mathfrak{W}(\chi) = m \quad \text{if} \quad 0 \neq \chi \in \mathcal{U}(\mathcal{R})_{-m}, \quad \forall m \in \mathbb{N}.$$

For $i \in \mathcal{S}$, set

$$g_{\mathbf{i}} = \cdots \left(G_{-k+1}^{i_{2k}} L_{-k}^{i_{2k-1}} \right) \cdots \left(G_{-2}^{i_{6}} L_{-3}^{i_{5}} \right) \left(G_{-1}^{i_{4}} L_{-2}^{i_{3}} \right) \left(G_{0}^{i_{2}} L_{-1}^{i_{1}} \right) \in \mathcal{U}(\mathcal{R}_{-} \oplus \mathcal{R}_{0}),$$
(2.1)

where $i_{2k-1} \in \mathbb{N}, i_{2k} \in \mathbb{X}, k \in \mathbb{Z}_+$. Then we have

$$\mathfrak{W}(\mathbf{i}) := \mathfrak{W}(g_{\mathbf{i}}) = \sum_{k=1}^{+\infty} k i_{2k-1} + \sum_{k=1}^{+\infty} (k-1) i_{2k}.$$

Denote

$$\mathbf{D}(\mathbf{i}) := \mathbf{D}(g_{\mathbf{i}}) = \sum_{k=1}^{+\infty} (i_{2k-1} + i_{2k}).$$

The following total order on \mathcal{S} can be found in [25].

Definition 2.3. We denote by > the *reverse lexicographical total order* on \mathcal{S} , defined as follows:

- (a) **0** is the minimum element;
- (b) for different nonzero $\mathbf{i}, \mathbf{j} \in \mathcal{S}$, we have

 $\mathbf{j} > \mathbf{i} \iff$ there exists $m \in \mathbb{Z}_+$ such that $(j_k = i_k, \forall 0 < k < m)$ and $j_m > i_m$.

Definition 2.4. By using the above reverse lexicographical total order, we can define the *principal total order* \succ on S: for different $\mathbf{i}, \mathbf{j} \in S$, set $\mathbf{i} \succ \mathbf{j}$ if and only if one of the following condition is satisfied:

- (a) $\mathfrak{W}(\mathbf{i}) > \mathfrak{W}(\mathbf{j});$
- (b) $\mathfrak{W}(\mathbf{i}) = \mathfrak{W}(\mathbf{j})$ and $\mathbf{D}(\mathbf{i}) > \mathbf{D}(\mathbf{j})$;
- (c) $\mathfrak{W}(\mathbf{i}) = \mathfrak{W}(\mathbf{j}), \mathbf{D}(\mathbf{i}) = \mathbf{D}(\mathbf{j}) \text{ and } \mathbf{i} > \mathbf{j}.$

For any simple module V over \mathcal{R} or one of its subalgebra containing the central element C, we denote that the action of C on V is a scalar c. Write $\mathcal{B} := \mathcal{R}_+ \oplus \mathbb{C}L_0 \oplus \mathbb{C}C$. Let V be a simple \mathcal{B} -module. We have the following induced \mathcal{R} -module

$$\operatorname{Ind}_{\mathcal{B},c}(V) = \mathcal{U}(\mathcal{R}) \otimes_{\mathcal{U}(\mathcal{B})} V.$$

By the PBW Theorem (see [7]), and $G^2_{-m} = -L_{-2m}$ for $m \in \mathbb{N}$, every element of $\operatorname{Ind}_{\mathcal{B},c}(V)$ can be uniquely written as the following form

$$\sum_{\mathbf{i}\in\mathcal{S}}g_{\mathbf{i}}v_{\mathbf{i},c}\tag{2.2}$$

where $g_{\mathbf{i}}$ defined as (2.1), $v_{\mathbf{i},c} \in V$ and only finitely many of them are nonzero. For any $w \in \operatorname{Ind}_{\mathcal{B},c}(V)$ as in (2.2), we write $\operatorname{supp}(w)$ the set of all $\mathbf{i} \in \mathcal{S}$ such that $v_{\mathbf{i},c} \neq 0$. For $0 \neq w \in \operatorname{Ind}_{\mathcal{B},c}(V)$, we denote by $\operatorname{deg}(w)$ the maximal element in $\operatorname{supp}(w)$ (with respect to the principal total order on \mathcal{S}), which is called the *degree* of w. Note that $\operatorname{deg}(w)$ is defined only for $w \neq 0$. Let $\mathfrak{W}(w) = \mathfrak{W}(\operatorname{deg}(w))$ and $\mathfrak{W}(0) = -\infty$. For any $m \in \mathbb{N}$, set

$$\operatorname{supp}_m(w) = \{ \mathbf{i} \in \operatorname{supp}(w) \mid \mathfrak{W}(\mathbf{i}) = m \}.$$

3 Construction of simple *R*-modules

For $k \in \mathbb{Z}$, denote $\mathbb{Z}_{\geq k} = \{m \in \mathbb{Z} \mid m \geq k\}$. Now we give a characterization for simple \mathcal{B} -modules.

Lemma 3.1. Let $k, m \in \mathbb{Z}, t \in \mathbb{Z}_{\geq 2}$ and V be a simple \mathcal{B} -module with $L_k V = 0$ for all k > t. Then we get $G_m V = 0$ for all m > t.

Proof. Choose $m \in \mathbb{Z}$ with $m \geq t+1$. According to $G_m^2 V = -L_{2m}V = 0$, we know that $\Upsilon = G_m V$ is a subspace of V and $\Upsilon \neq V$. For any $l \in \mathbb{Z}_+$, one gets

$$G_{m+l}V = \frac{2}{3-m-l}(L_{m+l-1}G_1 - G_1L_{m+l-1})V = 0.$$

Then for any $n \in \mathbb{N}$, we check

$$L_n\Upsilon = L_n G_m V = G_m L_n V + (m - \frac{n}{2})G_{m+n} V \subset \Upsilon.$$

For $k \in \mathbb{Z}_+$, it is clear that $G_k \Upsilon \subset \Upsilon$. Thus, Υ is a submodule of V. It follows from the simplicity of V that we get $\Upsilon = G_m V = 0$ for all m > t.

Lemma 3.2. Let $m, k \in \mathbb{Z}, c \in \mathbb{C}, F_k = L_k$ or G_k . Let V be a simple \mathcal{B} -module and there exists $r \in \mathbb{Z}_{\geq 2}$ such that V satisfying the following two conditions

- (1) the action of L_r on V is injective;
- (2) $L_m V = G_r V = 0$ for all m > r.

Then for any $0 \neq w \in \text{Ind}_{\mathcal{B},c}(V)$ with $\mathfrak{W}(w) = q \in \mathbb{Z}, k \geq r$, we get

(i)
$$\operatorname{supp}_{q-k+r}(F_k w) \subset \{\mathbf{i} - \mathbf{j} \mid \mathbf{i} \in \operatorname{supp}_q(w), \mathfrak{W}(\mathbf{j}) = k - r\};$$

(ii) $\mathfrak{W}(F_k w) \leq q - k + r$.

Proof. (1) By Lemma 3.1 and (2), we have $G_m V = 0$ for all m > r - 1. Now suppose that $w = g_i v_{i,c}$ with $\mathfrak{W}(\mathbf{i}) = q \in \mathbb{Z}$. For k > r and any fixed F_k , by using Lie super-brackets in (1.1), we may transfer the only positive degree term in $[F_k, g_i]$ to the right side, i.e., $[F_k, g_i] \in \sum_{m \in \{q-k, \dots, q\}} \mathcal{U}(\mathcal{R}_- \oplus \mathcal{R}_0)_{-m} \mathcal{R}_{k+m-q}$. So

$$F_k w = [F_k, g_{\mathbf{i}}] v_{\mathbf{i},c} = g_{k-q} v_{\mathbf{i},c} + \sum_{\mathbf{j} \in \{\mathbf{k} | \mathbf{i} - \mathbf{k} \in \mathcal{S}, 0 \le \mathfrak{W}(\mathbf{k}) \le k-1\}} g_{\mathbf{i} - \mathbf{j}} v_{\mathbf{j},c}$$
(3.1)

for some $g_{k-q} \in \mathcal{U}(\mathcal{R})_{k-q}$.

(2) It follows from (1) that we have $\mathfrak{W}(F_k g_i v_{i,c}) \leq \mathfrak{W}(g_i v_{i,c}) - k + r$. The lemma clears. \Box

Lemma 3.3. Let $m \in \mathbb{Z}, c \in \mathbb{C}, \mathbf{i} \in S$ with $\hat{k} = \min\{m \mid i_m \neq 0\} \ge 0$. Assume that V is a simple \mathcal{B} -module and there exists $r \in \mathbb{Z}_{\ge 2}$ such that V satisfying the conditions

(1) the action of L_r on V is injective;

(2)
$$L_m V = G_r V = 0$$
 for all $m > r$.

Then

(b)
$$\mathbf{i} - \epsilon_{\hat{k}} \notin \operatorname{supp}(L_{k+r}g_{\tilde{\mathbf{i}}}v_{\tilde{\mathbf{i}},c})$$
 for all $\mathbf{i} \succ \tilde{\mathbf{i}}$.

(ii) if
$$\widehat{k} = 2k$$
 for some $k \in \mathbb{Z}_+$, we have

(a)
$$\deg(G_{k+r-1}g_{\mathbf{i}}v_{\mathbf{i},c}) = \mathbf{i} - \epsilon_{\widehat{k}};$$

(b) $\mathbf{i} - \epsilon_{\widehat{k}} \notin \operatorname{supp}(G_{k+r-1}g_{\widetilde{\mathbf{i}}}v_{\widetilde{\mathbf{i}},c})$ for all $\mathbf{i} \succ \widetilde{\mathbf{i}}$

Proof. (i) (a) To prove this, we write $L_{k+r}g_iv_{i,c}$ as (3.1). Clearly, the only way to give $g_{\mathbf{i}-\epsilon_{\widehat{k}}}v_{\mathbf{i},c}$ is to commute L_{r+k} with an L_{-k} , which implies $\mathbf{i}-\epsilon_{\widehat{k}} \in \operatorname{supp}(L_{k+r}g_{\mathbf{i}}v_{\mathbf{i},c})$. If there exists a G_{-k} in $g_{\mathbf{i}}$, we obtain $[L_{r+k}, G_{-k}]v_{\mathbf{i},c} = 0$. Then by Lemma 3.2, we conclude $\operatorname{deg}(L_{k+r}g_{\mathbf{i}}v_{\mathbf{i},c}) = \mathbf{i}-\epsilon_{\widehat{k}}$.

(b) Now we consider the following three cases.

First consider $\mathfrak{W}(\mathbf{i}) < \mathfrak{W}(\mathbf{i})$. According to Lemma 3.2, we have

$$\mathfrak{W}(L_{k+r}g_{\widetilde{\mathbf{i}}}v_{\widetilde{\mathbf{i}},c}) \leq \mathfrak{W}(\widetilde{\mathbf{i}}) - k < \mathfrak{W}(\mathbf{i} - \epsilon_{\widehat{k}}) = \mathfrak{W}(\mathbf{i}) - k.$$

Obviously, (b) follows in this case.

Assume $\mathfrak{W}(\widetilde{\mathbf{i}}) = \mathfrak{W}(\mathbf{i}) = p \in \mathbb{Z}$ and $\mathbf{D}(\widetilde{\mathbf{i}}) < \mathbf{D}(\mathbf{i})$. If the element $\mathbf{j} \in \operatorname{supp}(L_{k+r}g_{\widetilde{\mathbf{i}}}v_{\widetilde{\mathbf{i}},c})$ is such that $\mathbf{D}(\mathbf{j}) < \mathbf{D}(\widetilde{\mathbf{i}})$, then

$$\mathbf{D}(\mathbf{j}) < \mathbf{D}(\widetilde{\mathbf{i}}) \le \mathbf{D}(\mathbf{i}) - 1 = \mathbf{D}(\mathbf{i} - \epsilon_{\widehat{k}}).$$

This shows $\mathbf{j} \neq \mathbf{i} - \epsilon_{\hat{k}}$. If the element $\mathbf{j} \in \operatorname{supp}(L_{k+r}g_{\tilde{\mathbf{i}}}v_{\tilde{\mathbf{i}},c})$ is such that $\mathbf{D}(\mathbf{j}) = \mathbf{D}(\tilde{\mathbf{i}})$, then such \mathbf{j} can only be given by commuting L_{k+r} with some L_{-j} , where j > k + r. Then we check

$$\mathfrak{W}(\mathbf{j}) = \mathfrak{W}(\mathbf{\tilde{i}}) - k - r < \mathfrak{W}(\mathbf{\tilde{i}}) - k = \mathfrak{W}(\mathbf{i}) - k = \mathfrak{W}(\mathbf{i} - \epsilon_{\widehat{k}}),$$

which implies that $\mathbf{j} \neq \mathbf{i} - \epsilon_{\hat{k}}$. So, (b) also follows in this case. Let $\tilde{k} = \min\{k \mid \tilde{i}_k \neq 0\}$ be in $\tilde{\mathbf{i}}$. If $\tilde{k} = \hat{k}$, then by (1), $\deg(L_{k+r}g_{\tilde{\mathbf{i}}}v_{\tilde{\mathbf{i}},c}) = \tilde{\mathbf{i}} - \epsilon_{\hat{k}}$, we also have (b) in this case.

Consider the last case $\mathfrak{W}(\mathbf{\tilde{i}}) = \mathfrak{W}(\mathbf{i}) = p$, $\mathbf{D}(\mathbf{\tilde{i}}) = \mathbf{D}(\mathbf{i})$ and $\tilde{k} > \hat{k}$. Then from Lemma 3.2, we have $\mathfrak{W}(L_{k+r}g_{\mathbf{\tilde{i}}}v_{\mathbf{\tilde{i}},c}) . We complete the proof of (i).$

(ii) (a) Write $G_{k-1+r}g_{\mathbf{i}}v_{\mathbf{i},c}$ as the form of (3.1). We know that the only way to obtain $g_{\mathbf{i}-\epsilon_{\widehat{k}}}v_{\mathbf{i},c}$ is to commute G_{k-1+r} with a G_{-k+1} , which gives $\mathbf{i}-\epsilon_{\widehat{k}} \in \operatorname{supp}(G_{k-1+r}g_{\mathbf{i}}v_{\mathbf{i},c})$. Combining this with Lemma 3.2, we deduce $\deg(G_{k-1+r}g_{\mathbf{i}}v_{\mathbf{i},c}) = \mathbf{i}-\epsilon_{\widehat{k}}$.

(b) Now we have the following three cases.

First consider $\mathfrak{W}(\mathbf{i}) < \mathfrak{W}(\mathbf{i})$. It follows from Lemma 3.2 that we get

$$\mathfrak{W}(G_{k-1+r}g_{\widetilde{\mathbf{i}}}v_{\widetilde{\mathbf{i}},c}) \leq \mathfrak{W}(\widetilde{\mathbf{i}}) - k + 1 < \mathfrak{W}(\mathbf{i} - \epsilon_{\widehat{k}}) = \mathfrak{W}(\mathbf{i}) - k + 1.$$

Thus, (b) holds in this case.

Consider $\mathfrak{W}(\widetilde{\mathbf{i}}) = \mathfrak{W}(\mathbf{i}) = p \in \mathbb{Z}$ and $\mathbf{D}(\widetilde{\mathbf{i}}) < \mathbf{D}(\mathbf{i})$. If there exists $\mathbf{j} \in \operatorname{supp}(G_{k-1+r}g_{\widetilde{\mathbf{i}}}v_{\widetilde{\mathbf{i}},c})$ such that $\mathbf{D}(\mathbf{j}) < \mathbf{D}(\widetilde{\mathbf{i}})$, then we have

$$\mathbf{D}(\mathbf{j}) < \mathbf{D}(\mathbf{i}) \le \mathbf{D}(\mathbf{i}) - 1 = \mathbf{D}(\mathbf{i} - \epsilon_{\widehat{k}}).$$

This implies $\mathbf{j} \neq \mathbf{i} - \epsilon_{\hat{k}}$. If there exists $\mathbf{j} \in \operatorname{supp}(G_{k-1+r}g_{\mathbf{i}}v_{\mathbf{i},c})$ such that $\mathbf{D}(\mathbf{j}) = \mathbf{D}(\mathbf{i})$, then such \mathbf{j} can only be given by commuting G_{k-1+r} with some G_{-p} , where p > k - 1 + r. So we obtain

$$\mathfrak{W}(\mathbf{j}) = \mathfrak{W}(\widetilde{\mathbf{i}}) - k + 1 - r < \mathfrak{W}(\widetilde{\mathbf{i}}) - k + 1 = \mathfrak{W}(\mathbf{i}) - k + 1 = \mathfrak{W}(\mathbf{i} - \epsilon_{\widehat{k}}),$$

which gives $\mathbf{j} \neq \mathbf{i} - \epsilon_{\hat{k}}$. Obviously, (ii) also holds in this case. We denote by $\tilde{k} = \min\{k \mid \tilde{i}_k \neq 0\}$ for $\mathbf{\tilde{i}}$. If $\tilde{k} = \hat{k}$, then by (1), $\deg(G_{k-1+r}g_{\mathbf{\tilde{i}}}v_{\mathbf{\tilde{i}},c}) = \mathbf{\tilde{i}} - \epsilon_{\hat{k}}$, we also have (ii) in this case.

At last, we suppose $\mathfrak{W}(\tilde{\mathbf{i}}) = \mathfrak{W}(\mathbf{i}) = p \in \mathbb{Z}$, $\mathbf{D}(\tilde{\mathbf{i}}) = \mathbf{D}(\mathbf{i})$ and $\tilde{k} > \hat{k}$. Then it follows from Lemma 3.2 that we get $\mathfrak{W}(G_{k-1+r}g_{\tilde{\mathbf{i}}}v_{\tilde{\mathbf{i}},c}) . In conclusion, (b) holds.$

Now we present the main result of this section.

Theorem 3.4. Let $m \in \mathbb{Z}, c \in \mathbb{C}$. Let V be a simple \mathcal{B} -module and there exists $r \in \mathbb{Z}_{\geq 2}$ such that V satisfying the following two conditions

- (1) the action of L_r on V is injective;
- (2) $L_m V = G_r V = 0$ for all m > r.

Then we obtain that \mathcal{R} -module $\operatorname{Ind}_{\mathcal{B},c}(V)$ is simple.

Proof. Let $0 \neq w \in \operatorname{Ind}_{\mathcal{B},c}(V)$ and $\deg(w) = \mathbf{i}$ for $\mathbf{i} \in \mathcal{S}$. Write $\hat{k} = \min\{m \mid i_m \neq 0\} \ge 0$. Based on Lemma 3.3, we immediately get the following results. If $\hat{k} = 2k - 1$ for some $k \in \mathbb{Z}_+$, then $L_{r+k}w \neq 0$. If $\hat{k} = 2k$ for some $k \in \mathbb{Z}_+$, then $G_{r+k-1}w \neq 0$. Therefore, from any $0 \neq w \in \operatorname{Ind}_{\mathcal{B},c}(V)$ we always get a nonzero element in $\mathcal{U}(\mathcal{R})w \cap V \neq 0$, which shows the simplicity of $\operatorname{Ind}_{\mathcal{B},c}(V)$. This completes the proof.

4 Characterization of simple restricted *R*-modules

For $m \in \mathbb{Z}, r \in \mathbb{Z}_{\geq 2}$, we denote

$$\mathcal{R}^{(r)} = \bigoplus_{m>r} \mathbb{C}L_m \oplus \bigoplus_{m>r-1} \mathbb{C}G_m.$$

First, several equivalent conditions of simple restricted modules over \mathcal{R} are shown.

Proposition 4.1. Let \mathcal{P} be a simple module for \mathcal{R} . Then the following conditions are equivalent:

- (i) \mathcal{P} is a restricted \mathcal{R} -module.
- (ii) There exists $r \in \mathbb{Z}_{\geq 2}$ such that the actions of L_m, G_n for m > r, n > r 1 on \mathcal{P} are locally nilpotent.
- (iii) There exists $r \in \mathbb{Z}_{\geq 2}$ such that the actions of L_m, G_n for m > r, n > r 1 on \mathcal{P} are locally finite.
- (iv) There exists $r \in \mathbb{Z}_{\geq 2}$ such that \mathcal{P} is a locally nilpotent $\mathcal{R}^{(r)}$ -module.
- (v) There exists $r \in \mathbb{Z}_{\geq 2}$ such that \mathcal{P} is a locally finite $\mathcal{R}^{(r)}$ -module.

Proof. It is clear that $(iv) \Rightarrow (v) \Rightarrow (iii)$ and $(iv) \Rightarrow (ii) \Rightarrow (iii)$. So, we only need to prove $(i) \Rightarrow (iv)$ and $(iii) \Rightarrow (i)$.

(i) \Rightarrow (iv). From the definition of restricted module, for any nonzero element $v \in \mathcal{P}$, there exists $r' \in \mathbb{Z}_+$ such that $L_m v = G_n v = 0$ for m > r', n > r' - 1. By the simplicity of \mathcal{P} , we have $\mathcal{P} = \mathcal{U}(\mathcal{R})v$. Then from the PBW Theorem, we check that \mathcal{P} is a locally nilpotent module over $\mathcal{R}^{(r)}$ for r > r'. (iii) \Rightarrow (i). Since there exists $r \in \mathbb{Z}_{\geq 2}$ such that the actions of L_m and G_n for all m > r, n > r - 1 on \mathcal{P} are locally finite, then we can choose a nonzero element $w \in \mathcal{P}$ such that $L_{r+1}w = \mu w$ for some $\mu \in \mathbb{C}$.

Let $m, n \in \mathbb{Z}, r \in \mathbb{Z}_{\geq 2}$ with m > r, n > r - 1. We denote

$$\mathfrak{M}_L = \sum_{i \in \mathbb{N}} \mathbb{C}L_{r+1}^i L_m w = \mathcal{U}(\mathbb{C}L_{r+1}) L_m w \quad \text{and} \quad \mathfrak{M}_G = \sum_{i \in \mathbb{N}} \mathbb{C}L_{r+1}^i G_n w = \mathcal{U}(\mathbb{C}L_{r+1}) G_n w.$$

It follows from the definition of \mathcal{R} and $i \in \mathbb{N}$ that we get

$$L_{m+i(r+1)}w \in \mathfrak{M}_L \Rightarrow L_{m+(i+1)(r+1)}w \in \mathfrak{M}_L, \quad \forall \ m > r,$$

$$G_{n+i(r+1)}w \in \mathfrak{M}_G \Rightarrow G_{n+(i+1)(r+1)}w \in \mathfrak{M}_G, \quad \forall \ n > r-1$$

Then we have $L_{m+i(r+1)}w \in \mathfrak{M}_L$ and $G_{n+i(r+1)}w \in \mathfrak{M}_G$ for all m > r, n > r-1 by induction on $i \in \mathbb{N}$. In particular, $\sum_{i \in \mathbb{N}} \mathbb{C}L_{m+i(r+1)}w$ and $\sum_{i \in \mathbb{N}} \mathbb{C}G_{n+i(r+1)}w$ are both finite-dimensional for m > r, n > r-1. Thus,

$$\sum_{i\in\mathbb{N}} \mathbb{C}L_{r+1+i}w = \mathbb{C}L_{r+1}w + \sum_{m=r+2}^{2r+2} \left(\sum_{i\in\mathbb{N}} \mathbb{C}L_{m+i(r+1)}w\right),$$
$$\sum_{i\in\mathbb{N}} \mathbb{C}G_{r+i}w = \mathbb{C}G_rw + \sum_{n=r+1}^{2r+1} \left(\sum_{i\in\mathbb{N}} \mathbb{C}G_{n+i(r+1)}w\right)$$

are both finite dimensional. Then we can safely choose $t \in \mathbb{N}$ such that

$$\sum_{i \in \mathbb{N}} \mathbb{C}L_{r+1+i}w = \sum_{i=0}^{t} \mathbb{C}L_{r+1+i}w \quad \text{and} \quad \sum_{i \in \mathbb{N}} \mathbb{C}G_{r+i}w = \sum_{i=0}^{t} \mathbb{C}G_{r+i}w.$$

We denote

$$M' = \sum_{x_1, \dots, x_t \in \mathbb{N}, y_0, y_1, \dots, y_t \in \mathbb{X}} \mathbb{C}L_{r+1}^{x_1} \cdots L_{r+t}^{x_t} G_r^{y_0} G_{r+1}^{y_1} \cdots G_{r+t}^{y_t} w,$$

which is a (finite-dimensional) $\mathcal{R}^{(r)}$ -module by (i).

It follows that we can take a minimal $q \in \mathbb{N}$ such that

$$(L_m + \alpha_1 L_{m+1} + \dots + \alpha_q L_{m+q})M' = 0$$
(4.1)

for some $m \in \mathbb{Z}$ with m > r and $\alpha_i \in \mathbb{C}, i = 1, \ldots, q$. By applying L_m to (4.1), we immediately get

$$(\alpha_1[L_m, L_{m+1}] + \dots + \alpha_q[L_m, L_{m+q}])M' = 0.$$

This implies q = 0, i.e., $L_m M' = 0$ for some m > r. Thus, for any k > m, one checks

$$L_{m+k}M' = \frac{1}{m-k} (L_k L_m - L_m L_k) M' = 0.$$

Choosing any $l > \max\{\frac{m}{2}, r-1\}$, we obtain

$$G_{m+l}M' = \frac{2}{2l-m} (L_m G_l - G_l L_m) M' = 0.$$

Therefore, there exists a nonzero element $u \in \mathcal{P}$ such that $L_k u = G_k u = 0$ for all k > 2m. Note that $\mathcal{P} = \mathcal{U}(\mathcal{R})u$. By the PBW Theorem, we conclude that each element of \mathcal{P} can be written as a linear combinations of vectors

$$\cdots G_{2m-1}^{l_{2m-1}} G_{2m}^{l_{2m}} \cdots L_{2m-1}^{j_{2m-1}} L_{2m}^{j_{2m}} u.$$

Then for any $u' \in \mathcal{P}$, there exists sufficiently large $s \in \mathbb{Z}_+$ such that $L_n u' = G_n u' = 0$ for any n > s, which shows that \mathcal{P} is a restricted \mathcal{R} -module. Then (i) holds. We complete the proof.

Theorem 4.2. Let $c \in \mathbb{C}$ and \mathcal{P} be a simple restricted module for \mathcal{R} . Assume that there exists $k \in \mathbb{Z}_{\geq 2}$ such that the action of L_k on \mathcal{P} is injective.

(1) Then there exists the smallest $b \in \mathbb{Z}_{\geq 2}$ with $b \geq k$ such that

$$\mathfrak{M}_b = \left\{ w \in \mathcal{P} \mid L_m w = G_n w = 0 \quad \text{for all } m > b, n > b - 1 \right\} \neq 0.$$

In particular, $V := \mathfrak{M}_b$ is a \mathcal{B} -module.

- (2) More important, \mathcal{P} can be described by V as follows.
 - (i) If b = k, then V is a simple \mathcal{B} -module, and $\mathcal{P} \cong \operatorname{Ind}_{\mathcal{B},c}(V)$.
 - (ii) If b > k, suppose that L_b acts injectively on V. Then V is a simple \mathcal{B} -module, and $\mathcal{P} \cong \operatorname{Ind}_{\mathcal{B},c}(V)$.

Proof. (1) Note that there exists $k \in \mathbb{Z}_{\geq 2}$ such that the action of L_k on \mathcal{P} is injective. Then from the proof of Proposition 4.1, we see that

$$\mathfrak{M}_{b'} = \left\{ w \in \mathcal{P} \mid L_m w = G_n w = 0 \quad \text{for all } m > b', n > b' - 1 \right\} \neq 0$$

for sufficiently large $b' \ge k$. On the other hand, $\mathfrak{M}_{b'} = 0$ for all b' < k since the action of L_k on \mathcal{P} is injective. Thus we can find the smallest $b \in \mathbb{Z}_{\ge 2}$ with $b \ge k$ such that $V := \mathfrak{M}_b \neq 0$.

It follows from $m > b, n > b - 1, p \in \mathbb{N}, q \in \mathbb{Z}_+$ that we have

$$L_m(L_pw) = (p-m)L_{m+p}w = 0, \ G_n(L_pw) = (\frac{p}{2} - n)G_{n+p}w = 0,$$
$$L_m(G_qw) = (q - \frac{m}{2})G_{m+q}w = 0, \ G_n(G_qw) = -2L_{n+q}w = 0.$$

This shows that $L_p w, G_q w \in V$ for all $p \in \mathbb{N}, q \in \mathbb{Z}_+$. So, V is a \mathcal{B} -module.

(2) First, consider (i). Clearly, the action of L_b on V is injective.

Claim 1. For $s_1, s_2 \in \{1, \dots, b-1\}$ with $s_1 + s_2 = b$, we have $G_{s_1}V \neq 0$ and $G_{s_2}V \neq 0$. Suppose $G_{s_1}V = 0$ or $G_{s_2}V = 0$. For any $b \in \mathbb{Z}_{\geq 2}$, we have

$$L_b V = -\frac{1}{2} (G_{s_1} G_{s_2} + G_{s_2} G_{s_1}) V = 0,$$

which shows V = 0. This contradicts with $V \neq 0$ in (1). The claim holds. Then one can see that the elements in V are not the same as in

$$\left\{ w \in \mathcal{P} \mid L_m w = G_n w = 0 \quad \text{for all } m > b, n > a' \right\} \neq 0 \quad \text{for} \quad a' < b - 1.$$

Since \mathcal{P} is simple and generated by V, we know that there exists a canonical surjective map

$$\Phi: \operatorname{Ind}_{\mathcal{B},c}(V) \to \mathcal{P}, \quad \Phi(1 \otimes v) = v, \quad \forall v \in V.$$

Hence, we only need to show that Φ is an injective map. Let $\mathcal{K} = \ker(\Phi)$. For any $v \in V$ we have $\Phi(1 \otimes v) = v$, then $\mathcal{K} \cap (1 \otimes V) = 0$. If $\mathcal{K} \neq 0$, we can choose $0 \neq w \in \mathcal{K} \setminus (1 \otimes V)$ such that $\deg(w) = \mathbf{i}$ is minimal possible. Observe that \mathcal{K} is an \mathcal{R} -submodule of $\operatorname{Ind}_{\mathcal{B},c}(V)$. Consider that L_b acts injectively on V. Then by using the similar method in Lemma 3.3, a new vector $\eta \in \mathcal{K}$ with $\deg(\eta) \prec \mathbf{i}$ can be obtained. This shows a contradiction, namely, $\mathcal{K} = 0$. Then $\mathcal{P} \cong \operatorname{Ind}_{\mathcal{B},c}(V)$. By the property of induced modules, we see that V is simple as a \mathcal{B} -module.

Using an identical process of (i), we have (ii). This completes the proof.

5 Weak modules for vertex operator superalgebras

The Neveu-Schwarz algebra \mathcal{N} is the Lie superalgebra with a basis $\{L_m, G_p, C \mid m \in \mathbb{Z}, p \in \frac{1}{2} + \mathbb{Z}\}$ and the Lie super-bracket defined by

$$[L_m, L_n] = (n - m)L_{m+n} + \frac{n^3 - n}{12}\delta_{m+n,0}C,$$

$$[G_p, G_q] = -2L_{p+q} + \frac{4p^2 - 1}{12}\delta_{p+q,0}C,$$

$$[L_m, G_p] = (p - \frac{m}{2})G_{m+p}, \ [\mathcal{N}, C] = 0$$

for $m, n \in \mathbb{Z}$, $p, q \in \frac{1}{2} + \mathbb{Z}$. By its definition, we have the following decomposition:

$$\mathcal{N} = \mathcal{N}_{\bar{0}} \oplus \mathcal{N}_{\bar{1}},$$

where $\mathcal{N}_{\bar{0}} = \operatorname{span}\{L_m, C \mid m \in \mathbb{Z}\}, \ \mathcal{N}_{\bar{1}} = \operatorname{span}\{G_p \mid p \in \frac{1}{2} + \mathbb{Z}\}.$ Note that the \mathcal{N} is isomorphic to the subalgebra of \mathcal{R} spanned by $\{L_m \mid m \in 2\mathbb{Z}\} \cup \{G_p \mid p \in 2\mathbb{Z} + 1\} \cup \{C\}.$

It is clear that the \mathcal{N} has a $\frac{1}{2}\mathbb{Z}$ -grading by the eigenvalues of the adjoint action of L_0 . Then \mathcal{N} has the following triangular decomposition:

$$\mathcal{N} = \mathcal{N}_+ \oplus \mathcal{N}_0 \oplus \mathcal{N}_-,$$

where $\mathcal{N}_{+} = \text{span}\{L_{m}, G_{p} \mid m, p > 0\}, \mathcal{N}_{-} = \text{span}\{L_{m}, G_{p} \mid m, p < 0\} \text{ and } \mathcal{N}_{0} = \mathbb{C}\{L_{0}, C\}.$ Set

$$L(z) = \sum_{m \in \mathbb{Z}} L_m z^{-m-2}, \ G(z) = \sum_{n \in \mathbb{Z}} G_{n+\frac{1}{2}} z^{-n-2}.$$
 (5.1)

By Section 4.2 of [21], we have

$$\begin{split} [L(z_1), L(z_2)] &= z_1^{-1} \delta(\frac{z_2}{z_1}) \frac{\partial}{\partial z_2} (L(z_2)) + 2z_1^{-2} \frac{\partial}{\partial z_2} (\delta(\frac{z_2}{z_1})) L(z_2) + \frac{c}{12} z_1^{-4} (\frac{\partial}{\partial z_2})^3 \delta(\frac{z_2}{z_1}), \\ [L(z_1), G(z_2)] &= z_1^{-1} \delta(\frac{z_2}{z_1}) \frac{\partial}{\partial z_2} (G(z_2)) + \frac{3}{2} \left(\frac{\partial}{\partial z_2} z_1^{-1} \delta(\frac{z_2}{z_1})\right) G(z_2), \\ [G(z_1), G(z_2)] &= 2z_1^{-1} \delta(\frac{z_2}{z_1}) L(z_2) + \frac{c}{3} (\frac{\partial}{\partial z_2})^2 z_1^{-1} \delta(\frac{z_2}{z_1}). \end{split}$$

Definition 5.1. A vertex superalgebra denoted by a quadruple $(V, Y, \mathbf{1}, D)$ is a \mathbb{Z}_2 -graded vector space

$$V = V^{(0)} \oplus V^{(1)}.$$

and equipped with a linear map

$$V \longrightarrow (\operatorname{End}(V))[[z, z^{-1}]]$$
$$v \longmapsto Y(v, z) = \sum_{n \in \mathbb{Z}} v_n z^{-n-1} \quad (\text{where } v_n \in \operatorname{End}(V)),$$

and with a specified vector $\mathbf{1} \in V_0$ (the *vacuum vector*) and an endomorphism D of V, such that

(1) For any $u, v \in V$, $u_n v = 0$ for n sufficiently large;

(2)
$$[D, Y(v, z)] = Y(D(v), z) = \frac{\mathrm{d}}{\mathrm{d}z}Y(v, z)$$
 for any $v \in V$;

- (3) $Y(\mathbf{1}, z) = \mathrm{Id}_V$ (the *identity operator* of V);
- (4) $Y(v, z)\mathbf{1} \in \text{End}(V)[[z]]$ and $\lim_{z\to 0} Y(v, z)\mathbf{1} = v$ for any $v \in V$;

(5)
$$z_0^{-1}\delta(\frac{z_1-z_2}{x_0})Y(u,z_1)Y(v,z_2) - (-1)^{|u||v|}z_0^{-1}\delta(\frac{z_2-z_1}{-z_0})Y(v,z_2)Y(u,z_1))$$

= $z_2^{-1}\delta(\frac{z_1-z_0}{z_2})Y(Y(u,z_0)v,z_2)$ (the Jacobi identity), where $|v| = j$ if $v \in V^{(j)}$ for $j \in \mathbb{Z}_2$.

This completes the definition of vertex superalgebra.

A vertex superalgebra V is called a *vertex operator superalgebra* if there exists another distinguished vector ω of V satisfying the following conditions

(6) $[L(m), L(n)] = (n - m)L(m + n) + \frac{n^3 - n}{12}\delta_{m+n,0}C$ for $m, n \in \mathbb{Z}$, where $Y(\omega, z) = \sum_{n \in \mathbb{Z}} L(n)z^{-n-2}$;

(7)
$$L_{-1} = D$$
, i.e., $\frac{d}{dz}Y(v, z) = Y(L_{-1}v, z)$ for any $v \in V$;

(8) V is $\frac{1}{2}\mathbb{Z}$ -graded such that $V = \bigoplus_{n \in \frac{1}{2}\mathbb{Z}} V_{(n)}$, $L(0) \mid_{V_{(n)}} = n \operatorname{Id}_{V_{(n)}}$, $\dim(V_{(n)}) < \infty$ and $V_{(n)} = 0$ for n sufficiently negative.

For any $h, c \in \mathbb{C}$, assume that W(h, c) is the Verma module for \mathcal{N} with highest weight (h, c). Let **1** be a highest weight vector of W(0, c). We denote

$$\bar{W}(0,c) = W(0,c)/\langle G_{-\frac{1}{2}}\mathbf{1}\rangle,\tag{5.2}$$

where $\langle G_{-\frac{1}{2}} \mathbf{1} \rangle$ is the submodule generated by $G_{-\frac{1}{2}} \mathbf{1}$. It is well know that $\overline{W}(0,c)$ has a natural vertex operator superalgebra structure (see [16,21]).

Assume that \mathcal{V} is a vertex superalgebra. Define the following linear map

$$\psi: \mathcal{V} \longrightarrow \mathcal{V}$$
$$a+b \longmapsto a-b$$

for $a \in \mathcal{V}^{(0)}, b \in \mathcal{V}^{(1)}$. It is clear that ψ is an automorphism of \mathcal{V} (called the *canonical* automorphism (see [11])). Then $\operatorname{Aut}(\bar{W}(0,c)) = \mathbb{Z}_2 = \langle \psi \rangle$.

The following results can be found in [21, 22].

Lemma 5.2. Let $c \in \mathbb{C}$, $\overline{W}(0, c)$ defined as (5.2).

- (i) Any weak W
 (0, c)-module is a naturally restricted module for N with central charge c; and conversely, any restricted module for N with central charge c is a weak W
 (0, c)-module;
- (ii) Any weak ψ-twisted W
 (0, c)-module is a naturally restricted module for R with central charge c; and conversely, any restricted module for R with central charge c is a weak ψ-twisted W
 (0, c)-module.

The following result appeared in [18].

Theorem 5.3. Let H be a simple module of $\mathfrak{B} = \mathcal{N}_+ \oplus \mathcal{N}_0$. Assume that the action of C on H is a scalar c. If there exists $t \in \mathbb{N}$ such that H satisfying the following two conditions

(1) the action of L_t on H is injective;

(2) $L_m H = 0$ for all m > t,

then $\operatorname{Ind}_{\mathfrak{B},c}(H)$ is a simple \mathcal{N} -module.

For any $t \in \mathbb{N}$, let \mathfrak{N}_t be the set of simple \mathfrak{B} -modules satisfying the condition in Theorem 5.3. For any $r \in \mathbb{Z}_{\geq 2}$, let \mathfrak{R}_r be the set of simple \mathcal{B} -modules satisfying the condition in Theorem 3.4. Set $\mathfrak{N} = \bigcup_{t \in \mathbb{N}} \mathfrak{N}_t$ and $\mathfrak{R} = \bigcup_{r \in \mathbb{Z}_{\geq 2}} \mathfrak{R}_r$. By [18], Theorem 4.2 and Lemma 5.2, we have the following results.

- **Proposition 5.4.** (1) The set of $\{ \operatorname{Ind}_{\mathfrak{B},c}(H) \mid c \in \mathbb{C}, H \in \mathfrak{N} \}$ gives a complete list of simple weak $\overline{W}(0, c)$ -module under some conditions.
 - (2) The set of $\{ \operatorname{Ind}_{\mathcal{B},c}(V) \mid c \in \mathbb{C}, V \in \mathfrak{R} \}$ gives a complete list of simple weak ψ -twisted $\overline{W}(0,c)$ -modules under some conditions.

6 Examples

In this section, we show some examples of simple restricted \mathcal{R} -modules.

6.1 Simple induced modules

Let $\mathfrak{l} = \mathbb{C}x + \mathbb{C}y$ be the 2-dimensional solvable Lie algebra with basis $\{x, y\}$, which satisfies the non-trivial Lie bracket [x, y] = y. Basically, \mathfrak{l} is a subalgebra of classical 3-dimensional Lie algebra $\mathfrak{sl}(2)$. We construct a class of induced restricted \mathcal{R} -module by using a $\mathbb{C}[y]$ -torsionfree simple \mathfrak{l} -module $\mathfrak{k} = (\partial - 1)^{-1}\mathbb{C}[\partial^{\pm 1}]$ defined in [23, Example 13], whose structure is given by

$$x \cdot f(\partial) = \partial \frac{d}{d\partial} f(\partial) + \frac{f(\partial)}{\partial^2 (\partial - 1)}, \ y \cdot f(\partial) = \partial f(\partial), \ \forall f(\partial) \in \mathfrak{k}.$$

Denote $\widehat{\mathcal{B}} = \bigoplus_{m \ge 0} \mathbb{C}L_m \oplus \bigoplus_{n \ge 2} \mathbb{C}G_n$. Then we can extend \mathfrak{l} -module to a $\widehat{\mathcal{B}}$ -module $\widehat{V}_{\mathfrak{k}} = (\partial - 1)^{-1}\mathbb{C}[\partial^{\pm 1}] \oplus G_1(\partial - 1)^{-1}\mathbb{C}[\partial^{\pm 1}]$ by defining

$$L_{0} \cdot f(\partial) = 2x \cdot f(\partial), \ L_{0} \cdot (G_{1}f(\partial)) = G_{1}(2x \cdot f(\partial) + f(\partial)),$$

$$L_{m} \cdot f(\partial) = G_{n} \cdot f(\partial) = L_{m} \cdot (G_{1}f(\partial)) = G_{n} \cdot (G_{1}f(\partial)) = 0,$$

$$L_{2} \cdot f(\partial) = y \cdot f(\partial), \ L_{2} \cdot (G_{1}f(\partial)) = G_{1}(y \cdot f(\partial)), \ C \cdot f(\partial) = cf(\partial)$$

where $c \in \mathbb{C}, m \in \mathbb{Z}_{\geq 3} \bigcup \{1\}, n \in \mathbb{Z}_{\geq 2}, f(\partial) \in \mathfrak{k}$. Note that $G_1^2 f(\partial) = -y \cdot f(\partial)$. Clearly, $V_{\mathfrak{k}} = U(\mathcal{B}) \otimes_{U(\widehat{\mathcal{B}})} \widehat{V}_{\mathfrak{k}}$ is a simple \mathcal{B} -module. By Theorem 3.4, we get the simple induced \mathcal{R} -modules $\mathrm{Ind}_{\mathcal{B},c}(V_{\mathfrak{k}})$.

6.2 Whittaker modules

For $m, n \in \mathbb{Z}$, we denote

$$\widehat{\mathfrak{b}} = \bigoplus_{m \ge 1} \mathbb{C}L_m \oplus \bigoplus_{n \ge 2} \mathbb{C}G_n$$

Let $\phi : \hat{\mathfrak{b}} \to \mathbb{C}$ be a non-trivial Lie superalgebra homomorphism and $\phi(G_2) = 0$. Then we have $\phi(L_m) = \phi(G_n) = 0$ for m > 2, n > 1. Let $\mathfrak{s}_{\phi} = \mathbb{C}v_{\bar{0}} \oplus \mathbb{C}v_{\bar{1}}$ be a 2-dimensional vector space with

$$xv_{\bar{0}} = \phi(x)v_{\bar{0}}, \ v_{\bar{1}} = G_1v_{\bar{0}}, \ Cv_{\bar{0}} = cv_{\bar{0}}, \ Cv_{\bar{1}} = cv_{\bar{1}}$$

for all $x \in \hat{\mathfrak{b}}$. Clearly, if $\phi(L_2) \neq 0$, \mathfrak{s}_{ϕ} is a simple $\hat{\mathfrak{b}}$ -module and $\dim(\mathfrak{s}_{\phi}) = 2$. Now we consider the induced module

$$M_{\phi} = \mathcal{U}(\widehat{\mathcal{B}}) \otimes_{\mathcal{U}(\widehat{\mathfrak{b}})} \mathfrak{s}_{\phi} = \mathbb{C}[L_0]v_{\bar{0}} \oplus \mathbb{C}[L_0](G_1v_{\bar{0}}).$$

Let $V_{\phi} = \mathcal{U}(\mathcal{B}) \otimes_{\mathcal{U}(\widehat{\mathcal{B}})} M_{\phi}$. It is easy to check that V_{ϕ} is a simple \mathcal{B} -module if $\phi(L_2) \neq 0$. When $\phi(L_2) \neq 0$, the simple induced \mathcal{R} -modules $\operatorname{Ind}_{\mathcal{B},c}(V_{\phi})$ in Theorem 3.4 are so-called classical Whittaker modules (see [17]).

6.3 High order Whittaker modules

In this section, we show a generalization version of Whittaker modules of \mathcal{R} called the high order Whittaker modules.

For $m, n \in \mathbb{Z}$, $s \in \mathbb{Z}_{\geq 2}$, we denote

$$\Gamma(s) = \bigoplus_{m \ge s} \mathbb{C}L_m \oplus \bigoplus_{n \ge s} \mathbb{C}G_n.$$

Let ϕ_s be a Lie superalgebra homomorphism $\phi_s : \Gamma(s) \to \mathbb{C}$ for $s \in \mathbb{Z}_{\geq 2}$. Then we get $\phi_s(L_m) = \phi_s(G_n) = 0$ for m > 2s, n > 2s - 1. Assume that $\mathfrak{s}_{\phi_s} = \mathbb{C}v_{\bar{0}} \oplus \mathbb{C}v_{\bar{1}}$ is a 2-dimensional vector space with

$$xv_{\bar{0}} = \phi(x)v_{\bar{0}}, \ v_{\bar{1}} = G_1v_{\bar{0}}, \ Cv_{\bar{0}} = cv_{\bar{0}}, \ Cv_{\bar{1}} = cv_{\bar{1}}$$

for all $x \in \Gamma(s)$. If $\phi_s(L_{2s}) \neq 0$, \mathfrak{s}_{ϕ_s} is a simple $\Gamma(s)$ -module and $\dim(\mathfrak{s}_{\phi_s}) = 2$. Consider the induced module

$$M_{\phi_s} = \mathcal{U}(\widehat{\mathcal{B}}) \otimes_{\mathcal{U}(\Gamma(s))} \mathfrak{s}_{\phi_s}.$$

Denote $V_{\phi_s} = \mathcal{U}(\mathcal{B}) \otimes_{\mathcal{U}(\widehat{\mathcal{B}})} M_{\phi_s}$. It is clear that V_{ϕ_s} is a simple \mathcal{B} -module if $\phi(L_{2s}) \neq 0$. The corresponding simple \mathcal{R} -modules $\operatorname{Ind}_{\mathcal{B},c}(V_{\phi_s})$ in Theorem 3.4 are exactly the high order Whittaker modules.

Acknowledgements

This work was carried out during the author's visit to University of California, Santa Cruz. The author would like to thank Professor Chongying Dong for the warm hospitality during his visit. This work was supported by the National Natural Science Foundation of China (Grant Nos. 11801369, 12171129), the Overseas Visiting Scholars Program of Shanghai Lixin University of Accounting and Finance (Grant No. 2021161). The author thanked the Professor Hongyan Guo for telling him reference [22].

References

- [1] D. Adamović, R. Lü, K. Zhao, Whittaker modules for the affine algebra $A_1^{(1)}$, Adv. Math., **289** (2016), 438-479.
- [2] Y. Cai, D. Liu, R. Lü, Classification of simple Harish-Chandra modules over the N = 1Ramond algebra, J. Algebra, 567 (2021), 114-127.
- [3] G. Chen, D. Cheng, J. Han, Y. Su, Classification of irreducible weak modules over Virasoro vertex operator algebras, *Internat. J. Math.*, **30** (2019), 1950054.
- [4] H. Chen, X. Dai, D. Liu, From super Weyl modules to the super Virasoro modules, arXiv:2111.10691.
- [5] H. Chen, X. Guo, New simple modules for the Heisenberg-Virasoro algebra, J. Algebra, 390 (2013), 77-86.
- [6] G. Chen, J. Han, Y. Su, Some modules over Lie algebras related to the Virasoro algebra, J. Math. Soc. Japan, 72 (2020), 61-72.
- [7] S. Cheng, W. Wang, Dualities and representations of Lie superalgebras, Graduate Studies in Mathematics, 144. American Mathematical Society, Providence, RI, 2012.
- [8] V. K. Dobrev, Multiplet classification of the indecomposable highest weight modules over the Neveu-Schwarz and Ramond superalgebras, *Lett. Math. Phys.*, **11** (1986), 225-234.
- [9] V. K. Dobrev, Characters of the irreducible highest weight modules over the Virasoro and super-Virasoro algebras. Proceedings of the 14th winter school on abstract analysis (Srní, 1986). Rend. Circ. Mat. Palermo (2) Suppl. No. 14 (1987), 25-42.
- [10] V. K. Dobrev, A Weyl group for the Virasoro and N = 1 super-Virasoro algebras, Comm. Math. Phys., 124 (1989), 501-514.
- [11] A. J. Feingold, I. B. Frenkel, J. F. X. Ries, Spinor construction of vertex operator algebras, triality and $E_8^{(1)}$, Contemporary Math. Vol. 121, Amer. Math. Soc., Providence, 1991.
- [12] D. Gao, Simple restricted modules for the Heisenberg-Virasoro algebra, J. Algebra, 574 (2021), 233-251.
- [13] K. Iohara, Y. Koga, Representation theory of Neveu-Schwarz and Ramond algebras I: Verma modules, Adv. Math., 178 (2003), 1-65.

- [14] K. Iohara, Y. Koga, Representation theory of Neveu-Schwarz and Ramond algebras II: Fock modules, Ann. Inst. Fourier, 53 (2003), 1755-1818.
- [15] K. Iohara, Y. Koga, The structure of pre-Verma modules over the N = 1 Ramond algebra, Lett. Math. Phys., 78 (2006), 89-96.
- [16] V. G. Kac, W. Wang, Vertex operator superalgebras and representations, *Contemp. Math.*, 175 (1994), 161-191.
- [17] D. Liu, Y. Pei, L. Xia, Whittaker modules for the super-Virasoro algebras, J. Algebra Appl., 18 (2019), 1950211.
- [18] D. Liu, Y. Pei, L. Xia, Simple restricted modules for Neveu-Schwarz algebra, J. Algebra, 546 (2020), 341-356.
- [19] D. Liu, Y. Pei, L. Xia, K. Zhao, Irreducible modules over the mirror Heisenberg-Virasoro algebra, *Commun. Contemp. Math.*, (2021), 2150026.
- [20] J. Lepowsky and H. Li, Introduction to vertex operator algebras and their representations, Progress in Mathematics, Vol. 227 (Birkhäuser, Boston, 2004).
- [21] H. Li, Local systems of vertex operators, vertex superalgebras and modules, J. Pure Appl. Algebra, 109 (1996), 143-195.
- [22] H. Li, Local systems of twisted vertex operators, vertex operator superalgebras and twisted modules. Moonshine, the Monster, and related topics (South Hadley, MA, 1994), 203-236, Contemp. Math., 193, Amer. Math. Soc., Providence, RI, 1996.
- [23] R. Lü, V. Mazorchuk, K. Zhao, Classification of simple weight modules over the 1-Spatial ageing algebra, Algebr. Represent. Theory, 18 (2015), 381-395.
- [24] H. Tan, Y. Yao, K. Zhao, Simple restricted modules over the Heisenberg-Virasoro algebra as VOA modules, arXiv:2110.05714.
- [25] V. Mazorchuk, K. Zhao, Simple Virasoro modules which are locally finite over a positive part, Selecta Math. (N.S.), 20 (2014), 839-854.
- [26] Y. Su, Classification of Harish-Chandra modules over the super-Virasoro algebras, Comm. Algebra, 23 (1995), 3653-3675.
- [27] H. Yang, Y. Yao, L. Xia, A family of non-weight modules over the super-Virasoro algebras, J. Algebra, 547 (2020), 538-555.

Haibo Chen

1. School of Statistics and Mathematics, Shanghai Lixin University of Accounting and Finance, Shanghai 201209, China

2. Department of Mathematics, Jimei University, Xiamen, Fujian 361021, China Hypo1025@163.com