EQUIVARIANT GROTHENDIECK RING OF A COMPLETE SYMMETRIC VARIETY OF MINIMAL RANK

V. UMA

ABSTRACT. We describe the G-equivariant Grothendieck ring of a regular compactification X of an adjoint symmetric space G/H of minimal rank. This extends the results of Brion and Joshua for the equivariant Chow ring of wonderful symmetric varieties of minimal rank in [7] and generalizes the results on the regular compactification of an adjoint semisimple group in [17].

1. INTRODUCTION

For a semi-simple group G of adjoint type, we consider a regular compactification X of a symmetric space G/H of minimal rank i.e., $\operatorname{rk}(G/H) = \operatorname{rk}(G) - \operatorname{rk}(H)$. The main examples of symmetric spaces of minimal rank are the groups $G = G \times G/\operatorname{diag}(G)$ and the spaces PGL(2n)/PSp(2n).

We recall here that a normal complete variety X is called an *equivariant compactification* of G/H if X contains G/H as an open subvariety and the action of G on G/H extends to X. We further say that X is a *regular compactification* if X is an equivariant compactification which is regular as a G-variety (see [1, Section 2] and [5, Section 2.1]).

Indeed, the regular compactifications of an adjoint symmetric space are the complete regular embeddings of a symmetric space or the *complete symmetric varieties* considered by Bifet, De Concini and Procesi in [1]. In particular, X considered above is a complete symmetric variety of minimal rank.

The wonderful compactification X^{wond} of a symmetric space G/H constructed by De Concini and Procesi in [8], is the unique regular compactification of G/H with a unique closed orbit. These are called the wonderful symmetric varieties.

Wonderful symmetric varieties have been constructed in characteristic 0 by De Concini and Procesi in [8] and for an arbitrary characteristic by De Concini and Springer [10]. The geometry and topology of these varieties have been widely studied by means of their characterizing wonderful properties (see Section 2). For instance the equivariant cohomology of a wonderful group compactification has been described by Strickland [14]. Also see [12] for the study of the equivariant cohomology ring of a wonderful symmetric variety.

Let $T \subseteq G$ denote a maximal torus containing a maximal torus $T_H \subseteq H$ with Weyl group W_H . Among the wonderful symmetric varieties those of minimal rank (see Definition 2.1) have even better geometric properties. For instance, they have only finitely many *T*-fixed points and finitely many *T*-invariant curves which can be explicitly described (see [7, Lemma 2.1.1]). This enables the study of the cohomology theory of these varieties using a precise form of the localization theorem.

In particular, using the description of the T-fixed points and T-invariant curves, the T-equivariant Chow ring and hence the G-equivariant Chow ring of wonderful symmetric varieties of minimal rank have been described by Brion and Joshua in [7]. The property of finitely many T-fixed points and finitely many Tinvariant curves has also been exploited by Tchoudjem [16] to describe the cohomology groups of line bundles on wonderful varieties of minimal rank.

The complete symmetric varieties have been defined and studied in [1] and [9] where their equivariant cohomology ring structure has been described.

For a regular compactification of a reductive algebraic group (which includes the adjoint semisimple group) the equivariant Chow ring has been described in [5, Section 3] and its equivariant Grothendieck ring has been described in [17].

In this article our main aim is to give a description of the T-equivariant and the G-equivariant Grothendieck ring of algebraic equivariant vector bundles on the complete symmetric variety X of minimal rank.

²⁰²⁰ Mathematics Subject Classification. 14M27, 19E99, 19L99.

Key words and phrases. wonderful compactifications, complete symmetric variety, equivariant K-theory.

With this aim in view in Lemma 2.4 we show that any complete symmetric variety of minimal rank has only finitely many *T*-fixed points and *T*-invariant curves and describe them explicitly. This is an extension of [7, Lemma 2.1.1] for the wonderful symmetric variety of minimal rank. This is also a generalization of the corresponding description of the $T \times T$ -fixed points and the $T \times T$ -invariant curves for a regular compactification of a reductive algebraic group in [5, p. 160] which has been used to describe its $T \times T$ equivariant and the $G \times G$ -equivariant Chow ring in [5] and also to describe its $T \times T$ -equivariant and $G \times G$ -equivariant Grothendieck ring in [17].

Let $K_T(X)$ (resp. $K_G(X)$) denote the *T*-equivariant (resp. *G*-equivariant) Grothendieck ring of *T*equivariant (resp. *G*-equivariant) vector bundles on the complete symmetric variety *X*. Also $R(T) = K_T(pt)$ (resp. $R(G) = K_G(pt)$) denotes the Grothendieck ring of complex representations of *T* (resp. *G*). The structure morphism $X \longrightarrow$ Spec \mathbb{C} induces R(T) (resp. R(G))-module structure on $K_T(X)$ (resp. $K_G(X)$) (see [17, Section 1.2]). In particular, when *X* is the regular compactification of $G = G \times G/\text{diag}(G)$ then we consider $K_{T \times T}(X)$ (resp. $K_{G \times G}(X)$) as $R(T) \otimes R(T)$ (resp. $R(G) \otimes R(G)$)-module. We denote by $K_G(X^{wond})$ the Grothendieck ring of *G*-equivariant vector bundles on the wonderful compactification X^{wond} of G/H.

In Proposition 3.1 and Theorem 3.2, we describe the $K_T(X)$ and $K_G(X)$ using a precise form of localization theorem [17, Theorem 1.3] for X and the description of the T-fixed points and the invariant curves in Lemma 2.4. In particular, we show that $K_G(X) \simeq K_T(Y)^{W_H}$ where Y denotes the smooth complete toric variety which is the closure of T/T_H in X. These results are the K-theoretic analogue of the description of the equivariant Chow ring of wonderful symmetric varieties of minimal rank given in [7, Theorem 2.2.1]. These also generalize the analogous results in [17, Theorem 2.1, Corollary 2.2, Corollary 2.3] for a regular compactification of a complex reductive algebraic group (which includes the case of the regular compactification of an adjoint semisimple group $G = (G \times G)/diag(G)$) to all complete symmetric varieties of minimal rank.

Recall that in [17, Theorem 2.10] we also obtain a direct sum decomposition of $K_{G\times G}(X)$ as a $1 \otimes R(G)$ algebra and describe the multiplication of the graded pieces. In [17, Corollary 2.11] we further show the existance of a canonical multifiltration associated to the direct sum decomposition which in turn gives $K_{G\times G}(X)$ the structure of a $R(T) \otimes R(G)$ -algebra. Moreover, in [17, Corollary 2.12] we gave a geometric interpretation of the pieces occuring in the direct sum decomposition.

Our aim in this paper is to extend this kind of description to any complete symmetric variety of minimal rank.

In Theorem 3.12 we give a direct sum decomposition of $K_G(X)$ as a $1 \otimes R(H)$ -module and also describe the multiplicative structure. This extends [17, Theorem 2.10] where a similar result was proved for the regular group compactifications. We then give a canonical multifiltration for $K_G(X)$ arising out of the direct sum decomposition and show that each of the filtered pieces has a $R(T/T_H) \otimes R(H)$ -module structure (see Corollary 3.13 and Proposition 3.14). In particular, this shows that $K_G(X)$ has a structure of a $R(T/T_H) \otimes R(H)$ -algebra (see Proposition 3.14). This extends the corresponding results for regular group compactifications in [17, Corollary 2.11]. Further, Corollary 3.15 extends [17, Corollary 2.12].

In particular, $R(T/T_H) = R(T/T_H) \otimes 1$ can be identified with the subalgebra of $K_G(X^{wond})$ generated by the isomorphism classes of the G-linearized line bundles corresponding to the boundary divisors and R(H)can be identified with $K_G(G/H)$ which is the G-equivariant Grothendieck ring of the symmetric space G/Hof minimal rank.

The rational equivariant cohomology of complete symmetric varieties have been described by Bifet, De Concini and Procesi [1] in terms of Stanley-Reisner systems. Our results in Section 3 are an integral version of their results via K-theory and localization theorem for the complete symmetric varieties of minimal rank.

Let \widetilde{G} denote the semi-simple simply connected cover of G and \widetilde{T} denote its maximal torus which is a lift of T in \widetilde{G} . Then \widetilde{G} and \widetilde{T} act on X via their canonical projections to G and T respectively.

In Section 3.1 we extend Theorem 3.2 to $K_{\tilde{G}}(X)$ and use the results of [13] to describe the ordinary Grothendieck ring of X.

In Section 3.3 we also give a presentation of $K_G(X)$ as an algebra over $K_G(X^{wond})$ which extends [18, Corollary 4.1].

2. Complete symmetric varieties of minimal rank

2.1. Symmetric spaces of minimal rank. We briefly recall the definition and necessary structure of symmetric spaces of minimal rank. We refer to [6, Section 3] and [7] for details. We shall consider algebraic varieties and algebraic groups over the field of complex numbers.

Let G be a connected reductive algebraic group and let

$$\theta: G \longrightarrow G$$

be an involutive automorphism. Let

$$H := G^{\theta} \subset G$$

denote the subgroup of fixed points. The homogeneous space G/H is called a symmetric space.

It is known that H is reductive and H^0 is nontrivial unless G is a θ -split torus i.e., a torus such that $\theta(g) = g^{-1}$ for every $g \in G$.

Recall that any two maximal θ -split subtori of G are conjugate in H^0 , and their common dimension is called the *rank* of the symmetric space G/H denoted by $\operatorname{rk}(G/H)$. Also every maximal θ -fixed subtorus of G is a maximal torus of H. Any two maximal θ -fixed subtori of G are conjugate in H^0 and their common dimension is the rank of H denoted by $\operatorname{rk}(H)$.

Let T be any θ -stable maximal torus of G.

Let

$$T^{\theta} := \{ x \in T \mid \theta(x) = x \}$$

and

$$T^{-\theta} := \{ x \in T \ | \ \theta(x) = x^{-1} \}$$

Then we have

(2.1)
$$T = T^{\theta} \cdot T^{-\theta} \text{ and } T^{\theta} \cap T^{-\theta} \text{ is finite.}$$

By (2.1) it follows that in general we have

(2.2)
$$\operatorname{rk}(G/H) \ge \operatorname{rk}(G) - \operatorname{rk}(H).$$

If equality holds in (2.2) then we say that the symmetric space G/H is of minimal rank i.e., if rk(G/H) = rk(G) - rk(H). This is equivalent to the condition that $T^{\theta,0}$ is a maximal θ -fixed subtorus of G and to the condition that $A := T^{-\theta,0}$ is a maximal θ -split subtorus of G. This is also equivalent to the condition that any two θ -stable maximal tori of G are conjugate in H^0 (see [6, Section 3.1]).

We choose a θ -stable Borel subgroup B and a θ -stable maximal torus T of B and call (B, T) the standard pair. Also θ acts on the Weyl group W and the root system Φ and stabilizes Φ^+ , Φ^- and Δ which denote respectively the positive roots, negative roots and the simple roots of the root system Φ .

We shall assume from now on that G is semi-simple and adjoint so that Δ is a basis of $X^*(T)$. We call the symmetric space G/H adjoint as well. We shall also assume that G/H is of minimal rank. Since G is semi-simple and adjoint, $T_H := T^{\theta}$ and $B_H := B^{\theta}$ are connected. Further, H is connected, semi-simple and adjoint. Thus T_H is a maximal torus and B_H is a Borel subgroup of H. Moreover, the roots of (H, T_H) are the restrictions to T_H of the roots of (G, T). The Weyl group W_H of (H, T_H) can be identified with W^{θ} (see [6, Lemma 5]).

Recall that the centralizer $C_G(A)$ is a Levi subgroup L of a minimal θ -split parabolic subgroup P of G i.e., $\theta(P)$ is opposite to P and $L = P \cap \theta(P)$.

Let Φ_L denote the root system and $\Delta_L \subseteq \Delta$ denote the subset of simple roots of L. Under the natural action of θ on Φ we have $\Phi_L = \Phi^{\theta}$. Let $p: X^*(T) \longrightarrow X^*(A)$ denote the restriction map from the character group of T to the character group of A. Then $p(\Phi) \setminus \{0\}$ is a root system denoted $\Phi_{G/H}$ called the restricted root system. Also $\Delta_{G/H} := p(\Delta \setminus \Delta_L)$ is a basis of $\Phi_{G/H}$. The simple restricted roots $\Delta_{G/H}$ can further be identified with $\alpha - \theta(\alpha)$ for $\alpha \in \Delta \setminus \Delta_L$ under p.

The restricted Weyl group $W_{G/H} = N_G(A)/C_G(A)$ can further be identified with $N_W(A)/C_W(A)$. This yields the exact sequence

$$1 \longrightarrow W_L \longrightarrow W^{\theta} = W_H \longrightarrow W_{G/H} \longrightarrow 1$$

(see [7, Subsections 1.2 and 1.3]).

Let X^{wond} denote the wonderful compactification of the adjoint symmetric space G/H constructed by De Concini and Procesi [8] and De Concini and Springer [10], which has the following properties:

- V. UMA
- (1) X^{wond} is a nonsingular projective variety.
- (2) G acts on X^{wond} with an open orbit isomorphic to G/H.
- (3) The complement of the open orbit is the union of $r = \operatorname{rk}(G/H)$ nonsingular prime divisors D_1, \ldots, D_r with normal crossings called the *boundary divisors*.
- (4) The G-orbit closures in X^{wond} are exactly the partial intersections $X_I^{wond} := \bigcap_{i \in I} D_i$ where I runs
 - over the subsets of $\{1, \ldots, r\}$. In particular, each X_I^{wond} is smooth.
- (5) The unique closed orbit $D_1 \cap \cdots \cap D_r$ is isomorphic to $G/P \simeq G/\theta(P)$.

Definition 2.1. When G/H is of minimal rank, we call the wonderful compactification X^{wond} a wonderful symmetric variety of minimal rank.

The wonderful compactification of $G \times G/\text{diag}(G)$ is an example of a wonderful symmetric variety of minimal rank. For other examples and the complete classification of wonderful symmetric varieties we refer to [7, Example 1.4.4].

Let X^{wond} be the wonderful symmetric variety of minimal rank. The associated toric variety Y^{wond} is the closure in X^{wond} of T/T_H . Then Y^{wond} is a smooth toric variety associated with the Weyl chambers of the restricted root system $\Phi_{G/H}$. Let Y_0^{wond} denote the open affine toric subvariety associated with the positive Weyl chamber dual to $\Delta_{G/H}$. Then $Y^{wond} = W_{G/H} \cdot Y_0^{wond}$. The distinguished point z of the unique closed orbit G/P is the unique T-fixed point of Y_0^{wond} . Since $W_{G/H}$ acts simply transitively on the T-fixed points, the T-fixed points of Y^{wond} are precisely $w \cdot z$ for $w \in W_H/W_L \simeq W_{G/H}$. Also the T-fixed points of X^{wond} are precisely $w \cdot z$ for $w \in W/W_L$. It is known that any wonderful variety of minimal rank admits finitely many T-stable curves [16, Section 10]. The precise description of the T-stable curves in the wonderful symmetric variety of minimal rank X^{wond} as well as that of the T-stable curves lying in the associated toric variety Y^{wond} is given in [7, Lemma 2.1.1]. We recall this description below.

Lemma 2.2. [7, Lemma 2.1.1]

- (1) The T-fixed points in X^{wond} (resp., Y^{wond}) are exactly the points $w \cdot z$ where $w \in W$ (resp., W_H), and z denotes the unique T-fixed point of Y_0^{wond} . These fixed points are parametrized by W_G/W_L (resp., $W_H/W_L \simeq W_{G/H}$).
- (2) For any $\alpha \in \Phi^+ \setminus \Phi_L^+$, there exists a unique irreducible *T*-stable curve $C_{z,\alpha}$ which contains *z* and on which *T* acts through its character α . The *T*-fixed points in $C_{z,\alpha}$ are exactly *z* and $s_{\alpha} \cdot z$.
- (3) For any $\gamma = \alpha \theta(\alpha) \in \Delta_{G/H}$, there exists a unique irreducible T-stable curve $C_{z,\gamma}$ on which T-acts through its character γ . The T-fixed points in $C_{z,\gamma}$ are exactly z and $s_{\alpha}s_{\theta(\alpha)} \cdot z$.
- (4) The irreducible T-stable curves in X^{wond} are the W-translates of the curves $C_{z,\alpha}$ and $C_{z,\gamma}$. They are all isomorphic to \mathbb{P}^1 .
- (5) The irreducible T-stable curves in Y^{wond} are the $W_{G/H}$ -translates of the curves $C_{z,\gamma}$.

We now recall the definition of a regular G-variety due to Bifet, De Concini and Procesi [1]. (See [5, Section 1.4])

Definition 2.3. A G-variety X is said to be regular if it satisfies the following conditions:

(i) X is smooth and contains a dense G-orbit X_G^0 whose complement is a union of irreducible smooth divisors with normal crossings (the boundary divisors).

(ii) Any G-orbit closure in X is the transversal intersection of the boundary divisors which contain it.

(iii) For any $x \in X$, the normal space $T_x X/T_x(Gx)$ contains a dense orbit of the isotropy group G_x .

Consider the adjoint symmetric space G/H of minimal rank. Then G/H is a homogeneous space under G for the standard action from the left with base point $1 \cdot H$. A normal complete variety X is an equivariant compactification of G/H if X contains G/H as an open subvariety and the action of G on G/H by left multiplication extends to X. We say that X is a regular compactification of G/H if X is an equivariant compactification which is regular as a G-variety. The canonical wonderful compactification X^{wond} of G/H is the unique regular compactification of G/H with a unique closed orbit.

We have a complete classification of the regular compactifications of G/H in terms of smooth torus embeddings Y_0 of T/T_H lying over $Y_0^{wond} \simeq \mathbb{A}^r$ such that the map $Y_0 \longrightarrow Y_0^{wond}$ is proper ([1, Section 10, Definition 23]). Here Y_0 is the T/T_H -toric variety associated to a fan \mathcal{F}_+ which is a smooth subdivision of the positive Weyl chamber \mathcal{C}_+ in the lattice $X_*(T/T_H) \otimes \mathbb{R}$ generated by the coweight vectors dual to the simple restricted roots $\Delta_{G/H}$. Recall that $W_{G/H}$ acts on the coweight lattice $X_*(T/T_H)$ by reflection about the walls of the Weyl chambers. The closure of T/T_H in X is the smooth complete toric variety Y which is associated to the fan \mathcal{F} in $X_*(T/T_H) \otimes \mathbb{R}$ whose cones are $W_{G/H}$ -translates of the cones in \mathcal{F}_+ .

Let $\mathcal{F}_+(r)$ denote the maximal dimensional cones of \mathcal{F}_+ which parametrize the closed *G*-orbits in *X*. For $\sigma \in \mathcal{F}_+(r)$ we denote by $Z_{\sigma} \simeq G/P$ the corresponding closed orbit with base point z_{σ} (see [1, pp. 20-22, Proposition 24]).

We have the following precise description of the T-fixed points and the T-stable curves in the complete symmetric variety of minimal rank X as well as that of the T-fixed points and the T-stable curves lying in the associated toric variety Y.

Lemma 2.4.

- (1) The T-fixed points in X (resp., Y) are exactly the points $w \cdot z_{\sigma}$ where $w \in W$ (resp., W_H), and z_{σ} denotes the T-fixed point of Y_0 corresponding to the maximal cone $\sigma \in \mathcal{F}_+(r)$. These fixed points are parametrized by $\mathcal{F}_+(r) \times W/W_L$ (resp., $\mathcal{F}_+(r) \times W_H/W_L$) where $W_H/W_L \simeq W_{G/H}$.
- (2) For any $\alpha \in \Phi^+ \setminus \Phi_L^+$, there exists a unique irreducible *T*-stable curve $C_{z_{\sigma},\alpha}$ which contains z_{σ} and on which *T* acts through its character α . The *T*-fixed points in $C_{z_{\sigma},\alpha}$ are exactly z_{σ} and $s_{\alpha} \cdot z_{\sigma}$ for every $\sigma \in \mathcal{F}_+(r)$.
- (3) For any $\gamma = \alpha \theta(\alpha) \in \Delta_{G/H}$, there exists a unique irreducible T-stable curve $C_{z_{\sigma},\gamma}$ on which T-acts through its character γ . The T-fixed points in $C_{z_{\sigma},\gamma}$ are exactly z_{σ} and $s_{\alpha}s_{\theta(\alpha)} \cdot z_{\sigma}$. In this case the cone $\sigma \in \mathcal{F}_{+}(r)$ has a facet orthogonal to γ .
- (4) There is a unique irreducible T-stable curve $C_{z_{\sigma},z_{\sigma'}}$ in $Y_0 \subseteq Y \subseteq X$ namely the projective line joining z_{σ} and $z_{\sigma'}$, which are respectively the base points of the distinct closed orbits Z_{σ} and $Z_{\sigma'}$, whenever the cones σ and σ' in $\mathcal{F}_+(r)$ have a common facet.
- (5) The irreducible T-stable curves in X are the W-translates of the curves $C_{z_{\sigma},\alpha}$ for every $\sigma \in \mathcal{F}_{+}(r)$, $C_{z_{\sigma},\gamma}$ whenever $\sigma \in \mathcal{F}_{+}(r)$ has a facet orthogonal to γ and $C_{z_{\sigma},z_{\sigma'}}$ whenever the cones σ and σ' in $\mathcal{F}_{+}(r)$ have a common facet. They are all isomorphic to \mathbb{P}^{1} .
- (6) The irreducible T-stable curves in Y are the $W_{G/H}$ -translates of the curves $C_{z_{\sigma},\gamma}$ whenever $\sigma \in \mathcal{F}_{+}(r)$ has a facet orthogonal to γ and $C_{z_{\sigma},z_{\sigma'}}$ whenever the cones σ and σ' in $\mathcal{F}_{+}(r)$ have a common facet.

Proof: By [1] there exists a *G*-equivariant (hence *T*-equivariant) morphism from $\Phi: X \longrightarrow X^{wond}$ which restricts to the morphism of toric varieties $\varphi: Y \longrightarrow Y^{wond}$ induced by the map of fans $\mathcal{F}_+ \longrightarrow \mathcal{C}_+$.

Let x be a T-fixed point of X. Since Φ is T-equivariant $\Phi(x)$ is a T-fixed point of X^{wond} . Thus $\Phi(x) = w \cdot z$ for some $w \in W$. Here $z \in Y_0^{wond}$ is the base point of the unique closed orbit of X^{wond} . Since Φ is Gequivariant $\Phi(w^{-1} \cdot x) = z$. Now $\Phi^{-1}(Y_0^{wond}) = Y_0$. Thus $w^{-1} \cdot x$ is a T-fixed point of Y_0 and hence is equal to z_{σ} for some $\sigma \in \mathcal{F}_+(r)$. Thus it follows that $x = w \cdot z_{\sigma}$ for some $w \in W$ and $\sigma \in \mathcal{F}_+(r)$. Since W_L acts trivially on Y it follows that the T-fixed points of X are parametrized by $\mathcal{F}_+(r) \times W/W_L$. This proves (1) of Lemma 2.4.

Recall that W_H acts on Y and that the T-fixed point of Y are $w \cdot z_{\sigma}$ for $\sigma \in \mathcal{F}_+(r)$ and $w \in W_H$ is a representative of the coset of W_H/W_L . Thus there are $\mathcal{F}_+(r) \cdot |W_H|/|W_L|$ fixed points in Y.

Consider the translation $v \cdot Y$ where $v \in W$ is a representative of the coset W/W_H . Here $v \cdot Y$ is an irreducible variety isomorphic to Y with an appropriate twist for the T-action. Then the T-fixed points in $v \cdot Y$ are $v \cdot w \cdot z_{\sigma}$ for $\sigma \in \mathcal{F}_+(r)$ and $w \in W_H/W_L$.

Since the *T*-fixed points of *X* are parametrized by $\mathcal{F}_+(r) \times W/W_L$ where $W/W_L = \bigcup_{\overline{v} \in W/W_H} \overline{v} \cdot W_H/W_L$. It follows that the *T*-fixed points of $v \cdot Y$ as v varies over the coset representatives of W/W_H are all distinct and exhaust all the *T*-fixed points of *X*.

In particular, this implies that $v \cdot Y$ as v varies over the coset representatives of W/W_H are disjoint subvarieties of X. For, if they intersect then the intersection being a complete T-variety will contain a T-fixed point which is a contradiction to the above observation that the collection of T-fixed points in $v \cdot Y$ for distinct coset representatives v of W/W_H are distinct.

Thus a T-invariant curve C of X is one the following types.

(i) It lies in some $v \cdot Y$, in which case it is translate by $v \in W$ to a *T*-invariant curve *C'* in *Y*. If *C'* lies in Y_0 then *C'* is of the form (4) in Lemma 2.4. Thus *C* is translate by *v* of a curve of the form (4) in Lemma 2.4. Or else if *C'* does not lie in Y_0 then *C'* is conjugate by $w \in W_H/W_L$ to a *T*-invariant curve in *Y* joining z_{σ} and $w \cdot z_{\sigma}$. Thus $\Phi(C')$ is a *T*-invariant curve of Y^{wond} joining the two distinct *T*-fixed points *z* and $w \cdot z$. Thus $\Phi(C')$ is of the form (3) of Lemma 2.2. Since Φ is *G*-equivariant and hence *T*-equivariant it follows

V. UMA

that C' is of the form (3) of Lemma 2.4. Hence C is a translate by w of a curve of the form (3) of Lemma 2.4

(ii) It is translate by $v' \in W$ of a *T*-invariant curve C' joining a *T*-fixed point of *Y* with a *T*-fixed point of $v \cdot Y$ for some coset representative v of W/W_H different from 1. Since $\Phi(v \cdot Y) = v \cdot Y^{wond}$ for every v the projection $\Phi(C')$ is a *T*-invariant curve in X^{wond} which joins z with $v \cdot z$. Therefore $\Phi(C')$ is of the form (2) of Lemma 2.2. Since Φ is *G*-equivariant and hence *T*-equivariant, C' is of the form (2) in Lemma 2.4. Thus C is a translate by $v' \in W$ of a curve of the form (2) in Lemma 2.4. This proves (5) and (6) of Lemma 2.4. \Box

We denote by $\iota_{\sigma} : K_T(X) \longrightarrow K_T(Z_{\sigma}) \simeq K_T(G/P)$ the restriction map. For $f \in K_T(Z_{\sigma})$ we denote by f_w the restriction of f to the fixed point $w \cdot z_{\sigma}$ for $w \in W/W_L$.

3. The *G*-equivariant Grothendieck ring of X

Let G/H be a symmetric space of minimal rank. Let X^{wond} denote the canonical wonderful symmetric variety of minimal rank. Let X be a projective regular compactification of G/H which in addition is characterized by the fact that the map of T/T_H -toric varieties $Y_0 \longrightarrow Y_0^{wond} \simeq \mathbb{A}^r$ is a projective morphism.

The following proposition describes the T-equivariant Grothendieck ring of X. Also see [17, Theorem 2.1] for the corresponding result for the case of regular group compactifications.

Proposition 3.1. The map

$$\prod_{r \in \mathcal{F}_+(r)} \iota_{\sigma} : K_T(X) \longrightarrow \prod_{\sigma \in \mathcal{F}_+(r)} K_T(G/P) \hookrightarrow R(T)^{|W/W_L| \cdot |\mathcal{F}_+(r)|}$$

is injective and its image consists of all families $(f_{\sigma})_{\sigma \in \mathcal{F}_+(r)}$ in $K_T(G/P)$ such that:

(i) $f_{\sigma,w} \equiv f_{\sigma,w \cdot s_{\alpha} s_{\theta(\alpha)}} \pmod{1 - e^{-w(\gamma)}}$ whenever $\gamma = \alpha - \theta(\alpha) \in \Delta_{G/H}$ and $w \in W$.

(ii) $f_{\sigma,w} \equiv f_{\sigma',w} \pmod{1 - e^{-w(\chi)}}$ whenever $\chi \in X^*(T/T_H)$ and the cones σ and σ' of $\mathcal{F}_+(r)$ have a common facet orthogonal to χ and $w \in W$.

Proof: By [17, Theorem 1.3], the image of $K_T(X) \longrightarrow K_T(X^T)$ is defined by linear congruences $f_x \equiv f_y \pmod{1 - e^{-\chi}}$ whenever $x, y \in X^T$ are connected by a curve where T acts by the character χ .

Recall that the union of the closed orbits $\bigcup_{\sigma \in \mathcal{F}_+(r)} Z_{\sigma}$ contains all the *T*-fixed points of *X* (see [1], [6,

Lemma 3.4] [16, Proposition 5.1]). Moreover, *T*-fixed points in Z_{σ} are the *W*-translates of z_{σ} . Since W_L fixes z_{σ} for every $\sigma \in \mathcal{F}_+(r)$ they are parametrized by W/W_L . In particular, z_{σ} is the distinguished point in Z_{σ} which is fixed by *P*. We therefore have the inclusions $\prod_{\sigma \in \mathcal{F}_+(r)} \iota_{\sigma} : K_T(X) \hookrightarrow \prod_{\sigma \in \mathcal{F}_+(r)} K_T(Z_{\sigma}) =$

 $K_T(G/P) \hookrightarrow (R(T))^{|W/W_L| \cdot |\mathcal{F}_+(r)|}.$

By Lemma 2.4(5), the invariant curves in X are the W-translates of curves of type (2), (3) or (4). Now, the W-translates of curves of type (2) which are $C_{z_{\sigma},\alpha}$ for $\alpha \in \Phi^+ \setminus \Phi_L^+$ lie inside the closed orbit Z_{σ} for $\sigma \in \mathcal{F}_+(r)$ and define its T-equivariant K-ring inside $(R(T))^{|W/W_L|}$. Thus the image of $K_T(X)$ under $\prod_{\sigma \in \mathcal{F}_+(r)} \iota_{\sigma}$ consists of $(f_{\sigma}) \in \prod_{\sigma \in \mathcal{F}_+(r)} K_T(G/P)$ satisfying the congruences (i) and (ii) corresponding

respectively to the W-translates of the curves of type (3) and (4). Note that T acts by $w(\gamma)$ (resp. $w(\chi)$) on the translation by $w \in W$ of the curve $C_{z_{\sigma},\gamma}$ (resp. $C_{z_{\sigma},z_{\sigma'}}$). \Box

The G-equivariant Chow ring of X^{wond} was described in [7, Theorem 2.2.1] using the description of the T-fixed points and the T-stable curves given in Lemma 2.2.

We state below the corresponding theorem for the *G*-equivariant Grothendieck ring $K_G(X)$ where X is a regular compactification of a symmetric space G/H of minimal rank. The proof follows along similar lines as that of the equivariant Chow ring of X^{wond} by using [17, Theorem 1.3] in place of [4, Section 3.4] and by the description of the *T*-fixed points and *T*-stable curves of X given in Lemma 2.4. It further generalizes [17, Corollary 2.2, Corollary 2.3] to any complete symmetric variety of minimal rank.

Theorem 3.2.

(1) The ring $K_G(X)$ consists in all families $(f_{\sigma})_{\sigma \in \mathcal{F}_+(r)}$ of elements in $R(T)^{W_L}$ satisfying

- (i) $s_{\alpha}s_{\theta(\alpha)} \cdot f_{\sigma} \equiv f_{\sigma} \pmod{1 e^{-\gamma}}$ whenever $\gamma = \alpha \theta(\alpha) \in \Delta_{G/H}$.
- (ii) $f_{\sigma} \equiv f_{\sigma'} \pmod{1 e^{-\chi}}$ whenever $\chi \in X^*(T/T_H)$ and the cones σ and σ' of $\mathcal{F}_+(r)$ have a common facet orthogonal to χ .
- (2) The map

$$r: K_G(X) \longrightarrow K_T(X)^W \longrightarrow K_T(X)^{W_H} \longrightarrow K_T(Y)^{W_H}$$

obtained by composing the canonical maps is an isomorphism of R(G)-algebras where $R(G) = R(T)^W$.

Proof: (1) Note that by [17, Theorem 1.8] the restriction homomorphism $K_G(X) \longrightarrow K_T(X)$ induces an isomorphism $K_G(X) \simeq K_T(X)^W$. Again by [17, Theorem 1.8] the ring $K_G(Z_{\sigma}) = K_G(G/P)$ is isomorphic to $K_T(G/P)^W$. It is further isomorphic to $R(P) = R(T)^{W_L}$ via restriction to z_{σ} .

Moreover, if $f \in K_T(Z_\sigma)^W$, then for each $w \in W$ we have

where $f_{\sigma,1}$ denotes the restriction of f to z_{σ} and $f_{\sigma,w}$ denotes the restriction of f to $w \cdot z_{\sigma}$. Furthermore, since W_L acts trivially on the *T*-fixed points (3.3) implies that $w \cdot f_{\sigma,1} = f_{\sigma,1}$ for $w \in W_L$.

Recall that T acts by $w(\gamma)$ (resp. $w(\chi)$) on the translation of $C_{z,\gamma}$ (resp. $C_{z_{\sigma},z_{\sigma'}}$) by $w \in W$. Thus for $(f_{\sigma}) \in \prod K_T(G/P)^W$ the congruences (i) and (ii) can be rewritten as

$$\sigma \in \mathcal{F}_{\perp}(r)$$

(i') $w \cdot f_{\sigma,1} \equiv w \cdot s_{\alpha} s_{\theta(\alpha)} \cdot f_{\sigma,1} \pmod{1 - e^{-w(\gamma)}}$ whenever $\gamma = \alpha - \theta(\alpha) \in \Delta_{G/H}$ and $w \in W$.

(ii') $w \cdot f_{\sigma,1} \equiv w \cdot f_{\sigma',1} \pmod{1 - e^{-w(\chi)}}$ whenever $\chi \in X^*(T/T_H)$ and the cones σ and σ' of $\mathcal{F}_+(r)$ have a common facet orthogonal to χ and $w \in W$.

Moreover, since (i') and (ii') are consequences of the congruences (i) and (ii), this proves (1).

(2) Recall that T-fixed points of Y are the W_H translates of z_{σ} for $\sigma \in \mathcal{F}_+(r)$. Also W_L acts trivially on the T-fixed points. Furthermore, by Lemma 2.4 (6) and by observing that T acts by $w(\gamma)$ (resp. $w(\chi)$) on the translation of the curve $C_{z_{\sigma},\gamma}$ (resp. $C_{z_{\sigma},z_{\sigma'}}$) by $w \in W_H$, it can be seen that $K_T(Y)$ can be identified with the the tuples $(f_{\sigma,w})_{w \in W_H/W_L, \sigma \in \mathcal{F}_+(r)}$ in $R(T)^{|W_H/W_L| \cdot |\mathcal{F}_+(r)|}$ satisfying the congruences

(i") $f_{\sigma,w} \equiv f_{\sigma,w \cdot s_{\alpha}s_{\theta(\alpha)}} \pmod{1 - e^{-w(\gamma)}}$ whenever $\gamma = \alpha - \theta(\alpha) \in \Delta_{G/H}$ and $w \in W_H$.

(ii") $f_{\sigma,w} \equiv f_{\sigma',w} \pmod{1 - e^{-w(\chi)}}$ whenever $\chi \in X^*(T/T_H)$, the cones σ and σ' of $\mathcal{F}_+(r)$ have a common facet orthogonal to χ and $w \in W_H$.

Here $f_{\sigma,w}$ denotes the restriction of $f \in K_T(Y)$ to $w \cdot z_{\sigma}$ for $\sigma \in \mathcal{F}_+(r)$ and $w \in W_H/W_L$. Furthermore, if $f \in K_T(Y)^{W_H}$ then $f_{\sigma,w} = w \cdot f_{\sigma}$ where f_{σ} denotes the restriction of f to z_{σ} . Thus it follows that the ring $K_T(Y)^{W_H}$ is identified with the tuples $(f_{\sigma})_{\sigma \in \mathcal{F}_+(r)} \in \prod_{\sigma \in \mathcal{F}_+(r)} R(T)^{W_L}$ satisfying the congruences (i) and

(*ii*). Hence the theorem. \Box

We shall denote by γ the restricted root $\alpha - \theta(\alpha)$ in $\Delta_{G/H}$ for $\alpha \in \Delta_H \setminus \Delta_L$ and by s_{γ} the corresponding simple reflection which is the image in $W_{G/H}$ of the element $s_{\alpha} \cdot s_{\theta(\alpha)} \in W_H$. (Recall that $s_{\gamma}^2 = 1$ [16, Definition 6, Lemma 10.2].)

We isolate below the description of $K_G(X^{wond})$ as a particular case of the above theorem. This is analogous to [17, Lemma 3.2] in the case of the wonderful compactification of an adjoint semi-simple group.

Corollary 3.3. The ring $K_G(X^{wond})$ is identified with the subring of $R(T)^{W_L}$ defined by the congruences

$$f \equiv s_{\alpha} s_{\theta(\alpha)} \cdot f \pmod{1 - e^{\gamma}}$$

for $\gamma = \alpha - \theta(\alpha) \in \Delta_{G/H}, \ \alpha \in \Delta_H \setminus \Delta_L$.

Proof: In this case $|\mathcal{F}_+(r)| = 1$ since X^{wond} has a unique closed *G*-orbit. Thus by Theorem 3.2 (1) $K_G(X^{wond})$ can be identified with the subring of $R(T)^{W_L}$ satisfying the congruences (*i*). Hence the corollary. \Box

Let \widetilde{G} denote the simply connected cover of G with projection $\pi : \widetilde{G} \to G$ and $\widetilde{T} := \pi^{-1}(T)$. Then \widetilde{G} is a semi-simple simply connected algebraic group with \widetilde{T} as a maximal torus. Let $\widetilde{P} := \pi^{-1}(P)$. There exists an involution $\widetilde{\theta}$ of \widetilde{G} which induces the involution θ of the adjoint quotient G. Let \widetilde{H} denote the fixed points of \widetilde{G} under $\widetilde{\theta}$. Then $\widetilde{T}_H := \widetilde{H} \cap \widetilde{T}$ is a maximal torus of \widetilde{H} . We have the following theorem analogous to Theorem 3.2 obtained by replacing G by \tilde{G} and T by \tilde{T} and by considering the action of \tilde{G} and \tilde{T} on X and Y through their canonical surjections to G and T respectively. We omit the proof to avoid repetition.

Theorem 3.4. The map

 $r: K_{\widetilde{G}}(X) \longrightarrow K_{\widetilde{T}}(X)^W \longrightarrow K_{\widetilde{T}}(X)^{W_H} \longrightarrow K_{\widetilde{T}}(Y)^{W_H}$

obtained by composing the canonical maps is an isomorphism.

We further have the following corollary analogous to Corollary 3.3.

Corollary 3.5. The ring $K_{\widetilde{G}}(X^{wond})$ is identified with the subring of $R(\widetilde{T})^{W_L}$ defined by the congruences

$$f \equiv s_{\alpha} s_{\theta(\alpha)} \cdot f \pmod{1 - e^{\gamma}}$$

for $\gamma = \alpha - \theta(\alpha) \in \Delta_{G/H}, \ \alpha \in \Delta_H \setminus \Delta_L$.

Remark 3.6. For X^{wond} the set of simple restricted roots $\Delta_{G/H}$ are in bijection with the set of boundary divisors. Let D_{γ} denote the boundary divisor corresponding to $\gamma \in \Delta_{G/H}$. Let \mathcal{L}_{γ} denote the *G*-linearized (and hence \tilde{G} -linearized) line bundle on *X* corresponding to the *G*-stable boundary divisor D_{γ} for $\gamma \in \Delta_{G/H}$. Note that $X^*(T/T_H)$ has a \tilde{G} -invariant section whose zero locus is the boundary divisor D_{γ} for $\gamma \in \Delta_{G/H}$. Note that $X^*(T/T_H)$ has a basis consisting of the simple roots $\gamma \in \Delta_{G/H}$. Thus $R(T/T_H) = \mathbb{Z}[X^*(T/T_H)]$ is generated as a \mathbb{Z} -algebra by $e^{\gamma}, \gamma \in \Delta_{G/H}$. Thus we can identify $R(T/T_H)$ with the subalgebra of $K_G(X^{wond})$ generated by $[\mathcal{L}_{\gamma}]$ for $\gamma \in \Delta_{G/H}$ (see [17, Remark 3.7]). In particular, $\mathcal{L}_{\gamma} \mid_{Y_0^{wond}}$ is the T/T_H -linearized line bundle on the toric variety Y_0^{wond} corresponding to the ray in \mathcal{F}_+ generated by ω_{γ} .

Remark 3.7. (*Extension to other oriented cohomology theories*) The methods of [7] have earlier been used by V. Kiritchenko and A. Krishna in [11] to compute the rational equivariant algebraic cobordism ring of wonderful symmetric varieties of minimal rank. We wish to remark here that Lemma 2.4 above allows us to extend the description in [11, Theorem 6.4] to all regular symmetric varieties of minimal rank.

3.1. Ordinary K-ring of X. Note that we have the equalities $R(\tilde{T})^{W_H} = K_{\tilde{G}}(\tilde{G}/\tilde{B})^{W_H}$ is an $R(\tilde{G}) = R(\tilde{T})^W$ -subalgebra of $K_{\tilde{G}}(\tilde{G}/\tilde{B}) = R(\tilde{T})$.

Consider the augmentation map $\epsilon : R(\widetilde{G}) \longrightarrow \mathbb{Z}$ which takes an element [V] to dim(V). Thus \mathbb{Z} becomes a $R(\widetilde{G})$ -module via the augmentation map.

We have the isomorphisms ([13, Theorem 6.1.22])

(3.4)
$$K(X) \simeq \mathbb{Z} \otimes_{R(\widetilde{G})} K_{\widetilde{G}}(X)$$

(3.5)
$$K_{\widetilde{T}}(Y)^{W_H} = K_{\widetilde{G}}(\widetilde{G} \times_{\widetilde{B}} Y)^{W_H}.$$

Remark 3.8. Here $\widetilde{G} \times_{\widetilde{B}} Y$ is a toric bundle with fibre the T/T_H -toric variety Y and base $\widetilde{G}/\widetilde{B}$.

Proposition 3.9. We have the following isomorphism of ordinary K-rings:

$$K(X) \simeq K(\widetilde{G} \times_{\widetilde{B}} Y)^{W_H}.$$

Proof: Follows immediately from Theorem 3.2, (3.4) and (3.5). \Box

3.2. Determination of the structure of $K_G(X)$. We first fix some notations similar to [17, Section 2.1]. Let \mathcal{F} denote the (smooth projective) fan associated to Y. Recall that the Weyl group W_H acts on \mathcal{F} by reflection across the walls of the Weyl chambers and the cones in \mathcal{F} get permuted by this action of W_H and each cone is stabilized by the reflections corresponding to the walls of the Weyl chambers on which it lies. Let W_{τ} denote the subgroup of W_H which fixes the cone $\tau \in \mathcal{F}$. Then, in particular, $W_{\sigma} = W_L$ for $\sigma \in \mathcal{F}(r)$ where $r = \operatorname{rk}(G/H)$ and $W_{\{0\}} = W_H$. In particular, $W_H \supseteq W_{\tau} \supseteq W_L$ for every $\tau \in \mathcal{F}$. Indeed, W_H acts on \mathcal{F} via its projection to $W_H/W_L = W_{G/H}$ which is the Weyl group of the restricted root system with simple roots $\Delta_{G/H}$.

Let $\{\rho_j \mid j = 1, \ldots, d\}$ denote the set of edges of the fan \mathcal{F} and let $\tau(1)$ denote the set of edges of the cone τ for every $\tau \in \mathcal{F}$. Let v_j denote the primitive vector along the edge ρ_j . Let O_{τ} denote the T/T_H -orbit in Y corresponding to $\tau \in \mathcal{F}$. Let L_j denote the T/T_H -equivariant line bundle on Y corresponding to the

edge ρ_j . Note that L_j has a T/T_H -invariant section s_j whose zero locus is $\overline{O_{\rho_j}}$. Recall that Y_0 is the toric

variety associated to the fan \mathcal{F}_+ consisting of the cones $\tau \in \mathcal{F}$ which lie in the positive Weyl chamber. Let $X_F := \prod_{\rho_j \in F} (1 - X_j)$ in the Laurent polynomial algebra $\mathbb{Z}[X_1^{\pm 1}, \ldots, X_d^{\pm 1}]$ for every $F \subseteq \{\rho_j \mid j = 1, \ldots, d\}$. In particular we let $X_\tau := X_{\tau(1)} = \prod_{\rho_j \in \tau(1)} (1 - X_j)$ for every $\tau \in \mathcal{F}$.

Let $C_{\tau} := X_{\tau} \cdot \mathbb{Z}[X_{j}^{\pm 1} : \rho_{j} \in \tau(1)]$. Recall from [17, Lemma 2.8] that we have the additive decomposition

(3.6)
$$K_{T/T_H}(Y_0) = \bigoplus_{\tau \in \mathcal{F}_+} C_{\tau}$$

which follows from the Stanley-Reisner presentation of the T/T_H -equivariant Grothendieck ring of Y_0 , $K_{T/T_H}(Y_0) = \mathbb{Z}[X_j^{\pm 1} : \rho_j \in \mathcal{F}_+(1)] / \langle X_F \forall F \notin \mathcal{F}_+ \rangle \text{ (see [19, Theorem 6.4])}.$

Similarly by [19, Theorem 6.4] we also have the following Stanley Reisner presentation for the T/T_{H} equivariant Grothendieck ring of Y:

(3.7)
$$K_{T/T_H}(Y) \simeq \mathbb{Z}[X_1^{\pm 1}, \dots, X_d^{\pm 1}] / \langle X_F \; \forall \; F \notin \mathcal{F} \rangle$$

Furthermore, since W_H acts on \mathcal{F} we have an action of W_H on $\mathbb{Z}[X_1^{\pm 1}, \ldots, X_d^{\pm 1}]$, given by $w(X_{\rho_j}^{\pm 1}) =$ $X_{w(\rho_i)}^{\pm 1}$ for every $w \in W_H$. Thus $w(X_F) = X_{w(F)}$ for $F \subseteq \{\rho_j \mid j = 1, \ldots, d\}$ and $w \in W_H$, and since W_H permutes the cones of \mathcal{F} we further get an action of W_H on the Stanley-Reisner algebra $\mathbb{Z}[X_1^{\pm 1}, \ldots, X_d^{\pm 1}]/\langle X_F \forall F \notin \mathcal{F} \rangle$ so that (3.7) is an isomorphism of W_H -modules where the W_H -action on $K_{T/T_H}(Y)$ is induced from the W_H -action on Y.

We further have the additive decomposition

$$\mathbb{Z}[X_1^{\pm 1}, \dots, X_d^{\pm 1}] / \langle X_F \ \forall \ F \notin \mathcal{F} \rangle = \bigoplus_{\tau \in \mathcal{F}} X_\tau \cdot \mathbb{Z}[X_j^{\pm 1} : \rho_j \in \tau(1)],$$

where W_H acts on the right hand side as follows:

$$w \cdot (X_{\tau} \cdot \mathbb{Z}[X_j^{\pm 1} : \rho_j \in \tau(1)]) = X_{w(\tau)} \cdot \mathbb{Z}[X_j^{\pm 1} : \rho_j \in w(\tau)(1)]$$

for every $w \in W_H$.

Thus we can write

$$K_{T/T_H}(Y) = \bigoplus_{\tau \in \mathcal{F}_+} \bigoplus_{w \in W_H/W_\tau} X_{w(\tau)} \cdot \mathbb{Z}[X_j^{\pm 1} : \rho_j \in w(\tau)(1)]$$

where we recall that W_{τ} denotes the subgroup of W_H which fixes $\tau \in \mathcal{F}_+$.

This implies that as a W_H -module we can write

(3.8)
$$K_{T/T_H}(Y) = \bigoplus Ind_{W_{\tau}}^{W_H}C_{\tau}$$

where $C_{\tau} := X_{\tau} \cdot \mathbb{Z}[X_{j}^{\pm 1} : \rho_{j} \in \tau(1)]$. Further, since C_{τ} is fixed by W_{τ} ,

(3.9)
$$Ind_{W_{\tau}}^{W_H}C_{\tau} = \mathbb{Z}[W_H/W_{\tau}] \otimes C_{\tau}$$

Recall (see [13]) that the structure morphism $Y \longrightarrow pt$ induces canonical inclusions $R(T) \hookrightarrow K_T(Y)$ (resp. $R(T/T_H) \hookrightarrow K_{T/T_H}(Y)$ which gives an R(T)-algebra (resp. $R(T/T_H)$ -algebra) structure on $K_T(Y)$ (resp. $K_{T/T_H}(Y)).$

Note that we have an exact sequence

$$(3.10) 0 \longrightarrow X^*(T/T_H) \longrightarrow X^*(T) \longrightarrow X^*(T_H) \longrightarrow 0$$

of character groups, where $X^*(T/T_H)$ denotes the characters of T that are trivial when restricted to T_H . The exact sequence (3.10) splits by choosing a basis of $X^*(T_H)$ and by lifting every element of this basis to a character of T. This implies that we have the following isomorphism

(3.11)
$$R(T) \simeq R(T/T_H) \otimes R(T_H)$$

where $R(T) = \mathbb{Z}[X^*(T)], R(T_H) = \mathbb{Z}[X^*(T_H)], R(T/T_H) = \mathbb{Z}[X^*(T/T_H)]$ (see [7, page 485] for the analogous statement for Chow ring).

More generally, we have the following proposition whose proof is similar to that of [17, Lemma 1.7] (see [7, Lemma 2.3.2] for the analogous statement for Chow ring).

Proposition 3.10. We have the following isomorphisms of rings

(i) $K_T(Y) \simeq K_{T/T_H}(Y) \otimes_{\mathbb{Z}} R(T_H)$

(ii) $K_T(Y) \simeq K_{T/T_H}(Y) \otimes_{R(T/T_H)} R(T)$

Proof: It suffices to prove (i). For, (ii) will then follow from (3.11) since the right hand side of (ii) can be written as $K_{T/T_H}(Y) \otimes_{R(T/T_H)\otimes 1} R(T/T_H) \otimes R(T_H)$ which is isomorphic to the right hand side of (i).

We now show (i). Since $\chi \in X^*(T_H)$ can be lifted to a character $\chi' \in X^*(T)$ (by (3.10)), we have a canonical homomorphism of rings $K_{T/T_H}(Y) \otimes R(T_H) \longrightarrow K_T(Y)$ where $e^{\chi} \in R(T_H) = \mathbb{Z}[X^*(T_H)]$ maps to $e^{\chi'} \in R(T) = \mathbb{Z}[X^*(T)] \hookrightarrow K_T(Y)$ and the map from $K_{T/T_H}(Y)$ to $K_T(Y)$ is induced by the surjection $T \longrightarrow T/T_H$.

To define the inverse of the above homomorphism, we let E be a T-equivariant vector bundle on Y. Since Y is a T/T_H -variety, the T_H -action on Y is trivial. Thus we get a T_H -action on every fibre of E which gives a weight space decomposition on each fibre. Further, the T-equivariant vector bundle E is locally trivial so the weights of the restricted T_H -action are locally constant. Moreover, since Y is irreducible these weights are constant over Y. Thus we get an isotypical decomposition $E = \bigoplus E_i$ where E_i denotes the subbundle of E whose fibre is the eigenspace for the T_H -action corresponding to a character χ_i of T_H . We can write $E_i = e^{\chi'_i} \otimes E^i$ where E^i is a T-equivariant vector bundle with a trivial T_H -action and hence a T/T_H -equivariant vector bundle on Y. We therefore define the inverse map which sends $[E] \in K_T(Y)$ to $\sum_i e^{\chi_i} \otimes [E^i] \in R(T_H) \otimes_{\mathbb{Z}} K_{T/T_H}(Y)$. \Box

Remark 3.11. Note that the above proposition holds in a more general setting where we can take

$$1 \longrightarrow T' \longrightarrow T \longrightarrow T'' \longrightarrow 1$$

to be an exact sequence of tori and Y to be any irreducible T''-variety. The proof follows verbatim. We have stated it in the above form for our convenience.

With the above notations we have the following theorem which extends [17, Theorem 2.10].

Theorem 3.12. Let X be a complete symmetric variety of minimal rank and Y be the corresponding torus embedding of T/T_H . The ring $K_G(X)$ has the following direct sum decomposition as a $1 \otimes R(H)$ -module:

(3.12)
$$K_G(X) = \bigoplus_{\tau \in \mathcal{F}_+} C_\tau \otimes R(T_H)^{W_\tau}$$

where $R(H) = R(T_H)^{W_H}$ acts naturally on the second factor in each piece of the above decomposition. Further, the multiplicitive structure of $K_G(X)$ can be described from the above decomposition as follows: Let $a_{\tau} \otimes b_{\tau} \in C_{\tau} \otimes R(T_H)^{W_{\tau}}$ and let $a_{\sigma} \otimes b_{\sigma} \in C_{\sigma} \otimes R(T_H)^{W_{\sigma}}$. Then

$$(a_{\tau} \otimes b_{\tau}) \cdot (a_{\sigma} \otimes b_{\sigma}) = \begin{cases} a_{\tau} \cdot a_{\sigma} \otimes b_{\tau} \cdot b_{\sigma} & \text{if } \tau \text{ and } \sigma \text{ span the cone } \gamma \\ 0 & \text{if } \tau \text{ and } \sigma \text{ do not span a cone in } \mathcal{F}_{+} \end{cases}$$

Here $a_{\tau} \cdot a_{\sigma} \otimes b_{\tau} \cdot b_{\sigma} \in C_{\gamma} \otimes R(T_H)^{W_{\gamma}}$ when τ and σ span γ . The multiplication in the first factor is as in $K_{T/T_H}(Y_0)$ where $C_{\tau} \cdot C_{\sigma} \subseteq C_{\gamma}$.

Proof: We have the following isomorphism by Theorem 3.2:

$$K_G(X) \simeq K_T(Y)^{W_H}$$

This together with Proposition 3.10(i) will imply that

(3.13)
$$K_G(X) \simeq (K_{T/T_H}(Y) \otimes_{\mathbb{Z}} R(T_H))^{W_H}.$$

Now, by (3.8) and (3.9) we get:

$$K_{T/T_H}(Y) \otimes_{\mathbb{Z}} R(T_H) = \bigoplus_{\tau \in \mathcal{F}_+} \mathbb{Z}[W_H/W_\tau] \otimes C_\tau \otimes R(T_H).$$

Further, by taking W_H -invariants on either side we get:

(3.14)
$$(K_{T/T_H}(Y) \otimes_{\mathbb{Z}} R(T_H))^{W_H} = \bigoplus_{\tau \in \mathcal{F}_+} C_\tau \otimes R(T_H)^{W_\tau}$$

Thus from (3.13) and (3.14), the additive decomposition (3.12) follows.

Observe that $C_{\tau} \cdot C_{\sigma} \subseteq C_{\gamma}$ whenever τ and σ span a cone γ in \mathcal{F}_{+} and $C_{\tau} \cdot C_{\sigma} = 0$ if τ and σ do not span a cone in \mathcal{F}_{+} . Furthermore, whenever $\gamma = \langle \tau, \sigma \rangle$ in \mathcal{F}_{+} the product $R(T_{H})^{W_{\tau}} \cdot R(T_{H})^{W_{\sigma}} \subseteq R(T)^{W_{\gamma}}$ since $R(T_{H})^{W_{\tau}}$ and $R(T_{H})^{W_{\sigma}}$ are both subrings of $R(T)^{W_{\gamma}}$.

The multiplicative structure now follows exactly as in the proof of [17, Theorem 2.10]. \Box

We have the following corollary which extends [17, Corollary].

Corollary 3.13. The ring $K_G(X) \simeq \bigoplus_{\tau \in \mathcal{F}_+} C_\tau \otimes R(T_H)^{W_\tau}$ admits a multifiltration $\{F_\tau\}_{\tau \in \mathcal{F}_+}$ where the filtered pieces are

$$F_{\tau} = \bigoplus_{\tau \preceq \sigma} C_{\sigma} \otimes R(T_H)^{W_{\sigma}},$$

where $F_{\tau} \supseteq F_{\sigma}$ whenever $\tau \preceq \sigma$, and $F_{\{0\}} = K_G(X)$. Further, under the multiplication described in Theorem 3.12, we have $F_{\tau} \cdot F_{\sigma} \subseteq F_{\gamma}$ whenever $\gamma = \langle \tau, \sigma \rangle$ in \mathcal{F}_+ . In particular $F_{\{0\}} \cdot F_{\tau} \subseteq F_{\tau}$ for all $\tau \in \mathcal{F}_+$.

Proof: The existance of the filtration $\{F_{\tau}\}_{\tau \in \mathcal{F}_{+}}$ and the properties follow by definition. Further, since the filtered pieces multiply by the multiplication rule given in Theorem 3.12, it follows that $F_{\tau} \cdot F_{\sigma} \subseteq F_{\gamma}$ whenever $\gamma = \langle \tau, \sigma \rangle$ in \mathcal{F}_{+} and $F_{\tau} \cdot F_{\sigma} \subseteq 0$ whenever τ and σ do not span a cone in \mathcal{F}_{+} . \Box

In the following proposition we show that the above multifiltration establishes the existence of the structure of $K_{T/T_H}(Y_0) \otimes R(H)$ -algebra on $K_G(X)$. We further show that the multifiltration also implies the existence of certain canonical $K_{T/T_H}(Y_0) \otimes R(H)$ -submodules which will be subsequently used in the next section. As a consequence of Theorem 3.12, we further show that $K_G(X)$ has the structure of a canonical $K_{T/T_H}(Y_0) \otimes R(H)$ submodule of $K_{T/T_H}(Y_0) \otimes R(T_H)^{W_L}$.

Proposition 3.14.

- (i) We have a canonical inclusion $K_{T/T_H}(Y_0) \otimes R(H) \subseteq F_{\{0\}} = K_G(X)$ as $1 \otimes R(H)$ -subalgebra.
- (ii) We have a canonical inclusion of the $K_{T/T_H}(Y_0) \otimes R(H)$ -submodule

$$\prod_{p_j \in \tau(1)} X_\tau \cdot K_{T/T_H}(Y_0) \otimes R(T_H)^{W_\tau}$$

in $F_{\tau} \subseteq F_{\{0\}} = K_G(X)$.

(iii) We have a canonical inclusion of $K_G(X)$ in $K_{T/T_H}(Y_0) \otimes R(T_H)^{W_L}$ as a $K_{T/T_H}(Y_0) \otimes R(H)$ -subalgebra. (This is analogous to [17, Proposition 2.5] and [18, Proposition 2.1].)

Proof: (i) From (3.6) it follows that

(3.15)
$$K_{T/T_H}(Y_0) \otimes R(H) = \bigoplus_{\tau \in \mathcal{F}_+} C_\tau \otimes R(H).$$

Also

(3.16)
$$C_{\tau} \otimes R(H) = C_{\tau} \otimes R(T_H)^{W_H} \subseteq C_{\tau} \otimes R(T_H)^{W_{\tau}}$$

for every $\tau \in \mathcal{F}_+(1)$.

Since both (3.12) and (3.15) are decopositions as as $1 \otimes R(H)$ -algebras, (i) follows from (3.16).

(ii) Again from (3.6) we have

(3.17)
$$X_{\tau} \cdot K_{T/T_H}(Y_0) \otimes R(T_H)^{W_{\tau}} = \bigoplus_{\sigma \in \mathcal{F}_+} X_{\tau} \cdot C_{\sigma} \otimes R(T_H)^{W_{\tau}}$$

Moreover, by the relations in the Stanley Reisner presentation of $K_{T/T_H}(Y_0)$ we have $X_{\tau} \cdot C_{\sigma} \subseteq C_{\gamma}$ whenever τ and σ span a cone γ in \mathcal{F}_+ and $X_{\tau} \cdot C_{\sigma} = 0$ whenever τ and σ do not span a cone in \mathcal{F}_+ . Thus

$$X_{\tau} \cdot C_{\sigma} \otimes R(T_H)^{W_{\tau}} \subseteq C_{\gamma} \otimes R(T_H)^{W_{\gamma}}$$

since $R(T)^{W_{\tau}} \subseteq R(T_H)^{W_{\gamma}}$ whenever $\tau \preceq \gamma$.

Now, (ii) follows by (3.17) and by the definition of F_{τ} from Corollary 3.13.

(iii) Recall that for every $\tau \in \mathcal{F}_+$ we have $W_\tau \supseteq W_L$ which further implies that $R(T_H)^{W_\tau} \subseteq R(T_H)^{W_L}$. Thus for every $\tau \in \mathcal{F}_+$ we have the inclusion of $1 \otimes R(H)$ -submodules

$$C_{\tau} \otimes R(T_H)^{W_{\tau}} \subseteq C_{\tau} \otimes R(T_H)^{W_H}$$

Taking direct sum over all $\tau \in \mathcal{F}_+$ we get:

V. UMA

(3.18)
$$\bigoplus_{\tau \in \mathcal{F}_+} C_\tau \otimes R(T_H)^{W_\tau} \subseteq \bigoplus_{\tau \in \mathcal{F}_+} C_\tau \otimes R(T_H)^{W_L}.$$

We see by (3.6) that the right hand side of the above equation is isomorphic to $K_{T/T_H}(Y_0) \otimes R(T_H)^{W_L}$. Now, by (3.12) and (3.18) (iii) follows. \Box

In the following corollary we give a geometric interpretation of Theorem 3.12. This is analogous to [17, Corollary 2.12] for regular compactifications.

Corollary 3.15. Let $N_{\tau} \simeq \bigoplus_{\rho_j \in \tau(1)} L_j$ be the normal bundle of $V_{\tau} = \overline{O_{\tau}}$ in Y. Let $N_{\tau} \mid_{O_{\tau}}$ denote the restriction of the normal bundle to O_{τ} so that so that

$$\lambda_{-1}(N_{\tau} \mid_{O_{\tau}}) := \prod_{\rho_j \in \tau(1)} (1 - [L_j] \mid_{O_{\tau}}) \in K_{T/T_H}(O_{\tau}).$$

Then we have the following decomposition:

$$K_G(X) \simeq \bigoplus_{\tau \in \mathcal{F}_+} \lambda_{-1}(N_\tau \mid_{O_\tau}) \cdot K_{T/T_H}(O_\tau) \otimes R(T_H)^{W_\tau}$$

Let $P_{\tau} := \lambda_{-1}(N_{\tau} \mid_{O_{\tau}}) \cdot K_{T/T_H}(O_{\tau})$ for each $\tau \in \mathcal{F}_+$. Then the above decoposition is a ring isomorphism where the multiplication on the right hand side is given as follows:

$$P_{\tau} \cdot P_{\sigma} \subseteq \begin{cases} P_{\gamma} & \text{if } \tau \text{ and } \sigma \text{ span the cone } \gamma \in \mathcal{F}_{+}, \\ 0 & \text{if } \tau \text{ and } \sigma \text{ do not span a cone in } \mathcal{F}_{+} \end{cases}$$

The structure of rational equivariant cohomology of regular embeddings have been described in [1] and [12, Theorem 2.3]. This result for a complete symmetric variety of minimal rank can alternately be obtained by proving results analogous to Theorem 3.2 and Theorem 3.12 for the equivariant cohomology of a complete symmetric variety (see [12] for the corresponding results on wonderful symmetric varieties).

Let $X_F = \prod_{\rho_j \in F} X_j$ for every $F \subseteq \{\rho_j \mid j = 1, \dots, d\}$ in the polynomial ring $\mathbb{Q}[X_1, \dots, X_d]$. In particular $X_\tau := X_{\tau(1)} = \prod_{\rho_j \in \tau(1)} X_j$ for every $\tau \in \mathcal{F}$. Let $e(N_\tau)$ denote the equivariant Euler class of the normal bundle $V_\tau = \overline{O_\tau}$. Let $S := H^*_{T_H}(pt)$ By [1, Theorem 8, p.7] we have:

$$H^*_{T/T_H}(Y) \simeq \mathbb{Q}[X_1, \dots, X_d]/\langle X_F \forall F \notin \mathcal{F} \rangle$$

Let $e(N_{\tau})$ denote the equivariant Euler class of the normal bundle of $V(\tau) = \overline{O_{\tau}}$ which is equal to the top Chern class of $\bigoplus_{\rho_j \in \tau(1)} L_j$. We then have the following description of the equivariant cohomology of a complete

symmetric variety of minimal rank (see [12] and [1]). We consider cohomology ring with \mathbb{Q} -coefficients.

Theorem 3.16.

$$H_G^*(X) \simeq \bigoplus_{\tau \in \mathcal{F}_+} e(N_\tau \mid_{O_\tau}) \cdot H_{T/T_H}(O_\tau) \otimes S^{W_\tau}$$

Note that $S^{W_H} = H^*_H(pt) \simeq H^*_G(G/H)$ and $H^*_{T/T_H}(Y_0)$ can be identified with the subalgebra of $H^*_G(X)$ generated by the equivariant classes of the boundary divisors.

Remark 3.17. From the above decomposition, an analogue of Proposition 3.14 for the equivariant cohomology can also be proved which will show that $H^*_G(X)$ is a $H^*_{T/T_H}(Y_0) \otimes S^{W_H}$ -submodule of $H^*_{T/T_H}(Y_0) \otimes S^{W_L}$ (see [7, Proposition 2.3.3] for a similar statement on $A^*_G(X)_{\mathbb{Q}}$). Our aim is to show that $H^*_G(X)$ is a free module of rank $|W_{G/H}|$ over $H^*_{T/T_H}(Y_0) \otimes S^{W_H}$ and to find a basis of this free module. This will be an extension of the results of Strickland [14] for the compactification of adjoint semisimple group to any complete symmetric variety of minimal rank. This will be taken up in a future work. We also note here that similar description of the equivariant cohomology ring of any regular compactification of an adjoint semisimple group has been obtained by Strickland (see [15, Theorem 6.1]).

12

3.3. Comparison with wonderful symmetric variety. Let X be a regular compactification of the adjoint symmetric space G/H and X^{wond} be the canonical wonderful compactification.

Let $\gamma_1, \ldots, \gamma_r$ denote the simple restricted roots.

Recall that \mathcal{L}_{γ_i} are \widetilde{G} -linearized line bundles on X^{wond} such that \widetilde{P} operates on \mathcal{L}_{γ_i} by the character γ_i for $1 \le i \le r$ where z is the base point of the unique closed orbit.

Furthermore, since the centre Z of G acts trivially on X^{wond} and hence on the fibre by the character γ_i the line bundle \mathcal{L}_{γ_i} is actually $G = \widetilde{G}/Z$ -linearized. Moreover, \mathcal{L}_{γ_i} admits a G-invariant section s_i whose zero locus is the boundary divisor D_{γ_i} for $1 \leq i \leq r$.

We recall the following construction from [1, Section 3.5] and [9, Section 10].

The bundle $\mathcal{V} = \bigoplus \mathcal{L}_{\gamma_i}$ being a direct sum of line bundles on X^{wond} admits a natural action of the r-dimensional torus \mathbb{G}_m^r . Put

$$\mathcal{P} := \mathcal{V} \setminus \bigcup_{i=1}^{\prime} \mathcal{L}_{\gamma_1} \oplus \cdots \oplus \hat{\mathcal{L}}_{\gamma_i} \oplus \cdots \oplus \mathcal{L}_{\gamma_r}.$$

Then \mathcal{P} is the principal $T/T_H = \mathbb{G}_m^r$ -bundle associated with \mathcal{V} over X^{wond} . The section $s = s_1 \oplus \cdots \oplus s_r$ of \mathcal{V} clearly maps G to \mathcal{P} . Therefore using s we can embed X^{wond} in the bundle $\mathcal{P} \times_{T/T_H} \mathbb{A}^r$ which is nothing but \mathcal{V} . Since the bundles \mathcal{L}_{γ_i} are G-linearized it follows that \mathcal{P} is a left G-space whose bundle map $\mathcal{P} \longrightarrow X^{wond}$ is G-equivariant for the canonical G-action on X^{wond} . Moreover, the right T/T_H -action is compatible with the left G-action on \mathcal{P} .

Consider the proper morphism of toric varieties $\varphi: Y_0 \longrightarrow Y_0^{wond} \simeq \mathbb{A}^r$ corresponding to a smooth subdivision \mathcal{F}_+ of the positive Weyl chamber \mathcal{C}_+ . Since Y_0 is a T/T_H -toric variety we can form the associated toric bundle $\mathcal{P} \times_{T/T_H} Y_0$ over X^{wond} . The morphism φ further induces $\overline{\varphi} : \mathcal{P} \times_{T/T_H} Y_0 \longrightarrow \mathcal{P} \times_{T/T_H} Y_0^{wond} = \mathcal{V}$ which is isomorphic to the pull back of $\mathcal{P} \times_{T/T_H} Y_0$ to \mathcal{V} via the bundle projection $p: \mathcal{V} \longrightarrow X^{wond}$.

From [1, Definition 23] and [9, Section 3.2], $X = X_{Y_0} = \overline{\varphi}^{-1}(s(X_{word}))$ which can be identified with pull-back of $\mathcal{P} \times_{T/T_H} Y_0 \xrightarrow{\overline{\varphi}} \mathcal{P} \times_{T/T_H} \mathbb{A}^r$ via the section s. Thus $X = X_{Y_0} = s^*(p^*(\mathcal{P} \times_{T/T_H} Y_0))$. Now, $p \circ s = Id_{X^{wond}}$ hence $s^* \circ p^* = Id^*$. Thus

$$(3.19) X \simeq \mathcal{P} \times_{T/T_H} Y_0.$$

For $u \in X^*(T/T_H)$ let $\mathcal{L}_u := \mathcal{P} \times_{T/T_H} \mathbb{C}_u$ denote the *G*-equivariant line bundle on X^{wond} associated to the character u. Let $[\mathcal{L}_u]_G$ denote its class in $K_G(X^{wond})$.

Let X be a projective regular compactification of G/H so that Y_0 is a semi-projective T/T_H -toric variety. Then the following theorem follows by [18, Theorem 5.1] and (3.19).

Theorem 3.18. The G-equivariant Grothendieck ring of X has the following description as an algebra over the G-equivariant Grothendiek ring of X^{wond} :

(3.20)
$$K_G(X) \simeq \frac{K_G(X^{wond})[X_j^{\pm 1} : \rho_j \in \mathcal{F}_+(1)]}{\mathcal{J}}$$

where \mathcal{J} is the ideal in $K_G(X^{wond})[X_j^{\pm 1} : \rho_j \in \mathcal{F}_+(1)]$ generated by the elements X_F for $F \notin \mathcal{F}_+$ and $\prod \quad X_j^{\langle u, v_j \rangle} - [\mathcal{L}_u]_G \text{ for } u \in X^*(T/T_H).$ $\rho_j \in \mathcal{F}_+(1)$

Recall that the adjoint symmetric spaces of minimal rank are exactly the products of the symmetric spaces listed in [7, Examples 1.4.4]. In particular, [7, Examples 1.4.4](1) is the adjoint semisimple group $G = G \times G/\text{diag}(G)$ and [7, Examples 1.4.4](2) is the space G/H = PGL(2n)/PSp(2n). The $G \times G$ equivariant K-ring of (1) has already beed described in [17]. In the following example we shall describe the G-equivariant K-ring of the wonderful compactification X^{wond} of PGL(2n)/PSp(2n).

Example 3.19.

Consider the group G = PGL(2n) and the involution θ associated with the symmetry of the Dynkin diagram so that H = PSp(2n) and rk(G/H) = n - 1. We shall consider the case when n = 3 for simplicity. The general case is similar.

V. UMA

Let $\Delta_G = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5\}$ be the simple roots of the root system A_5 corresponding to PGL(6). Then $\Delta_L = \{\alpha_1, \alpha_3, \alpha_5\}$. Also $\Delta_G \setminus \Delta_L = \{\alpha_2, \alpha_4\}$. Thus $\Delta_{G/H} = \{\gamma_1 = \alpha_2 - \theta(\alpha_2), \gamma_2 = \alpha_4 - \theta(\alpha_4)\}$ are the simple restricted roots.

Since T_H is a maximal torus of the adjoint semisimple group of type C_3 it is known that $R(T_H) = \mathbb{Z}[X^*(T_H)]$ where $X^*(T_H)$ has a basis consisting of the $\operatorname{rk}(H) = 3$ simple roots $\Delta_H = \{\beta_1 = 2e_1, \beta_2 = e_2 - e_1, \beta_3 = e_3 - e_2\}$ where β_1 is a long root and β_2 and β_3 are short roots.

Note that $\theta(\alpha_1) = \alpha_1$, $\theta(\alpha_3) = \alpha_3$, $\theta(\alpha_5) = \alpha_5$, $\theta(\alpha_2) = -\alpha_1 - \alpha_2 - \alpha_3$ and $\theta(\alpha_4) = -\alpha_3 - \alpha_4 - \alpha_5$.

Thus under the restriction map $q: X^*(T) \longrightarrow X^*(T_H)$ (see [7, Lemma 1.4.2]) $q^{-1}(\beta_1)$ consists of the unique root α_1 , $q^{-1}(\beta_2)$ consists of the two strongly orthogonal roots $\alpha_2 + \alpha_3$ and $-\alpha_1 - \alpha_2$ and $q^{-1}(\beta_3)$ consists of the two strongly orthogonal roots $-\alpha_3 - \alpha_4$ and $\alpha_4 + \alpha_5$. The restricted simple roots are $\gamma_1 = \alpha_2 - \theta(\alpha_2) = \alpha_1 + 2\alpha_2 + \alpha_3$ and $\gamma_2 = \alpha_4 - \theta(\alpha_4) = \alpha_3 + 2\alpha_4 + \alpha_5$.

Here $W = S_6$, W_H is the semidirect product of $S_2 \times S_2 \times S_2$ with S_3 , $W_L = S_2 \times S_2 \times S_2$ and $W_{G/H} = S_3$. In this case the toric variety Y^{wond} is associated to the Weyl chambers of type A_2 and Y_0^{wond} is the toric variety associated to the positive Weyl chamber in the coweight lattice spanned by the fundamental coweights $\omega_{\gamma_1}^{\vee}$ and $\omega_{\gamma_2}^{\vee}$ dual to the simple roots γ_1 and γ_2 respectively.

The simple reflections in $W_{G/H}$ corresponding to γ_1 and γ_2 are s_{γ_1} and s_{γ_2} respectively. Here \mathcal{F}_+ consists of 4 cones namely {0}, ρ_1, ρ_2 and $\sigma = \langle \rho_1, \rho_2 \rangle$, where the coweight vectors $\omega_{\gamma_1}^{\vee}$ and $\omega_{\gamma_2}^{\vee}$ are the primitive vectors along the rays ρ_1 and ρ_2 respectively. Note that $W_{\sigma} = W_L = S_2 \times S_2 \times S_2$, $W_{\{0\}} = W_H$ which is the semidirect product of W_L with S_3 , W_{ρ_1} is the semidirect product of W_L with the subgroup of $W_{G/H} = S_3$ generated by s_{γ_2} , W_{ρ_2} is the semidirect product of W_L with the subgroup of $W_{G/H} = S_3$ generated by s_{γ_1} .

Let $\mathcal{R} := R(T_H)^{W_L}$. Further, $R(PSp(6)) = R(H) = R(T_H)^{W_H} = \mathcal{R}^{W_{G/H}}$. Furthermore, $K_{T/T_H}(Y_0) \simeq R(T/T_H) = \mathbb{Z}[e^{\pm \gamma_1}, e^{\pm \gamma_2}] \simeq \mathbb{Z}[X_{\rho_1}^{\pm 1}, X_{\rho_2}^{\pm 1}]$ where X_{ρ_1} (respectively X_{ρ_2})maps to the class of the trivial line bundle on Y_0 where T/T_H acts on the fibre via the character γ_1 (respectively γ_2). The following is an additive decomposition of $K_{PGL_6}(X)$ as a $1 \otimes R(PSp(6))$ -submodule of $R(T/T_H) \otimes \mathcal{R}$.

$$\mathbb{Z} \otimes R(PSp(6)) \oplus (1 - e^{\gamma_1}) \cdot \mathbb{Z}[e^{\pm \gamma_1}] \otimes \mathcal{R}^{s_{\gamma_2}} \oplus \\ \oplus (1 - e^{\gamma_2}) \cdot \mathbb{Z}[e^{\pm \gamma_2}] \otimes \mathcal{R}^{s_{\gamma_1}} \oplus (1 - e^{\gamma_1}) \cdot (1 - e^{\gamma_2}) \cdot \mathbb{Z}[e^{\pm \gamma_1}, e^{\pm \gamma_2}] \otimes \mathcal{R}$$

Remark 3.20. By the parametrization of line bundles on spherical varieties (see [3, Section 2.2]) the \tilde{G} -linearized line bundles on X correspond to $PL(\mathcal{F}_+)$ which denotes the piecewise linear functions on the fan \mathcal{F}_+ . Let \mathcal{L}_h be the \tilde{G} -linearized line bundle on X corresponding to $h = (h_\sigma)_{\sigma \in \mathcal{F}_+(r)}$ then $\tilde{P} = \pi^{-1}(P)$ acts on $\mathcal{L}|_{z_\sigma}$ by the character h_σ . Thus $\mathbf{L}_h = \mathcal{L}_h |_{Y_0}$ is a \tilde{T}/\tilde{T}_H -linearized line bundle on Y_0 corresponding to the piecewise linear function $h \in PL(\mathcal{F}_+)$ so that \tilde{T}/\tilde{T}_H acts on $\mathbf{L}_h |_{z_\sigma}$ by the character h_σ . Since it is known that $\operatorname{Pic}^{\tilde{T}/\tilde{T}_H}(Y_0)$ generates $K_{T/T_H}(Y_0)$ as a ring (see [19]) it follows that $K_{\tilde{T}/\tilde{T}_H}(Y_0)$ is the subring of $K_{\tilde{G}}(X)$ generated by $\operatorname{Pic}^{\tilde{G}}(X)$. We can further identify $K_{\tilde{G}}(G/H) \simeq R(\tilde{H})$. We wonder whether $K_{\tilde{G}}(X)$ has a canonical $K_{\tilde{T}/\tilde{T}_H}(Y_0) \otimes R(\tilde{H})$ -module structure so that $K_{\tilde{G}}(X)$ is a free module of rank $|W_{G/H}|$ over $K_{\tilde{T}/\tilde{T}_H}(Y_0) \otimes R(\tilde{H})$ and if there exists a basis for this free module generalizing the results in [17, Section 3] and [18] to the setting of all complete symmetric varieties of minimal rank. See [6, Theorem 4 (ii)] for a similar statement describing $K(X^{wond})_{\mathbb{Q}}$ as a free module of rank $|W_H/W_L|$ over the classes of the boundary divisors $[\mathcal{O}_{D_\gamma}]$ for $\gamma \in \Delta_{G/H}$ together with the classes of equivariant vector bundles $\Phi(\lambda)$ induced by the fundamental weights λ of H.

An analogue of [7, Proposition 2.3.1] for the \tilde{T} and \tilde{G} -equivariant Grothendieck ring may give us a way to extend [17, Theorem 3.3] to all wonderful symmetric varieties of minimal rank. However the proof of [7, Proposition 2.3.1] does not extend to the integral setting of the Grothendieck ring, due to the reason that there is no W_L -invariant splitting for short exact sequence of character groups

$$(3.21) 0 \longrightarrow X^*(\widetilde{T}/\widetilde{T}_H) \longrightarrow X^*(\widetilde{T}) \longrightarrow X^*(\widetilde{T}_H) \longrightarrow 0.$$

This can be seen from the following example of $PGL_4/PSp(4)$.

Example 3.21. In the case of G = PGL(4) and H = PSp(4), $\tilde{G} = SL(4)$ and $\tilde{H} = Sp(4)$. The maximal torus \tilde{T} is the diagonal torus $(t_1, t_2, t_3, t_4) \in (\mathbb{C}^*)^4$ satisfying $t_1 \cdot t_2 \cdot t_3 \cdot t_4 = 1$. The subtorus \tilde{T}_H of \tilde{T} consisting of (t_1, t_2, t_3, t_4) such that $t_1 \cdot t_2 = 1$ and $t_3 \cdot t_4 = 1$. Here $W_L \simeq S_2 \times S_2$ is the subgroup of S_4 generated by the transpositions $\{(1, 2), (3, 4)\}$. The existance of a W_L -invariant splitting for the restriction

15

homomorphism

$$(3.22) q: X^*(\widetilde{T}) \longrightarrow X^*(\widetilde{T}_H)$$

would mean the existance of a complementary subtorus of \widetilde{T}_H in \widetilde{T} which is W_L -invariant. A 1-dimensional subtorus of \widetilde{T} is a 1-parameter subgroup $(t^{a_1}, t^{a_2}, t^{a_3}, t^{a_4})$ for $t \in \mathbb{C}^*$ where $(a_1, a_2, a_3, a_4) \in \mathbb{Z}^4$ satisfies $a_1 + a_2 + a_3 + a_4 = 0$. If this is W_L invariant then it will be of the form $(t^{a_1}, t^{a_1}, t^{a_3}, t^{a_3})$ such that $a_1 + a_3 = 0$. Thus it will be of the form (t, t, t^{-1}, t^{-1}) for $t \in \mathbb{C}^*$. However, this subtorus is not a complement of \widetilde{T}_H since it intersects \widetilde{T}_H along the subgroup $\{\pm I\}$ of order 2. Thus in general there cannot exist a W_L -invariant splitting for (3.22).

Remark 3.22. (Assumptions on the base field) Although we work over the field of complex numbers all the results in this paper should hold over an algebraically closed field k of characteristic $\neq 2$ (see [7, p. 473] and [10, p. 273-274]).

Acknowledgments: I am very grateful to Prof. Michel Brion for his valuable comments and suggestions while preparing this manuscript. I thank the referee for a careful reading of the manuscript and for very helpful comments and suggestions. I wish to mention that Remark 3.7 has been added based on the referee's observation.

Statements and Declarations I hereby declare that this work has no related financial or non-financial conflict of interests.

References

- [1] E. Bifet, C. De Concini and C. Procesi, Cohomology of regular embeddings, Adv. in Math. 82 (1990), 1-34.
- [2] N. Bourbaki, Groupes et Algebres de Lie, Hermann, Paris 1968.
- [3] M. Brion, Groupe de Picard et nombres caracteristiques des varietes spheriques, Duke Math. J. 58 (1989), no.2, 397-424.
- [4] M. Brion, Equivariant Chow Groups for Torus Actions, Journal of Transformation Groups 2 (1997), 225-267.
- [5] M. Brion, Behaviour at infinity of the Bruhat decomposition, Comment Math. Helv. 73 (1998), 137-174.
- [6] M. Brion, Construction of equivariant vector bundles in: Proceedings of the International Colloquium on Algebraic Groups and Homogeneous Spaces (Mumbai 2004), Narosa, New Delhi, 2007, 83-111.
- M. Brion and R. Joshua, Equivariant Chow ring and Chern classes of wonderful symmetric varieties of minimal rank, Transformation Groups 13 (2008), 471-493.
- [8] C. De Concini and C. Procesi, Complete symmetric varieties, *Invariant Theory* (Proceedings, Montecatini 1982), *Lecture Note in Math.* 996, 1-44, Springer-Verlag, New York 1983.
- [9] C. De Concini and C. Procesi, Complete Symmetric Varieties II Intersection theory, Advanced Studies in Pure Mathematics 6, 1985, Algebraic Groups and Related Topics(Kyoto/Nagoya, 1983) pp. 481-513, Publisher: North-Holland.
- [10] C. De Concini and T. Springer, Compactifications of symmetric varieties, Transformation Groups 4 (1999), 273-300.
- [11] V. Kiritchenko and A. Krishna, Equivariant cobordism of flag varieties and of symmetric varieties, Transformation Groups, 18 (2013), no. 2, 391-413.
- [12] P. Littelmann and C. Procesi, Equivariant cohomology of wonderful compactifications, in: Operator Algebras, Unitary Representations, Enveloping Algebras, and Invariant Theory, Progress in Mathematics, Vol. 92, Birkhauser, Boston 1990, 219-262.
- [13] A.S. Merkurjev, Comparison of equivariant and ordinary K-theory of algebraic varieties, Algebra i Anliz 9 (1997), 175-214. Translation in St. Petersburg Math. J. 9 (1998), 815-850.
- [14] E. Strickland, Equivariant cohomology of the wonderful group compactification, J. Algebra 306 (2006), 610-621.
- [15] E. Strickland, The ring of conditions of a semisimple group, J. Algebra 320 (2008), 3069-3078.
- [16] A. Tchoudjem, Cohomologie des fibrés en droites sur les variétés magnifiques de rang minimal, Bulletin de la Société Mathématique de France, 135 (2007) no. 2, p. 171-214.
- [17] V. Uma, Equivariant K-theory of compactifications of algebraic groups, Transformation groups 12, No. 2,(2007), 371-406.
- [18] V. Uma, Equivariant K-theory of group compactifications: further developments, Izv. Math. 80 No. 2, (2016), 417-441.
- [19] G. Vezzosi and A. Vistoli, Higher algebraic K-theory for actions of diagonalizable groups, Invent. Math. 153 (2003), 1-44.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY-MADRAS, CHENNAI, INDIA *Email address:* vuma@iitm.ac.in