

# Tighter uncertainty relations based on $(\alpha, \beta, \gamma)$ modified weighted Wigner-Yanase-Dyson skew information of quantum channels

Cong Xu<sup>1</sup>, Zhaoqi Wu<sup>1\*</sup>, Shao-Ming Fei<sup>2,3</sup>

1. *Department of Mathematics, Nanchang University, Nanchang 330031, P R China*

2. *School of Mathematical Sciences, Capital Normal University, Beijing 100048, P R China*

3. *Max-Planck-Institute for Mathematics in the Sciences, 04103 Leipzig, Germany*

## Abstract

We use a novel formation to illustrate the  $(\alpha, \beta, \gamma)$  modified weighted Wigner-Yanase-Dyson  $((\alpha, \beta, \gamma)$  MWWYD) skew information of quantum channels. By using operator norm inequalities, we explore the sum uncertainty relations for arbitrary  $N$  quantum channels and for unitary channels. These uncertainty inequalities are shown to be tighter than the existing ones by a detailed example. Our results are also applicable to the modified weighted Wigner-Yanase-Dyson (MWWYD) skew information and the  $(\alpha, \gamma)$  modified weighted Wigner-Yanase-Dyson  $((\alpha, \gamma)$  MWWYD) skew information of quantum channels as special cases.

**Keywords:** Uncertainty relation;  $(\alpha, \beta, \gamma)$  MWWYD skew information; Quantum channel

## 1. Introduction

As an extremely important issue in quantum physics, the uncertainty principle has been widespread concerned since Heisenberg [1] proposed the notions of uncertainties in measuring non-commuting observables. Based on the variance of measurement outcomes the well-known Heisenberg-Robertson uncertainty relation [2] says that for arbitrary two observables  $A$  and  $B$  with respect to a quantum state  $|\psi\rangle$ , one has

$$\Delta A \Delta B \geq \frac{1}{2} |\langle \psi | [A, B] | \psi \rangle|, \quad (1)$$

where  $[A, B] = AB - BA$  and  $\Delta M = \sqrt{\langle \psi | M^2 | \psi \rangle - \langle \psi | M | \psi \rangle^2}$  is the standard deviation of an observable  $M$ .

---

\*Corresponding author. E-mail: wuzhaoqi\_conquer@163.com

There are also many ways to describe uncertainty relations, such as entropy [3–7], variance [8–11] and majorization techniques [12–15]. In particular, the quantum uncertainty can also be characterized by skew information. The skew information has been initially proposed by Wigner and Yanase [16], termed as *Wigner-Yanase* (WY) skew information. Then a more general quantity has been suggested by Dyson, called the *Wigner-Yanase-Dyson* (WYD) skew information [16]. This quantity has been further generalized in [17] and termed as *generalized Wigner-Yanase-Dyson* (GWYD) skew information. The uncertainty relations based on WY skew information, WYD skew information and GWYD skew information have been studied extensively [18–23].

For a quantum state  $\rho$  and an observable  $A$ , Furuichi, Yanagi and Kuriyama [24] defined another generalized Wigner-Yanase skew information,

$$K_{\rho}^{\alpha}(A) = -\frac{1}{2}\text{Tr} \left( \left[ \frac{\rho^{\alpha} + \rho^{1-\alpha}}{2}, A \right]^2 \right) = \frac{1}{2} \left\| \left[ \frac{\rho^{\alpha} + \rho^{1-\alpha}}{2}, A \right] \right\|^2, \quad 0 \leq \alpha \leq 1, \quad (2)$$

which, called as the *weighted Wigner-Yanase-Dyson* (WWYD) skew information in [23], is different from WYD skew information. Chen, Liang, Li and Wang [25] proposed then a generalized Wigner-Yanase skew information for arbitrary operator  $E$  (not necessarily Hermitian),

$$K_{\rho}^{\alpha}(E) = -\frac{1}{2}\text{Tr} \left( \left[ \frac{\rho^{\alpha} + \rho^{1-\alpha}}{2}, E^{\dagger} \right] \left[ \frac{\rho^{\alpha} + \rho^{1-\alpha}}{2}, E \right] \right) = \frac{1}{2} \left\| \left[ \frac{\rho^{\alpha} + \rho^{1-\alpha}}{2}, E \right] \right\|^2, \quad 0 \leq \alpha \leq 1, \quad (3)$$

which is termed as the *modified weighted Wigner-Yanase-Dyson* (MWWYD) skew information in [23]. By replacing the arithmetic mean of  $\rho^{\alpha}$  and  $\rho^{1-\alpha}$  with their convex combination, the *two-parameter extension of the Wigner-Yanase skew information* is introduced in [26],

$$\begin{aligned} K_{\rho, \gamma}^{\alpha}(A) &= -\frac{1}{2}\text{Tr} \left( [(1-\gamma)\rho^{\alpha} + \gamma\rho^{1-\alpha}, A]^2 \right) \\ &= \frac{1}{2} \left\| [(1-\gamma)\rho^{\alpha} + \gamma\rho^{1-\alpha}, A] \right\|^2, \quad 0 \leq \alpha \leq 1, \quad 0 \leq \gamma \leq 1, \end{aligned} \quad (4)$$

which is called the  $(\alpha, \gamma)$  *weighted Wigner-Yanase-Dyson*  $((\alpha, \gamma)$  WWYD) skew information in [27]. Note that Eq. (4) reduces to Eq. (2) when  $\gamma = \frac{1}{2}$ .

We defined the  $(\alpha, \beta, \gamma)$  *weighted Wigner-Yanase-Dyson*  $((\alpha, \beta, \gamma)$  WWYD) skew information as [27],

$$\begin{aligned} K_{\rho, \gamma}^{\alpha, \beta}(A) &= -\frac{1}{2}\text{Tr} \left( [(1-\gamma)\rho^{\alpha} + \gamma\rho^{\beta}, A]^2 \rho^{1-\alpha-\beta} \right) \\ &= \frac{1}{2} \left\| \left[ (1-\gamma)\rho^{\alpha} + \gamma\rho^{\beta}, A \right] \rho^{\frac{1-\alpha-\beta}{2}} \right\|^2, \quad \alpha, \beta \geq 0, \quad \alpha + \beta \leq 1, \quad 0 \leq \gamma \leq 1, \end{aligned} \quad (5)$$

which reduces to Eq. (4) when  $\beta = 1 - \alpha$ . We also defined the  $(\alpha, \beta, \gamma)$  *modified weighted Wigner-Yanase-Dyson*  $((\alpha, \beta, \gamma)$  MWWYD) skew information with respect to a quantum

state  $\rho$  and an arbitrary operator  $E$  (not necessarily Hermitian) in [27] as

$$\begin{aligned} K_{\rho,\gamma}^{\alpha,\beta}(E) &= -\frac{1}{2}\text{Tr}([(1-\gamma)\rho^\alpha + \gamma\rho^\beta, E^\dagger][(1-\gamma)\rho^\alpha + \gamma\rho^\beta, E]\rho^{1-\alpha-\beta}) \\ &= \frac{1}{2}\left\|[(1-\gamma)\rho^\alpha + \gamma\rho^\beta, E]\rho^{\frac{1-\alpha-\beta}{2}}\right\|^2, \quad \alpha, \beta \geq 0, \alpha + \beta \leq 1, 0 \leq \gamma \leq 1, \end{aligned} \quad (6)$$

which is the non-Hermitian extension of the  $(\alpha, \beta, \gamma)$  WWYD skew information. Eq. (6) reduces to Eq. (10) in [23] when  $\gamma = \frac{1}{2}$ . When  $\beta = 1 - \alpha$ , we obtain the  $(\alpha, \gamma)$  *modified weighted Wigner-Yanase-Dyson* ( $(\alpha, \gamma)$  MWWYD) skew information,

$$\begin{aligned} K_{\rho,\gamma}^\alpha(E) &= -\frac{1}{2}\text{Tr}([(1-\gamma)\rho^\alpha + \gamma\rho^{1-\alpha}, E^\dagger][(1-\gamma)\rho^\alpha + \gamma\rho^{1-\alpha}, E]) \\ &= \frac{1}{2}\left\|[(1-\gamma)\rho^\alpha + \gamma\rho^{1-\alpha}, E]\right\|^2, \quad 0 \leq \alpha \leq 1, 0 \leq \gamma \leq 1, \end{aligned} \quad (7)$$

which is the non-Hermitian extension of the  $(\alpha, \gamma)$  WWYD skew information. It reduces to Eq. (3) when  $\gamma = \frac{1}{2}$ .

Quantum channels characterize the general evolutions of quantum systems [28, 29], which play an essential role in quantum information processing. The uncertainty relations for quantum channels have been investigated from both the variance-based and entropic-based uncertainty measure [30, 31]. Specifically, the unitary channels are useful and commonly encountered in both quantum information theory and quantum computation [28]. Uncertainty relations for general unitary channels have been investigated both theoretically and experimentally [32–34]. Recently, the sum uncertainty relations for quantum channels have attracted considerable attention [27, 35–38]. Fu, Sun and Luo [35] investigated the uncertainty relations for two quantum channels based on WY skew information for arbitrary operators. Afterwards, Zhang, Gao and Yan [36] generalized the uncertainty relations for two quantum channels to arbitrary  $N$  quantum channels and proposed tighter lower bounds than the ones in [35] for two quantum channels. Zhang, Wu and Fei [37] proposed new bounds which are tighter than the results in [36]. Cai [38] confirmed that the results in [35] also hold for all metric-adjusted skew information. By employing the norm inequalities proposed in [37], we have established sum uncertainty relations for arbitrary  $N$  quantum channels based on  $(\alpha, \beta, \gamma)$  MWWYD skew information [27].

Following the idea in [39], the  $(\alpha, \beta, \gamma)$  MWWYD skew information of a state  $\rho$  with respect to a channel  $\Phi$  has been defined as [27],

$$K_{\rho,\gamma}^{\alpha,\beta}(\Phi) = \sum_{i=1}^n K_{\rho,\gamma}^{\alpha,\beta}(E_i), \quad (8)$$

where  $\alpha, \beta \geq 0, \alpha + \beta \leq 1, 0 \leq \gamma \leq 1$ , and  $E_i (i = 1, 2, \dots, n)$  are Kraus operators of the channel  $\Phi$ , i.e.,  $\Phi(\rho) = \sum_{i=1}^n E_i \rho E_i^\dagger$ . Very recently, we provided the following uncertainty relations for arbitrary  $N$  quantum channels  $\{\Phi_t\}_{t=1}^N$  with  $\Phi_t(\rho) = \sum_{i=1}^n E_i^t \rho (E_i^t)^\dagger, t =$

$1, 2, \dots, N$  ( $N > 2$ ) [27],

$$\sum_{t=1}^N K_{\rho, \gamma}^{\alpha, \beta}(\Phi_t) \geq \max_{\pi_t, \pi_s \in S_n} \frac{1}{N-2} \left\{ \sum_{1 \leq t < s \leq N} \sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t + E_{\pi_s(i)}^s) - \frac{1}{(N-1)^2} \left[ \sum_{i=1}^n \left( \sum_{1 \leq t < s \leq N} \sqrt{K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t + E_{\pi_s(i)}^s)} \right)^2 \right] \right\}, \quad (9)$$

$$\sum_{t=1}^N K_{\rho, \gamma}^{\alpha, \beta}(\Phi_t) \geq \max_{\pi_t, \pi_s \in S_n} \left\{ \frac{1}{N} \sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta} \left( \sum_{t=1}^N E_{\pi_t(i)}^t \right) + \frac{2}{N^2(N-1)} \left[ \sum_{i=1}^n \left( \sum_{1 \leq t < s \leq N} \sqrt{K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t - E_{\pi_s(i)}^s)} \right)^2 \right] \right\}, \quad (10)$$

$$\sum_{t=1}^N K_{\rho, \gamma}^{\alpha, \beta}(\Phi_t) \geq \max_{\pi_t, \pi_s \in S_n} \frac{1}{2(N-1)} \left\{ \frac{2}{N(N-1)} \left[ \sum_{i=1}^n \left( \sum_{1 \leq t < s \leq N} \sqrt{K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t \pm E_{\pi_s(i)}^s)} \right)^2 \right] + \sum_{1 \leq t < s \leq N} \sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t \mp E_{\pi_s(i)}^s) \right\}, \quad (11)$$

where  $\alpha, \beta \geq 0$ ,  $\alpha + \beta \leq 1$ ,  $0 \leq \gamma \leq 1$ ,  $S_n$  is the  $n$ -element permutation group and  $\pi_t, \pi_s \in S_n$  are arbitrary  $n$ -element permutations.

The remainder of this paper is structured as follows. In Section 2, we explore the  $(\alpha, \beta, \gamma)$  MWWYD skew information-based sum uncertainty relations for arbitrary  $N$  quantum channels. Especially, we show that when  $\beta = 1 - \alpha$ , i.e., when the  $(\alpha, \beta, \gamma)$  MWWYD skew information becomes the  $(\alpha, \gamma)$  MWWYD skew information, our new bounds are tighter than the existing ones by a detailed example. The uncertainty relations based on the  $(\alpha, \beta, \gamma)$  MWWYD skew information for unitary channels are discussed in Section 3. We conclude with a summary in Section 4.

## 2. Sum uncertainty relations for arbitrary $N$ quantum channels in terms of $(\alpha, \beta, \gamma)$ MWWYD skew information

In this section, by using a new formation we explore the uncertainty relations for arbitrary  $N$  quantum channels in terms of the  $(\alpha, \beta, \gamma)$  MWWYD skew information  $K_{\rho, \gamma}^{\alpha, \beta}(\Phi)$ .

Let  $\Phi$  be a quantum channel with Kraus representation,  $\Phi(\rho) = \sum_{i=1}^n E_i \rho E_i^\dagger$ . Following the idea in [37], we define the  $(\alpha, \beta, \gamma)$  MWWYD skew information of the channel as,

$$K_{\rho, \gamma}^{\alpha, \beta}(\Phi) = \frac{1}{2} \text{Tr}(u^\dagger u) = \frac{1}{2} \|u\|^2, \quad (12)$$

where  $\alpha, \beta \geq 0, \alpha + \beta \leq 1, 0 \leq \gamma \leq 1, u = ([ (1 - \gamma)\rho^\alpha + \gamma\rho^\beta, E_1 ] \rho^{\frac{1-\alpha-\beta}{2}}, [(1 - \gamma)\rho^\alpha + \gamma\rho^\beta, E_2 ] \rho^{\frac{1-\alpha-\beta}{2}}, \dots, [(1 - \gamma)\rho^\alpha + \gamma\rho^\beta, E_n ] \rho^{\frac{1-\alpha-\beta}{2}})$  characterizes some intrinsic features of both the quantum state and the quantum channel. By employing operator norm inequalities and Eq. (12), we have the following theorem for arbitrary  $N$  quantum channels.

**Theorem 1** Let  $\Phi_1, \dots, \Phi_N$  be  $N$  quantum channels with Kraus representations  $\Phi_t(\rho) = \sum_{i=1}^n E_i^t \rho (E_i^t)^\dagger, t = 1, 2, \dots, N (N > 2)$ . We have

$$\sum_{t=1}^N K_{\rho, \gamma}^{\alpha, \beta}(\Phi_t) \geq \max\{LB1, LB2, LB3\}, \quad (13)$$

where

$$LB1 = \max_{\pi_t, \pi_s \in S_n} \frac{1}{N-2} \left\{ \sum_{1 \leq t < s \leq N} \sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t + E_{\pi_s(i)}^s) - \frac{1}{(N-1)^2} \left[ \sum_{1 \leq t < s \leq N} \sqrt{\sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t + E_{\pi_s(i)}^s)} \right]^2 \right\}, \quad (14)$$

$$LB2 = \max_{\pi_t, \pi_s \in S_n} \left\{ \frac{1}{N} \sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta} \left( \sum_{t=1}^N E_{\pi_t(i)}^t \right) + \frac{2}{N^2(N-1)} \left[ \sum_{1 \leq t < s \leq N} \sqrt{\sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t - E_{\pi_s(i)}^s)} \right]^2 \right\}, \quad (15)$$

$$LB3 = \max_{\pi_t, \pi_s \in S_n} \frac{1}{2(N-1)} \left\{ \sum_{1 \leq t < s \leq N} \sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t \pm E_{\pi_s(i)}^s) + \frac{2}{N(N-1)} \left[ \sum_{1 \leq t < s \leq N} \sqrt{\sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t \mp E_{\pi_s(i)}^s)} \right]^2 \right\}, \quad (16)$$

$\alpha, \beta \geq 0, \alpha + \beta \leq 1, 0 \leq \gamma \leq 1, S_n$  is the  $n$ -element permutation group and  $\pi_t, \pi_s \in S_n$  are arbitrary  $n$ -element permutations.

*Proof* The proof is completed directly by using the following inequalities [36, 37, 40],

$$\begin{aligned} \sum_{t=1}^N \|u_t\|^2 &\geq \frac{1}{N-2} \left[ \sum_{1 \leq t < s \leq N} \|u_t + u_s\|^2 - \frac{1}{(N-1)^2} \left( \sum_{1 \leq t < s \leq N} \|u_t + u_s\| \right)^2 \right], \\ \sum_{t=1}^N \|u_t\|^2 &\geq \frac{1}{N} \left\| \sum_{t=1}^N u_t \right\|^2 + \frac{2}{N^2(N-1)} \left( \sum_{1 \leq t < s \leq N} \|u_t - u_s\| \right)^2, \\ \sum_{t=1}^N \|u_t\|^2 &\geq \frac{1}{2(N-1)} \left[ \frac{2}{N(N-1)} \left( \sum_{1 \leq t < s \leq N} \|u_t \pm u_s\| \right)^2 + \sum_{1 \leq t < s \leq N} \|u_t \mp u_s\|^2 \right], \end{aligned}$$

with  $\|u_t\|^2 = 2K_{\rho, \gamma}^{\alpha, \beta}(\Phi_t)$ ,  $\|u_t + u_s\|^2 = 2 \sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t + E_{\pi_s(i)}^s)$  and  $\|u_t - u_s\|^2 = 2 \sum_{i=1}^n K_{\rho, \gamma}^{\alpha, \beta}(E_{\pi_t(i)}^t - E_{\pi_s(i)}^s)$ .  $\square$

Note that when  $\alpha = \beta = \frac{1}{2}$ , Theorem 1 reduce to Theorem 1 in [37]. As a special case, we use the  $(\alpha, \gamma)$  MWWYD skew information to compare our lower bounds with the existing ones. For convenience, we denote by  $\overline{LB1}$ ,  $\overline{LB2}$ ,  $\overline{LB3}$  the right hand sides of (9), (10) and (11), respectively. The following example shows that our results give tighter lower bounds than  $\overline{LB1}$ ,  $\overline{LB2}$  and  $\overline{LB3}$ , see Figure 1.

**Example 1** Given a qubit state  $\rho = \frac{1}{2}(\mathbf{1} + \mathbf{r} \cdot \boldsymbol{\sigma})$ , where  $\mathbf{1}$  is the  $2 \times 2$  identity matrix,  $\mathbf{r} = (\frac{\sqrt{3}}{2} \cos \theta, \frac{\sqrt{3}}{2} \sin \theta, 0)$ ,  $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$  with  $\sigma_j$  ( $j = 1, 2, 3$ ) the Pauli matrices, and  $\mathbf{r} \cdot \boldsymbol{\sigma} = \sum_{j=1}^3 r_j \sigma_j$ . We consider the following three quantum channels:

(i) the amplitude damping channel  $\Phi_{AD}$ ,

$$\Phi_{AD}(\rho) = \sum_{i=1}^2 A_i \rho A_i^\dagger, \quad A_1 = |0\rangle\langle 0| + \sqrt{1-q}|1\rangle\langle 1|, \quad A_2 = \sqrt{q}|1\rangle\langle 1|;$$

(ii) the phase damping channel  $\Phi_{PD}$ ,

$$\Phi_{PD}(\rho) = \sum_{i=1}^2 B_i \rho B_i^\dagger, \quad B_1 = |0\rangle\langle 0| + \sqrt{1-q}|1\rangle\langle 1|, \quad B_2 = \sqrt{q}|0\rangle\langle 1|;$$

(iii) the bit flip channel  $\Phi_{BF}$ ,

$$\Phi_{BF}(\rho) = \sum_{i=1}^2 C_i \rho C_i^\dagger, \quad C_1 = \sqrt{q}|0\rangle\langle 0| + \sqrt{q}|1\rangle\langle 1|, \quad C_2 = \sqrt{1-q}(|0\rangle\langle 1| + |1\rangle\langle 0|)$$

with  $0 \leq q < 1$ , respectively.

For the case  $\alpha = \gamma = \frac{1}{4}$ ,  $q = 0.2$  and  $\theta = \frac{\pi}{2}$ , we have  $K_{\rho, \frac{1}{4}}^{\frac{1}{4}}(\Phi_{AD}) + K_{\rho, \frac{1}{4}}^{\frac{1}{4}}(\Phi_{PD}) + K_{\rho, \frac{1}{4}}^{\frac{1}{4}}(\Phi_{BF}) = 0.283955$ . The lower bounds  $\overline{LB1}$ ,  $\overline{LB2}$  and  $\overline{LB3}$  are 0.275596, 0.2644 and

0.256419, respectively, and the lower bounds  $LB1$ ,  $LB2$  and  $LB3$  are 0.260707, 0.26726 and 0.265758, respectively. Obviously,  $LB2$  is tightest among  $LB1$ ,  $LB2$  and  $LB3$ , which is also greater than  $\overline{LB2}$  and  $\overline{LB3}$  given in [27].

We also consider the case  $\alpha = \gamma = \frac{1}{4}$ . For  $q = 0.4$  and  $q = 0.9$ , the sum and the lower bounds  $\overline{LB1}$ ,  $\overline{LB2}$ ,  $\overline{LB3}$ ,  $LB1$ ,  $LB2$  and  $LB3$  are shown in Figure 1, respectively. Especially, for  $q = 0.4$ , the sum and the lower bounds are calculated for some special  $\theta$ , as listed in Table 1. It can be seen that for  $q = 0.4$ , our lower bounds  $LB2$  and  $LB3$  are tighter than  $\overline{LB1}$ ,  $\overline{LB2}$  and  $\overline{LB3}$ . While for  $q = 0.9$ , our lower bounds  $LB2$  and  $LB3$  are tighter than  $\overline{LB1}$ ,  $\overline{LB2}$  and  $\overline{LB3}$ .




Figure 1: The solid black line represents the sum  $= K_{\rho, \frac{1}{4}}^{\frac{1}{4}}(\Phi_{AD}) + K_{\rho, \frac{1}{4}}^{\frac{1}{4}}(\Phi_{PD}) + K_{\rho, \frac{1}{4}}^{\frac{1}{4}}(\Phi_{BF})$ . The solid blue, green and the red lines represent the lower bounds  $LB1$ ,  $LB2$  and  $LB3$  in Theorem 1, respectively. The dotted magenta, dashed blue and green lines are for the lower bounds  $\overline{LB1}$ ,  $\overline{LB2}$  and  $\overline{LB3}$ , respectively. (a)  $q = 0.4$ ; (b)  $q = 0.9$ .

Table 1. Comparison among the uncertainty lower bounds

q=0.4	$\overline{LB1}$	$\overline{LB2}$	$\overline{LB3}$	$LB1$	$LB2$	$LB3$	sum
$\theta = \pi/2$	0.234918	0.247658	0.241686	0.222065	0.252565	0.252654	0.258817
$\theta = \pi/3$	0.17968	0.204421	0.20082	0.168362	0.208841	0.208534	0.211782
$\theta = \pi/5$	0.0954994	0.13303	0.132687	0.0879256	0.135648	0.135459	0.135679
$\theta = \pi/7$	0.066361	0.104405	0.104922	0.0632504	0.106043	0.106062	0.106096

The above results show that Theorem 1 in this paper improve the existing results ones given in [27].

### 3. Sum uncertainty relations for $N$ unitary channels in terms of $(\alpha, \beta, \gamma)$ MWWYD skew information

In this section, we consider sum uncertainty relations for arbitrary  $N$  unitary channels. For a unitary channel  $\Phi_U$ , we have  $\Phi_U(\rho) = U\rho U^\dagger$ . According to Eq. (6), the  $(\alpha, \beta, \gamma)$

MWWYD skew information of an unitary operator  $U$  is given by

$$\begin{aligned} K_{\rho,\gamma}^{\alpha,\beta}(U) &= -\frac{1}{2}\text{Tr}([(1-\gamma)\rho^\alpha + \gamma\rho^\beta, U^\dagger][(1-\gamma)\rho^\alpha + \gamma\rho^\beta, U]\rho^{1-\alpha-\beta}) \\ &= \frac{1}{2}\left\| \left[ (1-\gamma)\rho^\alpha + \gamma\rho^\beta, U \right] \rho^{\frac{1-\alpha-\beta}{2}} \right\|^2, \quad \alpha, \beta \geq 0, \alpha + \beta \leq 1, 0 \leq \gamma \leq 1. \end{aligned} \quad (17)$$

The  $(\alpha, \beta, \gamma)$  MWWYD skew information of a unitary channel  $\Phi_U$  is defined as  $K_{\rho,\gamma}^{\alpha,\beta}(\Phi_U) = K_{\rho,\gamma}^{\alpha,\beta}(U)$ . For simplicity, in the following, we use  $K_{\rho,\gamma}^{\alpha,\beta}(U)$  to denote the quantity of the unitary channel  $\Phi_U$  determined by  $U$ . Similar to the proof of Theorem 1, we can prove the following theorem.

**Theorem 2** Let  $U_1, \dots, U_N$  be arbitrary  $N$  unitary operators. Then we have

$$\sum_{t=1}^N K_{\rho,\gamma}^{\alpha,\beta}(U_t) \geq \max\{Lb1, Lb2, Lb3\}, \quad (18)$$

where

$$Lb1 = \frac{1}{N-2} \left\{ \sum_{1 \leq t < s \leq N} K_{\rho,\gamma}^{\alpha,\beta}(U_t + U_s) - \frac{1}{(N-1)^2} \left[ \sum_{1 \leq t < s \leq N} \sqrt{K_{\rho,\gamma}^{\alpha,\beta}(U_t + U_s)} \right]^2 \right\}, \quad (19)$$

$$Lb2 = \frac{1}{N} K_{\rho,\gamma}^{\alpha,\beta} \left( \sum_{t=1}^N U_t \right) + \frac{2}{N^2(N-1)} \left[ \sum_{1 \leq t < s \leq N} \sqrt{K_{\rho,\gamma}^{\alpha,\beta}(U_t - U_s)} \right]^2, \quad (20)$$

$$\begin{aligned} Lb3 &= \max_{x \in \{0,1\}} \frac{1}{2(N-1)} \left\{ \sum_{1 \leq t < s \leq N} K_{\rho,\gamma}^{\alpha,\beta}(U_t + (-1)^x U_s) \right. \\ &\quad \left. + \frac{2}{N(N-1)} \left[ \sum_{1 \leq t < s \leq N} \sqrt{K_{\rho,\gamma}^{\alpha,\beta}(U_t + (-1)^{x+1} U_s)} \right]^2 \right\} \end{aligned} \quad (21)$$

and  $x \in \{0, 1\}$ ,  $\alpha, \beta \geq 0$ ,  $\alpha + \beta \leq 1$ ,  $0 \leq \gamma \leq 1$ .

Note that (19), (20) and (21) of Theorem 2 reduce to (13), (14) and (15) in [37] when  $\alpha = \gamma = \frac{1}{2}$ , respectively.

**Example 2** Given a qubit state  $\rho = \frac{1}{2}(\mathbf{1} + \mathbf{r} \cdot \boldsymbol{\sigma})$  with  $\mathbf{r} = (\frac{\sqrt{2}}{2} \cos \theta, \frac{\sqrt{2}}{2} \sin \theta, 0)$ . Consider the following three unitary operators,

$$U_1 = e^{\frac{i\pi\sigma_1}{8}} = \begin{pmatrix} \cos \frac{\pi}{8} & i \sin \frac{\pi}{8} \\ i \sin \frac{\pi}{8} & \cos \frac{\pi}{8} \end{pmatrix}, U_2 = e^{\frac{i\pi\sigma_2}{8}} = \begin{pmatrix} \cos \frac{\pi}{8} & \sin \frac{\pi}{8} \\ -\sin \frac{\pi}{8} & \cos \frac{\pi}{8} \end{pmatrix}, U_3 = e^{\frac{i\pi\sigma_3}{8}} = \begin{pmatrix} e^{i\frac{\pi}{8}} & 0 \\ 0 & -e^{i\frac{\pi}{8}} \end{pmatrix},$$



which correspond to the rotations around the  $z$  axis of the Bloch sphere. When  $\beta = 1 - \alpha$ , i.e., when the  $(\alpha, \beta, \gamma)$  MWWYD skew information reduces to the  $(\alpha, \gamma)$  MWWYD skew information, the comparison among the lower bounds of Theorem 2 is presented in Figure 2, from which one sees that the lower bound  $Lb3$  is tighter than  $Lb2$  and  $Lb1$  in this case.




Figure 2: For  $\alpha = \gamma = \frac{1}{4}$ , the solid black curve represents the sum  $= K_{\rho, \frac{1}{4}}^{\frac{1}{4}}(U_1) + K_{\rho, \frac{1}{4}}^{\frac{1}{4}}(U_2) + K_{\rho, \frac{1}{4}}^{\frac{1}{4}}(U_3)$ . The dotted blue, dashed green and dot-dashed red curves represent  $Lb1$ ,  $Lb2$  and  $Lb3$ , respectively.

#### 4. Conclusions

We have studied the sum uncertainty relations for  $N$  quantum channels based on the  $(\alpha, \beta, \gamma)$  MWWYD skew information. By detailed example it has been shown that our uncertainty inequalities are tighter than the existing ones. Since the MWWYD skew information and  $(\alpha, \gamma)$  MWWYD skew information are two special cases of the  $(\alpha, \beta, \gamma)$  MWWYD skew information, our results are also valid for the MWWYD skew information and the  $(\alpha, \gamma)$  MWWYD skew information. Finally, we have also explored sum uncertainty relations for unitary channels. These results may shed some new light on the study of skew information-based sum uncertainty relations for quantum channels.

#### Acknowledgements

This work was supported by National Natural Science Foundation of China (Grant Nos. 12161056, 12075159, 12171044); Jiangxi Provincial Natural Science Foundation (Grant No. 20202BAB201001); Beijing Natural Science Foundation (Grant No. Z190005); Academy for Multidisciplinary Studies, Capital Normal University; Shenzhen Institute for Quantum Science and Engineering, Southern University of Science and Technology (Grant No. SIQSE202001); the Academician Innovation Platform of Hainan Province.

## Conflict of interest

The authors declare that they have no conflict of interest.

## References

- [1] Heisenberg W 1927 Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik *Z. Phys.* **43** 172
- [2] Robertson H P 1929 The uncertainty principle *Phys. Rev.* **34** 163
- [3] Deutsch D 1983 Uncertainty in quantum measurements *Phys. Rev. Lett.* **50** 631
- [4] Maassen H and Uffink J B M 1988 Generalized entropic uncertainty relations *Phys. Rev. Lett.* **60** 1103
- [5] Wehner S and Winter A 2010 Entropic uncertainty relations-a survey *New J. Phys.* **12** 025009
- [6] Wu S, Yu S and Mølmer K 2009 Entropic uncertainty relation for mutually unbiased bases *Phys. Rev. A* **79** 022104
- [7] Rastegin A E 2013 Uncertainty relations for MUBs and SIC-POVMs in terms of generalized entropies *Eur. Phys. J. D* **67** 269
- [8] Gudder S 2007 Operator probability theory *Int. J. Pure Appl. Math.* **39** 511
- [9] Dou Y and Du H 2013 Generalizations of the Heisenberg and Schrödinger uncertainty relations *J. Math. Phys.* **54** 103508
- [10] Dou Y and Du H 2014 Note on the Wigner-Yanase-Dyson skew information *Int. J. Theor. Phys.* **53** 952
- [11] Sun Y and Li N 2021 The uncertainty of quantum channels in terms of variance *Quantum Inf. Process.* **20** 25
- [12] Rudnicki L, Puchała Z and Życzkowski K 2014 Strong majorization entropic uncertainty relations *Phys. Rev. A* **89** 052115
- [13] Puchała Z, Rudnicki L and Życzkowski K 2013 Majorization entropic uncertainty relations *J. Phys. A: Math. Theor.* **46** 272002
- [14] Rudnicki L 2015 Majorization approach to entropic uncertainty relations for coarse-grained observables *Phys. Rev. A* **91** 032123
- [15] Friedland S, Gheorghiu V and Gour G 2013 Universal uncertainty relations *Phys. Rev. Lett.* **111** 230401
- [16] Wigner E P and Yanase M M 1963 Information contents of distributions *Proc. Natl. Acad. Sci.* **49** 910-918
- [17] Chen P and Luo S 2007 Direct approach to quantum extensions of Fisher information *Front. Math.* **2** 359
- [18] Luo S and Zhang Q 2004 On skew information *IEEE Trans. Inf. Theory* **50** 1778

- [19] Cai L and Luo S 2008 On convexity of generalized Wigner-Yanase-Dyson information *Lett. Math. Phys.* **83** 253
- [20] Yanagi K 2010 Uncertainty relation on Wigner-Yanase-Dyson skew information *J. Math. Anal. Appl.* **365** 12
- [21] Yanagi K 2010 Wigner-Yanase-Dyson skew information and uncertainty relation *J. Phys. Conf. Ser.* **201** 012015
- [22] Wu Z, Zhang L, Fei S-M and Li-Jost X 2020 Coherence and complementarity based on modified generalized skew information *Quantum Inf. Process.* **19** 154
- [23] Wu Z, Zhang L, Wang J, Li-Jost X, and Fei S-M 2020 Uncertainty relations based on modified Wigner-Yanase-Dyson skew information *Int. J. Theor. Phys.* **59** 704
- [24] Furuichi S, Yanagi K and Kuriyama K 2009 Trace inequalities on a generalized Wigner-Yanase skew information *J. Math. Anal. Appl.* **356** 179
- [25] Chen Z, Liang L, Li H and Wang W 2016 Two generalized Wigner-Yanase skew information and their uncertainty relations *Quantum Inf. Process.* **15** 5107
- [26] Zhang Z 2021 Trace inequalities based on two-parameter extended Wigner-Yanase skew information *J. Math. Anal. Appl.* **497** 124851
- [27] Xu C, Wu Z and Fei S-M 2022 Sum uncertainty relations based on  $(\alpha, \beta, \gamma)$  weighted Wigner-Yanase-Dyson skew information *Int. J. Theor. Phys.* **61** 185
- [28] Nielsen M A and Chuang I L 2010 *Quantum Computation and Quantum Information* (Cambridge: Cambridge University Press)
- [29] Busch P, Grabowski M and Lahti P 1997 *Operational Quantum Physics* 2nd edn (Berlin: Springer)
- [30] Krishna M and Parthasarathy K R 2002 An entropic uncertainty principle for quantum measurements *Sankhyā A* **64** 842
- [31] Massar S 2008 Uncertainty relations for positive-operator-valued measures *Phys. Rev. A* **76** 042114
- [32] Bagchi S and Pati A K 2016 Uncertainty relations for general unitary operators *Phys. Rev. A* **94** 042104
- [33] Tajima H, Shiraishi N and Saito K 2018 Uncertainty relations in implementation of unitary operations *Phys. Rev. Lett.* **121** 110403
- [34] Bong K-W, Tischler N, Patel R B, Wollmann S, Pryde G J and Hall M J W 2018 Strong unitary and overlap uncertainty relations: theory and experiment *Phys. Rev. Lett.* **120** 230402
- [35] Fu S, Sun Y and Luo S 2019 Skew information-based uncertainty relations for quantum channels *Quantum Inf. Process.* **18** 258
- [36] Zhang L, Gao T and Yan F 2021 Tighter uncertainty relations based on Wigner-Yanase skew information for observables and channels *Phys. Lett. A* **387** 127029

- [37] Zhang Q, Wu J and Fei S-M 2021 A note on uncertainty relations of arbitrary  $N$  quantum channels *Laser Phys. Lett.* **18** 095204
- [38] Cai L 2021 Sum uncertainty relations based on metric-adjusted skew information *Quantum Inf. Process.* **20** 72
- [39] Luo S and Sun Y 2018 Coherence and complementarity in state-channel interaction *Phys. Rev. A.* **98** 012113
- [40] Chen B and Fei S-M 2015 Sum uncertainty relations for arbitrary  $N$  incompatible observables *Sci. Rep.* **5** 14238