Impact of the ligand deformation on the \mathcal{P} , \mathcal{T} -violation effects in the YbOH molecule

Anna Zakharova^{1, 2, a)} and Alexander $Petrov^{1, 2, b)}$

¹⁾St. Petersburg State University, St. Petersburg, 7/9 Universitetskaya nab., 199034, Russia

²⁾ Petersburg Nuclear Physics Institute named by B.P. Konstantinov of National Research Centre "Kurchatov Institute", Gatchina, 1, mkr. Orlova roshcha, 188300, Russia

(Dated: 16 August 2022)

The ytterbium monohydroxide is a promising molecule for a new physics searches. It is well known that levels of the opposite parity, separated by the energy split, so-called *l*-doublets, define the experimental electric field strength required for the molecule polarization. In addition, in our previous paper¹ we have shown that the value of *l*-doubling directly influences the sensitivity of linear triatomic molecules to the \mathcal{P},\mathcal{T} -odd effects. In our work² we have calculated the value of *l*-doubling for the YbOH molecule with approximation of fixed O-H bond length. Accounting the importance of this property, in the present study, we consider the additional degree of freedom corresponding to the ligand (OH) deformation.

I. INTRODUCTION

Symmetry breaking with respect to the spatial reflection (\mathcal{P}) , the time inversion (\mathcal{T}) , and the charge conjugation (\mathcal{C}) are embedded in the Standard model $(SM)^{3,4}$. One of the possible sources of the strong \mathcal{P}, \mathcal{T} -violation in the SM is θ term, that however is negligible^{5,6}. The observed source of the \mathcal{P} and \mathcal{C} nonconservation comes from the weak interaction, that affects only the left components of the lepton and quark spinors. This fact means that \mathcal{P} , and \mathcal{C} symmetries are broken separately⁷. However, the combined \mathcal{CP} parity could be conserved. But in SM there is a Cabibbo-Kobayashi-Maskawa (CKM)^{8,9} matrix, which is associated with the interaction of quarks and W^{\pm} bosons, and Pontecorvo-Maki-Nakagawa-Sakata matrix $(PMNS)^{10,11}$, which is associated with the interaction of leptons with W^{\pm} -bosons. When these matrices have complex components, they violate \mathcal{CP} , which is manifested in neutral kaon and B^0 -meson decays³. Because of the \mathcal{CPT} theorem, \mathcal{CP} combination is equivalent to time reversal \mathcal{T} .

The probable effect of the \mathcal{CP} -symmetry violation in the Standard model can be found in the electron electric dipole moment (eEDM) and scalar-pseudoscalar electron-nucleon interaction (S-PS)¹², parametrized correspondingly by d_e and k_s constants. In the SM the predicted values for these constants are very small. Nevertheless, there are many extensions of the SM that can lead to the considerable increase of the d_e and k_s constants^{13–16}.

To investigate the CP-violating physics, tabletop experiments with atoms and molecules can be employed^{17–24}. The current limit on the electron electric dipole moment (the ACME II experiment), $|d_e| <$ 1.1×10^{-29} e·cm (90% confidence), was set by measuring the spin precession using thorium monoxide (ThO) molecules in the metastable electronic H³ Δ_1 state¹⁹.

In its turn, cold polar molecules provide unique opportunities for further progress in search for effects of symmetry violation²⁵. The strength of polyatomic spices as probes of the parity-violating physics lies in a possibility of laser-cooling that does not interfere with the existence of the parity doublets²⁶. Unlike the diatomic molecules, polyatomic ones possess the distinctive rovibrational spectrum with additional energy levels of opposite parity. The transverse vibrations of triatomic molecules result in *l*-doublets^{27,28}. For symmetric top molecules nonvibrational *K*-doublets, associated with rotation of the molecule around its axis, appear^{27,29}.

One dimensional laser-cooling was made for SrOH, CaO, CaOCH₃³⁰⁻³², and the YbOH^{33,34} molecules. Recently, the magneto-optical trapping and cooling of the CaOH molecule was achieved³⁵. The molecule we are considering, ytterbium monohydroxide, is a good candidate for the eEDM search^{1,2,27}.

For triatomic molecules, the external fields are mixing the opposite parity levels, and due to this, the molecule is polarized to a degree determined by a coefficient P^1 . If the electron has eEDM and interacts with the nucleus by the S-PS, the \mathcal{P} , \mathcal{T} -nonconservation emerges in the energy difference between levels with opposite projection of the total angular momentum on the field's axis:

$$\Delta E_{\mathcal{P},\mathcal{T}} = P(2E_{\text{eff}}d_e + 2E_{\text{s}}k_s),\tag{1}$$

Knowing the enhancement coefficients E_{eff} , E_{s} and P one may extract the value of the constants d_e and k_s from this energy splitting^{36–38}.

Recently, we have calculated the enhancement factors $E_{\rm eff}$ and $E_{\rm s}$ for the RaOH, YbOH, and the symmetric top molecule RaOCH₃, the rovibrational wavefunctions were obtained^{2,29,39} on the CCSD level. Our computational approach, that does take into account the rotational and anharmonic effects, allowed us to calculate *l*-doubling for the triatomic molecules RaOH and YbOH. We have

^{a)}Electronic mail: zakharova.annet@gmail.com

^{b)}Electronic mail: petrov_an@pnpi.nrcki.ru

also determined the value of the polarization coefficient P for the YbOH molecule¹. In our work¹ we have shown that the *l*-doubling structure is, in general, different from Ω -doubling of diatomics, and the P value tends to approach 50% value for most of the ground rotational levels of the first excited $\nu_2 = 1$ bending mode of YbOH. For some levels a maximum of the polarization P^{\max} value as a function of electric field with $1/2 < |P|^{\max} < 1$ is observed. The smaller the *l*-doubling, the larger is the $|P|^{\max,1}$. Thus, the value of *l*-doubling directly influences the sensitivity of linear triatomic molecules to the \mathcal{P},\mathcal{T} -odd effects.

However, when calculating l-doubling, in our preceding work² we assumed that the deformations of the ligand may be neglected as the corresponding vibrational frequency is much higher than the frequencies for bending and stretching of the whole molecule. The aim of the present paper is to take into account the change of the ligand's bond length (i.e., of the equilibrium ligand size and its vibrational frequency) due to the interaction with bending and stretching modes of the YbOH.

II. POTENTIAL SURFACE INTERPOLATION

To obtain the potential surface we used the Dirac 19 program suite⁴⁰. The PNPI⁴¹ 42-valence electron basis was used to describe the Ytterbium atom. Moreover, to simplify calculations with a heavy atom, we employed the 28-electron generalized relativistic effective potential, also developed by the PNPI Quantum Chemistry Laboratory^{42–44}. For oxygen and hydrogen atoms, we used cc-pVTZ basis sets. Active space for the coupled cluster computations comprises 21 active and 30 frozen electrons.

We obtain the CCSD and CCSD(T) potential surface V on the grid of coordinates (R_i, r_j, θ_k) ,

$$\{R_i\} = 3.3, 3.5, \dots 4.3 \text{ a.u.}$$
 (2)

$$\{r_j\} = 1.632, 1.732, \dots 2.032 \text{ a.u.}$$
 (3)

$$\{\theta_k\} = 0^{\circ}, 5^{\circ}, 10^{\circ}, 15^{\circ}, 20^{\circ}, 25^{\circ}, 55^{\circ}, 90^{\circ}, 122^{\circ}, 155^{\circ} \quad (4)$$

At the first step we approximate the dependence on r for each (R_i, θ_k) by the Morse potential

$$V(r) = V_0 + D(1 - e^{-\alpha(r - r^{eq})})^2$$
(5)

where r^{eq} is the equilibrium ligand size, D – dissociation energy, α is the exponent parameter, and V_0 is the energy of the minimum. The values for V_0 and r^{eq} parameters for various configurations of YbOH are presented on Fig. 1 and Fig. 2 correspondingly.

At the second step, for each R_i we interpolate the dependence of this set of parameters $V_0(R_i, \theta_k)$, $r^{eq}(R_i, \theta_k)$, $D(R_i, \theta_k)$, and $\alpha(R_i, \theta_k)$, on θ by the Akima splines.

Then, using Akima splines, obtained at the previous step, we take the values at the zeros of the $(\lambda_{max} + 1)$ -th Legendre polynomial $P_{\lambda_{max}+1}(x)$, where $x = \cos \theta$. For

FIG. 1. Morse potential parameter V_0 as a function of R and θ coordinates



each R_i , we then use the Gauss-Legendre method to rep-





resent the parameters as sums of Legendre polynomials,

$$V_0(R_i,\theta) = \sum_{\lambda=0}^{\lambda_{max}} V_{0,\lambda}(R_i) P_\lambda(\cos\theta), \qquad (6)$$

$$D(R_i, \theta) = \sum_{\lambda=0}^{\lambda_{max}} D_{\lambda}(R_i) P_{\lambda}(\cos \theta), \qquad (7)$$

$$r^{eq}(R_i,\theta) = \sum_{\lambda=0}^{\lambda_{max}} r_{\lambda}^{eq}(R_i) P_{\lambda}(\cos\theta), \qquad (8)$$

$$\alpha(R_i,\theta) = \sum_{\lambda=0}^{\lambda_{max}} \alpha_\lambda(R_i) P_\lambda(\cos\theta), \qquad (9)$$

In this paper we have taken $\lambda_{max} = 40$. Such a representation is guaranteed to be smooth and have good periodicity in θ .

At the final step, we interpolate the dependence of the coefficients of the Legendre polynomials on R using Akima's splines.

III. METHODS

We use the Born-Oppenheimer approximation to separate wavefunction to Ψ_{nuc} , responsible for the nuclear motion in the adiabatic potential of the electrons, and the Ψ_{elec} , corresponding to the electronic motion in the field of heavy nuclei

$$\Psi_{\text{total}} \approx \Psi_{\text{nuc}}(\vec{R}, \vec{r}) \psi_{\text{elec}}(\vec{R}, \vec{r}|q), \qquad (10)$$

where q are the generalized coordinates of the electrons, \vec{R} is the vector from the Ytterbium atom to the center of mass of the OH ligand, \vec{r} is the vector from the Oxygen to the Hydrogen atom.

The Hamiltonian of nuclei takes the form,

$$\hat{H}_{\text{nuc}} = -\frac{1}{2\mu} \frac{\partial^2}{\partial R^2} - \frac{1}{2\mu_{OH}} \frac{\partial^2}{\partial r^2} + \frac{\hat{L}^2}{2\mu R^2} + \frac{\hat{j}^2}{2\mu_{OH}r^2} + V(R, r, \theta)$$
(11)

where θ is the angle between vectors \vec{R} and \vec{r} (so that $\theta = 0$ corresponds to the Yb-O-H linear configuration), μ is the Yb-OH reduced mass, μ_{OH} is the reduced mass of the ligand, \hat{L} and \hat{j} are the angular momentum of the whole system rotation and the OH angular momentum correspondingly. Directions of axes \hat{r} and \hat{R} are represented in Fig. 3. V is the electronic potential surface obtained on the CCSD or CCSD(T) level and depending only on the relative Jacobi coordinates (R, r, θ) .

The nuclear wavefunction $\Psi_{nuc}(\vec{R}, \vec{r})$ is the solution of the Schrödinger equation

$$\hat{H}_{nuc}\Psi_{nuc}(\vec{R},\vec{r}) = E\Psi_{nuc}(\vec{R},\vec{r}).$$
(12)

To solve eq. (12) we use the expansion

$$\Psi_{\rm nuc}(\vec{R},\vec{r}) = \sum_{L=0}^{L_{max}} \sum_{j=0}^{j_{max}} \sum_{n=1}^{n_{max}} F_{JjLn}(R) \Phi_{JjLM}(\hat{R},\hat{r}) f_n(r),$$
(13)

where

$$\Phi_{JjLM}(\hat{R}, \hat{r}) = \sum_{m_L, m_j} C_{Lm_L, jm_j}^{JM} Y_{Lm_L}(\hat{R}) Y_{jm_j}(\hat{r}) \quad (14)$$

is coupled to conserved total angular momentum J basis set, Y_{Lm_L} is a spherical function, $f_n(r)$ is a solution of

$$\left(-\frac{1}{2\mu_{OH}}\frac{\partial^2}{\partial r^2} + V(R_i, r, \theta_i)\right)f_n(r) = e_n f_n(r), \quad (15)$$

where R_i and θ_i are some fixed values. To test the approximation (13) we perform our computations for different sets of (R_i, θ_i) .

Substituting wavefunction (13) to eq. (12) one gets the system of close-coupled equations for $F_{JjLn}(R)^{45}$.



FIG. 3. Jacobi coordinates

IV. RESULTS

The parameters obtained from the spectrum of the nu-), clear wavefunctions Ψ_{nuc} compared with the results from the rigid ligand approximation and experimental values are presented in the Table I. One can see that taking into account the ligand deformation improves the agreement with experiments. To test our method and codes, which now take into account the ligand deformation, we performed calculations with two different sets of the parameters R_i and θ_i from eq.(15). The Set 1 corresponds to $R_1 = 3.9$ a.u, $\theta_1 = 0$ and and Set 2 corresponds to $R_2 = 4.2$ a.u, $\theta_1 = 0.5$ radians. In Table II one can see that results with Set 1 and Set 2 converge to each other as quantum number n_{max} (see eq. (13)) increases. Table II also show that for purposes of the present paper it is sufficient to use $n_{max} = 2$. Probably inclusion of the noniterative triple excitations to the coupled cluster computations and using more extensive basis sets are necessary to reach the accuracy of the calculation on the order of $\sim 5 \,\mathrm{cm}^{-1}$ for the vibrational frequencies, what is out of scope of the current work.

However, to estimate the actual value of the l-doubling one can note the following. The decrease of the l-doubling value together with the increase of the ν_2 frequency is consistent with the estimate⁴⁶,

$$q \simeq \frac{B^2}{\nu_2} \Big(1 + 4 \sum_{k=1,3} \frac{\zeta_{2k}^2 \nu_2^2}{\nu_k^2 - \nu_2^2} \Big) (v+1), \quad \Delta E = 2q, \quad (16)$$

where ζ_{2k} are Coriolis coefficients. As the current experimental data for the ν_2 frequency fall into a rather wide range 319 - 339, cm⁻¹, we sum up our results for *l*-doubling with the estimate of $\Delta E_{J=1} \simeq 24 - 26$ MHz.

ACKNOWLEDGMENTS

The work is supported by the Russian Science Foundation grant No. 18-12-00227.

AUTHOR DECLARATIONS

Conflict of interest

The authors have no conflicts to disclose.

AVAILABILITY OF DATA

The data that support the findings of this study are available from the corresponding author upon reasonable request.

- ¹A. Petrov and A. Zakharova, "Sensitivity of the yboh molecule to p t-odd effects in an external electric field," Physical Review A **105**, L050801 (2022).
- ²A. Zakharova, I. Kurchavov, and A. Petrov, "Rovibrational structure of the ytterbium monohydroxide molecule and the p, t-violation searches," The Journal of Chemical Physics **155**, 164301 (2021).
- ³M. D. Schwartz, *Quantum field theory and the standard model* (Cambridge University Press, 2014).
- ⁴Particle Data Group, P. Zyla, R. Barnett, J. Beringer, O. Dahl, D. Dwyer, D. Groom, C.-J. Lin, K. Lugovsky, E. Pianori, *et al.*, "Review of particle physics," Progress of Theoretical and Experimental Physics **2020**, 083C01 (2020).
- ⁵H.-Y. Cheng, "The strong CP problem revisited," Physics Reports **158**, 1–89 (1988).
- ⁶J. E. Kim and G. Carosi, "Axions and the Strong CP Problem," Rev. Mod. Phys. **82**, 557–602 (2010), [Erratum: Rev.Mod.Phys. 91, 049902 (2019)], arXiv:0807.3125 [hep-ph].
- ⁷I. B. Khriplovich and S. K. Lamoreaux, *CP* violation without strangeness: electric dipole moments of particles, atoms, and molecules (Springer Science & Business Media, 2012).
- ⁸N. Cabibbo, "Unitary Symmetry and Leptonic Decays," Phys. Rev. Lett. **10**, 531–533 (1963).
- ⁹M. Kobayashi and T. Maskawa, "CP Violation in the Renormalizable Theory of Weak Interaction," Prog. Theor. Phys. 49, 652–657 (1973).
- ¹⁰B. Pontecorvo, "Inverse beta processes and nonconservation of lepton charge," Zh. Eksp. Teor. Fiz. **34**, 247 (1957).
- ¹¹Z. Maki, M. Nakagawa, and S. Sakata, "Remarks on the unified model of elementary particles," Prog. Theor. Phys. 28, 870–880 (1962).

- ¹²S. M. Barr, "T-and p-odd electron-nucleon interactions and the electric dipole moments of large atoms," Physical Review D 45, 4148 (1992).
- ¹³T. FÙKUYAMA, "SEARCHING FOR NEW PHYSICS BE-YOND THE STANDARD MODEL IN ELECTRIC DIPOLE MOMENT," International Journal of Modern Physics A 27, 1230015 (2012).
- ¹⁴M. Pospelov and A. Ritz, "CKM benchmarks for electron electric dipole moment experiments," Phys. Rev. D 89, 056006 (2014), arXiv:1311.5537 [hep-ph].
- ¹⁵Y. Yamaguchi and N. Yamanaka, "Large long-distance contributions to the electric dipole moments of charged leptons in the standard model," Phys. Rev. Lett. **125**, 241802 (2020), arXiv:2003.08195 [hep-ph].
- ¹⁶Y. Yamaguchi and N. Yamanaka, "Quark level and hadronic contributions to the electric dipole moment of charged leptons in the standard model," Phys. Rev. D **103**, 013001 (2021), arXiv:2006.00281 [hep-ph].
- ¹⁷J. Baron, W. C. Campbell, D. DeMille, J. M. Doyle, G. Gabrielse, Y. V. Gurevich, P. W. Hess, N. R. Hutzler, E. Kirilov, I. Kozyryev, *et al.*, "Order of magnitude smaller limit on the electric dipole moment of the electron," Science **343**, 269–272 (2014).
- ¹⁸D. DeMille, J. M. Doyle, and A. O. Sushkov, "Probing the frontiers of particle physics with tabletop-scale experiments," Science **357**, 990–994 (2017).
- ¹⁹V. Andreev, D. Ang, D. DeMille, J. Doyle, G. Gabrielse, J. Haefner, N. Hutzler, Z. Lasner, C. Meisenhelder, B. O'Leary, *et al.*, "Improved limit on the electric dipole moment of the electron," Nature **562**, 355–360 (2018).
- ²⁰V. Cirigliano, A. Crivellin, W. Dekens, J. de Vries, M. Hoferichter, and E. Mereghetti, "C p violation in higgs-gauge interactions: From tabletop experiments to the lhc," Physical Review Letters **123**, 051801 (2019).
- ²¹N. R. Hutzler, A. Borschevsky, D. Budker, D. DeMille, V. Flambaum, G. Gabrielse, R. Ruiz, A. Jayich, L. Orozco, M. Ramsey-Musolf, *et al.*, "Searches for new sources of cp violation using molecules as quantum sensors," arXiv preprint arXiv:2010.08709 (2020).
- ²²W. B. Cairncross, D. N. Gresh, M. Grau, K. C. Cossel, T. S. Roussy, Y. Ni, Y. Zhou, J. Ye, and E. A. Cornell, "Precision measurement of the electron's electric dipole moment using trapped molecular ions," Phys. Rev. Lett. **119**, 153001 (2017).
- ²³A. N. Petrov, "Systematic effects in the HfF⁺-ion experiment to search for the electron electric dipole moment," Phys. Rev. A 97, 052504 (2018).
- ²⁴R. Alarcon, J. Alexander, V. Anastassopoulos, T. Aoki, R. Baartman, S. Baeßler, L. Bartoszek, D. H. Beck, F. Bedeschi, R. Berger, M. Berz, H. L. Bethlem, T. Bhattacharya, M. Blaskiewicz, T. Blum, T. Bowcock, A. Borschevsky, K. Brown, D. Budker, S. Burdin, B. C. Casey, G. Casse, G. Cantatore, L. Cheng, T. Chupp, V. Cianciolo, V. Cirigliano, S. M. Clayton, C. Crawford, B. P. Das, H. Davoudiasl, J. de Vries, D. DeMille, D. Denisov, M. V. Diwan, J. M. Doyle, J. Engel, G. Fanourakis, R. Fatemi, B. W. Filippone, V. V. Flambaum, T. Fleig, N. Fomin, W. Fischer, G. Gabrielse, R. F. G. Ruiz, A. Gardikiotis, C. Gatti, A. Geraci, J. Gooding, B. Golub, P. Graham, F. Gray, W. C. Griffith, S. Haciomeroglu, G. Gwinner, S. Hoekstra, G. H. Hoffstaetter, H. Huang, N. R. Hutzler, M. Incagli, T. M. Ito, T. Izubuchi, A. M. Javich, H. Jeong, D. Kaplan, M. Karuza, D. Kawall, O. Kim, I. Koop, W. Korsch, E. Korobkina, V. Lebedev, J. Lee, S. Lee, R. Lehnert, K. K. H. Leung, C.-Y. Liu, J. Long, A. Lusiani, W. J. Marciano, M. Maroudas, A. Matlashov, N. Matsumoto, R. Mawhorter, F. Meot, E. Mereghetti, J. P. Miller, W. M. Morse, J. Mott, Z. Omarov, L. A. Orozco, C. M. O'Shaughnessy, C. Ozben, S. Park, R. W. Pattie, A. N. Petrov, G. M. Piacentino, B. R. Plaster, B. Podobedov, M. Poelker, D. Pocanic, V. S. Prasannaa, J. Price, M. J. Ramsey-Musolf, D. Raparia, S. Rajendran, M. Reece, A. Reid, S. Rescia, A. Ritz, B. L. Roberts, M. S.

TABLE I. Rovibrational spectrum parameters

Parameter	Rigid ligand, $CCSD(T)^2$	CCSD	$\operatorname{CCSD}(T)$	Experiment ^a
Stretching mode ν_1 , cm ⁻¹	550	545	536	$529.341(1)^{47}$
Bending mode ν_2 , cm ⁻¹	319	351	342	$319(5)^{48}, 329(5)^{49}, 339(5),^{47}$
Ligand mode ν_3 , cm ⁻¹	-	4055	4030	-
Rotational constant $B(\nu_1 = 0, \nu_2 = 0, \nu_3 = 0)$, cm ⁻¹	0.2461	0.2456	0.2468	$0.245434(13)^{47}$
Rotational constant $B(\nu_1 = 1, \nu_2 = 0, \nu_3 = 0), \text{ cm}^{-1}$	-	0.2437	0.2443	$0.2439771(35)^{47}$
<i>l</i> -doubling $\Delta E_{J=1} = 2q$, MHz	26	23	24	-

 $^{\rm a}$ The number in parentheses denotes 2σ deviation

TABLE II. Convergence of the results with n_{max} (see eq. (13)) for different sets of parameters R_i and θ_i (see eq.(15)) for CCSD(T) calculations

parameter	Set of R_i, θ_i	$n_{max} = 1$	$n_{max} = 2$	$n_{max} = 3$
Bending mode ν_2 , cm ⁻¹	1	344.0892	341.9777	341.9665
	2	340.9922	341.9698	341.9671
<i>l</i> -doubling $\Delta E_{J=1} = 2q$, MHz	1	23.520	24.247	24.257
	2	23.712	24.235	24.256

Safronova, Y. Sakemi, P. Schmidt-Wellenburg, A. Shindler, Y. K. Semertzidis, A. Silenko, J. T. Singh, L. V. Skripnikov, A. Soni, E. Stephenson, R. Suleiman, A. Sunaga, M. Syphers, S. Syritsyn, M. R. Tarbutt, P. Thoerngren, R. G. E. Timmermans, V. Tishchenko, A. V. Titov, N. Tsoupas, S. Tzamarias, A. Variola, G. Venanzoni, E. Vilella, J. Vossebeld, P. Winter, E. Won, A. Zelenski, T. Zelevinsky, Y. Zhou, and K. Zioutas, "Electric dipole moments and the search for new physics," (2022).

- ²⁵T. A. Isaev and R. Berger, "Polyatomic candidates for cooling of molecules with lasers from simple theoretical concepts," Phys. Rev. Lett. **116**, 063006 (2016).
- ²⁶T. A. Isaev, A. V. Zaitsevskii, and E. Eliav, "Laser-coolable polyatomic molecules with heavy nuclei," Journal of Physics B: Atomic, Molecular and Optical Physics **50**, 225101 (2017).
- ²⁷I. Kozyryev and N. R. Hutzler, "Precision measurement of time-reversal symmetry violation with laser-cooled polyatomic molecules," Phys. Rev. Lett. **119**, 133002 (2017).
- ²⁸N. R. Hutzler, "Polyatomic molecules as quantum sensors for fundamental physics," Quantum Science and Technology 5, 044011 (2020).
- ²⁹A. Zakharova, "Rotating and vibrating symmetric-top molecule raoch 3 in fundamental p, t-violation searches," Physical Review A **105**, 032811 (2022).
- ³⁰I. Kozyryev, L. Baum, K. Matsuda, B. L. Augenbraun, L. Anderegg, A. P. Sedlack, and J. M. Doyle, "Sisyphus laser cooling of a polyatomic molecule," Physical review letters **118**, 173201 (2017).
- ³¹L. Baum, N. B. Vilas, C. Hallas, B. L. Augenbraun, S. Raval, D. Mitra, and J. M. Doyle, "1d magneto-optical trap of polyatomic molecules," Physical review letters **124**, 133201 (2020).
- ³²D. Mitra, N. B. Vilas, C. Hallas, L. Anderegg, B. L. Augenbraun, L. Baum, C. Miller, S. Raval, and J. M. Doyle, "Direct laser cooling of a symmetric top molecule," Science **369**, 1366–1369 (2020).
- ³³T. C. Steimle, C. Linton, E. T. Mengesha, X. Bai, and A. T. Le, "Field-free, Stark, and Zeeman spectroscopy of the $\tilde{A}^2\Pi_{1/2} \tilde{X}^2\Sigma^+$ transition of ytterbium monohydroxide," Physical Review A **100**, 052509 (2019).
- ³⁴B. L. Augenbraun, Z. D. Lasner, A. Frenett, H. Sawaoka, C. Miller, T. C. Steimle, and J. M. Doyle, "Laser-cooled polyatomic molecules for improved electron electric dipole moment searches," New Journal of Physics **22**, 022003 (2020).
- ³⁵N. B. Vilas, C. Hallas, L. Anderegg, P. Robichaud, A. Winnicki, D. Mitra, and J. M. Doyle, "Magneto-optical trapping and sub-

doppler cooling of a polyatomic molecule," Nature **606**, 70–74 (2022).

- ³⁶M. G. Kozlov and L. N. Labzowsky, "Parity violation effects in diatomics," Journal of Physics B: Atomic, Molecular and Optical Physics **28**, 1933 (1995).
- ³⁷A. Titov, N. Mosyagin, A. Petrov, T. Isaev, and D. DeMille, "P, t-parity violation effects in polar heavy-atom molecules," in *Re*cent Advances in the Theory of Chemical and Physical Systems (Springer, 2006) pp. 253–283.
- ³⁸M. S. Safronova, D. Budker, D. DeMille, D. F. J. Kimball, A. Derevianko, and C. W. Clark, "Search for New Physics with Atoms and Molecules," Rev. Mod. Phys. **90**, 025008 (2018), arXiv:1710.01833 [physics.atom-ph].
- ³⁹A. Zakharova and A. Petrov, "*P*,*T*-odd effects for the RaOH molecule in the excited vibrational state," Phys. Rev. A 103, 032819 (2021), arXiv:2012.08427 [physics.atom-ph].
- ⁴⁰DIRAC, a relativistic ab initio electronic structure program, Release DIRAC19 (2019), written by A. S. P. Gomes, T. Saue, L. Visscher, H. J. Aa. Jensen, and R. Bast, with contributions from I. A. Aucar, V. Bakken, K. G. Dyall, S. Dubillard, U. Ekström, E. Eliav, T. Enevoldsen, E. Faßhauer, T. Fleig, O. Fossgaard, L. Halbert, E. D. Hedegård, B. Heimlich-Paris, T. Helgaker, J. Henriksson, M. Iliaš, Ch. R. Jacob, S. Knecht, S. Komorovský, O. Kullie, J. K. Lærdahl, C. V. Larsen, Y. S. Lee, H. S. Nataraj, M. K. Nayak, P. Norman, G. Olejniczak, J. Olsen, J. M. H. Olsen, Y. C. Park, J. K. Pedersen, M. Pernpointner, R. di Remigio, K. Ruud, P. Sałek, B. Schimmelpfennig, B. Senjean, A. Shee, J. Sikkema, A. J. Thorvaldsen, J. Thyssen, J. van Stralen, M. L. Vidal, S. Villaume, O. Visser, T. Winther, and S. Yamamoto (available at http://dx.doi.org/10.5281/zenodo.3572669, see also http://www.diracprogram.org).
- ⁴¹URL: http://www.qchem.pnpi.spb.ru/Basis/~, GRECPs and basis sets.
- ⁴²A. Titov and N. Mosyagin, "Generalized relativistic effective core potential: Theoretical grounds," International journal of quantum chemistry **71**, 359–401 (1999).
- ⁴³N. S. Mosyagin, A. Zaitsevskii, and A. V. Titov, "Shapeconsistent relativistic effective potentials of small atomic cores," International Review of Atomic and Molecular Physics 1, 63–72 (2010).
- ⁴⁴N. S. Mosyagin, A. V. Zaitsevskii, L. V. Skripnikov, and A. V. Titov, "Generalized relativistic effective core potentials for actinides," International Journal of Quantum Chemistry **116**, 301–

315 (2016).

- ⁴⁵P. McGuire and D. J. Kouri, "Quantum mechanical close coupling approach to molecular collisions. j_z -conserving coupled states approximation," The Journal of Chemical Physics **60**, 2488–2499 (1974).
- ⁴⁶G. Herzberg, Molecular spectra and molecular structure. Vol. 3: Electronic spectra and electronic structure of polyatomic molecules (New York: Van Nostrand, 1966).
- $^{47}\text{T.}$ C. Melville and J. A. Coxon, "The visible laser excitation spectrum of YbOH: The $\tilde{A}^2\Pi-\tilde{X}^2\Sigma^+$ transition," The Journal of Chemical Physics **115**, 6974–6978 (2001).
- ⁴⁸C. Zhang, B. L. Augenbraun, Z. D. Lasner, N. B. Vilas, J. M. Doyle, and L. Cheng, "Accurate prediction and measurement of vibronic branching ratios for laser cooling linear polyatomic molecules," The Journal of Chemical Physics **155**, 091101 (2021).
- ⁴⁹E. T. Mengesha, A. T. Le, T. C. Steimle, L. Cheng, C. Zhang, B. L. Augenbraun, Z. Lasner, and J. Doyle, "Branching ratios, radiative lifetimes, and transition dipole moments for yboh," The Journal of Physical Chemistry A **124**, 3135–3148 (2020).