Nesterov smoothing for sampling without smoothness

Jiaojiao Fan, Bo Yuan, Jiaming Liang, Yongxin Chen Georgia Tech {jiaojiaofan,boyuan48,jiaming.liang,yongchen@gatech.edu}

Abstract

We study the problem of sampling from a target distribution in \mathbb{R}^d whose potential is not smooth. Compared with the sampling problem with smooth potentials, this problem is more challenging and much less well-understood due to the lack of smoothness. In this paper, we propose a novel sampling algorithm for a class of non-smooth potentials by first approximating them by smooth potentials using a technique that is akin to Nesterov smoothing. We then utilize sampling algorithms on the smooth potentials to generate approximate samples from the original nonsmooth potentials. With a properly chosen smoothing intensity, the accuracy of the algorithm is guaranteed. For strongly log-concave distributions, our algorithm can achieve \mathcal{E} error in Wasserstein-2 distance with complexity $\widetilde{\mathcal{O}}\left(\frac{d^{1/3}}{\mathcal{E}^{5/3}}\right)$. For log-concave distributions, we achieve \mathcal{E} error in total variation with complexity $\mathcal{O}\left(\frac{M_{\pi}d}{\mathcal{E}^3}\right)$ in expectation with M_{π} being the second moment of the target distribution. For target distributions satisfying the logarithmic-Sobolev inequality, our algorithm has complexity $\mathcal{O}\left(\frac{d}{c}\right)$.

1 Introduction

Sampling from a target distribution $\pi(x) \propto \exp(-s(x))$ known up to a normalization constant is an important problem in many areas such as machine learning, due to its pivotal role in Bayesian statistics and inference [22, 16, 27, 21]. Over the past decades, there has been a vast amount of research carried out on sampling problems from smooth distributions [13, 49], i.e., the gradient of the potential s(x) is Lipschitz. In contrast, the understanding of non-smooth sampling is still quite limited [7, 45, 32, 33, 12].

In this work, we consider the task of sampling from a specific class of non-smooth distributions $\pi(x) \propto \exp(-s(x))$ whose potentials s(x) permit an explicit max-structure as

$$s(x) = f(x) + \max_{y \in \mathcal{Y}} \{ \langle h(x), y \rangle - g(y) \}$$
(1)

where f is smooth, g is convex, \mathcal{Y} is convex and bounded, and h is Lipschitz and smooth. s(x) can be non-smooth. For instance, when f(x) = 0 and h(x) = Ax, s(x) becomes a piece-wise affine function and is clearly non-smooth. A special instance of particular interest is

$$s(x) = f(x) + \max\{h_1(x), h_2(x), \dots, h_n(x)\}.$$
(2)

The optimization of minimizing s(x) has been widely used for science and engineering problems to optimize the worst-case performance [3, 15]. Its Bayesian counterpart that accounts for uncertainties can be captured by a sampling problem from the potential s(x).

Compared to other recent work [7, 45, 32, 33, 12] for sampling from black box non-smooth distribution, we assume certain structures in the target distribution and hope to leverage these structures to achieve better performance. Indeed, in practice, it is rare to have a black box model as a

Preprint.

target distribution; we almost always know some structures of the problem. It could be advantageous to utilize such structures in the algorithm design. With this in mind, we take one step forward in the direction of sampling beyond black box models and propose a sampling method for non-smooth distribution that utilizes the specific max-structure (1). We develop a sampling algorithm for non-smooth potentials whose complexity has the same dimensional dependency as its smooth counterparts.

Our strategy is akin to Nesterov smoothing [40] in optimization that constructs a smooth function $s_{\beta}(x)$ whose difference with s(x) can be controlled through the smoothing intensity β . In Nesterov smoothing, the idea is to use a fast smooth optimization algorithm to minimize $s_{\beta}(x)$ for the purpose of optimizing the non-smooth function s(x). In our problem, instead, we sample from $\pi_{\beta}(x) \propto \exp(-s_{\beta}(x))$ for the purpose of sampling from $\pi(x) \propto \exp(-s(x))$. Thanks to the smoothness of $s_{\beta}(x)$, we can take advantage of the many existing sampling algorithms for smooth potentials to sample from $\pi_{\beta}(x) \propto \exp(-s_{\beta}(x))$. Our smoothing strategy is compatible with any such algorithms. With state-of-the-art sampling algorithms for smooth potentials and a proper smoothing intensity β , we obtain a series algorithms with competitive complexity bounds. Surprisingly, the complexity bounds we obtained with Nesterov smoothing have exactly the same dimension dependency as their smooth counterparts. This is crucial for high dimensional problems. In a high level, this work is along the direction of the recent line of research that tries to bridge optimization and sampling [51, 2, 33]. Our results prove that some smoothing techniques used in optimization are equally effective in sampling.

Related Works: Over the last few years, several algorithms for sampling from potentials that are not smooth have been developed. A majority of them is devoted to the analysis of original LMC in the non-smooth setting [19, 41, 12]. These theorectical results are challenging and significantly different from those in the smooth setting. In Bernton [4], Durmus et al. [17], Salim & Richtárik [45], the non-smooth sampling problem is studied and analyzed from an optimization point of view. Another algorithm that is effective for non-smooth sampling is the proximal sampler developed in the sequence of work [30, 32, 33, 9]. The methods that are most related to ours are Chatterji et al. [7], Durmus et al. [16], Lau & Liu [28], in which the authors use Gaussian smoothing and Moreau envelope. Gaussian smoothing can be applied to any convex potential, however the evaluation of the smoothed potential is difficult. Normally one can only get unbiased estimation of it through Gaussian sampling, inducing additional variance in the algorithm. Moreau envelope can be applied to any weakly-convex potential. To evaluate the smoothed potential or its gradient, one needs to solve an optimization problem (to get the proximal map), inducing additional complexity, especially when the potential is not convex. We include a more detailed comparison with these two smoothing techniques in the end of Section 3.2.

Contributions: We summarize our contributions as follows.

i) Inspired by Nesterov smoothing in optimization, we develop a smoothing technique, different from existing ones like Gaussian smoothing and Moreau envolep, for sampling problems for a class of potentials that permit the max-structure (1).

ii) Our smoothing technique is compatible with any first-order sampling algorithm for smooth potentials and the performance is guaranteed via a proper smoothing intensity β .

iii) Combining this smoothing technique with start-of-the-art sampling algorithms for smooth potentials we obtain competitive complexity bounds for strongly-log-concave distributions, log-concave distributions, and distributions satisfying log-Sobolev inequality (see Table 1). In particular, we achieve $d^{1/3}$ dependence in sampling non-smooth strongly-log-concave distributions, better than all existing methods. Besides, our analysis covers certain composite potentials with non-smooth (Lipschitz continuous) components and the distribution satisfies LSI. The complexity bound in this setting has not been established before.

Remark 1. In Table 1, our result contains one additional problem-dependent parameter D defined in (5). It can hide dimension dependency, like any other parameter λ_h, λ_ρ or L_f . In Abstract and Contribution, our claim in terms of dimension dependency assumes other parameters are dimensionfree. In realistic examples, this is unlikely to be true, so one need to include the dimension dependency hidden in all parameters for a fair comparison. An existing caveat is if D hides dimension dependency, then our algorithm is clearly not state-of-art. Interestingly, for the class of problems with potentials of the form (2), $D = \log(n)$ will be seen as in Section 4.4.

Table 1: Convergence comparison for state-of-the-art non-smooth sampling. We highlight the dependency on dimension d, accuracy \mathcal{E} , smoothness L_f of f(x), Lipschitz λ_h of h(x), Lipschitz λ_ρ of $\rho(x) := \max_{y \in \mathcal{Y}} \{ \langle h(x), y \rangle - g(y) \}$, boundness D defined in (5), and log-Sobolev constant C_{π} . A trivial upper bound for λ_{ρ} is $\lambda_h \max ||y||_2$. Other than Durmus et al. [17], all results are the last-iterate convergence.

Assumption	Source	Complexity	Metric	Note
	[33]	$\widetilde{\mathcal{O}}\left(rac{(\lambda_{ ho}^2 ee L_f)d}{lpha} ight)$	Rényi	
	[45]	$\widetilde{\mathcal{O}}\left(rac{\lambda_{ ho}^2+L_fd}{lpha^2\mathcal{E}^2} ight)$	W_2	
α -strongly	[7]	$\widetilde{\mathcal{O}}\left(rac{(\lambda_ ho+L_f)^2d^{5/2}}{lpha^4\mathcal{E}^4} ight)$	W_2	Gaussian smoothing
log-concave	Theorem 1 (1)	$\widetilde{\mathcal{O}}\left(\frac{\lambda_{h}^{2}Dd^{1/3}}{\alpha^{11/6}\mathcal{E}^{5/3}} + \frac{\lambda_{h}^{7/3}D^{7/6}d^{1/6}}{\alpha^{23/12}\mathcal{E}^{3/2}}\right)$	W_2	
	Theorem 1 (2)	$\widetilde{\mathcal{O}}\left(rac{\lambda_h^2 D d^{1/2}}{lpha \mathcal{E}} ight)$	TV	Assume warm-start
	[33]	$\mathcal{O}\left(rac{(\lambda_{ ho}^2 ee L_f) d^2}{\mathcal{E}} ight)$	Rényi	
	[31]	$\mathcal{O}\left(rac{\lambda_{ ho}^2 d^2}{\mathcal{E}^4} ight)$	W_2	Assume $M_{\pi} = \mathcal{O}(d)$
log-concave	[17]	$\mathcal{O}\left(rac{L_f d^2 + \lambda_ ho^2 d}{\mathcal{E}^2} ight)$	KL	
	[16]	$\mathcal{O}\left(rac{\lambda_{ ho}^2 d^5}{\mathcal{E}^2} ight)$	TV	Moreau envelope
	Theorem 2	$\mathcal{O}\left(rac{\lambda_h^2 D M_\pi d}{\mathcal{E}^3} ight)$	TV	
log-Sobolev	Theorem 3	$\widetilde{\mathcal{O}}\left(rac{\lambda_h^2 D C_\pi d}{\mathcal{E}} ight)$	TV	

2 Background

Before we proceed, we briefly introduce the notation basics. We assume the norm in the space of x and y can be different and denote them as $\|\cdot\|_{\mathcal{X}}$ and $\|\cdot\|_{\mathcal{Y}}$ respectively. We use $J_h(x) : \mathbb{R}^d \to \mathbb{R}^{n \times d}$ to represent the Jacobian matrix of the function h at x. The norm of a matrix $A : \mathbb{R}^d \to \mathbb{R}^{n \times d}$ is defined as $\|A\|_{\mathcal{X},\mathcal{Y}} = \max_{x,y}\{\langle Ax, y \rangle_{\mathcal{Y}} : \|x\|_{\mathcal{X}} = 1, \|y\|_{\mathcal{Y}} = 1\}$. Given a primal space equipped with norm $\|\cdot\|$, we denote the norm in its dual space as $\|\cdot\|^*$. In this article, we assume the probability measure have density with respect to the Lebesgue measure.

2.1 Nesterov smooothing

It is well-known that with subgradient methods, the optimal bound of non-smooth convex minimization complexity is $\mathcal{O}(1/\epsilon^2)$ to achieve ϵ error in the function value [38, 39]. However, beyond the black-box oracle model, there are various structures that can be used to improve this bound. Nesterov smoothing [40] is an interesting technique for functions with explicit max structure (1). In doing so, the complexity order can be improved to $\mathcal{O}(1/\mathcal{E})$. The underlying principle of Nesterov smoothing is that

The dual of a strongly-convex function is smooth and vice versa [26].

Specifically, it adds a σ -strongly-convex penalty function $\ell(y)$ to the original maximization such that

$$s_{\beta}(x) = f(x) + \max_{y \in \mathcal{Y}} \{ \langle h(x), y \rangle - g(y) - \beta \ell(y) \},\$$

where β is the smoothing intensity. In the original paper of Nesterov smoothing, they consider the specific task h(x) = Ax and find s_{β} is smooth with constant $L_f + ||A||^2_{\mathcal{X},\mathcal{Y}}/\beta\sigma$. Then they apply the Nesterov accelerated gradient descent algorithm to minimize s_{β} , an optimal scheme for smooth optimization. Our idea is very similar on the high level: use smoothing for the non-smooth potential first, and then apply the existing sampling algorithm for the smoothed distribution.

2.2 Smooth sampling

In this section, we summarize several state-of-art sampling algorithms for smooth distributions. In general, there are several dominant popular algorithms: Unadjusted Langevin Monte Carlo (LMC) [43], Underdamped/Kinetic Langevin Monte Carlo (KLMC) [11] Metropolis-Adjusted Langevin Algorithm (MALA) [44], and Hamiltonian Monte Carlo (HMC) [37]. Suppose we are given the target distribution $\pi \propto \exp(-V(x))$ for some smooth potential V(x). The LMC algorithm runs the update

$$x_{t+1} = x_t - \gamma \nabla V(x) + \sqrt{2\gamma}\epsilon_t, \tag{3}$$

where ϵ_t is the standard Gaussian random variable in \mathbb{R}^d , x_0 is a user-defined starting point and $\gamma > 0$ is the step size that is small enough. Indeed, (3) is the Euler discretization of Langevin diffusion dynamics,

$$\mathrm{d}x_t = -\nabla V(x_t)\mathrm{d}t + \sqrt{2}\mathrm{d}B_t. \tag{4}$$

This dynamics has an invariant density π no matter what x_0 is. The underdamped Langevin diffusion adds an Hamiltonian ingredient to the standard Langevin diffusion (4), so that an additional random variable $v_t := dx_t/dt$ that controls the velocity in the dynamics. Similarly, HMC maintains a velocity term, but with a different randomization manner. MALA is also a variant of LMC. As its name implies, it adds a Metropolis-Hastings [36, 24] rejection step to LMC, in order to eliminate the bias induced by discretization. We refer the readers to MacKay et al. [34] for the an overview of these algorithms.

In spite of the canonical role of LMC in sampling, it is not the fastest algorithm empirically. Theoretically, in the standard strongly-convex potential case, the complexity of LMC can achieve $\widetilde{O}(d)$ [13, 49, 20], whereas KLMC, HMC algorithms can achieve $\widetilde{O}(d^{1/2})$ [14, 10]. The complexity $\widetilde{O}(d^{1/2})$ also holds for MALA, however, under a warm start [52]. Regarding the accuracy dependency, MALA is notably the best with a benefit from Metropolis step. With a considerable research accumulated for decades, several novel sampling algorithms or the variants are proposed recently, such as the proximal algorithm [30, 9, 33] based on restricted Gaussian oracle, and the randomized midpoint method for KLMC [47]. The proximal algorithm has been proved to have the state-of-art dimension dependency for log-concave distribution [9].

There exist some accelerating algorithms that can achieve even lower dimension order in complexity [35, 46]. However, they require stronger regularity assumptions, e.g. bounded third-order derivative. Although we do not consider these higher-order algorithms in this paper for the sake of generality, one can still apply them in our framework if the higher-order smoothness of $s_{\beta}(x)$ can be verified in practice.

3 Nesterov smoothing for sampling

In this section, we will introduce our Nesterov smoothing sampling algorithm in Section 3.1 and bound the error caused by smoothing in Section 3.2.

3.1 Problem setup and algorithm

Formally, we consider the task to sample from a non-smooth distribution $\pi(x) \propto \exp(-s(x))$ where

$$s(x) = f(x) + \max_{y \in \mathcal{Y}} \{ \langle h(x), y \rangle - g(y) \}.$$

We will assume $f(x) : \mathcal{X} \to \mathbb{R}$ is L_f -smooth; $h(x) : \mathcal{X} \to \mathcal{Y}$ is λ_h -Lipschitz and L_h -smooth; $g(y) : \mathcal{Y} \to \mathbb{R}$ is continuous and convex through out the paper. A function f(x) is L-smooth if $\|\nabla f(x_1) - \nabla f(x_2)\|_{\mathcal{X}}^* \leq L \|x_1 - x_2\|_{\mathcal{X}}$ for $\forall x_1, x_2 \in \mathbb{R}^d$. Correspondingly, we say a multivariate function h(x) is L-smooth if $\|J_h(x_1) - J_h(x_2)\|_{\mathcal{X},\mathcal{Y}} \leq L \|x_1 - x_2\|_{\mathcal{X}}$ for $\forall x_1, x_2 \in \mathcal{X}$. We say a multivariate function h(x) is λ -Lipschitz if $\|h(x_1) - h(x_2)\|_{\mathcal{Y}}^* \leq \lambda \|x_1 - x_2\|_{\mathcal{X}}$, which is equivalent to $\|J_h(x)\|_{\mathcal{X},\mathcal{Y}} \leq \lambda$ for $\forall x \in \mathcal{X}$ if h(x) is differentiable. This boils down to that the operator norm of A is bounded when h(x) = Ax. And this allows us to compare the smoothing effect with the case considered in Nesterov [40]. Without more specification, in the rest of paper, we assume $\|\cdot\|_{\mathcal{X}}^* = \|\cdot\|_{\mathcal{X}} = \|\cdot\|_{\mathcal{X}} = \|\cdot\|_2$ is Euclidean norm to be consistent with the sampling literature. Algorithm 1 Nesterov smoothing sampling framework

Input: A first-order based smooth sampling algorithm, the target distribution $\pi \propto \exp(-s(x))$, number of iterations K, initial point $x_0 \sim \mu_0$ **Initialization:** Choose the smoothing intensity β for $k = 0, \ldots, K - 1$ do *# Calculate* $\nabla s_{\beta}(x)$ *according to* (6) Iterate x_{k+1} by the smooth sampling algorithm end for Output: x_K

It is very likely that s(x) is not differentiable since it is a "piece-wise" function segmented by different maximizer y. As such, it is not feasible to directly apply the smooth sampling algorithms. Inspired by the Nesterov smoothing technique, our strategy is to

1) smooth s(x) as $s_{\beta}(x)$, 2) apply smooth sampling algorithm to $\pi_{\beta}(x)$.

For smoothing, we consider the surrogate function

$$s_{\beta}(x) = f(x) + \max_{y \in \mathcal{Y}} \{ \langle h(x), y \rangle - g(y) - \beta \ell(y) \},\$$

where $\ell(y)$ is a σ -strongly-convex prox-function

$$\ell(y) \ge \frac{\sigma}{2} \|y - y_0\|_{\mathcal{Y}}^2.$$

The center of \mathcal{Y} is defined as $y_0 = \arg \min_y \{\ell(y) : y \in \mathcal{Y}\}$ and we can assume that $\ell(y_0) = 0$. The compactness of \mathcal{Y} also allows us to define

$$D = \max_{y} \{\ell(y) : y \in \mathcal{Y}\}$$
(5)

and $R = \max_{\mathcal{Y}} \{ \|y\|_{\mathcal{Y}} : y \in \mathcal{Y} \}$. From the definition, D is the squared diameter of the set \mathcal{Y} , and the R is the furthest distance of any point in the set to the origin. By the triangular inequality, it holds that $R \leq \|y_0\| + \sqrt{\frac{2D}{\sigma}}$. In this way, $s_\beta(x)$ is a smooth function, which is justified by Lemma 1. The proof is postponed to Appendix.

Lemma 1. Let
$$\phi_{\beta}(x,y) := \langle h(x), y \rangle - g(y) - \beta \ell(y)$$
, and $y_{\beta}(x) := \arg \max_{y \in \mathcal{Y}} \phi_{\beta}(x,y)$, then

$$\nabla s_{\beta}(x) = \nabla f(x) + J_h^{\dagger}(x) y_{\beta}(x).$$
(6)

Moreoever, $s_{\beta}(x)$ is $L_{s_{\beta}}$ -smooth with

$$L_{s_{\beta}} = L_f + RL_h + \frac{\lambda_h^2}{\beta\sigma}.$$

If h(x) = Ax, then $L_h = 0$ and $\lambda_h = ||A||_{\mathcal{X},\mathcal{Y}}$, we get $L_{s_\beta} = L_f + \frac{1}{\beta\sigma} ||A||_{\mathcal{X},\mathcal{Y}}^2$ in this special case. This recovers the smoothness constant in Nesterov [40, Theorem 1]. In many cases, λ_h is dimensional-free. For example, when h(x) = Ax and $A = cI_m$, then $\lambda_h = c$ regardless of the dimensionality.

We will also assume that we can solve $\max_y \phi_\beta(x, y)$ for any $x \in \mathcal{X}$ in a fast dimension-free manner, which has been verified by the complexity analysis [6, Section 3] and especially several examples in Nesterov [40, Section 4]. Thus we can query the gradient of $s_\beta(x)$ according to (6) and resort to first-order sampling algorithm to sample from $\pi_\beta \propto \exp(-s_\beta(x))$. The pseudo-code of our scheme is presented in Algorithm 1.

3.2 Bounded distance between π and π_{β}

When $\beta \to 0$, it holds that $\pi_{\beta} \to \pi$. In this section, we rigorously bound the distance between π and π_{β} . A key observation is that $s_{\beta}(x)$ is a uniform smooth approximation of s(x), i.e., $s_{\beta}(x) \le s(x) \le s_{\beta}(x) + \beta D$ (see (13)) [40, Eq. 2.7]. To bound the difference between π and π_{β} , we

select two widely-used distributional distance as the error criteria. The first one is the total variation $TV(p,q) = \frac{1}{2} ||p - q||_1$, which can be bounded by Kullback–Leibler divergence KL(p||q) through Pinsker's inequality

$$\mathrm{TV}(p,q) \le \sqrt{\frac{1}{2}\mathrm{KL}(p\|q)}.$$
(7)

The second one is the Wasserstein-2 distance [50]

$$W_2(p,q) := \sqrt{\inf_{\nu \in \Pi(p,q)} \int_{\mathbb{R}^d \times \mathbb{R}^d} \|a - b\|_2^2 d\nu(a,b)},$$

where $\Pi(p,q)$ denotes the set of the all joint distributions of p and q. The W_2 distance can also be bounded by KL(p||q) through the Talagrand inequality [42]

$$W_2(p,q) \le \sqrt{2C_{\text{LSI}}\text{KL}(p\|q)}$$

if q satisfies the following logarithmic-Sobolev inequality (LSI) with constant C_{LSI} .

Definition 1 (Log-Sobolev inequality). A probability distribution p satisfies a logarithmic-Sobolev inequality with constant C_{LSI} if, for all smooth functions $u : \mathbb{R}^d \to \mathbb{R}$, it holds that

$$\mathbb{E}_p[u^2 \log u^2] - \mathbb{E}_p[u^2] \log \mathbb{E}_p[u^2] \le 2C_{\text{LSI}} \mathbb{E}_p[\|\nabla u\|^2].$$
(LSI)

LSI is a powerful tool for the sampling convergence analysis when there is no log-concavity. It has nice properties, for instance, it implies the sub-Gaussian concentration property [29]. Moreoever, if s(x) is α -strongly-convex, then $\pi(x)$ satisfies (LSI) with constant $1/\alpha$ [1]. With the help of the above inequalities, we are able to show that Proposition 1, 2 hold.

Proposition 1. $TV(\pi, \pi_{\beta}) \leq \frac{\beta D}{2}$.

Proposition 2. If π satisfies (LSI) with constant C_{π} , then $W_2(\pi, \pi_{\beta}) \leq \sqrt{C_{\pi}\beta D}$.

We next compare the smoothing effect of Nesterov smoothing, Moreau envelope and Gaussian smoothing. Recall that we define the non-smooth part in our potential as $\rho(x)$, and Lipschitz constant of $\rho(x)$ to be λ_{ρ} . All of three smoothing methods can deal with composite smooth+non-smooth potential, but *they require the convexity or weakly-convexity of* $\rho(x)$, while Nesterov smoothing does not require so.

Moreau envelope The Moreau-Yosida envelope of a l.s.c. convex function $\rho(x)$ is defined as

$$\rho_{\beta}(x) := \min_{y \in \mathbb{R}^d} \left\{ \rho(y) + \frac{1}{2\beta} \|x - y\|_2^2 \right\}.$$

 ρ_{β} is L_{ρ} -smooth with $L_{\rho} = \frac{1}{\beta}$. The smoothed composite potential is then written as $s_{\beta}(x) = f(x) + \rho_{\beta}(x)$. The smoothing effect of Moreau envelope applies to not only l.s.c. convex functions, but also l.s.c. weakly-convex functions with an upper bound constraint on β [5, 25]. According to Proposition 1 in Durmus et al. [16], there is $\text{TV}(\pi, \pi_{\beta}) \leq \frac{1}{2}\beta\lambda_{\rho}^2$, which is implied by $0 \leq s(x) - s_{\beta}(x) \leq \beta \|\lambda_{\rho}\|^2/2$.

Gaussian smoothing Unlike the proposed method in Chatterji et al. [7], where they apply Gaussian smoothing to both the smooth part f(x) and the non-smooth part $\rho(x)$ in s(x), our discussion is based on only smoothing the $\rho_{\beta}(x)$ here. We find smoothing both parts would introduce a worse dimension dependency in bounding the difference between $\pi(x)$ and $\pi_{\beta}(x)$. The Gaussian smoothing for $\rho(x)$ is defined as $\rho_{\beta}(x) = \mathbb{E}_{\xi}[\rho(x + \beta\xi)]$, where $\xi \sim \mathcal{N}(0, I_{d \times d})$. The smoothing effect of Gaussian smoothing only applies to convex functions. Indeed, according to Lemma 2.2 in Chatterji et al. [7], if $\rho(x)$ is convex, then $\rho_{\beta}(x)$ is L_{ρ} -smooth with $L_{\rho} = \lambda_{\rho} d^{1/2} / \beta$. To bound the difference between π and π_{β} , one doesn't need a stronger assumption. By Lemma E.2 in Chatterji et al. [7], $\mathrm{TV}(\pi, \pi_{\beta}) \leq \frac{1}{2}\beta\lambda_{\rho}d^{1/2}$. Additionally, if f(x) is α -strongly convex, by Lemma 3.3 in Chatterji et al. [7], $W_2(\pi, \pi_{\beta}) = \widetilde{O}\left(\frac{\beta\lambda_{\rho}d}{\alpha}\right)$.

Now, if we want to achieve $TV(\pi, \pi_{\beta}) = \mathcal{E}/2$, then Moreau envelope results in $L_{\rho} = \frac{\lambda_{\rho}^2}{\mathcal{E}}$; Gaussian smoothing results in $L_{\rho} = \frac{\lambda_{\rho}^2 d}{\mathcal{E}}$; Nesterov smoothing results in $L_{\rho} = RL_h + \frac{\lambda_h^2 D}{\mathcal{E}\sigma}$. Given the same

smooth sampling algorithm, the dimension dependency in L_{ρ} determines the dimension dependency in the final convergence rate. Normally, $\sigma = 1$. Assuming $\lambda_{\rho} = \lambda_h, D = \widetilde{\mathcal{O}}(1)$, then Nesterov smoothing effect is the same order as Moreau envelope, and better than Gaussian smoothing. If Dhides dimension dependency, then Nesterov smoothing is worse than Moreau envelope. However, Moreau envelope requires more strict assumption on $\rho(x)$, i.e. weakly-convexity; and since the potential is not smooth, solving the proximal operator of Moreau envelope requires additional complexity in each sampling step, even for structured potential in (1).

4 Non-asymptotic convergence analysis

In this section, we present our convergence analysis for several fundamental scenarios: strongly log-concavity, log-concavity, and log-Sobolev inequality. Denote the underlying distribution of x_k as μ_k . We will give the sufficient number of gradient queries for μ_k to reach a given error tolerance. As discussed in Section 2.2, there are multiple algorithm candidates to use for the smooth sampling in Algorithm 1. Our strategy is to test every algorithm among them and select the one with the best complexity. In the main theorems of this section, each complexity is accompanied by a state-of-art smoothing algorithm, which we will specify concretely.

4.1 Strong log-concavity

The strong log-concavity condition is mostly-widely studied for sampling convergence since it can be viewed as the strongest assumption. To our best knowledge, under this condition and without higher-order smoothness,¹ the randomized midpoint KLMC has the best dimension dependency $\widetilde{\mathcal{O}}(d^{1/3})$ among the first-order sampling algorithms. Whereas MALA doesn't have the best dependency on dimension, it has best accuracy dependency. So we will adopt these two algorithms as our smooth sampling algorithm.

We also need to point out that, to have such good complexity, MALA needs a warm-start initialization. The initial distribution μ_0 is called *G*-warm if the $\sup_{S \in \mathcal{B}(\mathbb{R}^d)} \mu_0(S)/\pi(S) \leq G$, where \mathcal{B} means the Borel sigma algebra. If *G* does not scale with dimension or condition number, then μ_0 is called a warm start. This warm-start may not hold without a delicate initialization. For example, if Gaussian initialization is used, $G = \mathcal{O}(\exp(d))$.

Theorem 1. Assume f is α -strongly-convex, and $\langle h(\cdot), y \rangle$ is convex for any $y \in \mathcal{Y}$. 1) In Algorithm 1, if we choose the smooth sampling algorithm to be randomized midpoint KLMC [47] with initial point $x_0 = \arg \min s(x)$, $\beta = \frac{\sqrt{\alpha}\mathcal{E}}{2D}$, then for any $0 < \mathcal{E} < 2\sqrt{\frac{d}{\alpha}}$, the iterate μ_K satisfies $W_2(\mu_K, \pi) \leq \mathcal{E}$ for $K = \widetilde{\mathcal{O}}\left(\left(\frac{L_{s_\beta}}{\alpha}\right)^{7/6} \left(\frac{2}{\mathcal{E}}\sqrt{\frac{d}{\alpha}}\right)^{1/3} + \left(\frac{L_{s_\beta}}{\alpha}\right) \left(\frac{2}{\mathcal{E}}\sqrt{\frac{d}{\alpha}}\right)^{2/3}\right)$. If \mathcal{E} is sufficiently small such that $\frac{L_f + RL_h}{\lambda_h^2 D} = \mathcal{O}(1/\mathcal{E})$, we obtain

$$K = \widetilde{\mathcal{O}}\left(\frac{\lambda_h^2 D d^{1/3}}{\alpha^{11/6} \mathcal{E}^{5/3}} + \frac{\lambda_h^{7/3} D^{7/6} d^{1/6}}{\alpha^{23/12} \mathcal{E}^{3/2}}\right).$$

Note that L_f and L_h show up in the coefficient of lower order of $1/\mathcal{E}$. 2) In Algorithm 1, if we choose the smooth sampling algorithm to be MALA [52] with the initial μ_0 being a warm start, $\beta = \frac{\mathcal{E}}{D}$, then for any $0 < \mathcal{E} < 2$, the iterate μ_K satisfies $\text{TV}(\mu_K, \pi) \leq \mathcal{E}$ for some $K = \widetilde{\mathcal{O}}\left(\frac{L_{s_\beta}}{\alpha}\sqrt{d}\right)$. If \mathcal{E} is sufficiently small such that $\frac{L_f + RL_h}{\lambda_h^2 D} = \mathcal{O}(1/\mathcal{E})$, we get

$$K = \widetilde{\mathcal{O}}\left(\frac{\lambda_h^2 D\sqrt{d}}{\alpha \mathcal{E}}\right).$$

We need the assumption $\langle h(\cdot), y \rangle$ is convex for any $y \in \mathcal{Y}$ since it not only promises s(x) is convex, but also $s_{\beta}(x)$. Thus we can utilize the convergence result for strong log-concave smooth sampling. However, this assumption does not necessarily restrict h(x) to be linear. Here, we give several

¹Here, higher-order smoothness means the condition that is higher than bounded Hessian.

examples where this assumption hold: (i) h(x) = Ax, (ii) $h_i(x)$ is convex for $\forall i$, and $\mathcal{Y} \subset \mathbb{R}^n_+$, (iii) $h_i(x)$ is concave for $\forall i$, and $\mathcal{Y} \subset \mathbb{R}^n_-$, where \mathbb{R}^n_+ means the non-negative part of \mathbb{R}^n and \mathbb{R}^n_- means the non-positive part.

4.2 Log-concavity

Denote the second moment of π is M_{π} . Under the log-concavity condition, the smooth sampling algorithm with best dimension dependency [9, Corollary 6] is the proximal algorithm. Similar to Section 4.1, MALA can achieve the better accuracy order [8] and worse dimension order compared to the proximal algorithm. For the sake of conciseness, we do not list MALA result in this section.

Theorem 2. Assume f is convex, and $\langle h(\cdot), y \rangle$ is convex for any $y \in \mathcal{Y}$. In Algorithm 1, if we choose the smooth sampling algorithm to be proximal algorithm [30] with $\beta = \frac{\mathcal{E}}{D}$, then for any $\mathcal{E} > 0$, the iterate μ_K satisfies $\text{TV}(\mu_K, \pi) \leq \mathcal{E}$ within $K = \mathcal{O}\left(\frac{L_{s_\beta} dW_2^2(\mu_0, \pi_\beta)}{\mathcal{E}^2}\right)$ iterations in expectation. For simplification, assume that $W_2^2(\mu_0, \pi_\beta) = \mathcal{O}(M_\pi)$, and \mathcal{E} is sufficiently small such

that $\frac{L_f + RL_h}{\lambda_h^2 D} = O(1/\mathcal{E})$, then we have

$$K = \mathcal{O}\left(\frac{\lambda_h^2 D M_\pi d}{\mathcal{E}^3}\right). \tag{8}$$

One may notice that an additional term M_{π} shows up in Theorem 2. This is because in Theorem 1, this term is absorbed into logarithm. Generally, M_{π} would scale with dimension d. For example $M_{\pi} = \mathcal{O}(d)$ for a product measure. One needs to take this additional dimension dependency into account when applying Theorem 2.

The randomness of K in (8) comes from the rejection sampling step in proximal algorithm Lee et al. [30]. The bound is in expectation but one can also derive a high probability bound by Multiplicative Chernoff bound since each iteration in rejection sampling is independent.

4.3 Log-Sobolev

A notable condition when there is no log-concavity is the LSI condition. For example, the double-well potential $s(x) = ax^4 - bx^2$ would satisfy the LSI condition but it is not convex. For the LSI condition, the convergence analysis still concentrate on LMC [49, 12]. The recent study on proximal algorithm also has the convergence guarantee [9]. After the comparison on their convergence, no one is strictly better than another. So we list the complexity for both algorithms here.

In this part, we will use the second order Rényi divergence, defined as $\mathcal{R}_2(p||q) := \log \left\| \frac{dp}{dq} \right\|_{L^2(q)}^2$, to control the initialization error. It is a strong divergence since it can upper bound both KL divergence, and squared total variation. It has been a popular analysis tool in sampling [12].

Lemma 2. If $\pi(x)$ satisfies (LSI) with constant C_{π} , then π_{β} satisfies (LSI) with constant $C_{\beta} = C_{\pi} \exp{(4\beta D)}$.

Remark 2. Although C_{β} has exponential dependence on D, we will choose $\beta = \frac{\mathcal{E}}{D}$ in the algorithm such that $\exp(4\beta D) = \exp(4\mathcal{E}) \approx 1$. In this manner, C_{β} has the same order of C_{π} .

With Lemma 2, we are able to apply the convergence of smooth sampling algorithms in our scheme and reach Theorem 3.

Theorem 3. Assume $\pi(x)$ satisfies (LSI) with constant C_{π} . In Algorithm 1, if we choose the smooth sampling algorithm to be proximal algorithm [30], $\beta = \frac{\mathcal{E}}{D}$, then for any $\mathcal{E} > 0$, the iterate μ_K satisfies $\text{TV}(\mu_K, \pi) \leq \mathcal{E}$ within $K = \mathcal{O}\left(L_{s_\beta}C_\beta d\log(2/\mathcal{E}^2)\right)$ iterations in expectation. Assume \mathcal{E} is sufficiently small such that $\frac{L_f + RL_h}{\lambda_h^2 D} = \mathcal{O}(1/\mathcal{E})$, we obtain

$$K = \widetilde{\mathcal{O}}\left(\frac{\lambda_h^2 D C_\pi d}{\mathcal{E}}\right).$$

4.4 Example

In this section, we present case study to illustrate our algorithm. We consider the case where s(x) is the summation of a quadratic term and the maximum of absolute values (also see Section 4.4 of [40])

$$s(x) = \|x\|_{\mathcal{X}}^2 + \max_{1 \le j \le m} \{ |\langle a_j, x \rangle - b^{(j)}| \}, \quad a_j \in \mathbb{R}^d, \ b^{(j)} \in \mathbb{R},$$
(9)

where we choose $\|\cdot\|_{\mathcal{X}}$ as Euclidean norm $\|\cdot\|_2$. It is not difficult to find that $\lambda_{\rho} = \max_{1 \le j \le m} \|a_j\|_2$ since it is a piece-wise affine function. To write (9) in the form of (1), we denote the matrix A with rows $a_j, j = 1, \ldots, m$, and $b = (b_1, \ldots, b_m)$. We also choose n = 2m, $\mathcal{Y} = \Delta_n$, where Δ_n is the probability simplex on \mathbb{R}^n . Then with $\widehat{A} = \begin{pmatrix} A \\ -A \end{pmatrix}$ and $\widehat{b} = \begin{pmatrix} b \\ -b \end{pmatrix}$,

$$s(x) = \|x\|_2^2 + \max_{y \in \Delta_n} \{ \langle \widehat{A}x, y \rangle - \langle \widehat{b}, y \rangle \}.$$

This potential is associated with a Bayesian inference problem with a Gaussian prior and likelihood consists of (absolute values of) linear functions. The goal is to infer a posterior distribution that would optimize the worst-case performance.

To construct the smoothed $s_{\beta}(x)$, we choose $\|\cdot\|_{\mathcal{Y}}$ as 1-norm $\|\cdot\|_1$, y_0 is the uniform distribution, and $\ell(y) = \log(n) + \sum_{j=1}^n y^{(j)} \log y^{(j)}$. Thus by Lemma 3 in Nesterov [40], we result in $\sigma = 1$ and $D = \log(n)$. Clearly, s(x) is strongly-convex, and we have strongly-convex constant $\alpha = 2$, smoothness $L_f = 2$, $L_h = 0$, and Lipschitz

$$\lambda_h = \|\widehat{A}\|_{\mathcal{X}, \mathcal{Y}} = \max_x \{ \max_{1 \le j \le m} |\langle a_j, x \rangle| : \|x\|_2 = 1 \} = \max_{1 \le j \le m} \|a_j\|_2 = \lambda_\rho$$

Putting these together in Theorem 1 bound, with randomized midpoint KLMC, it is enough to have $\widetilde{O}\left(\frac{\lambda_h^{7/3}d^{1/3}}{\mathcal{E}^{5/3}}\right)$, however, the best bound up to now for strongly-convex non-smooth sampling is $\widetilde{O}\left(\lambda_h^2d\right)$ [33]. If we assume λ_h is dimensional free, we have better convergence result in terms of dimension.

5 Conclusion

We consider sampling from a non-smooth distribution $\pi(x) \propto \exp(-s(x))$ that permits the explicit max structure (1). Inspired by Nesterov smoothing, we firstly transfer the non-smooth potential s(x)to a smooth potential $s_{\beta}(x)$ by adding a strongly convex penalty in the max structure, then we apply the smooth sampling algorithms directly to $\pi_{\beta}(x) \propto \exp(-s_{\beta}(x))$. Our smoothing technique is compatible with any first-order sampling algorithm for smooth potentials. We also show that the error introduced by smoothing can be controlled by a careful choice of smoothing intensity β . Finally, we systematically study the convergence rate under several common conditions, and compare them with the state-of-the-art results in Table 1. Our convergence rates are better than or on par with existing methods if no parameter is hiding dimension dependency.

Limitation and future work In this paper, we only consider one class of non-smooth potential with max structure (1) for smoothing. It is interesting to explore more structures that can benefit from smoothing techniques. In addition, from Table 1, one can see that our accuracy dependency does not achieve state-of-art, since we need to compensate the error caused by smoothing. If some parameter like D is hiding dimension dependency, then our method may not be state-of-art anymore. Finally, in the Nesterov smoothing work [40], the optimal estimate for smooth convex optimization is proved. However, the lower bound of smooth sampling is still unclear and seems more challenging. Only a specific class of sampling algorithms has been shown to have lower bound complexity [52]. It would also be pivotal to derive the lower bound of smooth sampling.

References

- [1] Bakry, D. and Emery, M. Diffusions hypercontractives. In *Seminaire de probabilités XIX* 1983/84, pp. 177–206. Springer, 1985. (Cited on page 6.)
- [2] Barp, A., Da Costa, L., França, G., Friston, K., Girolami, M., Jordan, M. I., and Pavliotis, G. A. Geometric methods for sampling, optimisation, inference and adaptive agents. *arXiv preprint arXiv:2203.10592*, 2022. (Cited on page 2.)
- [3] Ben-Tal, A., El Ghaoui, L., and Nemirovski, A. *Robust optimization*, volume 28. Princeton university press, 2009. (Cited on page 1.)
- [4] Bernton, E. Langevin monte carlo and jko splitting. In *Conference On Learning Theory*, pp. 1777–1798. PMLR, 2018. (Cited on page 2.)
- [5] Böhm, A. and Wright, S. J. Variable smoothing for weakly convex composite functions. *Journal* of optimization theory and applications, 188(3):628–649, 2021. (Cited on page 6.)
- [6] Bubeck, S. et al. Convex optimization: Algorithms and complexity. *Foundations and Trends*® *in Machine Learning*, 8(3-4):231–357, 2015. (Cited on page 5.)
- [7] Chatterji, N., Diakonikolas, J., Jordan, M. I., and Bartlett, P. Langevin monte carlo without smoothness. In *International Conference on Artificial Intelligence and Statistics*, pp. 1716–1726. PMLR, 2020. (Cited on pages 1, 2, 3, and 6.)
- [8] Chen, Y., Dwivedi, R., Wainwright, M. J., and Yu, B. Fast mixing of metropolized hamiltonian monte carlo: Benefits of multi-step gradients. J. Mach. Learn. Res., 21:92–1, 2020. (Cited on page 8.)
- [9] Chen, Y., Chewi, S., Salim, A., and Wibisono, A. Improved analysis for a proximal algorithm for sampling. *arXiv preprint arXiv:2202.06386*, 2022. (Cited on pages 2, 4, 8, and 16.)
- [10] Chen, Z. and Vempala, S. S. Optimal convergence rate of hamiltonian monte carlo for strongly logconcave distributions. *Theory of Computing*, 18(1):1–18, 2022. (Cited on page 4.)
- [11] Cheng, X., Chatterji, N. S., Bartlett, P. L., and Jordan, M. I. Underdamped langevin mcmc: A non-asymptotic analysis. In *Conference on learning theory*, pp. 300–323. PMLR, 2018. (Cited on page 4.)
- [12] Chewi, S., Erdogdu, M. A., Li, M. B., Shen, R., and Zhang, M. Analysis of langevin monte carlo from poincaré to log-sobolev. *arXiv preprint arXiv:2112.12662*, 2021. (Cited on pages 1, 2, and 8.)
- [13] Dalalyan, A. S. Theoretical guarantees for approximate sampling from smooth and log-concave densities. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(3): 651–676, 2017. (Cited on pages 1, 4, and 15.)
- [14] Dalalyan, A. S. and Riou-Durand, L. On sampling from a log-concave density using kinetic langevin diffusions. *Bernoulli*, 26(3):1956–1988, 2020. (Cited on page 4.)
- [15] Diehl, M., Bock, H. G., and Kostina, E. An approximation technique for robust nonlinear optimization. *Mathematical Programming*, 107(1):213–230, 2006. (Cited on page 1.)
- [16] Durmus, A., Moulines, E., and Pereyra, M. Efficient bayesian computation by proximal markov chain monte carlo: when langevin meets moreau. *SIAM Journal on Imaging Sciences*, 11(1): 473–506, 2018. (Cited on pages 1, 2, 3, and 6.)
- [17] Durmus, A., Majewski, S., and Miasojedow, B. Analysis of langevin monte carlo via convex optimization. *The Journal of Machine Learning Research*, 20(1):2666–2711, 2019. (Cited on pages 2 and 3.)
- [18] Dwivedi, R., Chen, Y., Wainwright, M. J., and Yu, B. Log-concave sampling: Metropolishastings algorithms are fast! In *Conference on learning theory*, pp. 793–797. PMLR, 2018. (Cited on page 13.)

- [19] Erdogdu, M. A. and Hosseinzadeh, R. On the convergence of langevin monte carlo: the interplay between tail growth and smoothness. In *Conference on Learning Theory*, pp. 1776–1822. PMLR, 2021. (Cited on page 2.)
- [20] Erdogdu, M. A., Hosseinzadeh, R., and Zhang, S. Convergence of langevin monte carlo in chi-squared and rényi divergence. In *International Conference on Artificial Intelligence and Statistics*, pp. 8151–8175. PMLR, 2022. (Cited on page 4.)
- [21] Fan, J., Taghvaei, A., and Chen, Y. Variational wasserstein gradient flow. *arXiv preprint arXiv:2112.02424*, 2021. (Cited on page 1.)
- [22] Gelman, A., Carlin, J. B., Stern, H. S., and Rubin, D. B. *Bayesian data analysis*. Chapman and Hall/CRC, 1995. (Cited on page 1.)
- [23] Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., Gérard-Marchant, P., Sheppard, K., Reddy, T., Weckesser, W., Abbasi, H., Gohlke, C., and Oliphant, T. E. Array programming with NumPy. *Nature*, 585(7825):357–362, September 2020. doi: 10.1038/s41586-020-2649-2. URL https://doi.org/10.1038/s41586-020-2649-2. (Cited on page 13.)
- [24] Hastings, W. K. Monte carlo sampling methods using markov chains and their applications. 1970. (Cited on page 4.)
- [25] HOHEISEL, T., LABORDE, M., and OBERMAN, A. On proximal point-type algorithms for weakly convex functions and their connection to the backward euler method. (Cited on page 6.)
- [26] Kakade, S., Shalev-Shwartz, S., Tewari, A., et al. On the duality of strong convexity and strong smoothness: Learning applications and matrix regularization. *Unpublished Manuscript*, *http://ttic. uchicago. edu/shai/papers/KakadeShalevTewari09. pdf*, 2(1):35, 2009. (Cited on page 3.)
- [27] Krauth, W. *Statistical mechanics: algorithms and computations*, volume 13. OUP Oxford, 2006. (Cited on page 1.)
- [28] Lau, T. T.-K. and Liu, H. Bregman proximal langevin monte carlo via bregman-moreau envelopes. In *International Conference on Machine Learning*, pp. 12049–12077. PMLR, 2022. (Cited on page 2.)
- [29] Ledoux, M. Concentration of measure and logarithmic sobolev inequalities. In *Seminaire de probabilites XXXIII*, pp. 120–216. Springer, 1999. (Cited on page 6.)
- [30] Lee, Y. T., Shen, R., and Tian, K. Structured logconcave sampling with a restricted gaussian oracle. In *Conference on Learning Theory*, pp. 2993–3050. PMLR, 2021. (Cited on pages 2, 4, and 8.)
- [31] Lehec, J. The langevin monte carlo algorithm in the non-smooth log-concave case. *arXiv* preprint arXiv:2101.10695, 2021. (Cited on page 3.)
- [32] Liang, J. and Chen, Y. A proximal algorithm for sampling from non-smooth potentials. *arXiv* preprint arXiv:2110.04597, 2021. (Cited on pages 1 and 2.)
- [33] Liang, J. and Chen, Y. A proximal algorithm for sampling. *arXiv preprint arXiv:2202.13975*, 2022. (Cited on pages 1, 2, 3, 4, and 9.)
- [34] MacKay, D. J., MacKay, D. J., et al. *Information theory, inference and learning algorithms*. Cambridge university press, 2003. (Cited on page 4.)
- [35] Mangoubi, O. and Vishnoi, N. Dimensionally tight bounds for second-order hamiltonian monte carlo. *Advances in neural information processing systems*, 31, 2018. (Cited on page 4.)
- [36] Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., and Teller, E. Equation of state calculations by fast computing machines. *The journal of chemical physics*, 21(6): 1087–1092, 1953. (Cited on page 4.)

- [37] Neal, R. M. et al. Mcmc using hamiltonian dynamics. *Handbook of markov chain monte carlo*, 2(11):2, 2011. (Cited on page 4.)
- [38] Nemirovski, A. and Yudin, D. Problem complexity and method efficiency in optimization. 1983. (Cited on page 3.)
- [39] Nesterov, Y. Introductory lectures on convex optimization: A basic course, volume 87. Springer Science & Business Media, 2003. (Cited on page 3.)
- [40] Nesterov, Y. Smooth minimization of non-smooth functions. *Mathematical programming*, 103 (1):127–152, 2005. (Cited on pages 2, 3, 4, 5, and 9.)
- [41] Nguyen, D., Dang, X., and Chen, Y. Unadjusted langevin algorithm for non-convex weakly smooth potentials. arXiv preprint arXiv:2101.06369, 2021. (Cited on page 2.)
- [42] Otto, F. and Villani, C. Generalization of an inequality by talagrand and links with the logarithmic sobolev inequality. *Journal of Functional Analysis*, 173(2):361–400, 2000. (Cited on page 6.)
- [43] Parisi, G. Correlation functions and computer simulations. *Nuclear Physics B*, 180(3):378–384, 1981. (Cited on page 4.)
- [44] Roberts, G. O. and Tweedie, R. L. Exponential convergence of langevin distributions and their discrete approximations. *Bernoulli*, pp. 341–363, 1996. (Cited on page 4.)
- [45] Salim, A. and Richtárik, P. Primal dual interpretation of the proximal stochastic gradient langevin algorithm. *Advances in Neural Information Processing Systems*, 33:3786–3796, 2020. (Cited on pages 1, 2, and 3.)
- [46] Sanz-Serna, J. M. and Zygalakis, K. C. Wasserstein distance estimates for the distributions of numerical approximations to ergodic stochastic differential equations. *Journal of Machine Learning Research*, 22(242):1–37, 2021. (Cited on page 4.)
- [47] Shen, R. and Lee, Y. T. The randomized midpoint method for log-concave sampling. Advances in Neural Information Processing Systems, 32, 2019. (Cited on pages 4, 7, 13, and 15.)
- [48] Sinha, A., Namkoong, H., and Duchi, J. Certifying some distributional robustness with principled adversarial training. In *International Conference on Learning Representations*, 2018. URL https://openreview.net/forum?id=Hk6kPgZA-. (Cited on page 14.)
- [49] Vempala, S. and Wibisono, A. Rapid convergence of the unadjusted langevin algorithm: Isoperimetry suffices. Advances in neural information processing systems, 32, 2019. (Cited on pages 1, 4, and 8.)
- [50] Villani, C. Topics in optimal transportation. 2003. (Cited on pages 6 and 14.)
- [51] Wibisono, A. Sampling as optimization in the space of measures: The langevin dynamics as a composite optimization problem. In *Conference on Learning Theory*, pp. 2093–3027. PMLR, 2018. (Cited on page 2.)
- [52] Wu, K., Schmidler, S., and Chen, Y. Minimax mixing time of the metropolis-adjusted langevin algorithm for log-concave sampling. *arXiv preprint arXiv:2109.13055*, 2021. (Cited on pages 4, 7, 9, and 16.)

A Numerical experiments

We consider sampling from strongly log-concave distribution $\pi(x) \propto \exp(-s(x))$ in Section 4.4

$$s(x) = \|x\|_2^2 + \max_{1 \le j \le m} \{ |\langle a_j, x \rangle - b^{(j)}| \}, \quad a_j \in \mathbb{R}^d, \ b^{(j)} \in \mathbb{R}$$
$$= \|x\|_2^2 + \max_{y \in \Delta_n} \{ \langle \widehat{A}x, y \rangle - \langle \widehat{b}, y \rangle \}.$$

We normalize a_j , $\forall j$ such that $\lambda_h = 4$. We choose the accuracy $\mathcal{E} = 0.1$, the dimension of y is n = 10. In this example, with prox-function $\ell(y) = \log(n) + \sum_{j=1}^{n} y^{(j)} \log y^{(j)}$, we can solve y_β exactly:

$$y_{\beta} \propto \exp\left(\frac{\widehat{A}x - \widehat{b}}{\beta}\right).$$

This aligns with our assumption that we can solve y_β in $\mathcal{O}(1)$. We select randomized midpoint KLMC Shen & Lee [47] as the smooth sampling method in Algorithm 1. Since this example involves a considerable amount of matrix multiplication, we conduct experiments on a GPU card NVIDIA GeForce 2080Ti with 11GB memory.

A.1 Dimension dependency

Since the total variation error is difficult to calculate in high dimension, we follow the setup of Section 4.1 in Dwivedi et al. [18] and consider using the quantile as the stopping criteria. Specifically, we select 25%, 50%, 75% quantiles on the first coordinate direction to compare. Unlike the Gaussian example in Dwivedi et al. [18], we do not have access to the ground truth marginal distribution for x_1 and can not calculate the real quantile easily. To solve this issue, we follow the experiment setup in Shen & Lee [47] and run the sampling algorithm for K = 50000 steps, which is much larger than the mixing time of this example. Then we take the sample quantile at the step K = 50000 as the ground truth. We use the function numpy.quantile [23] with 5000 samples to calculate the sample quantile. We use step size 0.1 throughout all dimensions.

We repeat the experiments for 5 times with different random seeds and report the error bar in Figure 1. The experimental results align with the reference well, which confirms our complexity analysis for strongly log-concave sampling in Theorem 1 (1).




Figure 1: We report the number of iterations to reach a fixed error tolerance. The reference dashed line is $\tau(d) = 2700 \ d^{1/3} \cdot \log(d)$.

A.2 Trace plot

To investigate the rate of convergence in a single run, we also give the trace plot of the first coordinate for both our method and Langevin Monte Carlo (3) in Figure 2. We conduct the experiments at the dimension d = 17. The comparison shows that our algorithm converges faster than LMC in high dimension.




Figure 2: The trajectories of 10 independent runs simulated by LMC and our algorithm at dimension d = 17.

B Proof of technical lemmas and propositions

B.1 Proof of Lemma 1

Proof. Define $S(x,y) = f(x) + \langle h(x), y \rangle - g(y) - \beta \ell(y)$. According to the definition of $\ell(y)$, $S(x, \cdot)$ is $\beta \sigma$ -strongly-concave with respect to $\|\cdot\|_{\mathcal{Y}}$.

Moreoever, since f and h are smooth, we can get

$$\begin{aligned} \|\nabla_{x}S(x_{1},y) - \nabla_{x}S(x_{2},y)\|_{\mathcal{X}}^{*} &= \|\nabla f(x_{1}) - \nabla f(x_{2}) + J_{h}^{\top}(x_{1})y - J_{h}^{\top}(x_{2})y\|_{\mathcal{X}}^{*} \\ &\leq \|\nabla f(x_{1}) - \nabla f(x_{2})\|_{\mathcal{X}}^{*} + \|J_{h}^{\top}(x_{1})y - J_{h}^{\top}(x_{2})y\|_{\mathcal{X}}^{*} \\ &\leq L_{f}\|x_{1} - x_{2}\|_{\mathcal{X}} + \|J_{h}(x_{1}) - J_{h}(x_{2})\|_{\mathcal{X},\mathcal{Y}}\|y\|_{\mathcal{Y}} \\ &\leq (L_{f} + L_{h}R)\|x_{1} - x_{2}\|_{\mathcal{X}}, \end{aligned}$$
(10)

where we use the definition of matrix norm $\|\cdot\|_{\mathcal{X},\mathcal{Y}}$ in the second last inequality and $\|y\| \leq R$ in the last inequality. On the other hand, by h is Lipschitz, we have

$$\|\nabla_{x}S(x,y_{1}) - \nabla_{x}S(x,y_{2})\|_{\mathcal{X}}^{*} = \|J_{h}^{\top}(x)(y_{1} - y_{2})\|_{\mathcal{X}}^{*}$$

$$\leq \|J_{h}^{\top}(x)\|_{\mathcal{X},\mathcal{Y}}\|y_{1} - y_{2}\|_{\mathcal{Y}}$$

$$\leq \lambda_{h}\|y_{1} - y_{2}\|_{\mathcal{Y}}.$$
 (11)

Similarly, it holds that

$$|\nabla_y S(x_1, y) - \nabla_y S(x_2, y)|| = ||h(x_1) - h(x_2)|| \le \lambda_h ||x_1 - x_2||.$$
(12)

Together with Sinha et al. [48, Lemma 1], (10) (11) (12) imply that $\nabla s_{\beta}(x) = \nabla f(x) + J_h^{\top}(x)y_{\beta}(x)$ and $\|\nabla s_{\beta}(x_1) - \nabla s_{\beta}(x_2)\| \leq \left(L_f + RL_h + \frac{\lambda_h^2}{\beta\sigma}\right)\|x_1 - x_2\|.$

B.2 Proof of Lemma 2

Proof. We can write

$$\pi_{\beta}(x) = \frac{Z_0}{Z_{\beta}} \exp(-(s_{\beta}(x) - s(x)))\pi(x) = \exp\left(-\left(s_{\beta}(x) - s(x) + \log\left(\frac{Z_{\beta}}{Z_0}\right)\right)\right)\pi(x),$$

where $Z_0 = \int_{\mathbb{R}^d} \exp(-s(x)) dx$ and $Z_\beta = \int_{\mathbb{R}^d} \exp(-s_\beta(x)) dx$ are the normalization constants. By (13), we have $-\beta D \le s_\beta(x) - s(x) \le 0$, and

$$Z_0 \le Z_\beta = \int \exp(-s(x)) \exp(s(x) - s_\beta(x)) dx \le Z_0 \exp(\beta D)$$

$$\Rightarrow \ 0 \le \log\left(\frac{Z_\beta}{Z_0}\right) \le \beta D.$$

This implies that $-\beta D \leq s_{\beta}(x) - s(x) + \log\left(\frac{Z_{\beta}}{Z_{0}}\right) \leq \beta D$. Then the result is immediate once noticing that LSI is stable under bounded perturbation [50, Theorem 9.9 (ii)].

B.3 Proof of Proposition 1

Proof. According to the definition $\phi_{\beta}(x, y) := \langle h(x), y \rangle - g(y) - \beta \ell(y)$, it's clear that $\phi_{\beta}(x, y)$ is strongly-concave w.r.t. y when $\beta > 0$. Thus $\phi_{\beta}(x, y_0(x)) - \phi_{\beta}(x, y_{\beta}(x)) \le 0$. This implies

$$0 \le s(x) - s_{\beta}(x) = \phi_{\beta}(x, y_0(x)) + \beta \ell(y_0(x)) - \phi_{\beta}(x, y_{\beta}(x)) \le \beta \ell(y_0(x)) \le \beta D.$$
(13)

This further leads to

$$\int (s(x) - s_{\beta}(x))^2 \pi_{\beta}(x) \mathrm{d}x \leq \int \beta^2 D^2 \pi_{\beta}(x) \mathrm{d}x = \beta^2 D^2.$$

Then by Pinsker's inequality (7) and Lemma 3 in Dalalyan [13],

$$\mathrm{TV}(\pi,\pi_{\beta}) \leq \sqrt{\frac{1}{2}\mathrm{KL}(\pi_{\beta}\|\pi)} \leq \frac{1}{2}\sqrt{\int (s(x) - s_{\beta}(x))^2 \pi_{\beta}(x) dx} \leq \frac{\beta D}{2}.$$

B.4 Proof of Proposition 2

Proof. According to the definition $\phi_{\beta}(x, y) := \langle h(x), y \rangle - g(y) - \beta \ell(y)$, it's clear that $\phi_{\beta}(x, y)$ is strongly-concave w.r.t. y when $\beta > 0$. Thus $\phi_{\beta}(x, y_0(x)) - \phi_{\beta}(x, y_{\beta}(x)) \le 0$. This implies

$$0 \le s(x) - s_{\beta}(x) = \phi_{\beta}(x, y_0(x)) + \beta \ell(y_0(x)) - \phi_{\beta}(x, y_{\beta}(x)) \le \beta \ell(y_0(x)) \le \beta D.$$

This further leads to

$$\int (s(x) - s_{\beta}(x))^2 \pi_{\beta}(x) \mathrm{d}x \le \int \beta^2 D^2 \pi_{\beta}(x) \mathrm{d}x = \beta^2 D^2.$$

Then by Talagrand's inequality and Lemma 3 in Dalalyan [13],

$$W_2(\pi_\beta, \pi) \le \sqrt{2C_\pi \operatorname{KL}(\pi_\beta \| \pi)} \le \sqrt{C_\pi \int (s(x) - s_\beta(x))^2 \pi_\beta(x) dx} \le \sqrt{C_\pi} \beta D.$$

C Proof of Theorems

We provide the full proof for Theorem 1 together with other Theorems in this section.

C.1 Proof of Theorem 1

Proof. Since $\langle h(\cdot), y \rangle$ is convex for any $y \in \mathcal{Y}$, it holds that $\psi_{\beta}(x) := \max_{y \in \mathcal{Y}} \{\langle h(x), y \rangle - g(y) - \beta \ell(y)\}$ is convex for both $\beta > 0$ and $\beta = 0$. This is because the maximization of convex function is still convex.

1) Remember that we have the relationship: α -strongly-convex s(x) implies $\pi(x)$ satisfies (LSI) with constant $1/\alpha$. Thus we can bound $W_2(\pi_\beta, \pi) \leq \beta D/\sqrt{\alpha}$ according to Proposition 2. Plugging in the choice $\beta = \frac{\sqrt{\alpha}\mathcal{E}}{2D}$ further gives $W_2(\pi_\beta, \pi) \leq \frac{\mathcal{E}}{2}$.

On the other hand, with the choice

$$K = \widetilde{\mathcal{O}}\left(\left(\frac{L_{s_{\beta}}}{\alpha}\right)^{7/6} \left(\frac{2}{\mathcal{E}}\sqrt{\frac{d}{\alpha}}\right)^{1/3} + \left(\frac{L_{s_{\beta}}}{\alpha}\right) \left(\frac{2}{\mathcal{E}}\sqrt{\frac{d}{\alpha}}\right)^{2/3}\right),\tag{14}$$

we apply the convergence result in Theorem 3 in Shen & Lee [47] and have $W_2(\mu_K, \pi_\beta) \leq \frac{\varepsilon}{2}$. Putting these two inequalities together gives

$$W_2(\mu_K, \pi) \le W_2(\mu_K, \pi_\beta) + W_2(\pi_\beta, \pi) \le \frac{\mathcal{E}}{2} + \frac{\mathcal{E}}{2} = \mathcal{E}.$$

To simplify (14), we plug in $L_{s_{\beta}} = L_f + RL_h + \frac{\lambda_h^2}{\beta\sigma} = L_f + RL_h + \frac{2\lambda_h^2 D}{\sqrt{\alpha\sigma\mathcal{E}}}$ and assume \mathcal{E} is sufficiently small such that $\frac{L_f + RL_h}{\lambda_h^2 D} = \mathcal{O}(1/\mathcal{E})$. Then $L_{s_{\beta}} = \mathcal{O}\left(\frac{\lambda_h^2 D}{\sqrt{\alpha\sigma\mathcal{E}}}\right)$ and

$$\begin{split} \widetilde{\mathcal{O}}\left(\left(\frac{L_{s_{\beta}}}{\alpha}\right)^{7/6} \left(\frac{2}{\mathcal{E}}\sqrt{\frac{d}{\alpha}}\right)^{1/3} + \left(\frac{L_{s_{\beta}}}{\alpha}\right) \left(\frac{2}{\mathcal{E}}\sqrt{\frac{d}{\alpha}}\right)^{2/3}\right) \\ &= \widetilde{\mathcal{O}}\left(\frac{\left(\frac{\lambda_{h}^{2}D}{\sqrt{\alpha\sigma\mathcal{E}}}\right)^{7/6}}{\alpha^{4/3}} \cdot \frac{d^{1/6}}{\mathcal{E}^{1/3}} + \frac{\lambda_{h}^{2}D}{\alpha\mathcal{E}^{2/3}} \cdot \left(\frac{d}{\alpha}\right)^{1/3}\right) \\ &= \widetilde{\mathcal{O}}\left(\frac{\lambda_{h}^{7/3}D^{7/6}d^{1/6}}{\alpha^{23/12}\mathcal{E}^{3/2}\sigma^{7/6}} + \frac{\lambda_{h}^{2}Dd^{1/3}}{\alpha^{11/6}\mathcal{E}^{5/3}\sigma}\right). \end{split}$$

2) With the choice

$$K = \widetilde{\mathcal{O}}\left(\frac{L_{s_{\beta}}}{\alpha}\sqrt{d}\right) \tag{15}$$

we apply the convergence result in Theorem 1 in Wu et al. [52] and have $\text{KL}(\mu_K || \pi_\beta) \leq \mathcal{E}^2/2$. By Proposition 1 and the choice $\beta = \frac{\mathcal{E}}{D}$, there is $\text{TV}(\pi_\beta . \pi) \leq \frac{\mathcal{E}}{2}$. Further, by triangular inequality, and Pinsker's inequality,

$$\mathrm{TV}(\mu_K, \pi) \leq \mathrm{TV}(\mu_K, \pi_\beta) + \mathrm{TV}(\pi_\beta . \pi) \stackrel{(7)}{\leq} \sqrt{\frac{1}{2} \mathrm{KL}(\mu_K \| \pi_\beta)} + \frac{\mathcal{E}}{2} \leq \frac{\mathcal{E}}{2} + \frac{\mathcal{E}}{2} = \mathcal{E}.$$

To simplify (15), we plug in $L_{s_{\beta}} = L_f + RL_h + \frac{\lambda_h^2}{\beta\sigma} = L_f + RL_h + \frac{\lambda_h^2 D}{\sigma\mathcal{E}}$ and assume \mathcal{E} is sufficiently small such that $\frac{L_f + RL_h}{\lambda_h^2 D} = \mathcal{O}(1/\mathcal{E})$. Then $L_{s_{\beta}} = \mathcal{O}\left(\frac{\lambda_h^2 D}{\sigma\mathcal{E}}\right)$ and

$$\widetilde{\mathcal{O}}\left(\frac{L_{s_{\beta}}}{\alpha}\sqrt{d}\right) = \widetilde{\mathcal{O}}\left(\frac{\lambda_{h}^{2}D\sqrt{d}}{\alpha\mathcal{E}}\right).$$

C.2 Proof of Theorem 2

Proof. Since $\langle h(\cdot), y \rangle$ is convex for any $y \in \mathcal{Y}$, it holds that $\psi_{\beta}(x) := \max_{y \in \mathcal{Y}} \{\langle h(x), y \rangle - g(y) - \beta \ell(y) \}$ is also convex for both $\beta > 0$ and $\beta = 0$. This is because the maximization of convex function is still convex.

With the choice of K, we apply the convergence result in Corollary 6 in Chen et al. [9] and have $\operatorname{KL}(\mu_K \| \pi_\beta) \leq \mathcal{E}^2/2$. By Proposition 1 and the choice $\beta = \frac{\mathcal{E}}{D}$, there is $\operatorname{TV}(\pi_\beta.\pi) \leq \frac{\mathcal{E}}{2}$. The rest of the proof follows part 2) proof of Theorem 1.

C.3 Proof of Theorem 3

Proof. Since $\pi(x)$ satisfies (LSI) with constant C_{π} , by Lemma 2, we have π_{β} satisfies (LSI) with constant $C_{\pi} \exp(4\beta D)$. With the choice

$$K = \mathcal{O}\left(L_{s_{\beta}}C_{\beta}d\log(2/\mathcal{E}^{2})\right),\tag{16}$$

we apply the convergence result in Corollary 7 in Chen et al. [9] and have $\text{KL}(\mu_K || \pi_\beta) \leq \mathcal{E}^2/2$. By Proposition 1 and the choice $\beta = \frac{\mathcal{E}}{D}$, there is $\text{TV}(\pi_\beta . \pi) \leq \frac{\mathcal{E}}{2}$. Further, by triangular inequality, and Pinsker's inequality,

$$\mathrm{TV}(\mu_K, \pi) \leq \mathrm{TV}(\mu_K, \pi_\beta) + \mathrm{TV}(\pi_\beta, \pi) \stackrel{(7)}{\leq} \sqrt{\frac{1}{2}\mathrm{KL}(\mu_K \| \pi_\beta)} + \frac{\mathcal{E}}{2} \leq \frac{\mathcal{E}}{2} + \frac{\mathcal{E}}{2} = \mathcal{E}.$$

To simplify (16), we plug in $L_{s_{\beta}} = L_f + RL_h + \frac{\lambda_h^2}{\beta\sigma} = L_f + RL_h + \frac{\lambda_h^2 D}{\sigma \mathcal{E}}$ and assume \mathcal{E} is sufficiently small such that $\frac{L_f + RL_h}{\lambda_h^2 D} = \mathcal{O}(1/\mathcal{E})$. Then $L_{s_{\beta}} = \mathcal{O}\left(\frac{\lambda_h^2 D}{\sigma \mathcal{E}}\right)$ and β is also small enough to do the Taylor approximation $C_{\beta} = C_{\pi} \exp(4\beta D) \approx C_{\pi}$. Finally, plugging in these pieces results in

$$K = \mathcal{O}\left(L_{s_{\beta}}C_{\beta}d\log(2/\mathcal{E}^{2})\right) = \widetilde{\mathcal{O}}\left(\frac{\lambda_{h}^{2}DC_{\pi}d}{\sigma\mathcal{E}}\right).$$