

Connecting Cosmic Inflation to Particle Physics with LiteBIRD, CMB-S4, EUCLID and SKA

Marco Drewes^{1*} and Lei Ming^{2,1,3†}

¹Centre for Cosmology, Particle Physics and Phenomenology,

Université catholique de Louvain, Louvain-la-Neuve B-1348, Belgium

²School of Physics, Sun Yat-Sen University, Guangzhou 510275, China and

³School of Physics, Nanjing University, Nanjing, 210093, China

We show that next generation Cosmic Microwave Background experiments will be capable of the first ever measurement of the inflaton coupling to other particles, opening a new window to probe the connection between cosmic inflation and particle physics. This sensitivity is based on the impact that the reheating phase after cosmic inflation has on the redshifting of cosmic perturbations. For our analysis we introduce a simple analytic method to estimate the sensitivity of future CMB observations to the reheating temperature and the inflaton coupling. Applying our method to LiteBIRD and CMB-S4 we find that, within a given model of inflation, these missions have the potential to impose both an upper and a lower bound on the inflaton coupling. Further improvement can be achieved if CMB data is combined with optical and 21cm surveys. Our results demonstrate the potential of future observations to constrain microphysical parameters that can provide an important clue to understand how a given model of inflation may be embedded in a more fundamental theory of nature.

Introduction: The current concordance model of cosmology, known as Λ CDM model, can explain almost all properties of the observable universe at an astonishing accuracy with only a handful of free parameters [1, 2].¹ Leaving aside the composition of the Dark Matter (DM), the model is firmly based on the Standard Model (SM) of particle physics and the theory of General Relativity (GR), implying that the most fundamental laws of nature that we know from earth [4] hold in the most distant regions of the observable universe.²

However, to date it is unknown what mechanism set the initial conditions for the hot big bang, including the initial overall geometry of the observable Universe and the temperature T_{re} of the primordial plasma at the onset of the radiation dominated epoch.³ The former – in particular overall homogeneity, isotropy, and spacial flatness reflected in the Cosmic Microwave Background (CMB) – is amongst the most compelling mysteries of modern cosmology. *Cosmic inflation* [7–9] offers an elegant solution for these problems and can in addition explain the observed correlations amongst the small perturbations in the CMB. However, very little is known about the mechanism that may have driven the exponential growth of the scale

factor a . A wide range of theoretical models of inflation exist (see e.g. [10] for a partial list), but the observational evidence is not conclusive enough to clearly single out one of them. Moreover, even less is known about the embedding of cosmic inflation into a more fundamental theory of nature and its connection to theories of particle physics beyond the SM. In the next decade the observational situation will change drastically. Upgrades at the South Pole Observatory [11] and the Simmons Observatory [12] aim at pushing the uncertainty in the scalar-to-tensor ratio r down to $\delta r \sim 3 \times 10^{-3}$. In the 2030s JAXA’s LiteBIRD satellite [13] and the ground-based CBM-S4 program [14] are expected to further reduce this to $\delta r < 10^{-3}$ for $r = 0$.

In the present work we for the first time quantify the ability of these missions to probe the connection between inflation and particle physics. We utilise the impact of *cosmic reheating* [15–21] after inflation on the expansion history, i.e., the dissipative transfer of energy from the inflationary sector to other degrees of freedom that filled the universe with particles and determined T_{re} . While the only known direct messenger from the reheating epoch would be gravitational waves (cf. [22]),⁴ it can be studied indirectly with CMB observables [27–29] due to the impact of the modified equation of state w during reheating [30, 31] on the post-inflationary expansion history. This has motivated studies in various models.⁵ Reheating

* marco.drewes@uclouvain.be

† minglei@mail.sysu.edu.cn (Corresponding author)

¹ See [3] for a recent summary of tensions and anomalies in the Λ CDM model.

² The value of the cosmological constant also has no microscopic explanation in the SM, but it can be regarded as a free parameter in GR and quantum field theory, reducing this issue to a theoretical problem of naturalness (cf. [5]) rather than a failure to describe data.

³ Another initial condition that cannot be explained within the SM but is not addressed in the present work is the initial matter-antimatter asymmetry (cf. [6]).

⁴ The thermal graviton background [23] can in principle be used to probe T_{re} [24, 25], but this is very challenging in practice [26].

⁵ Examples include natural inflation [32–35], power law [36–40] and polynomial potentials [33, 41–43], Starobinski inflation [33], Higgs inflation [33, 36, 44], hilltop type inflation [33, 36], axion inflation [36, 45], curvaton models [46], α -attractor inflation [47–55], tachyon inflation [56], inflection point inflation [57], fiber inflation [58], Kähler moduli inflation [59, 60], and other SUSY models [36, 61].

is inherently sensitive to the inflaton couplings to other fields, i.e., microphysical parameters that connect inflation to particle physics [62–64], as these interactions mediated the energy transfer. However, past studies have almost exclusively focused on T_{re} , ignoring the possibility to constrain microphysical (particle physics) parameters. The fundamental limitations on the possibility of constraining the microphysical coupling constant \mathbf{g} associated with the interaction through which the universe was primarily reheated were laid out in [64], where it was estimated that such a measurement may be within reach of next-generation instruments. However, neither there nor in any of the few phenomenological works addressing the relation to microphysical parameters [47, 50, 55] a systematic study was performed to quantify the feasibility of such a measurement with realistic instrumental sensitivities. In the present work we introduce a simple analytic method to quantify the sensitivity of observations to \mathbf{g} for given instrumental sensitivities to the amplitude of the scalar perturbations in the CMB A_s , the spectral index n_s and the tensor-to-scalar ratio r .⁶ We apply this method to show for the first time that upcoming observations will be capable of the first ever measurement of both \mathbf{g} and T_{re} , where we define a measurement as the ability to impose both an upper and a lower bound on the respective quantity. In this Letter we present the main results of our research, a more detailed analysis is presented in [65].

Imprint of reheating in the CMB: The primary goal of this work is to quantify constraints on \mathbf{g} and T_{re} from current and future CMB data. We consider inflationary models that can effectively be described by a single field Φ and assume that the effective single field description holds throughout both, inflation and the reheating epoch.⁷ Defining $\varphi = \langle \Phi \rangle$ as the quantum statistical expectation value of $\Phi = \varphi + \phi$ with fluctuations ϕ and φ_{end} as its value at the end of inflation, the energy density at the end of inflation $\rho_{\text{end}} \simeq \frac{4}{3}\mathcal{V}(\varphi_{\text{end}}) \equiv \frac{4}{3}\mathcal{V}_{\text{end}}$ and the spectrum of primordial perturbations are fixed by the effective potential $\mathcal{V}(\varphi)$.⁸ Assuming a standard cosmic history after reheating (and leaving aside foreground effects), the observable spectrum of CMB perturbations can be predicted from $\mathcal{V}(\varphi)$ once the expansion history during reheating is known. The latter requires knowledge of the duration of the reheating epoch N_{re} in terms of expansion e -folds N (defined as the logarithm of the scale factor growth) and the average equation of state during reheating $\bar{w}_{\text{re}} = \frac{1}{N_{\text{re}}} \int_0^{N_{\text{re}}} w(N) dN$. Since the total energy density of

the universe ρ is still dominated by the energy density ρ_φ of φ during reheating, \bar{w}_{re} is determined by specification of $\mathcal{V}(\varphi)$, and N_{re} is the only relevant quantity that is not fixed by the choice of $\mathcal{V}(\varphi)$, i.e., is sensitive to \mathbf{g} . Within a given model of inflation one can obtain information about N_{re} by comparing the observed CMB spectrum to the model's prediction. Using the general redshifting relation $\rho \propto \exp(-3N(1+w))$, this can then be translated into a constraint on the energy density at the end of reheating $\rho_{\text{re}} = \rho_{\text{end}} \exp(-3N_{\text{re}}(1+\bar{w}_{\text{re}}))$ [68], often expressed in terms of an effective reheating temperature defined as $\frac{\pi^2 g_*}{30} T_{\text{re}}^4 \equiv \rho_{\text{re}}$ with g_* the effective number of relativistic degrees of freedom,

$$T_{\text{re}} = \exp \left[-\frac{3(1+\bar{w}_{\text{re}})}{4} N_{\text{re}} \right] \left(\frac{40 \mathcal{V}_{\text{end}}}{g_* \pi^2} \right)^{1/4}. \quad (1)$$

In order to further translate knowledge on N_{re} into knowledge on microphysical parameters, we utilise the fact that reheating ends when $H = \Gamma$, where Γ is an effective dissipation rate for φ and $H = \dot{a}/a$ is the Hubble rate. Together with the Friedmann equation $H^2 = \rho/(3M_{\text{pl}}^2)$ this yields [50, 62]

$$\Gamma|_{\Gamma=H} = \frac{1}{M_{\text{pl}}} \left(\frac{\rho_{\text{end}}}{3} \right)^{1/2} e^{-3(1+\bar{w}_{\text{re}})N_{\text{re}}/2} \quad (2)$$

with $M_{\text{pl}} = 2.435 \times 10^{18}$ GeV the reduced Planck mass. The RHS of (1) and (2) only contain quantities that are either calculable for given $\mathcal{V}(\varphi)$ or can be obtained from CMB observations; we summarise the relations to CMB observables in the appendix. Meanwhile Γ on the LHS depends on microphysical parameters of the particle physics model in which $\mathcal{V}(\varphi)$ is realised.

Measuring the inflaton coupling in the CMB: We classify microphysical parameters in three categories. A *model of inflation* is defined by specifying the effective potential $\mathcal{V}(\varphi)$. Ignoring quantum corrections to the φ -trajectory, this is equivalent to fixing the set of coefficients $\{v_i\}$ of all operators in the action that can be constructed from Φ alone, e.g., by Taylor expanding the inflaton potential around its minimum as $\mathcal{V}(\varphi) = \sum_j \frac{v_j}{j!} \frac{\varphi^j}{\Lambda^{j-4}} = \frac{1}{2} m_\phi^2 \varphi^2 + \frac{g_\phi}{3!} \varphi^3 + \frac{\lambda_\phi}{4!} \varphi^4 + \dots$. The set of *inflaton couplings* $\{\mathbf{g}_i\}$ comprises coupling constants (or Wilson coefficients) associated with operators that are constructed from Φ and other fields. A complete particle physics model contains a much larger set of parameters than the combined set $\{v_i\} \cup \{\mathbf{g}_i\}$, including the masses of the particles produced during reheating as well as their interactions amongst each other and with all other fields. We refer to the set of all parameters in the action that are not contained in $\{v_i\} \cup \{\mathbf{g}_i\}$ as $\{\mathbf{a}_i\}$. This set, e.g., contains the parameters of the SM.

Γ in (2) necessarily depends on the $\{\mathbf{g}_i\}$ and $\{v_i\}$. For instance, for reheating through elementary particle decays, one typically finds $\Gamma = \mathbf{g}^2 m_\phi / c$, with $\mathbf{g} \in \{\mathbf{g}_i\}$

⁶ Throughout we fix $A_s = 10^{-10} e^{3.043}$ [1] and neglect the uncertainty σ_{A_s} ; we checked in [65] that this does not affect the conclusions.

⁷ See [64, 66, 67] and references therein for a discussion of effects that limit the validity of this assumption during reheating.

⁸ Throughout we work at leading order in the slow roll parameters ϵ and η defined after (13); we confirmed in [65] that this approximation does not affect our conclusions.

a coupling constant, $m_\phi \in \{v_i\}$ the inflaton mass, and c a numerical factor. However, in general feedback effects from produced particles on the ongoing reheating process introduce a dependence of Γ on a large sub-set of $\{\mathbf{a}_i\}$, making it impossible to determine \mathbf{g} from the CMB in a model-independent way, i.e., without having to specify the details of the underlying particle physics model and the values of the parameters $\{\mathbf{a}_i\}$.⁹ The conditions under which \mathbf{g} can be constrained model-independently have been studied in detail in [64], a conservative estimate is

$$|\mathbf{g}| \ll \left(\frac{m_\phi}{\varphi_{\text{end}}}\right)^{j-\frac{1}{2}} \min\left(\sqrt{\frac{m_\phi}{M_{pl}}}, \sqrt{\frac{m_\phi}{\varphi_{\text{end}}}}\right) \left(\frac{m_\phi}{\Lambda}\right)^{4-D} \quad (3)$$

$$|v_i| \ll \left(\frac{m_\phi}{\varphi_{\text{end}}}\right)^{j-\frac{5}{2}} \min\left(\sqrt{\frac{m_\phi}{M_{pl}}}, \sqrt{\frac{m_\phi}{\varphi_{\text{end}}}}\right) \left(\frac{m_\phi}{\Lambda}\right)^{4-j} \quad (4)$$

with D the mass dimension of the interaction term under consideration, j the power at which Φ appears in that operator, and Λ a scale that can be identified with m_ϕ for $D \leq 4$ and represents a UV cutoff of the effective theory for $D > 4$. The conditions (3) and (4) ensure that the production of particles proceeds slow enough that redshifting spreads their momenta over a sufficiently broad phase space volume for the occupation numbers in each mode to remain low enough to avoid sizeable feedback effects, such as a parametric resonance. The condition (4) practically restricts the possibility to constrain \mathbf{g} model-independently to scenarios where the φ -oscillations occur in a mildly non-linear regime.

Application to specific models: In the following we apply the previous considerations to two models of inflation, namely radion gauge inflation (RGI) [78, 79] and α -attractor T-models (α -T) [80–83], with the potentials

$$\text{RGI: } \mathcal{V}(\varphi) = M^4 \frac{(\varphi/M_{pl})^2}{\alpha + (\varphi/M_{pl})^2} \quad (5)$$

$$\alpha\text{-T: } \mathcal{V}(\varphi) = M^4 \tanh^{2n} \left(\frac{\varphi}{\sqrt{6\alpha} M_{pl}} \right) \quad (6)$$

The scale M can be expressed in terms of other parameters with the help of (14),

$$\text{RGI: } M = M_{pl} \left(\frac{3\pi^2}{2} r A_s \left(1 + \alpha \frac{M_{pl}^2}{\varphi_k^2} \right) \right)^{1/4}, \quad (7)$$

$$\alpha\text{-T: } M = M_{pl} \left(\frac{3\pi^2}{2} A_s r \right)^{1/4} \tanh^{-\frac{n}{2}} \left(\frac{\varphi_k}{\sqrt{6\alpha} M_{pl}} \right), \quad (8)$$

and condition (4) implies $n = 1$. Within these families of models (16) implies the relations $\alpha =$

$\frac{432r^2}{(8(1-n_s)+r)^2(4(1-n_s)-r)}$ and $\alpha = \frac{4r}{3(1-n_s)(4(1-n_s)-r)}$ for the RGI and α -T models, respectively. This defines a line in the n_s - r plane, the position along which is given by N_{re} (and hence \mathbf{g}), cf. Fig. 1. Condition (4) implies $\alpha > 2.4$ in (5) and $\alpha > 1/4$ in (6). For our analysis we pick $\alpha = 19$ in (5) and $\alpha = 6$ in (6). When conditions (3) and (4) are fulfilled we may parameterise $\Gamma = \mathbf{g}^2 m_\phi / c$ [64] with $(\mathbf{g}, c) = (g/m_\phi, 8\pi)$ for a scalar coupling $g\Phi\chi^2$ [84], $(\mathbf{g}, c) = (y, 8\pi)$ for a Yukawa coupling $y\Phi\psi\psi$ [73], and $(\mathbf{g}, c) = (\sigma m_\phi / \Lambda, 4\pi)$ for an axion-like coupling $\frac{\sigma}{\Lambda} \Phi F_{\mu\nu} \tilde{F}^{\mu\nu}$ [85], where we neglected the produced particles' rest masses. We shall assume a Yukawa coupling y in the following, bounds on other interactions can be obtained by simple rescaling according to c [65].




Figure 1: The diagonal line represents values of n_s and r predicted by the RGI model (upper panel) and α -T model (lower panel) for fixed α , with the inflaton coupling varying along the curve. The black discs indicate the predictions for specific values of a Yukawa coupling y , with $(\log_{10} y, \log_{10} T_{\text{re}}/\text{GeV})$ given in the legend. Conditions (3) and (4) are fulfilled in the green part, the gray parts are ruled out by the conditions $N_{\text{re}} > 0$ and $T_{\text{re}} < T_{\text{BBN}}$. Ellipses indicate current constraints and future sensitivities to n_s and r , N_k is the number of e -folds between the horizon crossing of a perturbation with wave number k and the end of inflation (12).

CMB constraints on the inflaton coupling: With the above considerations and the relations given in the appendix, n_s and r in a given model of inflation are simple functions of $x \equiv \log_{10} \mathbf{g}$. Prior to any measurement of (A_s, n_s, r) it is known that $N_{\text{re}} > 0$ and that there is a lower bound $T_{\text{re}} > T_{\text{BBN}}$ to allow for success-

⁹ This tends to happen due to resonant particle production during the so-called preheating phase (c.f. [69] for a review), but can in principle also occur due to thermal feedback during perturbative reheating [50, 62, 64, 70–77].

ful big bang nucleosynthesis (BBN). When (3) and (4) are fulfilled one can use the standard estimate $T_{\text{re}} \simeq \sqrt{\Gamma M_{\text{pl}}} \times (90/(\pi^2 g_*))^{1/4}$ to obtain a lower bound on the coupling $\mathbf{g} > g_*^{1/4} \sqrt{c T_{\text{BBN}}} / \sqrt{m_\phi M_{\text{pl}}}$, which for plateau models translates into $\mathbf{g} > \sqrt{c} (T_{\text{BBN}}/M_{\text{pl}}) \times (\frac{g_*}{A_{sr}})^{1/4}$ [64]. Hence, we use the prior probability density function (PDF)

$$P(x) = C_1 \theta(T_{\text{re}}(x) - T_{\text{BBN}}) \gamma(x) \theta(N_{\text{re}}(x)), \quad (9)$$

with θ the Heaviside function, and γ a function that allows for a re-weighting of the prior $P(x)$. The constant C_1 can be fixed from the requirement $\int dx P(x) = 1$. We now quantify the gain in knowledge about x that can be obtained from data \mathcal{D} . This gain can be quantified by the posterior distribution $P(x|\mathcal{D}) = P(\mathcal{D}|x)P(x)/P(\mathcal{D})$ with $P(\mathcal{D}) = \int dx P(\mathcal{D}|x)P(x)$. Current constraints from the data \mathcal{D} obtained by Planck and BICEP/Keck [2] can be approximated by the likelihood function

$$P(\mathcal{D}|x) = C_2 \mathcal{N}(n_s, r | \bar{n}_s, \sigma_{n_s}; \bar{r}, \sigma_r) \theta(r) \tilde{\gamma}(x), \quad (10)$$

with $\mathcal{N}(n_s, r | \bar{n}_s, \sigma_{n_s}; \bar{r}, \sigma_r)$ a two-dimensional Gaussian¹⁰ and $\tilde{\gamma}$ another weighting function. The constant C_2 is fixed by normalising $P(\mathcal{D}|x)$ to unity. We fix the fiducial values to $\bar{n}_s = 0.967$, $\bar{r} = 0.01$ and estimate the errors $\sigma_{n_s} = 0.005$, $\sigma_r = 0.018$ based on Fig. 5 in [2]. The result is shown in Fig. 2. While it is known that present data already provides information about the reheating epoch [86], current CMB observations do not provide a significant information gain on x with respect to $P(x)$.

The scalar-to-tensor ratio will be constrained with much higher accuracy in the future [87]. To quantify the expected information gain on x we repeat the analysis for $\bar{r} = 0.02$ with $\sigma_{n_s} = 0.002$ and $\sigma_r = 0.0012$, which reflects the sensitivity anticipated by LiteBIRD [13] or CMB-S4 [14].¹¹ Fig. 2 shows that in both models future data can rule out previously allowed values of T_{re} . In the α -T model the posterior peaks in a region where condition (3) is violated, implying that Γ depends on a potentially large number of model parameters $\{\mathbf{a}_i\}$, and it is impossible to translate a constraint on N_{re} into a model-independent constraint on \mathbf{g} . This is a result of the fact that the currently allowed region in Fig. 1 is very close to the $N_{\text{re}} = 0$ line. One can nevertheless obtain constraints $\log_{10}(T_{\text{re}}/\text{GeV}) = 13.1 \pm 1.4$ and

$0.00515 < M/M_{\text{pl}} < 0.00526$ (the latter from (14)). In the RGI model, on the other hand, the posterior peaks in a region where condition (3) is fulfilled, so that future CMB data will permit measuring \mathbf{g} independently of the $\{\mathbf{a}_i\}$. For the fiducial parameters chosen here, the mean values and variances for the posteriors read $\log_{10} y = -6.5 \pm 2.2$, $\log_{10}(T_{\text{re}}/\text{GeV}) = 8.4 \pm 2.1$ and $M/M_{\text{pl}} = 0.00529 \pm 0.00007$. Finally, we estimate the improvement that can be made with data from the EUCLID satellite [88] and Square Kilometre Array (SKA) [89] by using $\sigma_{n_s} = 0.00085$ [90]. The resulting posteriors in Fig. 2 for the chosen values of \bar{n}_s and \bar{r} give $\log_{10}(T_{\text{re}}/\text{GeV}) = 13.5 \pm 1.1$ in the α -T model and $\log_{10}(T_{\text{re}}/\text{GeV}) = 8.3 \pm 1.4$ in the RGI model. The latter corresponds to $\log_{10} y = -6.6 \pm 1.4$.




Figure 2: Prior $P(x)$ and posteriors $P(x|\mathcal{D})$ for $x = \log_{10} \mathbf{g}$ with the different choices of \bar{n}_s , \bar{r} , σ_{n_s} , and σ_r in the RGI model (upper panel) and α -T model (lower panel). We assumed $\gamma = \tilde{\gamma} = 1$, but checked that the conclusions remain unchanged when using $\gamma = N'_k$ or $\tilde{\gamma} = (n_s'^2 + r'^2)^{1/2}$, with $'$ indicating a derivative with respect to x .

Conclusions: We introduced a simple analytic method to quantify the information gain on the inflaton coupling \mathbf{g} and the reheating temperature T_{re} from observational constraints on n_s and r . When applying it to future CMB observations with LiteBIRD and CMB-S4 we showed for

¹⁰ The parametric dependencies presented in the appendix imply that using the full information about the data made public at <http://bicepkeck.org/> is very unlikely to change our conclusions.

¹¹ Due to the different designs, sky coverages and foregrounds affecting the ground-based CMB-S4 program and the LiteBIRD satellite estimates of σ_r as a function of \bar{r} vary for both observatories. The values used here, which are based on Fig. 8 in [14] and is roughly consistent with Fig. 44 in [13], are sufficiently accurate for our proof of principle. A more detailed discussion can be found in [65].

the first time that these missions will be capable of performing the first ever measurement of T_{re} in both models considered here. For the chosen fiducial values this can directly be translated into a measurement of \mathbf{g} in the RGI model, while in the α -T model such a translation would require a specification of further parameters $\{\mathbf{a}_i\}$. Adding information from optical and 21cm surveys can further reduce the error bar on \mathbf{g} , and may help to constrain α and \mathbf{g} simultaneously from data by including observational knowledge on quantities not considered here, such as non-Gaussianities or the running of n_s . The inflaton coupling \mathbf{g} did not only crucially shape the evolution of the observable universe through its impact on T_{re} , but it is also a key parameter that connects models of inflation to theories of particle physics. Measuring this microphysical parameter, even with large error bars, will open up a new window to probe the connection between cosmology and fundamental physics. Hence, our findings add a qualitatively new dimension to the physics cases of future observatories.

Acknowledgements: We would like to thank Marcos Garcia, Jan Hamann, Jin U Kang, Eiichiro Komatsu, Isabel Oldengott, Christophe Ringeval, Evangelos Sfakianakis, and Yvonne Wong for helpful discussions. We also thank Björn Garbrecht, Andrea Giammanco, Jan Hamann and Fabio Maltoni for feedback on the revised version of this manuscript. L.M. acknowledges the State Scholarship Fund managed by the China Scholarship Council (CSC) and the Project funded by China Postdoctoral Science Foundation (2022M723677). MaD would like to thank the University of New South Wales for their hospitality under the Gordon Godfrey visitor program during part of the work on this project.

Appendix: Relation to observables: In this appendix we give the relations between the RHS of (2) and observables. A detailed derivation can be found in [47] and has been adapted to our notation in [50]. N_{re} can be obtained from

$$N_{\text{re}} = \frac{4}{3\bar{w}_{\text{re}} - 1} \left[N_k + \ln \left(\frac{k}{a_0 T_0} \right) + \frac{1}{4} \ln \left(\frac{40}{\pi^2 g_*} \right) + \frac{1}{3} \ln \left(\frac{11g_{s*}}{43} \right) - \frac{1}{2} \ln \left(\frac{\pi^2 M_{pl}^2 r A_s}{2\sqrt{\mathcal{V}_{\text{end}}}} \right) \right], \quad (11)$$

with $g_{s*} \approx g_*$, a_0 and $T_0 = 2.725$ K the scale factor and the temperature of the CMB at the present time, respectively, and

$$N_k = \ln \left(\frac{a_{\text{end}}}{a_k} \right) = \int_{\varphi_k}^{\varphi_{\text{end}}} \frac{H d\varphi}{\dot{\varphi}} \approx \frac{1}{M_{pl}^2} \int_{\varphi_{\text{end}}}^{\varphi_k} d\varphi \frac{\mathcal{V}}{\partial_{\varphi} \mathcal{V}}. \quad (12)$$

The subscript notation H_k, φ_k etc. indicates the value of the quantities H, φ etc. at the moment when a pivot-scale k crosses the horizon. φ_k can be expressed in terms of n_s and r by solving the relations

$$n_s = 1 - 6\epsilon_k + 2\eta_k, \quad r = 16\epsilon_k \quad (13)$$

with the slow roll parameters $\epsilon = (\partial_{\varphi} \mathcal{V} / \mathcal{V})^2 M_{pl}^2 / 2$ and $\eta = M_{pl}^2 \partial_{\varphi}^2 \mathcal{V} / \mathcal{V}$. In the slow roll regime, we find

$$H_k^2 = \frac{\mathcal{V}(\varphi_k)}{3M_{pl}^2} = \pi^2 M_{pl}^2 \frac{r A_s}{2}. \quad (14)$$

with $A_s = 10^{-10} e^{3.043}$ [1]. T_{re} can be expressed in terms of the observables (n_s, A_s, r) by plugging (11) with (12) into (1), φ_k is found by solving (13) for φ_k , and \mathcal{V}_{end} , and φ_{end} can be determined by solving $\epsilon = 1$ for φ . From (13) we obtain

$$\epsilon_k = \frac{r}{16}, \quad \eta_k = \frac{n_s - 1 + 3r/8}{2}, \quad (15)$$

from which we find

$$\left. \frac{\partial_{\varphi} \mathcal{V}(\varphi)}{\mathcal{V}(\varphi)} \right|_{\varphi_k} = \sqrt{\frac{r}{8M_{pl}^2}}, \quad \left. \frac{\partial_{\varphi}^2 \mathcal{V}(\varphi)}{\mathcal{V}(\varphi)} \right|_{\varphi_k} = \frac{n_s - 1 + 3r/8}{2M_{pl}^2} \quad (16)$$

by using the definitions of ϵ and η . Together with (14) this provides three equations that can be used to relate the effective potential and its derivatives to the observables (n_s, A_s, r) . That is sufficient to express \bar{w}_{re} and N_{re} in (11) in terms of observables, which is all that is needed to determine the RHS of (2).

-
- [1] N. Aghanim *et al.* (Planck), *Astron. Astrophys.* **641**, A6 (2020), [Erratum: *Astron. Astrophys.* 652, C4 (2021)], arXiv:1807.06209 [astro-ph.CO].
 - [2] P. A. R. Ade *et al.* (BICEP, Keck), *Phys. Rev. Lett.* **127**, 151301 (2021), arXiv:2110.00483 [astro-ph.CO].
 - [3] E. Abdalla *et al.*, *JHEAp* **34**, 49 (2022), arXiv:2203.06142 [astro-ph.CO].
 - [4] R. L. Workman (Particle Data Group), *PTEP* **2022**, 083C01 (2022).
 - [5] G. F. Giudice, *Phys. Rev. D* **58**, 105002 (1998), arXiv:0801.2562 [hep-ph].
 - [6] L. Canetti, M. Drewes, and M. Shaposhnikov, *New J. Phys.* **14**, 095012 (2012), arXiv:1204.4186 [hep-ph].
 - [7] A. A. Starobinsky, *Phys. Lett. B* **91**, 99 (1980).
 - [8] A. H. Guth, *Phys. Rev. D* **23**, 347 (1981).
 - [9] A. D. Linde, *Phys. Lett. B* **108**, 389 (1982).
 - [10] J. Martin, C. Ringeval, and V. Vennin, *Phys. Dark Univ.* **5-6**, 75 (2014), arXiv:1303.3787 [astro-ph.CO].
 - [11] L. Moncelli *et al.*, *Proc. SPIE Int. Soc. Opt. Eng.* **11453**, 1145314 (2020), arXiv:2012.04047 [astro-ph.IM].
 - [12] P. Ade *et al.* (Simons Observatory), *JCAP* **02**, 056 (2019), arXiv:1808.07445 [astro-ph.CO].
 - [13] E. Allys *et al.* (LiteBIRD), (2022), arXiv:2202.02773 [astro-ph.IM].
 - [14] K. Abazajian *et al.* (CMB-S4), *Astrophys. J.* **926**, 54 (2022), arXiv:2008.12619 [astro-ph.CO].
 - [15] A. Albrecht, P. J. Steinhardt, M. S. Turner, and F. Wilczek, *Phys. Rev. Lett.* **48**, 1437 (1982).
 - [16] A. D. Dolgov and D. P. Kirilova, *Sov. J. Nucl. Phys.* **51**, 172 (1990).
 - [17] J. H. Traschen and R. H. Brandenberger, *Phys. Rev. D* **42**, 2491 (1990).

- [18] Y. Shtanov, J. H. Traschen, and R. H. Brandenberger, *Phys. Rev. D* **51**, 5438 (1995), arXiv:hep-ph/9407247.
- [19] L. Kofman, A. D. Linde, and A. A. Starobinsky, *Phys. Rev. Lett.* **73**, 3195 (1994), arXiv:hep-th/9405187.
- [20] D. Boyanovsky, H. J. de Vega, R. Holman, and J. F. J. Salgado, *Phys. Rev. D* **54**, 7570 (1996), arXiv:hep-ph/9608205.
- [21] L. Kofman, A. D. Linde, and A. A. Starobinsky, *Phys. Rev. D* **56**, 3258 (1997), arXiv:hep-ph/9704452.
- [22] C. Caprini and D. G. Figueroa, *Class. Quant. Grav.* **35**, 163001 (2018), arXiv:1801.04268 [astro-ph.CO].
- [23] J. Ghiglieri and M. Laine, *JCAP* **07**, 022 (2015), arXiv:1504.02569 [hep-ph].
- [24] J. Ghiglieri, G. Jackson, M. Laine, and Y. Zhu, *JHEP* **07**, 092 (2020), arXiv:2004.11392 [hep-ph].
- [25] A. Ringwald, J. Schütte-Engel, and C. Tamarit, *JCAP* **03**, 054 (2021), arXiv:2011.04731 [hep-ph].
- [26] M. Drewes, Y. Georis, J. Klaric, and P. Klose, *JCAP* **06**, 073 (2024), arXiv:2312.13855 [hep-ph].
- [27] J. Martin and C. Ringeval, *Phys. Rev. D* **82**, 023511 (2010), arXiv:1004.5525 [astro-ph.CO].
- [28] P. Adshead, R. Easther, J. Pritchard, and A. Loeb, *JCAP* **02**, 021 (2011), arXiv:1007.3748 [astro-ph.CO].
- [29] R. Easther and H. V. Peiris, *Phys. Rev. D* **85**, 103533 (2012), arXiv:1112.0326 [astro-ph.CO].
- [30] K. D. Lozanov and M. A. Amin, *Phys. Rev. Lett.* **119**, 061301 (2017), arXiv:1608.01213 [astro-ph.CO].
- [31] K. D. Lozanov and M. A. Amin, *Phys. Rev. D* **97**, 023533 (2018), arXiv:1710.06851 [astro-ph.CO].
- [32] J. B. Munoz and M. Kamionkowski, *Phys. Rev. D* **91**, 043521 (2015), arXiv:1412.0656 [astro-ph.CO].
- [33] J. L. Cook, E. Dimastrogiovanni, D. A. Easson, and L. M. Krauss, *JCAP* **04**, 047 (2015), arXiv:1502.04673 [astro-ph.CO].
- [34] N. Zhang, Y.-B. Wu, J.-W. Lu, C.-W. Sun, L.-J. Shou, and H.-Z. Xu, *Chin. Phys. C* **44**, 095107 (2020), arXiv:1807.03596 [astro-ph.CO].
- [35] N. K. Stein and W. H. Kinney, *JCAP* **01**, 022 (2022), arXiv:2106.02089 [astro-ph.CO].
- [36] R.-G. Cai, Z.-K. Guo, and S.-J. Wang, *Phys. Rev. D* **92**, 063506 (2015), arXiv:1501.07743 [gr-qc].
- [37] A. Di Marco, G. Pradisi, and P. Cabella, *Phys. Rev. D* **98**, 123511 (2018), arXiv:1807.05916 [astro-ph.CO].
- [38] D. Maity and P. Saha, *JCAP* **07**, 018 (2019), arXiv:1811.11173 [astro-ph.CO].
- [39] D. Maity and P. Saha, *Class. Quant. Grav.* **36**, 045010 (2019), arXiv:1902.01895 [gr-qc].
- [40] S. Antusch, D. G. Figueroa, K. Marschall, and F. Torrenti, *Phys. Lett. B* **811**, 135888 (2020), arXiv:2005.07563 [astro-ph.CO].
- [41] L. Dai, M. Kamionkowski, and J. Wang, *Phys. Rev. Lett.* **113**, 041302 (2014), arXiv:1404.6704 [astro-ph.CO].
- [42] V. Domcke and J. Heisig, *Phys. Rev. D* **92**, 103515 (2015), arXiv:1504.00345 [astro-ph.CO].
- [43] I. Dalianis, G. Koutsoumbas, K. Ntrekis, and E. Papantonopoulos, *JCAP* **02**, 027 (2017), arXiv:1608.04543 [gr-qc].
- [44] J.-O. Gong, S. Pi, and G. Leung, *JCAP* **05**, 027 (2015), arXiv:1501.03604 [hep-ph].
- [45] F. Takahashi and W. Yin, *JHEP* **07**, 095 (2019), arXiv:1903.00462 [hep-ph].
- [46] R. J. Hardwick, V. Vennin, K. Koyama, and D. Wands, *JCAP* **08**, 042 (2016), arXiv:1606.01223 [astro-ph.CO].
- [47] Y. Ueno and K. Yamamoto, *Phys. Rev. D* **93**, 083524 (2016), arXiv:1602.07427 [astro-ph.CO].
- [48] K. Nozari and N. Rashidi, *Phys. Rev. D* **95**, 123518 (2017), arXiv:1705.02617 [astro-ph.CO].
- [49] A. Di Marco, P. Cabella, and N. Vittorio, *Phys. Rev. D* **95**, 103502 (2017), arXiv:1705.04622 [astro-ph.CO].
- [50] M. Drewes, J. U. Kang, and U. R. Mun, *JHEP* **11**, 072 (2017), arXiv:1708.01197 [astro-ph.CO].
- [51] D. Maity and P. Saha, *Phys. Rev. D* **98**, 103525 (2018), arXiv:1801.03059 [hep-ph].
- [52] N. Rashidi and K. Nozari, *Int. J. Mod. Phys. D* **27**, 1850076 (2018), arXiv:1802.09185 [astro-ph.CO].
- [53] G. German, *Int. J. Mod. Phys. D* **31**, 2250081 (2022), arXiv:2010.09795 [astro-ph.CO].
- [54] S. S. Mishra, V. Sahni, and A. A. Starobinsky, *JCAP* **05**, 075 (2021), arXiv:2101.00271 [gr-qc].
- [55] J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, K. A. Olive, and S. Verner, *Phys. Rev. D* **105**, 043504 (2022), arXiv:2112.04466 [hep-ph].
- [56] A. Nautiyal, *Phys. Rev. D* **98**, 103531 (2018), arXiv:1806.03081 [astro-ph.CO].
- [57] S.-M. Choi and H. M. Lee, *Eur. Phys. J. C* **76**, 303 (2016), arXiv:1601.05979 [hep-ph].
- [58] P. Cabella, A. Di Marco, and G. Pradisi, *Phys. Rev. D* **95**, 123528 (2017), arXiv:1704.03209 [astro-ph.CO].
- [59] R. Kabir, A. Mukherjee, and D. Lohiya, *Mod. Phys. Lett. A* **34**, 1950114 (2019), arXiv:1609.09243 [gr-qc].
- [60] S. Bhattacharya, K. Dutta, and A. Maharana, *Phys. Rev. D* **96**, 083522 (2017), [Addendum: *Phys. Rev. D* **96**, 109901 (2017)], arXiv:1707.07924 [hep-ph].
- [61] I. Dalianis and Y. Watanabe, *JHEP* **02**, 118 (2018), arXiv:1801.05736 [hep-ph].
- [62] M. Drewes, *JCAP* **03**, 013 (2016), arXiv:1511.03280 [astro-ph.CO].
- [63] J. Martin, C. Ringeval, and V. Vennin, *Phys. Rev. D* **94**, 123521 (2016), arXiv:1609.04739 [astro-ph.CO].
- [64] M. Drewes, *JCAP* **09**, 069 (2022), arXiv:1903.09599 [astro-ph.CO].
- [65] M. Drewes, L. Ming, and I. Oldengott, *JCAP* **05**, 081 (2024), arXiv:2303.13503 [hep-ph].
- [66] S. Renaux-Petel and K. Turzyński, *Phys. Rev. Lett.* **117**, 141301 (2016), arXiv:1510.01281 [astro-ph.CO].
- [67] S. Passaglia, W. Hu, A. J. Long, and D. Zegeye, *Phys. Rev. D* **104**, 083540 (2021), arXiv:2108.00962 [hep-ph].
- [68] J. E. Lidsey, A. R. Liddle, E. W. Kolb, E. J. Copeland, T. Barreiro, and M. Abney, *Rev. Mod. Phys.* **69**, 373 (1997), arXiv:astro-ph/9508078.
- [69] M. A. Amin, M. P. Hertzberg, D. I. Kaiser, and J. Karouby, *Int. J. Mod. Phys. D* **24**, 1530003 (2014), arXiv:1410.3808 [hep-ph].
- [70] E. W. Kolb, A. Notari, and A. Riotto, *Phys. Rev. D* **68**, 123505 (2003), arXiv:hep-ph/0307241.
- [71] K. Mukaida and K. Nakayama, *JCAP* **01**, 017 (2013), arXiv:1208.3399 [hep-ph].
- [72] K. Mukaida and K. Nakayama, *JCAP* **03**, 002 (2013), arXiv:1212.4985 [hep-ph].
- [73] M. Drewes and J. U. Kang, *Nucl. Phys. B* **875**, 315 (2013), [Erratum: *Nucl. Phys. B* **888**, 284–286 (2014)], arXiv:1305.0267 [hep-ph].
- [74] K. Mukaida, K. Nakayama, and M. Takimoto, *JHEP* **12**, 053 (2013), arXiv:1308.4394 [hep-ph].
- [75] M. Drewes, *JCAP* **11**, 020 (2014), arXiv:1406.6243 [hep-ph].

- [76] R. T. Co, E. Gonzalez, and K. Harigaya, JCAP **11**, 038 (2020), arXiv:2007.04328 [astro-ph.CO].
- [77] M. A. G. Garcia, K. Kaneta, Y. Mambrini, and K. A. Olive, JCAP **04**, 012 (2021), arXiv:2012.10756 [hep-ph].
- [78] M. Fairbairn, L. Lopez Honorez, and M. H. G. Tytgat, Phys. Rev. D **67**, 101302 (2003), arXiv:hep-ph/0302160.
- [79] J. Martin, C. Ringeval, R. Trotta, and V. Vennin, JCAP **03**, 039 (2014), arXiv:1312.3529 [astro-ph.CO].
- [80] R. Kallosh and A. Linde, JCAP **10**, 033 (2013), arXiv:1307.7938 [hep-th].
- [81] R. Kallosh and A. Linde, JCAP **07**, 002 (2013), arXiv:1306.5220 [hep-th].
- [82] J. J. M. Carrasco, R. Kallosh, and A. Linde, JHEP **10**, 147 (2015), arXiv:1506.01708 [hep-th].
- [83] J. J. M. Carrasco, R. Kallosh, and A. Linde, Phys. Rev. D **92**, 063519 (2015), arXiv:1506.00936 [hep-th].
- [84] D. Boyanovsky, K. Davey, and C. M. Ho, Phys. Rev. D **71**, 023523 (2005), arXiv:hep-ph/0411042.
- [85] P. Carenza, A. Mirizzi, and G. Sigl, Phys. Rev. D **101**, 103016 (2020), arXiv:1911.07838 [hep-ph].
- [86] J. Martin, C. Ringeval, and V. Vennin, Phys. Rev. Lett. **114**, 081303 (2015), arXiv:1410.7958 [astro-ph.CO].
- [87] M. Kamionkowski and E. D. Kovetz, Ann. Rev. Astron. Astrophys. **54**, 227 (2016), arXiv:1510.06042 [astro-ph.CO].
- [88] R. Laureijs *et al.* (EUCLID), (2011), arXiv:1110.3193 [astro-ph.CO].
- [89] R. Maartens, F. B. Abdalla, M. Jarvis, and M. G. Santos (SKA Cosmology SWG), PoS **AASKA14**, 016 (2015), arXiv:1501.04076 [astro-ph.CO].
- [90] T. Sprenger, M. Archidiacono, T. Brinckmann, S. Clesse, and J. Lesgourgues, JCAP **02**, 047 (2019), arXiv:1801.08331 [astro-ph.CO].