

# GEOMETRIC DIRAC OPERATOR ON NONCOMMUTATIVE TORUS AND $M_2(\mathbb{C})$

E. LIRA-TORRES AND S. MAJID

**ABSTRACT.** We construct quantum geometrically realised spectral triples or ‘Dirac operators’ on the noncommutative torus  $\mathbb{C}_\theta[T^2]$  with its standard quantum metric and on the algebra  $M_2(\mathbb{C})$  of  $2 \times 2$  matrices with two standard quantum metrics and associated quantum Levi-Civita connections. We derive the required form of Clifford algebra and 2-spinor bundle connection compatible with the quantum Riemannian geometry. In the case of the noncommutative torus, we obtain a standard even spectral triple but now uniquely determined by geometric realisability. In the case of  $M_2(\mathbb{C})$ , one metric with its flat quantum Levi-Civita connection results in a natural even spectral triple. The other metric has a Lorentzian signature and we show that it admits a natural 2-parameter almost spectral triple where  $\not{D}$  is neither hermitian nor antihermitian in general and the quantum Levi-Civita connection is curved. In all cases, we split the construction into a ‘local’ tensorial part related to the quantum geometry where we classify the results more broadly, and the further requirements relating to the Hilbert space structure.

## 1. INTRODUCTION

Noncommutative geometry, the idea that coordinates of a space could be noncommutative as in quantum mechanics, has gained interest in recent years as a plausibly better description of spacetime and phase space that accounts for quantum gravity effects[28, 22, 23, 15, 18], as well as clearly having potential for diverse applications in quantum theory. Probably the best known approach is the one coming out of cyclic cohomology and operator algebras[10, 17] with the noncommutative torus as a prime example[8] and leading up to the notion of ‘Connes spectral triple’ in the role of Dirac operator[9, 11]. Meanwhile, since the arrival of true quantum groups in the 1980s, there has emerged a constructive ‘quantum groups’ approach which starts with a unital but possibly noncommutative ‘coordinate algebra’  $A$  with quantum or classical symmetries to narrow down the choice of a bimodule of 1-forms  $\Omega^1$ . In recent years, this was extended to a systematic theory of ‘quantum Riemannian geometry’ (QRG)[2] with a quantum metric  $g \in \Omega^1 \otimes_A \Omega^1$ , a quantum Levi-Civita connection (QLC)  $\nabla : \Omega^1 \rightarrow \Omega^1 \otimes_A \Omega^1$ , etc. This framework of curved quantum spacetimes has been applied to baby models of quantum gravity and quantum geodesics, and is also beginning to be applied to particle physics, see [1] for a review.

Of interest in this paper and linking the two approaches to noncommutative geometry is a framework [5][2, Chapter 8.5] to extend a QRG with (a) a ‘spinor’ bimodule

---

2000 *Mathematics Subject Classification.* Primary 81R50, 58B32, 83C57.

*Key words and phrases.* noncommutative geometry, quantum groups, quantum gravity.

$\mathcal{S}$  equipped with a bimodule connection  $\nabla_S$ , (b) a ‘Clifford action’ bimodule map  $\triangleright : \Omega^1 \otimes_A \mathcal{S} \rightarrow \mathcal{S}$  and (c) a ‘charge conjugation’ bimodule map  $\mathcal{J} : \mathcal{S} \rightarrow \bar{\mathcal{S}}$  to the conjugate bimodule (so  $\mathcal{J}$  is antilinear), so as to arrive at a quantum geometric Dirac operator  $\not{D} = \triangleright \circ \nabla_S : \mathcal{S} \rightarrow \mathcal{S}$  obeying the ‘local’ part of Connes axioms for a spectral triple (meaning before consideration of a Hilbert space structure on  $\mathcal{S}$ ). If the latter step can also be completed then we obtain a *quantum geometrically realised* spectral triple. Not every Connes spectral triple can be geometrically realised like this but, rather, geometric realisability, picks out natural spectral triples where we have this additional quantum geometry in an explicit form. Moreover, [5] proposed two further (optional) restrictions on the Clifford action  $\triangleright$  which are not needed for a spectral triple but link the spinor structures to the metric as would be the case classically. These are certain ‘Clifford relations’, see (2.2), and that  $\triangleright$  should be covariantly constant, i.e. covariant with respect to the connections on the two sides of the map, see (2.1).

In terms of examples, [5] constructed a natural q-deformed Dirac operator on the standard q-sphere  $C_q[S^2]$  obeying the local conditions but with  $\mathcal{J}$  being not quite an antilinear isometry (but a q-deformed one), and also gave an example of a spectral triple on  $M_2(\mathbb{C})$  but with Clifford action  $\triangleright$  not being covariantly constant, so not a full geometric realisation. [20] treated the fuzzy sphere and found a natural full quantum-geometrically realised spectral triple as member of a previous family proposed in [13]. In the present sequel, Section 3 provides a similar analysis for the algebraic noncommutative torus  $\mathbb{C}_\theta[\mathbb{T}^2]$ . Because the noncommutative torus has been so extensively studied, we do not expect to obtain, and don’t, a new spectral triple. Rather, we obtain the ‘obvious’ construction

$$(\not{D}\psi)_\beta = \partial_i \psi_\alpha \sigma^{i\alpha}_\beta$$

where  $\sigma^i$  are Pauli matrices for  $i = 1, 2$ ,  $\partial_i$  are certain natural derivations[8] and  $\psi_\alpha$  is a 2-spinor with values in the noncommutative torus. Up to conventions, this is also the Euclidean metric case in the exposition [7]. Our result is that this  $\not{D}$  is unique up to unitary equivalence among quantum-geometrically realised spectral triples with  $\mathcal{S}$  free and 2-dimensional over the algebra. We let  $\nabla$  be any weak QLC but the construction forces this to be the unique QLC on this algebra (given simply by  $\nabla s^i = 0$  on a canonical left-invariant self-adjoint basis  $\{s^i\}$ ). Including a possible even structure  $\gamma$  we actually have two spectral triples which one could view as  $n = 1, 2$  respectively in the period 8 classification. It is known that there are four (equivariant) spectral triples on the noncommutative torus[26] and presumably we obtain two of them. We then show in Section 3.3 that we get a much bigger moduli of  $\nabla_S, \nabla$  if we drop that  $\triangleright$  is covariantly constant. Remarkably, the final  $\not{D}$  itself remains of the above form but with the possible addition of a constant mass term or chiral ‘mass’ term. Throughout, to be concrete, we fix the Euclidean quantum metric  $g = s^1 \otimes s^1 + s^2 \otimes s^2$  in our basis.

Sections 4 and Section 5 then provide a similar analysis for  $M_2(\mathbb{C})$ , using the two standard quantum geometries in [2, Example 8.13] and [2, Exercise 8.5] respectively. In both cases, we convert to a self-adjoint basis  $\{s^i\}$  for  $\Omega^1$  and, in these terms, the first quantum metric is

$$g = \iota(s^2 \otimes s^1 - s^1 \otimes s^2)$$

and we work with a natural 1-parameter sub-moduli of the known 4-parameter QLCs for this. We then find, Theorem 4.3, a generic construction for a large

class of spectral triples at the ‘local’ level of  $\nabla_S, \triangleright, \mathcal{J}$  provided we are at the unique flat QLC in this class. With some simplifying assumptions we are led to a solution (Proposition 4.8) for which  $\not{D}$  is furthermore compatible with the usual inner product given by  $\text{Tr} : M_2(\mathbb{C}) \rightarrow \mathbb{C}$  on the spinor components to give an  $n = 0$  or  $n = 1$  quantum geometrically realised spectral triple (depending on a sign  $\epsilon' = \pm 1$ ). Moreover, the ‘Clifford relations’ hold in a slightly generalised form with an automorphism [5] and  $\triangleright$  is covariantly constant and therefore more canonical than previously. After a unitary change of basis in Corollary 4.9, we obtain

$$(\not{D}\psi)_\beta = \frac{i}{2\sqrt{2}} \left( (\sigma^1\psi_\alpha + \psi_\alpha\sigma^2)\sigma^{1\alpha}_\beta + \epsilon'(\sigma^2\psi_\alpha + \psi_\alpha\sigma^1)\sigma^{3\alpha}_\beta \right)$$

as natural geometrically realised spectral triples for this quantum metric.

By contrast, the second quantum metric in our self-adjoint basis has the diagonal but ‘Lorentzian’ form

$$g = -s^1 \otimes s^1 + s^2 \otimes s^2,$$

where the - sign in the metric is needed for quantum symmetry. We find that this does not admit solutions for the spinor  $\nabla_S, \triangleright, \mathcal{J}$  if we impose the full Clifford relations but does admit solutions if we impose only half of these relations. These lead to a natural real 2-parameter family of geometrically realised almost spectral triples on  $M_2(\mathbb{C})$ , see Theorem 5.4 and Proposition 5.5. The two real parameters  $s, t$  here fall on a circle  $s^2 + t^2 = 2$  and the QLC  $\nabla$  is forced to have a specific value with curvature. These are ‘almost’ spectral triples in that  $\not{D}$  turns out in the construction to be neither antihermitian nor hermitian but to be built in a simple way from two hermitian operators  $\not{D}^{(i)}$ ,  $i = 1, 2$ . We attribute this to the Lorentzian nature of the quantum metric.

The  $M_2(\mathbb{C})$  case needed significant development of the previous analysis[20], done in Section 2.2, so as to cover central bases of  $\Omega^1$  that need not obey the Grassmann algebra as differential forms and Christoffel symbols of  $\nabla, \nabla_S$  that need not be  $\mathbb{C}$ -valued. This and the required simplifications in the inner case in Section 2.3 are some general results of the paper and reduce everything to solving nonlinear matrix equations, which we then solved using Mathematica. Section 2.1 provides a very brief introduction to quantum Riemannian geometry as in [2] and to the geometric realisation programme [5]. The paper concludes in Section 6 with directions for further work.

## 2. PRELIMINARIES AND FORMALISM FOR CENTRAL BASES

Here, we very briefly outline the formalism of [5][2, Chap 8.5] for the construction of Connes spectral triples from within ‘quantum Riemannian geometry’ (QRG). Then, in Section 2.2, we extend the case of a central basis  $s^i$  of  $\Omega^1$  in [20] to now allow the  $s^i$  to not be Grassmann and the Christoffel symbols of the quantum Levi-Civita connection to not be constants. In Section 2.3, we look at the special features when  $\Omega^1$  is inner. We will need these more general results for the matrix algebra  $M_2(\mathbb{C})$ .

**2.1. Elements of QRG.** We work with  $A$  a unital algebra, typically a  $*$ -algebra over  $\mathbb{C}$ , in the role of ‘coordinate algebra’. Differentials are formally introduced as a bimodule  $\Omega^1$  of 1-forms equipped with a map  $d : A \rightarrow \Omega^1$  obeying the Leibniz rule  $d(ab) = (da)b + adb$ . We assume this extends to an exterior algebra  $(\Omega, d)$  with

$d^2 = 0$  and  $d$  obeying the graded-Leibniz rule, and where  $A, dA$  generate  $\Omega$  (this is more restrictive than a differential graded algebra). The exterior algebra is called *inner* if there exists  $\theta \in \Omega^1$  such that  $d = [\theta, \ ]$  is a graded derivation. We also use the term for  $\Omega^1$  to be inner with  $d = [\theta, \ ]$ .

A quantum metric is  $g \in \Omega^1 \otimes_A \Omega^1$  and a bimodule map inverse  $(\ , \ ) : \Omega^1 \otimes_A \Omega^1 \rightarrow A$  in an obvious sense, together with some form of quantum symmetry condition. The general functorial definition of the latter is  $\wedge(g) = 0$ , but in some contexts it seems better to modify this. Following [2], we say  $g$  is a generalised quantum metric if no form of symmetry is imposed. A (left) bimodule connection [16, 25] on  $\Omega^1$  is  $\nabla : \Omega^1 \rightarrow \Omega^1 \otimes_A \Omega^1$  obeying

$$\nabla(a.\omega) = a.\nabla\omega + da \otimes \omega, \quad \nabla(\omega.a) = (\nabla\omega).a + \sigma(\omega \otimes da)$$

for all  $a \in A, \omega \in \Omega^1$ , for some ‘generalised braiding’ bimodule map  $\sigma : \Omega^1 \otimes_A \Omega^1 \rightarrow \Omega^1 \otimes_A \Omega^1$ . The latter, if it exists, is uniquely determined and not additional data. But for an inner calculus, connections are given by a free choice of module map  $\sigma$  and an additional bimodule map  $\alpha : \Omega^1 \rightarrow \Omega^1 \otimes_A \Omega^1$ , see [2, Prop. 8.11]. A connection is torsion free if  $T_\nabla := \wedge\nabla - d$  vanishes, and metric compatible if

$$\nabla g = (\nabla \otimes \text{id} + (\sigma \otimes \text{id})(\text{id} \otimes \nabla))g$$

vanishes. When both vanish, we have a *quantum Levi-Civita connection* (QLC). A weaker concept is a WQLC where we demand zero torsion and zero *cotorsion* in the sense

$$(d \otimes \text{id} - \text{id} \wedge \nabla)g = 0$$

and we say in this case that we have a weak quantum Riemannian geometry (WQRG). The curvature of  $\nabla$  is (similarly) defined as

$$R_\nabla = (d \otimes \text{id} - \text{id} \wedge \nabla)\nabla : \Omega^1 \rightarrow \Omega^2 \otimes_A \Omega^1.$$

Finally, working over  $\mathbb{C}$ , we need  $(\Omega, d)$  to be a  $*$ -calculus,  $g$  ‘real’ in the sense  $g^\dagger = g$  where  $\dagger = \text{flip}(* \otimes *)$  is well defined on  $\Omega^1 \otimes_A \Omega^1$ , and  $\nabla$   $*$ -preserving in the sense  $\nabla \circ * = \sigma \circ \dagger \circ \nabla$ , see [2].

The formulation of a bimodule connection above applies equally well to any  $A$ -bimodule  $\mathcal{S}$  in the role of sections of a ‘vector bundle’, now with  $\nabla : \mathcal{S} \rightarrow \Omega^1 \otimes_A \mathcal{S}$  and  $\sigma_\mathcal{S} : \mathcal{S} \otimes_A \Omega^1 \rightarrow \Omega^1 \otimes_A \mathcal{S}$ , and the above formulae now with  $\omega \in \Omega^1$  replaced by  $\phi \in \mathcal{S}$ . We also need a *Clifford bimodule map*  $\triangleright : \Omega^1 \otimes_A \mathcal{S} \rightarrow \mathcal{S}$  which generalises the role of the gamma-matrices when constructing Dirac operators [5]. In this case, we define  $\not{D} = \triangleright \circ \nabla_\mathcal{S} : \mathcal{S} \rightarrow \mathcal{S}$  as a minimal ‘geometric Dirac operator’ in the formalism. We ideally would like  $\nabla(\triangleright) = 0$  i.e.,

$$(2.1) \quad (\text{id} \otimes \triangleright) \circ (\nabla \otimes \text{id} + (\sigma \otimes \text{id})(\text{id} \otimes \nabla_\mathcal{S})) = \nabla \circ \triangleright,$$

which says that  $\triangleright$  is covariantly constant in the sense that it intertwines the tensor product connection on  $\Omega^1 \otimes_A \mathcal{S}$  and the connection on  $\mathcal{S}$ . [5] also tentatively proposed a ‘Clifford relation’ that  $\triangleright$  extends to  $\Omega^2 \otimes_A \mathcal{S} \rightarrow \mathcal{S}$  in such a way that

$$(2.2) \quad (\omega \wedge \eta) \triangleright \phi := \omega \triangleright (\eta \triangleright \phi) + (\omega, \eta) \phi$$

for all  $\omega, \eta \in \Omega^1$  and  $\phi \in \mathcal{S}$ , or a generalisation of this to allow an automorphism and a scaling of the inner product. The motivation here is that this reduces to the Clifford algebra relations in the classical case. Both of these geometric conditions are part of the quantum Riemannian geometry and *not* part of the requirements for  $\not{D}$  to be part of a Connes spectral triple.

For a Connes spectral triple, we instead need that  $\mathcal{S}$  has an antilinear skew-bimodule map  $\mathcal{J} : \mathcal{S} \rightarrow \mathcal{S}$  in the role of ‘charge conjugation’ and, if it exists, a bimodule map  $\gamma : \mathcal{S} \rightarrow \mathcal{S}$ , both with certain properties needed to fit the algebraic part of Connes axioms[10] for free sign parameters  $\epsilon, \epsilon', \epsilon''$ . Finally, we need a compatible sesquilinear inner product  $\langle \cdot, \cdot \rangle$  on  $\mathcal{S}$  with respect to which we complete the latter into a Hilbert space as explained in [5][2, Chap. 8.5], where the inner product is denoted  $\langle\langle \cdot, \cdot \rangle\rangle$ , such that  $\not{D}$  is antihermitian,  $\mathcal{J}$  an (antilinear) isometry and (if it exists)  $\gamma$  hermitian. Here,  $\not{D}$  is the hermitian Dirac operator in Connes axioms. Note that [5][2, Chap. 8.5] uses a more careful notation in which antilinearity is expressed as linearity by the use of the conjugate bimodule  $\bar{\mathcal{S}}$  (defined as the same set as  $\mathcal{S}$  but conjugated actions of the field and of  $A$ ) and  $j(\phi) = \overline{\mathcal{J}(\phi)}$ , etc. The overline here means to view the element of  $\mathcal{S}$  in  $\bar{\mathcal{S}}$ . One should similarly distinguish between a sesquilinear map and a linear map on  $\bar{\mathcal{S}} \otimes \mathcal{S}$ . These distinctions are, however, only for book-keeping and we suppress them here.

**2.2. QRG realisation of Connes spectral triples for a central basis.** We will be interested in the case where  $A$  has trivial centre,  $\Omega^1$  has a central basis  $\{s^i\}$  and  $\mathcal{S}$  has a central basis  $\{e^\alpha\}$ . Some general analysis for quantum Riemannian geometry on  $\Omega^1$  in this parallelizable case is in [2, Ex. 8.2] and was extended to construct QRG-realised spectral triples in [20] but under more restrictive conditions than needed now. Over  $\mathbb{C}$ , we assume that  $s^{i*} = s^i$  are self-adjoint. Then ‘reality’ of the metric translates to the matrix  $g_{ij}$  of metric coefficient in the basis being hermitian. If  $\sigma$  is the flip map then the  $*$ -preserving condition on  $\nabla$  translates to the Christoffel symbols in the basis being real, otherwise the condition is more complicated.

The further data for a Dirac operator starts with a ‘Clifford action’  $\triangleright : \Omega^1 \otimes_A \mathcal{S} \rightarrow \mathcal{S}$  of the form

$$(2.3) \quad s^i \triangleright e^\alpha = C^{i\alpha}{}_\beta e^\beta, \quad C^{i\alpha}{}_\beta \in \mathbb{C}$$

due to centrality of the bases and being a bimodule map. We will also consider this as a collection of matrices  $(C^i)^\alpha{}_\beta = C^{i\alpha}{}_\beta$ . Next, we define the ‘charge conjugation’ operator according to

$$(2.4) \quad \mathcal{J}(ae^\alpha) = a^* J^\alpha{}_\beta e^\beta$$

$$J^\alpha{}_\beta * J^\beta{}_\gamma = \epsilon \delta^\alpha{}_\gamma; \quad \overline{J}J = \text{id}, \quad \epsilon = \pm 1, \quad J^\alpha{}_\beta \in \mathbb{C},$$

where we also write the equation in a compact form for a matrix  $J$ . The overline denotes complex conjugation of the entries.

We similarly need a bimodule connection which we write in the form

$$\nabla_S e^\alpha = S^\alpha{}_{i\beta} s^i \otimes e^\beta, \quad \sigma_S(e^\alpha \otimes s^j) = \sigma_S^{\alpha j}{}_{i\beta} s^i \otimes e^\beta, \quad \sigma_S^{\alpha j}{}_{i\beta} \in \mathbb{C},$$

where now  $S^\alpha{}_{i\beta}$  will have entries in  $A$ , not constants as in [20] and  $\sigma_S$  need not be the flip map (this is the same story as  $\Gamma, \sigma$ ). The abstract formulation of the ‘reality condition’ in [5] then becomes

$$(2.5) \quad S^\alpha{}_{i\beta} * J^\beta{}_\gamma \sigma_S^{\gamma i}{}_{j\delta} = J^\alpha{}_\beta S^\beta{}_{j\delta}; \quad \overline{S_j} J \sigma_{S^j}^j = J S_i.$$

Compatibility of  $\mathcal{J}$  with the Clifford action in the general scheme of [5] now becomes

$$(2.6) \quad C^{i\alpha}{}_\beta * J^\beta{}_\delta = \epsilon' J^\alpha{}_\beta \sigma_S^{\beta i}{}_{j\gamma} C^{j\gamma}{}_\delta; \quad \epsilon' = \pm 1, \quad \overline{C^i} J = \epsilon' J \sigma_{S^j}^i C^j.$$

Finally, for an ‘even’ spectral triple, we also set  $\gamma(e^\alpha) = \gamma^\alpha_\beta e^\beta$ , here again with  $\gamma^\alpha_\beta \in \mathbb{C}$  under our central basis and centre assumptions. Then the properties needed in [5] reduce to

$$\begin{aligned}
(2.7) \quad & \gamma^\alpha_\beta \gamma^\beta_\nu = \delta^\alpha_\nu; & \gamma^2 &= \text{id}, \\
(2.8) \quad & C^{i\alpha}_\beta \gamma^\beta_\nu = -\gamma^\alpha_\beta C^{i\beta}_\nu; & \{C^i, \gamma\} &= 0, \\
(2.9) \quad & \gamma^\alpha_\beta {}^* J^\beta_\nu = \epsilon'' J^\alpha_\beta \gamma^\beta_\nu; & \bar{\gamma} J &= \epsilon'' J \gamma, \quad \epsilon'' = \pm 1, \\
(2.10) \quad & S^\alpha_{i\beta} \gamma^\beta_\nu = \gamma^\alpha_\beta S^\beta_{i\nu}; & [S_i, \gamma] &= 0.
\end{aligned}$$

Here (2.8) assumes the pattern of matches an even  $n$ . We note that the full set of conditions for  $J$  is invariant under multiplication by a phase. Beyond the above local data for a spectral triple, we have two further, quantum-geometric, requirements not present in Connes’ formalism. The first is the covariance  $\nabla(\triangleright) = 0$  condition:

$$(2.11) \quad C^{i\alpha}_\beta S^\beta_{j\gamma} - \sigma^{ik}_{jl} S^\alpha_{k\beta} C^{l\beta}_\gamma = -\frac{1}{2} \Gamma^i_{jk} C^{k\alpha}_\gamma; \quad C^i S_j - \sigma^{ik}_{jl} S_k C^l = -\frac{1}{2} \Gamma^i_{jk} C^k,$$

where  $\nabla s^i = -\frac{1}{2} \Gamma^i_{jk} s^j \otimes s^k$  and  $\sigma(s^i \otimes s^j) = \sigma^{ij}_{kl} s^k \otimes s^l$ . Unlike [20], we allow  $\Gamma^i_{jk} \in A$  and  $\sigma$  need not be the flip. The second is a tentative proposal for compatibility of the Clifford action with  $\Omega^2$ , namely (in the simplest case),

$$(2.12) \quad (s^i \wedge s^j) \triangleright e^\alpha := s^i \triangleright (s^j \triangleright e^\alpha) - g^{ij} e^\alpha$$

is required to be well-defined. Here  $g = g_{ij} s^i \otimes s^j$  is the quantum metric with coefficients forming a hermitian matrix and  $g^{ij}$  is its inverse. But again unlike [20], we do not assume that the  $s^i$  enjoy the usual Grassmann algebra, so this does not necessarily impose that the  $C^i$  obey something resembling the usual Clifford algebra relations.

This completes our analysis of the ‘local’ side of the geometric realisation problem[5] for a slightly larger class of quantum geometries with central bases than in [20]. Note that the equation (2.4) is a self-contained problem and can already be analysed in each dimension. When  $\mathcal{S}$  is 2-dimensional over the algebra, which will be the case of interest, it is easy enough to describe the possible  $J$  explicitly up to a phase (which is not determined by the equations).

**Lemma 2.1.** *Up to a phase in the 2D spinor bundle case,  $J$  can be obtained with  $r > 0$  and  $z \in \mathbb{C}$  as either:*

$$(1): \quad J = \begin{pmatrix} z & r \\ \frac{\epsilon - |z|^2}{r} & -\bar{z} \end{pmatrix}, \quad (2): \quad J = \begin{pmatrix} 1 & \frac{\epsilon - 1}{|z|^2} z \\ z & -\frac{z}{\bar{z}} \end{pmatrix}$$

or its transpose. The  $\epsilon = -1$  case of (2) needs  $z \neq 0$  and up to a phase is also an instance to type (1).

*Proof.* Up to a phase we can assume the top right corner is real or zero, so  $J = \begin{pmatrix} a & r \\ c & d \end{pmatrix}$

or  $J = \begin{pmatrix} a & 0 \\ c & d \end{pmatrix}$  for complex parameters and a real  $r > 0$ . Writing out (2.4) we have in the first case looking off-diagonal in  $\bar{J}J$  that  $d = -\bar{a}$  and  $(\bar{c} - c)a = 0$ . If  $a \neq 0$ , we need  $c$  real and  $c = (\epsilon - |a|^2)/r$  which gives the form shown. If  $a = 0$  then we are led to  $c = \epsilon/r$ . For the other case, from the diagonal of  $\bar{J}J$  we have  $|a|^2 = |d|^2 = \epsilon$

hence  $\epsilon = 1$  and, up to a phase, we can assume  $a = 1$  also. In this case off diagonal we need that  $d = -c/\bar{c}$  as shown. We note, however, that those cases of type (1) for general  $r$  and  $\epsilon = -1$  that have an element of modulus 1 in the top left are equal up to a phase to the case (2) with  $\epsilon = -1$ , so this is redundant if we want an exclusive classification. Also for  $\epsilon = 1$  in type (2), we can send  $z \rightarrow 0$  along a ray, which up to a phase then overlaps with  $z$  a phase and  $r = 0$  as a limit of type (1).  $\square$

Note that our equations are invariant under global conjugation by an invertible  $u$ ,

$$(2.13) \quad C^i \mapsto uC^i u^{-1}, \quad S_i \mapsto uS_i u^{-1}, \quad \sigma_{S^j}^i \mapsto u\sigma_{S^j}^i u^{-1}, \quad J \mapsto \bar{u}J u^{-1},$$

where  $\sigma_{kl}^{ij} \in \mathbb{C}$  and hence invariant. If we allow this then we can conjugate the type (1)  $J$  by a diagonal matrix of determinant 1 to set  $r = 1$ . Hence, in calculations for the type (1) case, we will generally only consider  $r = 1$  for simplicity. Then for both types,  $J$  is given by a single complex parameter  $z$ .

This completes the ‘local’ side. For the inner product, we will assume a reference positive linear functional  $\int : A \rightarrow \mathbb{C}$  and set

$$(2.14) \quad \langle \phi_\alpha e^\alpha, \psi_\beta e^\beta \rangle = \int \phi_\alpha^* \mu^{\alpha\beta} \psi_\beta$$

for some positive hermitian matrix  $\mu$  as ‘measure’ and  $\phi, \psi \in \mathcal{S}$ . We use this to complete  $\mathcal{S}$  to a Hilbert space. One then needs to check that  $\mathcal{J}$  is an antilinear isometry, which, adapting the calculation in [20] and if we assume that  $\int$  is a trace, amounts to

$$(2.15) \quad \epsilon \mu^{\alpha\gamma} J^\beta_\gamma = J^\alpha_\gamma \mu^{\gamma\beta}.$$

The simplest case, which will also suffice for our purposes, is the identity matrix  $\mu^{\alpha\beta} = \delta^{\alpha\beta}$  as in [20]. Then the condition on  $J$  from this is that it be symmetric if  $\epsilon = 1$  and antisymmetric if  $\epsilon = -1$ . Finally,

$$\not{D}(\psi_\alpha e^\alpha) = \triangleright(\nabla_S(\psi)) = \triangleright(\partial_i \psi_\alpha s^i \otimes e^\alpha + \psi_\beta \nabla_S e^\beta) = (\partial_i \psi_\alpha + \psi_\beta S^\beta_{i\alpha}) C^{i\alpha}_\gamma e^\gamma$$

gives the resulting Dirac operator

$$(2.16) \quad (\not{D}\psi)_\gamma = (\partial_i \psi_\alpha + \psi_\beta S^\beta_{i\alpha}) C^{i\alpha}_\gamma$$

in our conventions. For a spectral triple, we need that  $\not{D}$  and the operator  $\gamma$  (if present) are antihermitian and hermitian respectively, which it is easiest at this point to check directly.

**2.3. Case of inner calculus.** We now specialise to the case where the calculus  $\Omega^1$  is inner with  $\theta = \theta_i s^i$  for  $\theta_i \in A$ . Then by [2, Prop. 8.11], we have  $\nabla s^i = \theta \otimes s^i - \sigma(s^i \otimes \theta) - \alpha(s^i)$  for some bimodule map  $\alpha(s^i) = \alpha^i_{jk} s^j \otimes s^k$ , which translates to

$$(2.17) \quad -\frac{1}{2} \Gamma^i_{jk} = \theta_j \delta^i_k - \theta_m \sigma^{im}_{jk} + \alpha^i_{jk}; \quad \alpha^i_{jk} \in \mathbb{C}.$$

Similarly for  $\nabla_S$  (see [24, Sec. 6.4]), we will have  $\nabla_S = \theta \otimes (\ ) - \sigma_S((\ ) \otimes \theta) + \alpha_S$  for  $\sigma_S$  as above and another bimodule map,

$$(2.18) \quad \alpha_S(e^\alpha) = A^\alpha_{i\beta} s^i \otimes e^\beta, \quad A^\alpha_{i\beta} \in \mathbb{C}, \quad S^\alpha_{i\beta} = \theta_i \delta^\alpha_\beta - \theta_j \sigma_S^{\alpha j}_{i\beta} + A^\alpha_{i\beta}.$$

For an inner calculus, one says that  $\nabla, \nabla_S$  are inner if they are given entirely by  $\sigma, \sigma_S$  respectively, i.e.  $\alpha^i_{jk} = 0$  and  $A^\alpha_{i\beta} = 0$  respectively.

**Lemma 2.2.** *Let  $\nabla$  be inner and write  $(\sigma_{Sj}^i)^\alpha_\beta := \sigma_{Sj\beta}^{\alpha i}$  for the matrix versions.  
(1) If  $\nabla_S$  is inner then equation (2.11) holds if and only if*

$$C^{\alpha i}_\beta \sigma_{Sj\gamma}^{\beta k} = \sigma_{j\gamma}^{im} \sigma_{S\beta}^{\alpha k} C^{l\beta}_\gamma; \quad C^i \sigma_{Sj}^k = \sigma_{j\gamma}^{im} \sigma_{Sm}^k C^l$$

and equation (2.5) holds if and only if

$$J^\alpha_\delta \delta_{jk} = \overline{\sigma_{S\beta}^{\alpha k}} J^\beta_\gamma \sigma_{Sj\delta}^{\gamma i}; \quad \delta_{jk} J = \overline{\sigma_{Si}^k} J \sigma_{Sj}^i.$$

(2) If  $\nabla_S$  is general with  $A^\alpha_{i\beta}$  then (2.11)-(2.5) are equivalent to the conditions in (1) and additionally

$$C^i A_j = \sigma^{ik}_{jl} A_k C^l, \quad J A_i = \overline{A_i} J \sigma_{Sj}^i.$$

*Proof.* (1) We take  $A_i = 0$  and the inner form of  $\Gamma^i_{jk}$  and substitute into (2.11) to obtain

$$\theta_m C^{\alpha i}_\beta \sigma_{Sj\gamma}^{\beta m} = \theta_m \sigma^{ik}_{jl} \sigma_{S\beta}^{\alpha m} C^{l\beta}_\gamma.$$

Now note that by the surjectivity assumption on a calculus, for each  $i$  there exist  $a_\mu^i, b_\mu^i \in A$  such that  $\sum_\mu a_\mu^i [\theta_j, b_\mu^i] s^j = s^i$ . Since the  $\{s^i\}$  are a basis, this means  $\sum_\mu a_\mu^i [\theta_j, b_\mu^i] = \delta_j^i$ . Hence, if  $\sum_j c_j \theta_j = 0$  in  $A$  for some  $c_i \in \mathbb{C}$  then  $c_j = 0$ , so the  $\theta_j$  are linearly independent whenever we have a central basis. Hence, we can cancel the  $\theta_m$  to obtain the result. Similarly, (2.5) becomes the condition stated on putting in  $A = 0$  and noting that  $\theta^* = -\theta$  so  $\theta_j^* = -\theta_j$  given our central self-adjoint basis. (2) For the last part, we repeat the above with the extra  $A$  terms on both sides. These are valued in multiples of 1 and this can be taken as linearly independent of the  $\theta_i$  since, given the central basis, any constant parts of  $\theta_i$  would not affect commutators in computing  $df$ . Hence, these parts of the equations decouple and appear as shown in matrix form.  $\square$

For an even spectral triple, the condition (2.10) similarly translates in the inner case to

$$(2.19) \quad [\sigma_{Sj}^i, \gamma] = 0, \quad [A_i, \gamma] = 0$$

again using that the  $\theta_i \in A$  are linearly independent and can be taken as linearly independent of 1.

### 3. DIRAC OPERATOR ON NONCOMMUTATIVE TORUS $\mathbb{C}_\theta[\mathbb{T}^2]$

The well-known noncommutative torus at an algebraic level is the  $*$ -algebra  $\mathbb{C}_\theta[\mathbb{T}^2]$  generated by unitary  $u, v$  with relations  $vu = e^{i\theta} uv$ . Here  $u^* = u^{-1}$  and  $v^* = v^{-1}$  and  $\theta$  a real parameter (theres is no confusion with inner structures as the calculus will not be inner). The Dirac operator here was one of the first constructions on the noncommutative torus, but here we revisit with the question of which quantum Riemannian geometries can realise it. We assume that  $\theta$  is irrational, in which case the algebra has trivial centre.



**3.1. Recap of WQRG of  $\mathbb{C}_\theta[\mathbb{T}^2]$ .** For  $\Omega^1(\mathbb{C}_\theta[\mathbb{T}^2])$ , the standard differential calculus has

$$(du)u = udu, \quad (dv)v = vdv, \quad (dv)u = e^{i\theta}udv, \quad (du)v = e^{-i\theta}vdu$$

with maximal prolongation

$$du \wedge du = 0, \quad dv \wedge dv = 0, \quad dv \wedge du = -e^{i\theta}du \wedge dv$$

as explained e.g. in [2, Chap. 1]. The first thing we do is choose a self-adjoint basis

$$s^1 = -iu^{-1}du, \quad s^2 = -iv^{-1}dv.$$

Then one can check that these are central and Grassmann with  $ds^i = 0$ . The partial derivatives defined by  $df = (\partial_i f)s^i$  for all  $f \in \mathbb{C}_\theta[\mathbb{T}^2]$  are the standard derivations

$$\partial_1(u^m v^n) = mu^m v^n, \quad \partial_2(u^m v^n) = nu^m v^n.$$

Next, a (generalised, i.e. not-necessarily quantum symmetric) quantum metric has the form

$$g = c_1 s^1 \otimes s^1 + c_2 s^2 \otimes s^2 + c_3 s^1 \otimes s^2 + c_4 s^2 \otimes s^1$$

for  $c_1, c_2, c_3, c_4 \in \mathbb{C}$ , since the metric has to be central to be bimodule-invertible. We suppose non-degeneracy, which amounts to  $c_1 c_2 \neq c_3 c_4$ , and focus on the quantum symmetric case  $c_4 = c_3$ . In this case the ‘reality’ of the metric says that  $c_i$  are real.

Similarly, it is shown [2, Example 8.16] that a torsion free bimodule connection must have the form

$$(3.1) \quad \nabla s^1 = h^1_1 s^1 \otimes s^1 + h^1_2 s^2 \otimes s^2 + h^1_3 (s^1 \otimes s^2 + s^2 \otimes s^1),$$

$$(3.2) \quad \nabla s^2 = h^2_1 s^1 \otimes s^1 + h^2_2 s^2 \otimes s^2 + h^2_3 (s^1 \otimes s^2 + s^2 \otimes s^1)$$

for constants  $h^i_j \in \mathbb{R}$  and  $\sigma = \text{flip}$  on the basis 1-forms. Putting this form in the requirements for a QLC, we find that  $\nabla s^i = 0$  is the only solution in the quantum symmetric case. More generally, we can ask only for the torsion and cotorsion to vanish, which happens in the quantum symmetric case when

$$(3.3) \quad c_2 h^2_1 = c_1 h^1_3 + c_3 (h^2_3 - h^1_1), \quad c_2 h^2_3 = c_1 h^1_2 + c_3 (h^2_2 - h^1_3),$$

giving a 4-parameter moduli of WQLC’s[2]. These have curvature

$$R_\nabla(s^1) = \rho^1_j s^1 \wedge s^2 \otimes s^j, \quad R_\nabla(s^2) = \rho^2_j s^1 \wedge s^2 \otimes s^j,$$

where[2]

$$\rho = S \begin{pmatrix} c_3 & c_2 \\ -c_1 & -c_3 \end{pmatrix}, \quad S = \begin{cases} \frac{h^1_2 h^2_1 - h^1_3 h^2_3}{c_1} & \text{if } c_3 \neq 0 \\ \frac{h^2_3 (h^2_3 - h^1_1)}{c_1} + \frac{h^1_3 (h^1_3 - h^2_2)}{c_2} & \text{if } c_3 = 0. \end{cases}$$

Here  $S$  is the Ricci scalar and the Ricci  $= \frac{Sq}{2}$  so that the ‘obvious’ Einstein tensor vanishes. We used the usual antisymmetric lift of  $s^1 \wedge s^1$ . We also will need a spinor connection:

**Lemma 3.1.** *For a bimodule connection on  $\mathcal{S}$  with central basis  $\{e^\alpha\}$  over  $\mathbb{C}_\theta[\mathbb{T}^2]$ ,  $\sigma_S$  has to be the flip map and  $S^\alpha_i \beta \in \mathbb{C}$ .*

*Proof.* Computing  $\nabla_S(e^\alpha a) = \nabla_S(ae^\alpha)$  using the two Leibniz rules, we have that

$$(\sigma_S^{\alpha j}{}_{i\beta} - \delta^j_i \delta^\alpha_\beta) \partial_j a = [a, S_i^\alpha{}_\beta]$$

for all  $a = u^m v^n$ , say. But  $\mathbb{C}_\theta[\mathbb{T}^2]$  has a  $\mathbb{Z} \times \mathbb{Z}$  grading according to the powers of  $u, v$  and the left hand side has the same grade  $(m, n)$  as  $a$ . If we write  $S_i^\alpha{}_\beta = \sum_{a,b} S_i^{(a,b)\alpha}{}_\beta u^a v^b$  then the  $(a, b)$  term has grade  $(m+a, n+b)$  on the right hand side of the displayed equation. Hence, we conclude that  $S_i^{(a,b)}$  can only contribute to the right hand side if  $(a, b) = (0, 0)$  and in this case the right hand side is zero. Hence,  $\sigma_S$  is necessarily the flip map. It then follows that  $S_i^\alpha{}_\beta \in \mathbb{C}$  as the centre of the algebra is trivial.  $\square$

This also requires that  $\nabla$  has to have  $\sigma$  the flip map and  $\mathbb{C}$ -number Christoffel symbols as in [2], but now argued more generally.

**3.2. Uniqueness of geometric spectral triple on  $\mathbb{C}_\theta[\mathbb{T}^2]$ .** To be concrete, we take  $c_1 = c_2 = 1$  and  $c_3 = 0$  so that  $g = s^1 \otimes s^1 + s^2 \otimes s^2$ . In this case, given the Grassmann form of  $\wedge$ , the Clifford relations (2.12) become  $C^i C^j + C^j C^i = 2g^{ij}$  which we solve for 2-dimensional  $\mathcal{S}$  and up to unitary equivalence by the usual choice

$$C^i = \sigma^i; \quad \sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

where we recalled the Pauli matrices used here for  $i = 1, 2$

**Lemma 3.2.** *For  $C^i = \sigma^i$ ,  $J$  has the type (1) form*

$$\epsilon = \epsilon' = 1: \quad J = q\sigma^1, \quad \epsilon = \epsilon' = -1: \quad J = q\sigma^2$$

*with  $|q| = 1$ . In either case, we also have  $\gamma = \sigma^3$  with  $\epsilon'' = -1$ , thereby solving all our equations at the ‘local’ level that do not directly involve  $S_i$ .*

*Proof.* Given that  $\sigma_S$  has to be the flip map, (2.6) reduces to  $\overline{C^i} J = \epsilon' J C^i$  as in [20]. Also (2.4) has the standard form  $\overline{J} J = \epsilon \text{id}$  with solutions in Lemma 2.1. In fact, the result is also easily found directly by writing  $J = j_0 \text{id} + j_i \sigma^i$  and noting that  $\overline{\sigma^2} = -\sigma^2$  while the other  $\sigma^i$  have real entries. So  $\sigma^1 J = \epsilon' J \sigma^1$  and  $\sigma^2 J = -\epsilon' J \sigma^2$ . We use the Pauli matrix algebra to reduce to the basis and thereby deduce that if  $\epsilon' = 1$  then only  $j_1$  is allowed, and if  $\epsilon' = -1$  then only  $j_2$  is allowed. We then check the other condition and find that  $\epsilon = \epsilon'$  and  $|j_1| = 1$  or  $|j_2| = 1$  respectively.  $J$  is always determined only up to a phase, which we denote  $q$ . It is then easy to solve (2.7)-(2.8) to find  $\gamma = \sigma^3$  as anticommuting with  $\sigma^1, \sigma^2$ . In this case (2.9) also holds provided  $\epsilon'' = -1$ , for either form of  $J$ .  $\square$

We next have to solve for  $\nabla_S$  to be compatible with  $\nabla, C^i, J$  so as to solve the algebraic part of the geometric spectral triple requirements.

**Lemma 3.3.** *For  $C^i = \sigma^i$ , the algebraic axioms for a geometrically realised spectral triple require the WQLC to be the QLC  $\nabla s^i = 0$  and  $\nabla s e^\alpha = d_i s^i \otimes e^\alpha$  for some real coefficients  $d_i$ . We also have  $\gamma$  at this level as in the preceding lemma, relevant for the even case.*

*Proof.* Since  $\sigma_S$  is the flip, (2.5) is  $\bar{S}_i J = J S_i$ . Next, since the braiding  $\sigma$  of the connection is also trivial, we need  $[C^i, S_j] = -\frac{1}{2}\Gamma^i_{jk}C^k$ . But writing  $\Gamma$  in terms of  $h^i_j$  in this case (where we solve the WQLC conditions (3.3) by  $h^1_3 = h^2_1$  and  $h^2_3 = h^1_2$ ), we have

$$-\frac{1}{2}\Gamma^i_{jk} = h^i_j \delta_{jk} + h^{\bar{i}}_i \delta_{\bar{j},k}$$

where  $\bar{1} = 2$  and  $\bar{2} = 1$  and  $i, j, k$  are now restricted to 1, 2. Hence the covariance condition (2.11) is

$$[C^i, S_j] = h^i_j C^j + h^{\bar{i}}_i C^{\bar{j}}.$$

Using Mathematica to search among this 4-parameter moduli of  $\nabla$ , we find that there are no solutions unless  $h^i_j = 0$ , i.e. the trivial QLC, and in this case  $S_1$  and  $S_2$  have to be real multiples of the identity since they have to commute with both  $C^i$  and commute up to complex conjugation with  $J$ . In this case, the remaining (2.10) is automatic.  $\square$

Hence the only geometrically realised Dirac operator for the standard Euclidean metric and  $\nabla$  a WQLC are

$$(3.4) \quad \not{D}(\psi_\alpha e^\alpha) = (\partial_i \psi_\alpha s^i) \triangleright e^\alpha + \psi_\alpha d_i s^i \triangleright e^\alpha = \sigma^{i\alpha}_\beta ((\partial_i + d_i) \psi_\alpha) e^\beta.$$

For a spectral triple, we still need to construct the Hilbert space, which we do with respect to the state  $\int u^m v^n = \delta_{m,0} \delta_{n,0}$  (this would be integration on the classical torus). We use this to complete  $\mathcal{S} = \mathbb{C}_\theta[\mathbb{T}^2] \otimes \mathbb{C}^2$  to a Hilbert space via (2.14) with  $\mu = \text{id}$  the identity matrix. As  $J$  is symmetric if  $\epsilon = 1$  and antisymmetric if  $\epsilon = -1$ , we already know that  $\mathcal{J}$  is an antilinear isometry in either case for this Hilbert space. The remaining arguments are similar to [20].

**Theorem 3.4.** *Up to unitary equivalence,  $(\not{D}\psi)_\beta = (\partial_i \psi_\alpha) \sigma^{i\alpha}_\beta$  is the only possibility for a geometrically realised spectral triple on the noncommutative torus for the Euclidean metric, a WQLC and the standard Hilbert space structure on  $\mathcal{S}$ . Moreover, we are forced to  $\nabla = \nabla_S = 0$  on the bases. Here  $\not{D}$  forms a spectral triple of KO-dimension  $n = 2$  if we choose  $\epsilon = \epsilon' = \epsilon'' = -1$  and  $n = 1$  if  $\epsilon = \epsilon' = 1$  ignoring  $\gamma$ .*

*Proof.* We have done the work in the two lemmas and just need to check if (3.4) is antihermitian and  $\gamma$  is hermitian. For the former,

$$\langle \not{D}\psi, \phi \rangle + \langle \psi, \not{D}\phi \rangle = \int (\sigma^{i\alpha}_\beta ((\partial_i + d_i) \psi_\alpha)^* \phi_\beta + \psi_\alpha^* \sigma^{i\beta}_\alpha (\partial_i + d_i) \phi_\beta) = 2 \int d_i \sigma^{i\beta}_\alpha \psi_\alpha^* \phi_\beta.$$

The terms without  $d_i$  vanish since  $\sigma^i$  are hermitian,  $\partial_i$  are derivations and  $\int \partial_i (\psi_\alpha^* \phi_\beta) = 0$  because  $\int \partial_i (u^m v^n) = n \int u^m v^n = 0$  for all  $m, n$ . For our expression to be zero for all  $\phi, \psi$ , this requires  $d_i = 0$ . For  $\gamma$ , we have more easily  $\langle \gamma\psi, \phi \rangle = \int (\sigma^{3\alpha}_\beta \psi_\alpha)^* \phi_\beta = \langle \psi, \gamma\phi \rangle = 0$  because  $\sigma^3$  is hermitian. Finally, looking in the period 8 table of signs, and allowing that  $\not{D}$  is antihermitian (the actual operator in the spectral triple is  $\iota \not{D}$ ) which flips the required value of  $\epsilon'$ , we see that we can meet the conditions for an even spectral triple with  $n = 2$  if we choose  $\epsilon = \epsilon' = -1$ . In the other case, we fit with  $n = 1$  and ignore  $\gamma$ .  $\square$

We have, in particular, recovered the standard  $n = 2$  spectral triple on the noncommutative torus, including a suitable charge conjugation  $J$ . Although not a new spectral triple, we see how it is forced from the quantum geometry up to unitarity for the given choice of sign (ditto for the other sign). Other symmetric

metrics will be equivalent after an orthogonal transformation to the Euclidean one and can therefore be handled in a manner similar to our treatment for the fuzzy sphere in [20], again giving a unique answer up to unitary equivalence for a given choice of signs. Asymmetric metrics, however, won't be compatible with (2.12) since, due to the Grassmann algebra form of the wedge product, we have to solve  $C^i C^j + C^j C^i = 2g^{ij}$ , which requires  $g^{ij}$  to be symmetric.

**3.3. Similar results for more general  $\nabla$ .** Returning to our Euclidean metric, which enters into the choice of  $C^i = \sigma^i$  for a 2-dimensional  $\mathcal{S}$ , we can let  $\nabla$  be any torsion free bimodule connection (3.1)-(3.2), which is a 6-parameter moduli of connections.

**Proposition 3.5.** *Extending the geometric realisation to allow any torsion free  $\nabla$  leads to the same answer as in Theorem 3.4 with  $\nabla = \nabla_S = 0$  on the bases. Allowing  $\nabla$  any bimodule connection leads to a real 4-parameter moduli of  $\nabla, \nabla_S$  but  $\mathcal{D}$  itself of the same form as in Lemma 3.3, resulting in the same spectral triple(s) as in Theorem 3.4.*

*Proof.* If we drop the WQLC condition but keep the torsion free form and repeat the calculation in Lemma 3.3 then we do not obtain any more solutions, we are again forced to  $\nabla = 0$  in the basis. If, however, we allow all 8 values of  $\Gamma^i_{jk}$  for a general bimodule connection, then we find solutions to (2.11) provided  $\nabla$  has the 2-parameter form

$$\nabla s^1 = h^1_2 s^2 \otimes s^2 - h^2_1 s^1 \otimes s^2, \quad \nabla s^2 = h^2_1 s^1 \otimes s^1 - h^1_2 s^2 \otimes s^1,$$

which we see is not torsion free unless it is zero, and

$$S_1 = \begin{pmatrix} a_1 & 0 \\ 0 & a_1 + \imath h^2_1 \end{pmatrix}, \quad S_2 = \begin{pmatrix} a_2 & 0 \\ 0 & a_2 - \imath h^1_2 \end{pmatrix}$$

for free parameters  $a_i \in \mathbb{C}$ . Then (2.5) for either choice of  $J$  requires the reality condition  $\bar{a}_1 = a_1 + \imath h^2_1$  and  $\bar{a}_2 = a_2 - \imath h^1_2$ , hence two further real parameters  $d_i \in \mathbb{R}$  with

$$a_1 = d_1 - \frac{\imath}{2} h^2_1, \quad a_2 = d_2 + \frac{\imath}{2} h^1_2; \quad S_1 = d_1 \text{id} - \frac{\imath}{2} h^2_1 \sigma^3, \quad S_2 = d_2 \text{id} + \frac{\imath}{2} h^1_2 \sigma^3$$

as the  $h^i_j$  are real. One can check that (2.10) still holds, so we have the algebraic properties for an even spectral triple, although only relevant for one of the sign choices.

Finally, we do a similar analysis as before for  $\mathcal{D}$ . From (2.16), we have

$$\begin{aligned} (\mathcal{D}\psi)_\gamma &= (\partial_i \psi_\alpha + d_i \psi_\alpha) \sigma^{i\alpha}_\gamma - \frac{\imath}{2} h^2_1 \psi_\beta (\sigma^3 \sigma^1)^\beta_\gamma + \frac{\imath}{2} h^1_2 \psi_\alpha (\sigma^3 \sigma^2)^\alpha_\gamma \\ &= (\partial_i \psi_\alpha + d_i \psi_\alpha) \sigma^{i\alpha}_\gamma + \frac{\psi_\alpha}{2} (h^2_1 \sigma^2 + h^1_2 \sigma^1)^\alpha_\gamma = (\partial_i \psi_\alpha + d'_i \psi_\alpha) \sigma^{i\alpha}_\gamma \end{aligned}$$

where  $d'_i = d_i + h^i_i/2$ . This is of the same form as in Lemma 3.3 so is not antihermitian unless  $d'_i = 0$ . The  $\gamma$  is the same as before and hermitian.  $\square$

We can go further and drop the  $\mathbb{W}(\triangleright) = 0$  covariance condition entirely. Then  $\nabla$  does not enter the construction at all and we can search among all bimodule connections  $\nabla_S$ . This extends the results of the preceding proposition to an even bigger moduli.

**Proposition 3.6.** *Geometric realisation without imposing  $\nabla(\triangleright) = 0$  leads to an real 8-parameter moduli of  $\nabla_S$  at the ‘local’ level, with a real 4-parameter moduli that are even with our chosen  $\gamma$ . The even case results in the same  $\not{D}$  as in Lemma 3.3 and the same spectral triple(s) as in Theorem 3.4. The odd case results in a 1-parameter moduli of spectral triples with real parameter  $m$ ,*

$$\epsilon = \epsilon' = 1 : \quad \not{D}\psi = \partial_i \psi \sigma^i + im\psi \sigma^3; \quad \epsilon = \epsilon' = -1, \quad \not{D}\psi = \partial_i \psi \sigma^i + im\psi.$$

*Proof.* If we search among all bimodule connections then we only have to solve the reality condition (2.5), which tells us that each of  $S_1, S_2$  are of the form

$$\epsilon = \epsilon' = 1 : \quad S_i = \begin{pmatrix} a_i & b_i \\ \bar{b}_i & \bar{a}_i \end{pmatrix}; \quad \epsilon = \epsilon' = -1 : \quad S_i = \begin{pmatrix} a_i & b_i \\ -\bar{b}_i & \bar{a}_i \end{pmatrix}$$

for the two choices of  $J$ . This is a real 8-parameter moduli of spinor connections. In either case, the condition (2.10) for an even structure  $\gamma = \sigma^3$  holds iff  $b_i = 0$ , which cuts us down to 4 real dimensions. We also write our general  $S_i$  in terms of Pauli matrices and combine with the  $C^i$  as in the proof of Proposition 3.5. The only contributions to  $\not{D}$  that are antihermitian are of the form stated with  $m = \text{Im}(b_1) + \text{Re}(b_2)$ , which is, however, zero if we want an even spectral triple.  $\square$

The odd spectral triples could be viewed as  $n = 1$  and  $n = 3$  respectively according to the signs (after flipping  $\epsilon'$ ). This proposition represents the most general spectral triple of geometric type on the noncommutative torus under the assumption of the standard Clifford structure for a spinor space of two dimensions.

#### 4. DIRAC OPERATOR ON MATRICES $M_2(\mathbb{C})$

We let  $\Omega^1(M_2(\mathbb{C}))$  be the standard 2D calculus as in [5][2, Chapter 1] with  $s, t$  central and  $s^* = -t$ . We first convert this to a more physical self-adjoint basis

$$s^1 = \frac{(s+t)}{i}, \quad s^2 = s-t; \quad s = \frac{1}{2}(is^1 + s^2), \quad t = \frac{1}{2}(is^1 - s^2),$$

so that  $s^{i*} = s^i$ . The extension to higher forms is by  $s^2 = t^2 = 0$ ,  $s \wedge t = t \wedge s$  with top form  $\text{Vol} := 2is \wedge t$  which is self-adjoint in the  $*$ -calculus. In terms of the  $s^i$ , the exterior algebra, which is not Grassmann, is

$$(4.1) \quad s^1 \wedge s^1 = s^2 \wedge s^2 = i\text{Vol}, \quad s^1 \wedge s^2 = s^2 \wedge s^1 = 0.$$

The exterior algebra is inner with  $d = [\theta, \cdot]$  (graded-commutator) which in our basis converts to

$$(4.2) \quad \theta = E_{12}s + E_{21}t = \frac{i}{2}(\sigma^1 s^1 + \sigma^2 s^2),$$

where we use some of the Pauli matrices. Using the three Pauli matrices and  $1 = \text{id}$  as a basis of  $M_2(\mathbb{C})$ , the differentials are then

$$(4.3) \quad d\sigma^1 = \sigma^3 s^2, \quad d\sigma^2 = -\sigma^3 s^1, \quad d\sigma^3 = \sigma^2 s^1 - \sigma^1 s^2, \quad ds^i = -\sigma^i \text{Vol}$$

as equivalent in our  $s^i$  terms to the formulae in [2] in terms of  $s, t$ . The corresponding partial derivatives defined by  $df = (\partial_i f)s^i$  are

$$(4.4) \quad \partial_i \sigma^j = -\epsilon_{ijk} \sigma^k, \quad \partial_i = \frac{i}{2}[\sigma^i, \cdot]$$

for  $i = 1, 2$  and  $j, k = 1, 2, 3$ . We will use this differential calculus for both the QRG in this section and the alternate QRG in the next section. In both cases,  $\Gamma^i_{jk}$  is *not constant* even though the metric has constant coefficients, which makes the analysis fundamentally harder than the fuzzy sphere case in [20] or the noncommutative torus case in Section 3.

**4.1. Standard quantum Riemannian geometry on  $M_2(\mathbb{C})$ .** The standard quantum metric  $M_2(\mathbb{C})$  with its 2D calculus in [2, Ex 8.11] (times 2) is  $g = 2(s \otimes t - t \otimes s)$ , which in our self-adjoint basis becomes

$$(4.5) \quad g = \iota(s^2 \otimes s^1 - s^1 \otimes s^2).$$

The QLC is not unique, but there is a known 4-parameter moduli of  $\ast$ -preserving QLCs. Within this, we focus for simplicity on 1-parameter sub-moduli space

$$\begin{aligned} \nabla s &= 2\theta \otimes s + \mu E_{21}(s \otimes t - t \otimes s) + \mu E_{12}(t \otimes t - s \otimes s) \\ \nabla t &= 2\theta \otimes t + \mu E_{12}(s \otimes t - t \otimes s) + \mu E_{21}(t \otimes t - s \otimes s) \\ \sigma(s \otimes s) &= (\mu - 1)s \otimes s - \mu t \otimes t, \quad \sigma(s \otimes t) = -\mu s \otimes t + (\mu - 1)t \otimes s \\ \sigma(t \otimes s) &= (-\mu - 1)s \otimes t + \mu t \otimes s, \quad \sigma(t \otimes t) = \mu s \otimes s + (-\mu - 1)t \otimes t \end{aligned}$$

with one real parameter  $\mu$ . This is given by setting  $\nu = -\mu$  and  $\beta = -\alpha \rightarrow 0$  in [2, Ex 8.13]. In terms of  $s^i$ , the connection appears as

$$\nabla s^i = 2\theta \otimes s^i - \mu \epsilon_{ik} \sigma^i s^i \otimes s^k, \quad \sigma(s^i \otimes s^j) = -s^j \otimes s^i - \delta_{ij} 2\mu \epsilon_{ik} s^i \otimes s^k$$

(sum over  $k$ ). We also compute the curvature:

**Proposition 4.1.** *The curvature of the above 1-parameter QLC for the metric (4.5) is*

$$R_\nabla(s^i) = -\mu \epsilon_{ij} \text{Vol} \otimes s^j$$

with Ricci curvature and Ricci scalar are

$$\text{Ricci} = \mu \frac{\iota}{2} (s^1 \otimes s^2 + s^2 \otimes s^1), \quad S = 0.$$

*Proof.* For the Riemann curvature we use that the calculus is inner to obtain

$$\begin{aligned} R_\nabla(s^i) &= (d \otimes \text{id} - \text{id} \wedge \nabla)(2\theta \otimes s^i - \mu \epsilon_{ik} \sigma^i s^i \otimes s^k) \\ &= 2\theta^2 \otimes s^i - 2\theta(2\theta \otimes s^i - \mu \epsilon_{ik} \sigma^i s^i \otimes s^k) - \mu \epsilon_{ik} d(\sigma^i s^i) \otimes s^k \\ &\quad + \mu \epsilon_{ik} \sigma^i s^i (2\theta \otimes s^k - \mu \epsilon_{km} \sigma^k s^k \otimes s^m) \\ &= \mu \epsilon_{ik} d(\sigma^i s^i) \otimes s^k - \mu^2 \epsilon_{ik} \epsilon_{km} \sigma^i \sigma^k s^i s^k \otimes s^m \end{aligned}$$

where in principle we sum over  $k, m$ . However,  $s^i s^k = 0$  unless  $k = i$  when  $\epsilon_{ik} = 0$ , so the second term vanishes. The first term simplifies using  $d(\sigma^i s^i) = -\text{Vol}$  to the result stated. For the Ricci curvature, we use the standard lift  $i(s \wedge t) = \frac{1}{2}(s \otimes t + t \otimes s)$  and the inverse metric  $(s, t) = 1 = -(t, s)$ , which translate to

$$i(\text{Vol}) = \frac{1}{2\iota} (s^1 \otimes s^1 + s^2 \otimes s^2), \quad (s^i, s^j) = -\iota \epsilon_{ij}$$

respectively. This then gives Ricci and  $S$  as stated.  $\square$

**4.2. Clifford action and spinor connection on  $M_2(\mathbb{C})$ .** Next we solve for the ‘local’ data for a geometrically realised Dirac operator for this metric. We take  $J$  from Lemma 2.1 as already analysed for a 2D calculus. Next, we analyse the content of the Clifford relations (2.12) for the matrices  $s^i \triangleright = C^i$  with respect to a central spinor basis  $\{e^\alpha\}$ . Given the form of the wedge product and of the metric, the full Clifford relations then come down to

$$(4.6) \quad C^2 C^1 = \text{id}, \quad C^1 C^2 + C^2 C^1 = 0, \quad (C^1)^2 = (C^2)^2.$$

The first two here have no joint solutions. However, we recall from [2, Prop. 8.45] that a more general version is to allow a nontrivial automorphism  $\phi : \mathcal{S} \rightarrow \mathcal{S}$  and a scale  $\kappa$  in the Clifford relations.

**Lemma 4.2.** *The full Clifford relations in a generalised form with automorphism hold if and only if  $C^2 C^1$  is invertible and the second half  $(C^1)^2 = (C^2)^2$ ,  $C^1 C^2 + C^2 C^1 = 0$  of (4.6) hold.*

*Proof.* The more general form of the Clifford relations (2.2) are that  $(\omega \wedge \eta) \triangleright := \kappa(\omega, \eta) \text{id} - \omega \triangleright (\eta \triangleright ( ))$  is well defined where  $\omega, \eta$  are 1-forms [2, Prop. 8.45]. Given the relations of  $\Omega^2$  in our case, and the off-diagonal form of the metric, this comes down in matrix terms in our basis to

$$\phi C^1 C^2 = \kappa g^{12} = -\iota \kappa, \quad \phi C^2 C^1 = \kappa g^{21} = \iota \kappa, \quad \phi((C^1)^2 - (C^2)^2) = \kappa(g^{11} - g^{22}) = 0$$

where  $g^{ij} = (s^i, s^j)$  is the inverse matrix to the coefficient matrix  $g_{ij}$  of the quantum metric. Without loss of generality, we can set  $\kappa = 1$  then we see that  $\phi = \iota(C^2 C^1)^{-1}$  is uniquely determined and the only conditions then are that the second half of (4.6) holds.  $\square$

In the first instance, we will look more broadly by imposing only the last of (4.6). Then one finds that there are principally 4 and 5 dimensional moduli spaces

$$\text{I: } C^2 = \pm C^1, \quad \text{II: } \text{Tr}(C^1) = \text{Tr}(C^2) = 0, \quad \det(C^1) = \det(C^2)$$

We focus on the more interesting type II case, which we parameterise as

$$C^1 = \begin{pmatrix} c_1 & c_2 \\ \frac{c_0 - c_1^2}{c_2} & -c_1 \end{pmatrix}, \quad C^2 = \begin{pmatrix} c_3 & c_4 \\ \frac{c_0 - c_3^2}{c_4} & -c_3 \end{pmatrix}; \quad c_i \in \mathbb{C},$$

where  $-c_0$  is the common value of the determinant. These include choosing the  $C^i$  as any two Pauli distinct matrices. Next we solve the covariance of  $\triangleright$  conditions.

**Theorem 4.3.** *For the  $C^i$  of type (II) as above and any  $J$  obeying (2.4), we can solve all the remaining local conditions (2.5) – (2.6) and (2.11) for a geometrically realised spectral triple of inner type if and only if  $\mu = 0$  in the 1-parameter QLC. In this case, solutions have the form*

$$\sigma_{S^j}^i = \zeta^i_j [C^1, C^2], \quad \overline{C^i} = \epsilon' \zeta^i_j J [C^1, C^2] C^j J^{-1}$$

for any invertible matrix  $\zeta \in M_2(\mathbb{C})$  obeying

$$\overline{\zeta} \zeta = \text{id}, \quad \det(\zeta) = -\frac{1}{\det([C^1, C^2])^2}.$$

Here  $[C^1, C^2]$  is the matrix commutator. Moreover, a canonical even  $\gamma$  obeying its local conditions (2.7)-(2.10) has the form

$$\gamma = \frac{[C^1, C^2]}{\sqrt{-\det([C^1, C^2])}}, \quad \bar{\gamma} = \epsilon'' J \gamma J^{-1}.$$

These solutions are subject to solving the stated reality conditions for  $C^i$  and  $\gamma$  respectively for some  $\epsilon', \epsilon''$ .

*Proof.* For the  $C^i$  of type (II) as above, we first solve (2.11) for  $\sigma_S$  in the inner form in Lemma 2.2 and find the generic solution which we recognise as stated but provided  $\mu = 0$ . We then insert this into the reality condition (2.6) and solve the resulting condition by setting  $\bar{C}^i$  as stated. Finally, we put the form of  $\sigma_S$  into (2.5) in the inner form in Lemma 2.2 to get

$$\delta_{jk} J = \bar{\zeta}^k_j [C^1, C^2] J \zeta^i_j [C^1, C^2].$$

We solve this as two equations, one for  $\bar{\zeta}^i_j$  as stated, which specifies the real and imaginary parts of  $\zeta^i_j$ , and the matrix equation

$$J = \overline{[C^1, C^2]} J [C^1, C^2].$$

We then solve the latter for  $J$  of type (2) in Lemma 2.1 to obtain just the constraint shown for  $\det(\zeta)$  and no constraint on the parameter  $z$  of  $J$ . We then check that in fact any  $J$  obeys this provided  $\det(\zeta)$  is as shown.

For the even structure  $\gamma$ , we put  $\sigma_S$  as found into (2.7),(2.8) to find generically a unique solution, which we identify as stated. We take (2.9) as a specification of  $\bar{\gamma}$  as stated. We automatically obey (2.10) in the form (2.19) since  $\sigma_{Sj}^i \propto [C^1, C^2]$  and this matrix commutes with  $\gamma$  since the  $C^i$  anticommute.  $\square$

This gives a canonical solution of our data for a spectral triple with 10 parameters: 5 for  $C^i$ , 3 for  $\zeta$  and 2 for  $J$  eg as in Lemma 2.1, modulo an overall unitary equivalence and for a choice of  $\epsilon, \epsilon'$ . These are subject to the reality constraints shown and we still need to solve for these. In the case of  $\zeta$ , the ‘reality condition’ is that of an antilinear isometry with respect to our basis and was already solved (with  $\epsilon = 1$ ) in Lemma 2.1 when we solved for  $J$ , i.e.

$$(1): \quad \zeta = e^{-i\zeta_0} \begin{pmatrix} \zeta_1 e^{i\zeta_3} & \zeta_2 \\ \frac{1-\zeta_1^2}{\zeta_2} & -\zeta_1 e^{-i\zeta_3} \end{pmatrix}, \quad (2): \quad \zeta = e^{-i\zeta_0} \begin{pmatrix} e^{i\zeta_2} & 0 \\ \zeta_1 & -e^{-i\zeta_2} \end{pmatrix}; \quad \zeta_i \in \mathbb{R}$$

in polar coordinate form with real parameters. The other difference is that we have explicitly included a phase factor at the front whereby  $\det(\zeta) = -e^{-2i\zeta_0}$ . For each  $\zeta$ , we still have to solve for  $C^i$  with

$$\det([C^1, C^2]) = 4c_0^2 - \frac{((c_2 c_3 - c_1 c_4)^2 - c_0 (c_2^2 + c_4^2))^2}{c_2^2 c_4^2} = e^{i\zeta_0}$$

and obeying the reality condition in the theorem. The system of equations for the latter is rather nonlinear and we do not give the most general solutions. Rather, we focus on solutions where the entries of  $J$  and  $C^i$  are real and some obvious choices of  $\zeta$ .



**Example 4.4.** We set  $\zeta = \sigma^3$  and  $J$  to be real of type (1),  $C^i$  with real entries and the simplifying assumption that  $c_2 = c_4 = 1$ . There are then no solutions with  $\epsilon = -1$ , while for  $\epsilon = 1$  we have:

(a) A 1-parameter family of solutions with  $\epsilon' = 1$ , namely

$$J = \begin{pmatrix} x & 2 \\ \frac{1-x^2}{2} & -x \end{pmatrix}, \quad C^1 = \begin{pmatrix} \frac{x+1}{2} & 1 \\ -(\frac{x+1}{2})^2 - \frac{x+1}{2} \end{pmatrix}, \quad C^2 = \begin{pmatrix} \frac{x-1}{2} & 1 \\ -(\frac{x-1}{2})^2 - \frac{1-x}{2} \end{pmatrix}$$

with a single real parameter  $x \in \mathbb{R}$ . Here  $\det(C^i) = 0$ , while  $\det([C^1, C^2]) = -1$  as required, and the relations

$$C^1 C^2 + C^2 C^1 = -\text{id}.$$

Moreover, the canonical form of even structure works and is given by

$$\gamma = [C^1, C^2] = J, \quad \epsilon'' = 1.$$

We have similar solutions with the  $C^i$  swapped and/or with the off-diagonals of  $J$  negated and  $C^i$  negated.

(b) A 1-parameter family with  $\epsilon' = -1$ , namely the same  $J$  as before and

$$C^1 = \begin{pmatrix} \frac{x+1}{2} & 1 \\ \frac{1-x(x+2)}{4} - \frac{x+1}{2} \end{pmatrix}, \quad C^2 = \begin{pmatrix} \frac{x-1}{2} & 1 \\ \frac{1-x(x-2)}{4} - \frac{1-x}{2} \end{pmatrix}$$

with  $\det(C^i) = -\frac{1}{2}$ . These have

$$C^1 C^2 + C^2 C^1 = 0$$

so that the full Clifford relations hold in a generalised form with automorphism. Moreover, the canonical form of even structure works and is given by

$$\gamma = -\imath[C^1, C^2] = -\imath \begin{pmatrix} x & 2 \\ -\frac{1+x^2}{2} & -x \end{pmatrix}, \quad \epsilon'' = 1.$$

There are similarly 3 variants as in case (a). Also note that  $J$  in the two cases can never be symmetric.

**Remark 4.5.** The partially-geometrically realised spectral triple on  $M_2(\mathbb{C})$  in [5, 2] has the same calculus and quantum metric as used above. Working with our self-adjoint basis  $s^i$  and taking  $e^1 = e, e^2 = f$  in our conventions, one has  $\epsilon = \epsilon' = -1$  and

$$J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = -\imath\sigma^2, \quad C^1 = -\imath\sigma^1, \quad C^2 = \imath\sigma^2, \quad [C^1, C^2] = \imath\sigma^3, \quad \sigma_{S^j}^i = \delta^i_j \text{id}.$$

Here  $C^1 C^2 + C^2 C^1 = 0$  and  $(C^1)^2 = (C^2)^2$  (both being  $-\text{id}$ ), so that the full Clifford relations hold with a suitable automorphism  $\phi \propto \sigma^3$ . But this example fundamentally did not have covariance of  $\triangleright$  i.e. (2.5) fails as  $S_i = 0$  while the  $\mu = 0$  QLC does not have zero Christoffel symbols. By contrast, we see that we can have solutions with  $\epsilon = 1, \epsilon' = -1$  with covariance of  $\triangleright$  as well as the full Clifford relations generalised to allow an automorphism  $\phi$ . The difference is that  $\zeta = \sigma_3$  and  $\sigma_S$  also has a nontrivial operation  $[C^1, C^2]$  as per Theorem 4.3 rather than the identity on the spinor indices.

The other obvious real choice of  $\zeta$  also has solutions as follows.

**Example 4.6.** We set  $\zeta = \sigma^1$  and  $J$  to be real of type (1) and  $C^i$  real with  $c_2 = c_4 = 1$ . There are then no solutions with  $\epsilon = -1$ , while for  $\epsilon = 1$  we find have:

(a) A 2-parameter family, for either sign of  $\epsilon'$ , namely

$$J = \begin{pmatrix} \frac{\epsilon' + x^2 - y^2}{\sqrt{2}} & \sqrt{2}(x-y) \\ \frac{1 - \frac{1}{2}(\epsilon' + x^2 - y^2)^2}{\sqrt{2}(x-y)} & -\frac{\epsilon' + x^2 - y^2}{\sqrt{2}} \end{pmatrix},$$

$$C^1 = \begin{pmatrix} x & 1 \\ \frac{(x-y)^4 + 1}{4(x-y)^2} - x^2 & -x \end{pmatrix}, \quad C^2 = \begin{pmatrix} y & 1 \\ \frac{(x-y)^4 + 1}{4(x-y)^2} - y^2 & -y \end{pmatrix}$$

with

$$C^1 C^2 + C^2 C^1 = \frac{1}{2} \left( \frac{1}{(x-y)^2} - (x-y)^2 \right) \text{id}, \quad \det(C^2 C^1) = \frac{((x-y)^4 + 1)^2}{16(x-y)^4}.$$

Hence, if we set  $x - y = \pm 1$  then  $C^1 C^2 + C^2 C^1 = 0$  and  $C^2 C^1$  is invertible, i.e. the full Clifford relations hold in the generalised form with automorphism. Moreover, the canonical form of even structure works and is given by

$$\gamma = -\imath[C^1, C^2] = -\imath \begin{pmatrix} x^2 - y^2 & 2(x-y) \\ -\frac{(x^2 - y^2)^2 + 1}{2(x-y)} & y^2 - x^2 \end{pmatrix}, \quad \epsilon'' = 1.$$

There is a similar solution with the opposite sign of  $\sqrt{2}$ . Note also that one can choose  $x, y$  such that  $J$  is symmetric but not at the points where  $x - y = \pm 1$ .

(b) There is a 1-parameter family for  $\epsilon' = -1$  only, namely

$$J = \begin{pmatrix} 0 & \sqrt{4x(x - \sqrt{x^2 - 1}) - 2} \\ \frac{1}{\sqrt{4x(x - \sqrt{x^2 - 1}) - 2}} & 0 \end{pmatrix}, \quad C^1 = \begin{pmatrix} x & 1 \\ -\frac{1}{2} & -x \end{pmatrix}, \quad C^2 = \begin{pmatrix} \sqrt{x^2 - 1} & 1 \\ \frac{1}{2} & -\sqrt{x^2 - 1} \end{pmatrix}$$

with

$$C^1 C^2 + C^2 C^1 = 2x\sqrt{x^2 - 1} \text{id}, \quad \det(C^2 C^1) = \frac{1}{4} + x^2(x^2 - 1).$$

Here  $|x| \geq 1$ , so the  $C^i$  anticommute only at  $x = \pm 1$ . Moreover, the canonical form of even structure works and is given by

$$\gamma = -\imath[C^1, C^2] = -\imath \begin{pmatrix} 1 & 2(x - \sqrt{x^2 - 1}) \\ -\sqrt{x^2 - 1} - x & -1 \end{pmatrix}, \quad \epsilon'' = 1.$$

There are 3 similar solutions with different signs for the two types of square roots independently. Note also that  $J$  is symmetric at  $x = \frac{3}{2\sqrt{2}}$ .

(c) A 1-parameter family of solutions, for either sign of  $\epsilon'$ , namely

$$J = \begin{pmatrix} \frac{1}{\sqrt{2}} & \epsilon' 2\sqrt{2}x \\ \frac{\epsilon'}{4\sqrt{2}x} & -\frac{1}{\sqrt{2}} \end{pmatrix}, \quad C^1 = \begin{pmatrix} x & 1 \\ \frac{1}{16x^2} & -x \end{pmatrix}, \quad C^2 = \begin{pmatrix} -x & 1 \\ \frac{1}{16x^2} & x \end{pmatrix}$$

$$C^1 C^2 + C^2 C^1 = \left( \frac{1}{8x^2} - 2x^2 \right) \text{id}, \quad \det(C^2 C^1) = \frac{1}{8} + x^4 + \frac{1}{256x^4}.$$

We see that  $C^1 C^2 + C^2 C^1 = 0$  at  $x = \pm \frac{1}{2}$  and  $C^2 C^1$  is always invertible, so in this case we have the full Clifford relations in the generalised form with automorphism. Moreover, the canonical form of even structure works and is given by

$$\gamma = -\imath[C^1, C^2] = -\imath \begin{pmatrix} 0 & 4x \\ -\frac{1}{4x} & 0 \end{pmatrix}, \quad \epsilon'' = 1.$$

There is a similar solution with the opposite sign of  $\sqrt{2}$ . Note also that  $J$  is symmetric at  $x = \pm \frac{1}{4}$ .

We can equally let  $J$  be of type (2), again restricting to real entries at least for the  $C^i$ . The algebra is easier and we do not need to impose further conditions on  $c_2, c_4$ .

**Example 4.7.** We set  $\zeta = \sigma^3$  and  $J$  to be type (2) with  $\epsilon = 1$  and real entries for  $C^i$ . Then for either sign of  $\epsilon'$ , we have two 2-parameter solutions:

$$J = \begin{pmatrix} 1 & 0 \\ -\frac{2x \pm 1}{y} & -1 \end{pmatrix}, \quad C^1 = \begin{pmatrix} x & y \\ \frac{\frac{1}{2} - x^2}{y} & -x \end{pmatrix}, \quad C^2 = \epsilon' \begin{pmatrix} x \pm 1 & y \\ \frac{\frac{1}{2} - (x \pm 1)^2}{y} & -(x \pm 1) \end{pmatrix},$$

$$C^1 C^2 + C^2 C^1 = 0, \quad \gamma = -\imath[C^1, C^2] = -\imath \epsilon' \begin{pmatrix} -1 \mp 2x & \mp 2y \\ \pm \frac{2x(x \pm 1) + 1}{y} & 1 \pm 2x \end{pmatrix}$$

with  $\epsilon'' = 1$ . Here  $C^1 C^2$  is invertible and hence this obeys the full Clifford relations in the general form with automorphism. Note that  $J$  is symmetric at  $x = \mp \frac{1}{2}$  and that the  $y$  parameter can be absorbed in a conjugation and hence we could set  $y = c_2 = c_4 = 1$  as we did before. There are also some 1-parameter solutions with only this  $y$  parameter, i.e. discrete if we ignore this freedom. The matrix entries of  $J$  here are also real but allowing the full complex case does not lead to fundamentally new solutions.

We find no solutions for  $\zeta = \sigma^1$  under similar assumptions. It should be clear that there are potentially many other solutions, because we do not need to assume  $c_2 = c_4 = 1$  in the  $J$  type (1) examples and we do not need to restrict to real entries for  $C^i$  or  $J$ . The former is not too impactful given the freedom to conjugate solutions, while the latter holds classically in two dimensions if one uses  $\sigma^1, \sigma^3$  for the Clifford algebra as these have real entries.

**4.3. Natural spectral triple on  $M_2(\mathbb{C})$ .** Looking at the ‘local’ solutions in Section 4.2 for the spectral triple geometric data (i.e. before considering the Hilbert space structure), all solutions under our assumptions have  $\epsilon = 1$ . Moreover, only ones for which the  $C^1, C^2$  anticommute (and the full Clifford relations hold with automorphism) and  $J$  is symmetric (so that  $\mathcal{J}$  is an antilinear isometry with respect to the usual trace inner product given by  $\mu^{\alpha\beta} = \delta^{\alpha\beta}$  for the measure) are  $x = \pm \frac{1}{2}$  in Example 4.7. This is therefore the natural solution under our simplifying assumptions, mainly of real entries in the  $C^i$ . The value of  $y$  not significant as it can be changed by conjugation.

To be concrete, henceforth we focus in particular on  $x = -\frac{1}{2}$  and  $y = \frac{1}{2}$  in Example 4.7 so that

$$(4.7) \quad J = \sigma^3 = \zeta, \quad C^1 = \frac{\sigma^1 - \sigma^3}{2}, \quad C^2 = \epsilon' \frac{\sigma^1 + \sigma^3}{2}, \quad \gamma = -\epsilon' \sigma^2, \quad \epsilon = 1 = \epsilon''$$

in terms of Pauli matrices. Then  $\mathcal{J}$  is an antilinear isometry. We also need  $\gamma$  to be hermitian. As an operator,  $\gamma(e^\alpha) = \imath \epsilon' \epsilon_{\alpha\beta} e^\beta$  (with sum over repeated indices), where  $\epsilon_{\alpha\beta}$  is the antisymmetric tensor with  $\epsilon_{12} = 1$ . Hence on the spinor components of  $\psi = \psi_\alpha e^\alpha$ , we have likewise

$$(4.8) \quad (\gamma\psi)_\beta = -\epsilon' \psi_\alpha \sigma^{2\alpha}_\beta = \imath \epsilon' \psi_\alpha \epsilon_{\alpha\beta}.$$

Then

$$\langle \gamma\phi, \psi \rangle = \text{Tr}(\gamma\phi)_\beta^\dagger \psi_\beta = \text{Tr}(\imath \epsilon' \epsilon_{\alpha\beta} \phi_\alpha)_\beta^\dagger \psi_\beta = \text{Tr} \phi_\alpha^\dagger (-\imath \epsilon' \epsilon_{\alpha\beta} \psi_\beta) = \text{Tr} \phi_\alpha^\dagger (\gamma\psi)_\alpha = \langle \phi, \gamma\psi \rangle$$

as required.

**Proposition 4.8.** *The solution (4.7) for the local geometric data gives a quantum geometrically realised  $n = 0$  even spectral triple if  $\epsilon' = -1$  with*

$$(\not{D}\psi)_\beta = -\frac{\imath}{4} ([-\sigma^1 + \sigma^2, \psi_\alpha] \sigma^{1\alpha}_\beta + \{\sigma^1 + \sigma^2, \psi_\alpha\} \sigma^{3\alpha}_\beta)$$

*and an  $n = 1$  odd spectral triple if  $\epsilon' = 1$  with*

$$(\not{D}\psi)_\beta = \frac{\imath}{4} (\{\sigma^1 + \sigma^2, \psi_\alpha\} \sigma^{1\alpha}_\beta + [-\sigma^1 + \sigma^2, \psi_\alpha] \sigma^{3\alpha}_\beta).$$

*Proof.* We use (2.16) and the form of  $\partial_i$  in (4.4) and of  $S^\alpha_{i\beta}$  in (2.18) gives the Dirac operator

$$(4.9) \quad (\not{D}\psi)_\gamma = \frac{\imath}{2} (\sigma^i \psi_\alpha - \psi_\beta \sigma^j \sigma_S^{\beta j}_{i\alpha}) C^{i\alpha}_\gamma.$$

For the chosen solution and its form of  $\sigma_{Sj}^i$ , and noting that its  $C^i$  obey

$$[C^1, C^2]C^1 = -C^2, \quad [C^1, C^2]C^2 = C^1,$$

we have

$$(4.10) \quad (\not{D}\psi)_\beta = \frac{\imath}{4} ((\sigma^1 \psi_\alpha + \psi_\alpha \sigma^2)(\sigma^1 - \sigma^3)^\alpha_\beta + \epsilon' (\sigma^2 \psi_\alpha + \psi_\alpha \sigma^1)(\sigma^1 + \sigma^3)^\alpha_\beta)$$

which, collecting terms, we can write as stated. Note that the  $\sigma^1, \sigma^2$  coming from  $\theta$  are simply elements of the algebra while the  $\sigma^1, \sigma^3$  coming from the  $C^1, C^2$  etc. act as shown explicitly on spinor indices. Hence

$$\begin{aligned} \langle \phi, \not{D}\psi \rangle &= \frac{\imath}{4} \text{Tr} \phi_\alpha^\dagger ((\sigma^1 \psi_\beta + \psi_\beta \sigma^2)(\sigma^1 - \sigma^3)^\beta_\alpha + \epsilon' (\sigma^2 \psi_\beta + \psi_\beta \sigma^1)(\sigma^1 + \sigma^3)^\beta_\alpha) \\ \langle \not{D}\phi, \psi \rangle &= -\frac{\imath}{4} \text{Tr} ((\sigma^1 \phi_\alpha + \phi_\alpha \sigma^2)(\sigma^1 - \sigma^3)^\alpha_\beta + \epsilon' (\sigma^2 \phi_\alpha + \phi_\alpha \sigma^1)(\sigma^1 + \sigma^3)^\alpha_\beta)^\dagger \psi_\beta \\ &= -\frac{\imath}{4} \text{Tr} ((\phi_\alpha^\dagger \sigma^1 + \sigma^2 \phi_\alpha^\dagger) \psi_\beta (\sigma^1 - \sigma^3)^\beta_\alpha + \epsilon' (\phi_\alpha^\dagger \sigma^2 + \sigma^1 \phi_\alpha^\dagger) \psi_\beta (\sigma^1 + \sigma^3)^\beta_\alpha) \\ &= -\langle \phi, \not{D}\psi \rangle \end{aligned}$$

so that  $\not{D}$  is antihermitian. In the second computation, we used that  $\sigma^1 \pm \sigma^3$  are real to move them out from under  $(\ )^\dagger$  and that they are symmetric to swap  $\alpha, \beta$ . We also used that  $\sigma^1, \sigma^2 \in M_2(\mathbb{C})$  are hermitian as elements of the algebra. We then compare the terms using cyclicity of the trace to obtain  $-\langle \phi, \not{D}\psi \rangle$ .

Finally, remembering that the Dirac operator in a Connes spectral triple is hermitian and hence corresponds to  $D = \imath \not{D}$ , the interpretation of  $\epsilon'$  is reversed. Hence our above construction is  $n = 0$  in the period 8 classification if  $\epsilon' = -1$ , as an even spectral triple, and  $n = 1$  ignoring  $\gamma$ , as an odd spectral triple if  $\epsilon' = 1$ .  $\square$

One can check directly that the stated  $D = \imath \not{D}$ ,  $\gamma$  in (4.8) and

$$(4.11) \quad (\mathcal{J}\psi)_\beta = \psi_\alpha^\dagger \sigma^{3\alpha}_\beta$$

indeed obeys all the required axioms for a Connes spectral triple. Here

$$(\mathcal{J}b\mathcal{J}^{-1}\psi)_\beta = \psi_\beta b^\dagger, \quad ([D, a]\psi)_\beta = \frac{1}{4}([\sigma^1 - \sigma^2, a]\psi_\alpha \sigma^{1\alpha}_\beta + [\sigma^1 + \sigma^2, a]\psi_\alpha \sigma^{3\alpha}_\beta)$$

in the even case, and a similar formula with  $\sigma^1, \sigma^3$  swapped and ignoring  $\gamma$  in the odd case. Moreover, underlying  $D$  is the bimodule connection  $\nabla_S$  which for our basis has Christoffel symbols

$$(4.12) \quad S_i^\alpha_\beta = \frac{\imath}{2}(\sigma^i - \sigma^j \sigma^{3j}_i [C^1, C^2]^\alpha_\beta); \quad S_1 = \frac{\imath}{2}\sigma^1 \begin{pmatrix} 1 & \epsilon' \\ -\epsilon' & 1 \end{pmatrix}, \quad S_2 = \frac{\imath}{2}\sigma^2 \begin{pmatrix} 1 & -\epsilon' \\ \epsilon' & 1 \end{pmatrix}.$$

We have seen that this arises naturally up to conjugation under some modest assumptions and the requirement of geometric realisability.

**Corollary 4.9.** *The above solution (4.7) for the geometric data is unitarily equivalent to*

$$C^1 = \frac{1}{\sqrt{2}}\sigma^1, \quad C^2 = \frac{\epsilon'}{\sqrt{2}}\sigma^3, \quad J = \frac{1}{\sqrt{2}}(\sigma^3 - \sigma^1)$$

with  $\gamma, S_1, S_2$  unchanged and gives  $n = 0, 1$  spectral triples with

$$(\not{D}\psi)_\beta = \frac{\imath}{2\sqrt{2}}((\sigma^1\psi_\alpha + \psi_\alpha\sigma^2)\sigma^{1\alpha}_\beta + \epsilon'(\psi_\alpha\sigma^1 + \sigma^2\psi_\alpha)\sigma^{3\alpha}_\beta).$$

*Proof.* Here we apply (2.13), i.e. a change of the  $\{e^\alpha\}$  basis, via the orthogonal real matrix

$$u = \begin{pmatrix} \frac{\sqrt{2+\sqrt{2}}}{2} & \frac{\sqrt{2-\sqrt{2}}}{2} \\ -\frac{\sqrt{2-\sqrt{2}}}{2} & \frac{\sqrt{2+\sqrt{2}}}{2} \end{pmatrix}$$

to obtain new  $C^i, J$  as stated and  $u\not{D}u^{-1}$  for  $\not{D}$  in (4.10) obtain the new Dirac operator stated. As a check, one obtains the same new  $\not{D}$  from (2.16) using  $\partial_i$  from (4.4), the new  $C^i, J$  and the connection  $\nabla_S$  with  $S_i$  given by (4.12).  $\square$

## 5. DIRAC OPERATOR FOR ALTERNATE QRG OF $M_2(\mathbb{C})$

In this section we repeat the analysis as in Section 4 with the same  $\Omega^1$  and self-adjoint basis  $s^i$ , but now for the alternate quantum metric [2, Exercise 8.5], which is the only other one for which the QLCs have been analysed.

**5.1. Alternate quantum Riemannian geometry on  $M_2(\mathbb{C})$ .** Here the alternate metric (times 2) is  $2(s \otimes s + t \otimes t)$ , which in the self-adjoint basis comes out as

$$(5.1) \quad g = -s^1 \otimes s^1 + s^2 \otimes s^2,$$

with a ‘Minkowski signature’ metric. The most natural 1-parameter QLC here (not the most general, but sufficiently nontrivial for our purposes) is from [2, Exercise 8.3],

$$\begin{aligned} \nabla s &= 2E_{21}t \otimes s - \rho E_{12}(s \otimes t - t \otimes s) + \rho E_{21}(t \otimes t - s \otimes s), \\ \nabla t &= 2E_{12}s \otimes t - \rho E_{21}(s \otimes t - t \otimes s) + \rho E_{12}(t \otimes t - s \otimes s), \end{aligned}$$

where  $\rho$  is an *imaginary* parameter. This is again of an inner form determined by

$$\begin{aligned}\sigma(s \otimes s) &= s \otimes s + \rho(s \otimes t - t \otimes s), & \sigma(t \otimes t) &= t \otimes t + \rho(s \otimes t - t \otimes s), \\ \sigma(s \otimes t) &= -t \otimes s + \rho(s \otimes s - t \otimes t), & \sigma(t \otimes s) &= -s \otimes t + \rho(s \otimes s - t \otimes t).\end{aligned}$$

The braiding converts to

$$\begin{aligned}\sigma(s^1 \otimes s^1) &= -s^2 \otimes s^2 - 2i\rho s^2 \otimes s^1, & \sigma(s^1 \otimes s^2) &= s^1 \otimes s^2, \\ \sigma(s^2 \otimes s^1) &= s^2 \otimes s^1, & \sigma(s^2 \otimes s^2) &= -s^1 \otimes s^1 - 2i\rho s^1 \otimes s^2,\end{aligned}$$

which we can write as

$$(5.2) \quad \sigma(s^i \otimes s^j) = \begin{cases} s^i \otimes s^j & \text{if } i \neq j \\ -s^{\bar{i}} \otimes s^{\bar{i}} - 2i\rho s^{\bar{i}} \otimes s^i & \text{if } i = j, \end{cases}$$

where  $\bar{1} = 2$  and  $\bar{2} = 1$ . We can then recompute the connection as

$$\begin{aligned}\nabla s^i &= \theta \otimes s^i - \sigma(s^i \otimes \theta) = \frac{i\sigma_j}{2}(s^j \otimes s^i - \sigma(s^i \otimes s^j)) \\ &= \frac{i}{2}\sigma^i(s^1 \otimes s^1 + s^2 \otimes s^2) + \frac{i}{2}\epsilon_{ij}\sigma^j(s^2 \otimes s^1 - s^1 \otimes s^2) - \rho\sigma^i s^{\bar{i}} \otimes s^i,\end{aligned}$$

which one can check agrees with what we had in terms of  $s, t$ . These coefficients define the  $\Gamma^i_{jk}$ .

The Riemann curvature and Ricci tensor for the canonical lift  $i(s \wedge t) = \frac{1}{2}(s \otimes t + t \otimes s)$  translate to

$$(5.3) \quad R_{\nabla} s^i = -i(1 + \rho^2)\text{Vol} \otimes s^i, \quad \text{Ricci} = -\frac{1}{2}(1 + \rho^2)(s^1 \otimes s^1 + s^2 \otimes s^2),$$

which is real in the sense of a metric for the  $*$ -structures, but not quantum symmetric. The Ricci scalar vanishes, so the QRG is Ricci-scalar flat but not necessarily flat.

**5.2. Clifford action and spinor connection for alternate QRG.** We next solve for  $C^i$ . Thus, the induced ‘Clifford relations’ (2.12) come down in matrix terms to

$$(5.4) \quad C^1 C^2 = C^2 C^1 = 0, \quad -(C^1)^2 + (C^2)^2 = 2\text{id}$$

with the largest component (2-parameter family) of solutions

$$(5.5) \quad C^1 = \pm i \begin{pmatrix} \sqrt{2} - a & b \\ \frac{a}{b}(\sqrt{2} - a) & a \end{pmatrix}, \quad C^2 = \pm \begin{pmatrix} a & -b \\ -\frac{a}{b}(\sqrt{2} - a) & \sqrt{2} - a \end{pmatrix}, \quad a, b \in \mathbb{C}.$$

One can check that these  $C^i$  admit no solutions for  $J, \sigma_S$  and we must look more generally. The proof involves solving the covariance of  $\triangleright$  for parameters  $a, b$ , leading to a particular form of  $\sigma_S$ , which then fails (2.6) using the fact that  $\text{Tr}(C^1) = \pm i\sqrt{2}$ ,  $\text{Tr}(C^2) = \pm\sqrt{2}$  are imaginary/real, which prescribes also the conjugated  $\text{Tr}(\bar{C}^i)$ . We therefore look more generally by dropping the first half of the relations (5.4).

**Proposition 5.1.** *If we impose only  $-(C^1)^2 + (C^2)^2 = 2\text{id}$  then there are principally 5D and 4D moduli and we parametrise these by  $c_{11}, c_{12}, d_{11}, d_{12}$  and where applicable  $d_{21}$  as three types of solutions,*

$$\text{I: } C^1 = \begin{pmatrix} c_{11} & c_{12} \\ \frac{-c_{11}^2 + d_{11}^2 + d_{12}d_{21} - 2}{c_{12}} & -c_{11} \end{pmatrix}, \quad C^2 = \begin{pmatrix} d_{11} & d_{12} \\ d_{21} & -d_{11} \end{pmatrix};$$

$$\begin{aligned} \text{II: } C^1 &= \begin{pmatrix} c_{11} & c_{12} \\ \frac{c_{12}(c_{11}^2-d_{11}^2+2)}{d_{12}^2-c_{12}^2} & -c_{11} \end{pmatrix}, \quad C^2 = \begin{pmatrix} d_{11} & d_{12} \\ \frac{d_{12}(c_{11}^2-d_{11}^2+2)}{d_{12}^2-c_{12}^2} & -d_{11} \end{pmatrix}; \\ \text{III: } C^1 &= \begin{pmatrix} c_{11} & c_{12} \\ \frac{c_{12}(c_{11}^2-d_{11}^2+2)}{d_{12}^2-c_{12}^2} & \frac{-2c_{12}d_{11}d_{12}+c_{11}(c_{12}^2+d_{12}^2)}{d_{12}^2-c_{12}^2} \end{pmatrix}, \\ C^2 &= \begin{pmatrix} d_{11} & d_{12} \\ \frac{d_{12}(c_{11}^2-d_{11}^2+2)}{d_{12}^2-c_{12}^2} & \frac{-2c_{11}c_{12}d_{12}+d_{11}(c_{12}^2+d_{12}^2)}{d_{12}^2-c_{12}^2} \end{pmatrix}. \end{aligned}$$

There are then no solutions for  $J, \sigma_S$  in type I or for type III in the case (5.5).

*Proof.* This is by calculations using Mathematica. In all types, solving for the covariance of  $\triangleright$  in Lemma 2.2 gives a 2-parameter moduli of the  $\sigma_S, r$  for generic  $C^i$  in the class. For the last part, we note that (2.6) says that  $\bar{C}^i$  is conjugate to  $C^i$ , from which it follows that

$$(5.6) \quad \text{Tr}(\sigma_{Sj}^i C^j) = 0, \quad -(\sigma_{Si}^1 C^i)^2 + (\sigma_{Si}^2 C^i)^2 = 2\text{id}$$

in type I,II for the trace identities and in all types for the other identity. In type I, we solve the covariance of  $\triangleright$  condition and the trace identities together. There are no solutions for general  $C^i$  but if we solve also for one of their entries then we obtain 1-parameter solutions where  $(\sigma_{S1}^1)_{11}$  is undetermined but  $d_{21}$  is prescribed. We are then not able to impose the other half of (5.6). The analysis for general type III appears rather more complex as we have only the 2nd half of (5.6), but (5.5) is a special case where the trace has a fixed property under complex conjugation as noted and similar methods then apply to establish no solutions.  $\square$

The case of type II therefore appears most natural and we are led to focus on this. In this case, we are forced to

$$\rho = -\frac{c_{12}^2 + d_{12}^2}{2c_{12}d_{12}}, \quad d_{11} = \frac{c_{11}d_{12}}{c_{12}}, \quad C^1 = \begin{pmatrix} c_{11} & c_{12} \\ -\frac{c_{12}(c_{11}^2-d_{11}^2+2)}{c_{12}^2-d_{12}^2} & -c_{11} \end{pmatrix}, \quad C^2 = \begin{pmatrix} d_{11} & d_{12} \\ \frac{d_{12}(c_{11}^2-d_{11}^2+2)}{d_{12}^2-c_{12}^2} & -d_{11} \end{pmatrix}$$

and a reduced 5-parameter moduli of the  $\sigma_S$  entries as the solution of the covariance of  $\triangleright$  condition and both parts of (5.6). We also need

$$(5.7) \quad \bar{\rho} + \rho = 0, \quad \overline{\det C^i} = \epsilon' \det(\sigma_{Sj}^i C^j)$$

as further constraints independently of  $J$ .

**Lemma 5.2.** *For the type II Clifford algebra, the covariance of  $\triangleright$  condition, the condition (2.6) and that  $\rho$  is imaginary lead to one fewer parameter in  $\sigma_S$  (so the latter drops to 4-dimensional),  $\epsilon' = 1$  and uniquely determined  $\bar{C}^i$  in terms of  $C^i$  for given  $\epsilon$  for both types of  $J$ .*

*Proof.* Building on the analysis from the previous lemma, we impose (5.7) which fixes one of the coefficients of  $\sigma_S$  and one of the 3 entries of  $\bar{C}^i$  as

$$\frac{\bar{d}_{12}}{d_{12}} = -\frac{\bar{c}_{12}}{c_{12}}$$

We then solve for (2.6) to find the remaining entries of  $\bar{C}^i$ . For both types of  $J$ , we need  $\epsilon' = 1$  and for type (1)  $J$ , we find

$$\begin{aligned}\overline{c_{12}} &= \frac{(c_{12}^2 - d_{12}^2)(c_{12}z - c_{11}) - 2c_{12}^2}{c_{12}\epsilon(c_{12}^2 - d_{12}^2)}, \\ \overline{c_{11}} &= \frac{(c_{12}^2 - d_{11}^2)(c_{12}z - c_{11})(c_{12}(\epsilon - z\bar{z}) + c_{11}\bar{z}) - 2c_{12}^2\bar{z}}{c_{12}\epsilon(c_{12}^2 - d_{12}^2)},\end{aligned}$$

while for type (2)  $J$ , the case of  $\epsilon = 1$  is simpler with

$$-\frac{\overline{c_{12}}}{c_{12}} = \frac{\bar{z}}{z}, \quad \overline{c_{11}} - c_{11} = c_{12}\bar{z}.$$

These conditions are solved by writing  $z := |z|e^{i\theta}$ , then

$$(5.8) \quad d_{12} = te^{i\theta} = \frac{tz}{|z|}, \quad c_{12} = \imath se^{i\theta} = \frac{\imath sz}{|z|}, \quad c_{11} = x - \frac{\imath}{2}|z|s, \quad x, s, t \in \mathbb{R}.$$

The full  $C^i$  matrices are then determined according to the above as

$$(5.9) \quad C^1 = \begin{pmatrix} x - \frac{\imath s|z|}{2} & \frac{\imath sz}{|z|} \\ -\frac{\imath \bar{z}((s^2+t^2)(s|z|+2\imath x)^2-8s^2)}{4s|z|(s^2+t^2)} & \frac{\imath s|z|}{2} - x \end{pmatrix},$$

$$(5.10) \quad C^2 = \begin{pmatrix} -\frac{t(s|z|+2\imath x)}{2s} & \frac{tz}{|z|} \\ -\frac{t\bar{z}((s^2+t^2)(s|z|+2\imath x)^2-8s^2)}{4|z|s^2(s^2+t^2)} & \frac{1}{2}t\left(|z| + \frac{2\imath x}{s}\right) \end{pmatrix},$$

and we also had to fix

$$(5.11) \quad \rho = \frac{\imath}{2}\left(\frac{t}{s} - \frac{s}{t}\right).$$

□

This lemma gives us relations between the  $c_{ij}, d_{ij}$  and their conjugates, i.e. reality conditions in the  $C^i$ . It remains to solve the  $\bar{\sigma}_S$  identity in Lemma 2.2 for some  $J$ . The second identity in Lemma 2.2 can be seen as defining the conjugates of the entries of  $\sigma_S$ ,

$$(5.12) \quad \overline{\sigma_{Sj}^i} = J\sigma_S^{-1i}{}_j J^{-1},$$

where the inverse is as an  $M_2(\mathbb{C})$ -valued matrix with respect to the  $i, j$  indices. This gives us relations between the entries of  $\sigma_S$  and their conjugates. We focus on the simplest case  $J$  of type (2) with  $\epsilon = 1$  identified in the proof of the last lemma.

**Lemma 5.3.** *For  $J$  of type (2) with  $\epsilon = 1$  and the 3 real parameters  $s, t, x$  of  $C^i$  identified in (5.8), (5.12) imposes two further relations of  $\sigma_S$ , so this drops down to two parameters subject to a reality constraint. Solutions of this include a natural  $\sigma_S$  determined uniquely by the parameters of  $C^i$  and the complex parameter  $z$  of  $J$ .*

*Proof.* In principle, the relations for  $\overline{\sigma_S}$  entail in particular

$$\overline{\text{Tr}\sigma_{Sj}^i} = \text{Tr}(\sigma_S^{-1i}{}_j), \quad \overline{\det\sigma_{Sj}^i} = (\det\sigma_S)^{-1i}{}_j,$$

where we trace and det over the suppressed spinor matrix indices. These are independent of  $J$ , hence it may be helpful to solve for these first. In practice, for our example, it is cleaner to look at all the 12 equations for the constrained entries of  $\sigma_S$  (since there are four parameters left), take their complex conjugates and



compare with  $\overline{\sigma_S}$  as defined by the right hand side of (5.12), solving the simplest (and  $J$ -independent) ones first. The end result is to reduce our moduli space of  $\sigma_S$  entries to two dimensions (we add two more relations) and find that then (5.12) is solved. We choose these parameters as

$$\eta := (\sigma_{S1}^2)^1_1, \quad \zeta := (\sigma_{S1}^2)^1_2$$

and their reality condition is

$$\begin{aligned} \bar{\eta} &= \frac{d_{12} (4\zeta^2 c_{12}^2 + (c_{12}^2 - d_{12}^2) (d_{12}(\eta c_{12} - \zeta 2c_{11} - \zeta c_{12}\bar{z}) + 2(-\zeta c_{11} + \eta c_{12})^2))}{c_{12} (8\zeta^2 c_{12}^2 + (c_{12}^2 - d_{12}^2)(-2c_{11}\zeta + 2\eta c_{12} + d_{12})^2)}, \\ \bar{\zeta} &= \frac{\zeta d_{12}^2 (c_{12}^2 - d_{12}^2) \bar{z}}{z (8\zeta^2 c_{12}^2 + (c_{12}^2 - d_{12}^2)(-2c_{11}\zeta + 2\eta c_{12} + d_{12})^2)}. \end{aligned}$$

Given the solution (5.8) for the  $C^i$ , a natural solution of these reality conditions is

$$\eta = -\frac{(s^2 + t^2)tx|z|}{2((s^2 + t^2)x^2 + 2s^2)}, \quad \zeta = \frac{tx(s^2 + t^2)}{(s^2 + t^2)x^2 + 2s^2} e^{i\theta}.$$

This was obtained by making a polar expansion  $\eta = ue^{i\phi}, \zeta = ve^{i\psi}$  then asking for the imaginary part of  $\bar{\zeta}/(ve^{-i\psi})$  to vanish (where  $\bar{\zeta}$  is given by the right hand side of the expression above). This gives a simple equation for  $v$  with two solutions if  $s, t$  are generic. Taking the one stated (the other does not lead to a solution of the whole system), we ask for the imaginary part of  $\bar{\eta}/(ue^{-i\phi})$  to vanish with  $\bar{\eta}$  from the right hand side of the above. This gives a complicated equation for  $u$  which, however, greatly simplifies if we assume  $\psi = \theta$  and  $\phi = 0$ .

The entire  $\sigma_S$  for this simplest choice of  $\eta, \zeta$  is

$$(5.13) \quad \sigma_{S1}^1 = \begin{pmatrix} 1 - \frac{isx|z|(s^2+t^2)}{2((s^2+t^2)x^2+2s^2)} & \frac{izsx(s^2+t^2)}{|z|((s^2+t^2)x^2+2s^2)} \\ -\frac{i\bar{z}|z|sx(s^2+t^2)}{4((s^2+t^2)x^2+2s^2)} + \frac{\bar{z}}{2} & \frac{isx|z|(s^2+t^2)}{2((s^2+t^2)x^2+2s^2)} \end{pmatrix},$$

$$(5.14) \quad \sigma_{S2}^1 = \begin{pmatrix} \frac{s^2x|z|(s^2+t^2)}{-2t((s^2+t^2)x^2+2s^2)} & \frac{zs^2x(s^2+t^2)}{|z|t((s^2+t^2)x^2+2s^2)} \\ -\frac{\bar{z}|z|s^2x(s^2+t^2)}{4t((s^2+t^2)x^2+2s^2)} - \frac{i\bar{z}s}{2t} & \frac{s^2x|z|(s^2+t^2)}{2t((s^2+t^2)x^2+2s^2)} + \frac{is}{t} \end{pmatrix},$$

$$(5.15) \quad \sigma_{S1}^2 = \begin{pmatrix} \frac{tx|z|(s^2+t^2)}{-2((s^2+t^2)x^2+2s^2)} & \frac{zt x(s^2+t^2)}{|z|(s^2+t^2)x^2+2s^2} \\ -\frac{t\bar{z}|z|x(s^2+t^2)}{4((s^2+t^2)x^2+2s^2)} - \frac{i\bar{z}t}{2s} & \frac{tx|z|(s^2+t^2)}{2((s^2+t^2)x^2+2s^2)} + \frac{it}{s} \end{pmatrix},$$

$$(5.16) \quad \sigma_{S2}^2 = \begin{pmatrix} -1 + \frac{isx|z|(s^2+t^2)}{2((s^2+t^2)x^2+2s^2)} & -\frac{izsx(s^2+t^2)}{|z|((s^2+t^2)x^2+2s^2)} \\ \frac{i\bar{z}|z|sx(s^2+t^2)}{4((s^2+t^2)x^2+2s^2)} - \frac{\bar{z}}{2} & -\frac{isx|z|(s^2+t^2)}{2((s^2+t^2)x^2+2s^2)} \end{pmatrix}.$$

□

**5.3. Construction of non-hermitian almost spectral triples on  $M_2(\mathbb{C})$ .** We next consider the Hilbert space structure, taking the obvious choice

$$(5.17) \quad \langle \phi_\alpha e^\alpha, \psi_\beta e^\beta \rangle = \frac{1}{2} \text{Tr}(\phi_\alpha^* \psi_\alpha),$$

corresponding to  $\int = \frac{1}{2} \text{Tr}$  and  $\mu^{\alpha\beta} = \delta^{\alpha\beta}$  in (2.14). Here, the spinor coefficients  $\phi_\alpha, \psi_\alpha \in M_2(\mathbb{C})$  with its usual (hermitian adjoint)  $*$ -operation.

**Theorem 5.4.** *On  $M_2(\mathbb{C})$  with its alternate 1-parameter quantum Riemannian geometry as in Section 5.1, the ‘local’ conditions for an even spectral triple have a natural geometric realisation with 4 real parameters  $s, t, x, y$ ,  $\epsilon = \epsilon' = \epsilon'' = 1$ ,*

$$J = \begin{pmatrix} 1 & 0 \\ 0 & e^{2iy} \end{pmatrix}, \quad \gamma = \pm \begin{pmatrix} 1 & 2 \frac{e^{iy} sx(s^2+t^2)}{(s^2+t^2)x^2+2s^2} \\ 0 & -1 \end{pmatrix}$$

*such that  $\mathcal{J}$  is an antilinear isometry with respect to (5.17). Moreover, the QLC has its parameter  $\rho$  fixed as (5.11) with curvature proportional to*

$$1 + \rho^2 = \frac{1}{4} \left( 6 - \frac{s^2}{t^2} - \frac{t^2}{s^2} \right).$$

*Proof.* We first bring together our lemmas above. We fix  $J$  of type (2) with  $\epsilon = 1$  as this gave the most natural solutions for  $C^i$ . Here  $z$  is a free parameter in  $J$  in case (2) of Lemma 2.1. The three parameters  $s, t, x$  go into solving for  $C^i$  to obey the 2nd half of (5.4), covariance of  $\triangleright$  and (2.6), and also determine  $\rho$ , i.e. the choice of QLC which was otherwise not fixed for the given metric. Existence of the  $C^i$  also cuts down the allowed  $\nabla_S$  to 4 parameters and among solutions we found a natural choice of these so as to obey the reality condition in Lemma 2.2 for  $\nabla_S$  to be  $\ast$ -preserving.

Next, we take the natural choice of Hilbert space structure on  $\mathcal{S} = M_2(\mathbb{C}) \otimes \mathbb{C}^2$  as stated. As  $\epsilon = 1$ , we then need  $J$  to be symmetric. We achieve this by taking a limiting value  $z \rightarrow 0$  along a ray such that  $-z/\bar{z} = e^{2iy}$  for some real  $y$ . Our solutions for  $C^i, \sigma_{Sj}^i$  then simplify considerably to

$$(5.18) \quad C^1 = \begin{pmatrix} x & se^{iy} \\ -\frac{(s^2+t^2)x^2+2s^2}{e^{iy}s(s^2+t^2)} & -x \end{pmatrix}, \quad C^2 = -\frac{it}{s} C^1,$$

$$(5.19) \quad \sigma_{S1}^1 = \begin{pmatrix} 1 & \frac{e^{iy}sx(s^2+t^2)}{(s^2+t^2)x^2+2s^2} \\ 0 & 0 \end{pmatrix} = -\sigma_{S2}^2, \quad \sigma_{S2}^1 = \frac{is}{t} \begin{pmatrix} 0 & -\frac{e^{iy}sx(s^2+t^2)}{(s^2+t^2)x^2+2s^2} \\ 0 & 1 \end{pmatrix} = \frac{s^2}{t^2} \sigma_{S1}^2.$$

For the last part, we first solve (2.7), (2.8) and (2.19) to find the unique answer for  $\gamma$  up to sign (which is not determined by the equations). Then (2.9) is solved automatically with  $\epsilon'' = 1$ .  $\square$

Note that we can set  $y = 0$  by conjugation by a unitary  $\text{diag}(1, e^{iy})$ . We also see that there are four points  $t = \pm(\sqrt{2} \pm 1)s$  where the QRG is flat. It remains to see if  $\mathcal{D}$  and  $\gamma$  constructed above are (anti)hermitian.

**Proposition 5.5.** *The geometrically realised Dirac operator in Theorem 5.4 has the form*

$$(\mathcal{D}\psi)_\gamma = \frac{i}{2}([\sigma^1, \psi_\alpha]C^{1\alpha}_\gamma + \{\sigma^2, \psi_\alpha\}C^{2\alpha}_\gamma)$$

*which we write as*

$$\mathcal{D} = \phi^{(1)} - \frac{it}{s}\phi^{(2)}, \quad (\phi^{(1)}\psi)_\gamma = \frac{i}{2}[\sigma^1, \psi_\alpha]C^{1\alpha}_\gamma, \quad (\phi^{(2)}\psi)_\gamma = \frac{i}{2}\{\sigma^2, \psi_\alpha\}C^{1\alpha}_\gamma.$$

*If  $x = \pm s\sqrt{1 - \frac{2}{s^2+t^2}}$  then  $\phi^{(i)}$  are both hermitian with respect to (5.17), while  $\gamma$  is hermitian if  $x = 0$ . All three are hermitian if  $s^2 + t^2 = 2$  and  $x = 0$ .*

*Proof.* We again use (4.9) given the inner form of the connection and the form of  $\theta$  for this calculus. This time, however, for our particular 4-parameter solutions, we have

$$\sigma_{S^i}^j C^i = \pm C^j$$

with  $+1$  if  $j = 1$  and  $-1$  if  $j = 2$ , which gives the result stated. Next, if  $C^1$  is *antihermitian*, which happens for  $x$  with the value stated, then

$$\langle \not{D}^{(1)} \psi, \phi \rangle - \langle \psi, \not{D}^{(1)} \phi \rangle = \frac{i C^{1\gamma} \alpha}{4} \text{Tr} \left( [\sigma^1, \psi_\alpha]^\dagger \phi_\gamma - \psi_\alpha^\dagger [\sigma^1, \phi_\gamma] \right) = 0,$$

$$\langle \not{D}^{(2)} \psi, \phi \rangle - \langle \psi, \not{D}^{(2)} \phi \rangle = \frac{i C^{1\gamma} \alpha}{4} \text{Tr} \left( \{\sigma^2, \psi_\alpha\}^\dagger \phi_\gamma - \psi_\alpha^\dagger \{\sigma^2, \phi_\gamma\} \right) = 0,$$

using the trace property.

For  $\gamma$  to be hermitian with respect to (5.17) amount to the matrix for  $\gamma$  hermitian, which is clearly not the case unless  $xs = 0$ . Our solutions suppose  $s \neq 0$  so the condition is  $x = 0$  and then  $\gamma = \pm \sigma^3$ . This is compatible with our condition for  $\not{D}^{(i)}$  when  $s^2 + t^2 = 2$ .  $\square$

We see that we see that our 4-parameter space of solutions in Theorem 5.4 reduces to a circle  $s^2 + t^2 = 2$  (with  $s, t \neq 0$ ) if we require  $\not{D}^{(i)}, \gamma$  hermitian. Choosing the  $+$  sign for square roots and setting  $y = 0$  by the freedom under unitary equivalence, we have in this case

$$(5.20) \quad C^1 = i s \sigma^2, \quad C^2 = t \sigma^2, \quad J = \text{id},$$

$$(5.21) \quad \sigma_{S^1}^1 = E_{11} = -\sigma_{S^2}^2, \quad \sigma_{S^2}^1 = \frac{is}{t} E_{22}, \quad \sigma_{S^1}^2 = \frac{it}{s} E_{22}, \quad \gamma = \sigma^3.$$

In fact, the pattern of signs would fit into the period 8 KO-dimension signs with  $n = 0$  if  $\not{D}$  were to be hermitian and would fit  $n = 1$  if  $\not{D}$  were to be antihermitian (then we would flip the value of  $\epsilon'$ ), but our  $\not{D}$  is neither: the first part of  $\not{D}$  is hermitian and the second part is antihermitian. Similary,  $C^1$  is antihermitian and  $C^2$  is hermitian in line with the Minkowski nature of the metric (5.1) on  $M_2(\mathbb{C})$ .

We do not claim that Theorem 5.4 is the only data for a geometrically realised spectral triples on  $M_2(\mathbb{C})$  for this quantum metric, but our results were obtained by solving the equations with the simplest nontrivial choices at the various branches as per the derivation above. Also note that by Lemma 2.2, we are free (for both QRGs on  $M_2(\mathbb{C})$ ) to add a bimodule map in the form of matrices  $A_i$  to the connection  $\nabla_S$  subject to the conditions in the lemma. This will appear as further freedom in the construction of a geometrically realised Dirac operator as fluctuations of the inner case that we solved for. On the other hand, it does not appear that looking more widely will solve the problem that  $\not{D}$  is not antihermitian for the quantum metric in this section.

## 6. CONCLUDING REMARKS

We have set up the formalism for geometric realisation of spectral triples with central bases for  $\Omega^1$  and the spinor bundle  $\mathcal{S}$  as in [20] but now extended to allow the Christoffel symbols to be nonconstant and the basic 1-forms not form a Grassmann algebra. We demonstrated how the formalism leads uniquely (up to unitary

equivalence) to the expected  $n = 2$  spectral triple on the noncommutative torus, just as we did for the fuzzy sphere in [20], as well as  $n = 1$  for another choice of signs. Interestingly, we saw in Section 3.3 that the spinor connection  $\nabla_S$  and the connection  $\nabla$  on  $\Omega^1$  need not be trivial and can be used to encode a dynamically generated mass term or chiral term for the Dirac operator. It would be interesting to see how our results extend to a noncommutative  $d$ -torus, where it is known that there are  $2^d$  spin structures matching their classical counterparts[29]. Moreover, the noncommutative torus in a functional analysis context admits genuinely projective modules[8], and some of these could potentially lead to more sophisticated examples using the geometric formalism of [5].

Looking beyond standard noncommutative torus, we note that the  $q$ -deformed noncommutative torus  $\mathbb{C}_{q,\theta}[\mathbb{T}^2]$  is the same algebra as before but now with a  $q$ -deformed differential structure[2],

$$(du)u = qudu, \quad (dv)v = qvdv, \quad (dv)u = e^{i\theta}udv, \quad (du)v = e^{-i\theta}vdu$$

with  $q$  real for a  $\ast$ -calculus. One might ask if one can extend the analysis we have done to construct spectral triples in this case. The problem here is the lack of a central self-adjoint basis, which makes calculations harder. On the other hand, the  $q$ -deformed case is known to be a comodule algebra cocycle twist of the standard noncommutative torus[2, Exercise 9.7] and hence can be constructed as a twist of the standard one above. Here, the quantum Riemannian geometry itself can be twisted using [4].

Alternatively, we can set  $q = e^{i\theta}$  and then  $s^1 = v^{-1}u^{-1}du$  and  $s^1 = uv^{-1}dv$  are a central basis [2, Exercise 1.5] but, because  $q$  is not real, this case is not a  $\ast$ -calculus so we leave the existing framework[5] for geometric realisation of spectral triples. However, this case is nevertheless interesting and if  $q = e^{i\theta}$  is a primitive  $n$ -th root of unity then one can define the reduced noncommutative torus as

$$c_q[\mathbb{T}^2] = \mathbb{C}_{q,\theta}[\mathbb{T}^2]/\langle u^n - 1, v^n - 1 \rangle,$$

which is compatible with the differential calculus [2, Exercise 1.5], just not as a  $\ast$ -calculus. Just as  $c_q[SL_2]$  are not usual Hopf  $\ast$ -algebras but ‘flip’ ones, one could expect a theory for the reduced noncommutative torus that covers this non-standard case and possibly geometrically realises the spectral triples on fuzzy tori in [3]. This is another interesting direction for further work.

Secondly, we demonstrated the formalism on  $M_2(\mathbb{C})$  with its standard 2D calculus and the two choices of quantum metric for which the QRG is known. The moduli of geometric Dirac operators at the local level turned out to be very rich and for one quantum metric the Hilbert space conditions led naturally to a pair of spectral triples depending on  $\epsilon'$ . For the other metric, as one might expect from its Lorentzian form, the most reasonable local Dirac operator was not antihermitian so we do not have exactly a spectral triple. Indeed, one might expect a Lorentzian spectral triple [14, 27], although our example does not appear to immediately fit into one of these frameworks either. This is another topic for further work.

Quantum Riemannian geometries have been used for baby quantum gravity models, see e.g. [19, 1] and the inclusion of a Dirac operator should allow the coupling of such models to fermionic matter, which remains to be investigated. There are also new tools such as quantum geodesics[6, 21] recently computed on the noncommutative

torus and  $M_2(\mathbb{C})$ . Another reason to be interested in a full classification of geometrically realised spectral triples on finite-dimensional algebras such as  $M_2(\mathbb{C})$  and  $c_q[\mathbb{T}^2]$  at roots of unity is the potential for applications to the structure of elementary particles by tensoring spacetime with such a finite quantum geometry[9, 11]. In the current state of the art, while one can choose the algebra, its representation and  $\not{D}$  to reproduce the standard model and obtain some predictions, having further (geometric realisation) constraints on the allowed Dirac operators would translate directly into further constraints on the particle physics. One can also come at the problem in a Kaluza-Klein manner without the Dirac operator[1] and then extend this to spinors and spectral triples. This will be looked at elsewhere.

## REFERENCES

- [1] J. Argota-Quiroz and S. Majid, Quantum gravity on finite spacetimes and dynamical mass, PoS (2022) 210 (41pp)
- [2] E.J. Beggs and S. Majid, *Quantum Riemannian Geometry*, Grundlehren der mathematischen Wissenschaften, Vol. 355, Springer (2020) 809pp
- [3] J.W. Barrett and J. Gaunt, Finite spectral triples for the fuzzy torus, arXiv:1908.06796 [math.QA]
- [4] E.J. Beggs and S. Majid, Nonassociative Riemannian geometry by twisting, J. Phys. Conf. Ser. 254 (2010) 012002 (29pp)
- [5] E.J. Beggs and S. Majid, Spectral triples from bimodule connections and Chern connections, J. Noncomm. Geom., 11 (2017) 669–701
- [6] E. Beggs and S. Majid, Quantum geodesics and curvature, arXiv: 2201.08244 (math.QA)
- [7] A.L. Carey, J. Phillips and A. Rennie, Spectral triples: examples and index theory, in *Noncommutative Geometry and Physics: Renormalisation, Motives, Index Theory* (European Mathematical Society: Zurich) 2011, pp175–265
- [8] A. Connes,  $C^*$ -algèbres et géométrie différentielle, C.R. Acad. Sci. Paris, Ser. A-B , 290, (1980)
- [9] A. Connes, Gravity coupled with matter and foundation of noncommutative geometry, Commun. Math. Phys. 182 (1996) 155
- [10] A. Connes, Noncommutative Geometry, Academic Press, Inc., San Diego, CA, 1994
- [11] A. Connes and M. Marcolli, *Noncommutative Geometry, Quantum Fields and Motives* (AMS Colloquium Publications Vol 55), Hindustan Book Agency, 2008.
- [12] L. Dabrowski and A. Sitarz, Fermion masses, mass-mixing and the almost commutative geometry of the Standard Model, JHEP (2019) 68
- [13] F. D’Andrea, F. Lizzi and J. C. Varilly, Metric properties of the fuzzy sphere, Lett. Math. Phys. 103 (2013) 183–205
- [14] A. Devastato, S. Farnsworth, F. Lizzi and P. Martinetti, Lorentz signature and twisted spectral triples, JHEP (2018) 89
- [15] S. Doplicher, K. Fredenhagen and J. E. Roberts, The quantum structure of spacetime at the Planck scale and quantum fields, Commun. Math. Phys. 172 (1995) 187–220
- [16] M. Dubois-Violette and P.W. Michor, Connections on central bimodules in noncommutative differential geometry, J. Geom. Phys. 20 (1996) 218–232
- [17] J. M. Gracia-Bondía, J.C. Varilly, H. Figueroa, *Elements of Noncommutative Geometry*, Birkhäuser, Boston, 2001
- [18] G. ’t Hooft, Quantization of point particles in 2+1 dimensional gravity and space- time discreteness, Class. Quant. Grav. 13 (1996) 1023
- [19] E. Lira-Torres and S. Majid, Quantum gravity and Riemannian geometry on the fuzzy sphere, Lett. Math. Phys. (2021) 111:29 (21pp)
- [20] E. Lira-Torres and S. Majid, Geometric Dirac operator on the fuzzy sphere, Lett. Math. Phys. (2022) 112:10
- [21] C. Liu and S. Majid, Quantum geodesics on quantum Minkowski spacetime, J. Phys. A 55 (2022) 424003 (35pp)
- [22] S. Majid, Hopf algebras for physics at the Planck scale, Class. Quant. Grav. 5 (1988) 1587–1607

- [23] S. Majid and H. Ruegg, Bicrossproduct structure of the  $\kappa$ -Poincare group and non-commutative geometry, *Phys. Lett. B.* 334 (1994) 348–354
- [24] S. Majid and F. Simao, Quantum jet bundles, *arXiv: 2202.03067 (math.QA)*
- [25] J. Mourad, Linear connections in noncommutative geometry, *Class. Quant. Grav.* 12 (1995) 965–974
- [26] M. Paschke and A. Sitarz, On spin structures and Dirac operators on the noncommutative torus, *Lett. Math. Phys.* 77 (2006), 317–327
- [27] M. Paschke and A. Sitarz, Equivariant Lorentzian spectral triples, *arXiv:math-ph/0611029*
- [28] H.S. Snyder, Quantized space-time, *Phys. Rev.* 71 (1947) 38–41
- [29] J.J. Venselaar, Classification of spin structures on the noncommutative n-torus, *J. Non-comm. Geom.* 7 (2013), 787–816

QUEEN MARY, UNIVERSITY OF LONDON, SCHOOL OF MATHEMATICS, MILE END RD, LONDON E1 4NS, UK

*Email address:* `s.majid@qmul.ac.uk`