

User Experience of Symptom Checkers: A Systematic Review

Yue You, MS¹, Renkai Ma, MS¹, Xinning Gui, PhD¹
¹Pennsylvania State University, University Park, PA, USA

Abstract

This review reports the user experience of symptom checkers, aiming to characterize users studied in the existing literature, identify the aspects of user experience of symptom checkers that have been studied, and offer design suggestions. Our literature search resulted in 31 publications. We found that (1) most symptom checker users are relatively young; (2) eight relevant aspects of user experience have been explored, including motivation, trust, acceptability, satisfaction, accuracy, usability, safety/security, and functionality; (3) future symptom checkers should improve their accuracy, safety, and usability. Although many facets of user experience have been explored, methodological challenges exist and some important aspects of user experience remain understudied. Further research should be conducted to explore users' needs and the context of use. More qualitative and mixed-method studies are needed to understand actual users' experiences in the future.

Introduction

The rapid development of digital technology and the healthcare industry has contributed to the increased popularity of symptom checkers (SCs) in application (app) markets. Some SC apps (e.g., Ada) have been downloaded from app stores tens of millions of times¹. These SCs utilize either a chatbot or a questionnaire-like form to solicit symptom input and provide users with a potential diagnosis. Some SCs describe themselves as akin to intelligent doctors who give accurate and reliable medical recommendations. For instance, Ada claims to “*think like a doctor*”². However, the reliability of SCs is questionable³. Healthcare consumers may put their health at risk if they uncritically accept the diagnoses from SCs⁴. Recognizing that individuals' perceptions and interpretations of SCs' responses may influence their subsequent actions, it is important to understand more fully the user experience (UX) of SCs⁵. UX is “a consequence of a user's internal state, the characteristics of the designed system, and the context (or the environment) within which the interaction occurs.”⁶ As Table 1 indicates, we adapted this definition of UX to characterize the UX of SCs along three dimensions. As the UX of SCs is highly related to the acceptance of SCs⁷ and the minimization of unintended consequences (e.g., anxiety and cyberchondria)⁸, it is crucial to improve the UX of SCs by understanding individuals' motivations for using SCs, users' expectations of SCs, the benefits as well as the limitations of SCs' features, and the usage contexts.

Table 1. Three dimensions of the UX of SCs.

Dimension	Definition
Consequence of a user's internal state	User's expectations of, needs for, and attitude towards SCs
Characteristics of the designed system	SCs' features, including usability, complexity, purpose, and functionality
Context or environment	Context or environment within which users interact with SCs

Nevertheless, few existing papers have systematically reviewed studies of the UX of SCs. Instead, previous systematic reviews of SCs have focused on the influence of SCs' use on the patient–doctor relationship⁹, the current evaluation methodologies for SCs¹, and the impact of SCs on urgent care seeking¹⁰ and primary care¹¹. To the best of our best knowledge, only one scoping review of SCs covered studies addressing the UX of SCs¹². This scoping review provided a broad view of research on SCs and points out that UX is one theme that has been explored. However, it did not offer an in-depth analysis of the aspects of the UX of SCs that have been explored. It is still unclear who SC users are, what factors motivate and influence the use of SCs, and what relevant design suggestions might be. To address this gap in existing reviews, we systematically reviewed papers that examine the UX of SCs. We choose the systematic review approach instead of doing a scoping review because a systematic review can allow us to synthesize the findings from previous literature and identify areas for future research¹³. We intended to address the following three questions: (1) Who uses SCs? (2) Which aspects of the UX of SCs have been studied based on the three dimensions of UX? and (3) What design suggestions have been proposed to improve the UX of SCs?

Materials and Methods

Search strategy and screening process. We developed a study protocol in compliance with the PRISMA guidelines. The PROSPERO website was searched and no previous systematic review on similar topics was found, which validated the novelty of our review. We searched six electronic databases from May 2021 to July 2021: PubMed,

CINAHL, ACM Digital Library, Scopus, Web of Science, and Google Scholar. We also conducted backward–forward searches by examining the references from relevant articles and systematic reviews. We utilized three groups of search terms: SC-related terms, internet technology-related terms, and health-related terms. The following search query was used for Scopus: (TITLE-ABS-KEY(“symptom checker” OR “self diagnosis” OR “self-diagnosis” OR “self triage” OR “self-assessment” OR “self assessment” OR “self diagnosing” OR “self-evaluation” OR “self evaluation” OR “self-appraisal” OR “check your symptom” OR “check their symptom” OR “online diagnosis” OR “web-based triage” OR “electronic triage” OR “etriage” OR “self referral” OR “assistance technology” OR “assistance technologies”) AND TITLE-ABS-KEY(“online” OR “technology” OR “website” OR “tool” OR “computer” OR “mhealth” OR “m-health” OR “ehealth” OR “ e-health” OR “app” OR “mobile application” OR “smartphone” OR “smart phone” OR “cell phone” OR “cellular phone” OR “mobile phone” OR “electronic” OR “automated” OR “internet” OR “digital” OR “mobile”) AND TITLE-ABS-KEY(“medical” OR “health” OR “medication” OR “healthcare”). The search strings for Scopus were adapted for other databases.

After completing the database and backward–forward searches, we conducted two rounds of screening to determine which articles to include in our review. First, we screened the titles and abstracts of the identified papers using three eligibility criteria: (1) the paper is an original research article published in a peer-reviewed journal or conference proceedings; (2) the paper was published in English; and (3) the paper’s title and abstract included our search terms. We then screened the full texts of those papers based on the following inclusion criteria: (1) the subject of the paper is an application rather than a model or algorithm or general online search; (2) consumers/laypersons are the primary users of the technology developed and/or evaluated in the paper; (3) the technology described in the paper is internet-enabled; (4) the technology studied in the paper is used for self-diagnosis; (5) the topic of the paper addresses an aspect of the UX of SCs; and (6) the full text is available. After each round of screening, three researchers performed a cross-review and settled any disagreements. Only those papers that the researchers deemed sufficiently relevant were included. In addition, quality assessment was conducted for all papers to assess whether these papers’ aims and methodology were clearly described, whether their findings align with their research questions, and whether they were peer-reviewed. All discrepancies were discussed and resolved. The quality assessment results were taken into account when synthesizing data. We finally identified 16,825 potentially relevant articles written in the English language. We also identified 3,425 papers using Google Scholar and backward–forward searching. Ultimately, 31 papers (24 from databases and 7 from other methods) were chosen to be included. Figure 1 shows the screening process.




Figure 1. Processes of identification and screening.

Data collection and analysis. Using the papers identified during the two rounds of screening, we conducted data extraction based on the PRISMA guidelines. We collected the following main information: (1) the source of the paper, including the author(s), date of publication, and publication venue; (2) background information, including the diseases examined in the paper, the research purposes, stakeholders, and the features/functions of SCs; (3) methodological information, including each author’s country and affiliation, the study setting and duration, the participants’ characteristics, the data collection and analysis methods; and (4) findings, including the participants’ motivations for using SCs, the participants’ characteristics, the dimensions/measurements of UX, and design suggestions. We then used thematic analysis and descriptive statistical analysis to synthesize the extracted data.

Results

Study characteristics. The 31 papers chosen for inclusion were published between 2007 and 2021 (see Table 2).

Table 2. List of studies included in the systematic review.

Source	Country	Study Design	Methodology	Diseases	Participants	System Features
Powley et al, 2016 ¹⁴	UK	Laboratories study	Quantitative; Experimental study	Inflammatory joint disease	N = 34; Age: 29.75-78.5; Female	Existing apps; Questionnaire-based
Bisson et al, 2016 ¹⁵	US	Laboratories study	Quantitative; Experimental study	Knee pain	N = 328; Age: 18-76; 49.7% male	Prototype; Questionnaire-based
Bisson et al, 2014 ¹⁶	US	Field study	Quantitative; Experimental study	Knee pain	N = 527; Age: avg: 47, 18-84; 51.6% male	Prototype
Hageman et al, 2015 ¹⁷	US	Laboratories study	Quantitative; Experimental study & questionnaire	Hand diseases	N = 86 patients; Avg age: 46; 51% male	Existing apps; Questionnaire-based
Knitzta et al, 2021 ¹⁸	Germany	Laboratories study	Quantitative; Experimental study	Inflammatory rheumatic disease	N = 164 patients	Existing apps; Chatbot & questionnaire-based; AI-driven & rule-based
Cross et al, 2021 ¹⁹	Australia	Laboratories study	Quantitative; Experimental study & questionnaire & log-based study	N/A	N = 64; Age: > 18, 87.49%: 18-34; 4.68%: 35-44; 1.56%: 45-54	A clone of existing app; Questionnaire-based
Verzantvoort et al, 2018 ²⁰	the Netherlands	Field study	Mixed methods; Experimental study & interview	Acute primary care	N = 4,456; Age: most 19-45 (56.4%)	Existing app
Fan et al, 2021 ²¹	China	Field study	Mixed methods; Log-based study	N/A	N = 16,519; Age: most 20-39; 54.8% male	Existing app; Chatbot; AI-driven
Price et al, 2013 ²²	US	Laboratories study	Quantitative; Experimental study & questionnaire	Influenza-Like Illness	N = 294 parents and caregivers; Age: > 18	Prototype; Questionnaire-based
You & Gui, 2020 ²³	US	Field study	Qualitative; App review analysis & interview	N/A	N = 10 (interview); Age: 23-40 (interview)	Existing apps; Chatbot
Aboueid et al, 2021 ⁷	Canada	Laboratories study	Qualitative; Think aloud & interview	N/A	N = 24; Age: 18-34	Existing apps; Questionnaire-based & chatbot
Aboueid et al, 2021 ²⁴	Canada	Laboratories study	Qualitative; Interview	COVID-19	N = 22; Age: 18-33; 54.5% female	Existing apps; Questionnaire-based & chatbot
Marco-Ruiz et al, 2017 ²⁵	Norway	Laboratories studies	Mixed methods; Experimental study & think aloud	Respiratory diseases	Phase 1: N = 53; Phase 2: N = 15	Existing apps
Ponnada, 2020 ²⁶	US	Laboratories study	Mixed methods; Usability testing & interview	COVID-19	N/A	Existing app & improved prototype; Chatbot
Schrager et al, 2020 ²⁷	US	Laboratories study	Quantitative; Prototyping & usability testing	COVID-19	N = 877; Age: avg: 32, 19-84; 65.3% female	Prototype; Questionnaire-based
Reilly & Austin, 2021 ²⁸	Australia	Field study	Quantitative; Online survey	Depression in pregnancy	N = 140; Age: N/A; Female	Existing app; Questionnaire-based
Nieroda et al, 2018 ²⁹	UK	Laboratories study	Qualitative; Think aloud & online survey	Cancer	The public: N=39, Age: >= 40; Practitioners: N = 20	Existing app
Daher et al, 2020 ³⁰	Switzerland	Laboratories study	Quantitative; Experimental study	N/A	N = 12; Age: N/A	Prototype; Chatbot
Tsai et al, 2021 ³¹	US	Field study & laboratories study	Mixed methods; Interview & experimental study	General/ COVID-19	(1) N = 25; Age: 19-54;(2) N = 20; Age: 19-37; 70% female	Existing apps & prototype; Chatbot & questionnaire-based
Li et al, 2020 ³²	China	Field study & laboratories study	Mixed methods; Interview & experimental study	N/A	(1) N = 13; Age: 21-61 (2) N = 48	Existing apps; Chatbot; AI-driven
You et al, 2021 ³³	China	Field study	Qualitative; Interview	N/A	N = 30; Age: 20-55 Most were in their 20s and 30s	Existing apps; Chatbot & questionnaire-based; AI-driven

Table 2. List of studies included in the systematic review (continued).

Source	Country	Study Design	Methodology	Diseases	Participants	System Features
Hwang et al, 2018 ³⁴	US, South Korea	Laboratories study	Qualitative; Think aloud	N/A	N = 6; Age: 20-39; 4 female, 2 male	Existing apps; Chatbot; AI-driven
Baldauf et al, 2020 ³⁵	Switzerland	Questionnaires	Mixed methods; Online survey	N/A	N = 106; Age: avg: 34, 19-60; 58.5% male	Existing apps; Using sensor data; AI-driven
Miller et al, 2020 ³⁶	UK	Laboratories study	Quantitative; Experimental study & questionnaire	N/A	N = 523; Age: avg: 39.79, 81.6%: 15-64; 62.1% female	Existing app; Chatbot; AI-driven
Meyer et al, 2020 ³⁷	US	Questionnaires	Mixed methods; Online survey	N/A	N = 329; Age: avg: 48.0; 75.7% female	Existing app; questionnaire, AI-based
Winn et al, 2019 ³⁸	US	Field study	Quantitative; Log analysis	N/A	N = 158,083; Age: avg: 40; 78% female	Existing app; Chatbot
DeForte et al, 2020 ³⁹	US	Field study	Qualitative; App review analysis	Depression	N/A	Existing apps
Nijland et al, 2010 ⁴⁰	the Netherlands	Field study & questionnaires	Quantitative; Log-based & survey	N/A	N = 192; Age: 16–35 (56%); 66% female	Existing app; Questionnaire-based
Luger et al, 2014 ⁴¹	US	Laboratories study	Qualitative; Think aloud & questionnaire	N/A	N = 79; Age: >= 50;	Existing app; Questionnaire-based
Lanseng et al, 2007 ⁴²	Norway	Questionnaires	Quantitative; Online survey	N/A	N = 160; Age: 18-65; 46.3% female	N/A
Hua & Hou, 2020 ⁴³	Vietnam	Field study & questionnaires	Mixed methods; Interview & survey	N/A	N = 482; Age: 18-50; 63.1% female	N/A

Characteristics of users. Four studies that we reviewed (12.9%)^{21,21,37,38} reported on the characteristics of individuals who had used SCs prior to participating in the studies. The age of SC users was found to be relatively young (most users ranged in age from 20 to 39 years old²¹ or the average age was 40 years old³⁸). Regarding the gender of SC users, the study conducted in China found male users initiated most consultations (54.80%)²¹, while the U.S. study reported most users were female (75.7%)³⁷. Other less commonly mentioned user characteristics in our corpus included health literacy⁷, education³⁷, and income³⁷. Young individuals with low health literacy and high technology literacy were found to be inclined to use SCs⁷. The household income of most users was less than \$100,000 USD³⁷.

UX aspects of SCs.

(1) Consequence of a consumer’s internal state

Motivations for using SCs: Five studies (16.1%)^{21,28,33,38,43} addressed users’ motivations for using SCs (see Table 3).

Table 3. Motivations for using SCs (n = 5).

Motivation	Details
Convenience	The researchers found that individuals were motivated to use SCs instead of going to a doctor because of the time and cost savings as well as location independence ³³
Supplementing offline medical visits	Individuals were motivated to use SCs to supplement in-person medical visits and gain assistance in making healthcare-related decisions ^{28,33,38}
Gaming	Individuals (8.03%) gamed SC by entering nonsensical words for nontherapeutic purposes ^{21,43}
Social influence	Social influence (i.e., “the extent to which an individual is aware that significant others recommend they use the self-diagnostic app”) could prompt consumers to use SCs ⁴³

Trust: Six of the reviewed papers (19.4%) explored users’ trust in SCs^{21,31–35}, reporting factors that may influence users’ trust, including the provision of explanations, interaction design, and accuracy (see Table 4).

Table 4. Factors that influence the users’ trust (n = 6).

Factor	Impact	Methods
Explanations	The information source or authority ^{31,33} , medical certification ³⁵ , security of data transmission ³⁵ , and explanations for presented questions and reports ^{31,34} could positively influence users’ trust	Interviews ³³ , think aloud ³⁴ , questionnaires (adapted from previous literature ³¹ , self-created ³⁵)
Interaction design	Repeated or irrelevant questions ³⁴ , vague options ³⁴ , limited input functions ³³ , and an overly anthropomorphic conversation design ³³ could negatively affect users’ trust in SCs	Interviews ³³ , think aloud ³⁴
Accuracy	Users tended to distrust SCs when the SCs’ recommendations were inaccurate ^{21,32,33}	Interviews ³³ , log analysis ²¹ , questionnaires (self-created) ³²

Acceptability: In our corpus, a total of four papers (12.9%) measured whether users accepted SCs^{18,28,29,36}. These studies relied on questionnaires^{18,28,29,36} and the thinking-aloud method²⁹. All of the papers described how most users accepted the use of SCs to support emotional wellness²⁸, primary care³⁶, and cancer diagnosis²⁹.

Satisfaction: Four of the studies (12.9%) evaluated individuals' satisfaction with using SCs via questionnaires^{19-21,31}. These studies found that some of the relevant factors affecting satisfaction are disease type²¹, accuracy²¹, and explanations³¹. One log-based study found that for common diseases, users were usually satisfied with the diagnostic recommendations, while for more complex diseases, users were less satisfied²¹. This study also reported that inaccurate diagnostic results contributed to dissatisfaction with SCs. One experimental study reported that providing different styles of explanations can increase users' satisfaction with SCs³¹.

(2) Characteristics of the SCs

Accuracy: Eleven papers included in our corpus examined the accuracy of SCs. Three papers stated that compared with physicians' diagnoses, recommendations from SCs were relatively accurate in diagnosing knee pain (correctly diagnose 58% of the time)¹⁵, acute primary care (correctly triage 89% of patients)²⁰, and influenza-like illness (correctly diagnose 93% of the time)²², though two of them admitted that SCs had low specificity^{16,22}. However, the other eight papers revealed that SCs are frequently inaccurate when providing diagnoses for inflammatory joint diseases¹⁴, hand diseases¹⁷, inflammatory rheumatic diseases¹⁸, knee pain¹⁵, and general diseases (e.g., 1,084 out of 3,832 diagnoses were rated by users as inaccurate²¹)^{19,21,37,41}.

Usability: Ten papers explored the usability of SCs. Three experiment-based papers reported that SCs were easy to use^{18,22,36} (e.g., 97.8% of the participants considered Ada easy to use³⁶) by using the standardized scales (e.g., System Usability Scale)^{18,36} or self-created web survey²². The remaining seven studies reported usability issues (see Table 5).

Table 5. Usability issues (n = 10).

Usability issues	Details	Methods
Neglect of user groups and diseases	SCs could not support diverse user groups (e.g., transgender individuals) or diseases ²³	Interviews ²³ , app review analysis ²³
Input limitations	Users perceived difficulties in inputting the information of their diseases completely (62%) ²⁰ , inputting sufficient patient history information ²³ , describing disease dimensions (e.g., frequency) ^{23,44} , and entering medical terms that could be recognized by the SCs ²³	Questionnaires (self-created) ²⁰ , think aloud ⁴⁴ , interviews ²³ , app review analysis ²³
Problematic questions	Users perceived that the probing questions of SCs were seemingly irrelevant based and were presented in an unreasonable sequence ²³	Interviews ²³ , app review analysis ²³
Interpretation issues	Users encountered challenges in understanding medical jargon ⁷ , complex questions ²³ , and diagnostic reports ^{21,31}	Interviews ^{7,23} , app review analysis ²³ , log analysis ²¹ , questionnaires (adapted from previous literature) ³¹
Overwhelming amounts of information	Users complained of too much information on the welcome page ⁴⁴ , excessive subsections to fill in ⁴⁴ , too many pieces of information presented at once ²⁶ , and redundant questions ²³	App review analysis ²³ , interviews ^{23,26} , usability testing ²⁶ , think aloud ⁴⁴

Safety/security: Three studies highlighted that using SCs may have potential risks for both the healthcare system and consumers^{22,28,33}. For instance, the specificity of the SC used for influenza-like illness is poor, which may lead more users to go to the emergency department, thereby putting pressure on healthcare systems²². Likewise, since most health consumers are laypersons who may not have sufficient medical knowledge, they cannot discern the correctness of SCs' recommendations. Using SCs uncritically may compromise users' health and safety^{28,33}. For example, one paper found only 53% to 61% of users with depression reported by the focal SC discussed their depression with medical professionals²⁸. This might result in untreated depression. Two papers emphasized the significance of data security^{35,44}, calling for more information about how SCs encrypt³⁵, anonymize³⁵, and store data⁴⁴.

Functionality: Two studies examined the functions of SCs using an app feature analysis²³, an app review analysis^{23,39}, and semi-structured interviews²³. The first study examined 11 apps released in the United States via the Apple App store and the Google Play store²³. The researchers found most apps support five procedures: creating a patient history for users, evaluating symptom input by users, providing recommendations, ordering further tests, and providing follow-up treatment (e.g., contacting healthcare practitioners). The other study showcased the multiple functions users desired, such as mood tracking, journaling, and providing educational materials³⁹.

(3) Context

Four studies^{20,21,38,40} mentioned the context within which consumers interacted with SCs. They reported that SCs were used to address myriad symptoms. Two of them reported users mostly used SCs to address non-emergency care issues, such as common cold symptoms⁴⁰, gastroenterology²¹, and dermatology²¹, while fewer individuals used SCs in emergent situations²¹. The SC DoctorBot was also used for diseases that may carry social stigma²¹. One of the studies found that the most common symptoms reported by users are pain and abdominal functioning³⁸.

Design suggestions for improving the UX of SCs. Based on their findings, the papers included in our corpus offered design recommendations for improving the UX of SCs. The details are presented in Table 6.

Table 6. Design suggestions.

Aspects	Design suggestions
Accuracy	Increase the accuracy of SCs by considering the probability of specific conditions ¹⁷ and the uniqueness of a variety of symptom clusters ¹⁷
Safety	<ul style="list-style-type: none"> • Supply extra healthcare information²⁰ and provide guidance as well as warnings to the public^{29,33} • Enact relevant policies or legislation to regulate online SC consultation, such as asking SC suppliers to clarify the responsibility of SCs⁴⁰, pre-test SCs⁴², and develop a consumer protection policy⁴³
Functions	<ul style="list-style-type: none"> • Provide functions that are similar to those of in-person medical visits (e.g., offering customized symptoms tracking, contacting medical professionals, and providing medications information)^{23,33} • Provide a comprehensive health profile and follow-up treatment²³
Broader healthcare information	<ul style="list-style-type: none"> • Introduction to SC functions, the consultation process²¹, and prediction accuracy²¹, information about diagnostic conditions³⁵, Treatment options^{21,35} • Endorsement of established authorities³³ • Explanation of data sources^{21,31}, probing questions^{31,34}, and the decision model^{21,31,34} • Treatment options^{21,35} and links to other information sources²¹
Usability	<ul style="list-style-type: none"> • Have flexible input functions (e.g., voice recordings²¹ and approximate string matching function²³) • Convey human empathy^{26,30,32} and use layperson language^{23,29}, and provide a good information architecture²⁶ (i.e., present key information logically instead of showing multiple pieces of complex information at once) • Use readable fonts as well as vivid color²⁶, visual summaries (e.g., tables and figures)⁴¹, and navigation aids⁴¹

Discussion

This systematic review is the first to address the UX of SCs. We reported on and analyzed the characteristics of the published literature, SC users' characteristics, the motivations of individuals to use SCs, the relevant aspects of the UX of SCs, and design implications. In this section, we discuss the implications of our results for future research.

Overall, the UX studies in our corpus focus primarily on the characteristics of the SCs rather than investigating consumers' internal states. The included studies put more emphasis on evaluating rather than designing SCs centering around users' needs. In healthcare, patient-centered care (i.e., "providing care that is respectful of and responsive to individual patient preferences, needs, and values"⁴⁵) is always highlighted. When designing healthcare technologies, users' expectations and needs should also be centered and considered⁴⁶. However, insufficient attention has been paid to fully understanding users' needs for SCs, especially their concerns and expectations in everyday practices. Our findings show that most studies have explored the use of SCs in laboratory or otherwise controlled settings (n=16). While we acknowledge the importance of laboratory experiments, we suggest that studies that are deployed in the wild are needed to capture individuals' unanticipated use of SCs and reveal UX issues that only arise in-situ⁴⁷.

In addition, patient-centered care suggests giving respect, showing emotional support, communication skills, and education, and offering explanations^{48,49}. However, whether and how this line of guidelines should be applied to SC design is still unclear. First, our systematic review found that few studies focus on how to provide emotional support to SC users and improve their hedonic experience. Only four studies highlighted that SCs should convey human empathy^{26,30,32}. Little is known about how to deliver emotional support to users. Moreover, although existing healthcare research suggests that humanoid features of healthcare technologies may improve communication and therefore UX^{50,51}, little research in our corpus put forward recommendations for how to embed humanoid features in SCs to improve the hedonic experience. Further research should be conducted to investigate what level of empathy should be provided and how the interface design of SCs can be improved. Second, though four studies we reviewed investigated the explanations^{21,31,34,35}, which emphasized the importance of providing explanations (e.g., explanations for presented questions and diagnostic reports^{31,34}) to improve the transparency of SCs and users' trust, it is still unclear how we

should deliver explanations to users. Future research might explore how to give users more control and customization over the offerings of explanations³¹.

Furthermore, few studies in our corpus focus on the context of use. Existing research has highlighted the significance of the context in which technology is embedded⁵², recognizing that technology is integrated into everyday practices⁵². Health information needs are especially likely to vary across locations⁵³. It is crucial to consider the geographic and other situated contexts within which the user interactions with SC happen. For instance, our review shows that most papers on SCs are Western in origin and their studies were conducted in developed countries. Few studies were conducted in developing countries such as Vietnam (n = 1). Since healthcare systems vary across geographic contexts (from country to country^{54,55} and from rural to urban areas⁵⁶), it is possible that the findings of our reviewed papers would change if these studies were conducted in new geographic contexts. Our review indicates that future research should be including more diverse geographic contexts, such as undeveloped and developing countries.

Implications for future SC research. **More qualitative and mixed-methods research is needed.** A key takeaway of our review is that the majority of papers used quantitative methods (n = 14). These studies measured UX via questionnaires or through clinical vignettes. Few of the studies we reviewed investigated in depth why individuals decide to use, trust, and are satisfied with SCs. We call for more qualitative research exploring the deeper reasons behind users' preferences. In addition, mixed methods are common in healthcare research as they can improve the research rigor and offer a more comprehensive portrayal of the studied phenomenon^{57,58}. Though the qualitative studies and mixed-method studies in our corpus revealed how individuals' trust in SCs, SCs' usability, and SCs' accuracy were measured, there were only a few such studies. To validate the qualitative data and produce more generalizable results, future research needs to complement the qualitative data with quantitative data from large samples⁵⁹.

More diverse user groups should be explored. Our review suggests that future UX research should consider diverse user groups. One paper we reviewed argued that when designing new SCs, programmers should consider a diversity of user groups, as existing SCs lack support for special user groups, such as transgender individuals²³. This is in line with the principles of inclusive design. Inclusive design calls for the consideration of all users' needs without adaptation⁶⁰. Prior research has highlighted the importance of inclusiveness in healthcare services⁶¹. The reviewed studies reported on some user characteristics, such as age, gender, education level, health literacy, and household income. However, the number of studies we captured is quite limited (n = 4), and there is a lack of research on how users with different characteristics (e.g., different age, gender, and health literacy levels) experience SCs. When developing healthcare apps, all health literacy levels should be thoroughly considered since individuals with different health literacy levels may have different needs⁶². For example, older adults may have more barriers to using healthcare technologies due to limited health and technology literacy⁶³. The gender differences may lead to different acceptance of SCs as people of different genders may have distinct attitudes toward healthcare applications⁶⁴. It is thus important to uncover the diverse needs and experiences of users of different age, gender, and health literacy levels. In the future, researchers might look into how special user groups experience SCs.

More focus on safety and ethics is needed. In healthcare, safety and security problems need to be prioritized since user information is sensitive⁶⁵. Recent research on the reliability and accountability of health applications has pointed out that there are no clear measures to assess and certify SCs and guarantee their reliability and quality⁶⁶, which may lead to medical errors and cause unintended harm to users due to the nature of smartphones⁶⁷. Several papers in our review also highlighted potential risks of using SCs, including poor specificity²² and data security issues^{35,44}. However, there were few suggestions regarding how to ensure the safety of current SCs. To deliver safe healthcare services, it is also necessary to consider the ethical challenges or the moral considerations⁶⁸. As noted in the included studies^{7,35}, users were always concerned about their health data privacy. However, ethical issues regarding SC's anonymity and data collection/storage were not considered in the included studies. Few studies in our corpus offered recommendations for solving ethical challenges and enacting new policies, such as establishing a data protection policy⁴³ or testing SCs before release^{7,39}. It is still unclear whether the developers of SC apps have sufficient credentials, disclosed enough information to ensure transparency, and guaranteed data security. Furthermore, previous research has pointed out that AI systems need regulation and strict testing before being released to the public⁶⁹ since AI technologies may introduce risks due to their opaque algorithms⁷⁰. Although eight papers in the corpus described their focal SCs as AI-driven, few put forward policies to ensure users' safety. Thus, future research should explore the kinds of policies that should be made to govern SCs, addressing SCs' accountability, data privacy and security.

Limitations of the review process. The present paper has several methodological shortcomings. First, papers not written in English were excluded. Second, due to resource constraints, we may have missed relevant contemporary papers spurred by the fast-moving development of SCs. In fact, the ACM Digital Library's search engine limits

database users to the first 2,000 papers that appear in the search engine's index, even if they modify their search keywords several rounds to acquire as many as possible articles as we did. To limit the risk of missing papers, we searched multiple databases as well as Google Scholar and conducted a backward–forward reference check.

Conclusion

This systematic review summarizes trends in the UX of SCs. Despite the limitations, our review contributes to the SC research by providing a thorough overview of prior UX research and discussing factors that influence SC usage. Our review presents SC users' characteristics, individuals' motivations to use SCs, aspects of the UX of SCs, and design implications for SCs. More user studies are required to narrow the research gaps identified in our study.

References

1. Millenson ML, Baldwin JL, Zipperer L, Singh H. Beyond Dr. Google: the evidence on consumer-facing digital tools for diagnosis. *Diagnosis*. 2018;5(3):95–105.
2. Ada in the Apple App Store [Internet]. Available from: <https://apps.apple.com/us/app/ada-check-your-health/id1099986434>
3. Lupton D, Jutel A. 'It's like having a physician in your pocket!' A critical analysis of self-diagnosis smartphone apps. *Soc Sci Med* [Internet]. 2015;133:128–35. Available from: <http://dx.doi.org/10.1016/j.socscimed.2015.04.004>
4. Fraser H, Coiera E, Wong D. Safety of patient-facing digital symptom checkers. *Lancet* [Internet]. 2018;392(10161):2263–4. Available from: [http://dx.doi.org/10.1016/S0140-6736\(18\)32819-8](http://dx.doi.org/10.1016/S0140-6736(18)32819-8)
5. Prakash AV, Das S. Intelligent Conversational Agents in Mental Healthcare Services: A Thematic Analysis of User Perceptions. *Pacific Asia J Assoc Inf Syst*. 2020;12(2):1.
6. Hassenzahl M, Tractinsky N. User experience-a research agenda. *Behav Inf Technol*. 2006;25(2):91–7.
7. Aboueid S, Meyer S, Wallace JR, Mahajan S, Chaurasia A. Young Adults' Perspectives on the Use of Symptom Checkers for Self-Triage and Self-Diagnosis: Qualitative Study. *JMIR Public Heal Surveill*. 2021;7(1):e22637.
8. Mueller J, Jay C, Harper S, Davies A, Vega J, Todd C. Web use for symptom appraisal of physical health conditions: a systematic review. *J Med Internet Res*. 2017;19(6):e6755.
9. Farnood A, Johnston B, Mair FS. A mixed methods systematic review of the effects of patient online self-diagnosing in the 'smart-phone society' on the healthcare professional-patient relationship and medical authority. *BMC Med Inform Decis Mak*. 2020;20(1):1–14.
10. Chambers D, Cantrell AJ, Johnson M, Preston L, Baxter SK, Booth A, et al. Digital and online symptom checkers and health assessment/triage services for urgent health problems: systematic review. *BMJ Open*. 2019;9(8):e027743.
11. Gottlieb K, Petersson G. Limited evidence of benefits of patient operated intelligent primary care triage tools: findings of a literature review. *BMJ Heal care informatics*. 2020;27(1).
12. Aboueid S, Liu RH, Desta BN, Chaurasia A. The Use of Artificially Intelligent Self-Diagnosing Digital Platforms by the General Public: Scoping Review. *JMIR Med informatics*. 2019;7(2):e13445.
13. Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol*. 2018;18(1):1–7.
14. Powley L, McIlroy G, Simons G, Raza K. Are online symptoms checkers useful for patients with inflammatory arthritis? *BMC Musculoskelet Disord*. 2016;17(1):1–6.
15. Bisson LJ, Komm JT, Bernas GA, Fineberg MS, Marzo JM, Rauh MA, et al. How accurate are patients at diagnosing the cause of their knee pain with the help of a web-based symptom checker? *Orthop J Sport Med*. 2016;4(2):2325967116630286.
16. Bisson LJ, Komm JT, Bernas GA, Fineberg MS, Marzo JM, Rauh MA, et al. Accuracy of a computer-based diagnostic program for ambulatory patients with knee pain. *Am J Sports Med*. 2014;42(10):2371–6.
17. Hageman MGJS, Anderson J, Blok R, Bossen JKJ, Ring D. Internet self-diagnosis in hand surgery. *Hand*. 2015;10(3):565–9.
18. Knitza J, Mohn J, Bergmann C, Kampylafka E, Hagen M, Bohr D, et al. Accuracy, patient-perceived usability, and acceptance of two symptom checkers (Ada and Rheport) in rheumatology: interim results from a randomized controlled crossover trial. *Arthritis Res Ther*. 2021;23(1):1–7.
19. Cross S, Mourad A, Zuccon G, Koopman B. Search engines vs. symptom checkers: A comparison of their effectiveness for online health advice. In: *Proceedings of the Web Conference 2021*. 2021. p. 206–16.
20. Verzantvoort NCM, Teunis T, Verheij TJM, van der Velden AW. Self-triage for acute primary care via a smartphone application: Practical, safe and efficient? *PLoS One*. 2018;13(6):e0199284.
21. Fan X, Chao D, Zhang Z, Wang D, Li X, Tian F. Utilization of self-diagnosis health chatbots in real-world

- settings: case study. *J Med Internet Res.* 2021;23(1):e19928.
22. Price RA, Fagbuyi D, Harris R, Hanfling D, Place F, Taylor TB, et al. Feasibility of web-based self-triage by parents of children with influenza-like illness: a cautionary tale. *JAMA Pediatr.* 2013;167(2):112–8.
 23. You Y, Gui X. Self-Diagnosis through AI-enabled Chatbot-based Symptom Checkers: User Experiences and Design Considerations. In: *AMIA Annual Symposium Proceedings*. American Medical Informatics Association; 2020.
 24. Aboueid S, Meyer SB, Wallace JR, Mahajan S, Nur T, Chaurasia A. Use of symptom checkers for COVID-19-related symptoms among university students: a qualitative study. *BMJ Innov.* 2021;7(2).
 25. Marco-Ruiz L, Bønes E, de la Asunción E, Gabarron E, Aviles-Solis JC, Lee E, et al. Combining multivariate statistics and the think-aloud protocol to assess Human-Computer Interaction barriers in symptom checkers. *J Biomed Inform.* 2017;74:104–22.
 26. Ponnada S. Reimagining the COVID-19 Digital Experience: The Value of User Empowerment and Accessibility in Risk Communication. In: *Proceedings of the 38th ACM International Conference on Design of Communication*. 2020. p. 1–3.
 27. Schrage JD, Schuler K, Isakov AP, Wright DW, Yaffee AQ, Jacobson KL, et al. Development and usability testing of a web-based COVID-19 Self-triage platform. *West J Emerg Med.* 2020;21(5):1054.
 28. Reilly N, Austin M-P. Attitudes and Engagement of Pregnant and Postnatal Women With a Web-Based Emotional Health Tool (Mummatters): Cross-sectional Study. *J Med Internet Res.* 2021;23(3):e18517.
 29. Nieroda ME, Lophatananon A, McMillan B, Chen L-C, Hughes J, Daniels R, et al. Online decision support tool for personalized cancer symptom checking in the community (REACT): acceptability, feasibility, and usability study. *JMIR cancer.* 2018;4(2):e10073.
 30. Daher K, Casas J, Khaled OA, Mugellini E. Empathic chatbot response for medical assistance. In: *Proceedings of the 20th ACM International Conference on Intelligent Virtual Agents*. 2020. p. 1–3.
 31. Tsai C-H, You Y, Gui X, Kou Y, Carroll JM. Exploring and Promoting Diagnostic Transparency and Explainability in Online Symptom Checkers. In: *Proceedings of the 2021 CHI Conference on Human Factors in Computing Systems*. 2021. p. 1–17.
 32. Li W, Fan X, Zhu H, Wu J, Teng D. Research on the Influencing Factors of User Trust Based on Artificial Intelligence Self Diagnosis System. In: *Proceedings of the ACM Turing Celebration Conference-China*. 2020. p. 197–202.
 33. You Y, Kou Y, Ding X, Gui X. The Medical Authority of AI: A Study of AI-enabled Consumer-Facing Health Technology. In: *Proceedings of the 2021 CHI Conference on Human Factors in Computing Systems*. 2021. p. 1–16.
 34. HWANG Y, KIM T, KIM J, LEE J, HONG H. Leveraging challenges of an algorithm-based symptom checker on user trust through explainable AI. 2018;
 35. Powell J. Trust me, I'm a chatbot: how artificial intelligence in health care fails the turing test. *J Med Internet Res.* 2019;21(10):e16222.
 36. Miller S, Gilbert S, Virani V, Wicks P. Patients' Utilization and Perception of an Artificial Intelligence--Based Symptom Assessment and Advice Technology in a British Primary Care Waiting Room: Exploratory Pilot Study. *JMIR Hum factors.* 2020;7(3):e19713.
 37. Meyer AND, Giardina TD, Spitzmueller C, Shahid U, Scott TMT, Singh H. Patient Perspectives on the Usefulness of an Artificial Intelligence--Assisted Symptom Checker: Cross-Sectional Survey Study. *J Med Internet Res.* 2020;22(1):e14679.
 38. Winn AN, Somai M, Fergestrom N, Crotty BH. Association of use of online symptom checkers with patients' plans for seeking care. *JAMA Netw open.* 2019;2(12):e1918561–e1918561.
 39. DeForte S, Huang Y, Bourgeois T, Hussain S-A, Lin S. The association between app-administered depression assessments and suicidal ideation in user comments: retrospective observational study. *JMIR mHealth uHealth.* 2020;8(8):e18392.
 40. Nijland N, Cranen K, Boer H, van Gemert-Pijnen JEW, Seydel ER. Patient use and compliance with medical advice delivered by a web-based triage system in primary care. *J Telemed Telecare.* 2010;16(1):8–11.
 41. Luger TM, Houston TK, Suls J. Older adult experience of online diagnosis: results from a scenario-based think-aloud protocol. *J Med Internet Res.* 2014;16(1):e2924.
 42. Lanseng EJ, Andreassen TW. Electronic healthcare: a study of people's readiness and attitude toward performing self-diagnosis. *Int J Serv Ind Manag.* 2007;
 43. Hua TV, Hou Y. Factors That Influence the Intention to Use Self-Diagnosis Apps in Vietnam. 2020;
 44. Marco-Ruiz L, Bønes E, de la Asunción E, Gabarron E, Aviles-Solis JC, Lee E, et al. Combining multivariate statistics and the think-aloud protocol to assess Human-Computer Interaction barriers in symptom checkers. *J Biomed Inform.* 2017;74:104–22.

45. Richardson WC, Berwick DM, Bisgard JC, Bristow LR, Buck CR, Cassel CK. Crossing the quality chasm: a new health system for the 21st century. Washington, DC: National Academy Press; 2001.
46. Cazabon D, Pande T, Sen P, Daftary A, Arsenault C, Bhatnagar H, et al. User experience and patient satisfaction with tuberculosis care in low-and middle-income countries: a systematic review. *J Clin Tuberc other Mycobact Dis*. 2020;19:100154.
47. Tolmie P, Crabtree A. Deploying research technology in the home. In: Proceedings of the 2008 ACM conference on Computer supported cooperative work. 2008. p. 639–48.
48. Smith SP, Barefield AC. Patients meet technology: the newest in patient-centered care initiatives. *Health Care Manag (Frederick)*. 2007;26(4):354–62.
49. Raja S, Hasnain M, Vadakumchery T, Hamad J, Shah R, Hoersch M. Identifying elements of patient-centered care in underserved populations: a qualitative study of patient perspectives. *PLoS One*. 2015;10(5):e0126708.
50. Liu B, Sundar SS. Should machines express sympathy and empathy? Experiments with a health advice chatbot. *Cyberpsychology, Behav Soc Netw*. 2018;21(10):625–36.
51. Li M, Mao J. Hedonic or utilitarian? Exploring the impact of communication style alignment on user's perception of virtual health advisory services. *Int J Inf Manage*. 2015;35(2):229–43.
52. Tolmie P, Pycok J, Diggins T, MacLean A, Karsenty A. Unremarkable computing. In: Proceedings of the SIGCHI conference on Human factors in computing systems. 2002. p. 399–406.
53. Boulos MNK. Location-based health information services: a new paradigm in personalised information delivery. *Int J Health Geogr*. 2003;2(1):1–11.
54. Drechsler D, Jutting J. Different countries, different needs: the role of private health insurance in developing countries. *J Health Polit Policy Law*. 2007;32(3):497–534.
55. Reibling N, Ariaans M, Wendt C. Worlds of healthcare: a healthcare system typology of OECD countries. *Health Policy (New York)*. 2019;123(7):611–20.
56. Qiu Y, Lu W, Guo J, Sun C, Liu X. Examining the Urban and Rural Healthcare Progress in Big Cities of China: Analysis of Monitoring Data in Dalian from 2008 to 2017. *Int J Environ Res Public Health*. 2020;17(4):1148.
57. Moffatt S, White M, Mackintosh J, Howel D. Using quantitative and qualitative data in health services research—what happens when mixed method findings conflict?[ISRCTN61522618]. *BMC Health Serv Res*. 2006;6(1):1–10.
58. Irizarry T, Shoemake J, Nilsen ML, Czaja S, Beach S, Dabbs AD. Patient portals as a tool for health care engagement: a mixed-method study of older adults with varying levels of health literacy and prior patient portal use. *J Med Internet Res*. 2017;19(3):e7099.
59. Bautista JR, Lin TTC, Theng Y-L. How and why users use social TV systems? A systematic review of user studies. In: 2016 49th Hawaii International Conference on System Sciences (HICSS). IEEE; 2016. p. 3868–77.
60. Abascal J, Nicolle C. Moving towards inclusive design guidelines for socially and ethically aware HCI. *Interact Comput*. 2005;17:484–505.
61. Beniuk C, Ward J, Clarkson PJ. Applying inclusive design principles in the assessment of healthcare services. In: Proceedings of conference on design. 2011. p. 22–36.
62. Kim H, Goldsmith J V, Sengupta S, Mahmood A, Powell MP, Bhatt J, et al. Mobile health application and e-health literacy: opportunities and concerns for cancer patients and caregivers. *J Cancer Educ*. 2019;34(1):3–8.
63. Harris L, Thomas V, Fox M. The influence of health care literacy on the use of PHRs among older adults. *Medsurg Nurs*. 2015;24(4):283–6.
64. Kuo IH, Rabindran JM, Broadbent E, Lee YI, Kerse N, Stafford RMQ, et al. Age and gender factors in user acceptance of healthcare robots. In: RO-MAN 2009-The 18th IEEE International Symposium on Robot and Human Interactive Communication. IEEE; 2009. p. 214–9.
65. Bhuyan SS, Kim H, Isehunwa OO, Kumar N, Bhatt J, Wyant DK, et al. Privacy and security issues in mobile health: Current research and future directions. *Heal Policy Technol [Internet]*. 2017;6(2):188–91. Available from: <http://dx.doi.org/10.1016/j.hlpt.2017.01.004>
66. Grundy QH, Wang Z, Bero LA. Challenges in Assessing Mobile Health App Quality. *Am J Prev Med [Internet]*. 2019;51(6):1051–9. Available from: <http://dx.doi.org/10.1016/j.amepre.2016.07.009>
67. Pine KH, Bossen C, Chen Y, Ellingsen G, Grisot M, Mazmanian M, et al. Data Work in Healthcare: Challenges for Patients, Clinicians and Administrators. In: Companion of the 2018 ACM Conference on Computer Supported Cooperative Work and Social Computing. 2018. p. 433–9.
68. Sun TQ, Medaglia R. Mapping the challenges of Artificial Intelligence in the public sector: Evidence from public healthcare. *Gov Inf Q*. 2019;36(2):368–83.
69. McCartney M. Margaret McCartney : AI in medicine must be rigorously tested. *Bmj [Internet]*. 2018;361:k1752. Available from: <http://dx.doi.org/doi:10.1136/bmj.k1752>
70. Macrae C. Governing the safety of artificial intelligence in healthcare. *BMJ Qual Saf*. 2019;28(6):495–8.