# One parameter generalization of BW inequality and its application to open quantum dynamics

Dariusz Chruściński<sup>a</sup>, Gen Kimura<sup>b</sup>, Hiromichi Ohno<sup>c,\*</sup>, Tanmay Singal<sup>a</sup>

<sup>a</sup>Institute of Physics, Faculty of Physics, Astronomy and Informatics Nicolaus Copernicus University, Grudziądzka 5/7, 87–100 Toruń, Poland

<sup>c</sup>Department of Mathematics, Faculty of Engineering, Shinshu University, 4-17-1 Wakasato, Nagano 380-8553, Japan.

### Abstract

In this paper, we introduce a one parameter generalization of the famous Böttcher-Wenzel (BW) inequality in terms of a q-deformed commutator. For  $n \times n$  matrices A and B, we consider the inequality

$$\operatorname{Re}\langle [B, A], [B, A]_q \rangle \le c(q) ||A||^2 ||B||^2,$$

where  $\langle A, B \rangle = \operatorname{tr}(A^*B)$  is the Hilbert-Schmidt inner product, ||A|| is the Frobenius norm, [A, B] = AB - BA is the commutator, and  $[A, B]_q = AB - qBA$  is the q-deformed commutator. We prove that when n = 2, or when A is normal with any size n, the optimal bound is given by

$$c(q) = \frac{(1+q) + \sqrt{2(1+q^2)}}{2}.$$

We conjecture that this is also true for any matrices, and this conjecture is perfectly supported for n up to 15 by numerical optimization. When q = 1, this inequality is exactly BW inequality. When q = 0, this inequality leads the sharp bound for the r-function which is recently derived for the application to universal constraints of relaxation rates in open quantum dynamics.

<sup>&</sup>lt;sup>b</sup>College of Systems Engineering and Science, Shibaura Institute of Technology, Saitama 330-8570, Japan

<sup>\*</sup>Corresponding author

Email addresses: darch@fizyka.umk.pl (Dariusz Chruściński), gen@shibaura-it.ac.jp (Gen Kimura), h\_ohno@shinshu-u.ac.jp (Hiromichi Ohno), tanmaysingal@gmail.com (Tanmay Singal)

*Keywords:* Hilbert-Schmidt Inner Product, Frobenius Norm, Commutator, BW inequality,

### 1. Introduction

For any complex matrices  $A, B \in M_n(\mathbb{C})$ , the sharp bound for the norm of commutator is known:

$$\|[A,B]\|^2 \le 2\|A\|^2 \|B\|^2, \tag{1}$$

where [A, B] := AB - BA denotes the commutator and the matrix norm is the Frobenius norm  $||A|| = \sqrt{\operatorname{tr}(A^*A)}$ . The inequality is often called the Böttcher-Wenzel (BW) inequality because of the following history: The inequality was first conjectured by Böttcher and Wenzel in [2] giving a proof for real 2 × 2 matrices and also for normal matrices. Later, Lásló proved BW inequality for 3 × 3 real matrices [12], Lu [13] and, independently, Wong and Jin [15] proved it for  $n \times n$  real matrices, after which the complex case was proved by Böttcher and Wenzel in [3].

On the other hand, motivated by a problem in the field of open quantum systems [1], we have introduced the following real-valued functional in [4]:

$$r(A,B) = \frac{1}{2} \left( \langle [B,A], BA \rangle + \langle [B,A^*], BA^* \rangle \right)$$
(2)

for any  $A, B \in M_n(\mathbb{C})$  where  $A^*$  is the Hermitian conjugation of A and  $\langle A, B \rangle = tr(A^*B)$  is the Hilbert-Schmidt inner product. The sharp bound for the *r*-function was shown in [4]:

$$r(A,B) \le \frac{1+\sqrt{2}}{2} \|A\|^2 \|B\|^2.$$
 (3)

This has been used to derive a universal constraint [6, 10, 11] for relaxation rates which is satisfied by any quantum Markovian dynamics described by completely positive dynamical semigroup, or GKLS master equation [7]. For this reason, we called the functional (2) the *r*-function, where "*r*" stands for relaxation. In [4], we also observed a close relationship between BW inequality and inequality (3). If B is normal, r-function reduces to the commutator:

$$r(A,B) = \frac{1}{2} \| [A,B] \|^2$$

Hence, the direct application of BW inequality leads the bound

$$r(A,B) \le ||A||^2 ||B||^2.$$

However, this is only true for a normal matrix B and does not hold for general matrices. On the other hand, applying the triangle inequality, Schwarz inequality, the norm inequality  $||AB|| \leq ||A|| ||B||$  and BW inequality for (2) implies the following general bound:

$$r(A,B) \le \sqrt{2} \|A\|^2 \|B\|^2.$$

However, this in turn is a loose inequality. In the end, the sharp bound for general matrices is given by (3) and, therefore the relation between BW inequality and (3) is not yet fully understood.

The purpose of this paper is to provide one answer to this problem by introducing the following function (referred in this paper *f*-function) for complex matrices  $A, B \in M_n(\mathbb{C})$  and one parameter  $q \in \mathbb{R}$ :

$$f(A, B; q) := \operatorname{Re}\langle [B, A], [B, A]_q \rangle$$
  
= tr  $\left( A^* B^* B A + q B^* A^* A B - \frac{1+q}{2} (A^* B^* A B + B^* A^* B A) \right), (4)$ 

where  $[A, B]_q := AB - qBA$  is the q-deformed commutator [8, 14]. When q = 1, the f-function is exactly the norm squared of the commutator:

$$f(A, B; 1) = \|[A, B]\|^2.$$
(5)

(See also [5] for another generalization  $||[A, B]_q||^2$  and its norm bound.) When q = 0, the *f*-function and *r*-function have the following relation:

$$r(A,B) = \frac{f(A,B;0) + f(A^*,B;0)}{2}.$$
(6)

In this sense, the f-function connects the r-function and the norm squared of the commutator through one parameter q.

We consider the sharp bound for the f-function, and obtain a partial answer.

**Theorem 1.** Let A and B be  $n \times n$  complex matrices. If A is a normal matrix or n = 2, then the inequality

$$f(A, B; q) \le c(q) \|A\|^2 \|B\|^2$$
(7)

holds, where

$$c(q) = \frac{(1+q) + \sqrt{2(1+q^2)}}{2}.$$

Moreover, this inequality is sharp, i.e., there are non-zero matrices A and B at which the bound is attained.

We conjecture that the theorem is also true for any matrices  $A, B \in M_n(\mathbb{C})$ . This is numerically supported for n up to 15 (See Fig. 1).

Importantly, the inequality (7) integrates BW inequality and the inequality (3) as follows. First, inequality (7) with q = 1 recovers BW inequality by noting c(1) = 2 and using relation (5). Hence, inequality (7) is considered as a one-parameter generalization of BW inequality. Second, when q = 0, we have  $c(0) = \frac{1+\sqrt{2}}{2}$ , and therefore, (7) and (6) imply inequality (3) for *r*-function. We also note that, the conjecture (7) for q = -1 is easily shown by noting c(-1) = 1 and  $f(A, B; -1) = ||BA||^2 - ||AB||^2$  and using the norm inequality  $||BA|| \leq ||B||||A||$ .

### 2. Proofs of the main theorem

First, we mention some general facts for the f-function and c(q).

**Proposition 1.** For any unitary matrix  $U \in M_n(\mathbb{C})$  and complex numbers  $\alpha, \beta \in \mathbb{C}$ , the following equations hold:

$$\begin{split} f(UAU^*, UBU^*; q) &= f(A, B; q), \\ f(\alpha A, \beta B; q) &= |\alpha|^2 |\beta|^2 f(A, B; q). \end{split}$$



Figure 1: Numerical evidences of (7) for n = 2, 5, 10, 15: By the quadratic nature of the *f*-function, our conjecture (7) can be rephrased as  $\sup_{A,B\neq 0\in M_n(\mathbb{C})} \frac{f(A,B;q)}{\|A\|^2 \|B\|^2} = c(q)$ . Numerically optimized values (Red points) of the left hand side perfectly coincide with  $c(q) = ((1+q) + \sqrt{2(1+q^2)})/2$  (blue curves).

The proof is obvious and will be omitted. This proposition says that the f-function is unitarily invariant.

In the following, we will write c(q) = c unless there is no confusion. Note that c is bigger than or equal to 1, q and 1 + q for any  $q \in \mathbb{R}$ .

**Lemma 1.** Let  $\varepsilon_1 = \varepsilon_1(q)$  and  $\varepsilon_2 = \varepsilon_2(q)$  be

$$\varepsilon_1 = \begin{cases} 1 & q \ge -1 \\ -1 & q < -1 \end{cases} \text{ and } \varepsilon_2 = \begin{cases} 1 & q \le 1 \\ -1 & q > 1 \end{cases}.$$

Then, the equations

$$2\varepsilon_1 \sqrt{(c-q)(c-1)} = (q+1)$$
 and  $2\varepsilon_2 \sqrt{c(c-q-1)} = (1-q)$ 

hold.

[Proof] Since  $\varepsilon_i^2 = 1$  (i = 1, 2), the solution for these equations is  $c = \frac{(1+q) + \sqrt{2(1+q^2)}}{2}$ .

Let  $A = (a_{ij})_{i,j=1}^n$  and  $B = (b_{ij})_{i,j=1}^n$  be arbitrary  $n \times n$  matrices. A straightforward calculation shows

$$c\|A\|^2\|B\|^2 - f(A, B; q)$$
  
=  $c\|A\|^2\|B\|^2 - \sum_{i,j,k,\ell=1}^n \left(a_{ij}\overline{a_{kj}}b_{\ell i}\overline{b_{\ell k}} + qa_{ij}\overline{a_{i\ell}}b_{jk}\overline{b_{\ell k}} - (1+q)\operatorname{Re}(a_{ij}\overline{a_{\ell k}}b_{jk}\overline{b_{i\ell}})\right).$ 

We define sets of indices as

$$\begin{aligned} D_0 &= \{(i, i, i, i) \colon 1 \le i \le n\}, \\ D_1 &= \{(i, j, j, i) \colon 1 \le i, j \le n, i \ne j\}, \\ D_2 &= \{(i, j, k, i) \colon 1 \le i, j, k \le n, j \ne k\}, \\ D_3 &= \{(i, j, j, k) \colon 1 \le i, j, k \le n, i \ne k\} \quad \text{and} \\ D_4 &= \{(i, j, k, \ell) \colon 1 \le i, j, k, \ell \le n\} \setminus (D_1 \cup D_2 \cup D_3) \end{aligned}$$

Note that  $D_1, D_2, D_3$  and  $D_4$  are mutually disjoint and  $D_0 \subset D_4$ . The term  $a_{ij}\overline{a_{kj}}b_{\ell i}\overline{b_{\ell k}}$  has the form  $|a_{**}|^2|b_{**}|^2$ , when i = k. In this case,

$$a_{ij}\overline{a_{ij}}b_{\ell i}\overline{b_{\ell i}} = |a_{ij}|^2 |b_{\ell i}|^2.$$

Therefore,

$$\sum_{i,j,\ell} a_{ij} \overline{a_{ij}} b_{\ell i} \overline{b_{\ell i}} = \sum_{i,j,\ell} |a_{ij}|^2 |b_{\ell i}|^2 = \sum_{(i,j,k,\ell) \in D_0 \cup D_1 \cup D_2} |a_{ij}|^2 |b_{k\ell}|^2.$$

Similarly, the term  $a_{ij}\overline{a_{i\ell}}b_{jk}\overline{b_{\ell k}}$  has the form  $|a_{**}|^2|b_{**}|^2$ , when  $j = \ell$ . Then,

$$\sum_{i,j,k} a_{ij}\overline{a_{ij}}b_{jk}\overline{b_{jk}} = \sum_{i,j,k} |a_{ij}|^2 |b_{jk}|^2 = \sum_{(i,j,k,\ell) \in D_0 \cup D_1 \cup D_3} |a_{ij}|^2 |b_{k\ell}|^2.$$

The term  $a_{ij}\overline{a_{\ell k}}b_{jk}\overline{b_{i\ell}}$  has the form  $|a_{**}|^2|b_{**}|^2$ , only when  $i=j=k=\ell$ .

Combining these, we have

$$\begin{split} c\|A\|^2 \|B\|^2 &- f(A, B; q) \\ = c \sum_{\substack{(i,j,k,\ell) \in D_4 \\ (i,j,k,\ell) \in D_3 \\ (i,j,k,\ell) = (1 - q) \sum_{\substack{(i,j,k,\ell) \in D_1 \\ (i,j,k,\ell) \\ (i,j,k,\ell) \in D_1 \\ (i,j,k,\ell) \\ (i,j,k,\ell) \in D_1 \\ (i,j,k,\ell) \\ (i,j,$$

Lemma 2. The next equation holds:

$$\sum_{\substack{i,k\\i\neq k}} |\sqrt{c-q} \sum_{j=1}^n a_{ij} b_{jk} + \varepsilon_1 \sqrt{c-1} \sum_{\ell=1}^n a_{\ell k} b_{i\ell}|^2$$
$$= (c-1) \sum_{D_2} |a_{ij}|^2 |b_{k\ell}|^2 + (c-q) \sum_{D_3} |a_{ij}|^2 |b_{k\ell}|^2$$
$$+ (c-1) \sum_{\substack{i,j,k,l\\i\neq k, j\neq \ell}} a_{ij} \overline{a_{kj}} b_{\ell i} \overline{b_{\ell k}} + (c-q) \sum_{\substack{i,j,k,l\\i\neq k, j\neq \ell}} a_{ij} \overline{a_{i\ell}} b_{jk} \overline{b_{\ell k}}$$
$$+ (1+q) \operatorname{Re} \sum_{\substack{i,j,k,l\\i\neq k}} a_{ij} \overline{a_{\ell k}} b_{jk} \overline{b_{i\ell}}.$$

[Proof] Using the equation  $\varepsilon_1 2\sqrt{(c-q)(c-1)} = (1+q)$  obtained in Lemma 1, we can prove this by straightforward calculation.

By this lemma, we can calculate  $c\|A\|^2\|B\|^2-f(A,B;q)$  as

$$\begin{split} c\|A\|^2 \|B\|^2 &- f(A, B; q) \\ &= \sum_{\substack{i,k\\i \neq k}} |\sqrt{c-q} \sum_j a_{ij} b_{jk} + \varepsilon_1 \sqrt{c-1} \sum_{\ell} a_{\ell k} b_{i\ell}|^2 \\ &+ c \sum_{D_4} |a_{ij}|^2 |b_{k\ell}|^2 + (c-1-q) \sum_{D_1} |a_{ij}|^2 |b_{k\ell}|^2 \\ &- c \sum_{\substack{i,j,k,\ell\\i \neq k, j \neq \ell}} (a_{ij} \overline{a_{kj}} b_{\ell i} \overline{b_{\ell k}} + a_{ij} \overline{a_{i\ell}} b_{jk} \overline{b_{\ell k}}) \\ &- \sum_{\substack{i,j,k\\i \neq k}} a_{ij} \overline{a_{kj}} b_{ji} \overline{b_{jk}} - q \sum_{\substack{i,j,\ell\\j \neq \ell}} a_{ij} \overline{a_{i\ell}} b_{ji} \overline{b_{\ell i}} + (1+q) \sum_{\substack{i,j,\ell\\except\\i=j=\ell}} \operatorname{Re}(a_{ij} \overline{a_{\ell i}} b_{ji} \overline{b_{i\ell}}). \end{split}$$

## 2.1. Proof for n = 2

In this subsection, the proof of Theorem 1 for n = 2 is given. When n = 2, a direct calculation shows:

$$\begin{aligned} |a_{11}\overline{b_{22}} + a_{22}\overline{b_{11}} - a_{12}\overline{b_{12}} - a_{21}\overline{b_{21}}|^2 \\ &= \sum_{D_4 \setminus D_0} |a_{ij}|^2 |b_{k\ell}|^2 - \sum_{\substack{i,j,k,\ell \\ i \neq k, j \neq \ell}} \left( a_{ij}\overline{a_{kj}}b_{\ell i}\overline{b_{\ell k}} + a_{ij}\overline{a_{i\ell}}b_{jk}\overline{b_{\ell k}} - a_{ij}\overline{a_{k\ell}}b_{ji}\overline{b_{\ell k}} \right). \end{aligned}$$

Using this, we have

$$\begin{aligned} c\|A\|^2\|B\|^2 - f(A, B; q) \\ &= \sum_{\substack{i,k\\i \neq k}} |\sqrt{c-q} \sum_j a_{ij} b_{jk} + \varepsilon_1 \sqrt{c-1} \sum_{\ell} a_{\ell k} b_{i\ell}|^2 \\ &+ c|a_{11}\overline{b_{22}} + a_{22}\overline{b_{11}} - a_{12}\overline{b_{12}} - a_{21}\overline{b_{21}}|^2 \\ &+ c \sum_{D_0} |a_{ij}|^2 |b_{k\ell}|^2 + (c-1-q) \sum_{D_1} |a_{ij}|^2 |b_{k\ell}|^2 - c \sum_{\substack{i,j,k,\ell\\i \neq k, j \neq \ell}} a_{ij}\overline{a_{kj}} b_{ji}\overline{b_{jk}} - q \sum_{\substack{i,j,\ell\\j \neq \ell}} a_{ij}\overline{a_{i\ell}} b_{ji}\overline{b_{\ell i}} + (1+q) \sum_{\substack{i,j,\ell\\except\\i=j=\ell}} \operatorname{Re}(a_{ij}\overline{a_{\ell i}} b_{ji}\overline{b_{i\ell}}). \end{aligned}$$
(8)

By (8) and  $2\varepsilon_2\sqrt{c(c-q-1)} = 1-q$  (see Lemma 1), we obtain

$$\begin{aligned} c\|A\|^2\|B\|^2 - f(A, B; q) &- \sum_{\substack{i,k\\i \neq k}} |\sqrt{c - q} \sum_j a_{ij} b_{jk} + \varepsilon_1 \sqrt{c - 1} \sum_{\ell} a_{\ell k} b_{i\ell}|^2 \\ &- c|a_{11}\overline{b_{22}} + a_{22}\overline{b_{11}} - a_{12}\overline{b_{12}} - a_{21}\overline{b_{21}}|^2 \\ &= c\left(|a_{11}|^2|b_{11}|^2 + |a_{22}|^2|b_{22}|^2\right) + (c - 1 - q)\left(|a_{12}|^2|b_{21}|^2 + |a_{21}^2|b_{12}|^2\right) \\ &- 2c\operatorname{Re}\left(a_{11}\overline{a_{22}}b_{11}\overline{b_{22}}\right) - 2(c - 1 - q)\operatorname{Re}\left(a_{12}\overline{a_{21}}b_{21}\overline{b_{12}}\right) \\ &+ (1 - q)\operatorname{Re}\left(-a_{11}\overline{a_{21}}b_{11}\overline{b_{12}} - a_{12}\overline{a_{22}}b_{21}\overline{b_{22}} + a_{11}\overline{a_{12}}b_{11}\overline{b_{21}} + a_{21}\overline{a_{22}}b_{12}\overline{b_{22}}\right) \\ &= \left|\sqrt{c}a_{11}b_{11} - \sqrt{c}a_{22}b_{22} + \varepsilon_2\left(\sqrt{c - 1 - q}a_{12}b_{21} - \sqrt{c - 1 - q}a_{21}b_{12}\right)\right|^2\end{aligned}$$

which implies the inequality (7).

To see that the inequality is sharp, one can easily find non-zero matrices A and B so that all the inequalities above are attained. For instance, let  $a_{11} = \sqrt{c-1}$ ,  $a_{22} = -\varepsilon_1 \sqrt{c-q}$ ,  $b_{12} = 1$  and the others are zero.

### 2.2. Proof for normal matrix

In this subsection, the proof of Theorem 1 for normal matrix is given. Suppose that A is a normal matrix. By the unitary invariance of the f-function, we can assume that A is a diagonal matrix. Letting  $A = \text{diag}[a_1, a_2, \ldots, a_n]$  and  $B = (b_{ij})$ , direct computation shows  $||A||^2 = \sum_i |a_i|^2$ ,  $||B||^2 = \sum_{i,j} |b_{ji}|^2$  and

$$f(A, B; q) = \sum_{i \neq j} |b_{ji}|^2 (|a_i|^2 + q|a_j|^2 - (1+q)\operatorname{Re}(a_i\overline{a_j})).$$

Hence, we have

$$\begin{aligned} c\|A\|^2 \|B\|^2 &- f(A, B; q) \\ &\geq \sum_{i \neq j} |b_{ji}|^2 (c(\sum_k |a_k|^2) - |a_i|^2 - q|a_j|^2 + (1+q) \operatorname{Re}(a_i \overline{a_j})) \\ &\geq \sum_{i \neq j} |b_{ji}|^2 ((c-1)|a_i|^2) + (c-q)|a_j|^2 + (1+q) \operatorname{Re}(a_i \overline{a_j})) \\ &= \sum_{i \neq j} |b_{ji}|^2 |\sqrt{c-1}a_i + \varepsilon_1 \sqrt{c-q} a_j|^2 \ge 0, \end{aligned}$$

where in the equality we have used Lemma 1. Therefore, the bound (7) is shown for a normal matrix A where B is any complex matrix.

One can prove that the inequality is sharp using the same example in the proof for n = 2.

#### 3. Conclusion and discussion

In this paper, we have introduced the f-function (4) which is a one parameter generalization of the norm square of the commutator and also r-function. The conjectured bound (7) for the f-function integrates BW inequality and our previously obtained inequality (3). The conjecture is numerically supported for n up to 15, and is proved when one of the matrix is normal, or for general  $2 \times 2$ matrices.

### Acknowledgements

D.C. was supported by the Polish National Science Centre Project No. 2018/30/A/ST2/00837.

### References

- R. Alicki, K. Lendi, Quantum Dynamical Semigroups and Applications, Springer, Berlin, 1987.
- [2] A. Böttcher, D. Wenzel, How big can the commutator of two matrices be and how big is it typically?, Linear Algebra Appl. 403, (2005) 216-228.
- [3] A. Böttcher, D. Wenzel, The Frobenius norm and the commutator, Linear Algebra Appl. 429, (2008) 1864-1185.
- [4] D. Chruściński, R. Fujii, G. Kimura, H. Ohno, Constraints for the spectra of generators of quantum dynamical semigroups, Linear Algebra Appl. 630, (2021) 293-305.
- [5] D. Chruściński, G. Kimura, H. Ohno, T. Singal, Bounding the Frobenius norm of a q-deformed commutator, Linear Algebra Appl. 646, (2022) 95-106.
- [6] D. Chruscinski, G. Kimura, A. Kossakowski, Y. Shishido, On the universal constraints for relaxation rates for quantum dynamical semigroup, Phys. Rev. Lett. **127**, (2021) 050401.
- [7] V. Gorini, A. Kossakowski, E. C. G. Sudarshan, Completely positive dynamical semigroups of N-level systems, J. Math. Phys. 17, (1976) 821.
- [8] R. Jagannathan, Some introductory notes on quantum groups, quantum algebras, and their appli-cations, arXiv:math-ph/0105002.
- [9] G. Lindblad, On the generators of quantum dynamical semigroups, Comm. Math. Phys. 48, (1976) 119-130.
- [10] G. Kimura, Restriction on relaxation times derived from the Lindbladtype master equations for two-level systems, Phys. Rev. A 66, (2002) 062113.

- [11] G. Kimura, S. Ajisaka, K. Watanabe, Universal Constraints on Relaxation Times for *d*-Level GKLS Master Equations, Open Syst. Inform. Dynam. 24, (2017) 1740009.
- [12] L. Laslo, Proof of Böttcher and Wenzel's conjecture on commutator norms for 3-by-3 matrices, Linear Algebra Appl. 422, (2007) 659–663.
- [13] Z. Lu, Normal Scalar Curvature Conjecture and its applications, Journal of Functional Analysis 261, (2011) 1284–1308.
- [14] S. Majid, Foundations of Quantum Group Theory, Cambridge University Press, Cambridge, 1995.
- [15] S. W. Vong, X. Q. Jin, "Proof of Böttcher and Wenzel's conjecture", Oper. Matrices 2, (2008) 435–442.